High-Throughput DNA melt measurements enable improved models of DNA folding thermodynamics

Yuxi Ke[1], Eesha Sharma[2,3], Hannah K. Wayment-Steele[4], Winston R. Becker[5], Anthony Ho[2], Emil Marklund[2,6], and William J. Greenleaf[2,7*]

[1]Department of Bioengineering, Stanford University, Stanford, CA 94305, USA

[2]Department of Genetics, Stanford University School of Medicine, Stanford, CA 94305, USA

[3]Lamar Health, Palo Alto, CA 94305, USA

[4]Department of Chemistry, Stanford University, Stanford, CA 94305, USA

[5]Program in Biophysics, Stanford University, Stanford, CA 94305, USA

[6]Science for Life Laboratory, Department of Biochemistry and Biophysics, Stockholm University, 17165, Solna, Sweden

[7]Department of Applied Physics, Stanford University, Stanford, CA 94305, USA

*Correspondence: wjg@stanford.edu (W.J.G.)

Abstract

DNA folding thermodynamics are central to many biological processes and biotechnological applications involving base-pairing. Current methods for predicting stability from DNA sequence use nearest-neighbor (NN) models that struggle to accurately capture the diverse sequence-dependency of elements other than Watson-Crick base pairs, likely due to insufficient experimental data. We introduce a massively parallel method–Array Melt–that uses fluorescence-based quenching signals to measure equilibrium stability of millions of DNA hairpins simultaneously on a repurposed Illumina sequencing flow cell. By leveraging this dataset of 27,732 sequences with two-state melting behavior, we derived a refined NUPACK-compatible NN model, a richer parameterization NN model that exhibits higher accuracy, and a graph neural network (GNN) model that identifies relevant interactions within DNA beyond nearest neighbors. All models provide improved accuracy in predicting DNA folding thermodynamics, providing improvements relevant for *in silico* design of qPCR primers, oligo hybridization probes, and DNA origami.

Introduction

The thermodynamics of DNA secondary structure formation is fundamental to understanding diverse biological processes, such as DNA replication[1] and repair[2], and is widely used in biotechnological applications, including the design of PCR primers[3], the engineering of in-situ hybridization (ISH) probes[4], the design of guide sequences for genome editing tools[5], and for DNA nanotechnology[6,7]. Numerous algorithms have been developed to predict DNA secondary structure thermodynamics, many of which rely on the nearest-neighbor (NN) models[8–10].

Briefly, NN models assume that energies from pairs of adjacent base-pairs can be linearly added to determine the overall folding energy of a DNA strand. With this assumption, the "folded state" can be defined as the minimum free energy (MFE) base pairing configuration, identified using dynamic programming[11]. However, the expected probability of finding other secondary structures can also be derived from these calculated free energies based on a calculated partition function. However, NN models often struggle to accurately capture the sheer diversity and complexity of DNA secondary structure motifs, including hairpin loops, mismatches, and bulges. One reason for this limitation is the lack of experimental data spanning those loop features upon which these models are built, leading to inaccuracies in prediction. For example, the most widely used parameter set from SantaLucia et al. 2004 used data from 108 sequences to derive 12 parameters for Watson-Crick (WC) base pairs, and 174 sequences to derive 44 parameters for internal single mismatches[8].

This data bottleneck has been due to the laborious nature of optical melting[12] and differential scanning calorimetry[13] experiments, traditionally considered the gold standards of DNA secondary structure dynamics. Several groups have attempted to overcome this bottleneck, usually with fluorescent melting in bulk solutions on plates[14–16], reaching a throughput on the scale of thousands. Still, due to the extremely large combinatorial DNA sequence space, these works are limited to a specific type of DNA structural element in a fixed sequence scaffold.

To address this data generation bottleneck, we developed a method for systematic, accurate, high-throughput measurements of nucleic acid secondary-structure motif thermal stability, enabling large scale, quantitative measurements of the thermodynamics of nucleic acid secondary structure. We demonstrate that NN thermodynamic parameters inferred from these data are nearly identical to those observed from bulk thermal melting experiments for WC base pairs, but are improved for loop sequence motifs. The improved NN model parameter sets are capable of accurately predicting DNA duplex folding energy on orthogonal unseen data sets. Furthermore, by deploying advanced computational methods including deep learning to model DNA thermodynamics, we develop a state of the art thermodynamic model for DNA hairpin folding with accuracy comparable to measurement uncertainties.

Results

Array melt: A high-throughput technique for quantifying the melting behavior of nucleic acids.

We developed the Array Melt technique, a method that allows for the rapid and comprehensive quantification of DNA folding thermodynamics in high throughput. This technique is based on an Illumina MiSeq chip repurposed for high-throughput measurements[17,18]. To assay melting of a nucleic acid construct, we engineered a common region for annealing a 3'-fluorophore-labeled oligonucleotide to the 5'-end of a hairpin to be investigated, and another for a 5'-quencher labeled oligonucleotide to the 3'-end of the hairpin (**Fig. 1a**). After determining the position and sequence of the variant on the MiSeq chip by sequencing, we anneal a Cy3- and a Black Hole Quencher (BHQ)-labeled oligonucleotide to these respective binding sites. These annealed oligos have predicted melting temperatures of 74°C, much higher than the highest temperature in our experiments (**Table S1**). At room temperature, this fluorophore-quencher pair results in low levels of fluorescence in each cluster, as the bulk of our

hairpin library is folded at room temperature. As the sequence-variable hairpin regions are exposed to increasing temperatures between 20 °C to 60 °C , the distance between Cy3 and BHQ increases, leading to brighter fluorescence signals (**Fig. 1a-b**). Using this measurement scheme, we synthesized a comprehensive library of 42,177 unique hairpin sequences for measurement (**Fig. 1c**). These sequences integrated diverse structural elements into a hairpin scaffold, including Watson-Crick pairing (WC), mismatches, bulges, hairpin loops of various lengths, and other control sequences, including super-stable stem, polynucleotide repeats, and single-strand fluorescence controls (see **Fig. 1c** and Methods).

We performed thorough quality control on the dataset by first requiring variants to accurately fit a two-state model with melting behavior observable within our temperature measurement range (see Methods for our filtering strategy). Total variants that pass these stringent thresholds include 6,393,050 individual melt curves (including technical replicates) spanning 27,732 sequence variants for subsequent analysis with a standard two-state melt model.

Sensitivity and precision of the Array Melt method

To calibrate the amount of fluorescence we observe as a function of distance between Cy3 and BHQ, we evaluated the fluorescence signals of a series of repeat sequence variants designed to systematically increase this distance by incorporating increasing numbers of mono-, di- or tri-nucleotide repeats. The fluorescent signals aggregated over same-length repeats showed a nearly linear response to distance variations within about 8 nucleotides (**Fig. 1d**), and our observed length-fluorescence curve fits well to a theoretical static quenching curve (Methods). We normalized the fluorescent signals to unfolded and folded controls designed in the library (**Fig. S1b**). Three example melt curves in **Fig. 1f** depict increasing observed melting temperatures and increasing GC content of the variable region. Technical replicate measurements from Array Melt data are also highly correlated with correlation R>0.94 (**Fig. S1c**). Melt curves were also highly correlated to anneal curves (R=0.964) confirming that the DNA molecules were measured at equilibrium (**Fig. 1e**). Analyzing our measurement precision (see Methods), we discerned that nearly 90% of the tested variants displayed ΔG_{37} uncertainties within 0.3 kcal/mol, or 0.5 k_BT (**Fig. 1h**). Overall, these data demonstrate that equilibrium, aggregate fluorescence signals obtained from Array Melt method provide highly accurate melt information and can be interpreted as a "molecular ruler" capable of resolving single nucleotide distance variations.

FIGURE 1

Figure 1. High-Throughput thermodynamic measurement of DNA secondary structures using a fluorophore–quencher system.

a. Schematic of the DNA molecules in a folded (quenched) and an unfolded (fluorescent) state.

b. Images of fluorescent DNA clusters immobilized on the sequencing chip surface. Top: Image with only fluorophore-conjugated oligo (Cy3), and with both fluorophore- and quencher-conjugated oligo at increasing temperatures. Bottom: Schematic of DNA molecules in each image. All images are normalized to super-stable stem and repeat control variants for temperature-dependent effects on fluorescence and quenching during the experiment as in **Fig. S1d**.

c. Design of the library variants between the constant-sequence binding sites for fluorophore- and quencher-conjugated oligonucleotides. Red represents scaffold nucleotides constant within each type, and blue the variable ones systematically permuted ('N'). Numbers under each class indicate the number of unique sequences in each class.

d. Fluorescence measurement of control constructs where single strand distance between the fluorophore and the quencher increases at single nucleotide steps. Orange line shows the theoretical fit.

e. Correlation of ΔG_{37} derived from library variants under increasing temperatures (melt curve, x axis) and decreasing temperatures (annealing curve, y axis)

f. Representative examples of melt curves for constructs that vary in GC content.

g. Pearson correlation of ΔG_{37} of replicates of three melt and one annealing curve experiment.

h. Standard error of ΔG_{37} for various construct classes as a function of ΔG_{37}.

SUPP FIGURE 1

Figure S1

    a.   Number of measurements in a single Array Melt experiment.

b. Pearson's R of dH, Tm, and ΔG_{37} between replicates.
c. Controls used to normalize fluorescence across different temperatures, including always unfolded constructs for f_{max} (maximum fluorescence) and folded constructs for f_{min} (minimum fluorescence). Some Watson-Crick curves are randomly selected to visualize after the normalization, colored by their expected ΔG_{37} values predicted by the NN model.
d. Estimation of f_{max} and f_{min} after the initial round of curve fitting. Of all the fitted variants (blue), only a fraction that passed quality control filters (orange) were used to estimate f_{max} and f_{min} parameters used for the next round of refinement fitting. Uncertainties of f_{max} and f_{min} were corrected as a function of number of clusters   per   variant   by   curve   fitting   (right).

UV melting validation

To further validate our method, we compared Array Melt results to established UV melting methods on a cloud lab platform (See Methods). A comparison of melting profiles of random library variants exhibited agreement between the two methods (**Fig. 2a-c,** R=0.85 for ΔG_{37}, or R^2= 0.62 after the linear adjustment described later). Melting temperature is independent of DNA oligo concentration (**Fig. S2a**), suggesting intramolecular folding. Noticeably, we observed a systematic offset between the UV measurements and our Array Melt results, perhaps due to the presence of the hairpins on a flow-cell surface, or due to the additional sequence elements we include to facilitate our quenching-based readout. To correct for these potential systematic differences, we applied a single offset to the melting points observed in our dataset and used the corrected results for subsequent analysis (Methods; on average, a correction of 9.35°C in Tm or approximately 0.7 kcal/mol in ΔG_{37} was applied). Compared to standard NN model predictions from NUPACK software, the measured values are moderately better than the NN model for ΔG and Tm, but are considerably better for dH (**Fig. 2d, Fig. S2b**). This shows the ability of the Array Melt method to accurately measure folding energies of DNA and generalize to in-solution systems.

FIGURE 2

a Example UV melting curve

b Mean absolute error

before correction
after correction

c

$\Delta G°_{37}$ R = 0.849

$\Delta H°$ (no correction)   R = 0.712

T_m R = 0.827

d

array measurement vs. UV melting
NN modelt prediction vs. UV melting

$\Delta G°_{37}$ $\Delta H°$ T_m

-0.001

Figure 2. Cloud lab UV melting validation of the method

a. Example UV melting curve. Pink dots, measured data points of cooling curve; green solid line, fitted line; green dotted line, fitted melting temperature.

b. Mean absolute error of ΔG, dH and Tm between Array Melt data and UV melting data before and after systematic error correction. Gray bars, errors prior to correction; colored bars, after applying correction with a single parameter offset.

c. Direct comparison of measured ΔG, dH and Tm between Array Melt and UV melting data. Each datapoint is one hairpin variant. Gray dots, measurements prior to correction; colored dots, measurements after applying correction with a single parameter offset.

d. Pearson's correlation R between UV melting data and either Array Melt data or NN model prediction. Colored bars, array measurements vs. UV melting; black bars, NN model predictions vs. UV melting.

SUPP FIGURE 2

Figure S2

a. UV melting Tm as a function of DNA hairpin oligo concentration. Oligo used is WC68.

b. Direct comparison of measured ΔG, dH and Tm between NN model prediction and UV melting data. Each datapoint is one hairpin variant.

Quantifying Nearest Neighbor model performance on non-WC hairpins

After validation of our experimental system, we aimed to examine the performance of the standard Nearest Neighbor (NN) model on hairpins whose stems are not perfect Watson-Crick pairs. When predicting folding free energies associated with different single mismatch types (**Fig. 3a**), the NN model achieved an R^2 of 0.61 on the average free energy for each of the types (implying it can explain 61% of the observed experimental variance), or $R^2 = 0.31$ on all individual variants. However, this model was unable to consistently predict significant variances in folding energies that arise from the same mismatch type with differing flanking sequences ($R^2 = -0.61$ for T>C mismatches, as shown on the right side panel, $R^2 = 0.09$ for A>G, $R^2 = -0.48$ for C>T, and $R^2 = -1.66$ for G>A) (**Fig. 3b, Fig. S3a**). Indeed, for three out of the four mismatches we analyzed, the NN model introduced extra variance (i.e. negative R^2 value) compared to the class-average prediction, implying that the variance in the NN model's prediction residual was greater than the actual variance present in the data (**Fig. 3c, Fig. S3b**). While the NN model succeeded in capturing variance for some variant groups with the same mismatch type and nearest neighbor base pairs, it is unable to capture others (**Fig. S3c**). This led us to conclude that there are significant variances in folding energy that are not accounted for by the NN model.

Influence of sequences on energy across different secondary structures

We hypothesized that some of the observed variance in melting behavior not explained by the nearest-neighbor model may be due to the influence of sequences adjacent to mismatches. To assess this possibility, we calculated the impact of each nucleotide position on folding energy. We first determined a nucleotide importance score at each nucleotide position in different sequence groups with the same target secondary structures by calculating the variance in either ΔG_{37} or $\Delta\Delta G_{37}$ values for the four different identities of the nucleotide at the target position (see Methods). While it is possible that variants might adopt alternative secondary structures yet maintain two-state behavior, our analysis still provides valuable insights into how sequences affect folding energy. For single mismatches, we used $\Delta\Delta G_{37}$ values relative to Watson-Crick pairing "parent" variants to regress out the energy expected from Watson-Crick stacks; for other secondary structures, we directly used ΔG_{37} values. As anticipated, nucleotides along the hairpin stem displayed consistent nucleotide importance scores for Watson-Crick pairing variants, and the two mismatching nucleotides have the highest contribution to this folding energy for single mismatch variants, both in the Array Melt data and the NN model (**Fig. 3d, Fig. S3d**). Extending our analysis to hairpin loops and bulges, we observed that specific sequences within these structures critically influence their thermodynamic energy (**Fig. 3e-f**). In triloops and tetraloops, loop center nucleotides had a more pronounced influence on folding energy in the Array Melt data than NN model predictions. We also noted a slight asymmetry between the 5' and 3' ends of the loop, regardless of loop size (**Fig. 3e, Fig. S3e**). In bulges, flanking Watson-Crick

pairs had greater influence than the bulge sequence itself, with double bulges showing more sequence dependence than singles (**Fig. 3f, Fig. S3f**).

These observations prompted us to examine the energetic interactions between bases by calculating the additional proportion of variance explained in a linear model that considered base interactions (i.e. allowing free parameters for pairwise combinations of bases in the two probed positions) compared to one that treats each position independently (i.e. only allowing free parameters for each individual position). Notably, nucleotides opposite the mismatch showed interactions unaccounted for in the NN model (**Fig. 3g, Fig. S3g**). This interaction score decays with distance along the stem (**Fig. S3h**). Our quantification showed that individual nucleotides, nearest neighbors, and longer-range interactions contribute 74%, 14%, and 12% respectively in this specific case of single mismatches (**Fig. 3h**). In hairpin loops, the interactions between the loop ends and the loop center nucleotides were stronger than expected in tetraloops, but not in triloops (**Fig. 3i, Fig. S3i-j**).

Subsequently, we fit linear regression models to Watson-Crick and single-mismatch variant Array Melt data, varying the length of stack features from one (base pairs only), to two (nearest neighbors) and three (next nearest neighbors). Although prediction errors were similar for nearest- and next-nearest-neighbor models in Watson-Crick variants,

next-nearest-neighbor models outperformed the nearest-neighbor ones in single-mismatch variants (**Fig. 4a**).

FIGURE 3

Figure 3. Decomposition of sources of variances in measured energies

a. Schematics of mismatch types and ΔΔG calculation for single mismatches.
b. Comparison of single mismatch ΔΔG between NN model prediction and Array Melt measurement. Each variant is colored by the identity of the mismatch (left) and the T>C mismatches are zoomed in on the right.
c. Unexplained expected biological variance for each group of mismatches and combined. Variance explained or introduced by the nearest neighbor (NN) model are shown in green and red respectively.
d. Nucleotide importance scores for each individual nucleotide in mismatch constructs, grouped by secondary structure. Importance score is calculated as the sum of squares (ss) introduced by varying the given nucleotide position.
e. Nucleotide importance scores for tetraloops and triloops.
f. Nucleotide importance scores for bulges.
g. Pairwise interaction scores for each pair of nucleotide locations for Watson-Crick and single mismatch variants. Yellow star is the anchor point in each subplot, and the color of each position indicates the strength of secondary interaction of that position with the anchor. Interaction score is calculated as R squared of a linear model given all combinations of the pair as features, subtracted by that of the two individual nucleotides.
h. Percentage of contribution of single nucleotides, nearest neighbors, and longer range interactions to total variance in $\Delta\Delta G_{37}$ for single mismatches.
i. Pairwise interaction scores for tetraloops and triloops.

SUPP FIGURE 3

SUPP FIGURE 3

i

pairwise interaction score *1e2 of (Array Melt data - NN model)

j

Array Melt data

NUPACK

Figure S3

a. R^2 values between Array Melt data and NN model predictions for all single mismatches, and for each mismatch type.
b. Relationship of biological variance to total variance in Array Melt data. Each dot is a group of variants with identical mismatch type and adjacent nearest neighbor base pairs as in **Fig. 3a**, colored by mismatched type. The biological/total variance ratio is shown on the right.
c. Examples of when the NN model succeeds or fails to capture total variance in data. Each subplot is calculated from a group of variants with identical mismatch type and adjacent base pairs. Green is the histogram of Array Melt measurements, and purple is what is expected from the NN model prediction with measurement uncertainty.
d. Comparison of nucleotide importance scores calculated from Array Melt data and NN model predictions for Watson-Crick and single mismatch variants.
e. Comparison of nucleotide importance scores calculated from either $\Delta\Delta G_{37}$ or ΔG_{37}, and on Array Melt data or NN model predictions, for tetraloops and triloops.
f. Comparison of nucleotide importance scores calculated from Array Melt data and NN model predictions of bulge variants.
g. Pairwise interaction scores calculated from NN model predictions for Watson-Crick and single mismatch variants.
h. Pairwise interaction scores as a function of distance between the two nucleotides in the pair, for Watson-Crick variants or single mismatch variants. Those for the latter are splitted by the distance from the anchor point to the single mismatch.
i. Difference between pairwise interaction scores from Array Melt data and NN model prediction.
j. Comparison of nucleotide importance scores calculated from Array Melt data and NN model predictions of triloop variants.

Revised parameter set for NUPACK integration

We next aimed to develop a refined set of parameters to integrate our findings with the NUPACK software[7,19], thereby enabling more accurate DNA thermodynamic predictions. We fit a compact linear regression model with only 178 parameters to Array Melt data, focusing on structural elements prevalent in our designed library. This revised parameter file was then incorporated into NUPACK. When we allowed Watson-Crick stack parameters to vary freely, we reproduced the established nearest-neighbor parameters set (**Fig. 4b**). We then fixed the values of Watson-Crick stack parameters and fitted the NUPACK-compatible NN model (**Fig. S4a**, parameter file available in supplementary material). When using NUPACK to predict free energy, we assumed a two-state model, since the ensemble model produced inferior results, either with the original *dna04* or newly-fitted parameter sets (**Fig. S4b**). Taking single mismatch parameters as an example, we note that the new and the original *dna04* parameters have similar overall trends, while our new parameters capture context-dependent variations of GT mismatches, as well as the instability of CC mismatches, which has been shown to be the most thermodynamically unstable DNA single mismatch[20] (**Fig. 4c, Fig. S4c**). Next, we splitted Array Melt data into training, validation, and test sets (24979, 1315, and 1438 variants respectively, stratified by variant type). We first tested our new NUPACK parameter set on held-out Array Melt test data and observed a low adjusted mean absolute error (MAE) of 0.56 kcal/mol for ΔG_{37} (**Fig. 4d-e, Methods**). This new parameter set also improved prediction on randomly designed hairpins measured with UV melting (**Fig. S4d**). We then asked whether the new parameter set, trained on DNA hairpin data, extends to

DNA duplexes. When predicting unseen data with NUPACK, either using the original *dna04* or the new parameter set, we observed a similar performance of Watson-Crick strands but improved performance on more complex variants. On a compiled UV melting data set consisting of 384 Watson-Crick pairing DNA duplexes[21], our parameter set performed similarly to *dna04* (**Fig. S4e**). In contrast, on a dataset from *Oliveira et al.* of DNA duplex melting temperatures[14] containing either no mismatch, single, double or triple mismatches, both parameter sets have a small MAE of around 0.6 °C for Watson-Crick duplexes; while on single mismatches, the new parameter set enables a much lower prediction error of 1.15 °C ($R^2 = 0.68$) compared to 1.66 °C ($R^2 = 0.25$) of *dna04*. The improvement is even greater on double and triple mismatches (**Fig. 4f-g**). We conclude that this updated parameter set, while simple, improves the prediction quality of NUPACK on DNA secondary structures, especially mismatches and hairpins.

Enhanced linear model with rich parameterization

Building on this NUPACK-compatible model, we next developed an enhanced linear model that incorporates a richer parameter set ("richer parameter model"), allowing still more accurate and comprehensive prediction of DNA thermodynamics on unseen validation data. Because this model has more parameters than are implemented in the NUPACK, it is not immediately compatible with the software. We fit this model with 1338 free parameters, incorporating still more context information (including interactions between the two stacks flanking a mismatch, bulge sequence identity, and full hairpin loop sequence for triloops and tetraloops, see **Methods**), to the Array Melt training data. This rich parameter NN model yields an even better performance both on the Array Melt test data ($MAE = 0.34 \, kcal/mol$, $R^2 = 0.81$) and the external duplex datasets (**Fig. 4d-g, Fig. S4e**). Notably, when comparing the prediction error on Array Melt validation dataset as we titrate the percentage of training data used to fit the model, the 178 parameter model plateaus with just a small fraction of training data used, while the 1338 parameter model requires more measurements to plateau, but still does so after being fit on less than half of the training data set (**Fig. 4h**). Given the relatively overdetermined nature of even this rich parameter NN model, this behavior is expected, and clearly demonstrates that increasing data size allows for more variance to be captured by a more expressive model.

FIGURE 4

a Error on Array Melt test set

ΔG°₃₇ / dH

MAE (kcal/mol) — length of stack features
Watson Crick / single mismatch

b NN parameters — Fitted / dna04

Nearest neighbors: AA/TT, AT/AT, TA/TA, CA/TG, GT/AC, CT/AG, GA/TC, CG/CG, GC/GC, GG/CC
ΔG°₃₇ (kcal/mol)

c Single mismatch NUPACK parameters — New / dna04

context vs mismatch
ΔG° (kcal/mol)

d NN models on Array Melt test set

all / Watson Crick / mismatch / bulge / tetraloop / triloop

all: MAE = 0.87 kcal/mol, R = 0.65, R² = -0.18
Watson Crick: MAE = 0.50, R = 0.86, R² = 0.66
mismatch: MAE = 1.09, R = 0.52, R² = -0.87
bulge: MAE = 0.75, R = 0.48, R² = -1.05
tetraloop: MAE = 0.54, R = 0.68, R² = 0.35
triloop: MAE = 0.48, R = 0.58, R² = 0.02

MAE = 0.56 kcal/mol, R = 0.81, R² = 0.49
Watson Crick: MAE = 0.43, R = 0.90, R² = 0.77
mismatch: MAE = 0.58, R = 0.84, R² = 0.38
bulge: MAE = 0.65, R = 0.51, R² = -0.47
tetraloop: MAE = 0.36, R = 0.85, R² = 0.68
triloop: MAE = 0.34, R = 0.77, R² = 0.51

MAE = 0.34 kcal/mol, R = 0.90, R² = 0.81
Watson Crick: MAE = 0.31, R = 0.95, R² = 0.89
mismatch: MAE = 0.41, R = 0.85, R² = 0.72
bulge: MAE = 0.32, R = 0.79, R² = 0.60
tetraloop: MAE = 0.27, R = 0.92, R² = 0.85
triloop: MAE = 0.23, R = 0.88, R² = 0.77

predicted ΔG°₃₇ (kcal/mol) / measured ΔG°₃₇ (kcal/mol)

e
MAE of ΔG°₃₇ (kcal/mol) / R² of ΔG°₃₇
NN model

original NN model
NUPACK-compatible NN model (178 features)
richer parameter NN model (1338 features)

f On an orthogonal duplex dataset
R² of Tm — number of mismatch

g
no mismatch: MAE = 0.60, R = 0.96, R² = 0.87
single mismatch: MAE = 1.66, R = 0.59, R² = 0.25
double mismatch: MAE = 2.17, R = 0.43, R² = 0.13
triple mismatch: MAE = 1.91, R = 0.16, R² = 0.03

no mismatch: MAE = 0.63, R = 0.96, R² = 0.86
single mismatch: MAE = 1.15, R = 0.85, R² = 0.68
double mismatch: MAE = 2.02, R = 0.71, R² = 0.26
triple mismatch: MAE = 1.67, R = 0.49, R² = 0.24

no mismatch: MAE = 0.47, R = 0.96, R² = 0.92
single mismatch: MAE = 0.95, R = 0.87, R² = 0.75
double mismatch: MAE = 2.09, R = 0.74, R² = 0.21
triple mismatch: MAE = 1.60, R = 0.53, R² = 0.27

predicted Tm (°C) / Tm (°C)

h Error on Array Test set
sequence variants — 12,490 / 24,979
MAE of ΔG°₃₇ (kcal/mol) — % training data used

NUPACK-compatible NN model (178 features)
richer parameter NN model (1338 features)

Figure 4. Improved linear regression models for DNA folding energetics

    a.   Mean absolute errors of linear regression models, with either single base pair, nearest neighbor, or next nearest neighbor stack features, on Array Melt data.

    b.   Comparison of nearest-neighbor parameters of Watson-Crick pairs fitted from array melt data and from literature[8].

    c.   Visualization of new and *dna04* free energy parameters for single mismatches. "Context" and "mismatch" are defined as in **Figure 3a**.

  d-e. Performance of the original NN model (*dna04*), NUPACK-compatible NN model, and richer parameter NN model on held-out Array Melt data.

    f-g. Performance of the original NN model (*dna04*), NUPACK-compatible NN model, and richer parameter NN model on an orthogonal dataset of DNA duplexes containing varying numbers of mismatches.

    h. Mean absolute error on Array Melt data as a function of percentage of training data used to fit the linear regression model.

SUPP FIGURE 4

Figure S4

    a.   Schematics of the NUPACK-compatible NN model.

Graph neural network (GNN) model for DNA energy prediction

Because our rich linear model likely had insufficient power to capture the biological variance in our data, we developed a graph neural network model[22,23] specifically designed for predicting DNA energy. In this model, a DNA molecule is represented as a graph where nodes represent nucleotides and edges represent chemical bonds. Nucleotides (A, T, C, or G) and chemical bonds (either 5' to 3' backbone, 3' to 5' backbone, or hydrogen bond) are one-hot encoded as node and edge features, respectively. Taking these graph representations as inputs, the GNN iteratively refines node embeddings across graph convolutional layers. Essentially, nodes exchange information through different edge types, allowing, over multiple layers, for the embedding of one node to encapsulate information from the entire graph. Investigation of various types of graph convolutional layers revealed that the graph transformer layers[24] outperformed other architectures, such as graph attention layers[25] (**Fig. S5a**). After graph convolution, each node possesses a learned embedded feature, reflecting not only the original node features, but also the edges and overarching graph structures. To embed this graph into a fixed-length vector representation, we employed a global pooling layer employing the Set2Set algorithm[26], which used a Long Short-Term Memory (LSTM) processing block. This block systematically ingests node embeddings while updating its internal state, resulting in a vector representation of the entire graph that is order-invariant to the input node features. This processed summary vector is then fed into a small, fully connected neural network, which regresses for dH and Tm values (**Fig. 5a**).

We trained this end-to-end model on Array Melt data. A portion of the Array Melt data was held out as validation and test set. We first explored the effect of the number of graph convolutional layers on prediction accuracy (**Fig. 5b, Fig. S5b**). Performance converged at around 4 graph convolutional layers, and did not improve beyond 4 layers, consistent with our observations in **Fig. S3h** that pairwise nucleotide interactions span around a distance of 4 nt. When evaluating model performance over an increasing percentage of training data used as in **Fig. 4h**, we observed a slower plateauing of the curve, indicating that the model can make use of this large-scale data set (**Fig. 5c**). The trained GNN achieved prediction error of 0.18 kcal/mol in dG_{37} ($R = 0.97$, $R^2 = 0.94$), compared to 0.34 kcal/mol of a baseline model that simply looks up the k=8 closest variants in the training set and gives a weighted average of the lookup values as prediction (**Fig. 5d-e, Fig. S5c-d**), and the GNN predictions are consistently high quality for different types of sequence variants (**Fig. S5e**). Interestingly, although the GNN was trained entirely on graph representations of DNA hairpins without knowing the nature of backbones and base pairing, it is able to

extend to the annealing and melting properties of DNA duplexes (**Fig. 5f**). While the GNN performance is slightly worse than our rich parameter NN model (**Fig. 4f-g, Fig. S4e**) on DNA duplexes, this level of performance suggests that the neural network has grasped the "gist" of the physical system of DNA folding thermodynamics, and can extrapolate to non-hairping annealing contexts. Compared to the slow decrease of prediction error on the Array Melt data, prediction error rapidly decreased within around the first 20 epochs and then plateaued for the two DNA duplex datasets (**Fig. S5f**), likely because of the relatively low complexity of those datasets.

To better understand how the GNN makes predictions, we analyzed the intermediate activation from the output of the Set2Set aggregation layer. The GNN projects each DNA molecular graph into a 250-dimensional embedding vector. Many of the top principal components of the embeddings correlate strongly with predicted thermodynamics parameters ($R = 0.8$ between PC1 and dH, $R = -0.77$ between PC5 and Tm, and $R = 0.79$ between PC1 and ΔG_{37}), but none with other qualities of the sequence variants including DNA length or GC content (**Fig. 5g-h**), underscoring the GNN's focus on thermodynamics. When examining embeddings from different datasets (Array Melt, literature UV, and Oliveira et al.), each dataset's distinct design structure mapped to unique areas in a two-dimensional UMAP visualization (**Fig. 5i**). This observation indicates that the embeddings capture more than just predicted thermodynamics like melting temperatures (**Fig. S5g**), and that the learned embedding space encompasses sequence and structure details beyond mere parameter prediction.

Lastly, we compared prediction errors across various models. The graph neural network is the top performer on Array Melt test data, while the 1778-parameter richer parameterization NN model performs best when applied to duplex data. Precision of the GNN is noteworthy, nearing the uncertainty range of bootstrapped experimental measurements (mean of uncertainty is 0.14 kcal/mol in ΔG_{37}), highlighting its effectiveness in capturing complex relationships in DNA sequences and structures (**Fig. 5j**).

FIGURE 5

Figure 5. Graph neural network model of DNA folding thermodynamics

    a.   Schematics of the graph neural network architecture.

b. Mean absolute error of melting temperature on Array Melt data as a function of the number of graph convolution layers in the GNN.

c. Mean absolute error of ΔG_{37} on Array Melt data as a function of percentage of training data used to fit the GNN model.

d. Pearson's R between GNN prediction and held out Array Melt, literature UV melting or Oliveira et al. duplex melting data.

e-f. Comparison between GNN prediction and held out Array Melt, literature UV melting or Oliveira et al. duplex melting data. Each dot is a single sequence variant.

g. Pearson's R between principal components of the learned variant embedding to measured thermodynamic parameters and variant properties.

h. Comparison between the most correlated principal components and dH or Tm.

i. UMAP visualization of learned variant embeddings after aggregation, colored by the dataset (Array Melt, literature UV melting or Oliveira et al. duplex melting data).

j. Benchmark of all models on held out Array Melt data, compared to Array Melt measurement error.

SUPP FIGURE 5

Figure S5

a. Mean absolute error of GNNs with either Graph Attention Network or Graph Transformer Network architecture at the graph convolutional layer.
b. Mean absolute error of enthalpy on Array Melt data as a function of the number of graph convolution layers in the GNN.
c. Mean absolute error of melting temperature on Array Melt data as a function of percentage of training data used to fit the GNN model.
d. Schematic of the baseline closest sequence lookup model.
e. GNN predicted ΔG_{37} vs. held out Array Melt, splitted by type of variant.
f. The objective function of GNN (RMSE of normalized dH and Tm), mean absolute error of Tm, and Pearson's R of Tm across the GNN model training process.
g. UMAP of learned embeddings of test data, colored by their predicted Tm.

Discussion

In this work, we developed an accurate, scalable fluorescence-based method for measuring the melting of DNA hairpins (Array Melt) and used it to generate the largest-to-date DNA folding thermodynamics dataset. The Array Melt method, with its high throughput and precise measurement capabilities, allows for a more detailed exploration of the thermodynamics of diverse DNA structures, including those with mismatches, bulges, and various loop motifs. The scale of this data set revealed the significant role played by sequences adjacent to mismatches and within loops to the thermodynamic stability of hairpins. This observation implies that traditional models that parameterize only the immediate nearest neighbors must be underpowered to capture the variance observed in these molecular variants. To this end, we developed three quantitative folding energy models of increasing complexity, including revised parameterization of the linear regression model compatible with NUPACK, a richer nearest neighbor model especially accurate for duplex annealing of mismatched molecules, and graph neural network model that accells at predicting the melting behaviors of hairpins. These results support the notion that while traditional nearest-neighbor (NN) models are accurate in predicting Watson-Crick stack behaviors, they fall short in accurately modeling the thermodynamics of more complex DNA structures, such as hairpins and interior loops.

Overall, by providing a more accurate model for DNA folding thermodynamics, we pave the way for improved design of biotechnological tools and applications. For example, our models will provide a more accurate starting point for *in silico* design in areas such as qPCR primer design, DNA oligo hybridization probe design, and DNA origami. However, the study's focus on two-state folding presents a notable limitation, although our comparison between the two-state and ensemble methods (**Fig. S4b**) revealed greater accuracy of the former in predicting Array Melt measured thermodynamic parameters. The inferior performance of the ensemble method may be attributed to both the inference methods and the experimental design of the datasets used to derive NN models. Therefore, interpreting NN model predictions and dynamic programming algorithm outputs as "true" energies associated with monolithic secondary structures warrants careful consideration. In this study, the two-state folding energies should be interpreted as an aggregated average across multiple folding states, rather than that of a single folded microstate

within this ensemble. We anticipate that future analysis incorporating experimental design, data preprocessing, and parameter inference[27] techniques that consider ensemble folding could result in further improvements.

Finally, while our measurements were carried out at a single monovalent salt concentration, large-scale measurements exploring the stability of DNA structure to the concentration and type of monovalent or divalent cation would be relatively straightforward. Furthermore, the potential application of our method to RNA structures presents an exciting avenue for future research. Given the diverse and critical biological functions of RNA, which are heavily dependent on its secondary structure, applying our high-throughput and precise thermodynamic measurement technique could unveil new dimensions in understanding RNA biology and facilitate the development of RNA-based therapeutics.

Methods

Library assembly and sequencing

Designed library variants were synthesized into DNA by Twist Biosciences (South San Francisco, CA). Sequence map of the library is provided in supplementary information as *array_melt_library.gb*, where the "filler sequence" is a 5' partial segment of AACAACAACAACATACTAACAACAACATAACAAATCAAAA, its length and the variable region adding up to 40 bp, such that all synthesized fragments have the same length despite different variable region length. Reverse complement of the sequence map was ordered as a synthesized oligo pool, amplified using internal primers to enrich for full-length library variants. The PCR reaction consisted of 1/100x dilution of the synthesized oligo pool (final concentration 0.01 nM), 200 nM of each primer (T7A1library, D-TruSeqR2 Table S1), 1x Phire Hot Start II PCR Master Mix (ThermoFisher Scientific F125L). The reaction proceeded for 9 cycles of 98˚C for 10 seconds, 56˚C for 30 seconds, and 72˚C for 30 seconds. Reaction mixtures were purified using QIAquick PCR Purification Kit (Qiagen 28104) to remove primers and proteins, then eluted into 20 uL dilution buffer.

After initial amplification, the library was amplified with primers to bring in sequences compatible with Illumina sequencing. This five-piece assembly PCR included two outside primers and two adapter sequences. The PCR reaction consisted of 1 μl of the previous reaction, 137 nM of outside primers (*short_C* and *short_D*; Table S1), 3.84 nM of the adapter sequences (*C-i7pr-bc-T7A1* and *D_TruSeqR2*; Table S1), 1x Phire Hot Start II PCR Master Mix (ThermoFisher Scientific F125L). The reaction proceeded for 14 cycles of 98˚C for 10 seconds, 56˚C for 30 seconds, and 72˚C for 30 seconds. Reactions were purified using the QIAquick PCR Purification Kit and quantified with a Qubit Fluorometer (ThermoFisher Scientific).

Imaging station setup

An imaging station was used to image the Miseq chip at increasing temperatures. This station was built from a combination of custom-designed parts from a disassembled Illumina genome analyzer IIx[17,28]. Two channels were employed: the "red" channel used the 660 nm laser and 664 nm long pass filter (Semrock), and the "green" channel

used the 50 nm laser and 590 nm band pass filter (Semrock). All images were taken with 600 ms exposure times at 150 mW fiber input laser power. Focusing at each temperature was achieved by sequentially adjusting the z-position and re-imaging the four corners of the flow cell; the adjusted z-$_{positions}$ were then fit to a plane.

Post-sequencing, the chip was washed with Cleavage buffer (100 mM Tris-HCl, 125 mM NaCl, 0.05% Tween20, 100 mM TCEP, pH 7.4) to remove residual fluorescence from the reversible terminators used in the sequencing reaction at 60°C for 5 minutes. Any strands of DNA not covalently attached to the surface of the chip was removed by washing in 100% formamide at 55°C. The resulting single-stranded DNA fragments were incubated with 500 nM of the oligo *Biotin_D_Read2* and *red_oligo* (Table S1) in Hybridization buffer (5x SSC buffer (ThermoFisher 15557036), 5 mM EDTA, 0.05% Tween20) for 15 minutes at 60 °C, subsequently the temperature was lowered to 40 °C for another 10 minutes.

Measuring melt curves on the chip

The Cy3 labeled *fluor_oligo* was annealed at 40°C for 5 minutes, then the quench-labeled oligo *quench_oligo* (**Table S1**)using the same protocol. The chip was imaged after each step to ensure that hybridization had occurred through increase or quenching of signal in the corresponding channel. The chip was then rinsed with melt buffer (50 mM Na-HEPES pH 8.0, 25 mM NaCl). To quantify the melt curves, the image station temperature was then lowered to 20°C. For each temperature point, the system was allowed to equilibrate to the new temperature (approximately 5 minutes) before refocusing and imaging. The temperature was raised in 2.5°C increments to a maximum temperature of 60°C.

Processing sequencing data

Sequencing data from Illumina Miseq was processed to extract tile and coordinates of each sequenced cluster. Forward and reverse paired-end reads were aligned using FLASH[29] with default settings. Consensus sequences from FLASH were aligned to the reverse complement sequences of *fluor_oligo* and *quench_oligo* using a Needleman-Wunsch alignment (nwalign3). For consensus sequences that successfully aligned to both with p-value <1e-3, evaluated as described in[30], the variable region was extracted as the region between the two flanking regions and aligned to sequences in the reference library. A reference sequence was assigned based on the best-scoring alignment with a p-value < 1e-6. The resulting clusters were used for fluorescence quantification in the downstream data analysis.

Fluorescence data processing and image fitting

Fluorescent melt curve images were mapped to sequencing data from the Illumina Miseq. First, sequencing data was processed to extract the tile and coordinates of each sequenced cluster. The coordinates were used to generate synthetic images of clusters, which were registered to the fluorescent images iteratively to map the coordinates to the images at sub-pixel resolution[31]. Once the identities of the clusters were determined, each cluster was fit to a 2D normal distribution to quantify its fluorescence[28].

Distance sensitivity curve

The repeat controls used to plot the distance sensitivity curve in Fig. 1d are repeats of varied lengths of the following sequences: poly-A, T, AT, AAC, AAG, AAT, AC, ACC, AG, AGG, TC, TG, TTA, TTC, and TTG.

The theoretical curve fitted to the distance-to-fluorescence curve in Fig. 1d is $f(x) = \frac{1}{1+(a \cdot (x+b)^{-3})}$, where x is the length in nt, $f(x)$ is the dimensionless fluorescence at length x, normalized to maximum fluorescence, b is a fitted parameter in nt, which accounts for systemic distance offset, and a is another fitted parameter:

$$a = \frac{2\pi}{3}^{2/3} (K_{D,BHQ} \cdot N_A \cdot k^3)^{-1} \qquad (1)$$

where $K_{D,BHQ}$ is the dissociation constant for the fluor-quench pair, N_A is Avogadro's number, and $k = 0.64 \times 10^{-9} m \cdot nt^{-1}$ is a distance conversion factor from nt to physical distance for ssDNA. The fitted parameter values are $a = 71.19$, $b = 0.08$, and $K_{D,BHQ} = 145.6 \, mol \cdot L^{-1}$. Using $\Delta G = RT \cdot \ln K_{D,BHQ}$, we have $dG_{37} = 3 \, kcal/mol$.

Fluorescence normalization

Size normalization

To reduce inter-cluster variation in fluorescence measurements, we normalized the Cy3 fluorescence during the melt curve measurements by the total amount of ssDNA in that cluster, measured as Cy3 fluorescence right after hybridization of *fluor_oligo* but before hybridization of *quench_oligo*. The normalizing factor Cy3 signal was clipped to the 1st and 99th percentile of the total distribution.

Temperature dependence and photobleaching normalization

Across the timespan of the experiment, fluorescent signals are dependent on temperature and undergo photobleaching. To correct for these factors, we further normalized the size-normalized signals of clusters to median signals of control variants designed to have constant distances between the Cy3 fluorophore and the quench, including 1) "super stable stem" variants with long GC stems that are not supposed to melt within the experimental temperature range and show minimum fluorescence signals, and 2) "long repeat" controls that are at least 39 nt long, not supposed to form a hairpin, and show maximum fluorescence signals. For both control variant groups, only the variants whose median signal across clusters fall within the 5th and 95th percentile of all median signals at each temperature point within that control variant group are used. Each variant is allowed to fall outside the percentile range at 1 temperature point at most. This step filters out the variants not exhibiting the expected behavior. Afterwards, the median values at each temperature point of both groups are taken and used to normalize the entire dataset. **(Fig. S1d)**

Direct melt curve fitting

The melt curve fitting pipeline was adapted from[31]. Briefly, the direct fitting process was carried out in the following steps: 1) Directly fit normalized signals of single clusters; 2) Estimate the distribution of f_{max} and f_{min} from single cluster fit results; 3) Refine the fit at the variant level with bootstrapping. The details are described below. The fitting process is available as a snakemake[32] pipeline at https://github.com/keyuxi/array_analysis.

1) Single cluster fitting

Equation (2) was directly fitted to normalized signals of single clusters with minimal constraints, where T and T_m are temperatures in Kelvin, and $k_B = 0.0019872$ is the Boltzmann constant. For initial values, Tm was set to the temperature point where the normalized signal is closest to 0.5, dH to -40, f_{max} to 1, and f_{min} to 0. The python package *lmfit* (v1.0.3)[33] was used for least-squares fitting.

$$f[T] = f_{min} + \frac{f_{max} - f_{min}}{1 + exp(\frac{\Delta H}{k_B} \cdot (\frac{1}{T_m} - \frac{1}{T}))} \qquad (2)$$

2) Estimating distributions of f_{max} and f_{min}

We used the following empirically gated criteria to decide if a single cluster fit was successful: R^2 of the fit at least 0.5, fitted f_{min} between -1 and 2, the estimated standard error of the best-fitted parameter value $\sigma_{f_{min}}$ less than $f_{min} + 1$, fitted f_{max} between 0 and 3, $\sigma_{f_{max}}$ less than f_{max}, σ_{Tm} less than 10 K, and $\sigma_{\Delta H}$ less than 100 kcal/mol. A p value is calculated for each variant with a null hypothesis of 25% successful single cluster fits as in[31]. The single clusters are then aggregated to the variant level. Median values of the parameters of each variant were calculated. To select for variants used for estimation of f_{max}, the previously mentioned single cluster criteria were combined with the following extra criteria: 1) be a good fitter, defined as having a success p value less than 0.01; 2) be unfolded at high temperatures, defined as having a fitted f_{max} value of at least 0.5, and $p_{unfolded}^{60°C}$, the portion of unfolded molecules in the cluster calculated from the fitted parameters, of at least 0.975. Similarly, for f_{min} we select for variants with a p value less than 0.01, f_{min} less than 0.05, and $p_{unfolded}^{20°C}$ at most 0.025. We then fitted cluster-per-variant-dependent Gamma distributions to f_{max} and f_{min} as in[31].

3) Fit refinement

For each variant, we separately decided whether to enforce f_{max} and f_{min} distributions to correct for variants that never become fully unfolded or folded during the experiment. Specifically, we calculated the lower bound for f_{max}

as $\mu_{f_{max}} - 10 \cdot \sigma_{f_{max}}$, where $\mu_{f_{max}}$ and $\sigma_{f_{max}}$ are the mean and standard deviation calculated from the Gamma distribution with the according number of clusters per variant from the previous step, and the upper bound for f_{min} as $\mu_{f_{min}} + \sigma_{f_{min}}$. Here, we set the margin for f_{max} larger to enforce global f_{max} less, as f_{max} is more variable between variants than f_{min} because of sequence-dependent effects. If the median signal at the highest temperature point is less than the lower bound for f_{max}, we enforce f_{max}; likewise, if the median signal at the lowest temperature point is greater than the upper bound for f_{min}, we enforce f_{min}.

We fitted each variant by bootstrapping the clusters within that variant. At each resampling step, we draw a f_{max} or f_{min} value from the global distribution if enforced or keep f_{max} or f_{min} floating otherwise, and fit equation (2) to the median signal of the drawn clusters, using the single cluster fit results as the initial parameters. Then, we calculate ΔG_{37} and ΔS for each resampling step:

$$\Delta G[T] = \Delta H - T \cdot \Delta S = \Delta H \cdot \left(1 - \frac{T}{T_m}\right) \qquad (3)$$

$$\Delta S = \frac{\Delta H}{Tm} \qquad (4)$$

where T is temperature in Kelvin. We repeated the process 100 times and obtained a 95% confidence interval and standard deviation for each parameter, ΔH, Tm, ΔG_{37} and ΔS.

ΔG Line fitting

We also fitted the data with an alternative line fitting to obtain heuristics for two-state behavior. We first re-normalize the signal $f[T]$, previously normalized to initial fluorescence after hybridization as aforementioned, with fitted f_{max} and f_{min}. We clip $f_{norm}[T] = \left(f[T] - f_{min}\right)/\left(f_{max} - f_{min}\right)$ between 0 and 1 with an $\epsilon = 0.01$ to avoid problems in the next step. We transformed $f_{norm}[T]$ to $\Delta G[T]$ with

$$\Delta G[T] = - k_B T ln\left(\frac{1}{f_{norm}[T]} - 1\right) \qquad (5)$$

Error propagation was calculated with the uncertainties python library. Finally, we fitted with a line with free parameters ΔH and Tm:

$$\Delta G[T] = \Delta H \cdot \left(1 - \frac{T}{T_m}\right) \qquad (6)$$

If the variant exhibits two-state behavior, $\Delta G[T]$ should fit well to a linear line. We first fit with RANSAC[34] to identify outliers, then re-fit the inliers with ordinary least squares linear regression for the final parameters.

Filtering and combining data replicates

Each variant measured and fitted to curves in a replicate experiment was first filtered by the criteria: at least measured in 5 clusters, bootstrapped error in ΔG_{37} is less than 2 kcal/mol, error in Tm less than 25°C, error in ΔH less than 25 kcal/mol, and curve fitting RMSE less than 0.5. They were then compared to the line fitting results. To be labeled as passing two-state criteria, the fitted dH value of a variant between the 2 methods need to agree within 20%, Tm within 2°C, the reduced $\chi 2$ of the line fit below 1.5, and the number of inlier data points in line fitting greater than 10. The replicates were then combined as described in[31]. Briefly, for each parameter (ΔH, Tm, ΔG_{37}, or

ΔS), its measurements X_i and bootstrapped error σ_i in each replicate, the combined error is $\sigma = \sqrt{\left(\sum_i \sigma_i^{-2} \right)^{-1}}$ and

the combined parameter $X = \sigma^2 \cdot \sum_i X_i / \sigma_i^2$, where the sum is over the replicates whose data are not missing for the given variant. The missing values were ignored. We then apply a second filter for two-state behavior. A variant needs to be labeled as passing two-state criteria in at least 2 replicates to be kept. We combined 4 such replicate experiments, including 3 melting and 1 cooling curve measurements from 2 different chips (see table for detail). We apply a final filter on the combined variants to only keep those whose Tm are between 25 and 55°C and fall within the dynamic range of the system.

UV melting

Designing sequences for UV melting and quality control

We randomly selected hairpin sequences from each category of the Array Melt library for validations in **Fig. 2**, and designed random 16-mers that would fold into a hairpin with the secondary structure ((((((....)))))) for **Fig. S4d**, filtering those with misfold secondary structures with NUPACK. DNA oligo samples are run through Ion Exchange High Pressure Liquid Chromatography (IE-HPLC) remotely on Emerald Cloud Lab (ECL) with DNAPac PA200, 9x250 mm Semi-Prep column (Thermo Fisher Scientific 063421), and the fractions were measured by a UV-Vis detector at 260 nm. Only the oligos with a single peak were used for data analysis.

Running cloud lab UV melting experiments

UV melting experiments were run remotely on ECL. The Manifold data objects could be found at the data object links in supplementary table 2. DNA oligos ordered from IDT were shipped to ECL and resuspended in high concentration stock solutions. For hairpins, the DNA stock solutions are first snap freezed right before each experiment, being placed at 90°C for 5 min and moved to 4°C for 30 min before diluted into working concentrations; for duplexes, the DNA stock solutions are first diluted into working concentrations, then placed on 90°C for 5min and left annealing for 30 min to 1 hour. Melt curves are measured at varied concentrations, where the range is 3 to 12 uM for hairpins and 72 to 96 uM for duplexes. Hairpin data used for method validation in **Fig. 2** and model comparison in **Fig. S4d** were all measured at 6 uM or 9 uM. Samples were measured on a Cary 3500 UV/Vis

spectrophotometer, using micro scale black walled UV quartz cuvettes with a pathlength of 10 mm. Samples were measured for 2 to 3 melt-cooling cycles with a temperature ramp rate of 0.8°C and 15 min equilibrium time between each melt and cool sweep. Temperatures were measured with an immersion probe. Data points were acquired every 1 °C at 260 nm from 5°C to 85 or 95 °C. All ECL objects, including samples and experiments, are available at the Airtable base https://airtable.com/apph4txt3ZhtDRk6l/shrTtTRL25OstczO7.

Analyzing UV melting curves

The UV melting curves were fitted to a modified version of equation (2) with an extra term to account for baseline slope:

$$f[T] = f_{min} + a \cdot T + \frac{f_{max} - f_{min}}{1 + exp(\frac{\Delta H}{k_B} \cdot (\frac{1}{T_m} - \frac{1}{T}))} \qquad (7)$$

where f[T] is the absorbance, a is a free parameter for the slope, and the rest the same as in equation (2). For each individual melt or cooling curve, we first found the temperature where the derivative $\frac{df[T]}{T}$ is maximum, that is, $T_m^{diff} = argmax_T \frac{df[T]}{T}$, as the initial value of T_m. We fixed the value of T_m when fitting equation (7) in the first round and let the parameters a, f_{max}, f_{min}, and ΔH float. Afterwards, we use the best-fitted values from the previous round as the initial values of the parameters and allow T_m to float in the second round of fitting. This two-step fitting process empirically improved the fitting RMSE. We selected the curves with an RMSE less than 0.015, $\sigma_{\Delta H} < 10$ And $\sigma_{T_m} < 5$ and aggregated the fitted dH and Tm values of the melt and cooling curves by each sample tube with

$X = \sigma^2 \cdot \sum_i X_i/\sigma_i^2$ and $\sigma = \sqrt{\left(\sum_i \sigma_i^{-2}\right)^{-1}}$ as for the array experiments. We then calculated ΔG_{37} and ΔS from the aggregated ΔH and T_m using equations (3) and (4), and used the uncertainty propagation equations

$\sigma_{\Delta S} = -\Delta S \cdot \sqrt{\left(\frac{\sigma_{\Delta H}}{\Delta H}\right)^2 + \left(\frac{\sigma_{T_m}}{T_m}\right)^2}$ and $\sigma_{\Delta G_T} = \sqrt{\sigma_{\Delta H}^2 + \left(T \cdot \sigma_{\Delta S}\right)^2}$ to calculate the measurement uncertainties.

Salt correction

All salt corrections used the equation from[35]:

$$\frac{1}{T_m(2)} = (\frac{1}{T_m(1)} + (4.29 \cdot f(GC) - 3.95) \times 10^{-5} \cdot ln\frac{[Na^+]_2}{[Na^+]_1} + 9.4 \times 10^{-6}(ln^2[Na^+]_2 - ln^2[Na^+]_1) \qquad (8)$$

where $T_m(1)$ and $T_m(2)$ are the melting temperatures before and after salt correction in Kelvin, $[Na^+]_1$ and $[Na^+]_2$ the corresponding sodium concentrations, and $f(GC)$ the fraction of GC in the sequence.

Variance analysis

Mismatch $\Delta\Delta G_{37}$ calculation

For each mismatch, we find its parent sequence variant, which is a hairpin with Watson-Crick pairing stem and can derive into the mismatch by mutating a single nucleotide. The mismatches are categorized by the identities of those nucleotide substitutions. We then calculated $\Delta\Delta G_{37}$ of mismatches by subtracting ΔG_{37} of the parent variant from that of the mismatch variant. C>G single mismatches were excluded from the variance calculations because of insufficient number of data points (only 2 variants).

Expected biological variance and model explained or introduced variance

For a dataset of n sequence variants with n corresponding $d\Delta$ values y_1, ..., y_n, measurement uncertainty values σ_1, ..., σ_n, and NN model prediction values $\hat{y}_1$, ..., $\hat{y}_n$, we have total variance $\sigma^2_{tot} = \sum_{i=1}^{n} (y_i - \bar{y})^2/n$ and technical variance $\sigma^2_{tech} = \sum_{i=1}^{n} \sigma_i^2/n$. The expected biological variance $\sigma^2_{bio} = \sigma^2_{tot} - \sigma^2_{tech}$. Model variance is the variance of prediction values $\sigma^2_{model} = \sum_{i=1}^{n} (\hat{y}_i - \bar{\hat{y}})^2/n$, and residual is $\sigma^2_{res} = \sum_{i=1}^{n} (y_i - \hat{y}_i)^2/n$. From these we calculate variance explained $\sigma^2_{explained} = \sigma^2_{tot} - \sigma^2_{res}$. In the case where $\sigma^2_{res} > \sigma^2_{tot}$, we have variance added by model $\sigma^2_{added} = -\sigma^2_{explained} = \sigma^2_{res} - \sigma^2_{tot}$.

Nucleotide importance score

We calculated nucleotide importance score (NIS) using $\Delta\Delta G_{37}$ for mismatches, and ΔG_{37} for triloop, tetraloop, and bulges. For a dataset of n sequence variants with n corresponding energy values y_1, ..., y_n, either ΔG_{37} or $\Delta\Delta G_{37}$, we group the n data points into four groups based on the identity of the nucleotide of interest. For each of the four groups, we calculated the mean, $\bar{y}_A$, $\bar{y}_T$, $\bar{y}_C$, and $\bar{y}_G$. The grand mean $\bar{y}_{tot}$ is the mean of y_1, ..., y_n. The NIS is simply logarithm of the between group sum of squares (SSB):

$$NIS = log[(\bar{y}_A - \bar{y}_{tot})^2 + (\bar{y}_T - \bar{y}_{tot})^2 + (\bar{y}_C - \bar{y}_{tot})^2 + (\bar{y}_G - \bar{y}_{tot})^2] \qquad (9)$$

Pairwise interaction score

For a pair of nucleotides of interest, M and N, we compare two linear models that predict the energy value Y, model A with independence assumption and model B without. Model A is in the form of $M + N \sim Y$, and model B $MN \sim Y$. The pairwise interaction score (PIS) is the difference of R^2 between the two linear models

$$PIS = R^2_{MN \sim Y} - R^2_{M+N \sim Y} \qquad (10)$$

Calculations with NUPACK

Melting temperature

Melting temperatures of DNA duplexes were calculated from the fraction of molecules unpaired. We first calculated structure free energies of the paired state at temperature T, $\Delta G_{paired,T}$, then got the equilibrium constant

$$K = \frac{[dsDNA]}{[ssDNA]^2} = exp(-\ \beta \cdot \Delta G_{paired,T}) \qquad (11)$$

where $\beta = 1/RT$ is the thermodynamic beta, $[dsDNA]$ is the concentration of double-strand DNA, and $[ssDNA]$ that of the single-strand DNA. At melting temperature, $[dsDNA] = [ssDNA] = [DNA]/2$ for distinguishable strands and $[dsDNA] = [DNA]/4$ for indistinguishable strands, where $[DNA]$ is the total concentration of DNA strand species. This gives $K = \frac{2}{[DNA]}$ or $K = \frac{1}{[DNA]}$ for distinguishable and indistinguishable strands respectively. Solving this equation with bisection gives us the melting temperature.

For DNA hairpins, melting temperature is independent of DNA strand concentration. The Boltzmann factor becomes:

$$e^{-\beta E} = exp(-\ \beta \cdot \Delta G_{folded}) \qquad (12)$$

T_m is then simply the temperature when $dG_{folded} = 0$. Therefore, we get structure free energies ΔG_1 and ΔG_2 at a lower temperature T_1 and a higher one T_2, then calculate $\Delta S = -\ (\Delta G_2 - \Delta G_1)/(T_2 - T_1)$, $\Delta H = \Delta G_1 + T_1 \cdot \Delta S$, and $T_m = \Delta H/\Delta S$.

$$T_m = \Delta H/(\frac{\Delta H - \Delta G_{37}}{T_{37}} - R \cdot \ln K) \qquad (13)$$

Data splitting for machine learning models

After filtering with the two-state criteria as described in "ΔG line fitting", we splitted the clean Array Melt into train, validation and test sets, stratified by variant type. We first splitted 5% of the data as the test set (1438 variants), then splitted 5% of the remaining data as the validation set (1315 variants), and 95% of the rest as the training set (24979 variants). We splitted the external 384-duplex dataset and Oliveira et al. dataset to a 1:1 split of validation and test data. The cloud lab UV melting data were used for correction between Array Melt and bulk solution, and were not used for model testing to prevent circularity.

Linear models

Weighted linear regression model

We used a singular value decomposition (SVD) solver to solve the linear regression solutions, taking measurement error of each sequence into account. For a training dataset of n sequence variants with n corresponding ΔH or ΔG_{37}

values $y_1, ..., y_n$, and measurement uncertainty values $\vec{\sigma} = [\sigma_1, ..., \sigma_n]$, we have a n by $n_{feature}$ feature count matrix X derived from the sequences. We then did SVD on matrix $A = X/\vec{\sigma}$ and get U, S, V. Any diagonal value in S that is smaller than a relative numeric threshold 10^{-15} was set to zero. Then the estimated parameter values for the features are $V^T \cdot S_{inv} \cdot U^T \cdot \vec{b}$, and the estimated errors for the features are $\sqrt{\sum V^T \cdot S^2_{inv}}$, where $\vec{b} = \vec{y}/\vec{\sigma}$, S_{inv} is S but taking $\frac{1}{s}$ for all diagonal elements, and S^2_{inv} is S but taking $\frac{1}{s^2}$ for all elements.

Regularized linear regression model

We regularized NN parameters to the original ones present in NUPACK by adding a quadratic penalty term to the loss function

$$loss = \overline{(y - \hat{y})^2} + \lambda \cdot \overline{(v - v_{prior})^2} \quad (14)$$

where v is the coefficient being fitted and v_{prior} is a "prior" taken from NUPACk. This loss function is then minimized using the L-BFGS algorithm implemented in the 'optimize.minimize' function in scipy python package.

The NUPACK-compatible NN model

We trained the improved NN model on the Array Melt training data set, containing 24,971 of the 27,732 sequences that passed quality control and two-state criteria, and tuned the model on a validation set of 1,315 sequences. We first adjusted salt concentration to 1M Na$^+$ to be compatible with NUPACK and other softwares, and then corrected systematic errors by linearly shifting Tm and ΔG, such that the medians of each construct type matches those of NUPACK predictions. The underlying assumption is that NUPACK predictions have minimal bias but could be improved by explaining more variance. Improved parameters for NUPACK were fitted with aforementioned weighted linear regression. We allowed an extra intercept parameter to adjust mean absolute errors. From sequence-structure pairs in the training set, we extracted features of stacks and loops. Stacks include NN Watson-Crick pairs and terminal stack penalty, as in NUPACK; loops include the following:

1) Hairpin loops, defined as a subsequence $\phi_{i:j}$ where $i \cdot j$ is the closing base pair. We extracted "hairpin_mismatch" features as in NUPACK, $\Delta G^{hairpinmm}_{\phi_{j-1},\phi_j,\phi_i,\phi_{i+1}}$, and introduced an intermediate "hairpin_loop_mid" feature, $\Delta G^{hairpinmid}_{\phi_{[i+2:j-2]}}$ to account for sequence variations in the middle of hairpin loops for triloops and tetraloops. The final energy of hairpin is $\Delta G^{hairpin}_{\phi_{[i:j]}} = \Delta G^{hairpinsize}_n + \Delta G^{hairpinseq}_{\phi_{[i:j]}}$ as in[19], where

$$\Delta G^{hairpinseq}_{\phi_{[i:j]}} = \Delta G^{hairpinmm}_{\phi_{j-1},\phi_j,\phi_i,\phi_{i+1}} + \Delta G^{hairpinmid}_{\phi_{[i+2:j-2]}} (+ \Delta G^{terminalbp}_{\phi_j,\phi_i} \; if \; triloop) \quad (15)$$

2) Interior loops, defined from 5' to 3' as two subsequences $\phi_{[i:d]}$ and $\phi_{[e:j]}$. Any other interior loop types were defined as in[19], apart from the $\Delta G^{interiormm}_{\phi_{j-1},\phi_j,\phi_i,\phi_{i+1}}$ were fitted separately instead of shared with terminal mismatches as in NUPACK, and that we introduced an intermediate parameter for the interaction between the two base pairs surrounding 1x1 mismatches $\Delta G^{interiormmstacks}_{\phi_i,\phi_j,\phi_{i+2},\phi_{j-2}}$.

This results in 178 features for ΔH and ΔG each. Parameters for hairpin and interior loop sizes were taken from NUPACK and clamped during fitting. Lookup tables of $\Delta G^{hairpinseq}_{\phi_{[i:j]}}$ and $\Delta G^{interiorseq}_{\phi_{[i:d]},\phi_{[e:j]}}$ for "interior_1_1", "interior_1_2", and "interior_2_2" were calculated from the fitted parameters. All updated parameters were then written to a NUPACK-compatible json file.

The richer parameter NN model

We trained the richer parameter model on the same Array Melt training data set as for the NUPACK-compatible NN model. For feature extraction, we extracted NN Watson-Crick pairs and terminal stack penalty as in NUPACK, and the following parameters for loops:

1) Hairpin loops. For triloops and tetraloops, $\Delta G^{hairpinseq}_{\phi_{[i:j]}} = \Delta G^{hairpinmm}_{\phi_{j-1},\phi_j,\phi_i,\phi_{i+1}} + \Delta G^{hairpinloop}_{\phi_{[i+1:j-1]}}$, where $\Delta G^{hairpinloop}_{\phi_{[i+1:j-1]}}$ is the full sequence of the hairpin loop, in contrast to only the middle of loops in the NUPACK-compatible model. For hairpin loops longer than tetraloop, we applied a single loop sequence parameter $\Delta G^{hairpinloop}_{long}$. The final energy of hairpin is also $\Delta G^{hairpin}_{\phi_{[i:j]}} = \Delta G^{hairpinsize}_n + \Delta G^{hairpinseq}_{\phi_{[i:j]}}$.

2) Interior loops. For single and double bulges or single mismatches, the sequence-dependent parameter is fitted for each sequence, $\Delta G^{interiorseq}_{\phi_{[i:d]},\phi_{[e:j]}}$. For double and triple mismatches,

$$\Delta G^{interiorseq}_{\phi_{[i:d]},\phi_{[e:j]}} = \Delta G^{interiormmstacks}_{\phi_i,\phi_j,\phi_{i+2},\phi_{j-2}} + \Delta G^{interiormm}_{\phi_{i+1},\phi_{d-1},\phi_{e+1},\phi_{j-1}}.$$

Deep learning models

Closest sequence lookup baseline model

We defined the distance between any two sequence-structure pairs by summing up the edit distance between the sequences and the dot-bracket secondary structure representation. For every query sequence-structure pair, we find the k closest sequence-structure pairs in the training set, whose distance-weighted sum becomes the predicted value for the query using python package *scipy*[36]. We found the best value for k using the held-out Array Melt validation data.

Graph neural network model

We implemented the graph neural network using python packages PyTorch and PyTorch Geometric[37]. The final model has 4 graph transformer layers with a 125-vec node attribute at each layer, stacked with dropout of 0.01 at each layer, a Set2Set module with 10 processing steps, and a 1 fully connected hidden layer with 128 channels and a dropout ratio of 0.49. We normalized dH and Tm of the training data to fall within 0 and 1, and used the sum of the

root mean square error (RMSE) of both dH and Tm as the loss function and trained with Adam optimizer for 200 epoches. The model has 287,136 parameters. The validation model registry was monitored and stored on weight and biases, available at https://wandb.ai/keyuxi/NNN_GNN/.

Usage of the NUPACK-compatible model

The parameter file for the NUPACK-compatible NN model is available in the supplementary material as 'NUPACK_compatible_NN_model.json'. To use this file, download and point to it when using the python package NUPACK 4[38]:

```
my_model =
nupack.Model(material='/path/to/parameter/file/NUPACK_compatible_NN_model.json', …)
```

Code availability

Code for array data preprocessing and curve fitting could be found at https://github.com/keyuxi/array_analysis, and all other analysis at https://github.com/keyuxi/nnn_paper.

References

1. Kunkel, T. A. DNA Replication Fidelity. *J. Biol. Chem.* **279**, 16895–16898 (2004).

2. Rajski, S. R., Jackson, B. A. & Barton, J. K. DNA repair: models for damage and mismatch recognition. *Mutat. Res. Mol. Mech. Mutagen.* **447**, 49–72 (2000).

3. Untergasser, A. *et al.* Primer3—new capabilities and interfaces. *Nucleic Acids Res.* **40**, e115 (2012).

4. Kibbe, W. A. OligoCalc: an online oligonucleotide properties calculator. *Nucleic Acids Res.* **35**, W43–W46 (2007).

5. Labun, K. *et al.* CHOPCHOP v3: expanding the CRISPR web toolbox beyond genome editing. *Nucleic Acids Res.* **47**, W171–W174 (2019).

6. Majikes, J. M. *et al.* Revealing thermodynamics of DNA origami folding via affine transformations. *Nucleic Acids Res.* **48**, 5268–5280 (2020).

7. Zadeh, J. N. *et al.* NUPACK: Analysis and design of nucleic acid systems. *J. Comput. Chem.* **32**, 170–173 (2011).

8. SantaLucia, J. & Hicks, D. The Thermodynamics of DNA Structural Motifs. *Annu. Rev. Biophys. Biomol. Struct.* **33**, 415–440 (2004).

9. SantaLucia, J. A unified view of polymer, dumbbell, and oligonucleotide DNA nearest-neighbor thermodynamics. *Proc. Natl. Acad. Sci.* **95**, 1460–1465 (1998).

10. SantaLucia, John, Allawi, H. T. & Seneviratne, P. A. Improved Nearest-Neighbor Parameters for Predicting DNA Duplex Stability. *Biochemistry* **35**, 3555–3562 (1996).

11. Zuker, M. On Finding All Suboptimal Foldings of an RNA Molecule. *Science* **244**, 48–52 (1989).

12. Schroeder, S. J. & Turner, D. H. Optical Melting Measurements of Nucleic Acid Thermodynamics. *Methods Enzymol.* **468**, 371–387 (2009).

13. Spink, C. H. Differential Scanning Calorimetry. in *Methods in Cell Biology* vol. 84 115–141 (Academic Press, 2008).

14. Oliveira, L. M., Long, A. S., Brown, T., Fox, K. R. & Weber, G. Melting temperature measurement and mesoscopic evaluation of single, double and triple DNA mismatches. *Chem. Sci.* **11**, 8273–8287 (2020).

15. Bae, J. H., Fang, J. Z. & Zhang, D. Y. High-throughput methods for measuring DNA thermodynamics. *Nucleic Acids Res.* **48**, e89–e89 (2020).

16. You, Y., Tataurov, A. V. & Owczarzy, R. Measuring thermodynamic details of DNA hybridization using fluorescence. *Biopolymers* **95**, 472–486 (2011).

17. Buenrostro, J. D. *et al.* Quantitative analysis of RNA-protein interactions on a massively parallel array reveals biophysical and evolutionary landscapes. *Nat. Biotechnol.* **32**, 562–568 (2014).

18. Marklund, E., Ke, Y. & Greenleaf, W. J. High-throughput biochemistry in RNA sequence space: predicting structure and function. *Nat. Rev. Genet.* **24**, 401–414 (2023).

19. Fornace, M. E., Porubsky, N. J. & Pierce, N. A. A Unified Dynamic Programming Framework for the Analysis of Interacting Nucleic Acid Strands: Enhanced Models, Scalability, and Speed. *ACS Synth. Biol.* **9**, 2665–2678 (2020).

20. Tikhomirova, A., Beletskaya, I. V. & Chalikian, T. V. Stability of DNA Duplexes Containing GG, CC, AA, and TT Mismatches. *Biochemistry* **45**, 10563–10571 (2006).

21. Khandelwal, G. & Bhyravabhotla, J. A Phenomenological Model for Predicting Melting Temperatures of DNA Sequences. *PLOS ONE* **5**, e12433 (2010).

22. Yan, Z., Hamilton, W. L. & Blanchette, M. Graph neural representational learning of RNA secondary structures for predicting RNA-protein interactions. *Bioinformatics* **36**, i276–i284 (2020).

23. Zhao, Q. *et al.* Review of machine learning methods for RNA secondary structure prediction. *PLOS Comput. Biol.* **17**, e1009291 (2021).

24. Shi, Y. *et al.* Masked Label Prediction: Unified Message Passing Model for Semi-Supervised Classification. Preprint at https://doi.org/10.48550/arXiv.2009.03509 (2021).

25. Veličković, P. *et al.* Graph Attention Networks. Preprint at https://doi.org/10.48550/arXiv.1710.10903 (2018).

26. Vinyals, O., Bengio, S. & Kudlur, M. Order Matters: Sequence to sequence for sets. Preprint at http://arxiv.org/abs/1511.06391 (2016).

27. Spasic, A. *et al.* Improving RNA nearest neighbor parameters for helices by going beyond the two-state model. *Nucleic Acids Res.* **46**, 4883–4892 (2018).

28. She, R. *et al.* Comprehensive and quantitative mapping of RNA–protein interactions across a transcribed eukaryotic genome. *Proc. Natl. Acad. Sci.* **114**, 3619–3624 (2017).

29. Magoč, T. & Salzberg, S. L. FLASH: fast length adjustment of short reads to improve genome assemblies. *Bioinformatics* **27**, 2957–2963 (2011).

30. Altschul, S. F. & Gish, W. [27] Local alignment statistics. in *Methods in Enzymology* vol. 266 460–480 (Academic Press, 1996).

31. Denny, S. K. *et al.* High-Throughput Investigation of Diverse Junction Elements in RNA Tertiary Folding. *Cell* **174**, 377-390.e20 (2018).

32. Köster, J. & Rahmann, S. Snakemake—a scalable bioinformatics workflow engine. *Bioinformatics* **28**, 2520–2522 (2012).

33. Newville, M., Stensitzki, T., Allen, D. B. & Ingargiola, A. LMFIT: Non-Linear Least-Square Minimization and Curve-Fitting for Python. (2014) doi:10.5281/zenodo.11813.

34. Fischler, M. A. & Bolles, R. C. Random sample consensus: a paradigm for model fitting with applications to image analysis and automated cartography. *Commun. ACM* **24**, 381–395 (1981).

35. Owczarzy, R. *et al.* Effects of Sodium Ions on DNA Duplex Oligomers: Improved Predictions of Melting Temperatures. *Biochemistry* **43**, 3537–3554 (2004).

36. Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat. Methods* **17**, 261–272 (2020).

37. Fey, M. & Lenssen, J. E. Fast Graph Representation Learning with PyTorch Geometric. Preprint at https://doi.org/10.48550/arXiv.1903.02428 (2019).

38. Fornace, M. E. *et al. NUPACK: Analysis and Design of Nucleic Acid Structures, Devices, and Systems.* https://chemrxiv.org/engage/chemrxiv/article-details/636c7089b588507d0045f283 (2022) doi:10.26434/chemrxiv-2022-xv98l.