Activating A1 adenosine receptor signaling boosts early pulmonary neutrophil recruitment in

aged mice in response to *Streptococcus pneumoniae* infection

Shaunna R. Simmons, Sydney E. Herring, Essi Y.I Tchalla, Alexsandra P. Lenhard, Manmeet

Bhalla, and Elsa N. Bou Ghanem[*]

Department of Microbiology and Immunology, School of Medicine, University at Buffalo,

Buffalo, NY, USA

Running title: A1 receptor signaling recruits PMNs to the lungs

Key words: extracellular adenosine, pneumococcus, PMNs, A1R, pneumonia, migration,

inflammation, immunosenescence, lung

* Corresponding author

E-mail: Elsaboug@buffalo.edu

ABSTRACT

Background: *Streptococcus pneumoniae* (pneumococcus) is a leading cause of pneumonia in

older adults. Successful control of pneumococci requires robust pulmonary neutrophil influx early

in infection. However, aging is associated with aberrant neutrophil recruitment and the

mechanisms behind that are not understood. Here we explored how neutrophil recruitment

following pneumococcal infection changes with age and the host pathways regulating this.

Results: Following pneumococcal infection there was a significant delay in early neutrophil

recruitment to the lungs of aged mice. Neutrophils from aged mice showed defects in trans-

endothelial migration *in vitro* compared to young controls. To understand the pathways involved,

we examined immune modulatory extracellular adenosine (EAD) signaling, that is activated upon

cellular damage. Signaling through the lower affinity A2A and A2B adenosine receptors had no

effect on neutrophil recruitment to infected lungs. In contrast, inhibition of the high affinity A1

receptor in young mice blunted neutrophil recruitment to the lungs following infection. A1 receptor

inhibition decreased expression of CXCR2 on circulating neutrophils, which is required for trans-

endothelial migration. Indeed, A1 receptor signaling on neutrophils was required for their ability

to migrate across endothelial cells in response to infection. Aging was not associated with defects

in EAD production or receptor expression on neutrophils. However, agonism of A1 receptor in

aged mice rescued the early defect in neutrophil migration to the lungs and improved control of

bacterial burden.

Conclusions: This study suggests age-driven defects in EAD damage signaling can be targeted to

rescue the delay in pulmonary neutrophil migration in response to bacterial pneumonia.

BACKGROUND

Streptococcus pneumoniae (pneumococcus) are encapsulated, Gram positive bacteria that are responsible for an estimated 1.6 million deaths each year (1). S. pneumoniae remain the leading cause of community acquired bacterial pneumonia in individuals over the age of 65 (2-4). Disease and mortality caused by these bacteria persist despite available vaccines and antibiotics (3, 5, 6). Older adults are at an increased risk of infection and serious disease from pneumococcal pneumonia due to progressive deleterious molecular, cellular and systemic changes termed the "Hallmarks of Aging" (7, 8), that can affect the ability of the host to mount an efficient and balanced immune response. These hallmarks include aberrant genetic changes, altered metabolic responses, cellular senescence and chronic inflammation(7, 8). Cellular senescence affects several cell types including immune cells, causing them to be less efficient at pathogen clearance (7, 8). Cellular senescence is also associated with increased production of pro-inflammatory cytokines leading to an increase in basal inflammation known as inflammaging (7, 8). Increased inflammation with age contributes to a reduction in the ability of immune cells to respond to acute stimuli, further dysregulating the immune response (7-9). A better understanding of how immune cell responses change with host aging upon pneumococcal infection, and the signaling pathways that control these responses, is necessary to promote better protection against disease.

An important mediator of host protection following pneumococcal pneumonia is the polymorphonuclear leukocyte (PMNs) response. PMNs are the first cell to be recruited to the lungs following pneumococcal infection (10, 11) and this response is necessary for host protection. Individuals with neutropenia are more susceptible to pneumonia and experimental depletion of PMNs early in infection result in reduced ability to control bacterial numbers and accelerated death in murine models of infection (12, 13). Upon pulmonary infection, PMNs are recruited from the

circulation into the lung tissue where they kill bacteria via several antimicrobial effector functions,

such as release of preformed antimicrobial granules, reactive oxygen species, phagocytosis, and

neutrophil extracellular traps (14). However, an effective PMN response in the pulmonary

environment must be balanced, where PMNs need to be recruited in quickly to clear bacteria and

then resolve. An exacerbated PMN response without resolution is damaging to host tissue due to

the antimicrobial products produced by PMNs, and without proper resolution PMNs contribute to

lung damage which worsens disease outcomes (12, 15). There is evidence that PMN influx to the

lungs later in infection is dysregulated with age as biopsy samples from older adults show an

increase in the percentage of PMNs in the lungs compared to younger patients following

pneumococcal pneumonia, and this can be recapitulated in mouse models (16, 17). However, how

neutrophil recruitment changes with age early in infection, when PMNs are beneficial to the host

in clearing the pathogen, and the pathways controlling PMN influx remain unexplored.

We previously found that the production of extracellular adenosine (EAD) is crucial for

regulating PMN influx to the lungs in young mice (10). Adenosine accumulates in the extracellular

space as a result of cellular damage. Upon tissue injury by a variety of insults, including infection,

ATP released from cells is metabolized to EAD by the sequential action of the extracellular

ectonucleotidases CD39 and CD73 (18). At baseline, EAD levels are very low/ undetectable in

extracellular spaces but can increase more than 10-fold upon infection (19). EAD can signal via 4

G-protein coupled receptors, A1, A2A, A2B and A3 that respond to EAD in a dose-dependent

manner (18, 20). A1 is high affinity and is activated by lower levels of adenosine while A2A and

A2B are lower affinity and require higher levels of adenosine to become activated (18). The

different adenosine receptors are also coupled to different G proteins that can inhibit (Gi) or

stimulate (Gs) adenylyl cyclase or activate phospholipase C (Gq), and therefore can have opposite

effects on downstream signaling and cell function. These receptors are ubiquitously expressed on

many cell types including PMNs (21).

Changes in EAD production and signaling occur with aging (22-27). Although the role of

this pathway in immunosenescence remains largely unexplored, we previously found that aging is

accompanied by changes in expression of EAD enzymes on murine PMNs at baseline and that

EAD signaling controlled PMN antibacterial function (21, 28, 29). However, how the levels of

extracellular adenosine and the receptors signaled through change after pneumococcal infection

and across host age, and their role in PMN migration, is unknown. In this study, we asked how

early PMN recruitment changes with host aging and if the extracellular adenosine pathway

regulates that during bacterial pneumonia. We tested the hypothesis that signaling through the high

affinity A1 adenosine receptor would be important early in pneumococcal infection when

adenosine levels are lower to help draw the PMNs to the lung, while signaling through the lower

affinity receptors may be required late in infection to help resolve PMN influx and that these

dynamics change during host aging.

METHODS

Mice

Young (8-12 weeks) and old (18-22 months) C57BL/6 (B6) male mice were obtained from the

National Institute on Aging colony or from Jackson Laboratories (Bar Harbor, ME). A2BR$^{-/-}$ mice

on a B6 background (B6.129P2-Adora2btm1Till/J) and A2AR$^{-/-}$ mice on a Balb/c background

(C;129S-Adora2atm1Jfc/J) were purchased from Jackson Laboratory and bred at our facility. Wild

type C57BL/6J and Balb/cJ mice were used as controls. Since males are more susceptible to

pneumococcal infection compared to females (30) and due to the limited availability of aged

cohorts, experiments were performed in male mice. All mice were housed at the University at

Buffalo in specific pathogen free housing for at least 2 weeks before use in experiments. All

experiments were conducted in accordance with the Institutional Animal Care and Use Committee

guidelines.

Bacteria

Streptococcus pneumoniae TIGR4 strain (serotype 4) was a kind gift from Andrew Camilli.

Briefly, bacteria were grown to mid-exponential phase at 37ºC at 5% CO_2 in Todd Hewitt broth

supplemented with 0.5% yeast extract and oxyrase as previously described (20).

PMN Isolation

Bone marrow cells were collected from the femurs and tibias of uninfected mice by flushing with

RPMI supplemented with 10% FBS and 2mM EDTA. Red blood cells were then lysed, and

remaining cells were washed, and resuspended in PBS. PMNs were separated via density

centrifugation using histopaque 1119 and 1077 (Sigma) as previously described (20, 21, 31, 32).

PMNs were then resuspended at the desired concentration in Hank's Balanced Salt Solution/0.1%

gelatin with no Ca^{2+} or Mg^{2+} and kept on ice until use. The purity of PMNs was confirmed using

flow cytometry and 85-90% of enriched cells were positive for Ly6G and CD11b.

Mouse Infections and Treatments

Mice were infected intratracheally (i.t.) with 50ul of *S. pneumoniae* TIGR4 at the indicated

concentrations. Mice were first anesthetized using isoflurane and the bacterial inoculum pipetted

into the trachea with the tongue gently pulled out to ensure direct delivery into the lungs. Following

infection, mice were euthanized at the indicated time points, blood was collected by portal vein

snip to determine bacteremia. Mice were then perfused with PBS through the right ventricle and

the lungs harvested. Following harvest, the lungs were minced into small pieces, mixed well and

half of each lung sample was homogenized in sterile PBS to plate for CFU, and the remaining half

was digested and used for flow cytometry analysis. Lung and blood samples were diluted in sterile

PBS and plated on blood agar to enumerate CFU. For experiments where the effect of A1 receptor

signaling was measured, the A1 receptor inhibitor, 8-Cyclopentyl-1,3-dipropylxanthine

(DPCPX)(28, 33, 34) or A1 receptor agonist 2-Chloro-N6-cyclopentyladenosine (35) were

purchased from Sigma Aldrich, dissolved in DMSO and filter sterilized by passing through a

0.22μm filter prior to use. Mice were given intraperitoneal (i.p.) injections of 1mg/kg of the A1

inhibitor or 0.1mg/kg of the A1 agonist or vehicle control at -1 and 0 (immediately before

challenge) days relative to infection. For experiments where the effect of A2A receptor signaling

was measured, the A2A receptor inhibitor, 3,7-Dimethyl-1-propargylxanthine (D134) (36), or A2A

receptor agonist CGS21680 (37), were purchased from Sigma, dissolved in DMSO and filter

sterilized by passing through a 0.22 μm filter prior to use. Mice were given i.p injections of 5mg/kg

of the A2A antagonist or 2mg/kg of the A2A agonist or vehicle control at -18 and 0 hours,

immediately before challenge for pre-treated (designated as PRE) and 18 hours post challenge for

post-treated groups (designated as POST). Control animals were treated with vehicle control.

Flow Cytometry

The lungs were digested into a single cell suspension in RPMI 1640 containing 10% FBS, 2mg/mL

type 2 collagenase and 50 U/mL DNase (Worthington) at 37°C/ 5% CO_2 as previously described

(20). Blood was collected by portal vein snips in EDTA as an anticoagulant. Red blood cells were

removed with a hypotonic lysis buffer (Lonza). Cells were resuspended in FACS buffer (HBSS/

1% FBS), treated with Fc block (anti-mouse clone 2.4G2) and stained with the following

antibodies purchased from BD Biosciences or Invitrogen: Anti-Ly6G (IA8), anti-CD11b (M1/70),

Anti-CD45 (30-F11). For adenosine receptor staining, the BD Cytofix/Cytoperm kit was used to

permeabilize the cells and the cells stained with the following unconjugated rabbit anti-adenosine

receptor antibodies from Abcam as previously described (21): A2a (ab3461), A2b (ab222901), A3

(ab203298) and A1 (ab82477). Rabbit polyclonal IgG (ab37415) was used as an isotype control

and secondary PE-conjugated anti-Rabbit IgG was used (12473981; Invitrogen). Fluorescent

intensities were measured on a BD Fortessa and at least 20,000 events were analyzed using FloJo.

Trans Endothelial Migration Assay

To determine trans-endothelial migration of PMNs *in vitro*, C57BL/6 murine primary lung

endothelial cells were grown according to manufacturer's protocol (Cell Biologics) and 5µm

transwells (Corning) were seeded at 10^6 cells/transwell and grown for 2 days at 37°C/ 5% CO_2

until confluent. The bottom of the transwell was infected with *S. pneumoniae* TIGR4 at an MOI

of 40. PMNs were isolated from the bone marrow of indicated mouse strains and $5x10^5$ PMNs

placed in the top of the transwell and incubated at 37°C at 5% CO2 for 1.5 hours. The number of

PMNs that migrated through the endothelial layer into the bottom of the transwell was determined

by collecting the cells from the bottom well and lysing them with 10% Triton X-100 to release

MPO. MPO was measured in each well by using 2,2′-Azino-bis(3-ethylbenzothiazoline-6-sulfonic

acid) diammonium salt (ABST) in the presence of hydrogen peroxide (10) and serial dilutions of

a known numbers of PMNs were used to establish a standard curve, which was then used to

quantify the number of migrated neutrophils as previously described (10).

Measurement of Adenosine

Mice were infected i.t with $2.5x10^4$ CFU *S. pneumoniae* TIGR4. Following infection, mice were

euthanized at indicated time points and blood was collected and placed into microtainer tubes (BD)

and centrifuged at 9000 rpm to collect sera. Circulating levels of adenosine were measured using

the Adenosine Assay Kit (Fluorometric) from BioVision per manufacturer's instructions. Results

were read using a Biotek plate reader.

Statistics

Statistics were analyzed using Prism 9 (GraphPad). Bar graph and scatter plot data are presented

as Mean +/- SD, and line graph data are presented as Mean +/- SEM. All data were tested for

normality using Shapiro-Wilk test prior to statistical analysis. Significant difference between

groups was determined by Student's t-test, Mann-Whitney, Kruskal Wallis or one-way ANOVA as

indicated. Difference in survival was determined by the log-rank (Mantel-Cox) test. *p* values *<0.05*

were considered significant. * indicates *p <0.05*, ** indicates *p <0.001*, *** indicates *p<0.0001*.

RESULTS

Rapid PMN influx to the lungs following *S. pneumoniae* infection is impaired with aging

To determine how PMN influx to the lungs changes with age throughout the course of

pneumococcal infection, young and aged mice were infected i.t. with 2.5×10^4 colony forming units

(CFU) of *S. pneumoniae* TIGR4 and PMN recruitment to the lungs was measured by flow

cytometry over time following infection. Young mice had significant PMN influx to the lungs

within 6 hours of infection, and further increase in pulmonary PMN numbers by 18 and 48 hours

post infection (hpi) (**Fig 1A**). With age there was a delay in PMN influx early in infection with no

increase from baseline by 6 hours, but a significant increase by 18 and 48 hpi (**Fig 1A**). When

compared to young controls, aged mice had approximately 4-fold lower PMNs in the lungs at 6

hpi (**Fig 1A**). By 18 and 48 hpi, PMNs were present in the lungs of aged mice at similar levels to

young mice. This delay in PMN influx to the lungs with age was associated with higher

pneumococcal CFU in the lungs throughout the course of infection (**Fig 1B**) and significantly

higher bacteremia at 18 hpi (**Fig 1C**). When normalized to bacterial burden, aged mice displayed

significantly reduced PMN numbers in the lungs compared to young controls at 6 hours and

significantly higher numbers of PMNs at 48 hpi (**Fig S1A**). These data indicate that with age there

is an initial delay in neutrophil influx to the lungs following pneumococcal infection that is

associated with impaired bacterial control.

PMN trans-endothelial migration in response to *S. pneumoniae* is impaired with aging

For PMNs to enter the lungs to control pneumococcal infection, these cells must be

recruited from the circulation and migrate through the endothelium to the infected pulmonary

tissue. To better understand the age-related impairment in PMN influx, we measured neutrophil

trans-endothelial migration. Using an *in vitro* transwell model, we measured neutrophil migration

across primary mouse lung endothelial cells (isolated from young mice) in response to *S.*

pneumoniae as a stimulus. At baseline there was no difference in neutrophil migration, however

upon infection, there was a significant decline in the number of PMNs that migrated through the

endothelial cell layer when PMNs were isolated from aged mice compared to young controls (**Fig**

2). These data suggest that there is an intrinsic defect in trans-endothelial migration by PMNs from

aged mice.

***S. pneumoniae* infection alters expression of extracellular adenosine pathway components on**

PMNs

To test if the extracellular adenosine (EAD) pathway regulates the age-driven changes in

PMN pulmonary influx following *S. pneumoniae* infection, we first needed to understand its role

in young hosts. To do so, we tracked the expression of the different pathway components over time

following infection (gating strategy Fig S2). As this is a pulmonary infection and the damage

response is initiated in the lungs, we first tracked expression of the EAD producing and degrading

enzymes in the lungs of young mice. We measured geometric mean florescence intensity (MFI) on

total lung cells as an indicator of expression levels and found that the EAD producing enzymes

CD39 and CD73 as well the EAD degrading enzyme ADA were expressed in the lung tissues and

that their levels were maintained constant over 18 hours of infection (**Fig S3**). We had challenges

measuring EAD levels in the bronchioalveolar lavage fluid (not shown), so instead we assessed

circulating levels and found overall levels of adenosine in the circulation did not significantly

change over the course of infection in young mice (**Fig 3**). We then tracked the expression of the

four adenosine receptors on the surface of PMNs in the lungs using flow cytometry. We found that

at baseline, the majority of PMNs in the lungs of young mice express A1, A2A, and A2B while

only about half of PMNs express A3 on the cell surface (**Fig 4A**). To determine if the level of

receptor expression on PMNs changes over time, we stained for adenosine receptor expression

following infection. We found that all four adenosine receptors showed a significant decrease in

expression within the first 18 hours of infection, followed by a recovery of levels comparable to

baseline by 48 hpi in young mice (**Fig 4B**). When we measured fold change from baseline, we

observed around 2-fold decrease by 6 hours for A2A and A2B and by 18 hours for A1 and A3

followed by recovery for all four receptors by 48 hpi in young mice (**Fig 4C**).

As PMNs are recruited from the blood, we also measured receptor expression in the

circulation. In circulating PMNs of uninfected young mice, A1, A2A, and A2B receptors were

highly expressed, however, very few PMNs expressed A3 (**Fig 5A**). When we measured how

receptor expression changes over time in circulating PMNs, we found that overall receptor

expression remains constant in young mice (**Fig 5B**). Unlike in the lungs, circulating PMNs in

young mice expressed A2A, A2B, and A3 at similar level over the course of infection (**Fig 5C**).

The expression of A1 in the circulation showed a slight, but significant 1.5-fold decrease at 6 hpi

but returned to baseline levels by 18 hpi) (**Fig 5C**). These data show that PMNs in the lungs and

circulation of young mice highly express adenosine receptors A1, A2A, and A2B on their surface,

and that there are dynamic changes in adenosine receptor expression on PMNs in response to

pneumococcal infection.

Age-driven changes in EAD-pathway components following *S. pneumoniae* infection

We next wanted to understand how the adenosine pathway components change with host

aging. We measured adenosine levels in the circulation of aged mice to see if the dysregulation in

PMN influx was due to a lack of extracellular adenosine production. We found that early in

infection, there was no significant difference in the amount of adenosine in the circulation when

compared to young mice (**Fig 3**). By 48 hpi, aged mice had significantly higher levels of adenosine

(**Fig 3**), suggesting exacerbated damage as the infection progressed in aged hosts, which fits with

the higher bacterial loads observed in aged hosts (**Fig 1**). When we measured adenosine pathway

enzymes in the lungs, we found that at baseline, the lungs of aged mice express significantly higher

amounts of the EAD producing enzymes CD73 and CD39, while the levels of the EAD degrading

enzyme ADA were higher, but did not reach statistical significance (**Fig S3**). When we tracked

enzyme expression through the course of infection, we found that unlike young controls, the

expression of all three enzymes had significantly decreased by 18 hpi in lungs of aged mice (**Fig

S3**).

Next, we analyzed receptor expression on PMNs in the lungs and circulation. We found

that in the lungs and blood at baseline there was no change in the percentage of PMNs expressing

A1, A2A, A2B, or A3 when compared to young mice with the majority of cells expressing A1,

A2A, and A2B (**Fig 4A and Fig 5A**). To look at receptor expression changes overtime, we

measured MFI and calculated fold change from baseline. In both lung (**Fig 4B**) and blood (**Fig 5B**)

there was no significant difference in receptor expression between young and aged mice and the

expression of A1 receptor trended to be higher on circulating PMNs at 6 hpi in aged mice compared

to young controls (**Fig 5B**). We did find that PMNs in the lungs of aged mice decreased levels of

adenosine receptors early in infection respective to their own baseline, however, unlike in young

mice where receptor expression returned to baseline levels by 48 hours, with age there was no

rebound of adenosine receptor expression at this later time point (**Fig 4C**). A similar pattern of

decreased receptor expression over time in aged mice on circulating PMNs was also seen, where

circulating PMNs maintained a 2-fold reduction in expression of A1, A2A and A2B following

infection (**Fig 5C**). These data show that there is no defect with age in production of extracellular

adenosine and or adenosine receptor expression on PMNs. Rather, the decrease in adenosine

receptor expression on PMNs during the course of infection is more pronounced and may result in

decrease responsiveness to adenosine signaling in the aged hosts.

A2B adenosine receptor plays no role in PMN pulmonary influx following *S. pneumoniae*

infection

To determine if the individual adenosine receptors played a role in PMN pulmonary

migration, we focused on the receptors highly expressed on PMNs (A1, A2A and A2B). We first

examined the A2B receptor. To examine the role of this receptor in PMN influx to the lungs *in*

vivo, WT C57BL/6 and A2BR$^{-/-}$ mice were infected intratracheally with 2.5x10^4 CFU *S.*

pneumoniae and the total number of PMNs in the lungs was determined by flow cytometry at 6,

18, and 48 hpi. We found that there was no difference in the total amount of PMNs in the lungs at

any of the time points tested between the mouse strains (**Fig 6**). Bacterial burden in the lungs also

did not differ (data not shown and supporting what was previously published (38)) and even when

normalized for bacterial numbers, PMN influx into the lungs in response to *S. pneumoniae*

infection was comparable between WT and A2BR$^{-/-}$ mice (**Fig S1B**). These data suggest that A2B

receptor signaling does not regulate PMN influx to the lung of *S. pneumoniae* infected mice.

A2A adenosine receptor plays no role in PMN pulmonary influx following *S. pneumoniae* infection

Next, we examined the role of A2A receptor signaling in PMN influx to the lungs following

infection. To examine the role of this receptor *in vivo*, A2AR$^{-/-}$ and WT BALB/c controls were

infected intratracheally with 2.5x10^4 CFU *S. pneumoniae* TIGR4 and total number of PMNs in the

lungs was determined at 6 and 48 hpi by flow cytometry. There was no difference in the number

of PMNs in the lungs of A2AR$^{-/-}$ or WT mice at either time point (**Fig 7A**). To confirm that this

result was not due to differences in bacterial burden, we analyzed *S. pneumoniae* CFU in both the

lung (**Fig 7B**) and blood (**Fig 7C**) and found no difference in bacterial burden across mouse strains.

Additionally, there was no difference in survival following infection between A2AR$^{-/-}$ and WT

mice (**Fig 7D**). Since these experiments were done using mice on a Balb/c background, we also

analyzed the role of A2A in C57BL/6 mice using a specific pharmacological A2A receptor

inhibitor, 3,7-Dimethyl-1-propargylxanthine. We first tested the role of A2A early in infection by

treating mice with the A2A receptor antagonist or vehicle control 18 hours prior to bacterial

challenge (PRE). We found no difference in total number of PMNs in the lungs (**Fig S4A**), total

bacterial burden in the lung (**Fig S4B**) or bacteremia (**Fig S4C**) by 48 hpi. Similarly, inhibition of

A2A prior to infection had no effect on mouse survival (**Fig S4D).** As A2A is a lower affinity

receptor that may play a role later in infection as damage and EAD levels build up, we also tested

its role later in infection, by timed inhibition 18 hours post (POST) bacterial challenge. Again, we

found no difference in total bacterial burden in the lung (**Fig S4B)** or bacteremia (**Fig S4C**) by 48

hpi in POST treated mice. However, there was a decrease in survival in the mice treated with A2A

receptor inhibitor post infection (**Fig S4D**) suggesting that A2A receptor may play a protective role

later in infection. Overall, these data suggest that A2A receptor signaling does not regulate PMN

influx to the lungs of young mice following *S. pneumoniae* challenge.

A1 receptor signaling is required for PMN pulmonary influx in response to *S. pneumoniae*

infection

To determine the role of A1 receptor signaling in PMN migration, young mice were treated

i.p. with a specific A1 receptor inhibitor (whose specificity we confirmed in prior publications (28,

33)) or vehicle control. Mice were then infected intratracheally with $2.5x10^4$ CFU *S. pneumoniae*

TIGR4 one day post drug treatment. Total PMN influx to the lungs of infected mice was measured

by flow cytometry at 6, 18, and 48 hours. We found that inhibition of A1 resulted in a significant

delay in PMN influx to the lungs at 6 hpi (**Fig 8A-B**). At 6 hpi, there was a significant decrease in

total PMN numbers in the lungs (**Fig 8A**) and the amount of MPO (a marker of PMNs) in lung

homogenates (**Fig 8B**) of A1 inhibited mice. To understand the mechanism of the decline in PMN

recruitment, we examined the expression of CD18 and CXCR2 on the surface of PMNs in the

circulation. We found no difference in the numbers of circulating PMNs in A1 receptor inhibited

versus vehicle treated mice (**Fig S5**). We further found that while A1 receptor inhibition had no

effect of CD18 expression (**Fig 8C**), it significantly reduced CXCR2 expression at 6 hpi (**Fig 8D**),

the time point at which PMN number in the lungs is lower (**Fig 8A**). As CXCR2 was reported to

be required for trans-endothelial migration in response to pneumococcal infection (11, 14, 39-41)

, we directly assessed the role of A1 on PMN trans-endothelial migration. Young mice were treated

i.p. with an A1 receptor inhibitor, and PMNs were then isolated, and trans-endothelial migration

was measured using the same *in vitro* transwell assay described above. A1 receptor inhibited PMNs

displayed a significant decrease in their ability to migrate trans-endothelially compared to VC

treated PMNs, indicating that A1 signaling on PMNs is required for efficient trans-endothelial

migration (**Fig 8E**). These data suggest that A1 adenosine receptor signaling is required for

recruitment of PMNs to the lungs in response to *S. pneumoniae* infection.

Activation of A1 receptor signaling rescues the ability of PMNs from old mice to be

recruited to the lungs following *S. pneumoniae* infection

To better understand if adenosine receptor signaling can be targeted to reverse the age

associated dysregulation in PMN pulmonary influx following pneumococcal infection, aged mice

were treated with specific adenosine receptor agonists to stimulate receptor signaling and then

infected intratracheally with 2.5×10^4 CFU *S. pneumoniae* TIGR4. We first tested A2A receptor as

it had a role in controlling survival of young hosts later in infection. We found that activating A2A

in aged mice at 18 hpi had no effect on the total number of PMNs in the lungs (**Sup. Fig 6A**), or

the number of bacteria in the lungs (**Sup. Fig 6B**), or blood (**Sup. Fig 6C**) by 48 hours post

challenge.

We then focused on the role of A1 as it was required for early influx of PMNs in young

hosts. We tested the effect of A1 agonism in aged mice and determined the total number of PMNs

in the lungs at 6 hpi by flow cytometry. We found that agonism of the A1 receptor significantly

increased the total number of PMNs in the lungs of aged mice 6 hpi when compared to uninfected

mice (**Fig 9A**). Additionally, at 6 hpi, a separate set of A1 agonist treated mice had significantly

higher levels of MPO in the lungs compared to vehicle control treated mice (**Fig 9B**). Importantly,

A1 receptor agonism decreases bacterial burden in the lungs of aged mice compared to untreated

controls (**Fig 9C**). Taken together, these data demonstrate that A1 adenosine receptor signaling can

be targeted to reverse the age-related decrease in PMN pulmonary influx and reduce bacterial

burden in the lungs of infected aged mice.

DISCUSSION

Cellular damage resulting from infection is proposed to shape the outcome of host-

pathogen interactions (42-44) and damage-associated molecular patters (DAMPS) can act as

signals that initiate the immune response (45). One such molecule is extracellular adenosine

(EAD). At baseline homeostatic levels, adenosine levels in extracellular microenvironments are

very low, however, upon tissue damage resulting from insults such as infection, EAD levels can

rise more than 10 fold (18). We previously found that EAD production and signaling are crucial

for host resistance against pneumococcal infection (10, 20, 21, 38, 46). We show here that EAD

signaling through the high affinity A1 receptor is required for initiating the recruitment of PMNs

into the lungs following bacterial challenge. This initiation of early PMN recruitment was defective

in aged hosts but could be reversed by activation of A1 receptor signaling, suggesting that with

age there are changes in extracellular adenosine signaling that impair the ability of PMNs to rapidly

respond to pneumococcal infection. One of the hallmarks of aging is altered intercellular

communication, including endocrine and neuronal signaling (7) and the findings here suggest that

damage signaling is also altered with aging and contributes to the age-driven changes in immune

response to infection.

PMNs have a dynamic role during pneumococcal pneumonia with their early presence in

the lungs being protective vs later persistence being detrimental (47). In murine models of

infection, PMN influx to the lungs within the first 12 hpi correlated with control of bacterial burden

and PMN depletion prior to infection resulted in increased bacterial burden in the lung and host

lethality (48, 49). In humans, patients with neutropenia are at an increased risk for pneumonia (13).

These findings indicate that early influx of PMNs to the lungs is required for host protection against

pneumococcal pneumonia. In contrast, PMN persistence in the pulmonary environment promotes

disease (10, 15, 17). Depletion of PMNs 18 hours after pneumococcal lung infection resulted in

reduced bacterial numbers and reduced lethality (10). These findings demonstrate that the ability

of PMNs to kill pneumococci is altered during infection (10). Here we found that with aging there

is a delay in PMN influx early within the first 6 hours of infection accompanied by persistent

presence in higher levels later in infection compared to young mice infected with the same dose of

S. pneumoniae. This delay left aged mice more susceptible to disease as it was accompanied by

increased bacterial burden in the lungs throughout the course of infection and higher incidence of

bacteremia compared to young mice. Therefore, with age there is a delay in the protective early

PMN recruitment to the site of infection. The majority of studies examining PMN influx in

response to infection in aged hosts have focused on later timepoints (9, 10, 17, 50, 51). In humans,

elderly pneumococcal pneumonia patients had higher percentages of neutrophils in terminal lung

biopsies compared to younger patients (16) and in several animal models of bacterial and viral

pneumonia, PMN numbers in the lungs of aged mice were higher than in young controls at later

timepoints (9, 10, 17, 50, 51), similar to what we found here. The dynamics of earlier influx and

how that is altered with aging have not been as thoroughly addressed in the literature and there are

no data from human pneumococcal pneumonia patients. Our findings suggest that the kinetics of

PMN influx to the lungs in response to pneumococcal pneumonia is altered with age.

Following lung challenge, *S. pneumoniae* localizes within the airways and binds to lung

tissues as well (10, 17). For PMNs to reach the site of infection in the lungs, they have to move

from the circulation across the endothelium, enter the interstitial space, then migrate across the

lung epithelium into the airways (11, 52). We found here that PMN migration across pulmonary

endothelial cells in impaired with aging. Prior work found that PMNs from elderly human donors

did not show defects in transepithelial migration in response to pneumococcal infection (53) but

rather that chemotaxis (or directional movement) of PMNs is diminished with aging (51, 54-58).

PMNs isolated from both healthy controls and pneumonia patients above 60 years of age

displayed diminished chemotaxis compared with younger individuals (54, 55) which was linked

to over activation of phosphoinositide-3-kinase (PI3K) signaling (54). PI3K is controlled

downstream of GPCRs including adenosine receptors (54, 59). Here we found that A1 receptor

signaling on PMNs is required for their migration across pulmonary endothelial cells in response

to infection and that activating A1 receptor reverses the defect in early PMN pulmonary influx

observed in aged hosts. These findings are supported by prior work showing that A1 receptor

signaling on PMNs is required for their adherence to the endothelium in response to N-

Formylmethionyl-leucyl-phenylalanine (fMLP) or Phorbol 12-myristate 13-acetate (PMA)

stimulation (60, 61). In this study we found that A1 receptor signaling was needed for the ability

of PMNs to upregulate the expression of the chemokine receptor CXCR2 early on in response to

infection. During *S. pneumoniae* infection, CXCR2 is required for PMN influx to the lungs (39)

and in aged mice, expression of CXCR2 on PMNs was reported to be lower than young hosts

(56). Therefore, it is possible that activating A1 receptor signaling in aged hosts boosts PMN

migration by upregulating expression of CXCR2.

The role of A1 receptor signaling in PMN recruitment to the lungs seem to play opposing

roles during pathogen driven vs sterile injury. In LPS or ischemia reperfusion lung injury, A1

receptor signaling attenuated PMN trafficking to the lungs (62-65). In contrast and similar to what

we observed here, A1 receptor signaling was required for PMN pulmonary influx during infection

with H1N1 influenza virus (66). This highlights that pathogens can alter the dynamics of the

immune response and that the role of the adenosine signaling may differ in sterile vs. infection

driven injury.

Extracellular adenosine is known to regulate PMN function (18) and play an important role

in host resistance against pulmonary infections (67). Aging is associated with aberrant PMN

responses and overall increased susceptibility to pulmonary infections (47) . However,

surprisingly, the role of EAD in immunosenescence remains largely unexplored. In prior work we

found that there are age driven changes in the expression of the enzymes that make and break EAD

on PMNs and that EAD signaling on PMNs could be targeted to reverse the decline in their

antibacterial activity (33). In this study, to examine the role extracellular adenosine signaling in

the age-related delay in PMN influx we tracked how this pathway changes with host aging. We

found that circulating levels of adenosine were no different in young and aged mice, in fact aged

mice had significantly more adenosine in circulation at 48 hpi. While this increase in adenosine

later on in infection may reflect the increased damage that is occurring in these mice (due to higher

bacterial burden), these data also indicate that it is not that adenosine is not being produced with

age, but that EAD signaling in PMNs of aged mice may be dysregulated. This is in line with prior

work showing that aging is associated with defects in overall GPCR signaling and that alterations

in EAD receptor signaling contribute to the age-associated decline of brain and heart organ

functions (25) (68) (23). When looking at adenosine receptor expression on the surface of PMNs

from aged mice following pneumococcal infection, there was a decrease in receptor expression

from baseline in both you and aged mice. However, in PMNs isolated from young mice these

receptor levels rebounded by 48hpi, while receptor expression on PMNs from aged mice did not

recover. Following activation by an agonist, GPCR signaling is desensitized by internalization of

the receptor into the cell, this is a result of interaction of the G protein with β-arrestins to stop

signaling and additional scaffold proteins which guide trafficking the internalized GPCR (69-71).

The decrease in adenosine receptor expression following pneumococcal infection may be due to

this process of GPCR desensitization. Following desensitization, GPCRs may be degraded but

may also recycle back to the cell membrane surface for resensitization, where the receptor can be

activated again (70). In PMNs isolated from aged mice adenosine receptor expression does not

rebound following the initial drop post infection. This would indicate that EAD receptors are not

being recycled back to the cell surface, and throughout infection, PMNs of aged mice may be less

responsive to activation by adenosine. This could be in part be driven by the consistently elevated

levels of EAD later in infection observed in old but not young hosts. Additionally, the differences

in receptor levels between young and aged mice later in infection could indicate that there are

changes in GPCR desensitization and re-sensitization in PMNs with host age.

A balanced PMN response where PMNs are recruited to the lungs and resolve at the proper

time is essential for effective clearance of pneumococcus and efficient host protection (12, 13, 15,

16). In prior work, we found that in young mice, EAD production by CD73 was needed for

regulation of PMN influx later on during infection, where by day 2 following pulmonary challenge,

CD73$^{-/-}$ mice had significantly more PMNs in the lungs compared to wildtype controls (10). We

hypothesized that extracellular adenosine signaling via the low affinity receptors A2A and A2B

receptors regulate PMN resolution from the lungs later in infection when adenosine levels are

higher due to increased inflammation and host damage. It has been shown previously that under

hypoxia conditions adenosine produced by the endothelium reduced PMN accumulation by

activating A2A and A2B receptors on the PMN surface (72). Surprisingly, we found here that

young mice lacking A2A or A2B receptors showed no changes in PMN presence in the lungs

thorough the course of pneumococcal infection. As both A2A or A2B receptors are expressed on

PMNs and are coupled to Gs (18), it is possible that they play a redundant in PMN movement and

that removal of both would be needed to observe a significant effect. An alternative explanation is

that the levels of adenosine receptors do decline on PMNs through infection even in young hosts,

suggesting that PMNs may become less responsive to adenosine signaling with time.

In summary, we have shown that A1 receptor signaling is required for early PMN influx to

the lungs following pneumococcal infection. Importantly, activation of the A1 receptor using A1

receptor agonist, 2-Chloro-N6-cyclopentyladenosine increased PMN influx to the lung in aged

mice, which resulted in lower bacterial burden in the lung. This enhanced ability to control

bacterial infection could be a combination of increase PMN influx as shown here and improved

function as previously reported (33). These findings suggest that extracellular adenosine signaling

may be dysregulated with host aging and that pharmacologically targeting this pathway can

enhance the immune response in aged mice leading to more effective clearance of pneumococcus.

Identifying a potential target to enhance the immune response to pneumococcal pneumonia in aged

hosts is important as this disease persists in older adults despite the availability of antibiotics and

vaccines (3, 5, 6). In addition to being a potential pharmacological target, the extracellular

adenosine pathway is an important pathway to understand in the context of host-pathogen

interaction as it could be a marker of host damage. This study adds dysregulation of the EAD

pathway with age under the aging hallmark of altered intercellular signaling, which in turn controls

immunosenescence.

DECLARATIONS

Ethics approval and consent to participate

All animal experiments were conducted in accordance with the Institutional Animal Care and Use

Committee guidelines under the approved protocol number MIC33018Y

Consent for publication

Not applicable

Availability of data and materials

All data generated or analyzed during this study are included in this published article [and its

supplementary information files].

Competing interests

The authors declare no conflict of interest.

Funding

This work was supported by National Institute of Health grants R00AG051784 and R01

AG068568-01A1 to ENBG and F31 AI169889-01A1 to SRS.

Authors' contributions

SRS designed research, conducted research, analyzed data, and wrote paper. SES designed

research, conducted research, and analyzed data. EYIT, APL and MB conducted research and

analyzed data. ENBG designed research, wrote the paper, and had responsibility for final content.

All authors read and approved the final manuscript.

Acknowledgements

We would like to thank Dr. Michael Battaglia for critical feedback on the manuscript.

REFERENCES:

1.   WHO. Pneumococcal disease. World Health Organization. 2014.

2.   Grudzinska FS, Brodlie M, Scholefield BR, Jackson T, Scott A, Thickett DR, et al.

Neutrophils in community-acquired pneumonia: parallels in dysfunction at the extremes of age.

Thorax. 2020;75(2):164-71.

3.   Simell B, Vuorela A, Ekstrom N, Palmu A, Reunanen A, Meri S, et al. Aging reduces the

functionality of anti-pneumococcal antibodies and the killing of Streptococcus pneumoniae by

neutrophil phagocytosis. Vaccine. 2011;29(10):1929-34.

4.   Gonçalves MT, Mitchell TJ, Lord JM. Immune ageing and susceptibility to Streptococcus

pneumoniae. Biogerontology. 2016;17(3):449-65.

5.   CDC. Pneumococcal Disease Global Pneumococcal Disease and Vaccine CDC: Centers

for Disease Control and Prevention; 2018 [updated 2018, November 15. Available from:

https://www.cdc.gov/pneumococcal/global.html.

6.   Matanock A LG, Gierke R, Kobayashi M, Leidner A, Pilishvili T. Use of 13-Valent

Pneumococcal Conjugate Vaccine and 23-Valent Pneumococcal Polysaccharide Vaccine Among

Adults Aged ≥65 Years: Updated Recommendations of the Advisory Committee on Immunization

Practices. MMWR Morb Mortal Wkly Rep 2019;68:1069–1075. . CDC; 2019.

7.   Lopez-Otin C, Blasco MA, Partridge L, Serrano M, Kroemer G. The hallmarks of aging.

Cell. 2013;153(6):1194-217.

8.   López-Otín C, Blasco MA, Partridge L, Serrano M, Kroemer G. Hallmarks of aging: An

expanding universe. Cell. 2023;186(2):243-78.

9.   Boyd AR, Shivshankar P, Jiang S, Berton MT, Orihuela CJ. Age-related defects in TLR2

signaling diminish the cytokine response by alveolar macrophages during murine pneumococcal

pneumonia. Exp Gerontol. 2012;47(7):507-18.

10.   Bou Ghanem EN, Clark S, Roggensack SE, McIver SR, Alcaide P, Haydon PG, et al.

Extracellular Adenosine Protects against Streptococcus pneumoniae Lung Infection by Regulating

Pulmonary Neutrophil Recruitment. PLoS Pathog. 2015;11(8):e1005126.

11.   Palmer CS, Kimmey JM. Neutrophil Recruitment in Pneumococcal Pneumonia. Frontiers

in Cellular and Infection Microbiology. 2022;12.

12.   Pechous RD. With Friends Like These: The Complex Role of Neutrophils in the

Progression of Severe Pneumonia. Front Cell Infect Microbiol. 2017;7:160.

13.   Rolston KV. The spectrum of pulmonary infections in cancer patients. Curr Opin Oncol.

2001;13(4):218-23.

14.   Kolaczkowska E, Kubes P. Neutrophil recruitment and function in health and

inflammation. Nat Rev Immunol. 2013;13(3):159-75.

15.   Bhowmick R, Tin Maung NH, Hurley BP, Ghanem EB, Gronert K, McCormick BA, et al.

Systemic disease during Streptococcus pneumoniae acute lung infection requires 12-lipoxygenase-

dependent inflammation. J Immunol. 2013;191(10):5115-23.

16.   Menter T, Giefing-Kroell C, Grubeck-Loebenstein B, Tzankov A. Characterization of the

inflammatory infiltrate in Streptococcus pneumoniae pneumonia in young and elderly patients.

Pathobiology. 2014;81(3):160-7.

17.   Bou Ghanem EN, Clark S, Du X, Wu D, Camilli A, Leong JM, et al. The alpha-tocopherol

form of vitamin E reverses age-associated susceptibility to streptococcus pneumoniae lung

infection by modulating pulmonary neutrophil recruitment. J Immunol. 2015;194(3):1090-9.

18.   Barletta KE, Ley K, Mehrad B. Regulation of neutrophil function by adenosine.

Arterioscler Thromb Vasc Biol. 2012;32(4):856-64.

19.   Hasko G, Linden J, Cronstein B, Pacher P. Adenosine receptors: therapeutic aspects for

inflammatory and immune diseases. Nat Rev Drug Discov. 2008;7(9):759-70.

20.   Siwapornchai N, Lee JN, Tchalla EYI, Bhalla M, Yeoh JH, Roggensack SE, et al.

Extracellular adenosine enhances the ability of PMNs to kill Streptococcus pneumoniae by

inhibiting IL-10 production. J Leukoc Biol. 2020.

21.   Bhalla M, Simmons SR, Abamonte A, Herring SE, Roggensack SE, Bou Ghanem EN.

Extracellular adenosine signaling reverses the age-driven decline in the ability of neutrophils to

kill Streptococcus pneumoniae. Aging Cell. 2020:e13218.

22.   Chen JF, Chern Y. Impacts of methylxanthines and adenosine receptors on

neurodegeneration: human and experimental studies. Handb Exp Pharmacol. 2011(200):267-310.

23.   Willems L, Ashton KJ, Headrick JP. Adenosine-mediated cardioprotection in the aging

myocardium. Cardiovasc Res. 2005;66(2):245-55.

24.   Headrick JP. Impact of aging on adenosine levels, A1/A2 responses, arrhythmogenesis, and

energy metabolism in rat heart. Am J Physiol. 1996;270(3 Pt 2):H897-906.

25.   Mackiewicz M, Nikonova EV, Zimmermann JE, Romer MA, Cater J, Galante RJ, et al.

Age-related changes in adenosine metabolic enzymes in sleep/wake regulatory areas of the brain.

Neurobiol Aging. 2006;27(2):351-60.

26.   Crosti F, Ciboddo GF, Barbieri MC, Inversi F, Pavoni D, Quarenghi S, et al. Evidence for

adenosine adenosinedeaminase lymphocyte system impairment in ageing. Boll Ist Sieroter Milan.

1987;66(4):282-8.

27.   Yeo EJ, Park SC. Age-dependent agonist-specific dysregulation of membrane-mediated

signal transduction: emergence of the gate theory of aging. Mech Ageing Dev. 2002;123(12):1563-

78.

28.   Bhalla M, Hui Yeoh J, Lamneck C, Herring SE, Tchalla EYI, Heinzinger LR, et al. A1

adenosine receptor signaling reduces Streptococcus pneumoniae adherence to pulmonary

epithelial cells by targeting expression of platelet-activating factor receptor. Cell Microbiol.

2019:e13141.

29.   Bhalla M, Heinzinger LR, Morenikeji OB, Marzullo B, Thomas BN, Bou Ghanem EN.

Transcriptome profiling reveals CD73 and age-driven changes in neutrophil responses against

Streptococcus pneumoniae. bioRxiv. 2021:2021.04.14.439887.

30.   Gutierrez F, Masia M, Mirete C, Soldan B, Rodriguez JC, Padilla S, et al. The influence of

age and gender on the population-based incidence of community-acquired pneumonia caused by

different microbial pathogens. J Infect. 2006;53(3):166-74.

31.   Swamydas M, Lionakis MS. Isolation, purification and labeling of mouse bone marrow

neutrophils for functional studies and adoptive transfer experiments. J Vis Exp. 2013(77):e50586.

32.   Tchalla EYI, Bhalla M, Wohlfert EA, Bou Ghanem EN. Neutrophils are required during

immunization with the pneumococcal conjugate vaccine for protective antibody responses and

host defense against infection. J Infect Dis. 2020.

33.   Bhalla M, Simmons SR, Abamonte A, Herring SE, Roggensack SE, Bou Ghanem EN.

Extracellular adenosine signaling reverses the age-driven decline in the ability of neutrophils to

kill Streptococcus pneumoniae. Aging Cell. 2020;19(10):e13218.

34.   Zhou Y, Tong L, Chu X, Deng F, Tang J, Tang Y, et al. The Adenosine A1 Receptor

Antagonist DPCPX Inhibits Tumor Progression via the ERK/JNK Pathway in Renal Cell

Carcinoma. Cell Physiol Biochem. 2017;43(2):733-42.

35.   Zhang X, Xia J, Qian D, Wang Y, Lin Y, Huang X, et al. An adenosine A1 agonist 2-chloro-

N6 cyclopentyladenosine inhibits the angiotensin II-induced cardiomyocyte hypertrophy through

the calcineurin pathway. Cardiology. 2014;129(3):153-62.

36.   Yao L, McFarland K, Fan P, Jiang Z, Ueda T, Diamond I. Adenosine A2a blockade prevents

synergy between mu-opiate and cannabinoid CB1 receptors and eliminates heroin-seeking

behavior in addicted rats. Proc Natl Acad Sci U S A. 2006;103(20):7877-82.

37.   Phillis JW. The selective adenosine A2 receptor agonist, CGS 21680, is a potent depressant

of cerebral cortical neuronal activity. Brain Res. 1990;509(2):328-30.

38.   Herring SE, Mao S, Bhalla M, Tchalla EYI, Kramer JM, Bou Ghanem EN. Mitochondrial

ROS production by neutrophils is required for host antimicrobial function against Streptococcus

pneumoniae and is controlled by A2B adenosine receptor signaling. PLoS Pathog.

2022;18(11):e1010700.

39.   Herbold W, Maus R, Hahn I, Ding N, Srivastava M, Christman JW, et al. Importance of

CXC chemokine receptor 2 in alveolar neutrophil and exudate macrophage recruitment in response

to pneumococcal lung infection. Infect Immun. 2010;78(6):2620-30.

40.   Reutershan J, Morris MA, Burcin TL, Smith DF, Chang D, Saprito MS, et al. Critical role

of endothelial CXCR2 in LPS-induced neutrophil migration into the lung. J Clin Invest.

2006;116(3):695-702.

41.   Williams MR, Azcutia V, Newton G, Alcaide P, Luscinskas FW. Emerging mechanisms of

neutrophil recruitment across endothelium. Trends Immunol. 2011;32(10):461-9.

42.   Casadevall A, Pirofski LA. The damage-response framework of microbial pathogenesis.

Nat Rev Microbiol. 2003;1(1):17-24.

43.   Pirofski LA, Casadevall A. The damage-response framework of microbial pathogenesis

and infectious diseases. Adv Exp Med Biol. 2008;635:135-46.

44.   Pirofski LA, Casadevall A. Pathogenesis of COVID-19 from the Perspective of the

Damage-Response Framework. mBio. 2020;11(4).

45.   Burgoyne RA, Fisher AJ, Borthwick LA. The Role of Epithelial Damage in the Pulmonary

Immune Response. Cells. 2021;10(10).

46.   Bhalla M, Hui Yeoh J, Lamneck C, Herring SE, Tchalla EYI, Heinzinger LR, et al. A1

adenosine receptor signaling reduces Streptococcus pneumoniae adherence to pulmonary

epithelial cells by targeting expression of platelet-activating factor receptor. Cell Microbiol.

2020;22(2):e13141.

47.   Simmons SR, Bhalla M, Herring SE, Tchalla EYI, Bou Ghanem EN. Older but Not Wiser:

the Age-Driven Changes in Neutrophil Responses during Pulmonary Infections. Infect Immun.

2021;89(4).

48.   Garvy BA, Harmsen AG. The importance of neutrophils in resistance to pneumococcal

pneumonia in adult and neonatal mice. Inflammation. 1996;20(5):499-512.

49.   Hahn I, Klaus A, Janze AK, Steinwede K, Ding N, Bohling J, et al. Cathepsin G and

neutrophil elastase play critical and nonredundant roles in lung-protective immunity against

Streptococcus pneumoniae in mice. Infect Immun. 2011;79(12):4893-901.

50.   Kulkarni U, Zemans RL, Smith CA, Wood SC, Deng JC, Goldstein DR. Excessive

neutrophil levels in the lung underlie the age-associated increase in influenza mortality. Mucosal

Immunol. 2019;12(2):545-54.

51.   Toapanta FR, Ross TM. Impaired immune responses in the lungs of aged mice following

influenza infection. Respir Res. 2009;10:112.

52.   Craig A, Mai J, Cai S, Jeyaseelan S. Neutrophil recruitment to the lungs during bacterial

pneumonia. Infect Immun. 2009;77(2):568-75.

53.   Bou Ghanem EN, Lee JN, Joma BH, Meydani SN, Leong JM, Panda A. The Alpha-

Tocopherol Form of Vitamin E Boosts Elastase Activity of Human PMNs and Their Ability to Kill

Streptococcus pneumoniae. Front Cell Infect Microbiol. 2017;7:161.

54.   Sapey E, Greenwood H, Walton G, Mann E, Love A, Aaronson N, et al. Phosphoinositide

3-kinase inhibition restores neutrophil accuracy in the elderly: toward targeted treatments for

immunosenescence. Blood. 2014;123(2):239-48.

55.   Sapey E, Patel JM, Greenwood HL, Walton GM, Hazeldine J, Sadhra C, et al. Pulmonary

Infections in the Elderly Lead to Impaired Neutrophil Targeting, Which Is Improved by

Simvastatin. Am J Respir Crit Care Med. 2017;196(10):1325-36.

56.   Nomellini V, Brubaker AL, Mahbub S, Palmer JL, Gomez CR, Kovacs EJ. Dysregulation

of neutrophil CXCR2 and pulmonary endothelial icam-1 promotes age-related pulmonary

inflammation. Aging Dis. 2012;3(3):234-47.

57.   Ren Z, Gay R, Thomas A, Pae M, Wu D, Logsdon L, et al. Effect of age on susceptibility

to Salmonella Typhimurium infection in C57BL/6 mice. J Med Microbiol. 2009;58(Pt 12):1559-

67.

58.   Van Avondt K, Strecker J-K, Tulotta C, Minnerup J, Schulz C, Soehnlein O. Neutrophils

in aging and aging-related pathologies. Immunological Reviews. 2023;314(1):357-75.

59.   Shi L, Wu Z, Miao J, Du S, Ai S, Xu E, et al. Adenosine interaction with adenosine receptor

A2a promotes gastric cancer metastasis by enhancing PI3K-AKT-mTOR signaling. Mol Biol Cell.

2019;30(19):2527-34.

60.   Cronstein BN, Levin RI, Philips M, Hirschhorn R, Abramson SB, Weissmann G.

Neutrophil adherence to endothelium is enhanced via adenosine A1 receptors and inhibited via

adenosine A2 receptors. J Immunol. 1992;148(7):2201-6.

61.   Felsch A, Stocker K, Borchard U. Phorbol ester-stimulated adherence of neutrophils to

endothelial cells is reduced by adenosine A2 receptor agonists. J Immunol. 1995;155(1):333-8.

62.   Sun CX, Young HW, Molina JG, Volmer JB, Schnermann J, Blackburn MR. A protective

role for the A1 adenosine receptor in adenosine-dependent pulmonary injury. J Clin Invest.

2005;115(1):35-43.

63.   Fernandez LG, Sharma AK, LaPar DJ, Kron IL, Laubach VE. Adenosine A1 receptor

activation attenuates lung ischemia-reperfusion injury. J Thorac Cardiovasc Surg.

2013;145(6):1654-9.

64.   Ngamsri KC, Wagner R, Vollmer I, Stark S, Reutershan J. Adenosine receptor A1 regulates

polymorphonuclear cell trafficking and microvascular permeability in lipopolysaccharide-induced

lung injury. J Immunol. 2010;185(7):4374-84.

65.   Gazoni LM, Walters DM, Unger EB, Linden J, Kron IL, Laubach VE. Activation of A1,

A2A, or A3 adenosine receptors attenuates lung ischemia-reperfusion injury. J Thorac Cardiovasc

Surg. 2010;140(2):440-6.

66.   Aeffner F, Woods PS, Davis IC. Activation of A1-adenosine receptors promotes leukocyte

recruitment to the lung and attenuates acute lung injury in mice infected with influenza A/WSN/33

(H1N1) virus. J Virol. 2014;88(17):10214-27.

67. Lee JS, Yilmaz O. Unfolding Role of a Danger Molecule Adenosine Signaling in

Modulation of Microbial Infection and Host Cell Response. Int J Mol Sci. 2018;19(1).

68. Rolband GC, Furth ED, Staddon JM, Rogus EM, Goldberg AP. Effects of age and

adenosine in the modulation of insulin action on rat adipocyte metabolism. J Gerontol.

1990;45(5):B174-8.

69. Waldo GL, Northup JK, Perkins JP, Harden TK. Characterization of an altered membrane

form of the beta-adrenergic receptor produced during agonist-induced desensitization. J Biol

Chem. 1983;258(22):13900-8.

70. Yuan L, Barbash S, Kongsamut S, Eishingdrelo A, Sakmar TP, Eishingdrelo H. 14-3-3

signal adaptor and scaffold proteins mediate GPCR trafficking. Scientific Reports.

2019;9(1):11156.

71. Shenoy SK, Lefkowitz RJ. β-Arrestin-mediated receptor trafficking and signal

transduction. Trends Pharmacol Sci. 2011;32(9):521-33.

72. Eltzschig HK, Thompson LF, Karhausen J, Cotta RJ, Ibla JC, Robson SC, et al.

Endogenous adenosine produced during hypoxia attenuates neutrophil accumulation: coordination

by extracellular nucleotide metabolism. Blood. 2004;104(13):3986-92.

FIGURES & LEGENDS

Figure 1: Old mice have delayed early pulmonary PMN influx. Young and old male C57BL/6 mice were infected i.t. with $5x10^4$ CFU *S. pneumoniae* TIGR4. (A) Lungs were harvested, digested, and stained for PMNs and analyzed by flow cytometry at the indicated timepoints following infection. (B) Lung and blood (C) were collected, and samples plated on blood agar and CFU was enumerated at the indicated timepoint post infection. Data are pooled from 6 separate experiments with a total of n=7 mice at 48-hour time point and n=12 mice at all other time points. Asterix indicates significant difference between indicated groups and as calculated by (A) Kruskal Wallis followed by Dunn's multiple comparison test and (B and C) one-way ANOVA followed by Sidak's multiple comparison test. ns indicates non-significant.

Figure 2: PMNs from old mice have a defect in trans-endothelial migration. PMNs were

isolated from the bone marrow of young and old male C57BL/6 mice. Transwells were seeded

with murine primary lung endothelial cells and the bottom of the transwell infected with *S.*

pneumoniae TIGR4 at an MOI of 40. PMNs were placed in the top of the transwell and incubated

at 37°C at 5% CO2 for 1.5 hours. PMNs migrated to the bottom of each transwell were enumerated.

Technical replicates are pooled from 4 separate experiments (n=4 mice). Asterix indicates

significance as determined by unpaired Student's t-test.

Figure 3: Adenosine levels in circulation change with age over time Young and old male

C57BL/6 mice were infected i.t. with 5x10^4 CFU *S. pneumoniae* TIGR4 and sera was collected at

6, 18, and 48 hpi. Adenosine levels in the circulation were measured from two separate

experiments with n=3 mice at each individual time point. Asterix indicates significantly different

from young at the indicated timepoint as calculated by Kruskal Wallis followed by Dunn's multiple

comparison test.

Figure 4: EAD pathway expression on pulmonary PMNs. Young and old male C57BL/6 mice

were infected i.t. with 5×10^4 CFU *S. pneumoniae* TIGR4. The lungs were harvested, and lung

digests were stained for PMNs and each adenosine receptor. Percent expression at baseline (A)

and MFI over time (B) of the indicated adenosine receptor on PMNs was determined. Replicates

are pooled from 6 separate experiments with n=6-12 mice per group. (B) # indicate significance

from uninfected baseline as determined by Kruskal Wallis followed by Dunn's Multiple

comparison test. (C) Asterix indicate significant difference from one as determined by a 1-sample

t-test.

Figure 5: EAD pathway expression on circulating PMNs. Young and old male C57BL/6 mice

were infected i.t. with 5×10^4 CFU *S. pneumoniae* TIGR4. Blood was collected and stained for

PMNs and adenosine receptors. Percent expression at baseline (A) and MFI over time (B) was

determined using flow cytometry. Replicates are pooled from 4 separate experiments with n=6-9

mice per group. Asterix indicate significant difference as determined by unpaired students t-test

(A) and significantly different from one as determined by 1-sample t-test (C). (B) # / p-values

indicate significance from uninfected baseline as determined by Kruskal Wallis followed by

Dunn's Multiple comparison test.

Figure 6: A2B receptor does not affect PMN pulmonary numbers *in vivo*. WT and A2BR[-/-]

mice were infected i.t with 5×10^4 CFU *S. pneumoniae* TIGR4. At the indicated timepoints

following bacterial challenge, lungs were harvested, digested, and stained for PMNs and analyzed

by flow cytometry. Data are pooled from 4 separate experiments with n=6-8 mice per group.

Figure 7: A2A receptor does not affect PMN pulmonary numbers or host resistance to infection. (A) WT and A2AR[-/-] mice were infected i.t with 5x10^4 CFU *S. pneumoniae* TIGR4. At the indicated timepoints following bacterial challenge, lungs were harvested, digested, and stained for PMNs and analyzed by flow cytometry. Data are pooled from 3 separate experiments with n=4-6 mice per group. Lungs (B) and Blood (C) were harvested at indicated timepoints and plated on blood agar plates and CFU was enumerated. Data were pooled from 2 separate experiments with (C) n=11-12 mice per group and (D) n=6 mice per group. (E) WT and A2AR[-/-] mice were infected as in (A) and monitored for survival. Data are pooled from 2 experiments with n=11-12 mice per group. No significant differences were found in survival as measured by Log-Rank (Mantle Cox) test.

Figure 8: A1 receptor signaling is required for PMN pulmonary influx. (A) Young, male mice

were treated i.p with A1 inhibitor DPCPX and infected i.t with $5x10^4$ CFU *S. pneumoniae* TIGR4.

At the indicated time points, lungs were harvested, digested, and stained for PMNs (CD45+,

Ly6G+ CD11b+) (A) and analyzed by flow cytometry. (B) Lungs of a separate set of treated and

infected mice were harvested 6 hpi, homogenized and MPO levels were determined by ELISA.

(C-D) Blood was harvested from infected and A1 treated mice (same mice as A) at the indicated

time points and stained for PMNs and CD18 (C) or CXCR2 (D). (A, C and D) Data are pooled

from 3 separate experiments with n=6-12 mice per group. (A and D) Asterix indicate significance

as determined by Kruskal Wallis followed by Dunn's multiple comparison test. (B) Asterix indicate

significance as determined by unpaired students t-test. (E) Young, male C57/B6 mice were treated

i.p. with A1 inhibitor DPCPX or vehicle control (VC) and PMNs were isolated from the bone

marrow. PMN migration across primary endothelial cells in response to *S. pneumoniae* TIGR4

820　　infection was measured. Technical replicates are pooled from 4 separate experiments. Asterix

821　　indicates significance as determined by Mann-Whitney test.

Figure 9: A1 receptor signaling rescues defect in early PMN pulmonary influx in aged mice.

(A) Old male mice were treated i.p. with A1 agonist 2-Chloro-N6-cyclopentyladenosine or Vehicle

Control and infected i.t. with 5×10^4 CFU *S. pneumoniae* TIGR4 for 6 hours. Lungs were harvested,

digested, and stained for PMNs and analyzed by flow cytometry. Asterix indicates significance as

determined by One- way ANOVA followed by Sidak's multiple comparison test. ns indicates non-

significant. (C) Lung homogenates from infected mice were plated on Blood agar and bacterial

CFU was enumerated. Data are pooled from two separate experiments and with n=3 uninfected

and n=6 infected mice per group. (B) Lungs of a separate set of treated and infected mice were

harvested 6 hpi, homogenized and MPO levels were determined by ELISA. Data are pooled from

n=3 mice. (B and C) Asterix indicates significance as determined by unpaired Student's t-test.