Evolution of plant cell-type-specific *cis*-regulatory elements

Haidong Yan[1,2,+], John P. Mendieta[1,+], Xuan Zhang[1], Alexandre P. Marand[1#], Yan Liang[3], Ziliang Luo[1], Mark A.A. Minow[1], Hosung Jang[1], Xiang Li[1], Thomas Roulé[4], Doris Wagner[4], Xiaoyu Tu[5], Yonghong Wang[3,6], Daiquan Jiang[7], Silin Zhong[7], Linkai Huang[2], Susan R. Wessler[8], and Robert J. Schmitz[1,*]

[1]Department of Genetics, University of Georgia, Athens, GA, 30602, USA

[2]College of Grassland Science and Technology, Sichuan Agricultural University, Chengdu, China

[3]College of Life Sciences, Shandong Agricultural University, Taian, Shandong 271018, China

[4]Department of Biology, University of Pennsylvania, Philadelphia, PA 19104, USA

[5]Joint Center for Single Cell Biology, School of Agriculture and Biology, Shanghai Jiao Tong University, Shanghai 200240, China

[6]Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing 100101, China

[7]State Key Laboratory of Agrobiotechnology, School of Life Sciences, The Chinese University of Hong Kong

[8]Department of Botany and Plant Sciences, University of California, Riverside, CA, 92521, USA

[+]These authors contributed equally.

[#]Current address: Department of Molecular, Cellular, and Developmental Biology, University of Michigan, Ann Arbor, MI, 48109, USA

*Corresponding author

Abstract:

Cis-regulatory elements (CREs) are critical in regulating gene expression, and yet understanding

of CRE evolution remains challenging. Here, we constructed a comprehensive single-cell atlas of

chromatin accessibility in *Oryza sativa*, integrating data from 103,911 nuclei representing 126

discrete cell states across nine distinct organs. We used comparative genomics to compare cell-

type resolved chromatin accessibility between *O. sativa* and 57,552 nuclei from four additional

grass species (*Zea mays, Sorghum bicolor, Panicum miliaceum*, and *Urochloa fusca*). Accessible

chromatin regions (ACRs) had different levels of conservation depending on the degree of cell-

type specificity. We found a complex relationship between ACRs with conserved noncoding

sequences, cell-type specificity, conservation, and tissue-specific switching. Additionally, we

found that epidermal ACRs were less conserved compared to other cell types, potentially

indicating that more rapid regulatory evolution has occurred in the L1-derived epidermal layer of

these species. Finally, we identified and characterized a conserved subset of ACRs that

overlapped the repressive histone modification H3K27me3, implicating them as potentially

silencer-like CREs maintained by evolution. Collectively, this comparative genomics approach

highlights the dynamics of plant cell-type-specific CRE evolution.

Main

    Cis-regulatory elements (CREs) function as pivotal hubs, facilitating the binding of transcription factors (TFs) and recruitment of chromatin-modifying enzymes, thereby fine-tuning gene expression in a spatiotemporal-specific manner[1]. CREs play important roles in developmental and environmental processes, and their functional divergence frequently drives evolutionary change[2,3]. Prior studies highlighted the dynamic nature of CREs throughout evolution and their involvement in regulating gene expression via distinct chromatin pathways[4-9]. Across diverse cell types, gene expression is intricately regulated by multiple distinct CREs, each exerting control within specific cell, tissue type, particular developmental stage, or environmental cue[10-12]. In plants, environmental sensing and adaptation relies heavily upon epidermal cells[13]. For example, grass epidermal bulliform cells change their turgor pressure to roll the leaf to slow water loss under stressful conditions, with the TF, ZINC FINGER HOMEODOMAIN 1 (ZHD1), modulating leaf rolling by influencing rice (*Oryza sativa*) bulliform cell development[14,15]. Several studies have identified CREs functioning in a cell-type-specific manner within diverse plant species[16-23]. Despite these findings, our understanding of CREs exhibiting evolutionarily conserved or divergent cell-type-specific activities remains limited.

    Through single-cell assay for transposase accessible chromatin sequencing (scATAC-seq), we constructed an expansive single-cell reference atlas (103,911 nuclei) of accessible chromatin regions (ACRs) within rice. We then leveraged these data in tandem with four additional scATAC-seq leaf datasets from diverse grasses [*Zea mays* (16,060 nuclei), *Sorghum bicolor* (15,301 nuclei), *Panicum miliaceum* (7,081 nuclei), and *Urochloa fusca* (19,110 nuclei)][24] allowing us to compare ACRs across species and cell types. We quantified the proportion of ACRs that were conserved in these monocots and found high rates of cell-type-specific ACR turnover, particularly in epidermal cells. This indicates that the ACRs associated with specific cell types are rapidly evolving. Finally, we used both conserved non-coding sequences (CNS) and H3K27me3 to find a series of conserved ACRs and the candidate CREs within them that are potentially important for recruitment of Polycomb-mediated gene silencing.

Construction of an ACR atlas at single-cell resolution in rice

To create a cell-type-resolved ACR rice atlas, we conducted scATAC-seq across a spectrum

of nine organs in duplicate (Fig. 1a and b). Data quality metrics, such as correlation between

biological replicates, transcription start site enrichment, fraction of reads in peaks, fragment size

distribution, and organelle content, revealed excellent data quality (Supplementary Fig. 1 and 2).

Following strict quality control filtering, we identified 103,911 high-quality nuclei, with an

average of 41,701 unique Tn5 integrations per nucleus. Based on a nine-step annotation strategy,

which included RNA *in situ* and spatial-omic (slide-seq) validation of cell-type specificity, we

identified a total of 126 cell states, encompassing 59 main cell types across various

developmental stages from all the organs sampled (Fig. 1b; Extended Data Fig. 1a;

Supplementary Note 1; Supplementary Fig. 3-13; Supplementary Tables 1-7).

By analyzing cell-type-aggregated chromatin accessibility profiles, we identified a total of

120,048 ACRs (Extended Data Fig. 1b and c). Among these ACRs, 30,796 were categorized as

'cell-type-specific ACRs', exhibiting cell-type-specific entropy signals of accessible chromatin

in less than 5% (3/59) of the main cell types, whereas approximately 89,252 were classified as

'broad ACRs' with chromatin accessibility in more than 5% of the cell types (Fig. 1c and d;

Extended Data Fig. 1d and e). When analyzing ACR proximity to genomic features in the rice

genome, about half of the ACRs were gene proximal (52%; located within 2 kb of genes; Fig.

1e). These proximal ACRs had higher but less variable chromatin accessibility than genic and

distal ACRs (Extended Data Fig. 1f). In contrast, about 19% of the ACRs overlapped genes,

mostly in introns, and the remaining 29% were categorized as distal (Fig. 1e; situated more than

2 kb away from genes). The greater chromatin accessibility variance in non-proximal ACRs

suggests these regions may act in select cellular contexts. To further investigate the association

of distal ACRs with gene activity, we examined the interactions between distal cell-type-specific

ACRs and genes using leaf bulk Hi-C data[25]. Among the 3,513 distal cell-type-specific ACRs in

leaf tissue, most (81.7%) were embedded within chromatin loops. More than one-third (37.7%)

of these ACRs interacted with promoters of cell-type-specific accessible genes, and 11.2% (392)

had both the ACRs and their interacting genes associated with the same cell type (Extended Data

Fig. 1g; Supplementary Table 8). As bulk Hi-C poorly measures rare cell types, we expect this

number to be a conservative count of the number of cell-type-specific ACRs associated with cell-

type-specific gene activity.

Fig. 1. Identifying cell types and characterizing ACRs in rice using scATAC-seq data. a, Overview of cell types in leaf, root, seed, and panicle organs. QC: Quiescent Center. SBM: Secondary Branch Meristem. PBM: Primary Branch Meristem. SM: Spikelet Meristem. FM: Floral Meristem. **b,** UMAP projection of nuclei, distinguished by assigned cell-type labels in axillary bud, early seedling (7 days after sowing), seedling (14 days after sowing), leaf (V4 stage; four leaves with visible leaf collars), seminal root, crown root, panicle, early seed development (6 DAP; days after pollination), and late seed development (10 DAP). SAM: Shoot Apical Meristem. VAS cells: vasculature-related cells. *VAS cells: vasculature-related cells, which were further distinguished as procambial meristem, developing phloem/xylem, developing phloem/xylem precursor, vascular parenchyma/sclerenchyma, xylem

parenchyma, and companion cell & sieve elements in Supplementary Fig. 6. CC & SE: Companion cell & sieve

elements. CSE: Central Starchy Endosperm. DSE: Dorsal Starchy Endosperm. LSE: Lateral Starchy Endosperm.

SVB: Scutellar Vascular Bundle. **c,** Evaluation of proportions of ACRs that are cell-type specific versus broad. **d,** A

screenshot illustrates the examples of cell-type-specific and broad ACRs. **e,** Accessible chromatin regions (ACRs)

show a bimodal distribution of distance to the nearest gene. The ACRs were categorized into three major groups

based on their locations to the nearest gene: genic ACRs (overlapping a gene), proximal ACRs (located within 2 kb

of genes), and distal ACRs (situated more than 2 kb away from genes).

The atlas uncovers key TFs, their motifs, and ACRs during rice development

To demonstrate the utility of this new resource, we associated the atlas ACRs with a set of

noncoding quantitative trait nucleotides (QTNs; Fig. 2a). Notably, we observed an enrichment of

agriculturally relevant QTNs[26], within ACRs (Fig. 2b), some of which were cell-type-specific

ACRs (Extended Data Fig. 2a and b; Supplementary Table 9). For instance, a QTN was within

an endosperm-specific ACR located at ~1 kb upstream of *GLUTELIN TYPE-A2 PRECURSOR*

(*OsGluA2*) (Fig. 2c), which is associated with increased seed protein content[27]. Exploring the

endosperm epigenome more, we observed an endosperm-specific reduction in cytosine

methylation at endosperm-specific ACRs, including an ACR linked to the DNA demethylase

OsROS1 (Extended Data Fig. 2c)[28]. We found that 5,159 ACRs had lower DNA methylation in

the endosperm compared to early seedling (Fig. 2d), and these ACRs were enriched for several

TF motif families such as MADS box factors, TEOSINTE

BRANCHED1/CYCLOIDEA/PROLIFERATING CELL FACTOR (TCP), Basic Leucine Zipper

(bZIP), and BARLEY B RECOMBINANT/BASIC PENTACYSTEINE (BBR/BPC), compared

to constitutively unmethylated ACRs (Fig. 2e). Therefore, established endosperm DNA

demethylation[29], coincides with endosperm-specific ACRs.

Fig. 2. Characterization of TF motifs essential for specific cell types using rice ACR atlas. a, The ratio of non-CDS QTNs overlapping with ACRs to all non-CDS QTNs. **b,** A bar plot showing a percentage of non-CDS QTNs overlapping with ACRs. The significance test was done by using the binomial test (alternative = 'two.sided'; See Methods: Construction of control sets for enrichment tests). **c,** Analysis of cell-type-aggregate chromatin accessibility across 12 seed cell types and eight non-seed-related cell types, showing signatures of a QTN within an endosperm-specific ACR situated at the promoter region of *OsGluA2*. An endosperm-specific reduction of cytosine methylation was identified over the endosperm-specific ACR. **d,** A total of 5,159 differentially methylated ACRs were identified, showing hypermethylation in early seedling tissue but hypomethylation within endosperm tissue. **e,** Enrichment of TF families based on their motifs within the 5,159 ACRs. The upper plot shows the TF motif count within the TF family indicated on the x-axis of the bottom plot. The *p* value was computed using a hypergeometric test (alternative = 'two.sided'). **f,** Deviations of motifs displaying enrichment in specific cell types (white frames) where their cognate TFs are known to be accumulated. **g,** UMAP visualizations depict the cell progression of root developing xylem in *O. sativa*. **h,** Relative motif deviations for 351 TF motifs (left). Four motifs enriched along the trajectory gradient are shown on the right. **i,** UMAP visualizations depict the cell progression of root developing xylem in *Z. mays*. **j,** The heatmap displays motif deviation scores ordered by pseudotime (x-axis). 'Conserved' indicates motifs that showed consistent patterns of motif deviation score changes along the pseudotime between *O. sativa* and *Z. mays*. 'Shift early' indicates higher motif deviation scores observed at the beginning of pseudotime. **k,**

Comparison of relative motif accessibility of TCP4 along the pseudotime between *O. sativa* and *Z. mays*. **l,**

Expression profiles of the *TCP4* gene using scRNA-seq datasets from *O. sativa* and *Z. mays*.

We further investigate the TF motifs that exhibited enrichment within the ACRs of specific

cell types (Supplementary Table 10). We observed consistency between known TF activity and

cognate motif enrichment (Fig. 2f). For example, ARABIDOPSIS THALIANA HOMEOBOX

PROTEIN 15 (ATHB-15) is associated with xylem differentiation[30] and we found a significant

enrichment of the ATHB-15 motif in seedling developing xylem or xylem precursor cells. Other

TFs, such as HOMEODOMAIN GLABROUS 1 (HDG1)[31], MYELOBLASTOSIS 46

(MYB46)[32], PLETHORA1 (PLT1)[33], INDETERMINATE DOMAIN 4 (IDD4)[34] are known to

accumulate in the epidermis, developing fiber, quiescent center (QC), and endosperm,

respectively, and their motifs were enriched in these cell types (Fig. 2f). Furthermore, the motif

analysis unveiled potential novel roles for certain cell types. For instance, AtTCP4 regulates

lignin and cellulose deposition and binds the promoter of *VASCULAR-RELATED NAC DOMAIN*

7 (VND7), a pivotal xylem development gene[35]. However, in *O. sativa*, the TCP4 motif was QC

enriched more so than in the developing root xylem, alluding to unknown QC roles. In sum, our

TF motif enrichment sheds light on both known and novel regulatory mechanisms underlying

cell differentiation and function.

To examine the ACR dynamics during cell fate progression, we organized nuclei along

pseudotime trajectories representing 14 developmental continuums (Fig. 2g; Extended Data Fig.

2d and e; Supplementary Fig. 14; Supplementary Table 11 and 12), focusing on root developing

xylem (RDX), we identified 16,673 ACRs, 323 of 2,409 TFs, and 351 of 540 TF motifs showing

differential chromatin accessibility along the xylem trajectory (Supplementary Table 13). Early

in the xylem trajectory, the TCP4 motif was notably enriched (Fig. 2h). To determine if TCP4

enrichment is also present in *Z. mays*, we aligned the RDX motifs of both species using a

dynamic time-warping algorithm (Extended Data Fig. 2f), which identified 62 motifs with

species-differential *cis*-regulatory dynamics during RDX developments (Extended Data Fig. 2f;

Supplementary Table 14). OsTCP4 decreased in motif accessibility during xylem development,

whereas ZmTCP4 increased along RDX trajectory (Fig. 2j and k). This mirrors the single-cell

RNA sequencing (scRNA-seq) expression patterns of *TCP4* during RDX development[36] (Fig.

2l). This unveiling of opposing developmental TCP4 motif accessibility gradients in *O. sativa*

and *Z. mays,* exemplifies how our atlas can merge with existing, and future data, to drive

discovery surrounding monocot development.

In sum, the *O. sativa* atlas provides a comprehensive ACR resource, capturing known

agronomic QTNs and bringing novel insights to seed and xylem development. Beyond the

discoveries outlined here, this atlas represents a potent reference for the rice research community

to answer diverse questions about cell type specific processes.

The landscape of cell-type-specific ACRs across grass species

We were interested in leveraging the *O. sativa* atlas to understand how ACRs change during

grass evolution. The *O. sativa* ACRs were overlapped with syntenic regions defined by their

relationship to four different grass species *Z. mays*, *S. bicolor*, *P. miliaceum*, and *U. fusca* that

have single-cell ATAC sequencing using combinatorial indexing (sciATAC-seq) data from

leaves[24]. The analysis revealed that 34% (40,477) of the *O. sativa* ACRs were within 8,199

syntenic regions (~86 Mb of the *O. sativa* genome) shared with at least one of the four examined

grass species (Extended Data Fig. 3a; Supplementary Fig. 15). In contrast, the majority of ACRs

(66%; 79,571) were in non-syntenic regions. Notably, the ACRs found in non-syntenic regions

were significantly enriched for cell-type specificity ACRs (p = 6e-248 to 0.0031; Fisher's exact

test), for both proximal and distal ACRs (Extended Data Fig. 3b and c). This reveals that most of

the *O. sativa* ACRs in the grass species examined occurred in non-syntenic regions.

To determine to what degree ACR number, genomic position and cell-type specificity differs

amongst grasses, we compared the composition and distribution of leaf ACRs across the five

species (Fig. 3a). Using previous cell-type annotations[24], we calculated the proportion of both

broad and cell-type-specific ACRs across all species. We categorized cell-type-specific ACRs as

those only accessible in 1-2 leaf cell-types (Extended Data Fig. 3d; Supplementary Table 15). An

average of ~53,000 ACRs were identified across the five species, with 15-35% of the ACRs

classified as cell-type specific (Fig. 3b; Supplementary Fig. 16 and 17). Broad and cell-type-

specific ACRs were equivalent in their distributions around promoters, distal, and genic regions

(Fig. 3c; Extended Data Fig. 3e-f).

Prior hypotheses suggested that large scale regulatory rewiring could play a key role in cell-

type environmental adaptation[8,37]. To explore instances where divergent TF activity occurred in

the same cell types, we associated TF gene body chromatin accessibility with their cognate TF

motifs across different species and cell types. Approximately 64% to 76% of the TFs (211 to 232) examined exhibited a positive correlation between the local chromatin accessibility of their gene body and global enrichment of their cognate binding motifs within ACRs (motif deviation) across all leaf cell types and all species (Extended Data Fig. 4a). The use of TF chromatin accessibility was supported by an analysis of TF expression and motif deviation in both seedling (Supplementary Fig. 7) and root data (Zhang et al. 2021), which uncovered a similar positive relationship across cells (Extended Data Fig. 4b). These results suggest a positive relationship between TF gene-body chromatin accessibility/expression and TF activity in the same cell type.

Moreover, the genomic sequences from all ACRs discovered in all species and cell types exhibited enrichment of TF motifs compared to a control set of sequences (Extended Data Fig. 4c). Furthermore, TF motif enrichment analysis revealed known TF-cell-type specificities (Fig. 3d). For example, the HDG1 TF is critical for epidermis and cuticle development[31] and its motif was enriched in epidermis cells in all five species (Extended Data Fig. 4d). We also observed motif enrichments of WRKY, HOMEODOMAIN-LEUCINE ZIPPER (HD-ZIP), and PLANT ZINC FINGER (PLINC) in epidermal cells across all five species examined (Fig. 3e; Extended Data Fig. 4e; Supplementary Fig. 18; Supplementary Table 16). This result indicates that these TFs play a conserved role in the development of the grass epidermis. Phloem companion, sieve element cell, and bundle sheath cell TFs exhibited similar enrichments across the species (Fig. 3e). However, species-specific motif patterns were also observed, with *O. sativa* being the most different. For example, the DNA-BINDING ONE ZINC FINGER (DOF) TF family motif exhibited higher enrichment scores in four C_4 photosynthesizing species (*Z. mays*, *S. bicolor*, *P. miliaceum*, and *U. fusca*), as opposed to C_3 photosynthesizing *O. sativa* (Fig. 3e; Extended Data Fig. 4f and g). The DOF TF family is involved in *Arabidopsis thaliana* vasculature development[38], and is important in the transition from C_3 to C_4 photosynthesis[23,39-42]. The enrichment of DOF motifs in *O. sativa* species-specific motifs is, therefore, an expected biological signal, indicating robust species-specific motif detection and high-quality single-cell data.

Taken together, our findings demonstrate the power of scATAC-seq data in a comparative framework to explore regulatory evolution, both based on the relationship of ACRs to TF motifs, as well as the relationship between TFs and their corresponding motifs.

Fig. 3. Position and motif enrichment of cell-type-specific ACRs across species. a, A phylogenetic tree illustrates
five species under examination. **b,** The count of broad and cell-type-specific ACRs. **c,** Broad and cell-type-specific
ACRs were classified into three main groups based on their proximity to the nearest gene: genic ACRs (overlapping
a gene), proximal ACRs (located within 2 kb of genes), and distal ACRs (situated more than 2 kb away from genes).
O. sativa, *P. miliaceum*, and *U. fusca* showed a higher percentage of proximal ACRs, but a lower percentage of
distal ACRs compared to *Z. mays* and *S. bicolor*, likely reflecting differences in intergenic space and overall genome
sizes. **d**, A heatmap illustrates nine TF motif enrichments, consistent with the known TF dynamics among cell types
[CIRCADIAN CLOCK ASSOCIATED 1 (CCA1), LATE ELONGATED HYPOCOTYL (LHY) and
JUNGBRUNNEN1 (JUB1): mesophyll[43]; CYCLING DOF FACTOR 3 (CDF3) and CDF5: companion cells[44];
DOF1.8: vascular-related cells[45]; ANTHOCYANINLESS2 (ANL2), HOMEODOMAIN GLABROUS 1 (HDG1),
and HDG11: epidermis[31,46,47]]. **e,** A heatmap illustrates collapsed TF motif enrichment patterns into super motif
families across various species for each cell type. The motif enrichment score cutoff was set to 0.05. The score for
each super TF motif family was calculated by averaging the enrichment scores of all the TF motif members within
that super family. The DOF TF motif family was highlighted by a red frame. To mitigate the impact of substantial
variations in cell numbers across species or cell types, we standardized (down-sampled) the cell counts by randomly
selecting 412 cells per cell type per species. This count represents the lowest observed cell count for a given cell
type across all species (See Methods: Linear-model based motif enrichment analysis).

Species-specific evolution of cell-type-specific ACRs

To understand how cell-type-specific and broad ACRs changed over evolution, we examined ACRs within syntenic regions among the studied species. To compare ACRs, we devised a synteny-based BLASTN pipeline that allowed us to compare sequences directly (See Methods: Identification of syntenic regions; Fig. 4a; Extended Data Fig. 5a; Supplementary Table 17). Using *O. sativa* ACRs as a reference, we identified three classes of cross-species ACR conservation: 1) ACRs with matching sequences that are accessible in both species (shared ACRs), 2) ACRs with matching sequences, but are only accessible in one species (variable ACRs), and 3) ACRs where the sequence is exclusive to a single species (species-specific ACRs; Fig. 4b; Extended Data Fig. 3d). The shared ACR BLASTN hits were often small syntenic sequences, highlighting the large divergence of grass ACRs sequences. However, the majority (92-94%) of these shared BLASTN sequences encoded known TF motifs (Supplementary Fig. 18), indicating that shared ACRs are conserved regulatory regions. In contrast, variable ACRs represent a blend of conserved and divergent regulatory elements, and species-specific ACRs likely indicate novel regulatory loci. We found that, on average, shared ACRs were enriched ($p = $ 9e-50 to 4e-17; Fisher's exact test) for broad ACRs, whereas the variable ($p = $ 2e-23 to 2e-04; Fisher's exact test) and species-specific ($p = $ 2e-06 to 1e-02; Fisher's exact test) classes were enriched for cell-type specificity (Fig. 4c; Extended Data Fig. 5b and c). Moreover, we observed that the genomic distribution of shared ACRs were biased towards proximal ACRs (Extended Data Fig. 5d). This contrasts with cell-type-specific ACRs, which are overrepresented in the species-specific class (Fig. 4c). The cell-type-specific ACRs within the species-specific class were more likely to reside in distal genomic regions compared to the ACRs within the shared and variable classes (Extended Data Fig. 5e).

We further investigated whether the cell-type-specific ACRs were conserved in their cell-type specificity by evolution. Pairwise comparison between *O. sativa* and the other grasses revealed that between 0.7% (137/19,941) to 1.8% (420/22,881) of the syntenic ACRs were shared ACRs retaining the same cell-type specificity in both species (Extended Data Fig. 5b and f). Of these few shared cell-type specific ACRs, the majority (62%-69%), were accessible in the identical cell-type in both *O. sativa* and the corresponding species (Fig. 4d; Extended Data Fig. 5f). For example, the promoter ACR associated with *LATERAL ROOT DEVELOPMENT 3* (*LRD3*), a gene critical in companion cell and sieve element development[48], showed sequence

conservation between *O. sativa* and *S. bicolor* (Fig. 4d). Interestingly, ACRs which were

mesophyll specific in *O. sativa* changed their cell-type specificity to bundle sheath 17%-41% of

the time, while bundle sheath ACRs changed to mesophyll 9%-25% of the time (Fig. 4d;

Extended Data Fig. 5g). This result is likely due to the functional divergence associated with the

shift from C_3 (*O. sativa*) to C_4 (all other species sampled) photosynthesis[23,24]. Within variable

ACRs, leaf companion cell and sieve elements cells were consistently over-represented (Fig. 4e),

potentially due to sampling leaves at different stages of the source-sink transition. Of all the

classes of cross-species ACR conservation, species-specific ACRs were the most predominant in

every cell type (Fig. 4e). These findings suggest a dynamic and rapid evolution of cell-type-

specific ACRs within the examined species. Notably, ACRs in L1-derived layer (epidermis and

protoderm) exhibited the highest proportion of species-specific ACRs (Fig. 4e; Extended Data

Fig. 6a-c). The high divergence of ACRs in L1-derived cells was also observed in the pairwise

comparison between *O. sativa* and *Z. mays* seedling (Extended Data Fig. 6d and e), as well as

root cells (Extended Data Fig. 6f and g), suggesting that grass ACRs in L1-derived cells are

surprisingly divergent compared to internal cell types. Further, we examined *O. sativa*

(Supplementary Fig. 7; Supplementary Table 18) and *Z. mays* single-nucleus RNA sequencing

(snRNA-seq) data[20], to investigate whether the L1-derived cell types exhibited the most

divergent transcriptomes. We found that, among the six examined cell types, protoderm showed

the lowest similarity in the gene expression levels between the two species (Extended Data Fig.

6h), which suggests ACR divergence in L1-derived cells likely drives transcriptional change.

We further investigated the divergence of ACRs in L1-derived cells between more closely

related species pairs: *Z. mays* and *S. bicolor*; *P. miliaceum* and *U. fusca*. More closely related

species shared more ACRs across all cell types, containing fewer species-specific ACRs

(Extended Data Fig. 6i). Unlike when compared to the *O. sativa* reference (Fig. 4e; Extended

Data Fig. 6a-c), we did not observe more species-specific ACRs in L1-derived cells compared to

other cell-types (Extended Data Fig. 6j). Taken together, we observed that ACRs in L1-derived

cells are the most divergent compared to other cell types, but only when comparing grasses over

long evolutionary distances.

Fig. 4. Cell-type-specific ACRs are frequently species-specific. a, A screenshot illustrating syntenic regions

capturing shared ACRs across five species. The red bars denote syntenic ACRs within regions flanked by

corresponding syntenic gene pairs, while the gray color highlights these syntenic gene pairs. **b,** Three classes

depicting variations in ACR conservation between two species. 'Shared ACRs': ACRs with matching sequences that

are accessible in both species; 'Variable ACRs': ACRs with matching sequences but are only accessible in one

species; 'Species-specific ACRs': ACRs where the sequence is exclusive to a single species. **c,** The percentage of

broad and cell-type-specific ACRs underlying three classes shown in panel **b**. The significance test was done by

using the Fisher's exact test (alternative = 'two.sided'). **d, Left**, the number and percentage of *O. sativa* shared

ACRs that retain or change cell-type specificity amongst the other four species. **Right**, a screenshot of a *O. sativa*

phloem-specific ACR that retains phloem specificity in *S. bicolor*. This ACR is situated at the promoter region of

LRD3 which is specifically expressed in companion cell and phloem sieve elements (Supplementary Table 2). The

gray shaded region highlights the syntenic gene pair. **e,** The percentage of cell-type-specific ACRs identified across

all cell types within each species pair split into three classes shown in panel **b**. The percentage for each cell type

within the three classes collectively sum to 100%.

Returning to *O. sativa* as the reference, we investigated the TF families underpinning the

species-specific ACRs in L1-derived cells. Within all species-specific syntenic ACRs, we

observed a predominance of TF motifs for the HD-ZIP, SQUAMOSA PROMOTER BINDING

PROTEIN (SBP), PLINC families (Extended Data Fig. 7a). Many of these, such as HDG1,

ZHD1, ATHB-20, SPL3, SPL4, and SPL5, function in epidermal cell development[15,49-52]. The

predominance of these motifs in the species-specific ACR class suggests that although L1 TF

motif sequences are well conserved (Fig. 3d and e), the ACRs which contain these motifs are not

conserved in grass genomes. Upon comparing TF-motif enrichment in syntenic and non-syntenic

ACRs, we observed the presence of these epidermal motif families in both groups (Extended

Data Fig. 7b; Supplementary Table 19), indicating their essential roles in both conserved

epidermal cell development and rapid gene-regulatory co-option in species-specific sequences.

Notably, some TF-motif families, such as WRKY, were more enriched in non-syntenic ACRs in

epidermal cells (Extended Data Fig. 7b), further supporting that this family may drive

phenotypic innovation in the epidermal layer.

To look for derived species-specific ACRs associated with the altered expression of

surrounding gene orthologs in epidermal cells, we integrated snRNA-seq data from *O. sativa*

(Supplementary Fig. 7), with a snRNA-seq data from *Z. mays*[20]. We identified 87 orthologous

genes, irrespective of synteny, which exhibited higher L1 *O. sativa* expression compared to *Z.*

mays (Extended Data Fig. 7c; Supplementary Table 20). A gene ontology enrichment test for

these 87 genes revealed eight genes that were significantly enriched in the lipid metabolic

process (Extended Data Fig. 7d), possibly related to cuticle metabolism. Among the eight genes,

one was orthologous to *A. thaliana GDSL LIPASE GENE* (*LIP1*), which is epidermis specific[53].

We further identified 102 L1 cell-type-specific ACRs from *O. sativa* that were the closest to the

87 orthologous genes and observed 11 TF motifs enriched (q = 3e-10 to 5e-04; Binomial test) in

these ACRs (Extended Data Fig. 7c). These included TF family motifs known for their roles in

epidermal cell development such as ZHD1[15], HDG11[54], ZHD5[55], HDG1[31], and WRKY25[56]. For

example, within the *OsLIP1* intron, we identified two ZHD1 motifs within a species-specific

ACR that was specifically accessible in L1-derived cells (Extended Data Fig. 7e). We also

flipped this comparison by identifying 166 orthologs with elevated *Z. mays* epidermal expression

compared to *O. sativa* (Supplementary Table 20). This revealed 196 L1 cell-type-specific ACRs

in *Z. mays*. Within these ACRs, the most enriched (q = 0.0129 to 0.0392; Binomial test) TF

motif was MYELOBLASTOSIS 17 (MYB17; Extended Data Fig. 7f and g). This R2R3 MYB

family TF is associated with epidermal cell development, specifically in the regulation of

epidermal projections[57]. Furthermore, we hypothesized that some of these novel motifs could be

related to *O. sativa* transposable element (TE) expansion. Notably, we found long terminal repeat

retrotransposon (LTR)-associated ACRs from the *Gypsy* family were enriched ($p = 0.0012$ to

0.0415; Fisher's exact test) in *O. sativa* epidermal cell-type-specific ACRs (Extended Data Fig.

7h). The ZHD1 motif was enriched within these *Gypsy*-associated ACRs ($p = 0.0011$ to 0.0052;

Binomial test) (Extended Data Fig. 7i). By linking snRNA-seq to scATAC-seq data, we tied

gene-proximal ACR changes to elevated epidermal expression of a small number of conserved

orthologs 50 million years derived[58]. These ACR changes are associated with variance in

species-specific L1-derived layer development, potentially contributing to species differences in

environmental adaptation.

CNS are enriched in cell-type-specific ACRs

To augment our syntenic ACR BLASTN approach, we intersected our ACRs with published

CNS[6,59-61]. Outside of untranslated regions (UTRs), CNS typically encompass transcriptional

regulatory sequences undergoing purifying selection, too critical to be lost during evolution[62].

Using the conservatory database

(https://conservatorycns.com/dist/pages/conservatory/about.php)[63], we extracted 53,253 and

284,916 CNS in *O. sativa* and *Z. mays*, respectively, for analysis. Excluding CNS overlapping

with UTRs, 30.8% and 21.3% of CNS overlapped with the leaf-derived ACRs in *O. sativa* and *Z.*

mays, respectively (Fig. 5a; Extended Data Fig. 8a). Using all ACRs in the *O. sativa* atlas, this

ratio increased to 65.0% (Fig. 5a), indicating that a significant portion of these CNS likely

function in specific cell types and tissues. One common assumption is that CNS which overlap

ACRs (CNS ACRs) retain similar function between species[64-70]. This potential of conserved

function makes CNS tempting targets for genome editing, as changes to these sequences can

yield morphological variation, some that are important for crop improvement[71,72]. To test

assumptions that CNS within ACR classes are ideal targets for genetic modification, we assessed

conservation of cell-type specificity associated with CNSs. We observed 39% to 51% of total

CNS ACRs within the 'shared CNS ACR' class (Extended Data Fig. 8b and c), suggesting these

ACRs have conserved cellular contexts between *O. sativa* and other species. We also compared

shared ACRs between *O. sativa* and *Z. mays*, both containing CNS. We found that ACRs with

identical cell-type specificity had significantly ($p = 0.04114$; Wilcoxon signed rank test) longer

alignments and higher ($p = 3e-05$; Wilcoxon signed rank test) TF motif numbers than ACRs with

differing cell-type specificity (Extended Data Fig. 8d). Within syntenic regions, CNS ACRs are

more cell-type specific than those without (Fig. 5b). This enrichment was consistent for all

classes of ACRs we identified, except for species-specific ACRs which were equivalently cell-

type specific and broad (Fig. 5b). The enrichment of CNS in cell-type-specific ACRs stresses the

importance of rare cell-type function, as the cell-type-specific ACRs are preferentially retained

during evolution.

However, the majority (49-61%) of all CNS ACRs differed in cell-type specificity between

O. sativa and *Z. mays* (Extended Data Fig. 8b and c). Specifically, we examined the CNS found

in *Z. mays* ACRs that did not have a corresponding leaf ACR in *O. sativa*. Leveraging the *O.*

sativa atlas, we identified these sequences had divergent cellular or tissue chromatin

accessibilities. 249 (75%) of the *Z. mays* leaf variable CNS ACRs were accessible in non-leaf

cell states (Fig. 5c-e; Extended Data Fig. 8e), highlighting the instability of the cellular context in

which CNS acts. This suggests frequent co-option of CNS ACRs into different tissues or cell

types. Investigating the CNS ACRs that lost leaf cell-type specificity, we observed that these

ACRs were accessible in many non-leaf cell types, uniformly distributed amongst the atlas cell

annotations (Extended Data Fig. 8f-i). Consistent with our findings that epidermal-specific ACRs

tend to have the most species-specific ACRs in syntenic regions (Fig. 4e; Extended Data Fig. 6a-

c), L1-derived cells showed a significantly lower ratio of non-syntenic CNS ACRs compared to

other cell types (Extended Data Fig. 8j). This lower ratio demonstrates the frequent loss of

epidermal CNS, further supporting the rapid evolution of epidermal transcriptional regulation.

Interestingly, we noticed a pattern where some CNS within ACRs also overlapped domains

of H3K27me3[6] (Fig. 5f). H3K27me3 is a histone modification associated with facultative

heterochromatin established by the POLYCOMB REPRESSIVE COMPLEX 2 (PRC2)[73-75].

Genes silenced by PRC2 and H3K27me3 are important regulators that are only expressed in

narrow developmental stages or under specific environmental stimuli, where they often initiate

important transcriptional changes[74,76]. This importance makes the identification of key CREs

controlling H3K27me3 silencing especially interesting. Upon closer examination of the ACRs

overlapping domains of H3K27me3 to ACRs away from H3K27me3, we observed that

H3K27me3 ACRs were significantly enriched for CNS (Fig. 5g). This enrichment supports that

some of these CNS underpin conserved, and critical components, of H3K27me3 silencing.

Fig. 5. Cell-type-specific ACRs exhibit an enrichment of CNS. a, The percentage of CNS overlapping with

ACRs. 'n' indicates the number of CNS. 'Atlas' means the ACRs were from the *O. sativa* atlas in Fig. 1b. **b, Left,**

the percentage of broad and cell-type-specific ACRs within syntenic and non-syntenic regions overlapping with the

CNS. **Right,** this panel presented similar meaning as the left panel but focusing on three classes within syntenic

regions shown in Fig. 4b. Broad: Broadly accessible ACR; CT: Cell-type-specific ACR. Significance testing was

performed using the t-test (alternative = 'two.sided'). **c,** The bar plot showcases the count of *Z. mays* variable ACRs

accessible in leaf cell types. **d,** A sketch illustrating whether variable ACRs containing CNS in *Z. mays* capture

ACRs derived from the *O. sativa* atlas. **e,** The bar plot represents the count of *O. sativa* atlas ACRs accessible in

non-leaf cell types. **f,** An example of a syntenic block containing *O. sativa*-to-*Z. mays* conserved ACRs within a

H3K27me3 region. CNS were highlighted using red color. **g,** The percentage of ACRs capturing CNS in and outside

of H3K27me3 regions. The percentage for each group within H3K27me3 and not within H3K27me3 regions

collectively sum to 100%. Significance testing was performed using Fisher's exact test (alternative = 'two.sided').

Candidate silencer CREs are enriched in broad ACRs

To assess the stability and change of H3K27me3 related ACRs across grass lineages, we

focused on comparing *O. sativa*, *Z. mays*, and *S. bicolor* using published H3K27me3 ChIP-seq

data[6]. We examined ACRs near or within H3K27me3 regions and classified them into two

groups: H3K27me3-broad, representing H3K27me3 associated ACRs with chromatin

accessibility in many cell types and H3K27me3-cell-type specific, those H3K27me3 associated

ACRs with chromatin accessibility in few cell types (Fig. 6a; Extended Data Fig. 3g). The

proportion of H3K27me3-broad and H3K27me3-cell-type-specific ACRs was consistent across

all species (Extended Data Fig. 9a). H3K27me3-broad ACRs exhibited a depletion of

H3K27me3 at the ACR (Fig. 6b), consistent with nucleosome absence in ACRs[77]. In contrast,

H3K27me3 depletion was not observed in H3K27me3-cell-type-specific ACRs, with most cells

in the bulk ChIP-seq likely containing H3K27me3-modified nucleosomes (Fig. 6b). This is

consistent with the H3K27me3-cell-type-specific ACRs potentially acting after the removal of

facultative heterochromatin in a specific cell type(s). However, the chromatin accessibility of the

H3K27me3-broad ACRs appears to be concurrent with H3K27me3, suggesting these ACRs may

regulate H3K27me3 maintenance and removal across most cellular contexts.

To assess the transcriptional state of genes near H3K27me3-broad ACRs, we evaluated

snRNA-seq/scRNA-seq from *O. sativa* seedling (Supplementary Fig. 7) and root[78], and *Z. mays*

seedling[20]. The results revealed significantly lower expression (p = 3e-34 to 4e-06; Wilcoxon

signed rank test) for H3K27me3-broad ACRs associated genes across most cell types (Fig. 6c;

Extended Data Fig. 9b and c; Supplementary Table 21). Moreover, 58 bulk RNA-seq libraries

from *O. sativa* organs[79], demonstrated the lower (p = 2e-07 to 0.0265; Wilcoxon signed rank

test) expression of genes near H3K27me3-broad ACRs than genes near H3K27me3-absent broad

ACRs (Extended Data Fig. 9d). To dissect the roles of H3K27me3-broad ACRs, we identified

2,164 H3K27me3-broad ACRs and measured neighboring gene expression in *O. sativa* cells

(Supplementary Table 22). About 926 (~42.8%) of the H3K27me3-broad ACRs were associated

with 838 genes that exhibited no expression across any sampled cell type, which was marginally,

but significantly (p = 1e-10; Fisher's exact test), higher than the other ACRs associated with the

unexpressed genes (Extended Data Fig. 9e; Supplementary Table 22), consistent with these

H3K27me3 proximal genes only being expressed under specific conditions. The 1,108 expressed

genes associated with H3K27me3-broad ACRs were enriched (p < 2e-16; Fisher's exact test) for

cell-type specificity compared to genes without H3K27me3 (Extended Data Fig. 9f). In

summary, single-cell expression analysis revealed that the genes linked to H3K27me3-broad

ACRs exhibited the hallmarks of facultative gene silencing.

We hypothesized that the H3K27me3-broad ACRs would be enriched for PRC2 silencer

elements, as their consistent chromatin accessibility provides an avenue to recruit PRC2 to

maintain H3K27me3 throughout development. Supporting the presence of silencer CREs with

these ACRs, a known silencer CRE ~5.3 kb upstream of *FRIZZY PANICLE* was within a

H3K27me3-broad ACR[80] (Extended Data Fig. 10a). To exploit the known Polycomb *A. thaliana*

targets, we used scATAC-seq[20] and H3K27me3[81] data from *A. thaliana* roots and annotated

H3K27me3-broad ACRs. The *A. thaliana* H3K27me3-broad ACRs significantly ($p < 2e\text{-}16$;

Binomial test) captured 53 of the 170 known Polycomb responsive elements compared to a

control class of ACRs, supporting their putative silencer function[74] (Extended Data Fig. 10b).

Furthermore, we implemented a *de novo* motif analysis on the 170 *A. thaliana* elements and

identified all reported Polycomb response element (PRE) motifs (CTCC, CCG, G-box, GA-

repeat, AC-rich, and Telobox)[74] (Fig. 6d). Using these motifs and our chromatin accessibility

data, we predicted putative binding sites in *O. sativa*, *Z. mays*, and *S. bicolor*, and observed that

five motifs were significantly ($p = 2e\text{-}178$ to $1e\text{-}05$; Binomial test) enriched in the H3K27me3-

broad ACRs compared to a genomic control (Fig. 6d; Extended Data Fig. 10c). Between 88.0%

and 92.7% of H3K27me3-broad ACRs contained at least one PRE motif, with few (0.1-0.2%)

ACRs having all six PRE motif types (Extended Data Fig. 10d). Surprisingly, we observed the

rates of PRE motif occurrence, and PRE motif counts, were comparable between of H3K27me3-

broad ACRs and H3K27me3-absent-broad ACRs (Extended Data Fig. 10c-e). To investigate

why the H3K27me3-absent-broad ACRs still contain these motifs, we analyzed ChIP-seq of

EMF2b, a crucial PRC2 component[74,82,83], revealing a significant overlap between EMF2b peaks

and the PRE motifs (Extended Data Fig. 10f). Both H3K27me3-broad and H3K27me3-absent-

broad ACRs exhibited high EMF2b signals, consistent with the PRE motifs within both ACR

groups (Extended Data Fig. 10g). However, H3K27me3-broad ACRs showed higher EMF2b

signal in the shoulder (non-peak) areas compared to the H3K27me3-absent-broad ACRs. The

EMF2b ChIP-seq suggests PRC2 is recruited to many broadly accessible ACRs, but additional

factors are required to activate PRC2 deposition of H3K27me3. In addition, we identified 236

genes and their adjacent H3K27me3-broad ACRs containing PRE motifs with SNPs or Indels in

'Zhenshan 97' genotype using 'Nipponbare' as a reference in *O. sativa* [84,85]. A total of 133 of

these genes were detected with transcriptional changes between the two genotypes

(Supplementary Table 23). In 'Zhenshan 97', we observed a significant ($p = 0.03$; Wilcoxon

signed rank test) increase in gene expression, accompanied by a decrease in H3K27me3 signal

(Extended Data Fig. 10h-j). This suggests that mutations in PRE motifs may preclude PRC2

targeting, potentially associating with higher expression levels of nearby genes within the

examined two genotypes. Beyond PRE motifs, we observed significant enrichment (p = 1e-16 to

0.0376; Hypergeometric test) of motifs from four TF families in H3K27me3-broad ACRs:

APETALA2-like (AP2), basic Helix-Loop-Helix (bHLH), Basic Leucine Zipper (bZIP), and

C2H2 zinc-finger (ZnF) (Extended Data Fig. 10k; Supplementary Table 24). AP2 and C2H2 are

known to recruit PRC2[74], and our motif enrichment supports all these TF families potentially

regulating H3K27me3 deposition and facultative heterochromatin formation.

Fig. 6. Discovery of candidate silencer CREs across species. a, A sketch graph illustrates the classification of

ACRs based on their proximity to H3K27me3 peaks. We classified the ACRs into two groups: H3K27me3-

associated ACRs (found within or surrounding H3K27me3 peaks) and H3K27me3-absent ACRs. The H3K27me3-

associated ACR were further divided into broad ACRs, characterized by chromatin accessibility in at least five cell

types, and cell-type-specific ACRs, accessible in less than three out of six examined cell types across all the species.

b, Alignment of H3K27me3 chromatin attributes at summits of distinct ACR groups. **c,** A comparative analysis of

expression levels and chromatin accessibility of genes surrounding broad ACRs under and outside of H3K27me3

peaks. ** indicate p value < 0.01, which was performed by the Wilcoxon signed rank test (alternative = 'two.sided').

The broad ACRs where the H3K27me3 region overlapped >50% of the gene body were positioned within 500 to

5,000 bp upstream of the transcriptional start site of their nearest gene. **d,** Percentage of H3K27me3-broad ACRs in

To investigate the relationship between these candidate silencers and species divergence, we

mirrored our previous syntenic ACR analysis by classifying *O. sativa* H3K27me3-broad ACRs

into shared, variable or species specific groups (Extended Data Fig. 11a-c; Supplementary Table

25). Between 54% and 61% of the H3K27me3-broad ACRs were present in the species-specific

class, with the H3K27me3-broad ACRs enriched for species-specificity compared to

H3K27me3-absent-broad ACRs (Extended Data Fig. 11d). Since these H3K27me3-broad ACRs

exhibit hallmarks of PRC2 recruitment, we suspect that altered silencer CREs use context to

drive species-specific developmental and environmental responses.

Since we identified L1-derived cells as being enriched in species-specific ACRs, we sought

to examine the changes in H3K27me3 regulation within this tissue. We examined our previously

identified 87 *O. sativa*-to-*Z. mays* orthologs to see if these genes contained H3K27me3. We

observed 18 of these 87 genes were close to H3K27me3-broad ACRs (Supplementary Table 26).

For example, we identified four H3K27me3-broad ACRs, and three ZHD1 motifs linking

bulliform cell development[14,15], within two species-specific ACRs surrounding *OsNCED5* that

were specifically accessible in L1-derived cells (Fig. 6e). OsNCED5 TF is known to regulate

tolerance to water stress and regulate leaf senescence in *O. sativa*[86]. These results highlight that

H3K27me3 mediated silencing may play a critical role in divergent regulation in L1-derived

cells.

Discussion

Our comparison of *O. sativa* with four other grasses revealed patterns in the evolutionary

dynamics of *O. sativa* ACRs within syntenic and non-syntenic regions and discovered that the

grass L1-derived cells exhibit elevated rates of transcriptional regulatory divergence, as well as

changes in *cis* regulatory architecture as compared to other cell types over large evolutionary

distance (Fig. 7a). However, we revealed a contrasting dichotomy; the epidermal TF motifs

were the most cell-type specific of all those studied, yet their cognate ACRs exhibited the

strongest target divergence amongst the measured species. This duality highlights tandem

conservation of core epidermal motifs, and the rapid co-option of novel regulatory regions into

these existing regulatory frameworks. This rapid regulatory evolution might relate to the

dynamic environmental pressures the epidermis has evolved to withstand, and may relate to the

higher mutation rate in L1-derived tissues compared to other somatic cell types[13,87]. Although to

a lesser extent than the epidermis, this interesting contrast, where the cell-type restricted TF

motifs are conserved and the cell-type-specific chromatin accessibility of cognate ACRs are not,

extends to other cell types. This supports a larger pattern of novel ACR evolution that co-opts

established cell-type-specific TF networks.

Fig. 7. Evolution of cell-type-specific ACRs and CREs. **a,** The analysis of leaf cell types across these species

revealed an enrichment of cell-type-specific ACRs in species-specific regions. Notably, these species-specific ACRs

were enriched within L1-derived cells compared to all others examined. Additionally, it was observed that epidermal

cell-specific ACRs significantly overlapped with LTR-*Gypsy* TEs, which were enriched for the ZHD1 motif known

to regulate leaf curling. The L1 ACRs within the ZHD1 motif are likely associated with species-specific elevated

expression of a small number of genes involved in L1-derived cell development. **b,** We found an enrichment of CNS

in cell-type-specific ACRs. Although some CNS ACRs retained the same cell-type specificity between *O. sativa* and

Z. mays, these CNS ACRs often switched tissue or cell-type accessibility between grass species. **c,** Despite being

within facultative heterochromatin, H3K27me3-broad ACRs are accessible in many cell types, providing a physical

entry point for PRC2 to bind. Several lines of evidence support that the H3K27me3-broad ACRs contain silencer

CREs. Specifically, these ACRs are linked to transcriptionally silent genes, are enriched for PRE motifs, enriched

for TF family motifs (AP2 and C2H2) reported to recruit PRC2, and enriched for PRC2 subunit (EMF2b) ChIP-seq

peaks.

Highlighting the rapid rate of regulatory evolution, ACRs, and the CREs within them,

underpin phenotypic variation within plant and mammalian species[10,20,88,89]. Despite the link

between CREs and phenotypic variation, how these transcriptional regulatory circuits have

changed during species divergence is challenging to address. This is partly due to rapid CRE

changes occluding pairwise comparison; even closely related plant species share but a fraction of

their ACR/CRE complements[6,90]. We used a comparative single-cell epigenomics approach to

characterize the evolution of cell-type-specific ACRs and CREs in grasses. We demonstrate that

grass cell-type-specific ACRs have changed tremendously over 50 million years[58], with

relatively few (0.7% to 1.8%) cell-type-specific ACRs remaining conserved across the examined

plant species (Extended Data Fig. 5b and f). This contrasts with mammals, where previous work

on liver specific enhancers found that ~5% (2,151/43,020) remain conserved over ~100 million

years[91]. The difference in ACR conservation highlights the speed at which plant CRE evolution

takes place compared to mammals, which has been supported previously[59,90,92-95]. The repeated

whole genome duplications in plant lineages[96,97], and the functional redundancy they provide,

may be the fuel for rapid CRE divergence driving plants' adaptation to diverse environments[98].

Integration of the *O. sativa* atlas with CNS revealed ~65% were accessible in at least one cell

type (Fig. 5a). We expect that most CNS not captured by an ACR in our study are likely

accessible in an unsampled cell type, environmental, or developmental condition. This stresses

the need for expanded accessible chromatin atlases using more tissues, segments of development,

and environmental conditions. Most ACRs containing CNS had variable cell-type specificity

between species (Fig. 7b), highlighting that deeply conserved grass ACRs readily evolve new

spatiotemporal usage. Although less frequently than grass ACRs, around one third of conserved

mouse DNAseI–hypersensitive sites (DHSs) are altered in human tissue contexts after ~90

million years of evolution[69]. Thus, although eukaryotic CNS exhibit sequence conservation, their

functional context is often altered[71], with novel spatiotemporal CNS usage appearing prevalent in

grass lineages. However, it remains possible that the main CNS function conserved between *O.*

sativa and *Z. mays* occurs in non-leaf tissues. Nonetheless, the switching of cell-type

accessibility highlights the importance of merging chromatin accessibility data with CNS

datasets, as the assumption of conserved CNS sequence equaling conserved CRE function is

often invalid.

Much focus has been placed on enhancer CREs within ACRs, yet silencers are equally

important, as they repress gene expression until the proper developmental or environmental cues.

Our prior research uncovered that some ACRs flanking H3K27me3 are linked to the suppression

of nearby genes[6,9,20]; however, the question remains whether these ACRs function as silencers.

Our *O. sativa* cell-type ACR atlas allowed the identification of ACRs within H3K27me3 regions

that were accessible in most cell types. Several lines of evidence support these H3K27me3-broad

ACRs as silencers of linked, transcriptionally repressed genes. Specifically, these putative

silencers were enriched for CNS, PREs and related TF motifs, and PRC2 *in vivo* occupancy[83]

(Fig. 7c). Additional comparison of two genotypes in *O. sativa* reveals the PRE motif mutations

within H3K27me3-broad ACRs may preclude PRC2 targeting (Extended Data Fig. 10f), but

future genome editing of these putative silencers will reveal more about their role in PRC2

recruitment and gene silencing. Similar H3K27me3-broad ACR putative silencers were also

present in the epigenomic landscape of *O. sativa*, *Z. mays*, and *S. bicolor*, suggesting this is a

conserved feature of grass genomes. H3K27me3 silencing is deeply conserved in eukaryotes[99],

and a recent study found that many H3K27me3-marked regions might function as silencer-like

regulatory elements in *O. sativa*[76]. We hypothesize that other single-cell comparative genomic

investigations will find this pattern of broadly accessible silencers in other angiosperm species

with H3K27me3.

Our rice atlas of cell-type-specific ACRs and this cross-species analysis provides a useful

resource to enhance our understanding of regulatory evolution more broadly. Highlighting the

rapid rate of regulatory evolution, we believe combining this data with that from more closely

related grasses in the future will reveal more nuanced evolutionary dynamics of CRE evolution

under different levels of evolutionary time. This resource, and these observations, will fuel

research into identifying key CREs controlling specific genes by demarcating high-confident

targets for genome editing.

Methods

Preparation of plant materials

Early seedlings, specifically seedling tissues above ground, were collected 7 and 14 days after

sowing. Flag leaf tissue was harvested at the V4 stage, characterized by collar formation on leaf

4 of the main stem. Axillary buds were obtained from rice plants grown in the greenhouse at

approximately the V8 stage. Rice seminal and crown root tips (bottom 2 cm) were gathered at

the same stage as seedling tissues, 14 days after sowing. Panicle tissue was acquired from rice

plants grown in the greenhouse. Inflorescence primordia (2-5 mm) were extracted from shoots

harvested at the R1 growth stage, where panicle branches had formed. Early seeds were

harvested at approximately six days after pollination (DAP), and late seeds at approximately ten

DAP. All tissues were collected between 8 and 9 am, and all fresh materials were promptly

utilized for library construction starting at 10 am.

Single-cell ATAC-seq library preparation

Nuclei isolation and purification were performed as described previously[100]. In brief, the tissue

was finely chopped on ice for approximately 2 minutes using 600 μL of pre-chilled Nuclei

Isolation Buffer (NIB, 10 mM MES-KOH at pH 5.4, 10 mM NaCl, 250 mM sucrose, 0.1 mM

spermine, 0.5 mM spermidine, 1 mM DTT, 1% BSA, and 0.5% TritonX-100). After chopping,

the entire mixture was passed through a 40-μm cell strainer and then subjected to centrifugation

at 500 rcf for 5 minutes at 4°C. The supernatant was carefully decanted, and the pellet was

reconstituted in 500 μL of NIB wash buffer, which consisted of 10 mM MES-KOH at pH 5.4, 10

mM NaCl, 250 mM sucrose, 0.1 mM spermine, 0.5 mM spermidine, 1 mM DTT, and 1% BSA.

The sample was filtered again, this time through a 10-μm cell strainer, and then gently layered

onto the surface of 1 mL of a 35% Percoll buffer, prepared by mixing 35% Percoll with 65%

NIB wash buffer, in a 1.5-mL centrifuge tube. The nuclei were subjected to centrifugation at 500

rcf for 10 minutes at 4°C. Following centrifugation, the supernatant was carefully removed, and

the pellets were resuspended in 10 μL of diluted nuclei buffer (DNB, 10X Genomics Cat#

2000207). About 5 μL of nuclei were diluted 10 times and stained with DAPI (Sigma Cat.

D9542) and then the nuclei quality and density were evaluated with a hemocytometer under an

epifluorescence microscope. The original nuclei were diluted with a DNB buffer to a final

concentration of 3,200 nuclei per uL. Finally, 5 uL of nuclei (16,000 nuclei in total) were used as

input for scATAC-seq library preparation. scATAC-seq libraries were prepared using the

Chromium scATAC v1.1 (Next GEM) kit from 10X Genomics (Cat# 1000175), following the

manufacturer's instructions. (10xGenomics,

CG000209_Chromium_NextGEM_SingleCell_ATAC_ReagentKits_v1.1_UserGuide_RevE).

Libraries were sequenced with Illumina NovaSeq 6000 in dual-index mode with eight and 16

cycles for i7 and i5 index, respectively.

Single-nuclei RNA-seq library preparation and data analysis

The protocol for nuclei isolation and purification was adapted from the previously described

scATAC-seq method. In summary, to minimize RNA degradation and leakage, the tissue was

finely chopped on ice for approximately 1 minute using 600 µL of pre-chilled Nuclei Isolation

Buffer containing 0.4U/µL RNase inhibitor (Roche, Protector RNase Inhibitor, Cat. RNAINH-

RO) and a comparatively low detergent concentration of 0.1% NP-40. Following chopping, the

entire mixture was passed through a 40-µm cell strainer and then subjected to centrifugation at

500 rcf for 5 minutes at 4°C. The supernatant was carefully decanted, and the pellet was

reconstituted in 500 µL of NIB wash buffer, comprising 10 mM MES-KOH at pH 5.4, 10 mM

NaCl, 250 mM sucrose, 0.5% BSA, and 0.2U/µL RNase inhibitor. The sample was filtered

again, this time through a 10-µm cell strainer, and gently layered onto the surface of 1 mL of a

35% Percoll buffer. The Percoll buffer was prepared by mixing 35% Percoll with 65% NIB wash

buffer in a 1.5-mL centrifuge tube. The nuclei were then subjected to centrifugation at 500 rcf for

10 minutes at 4°C. After centrifugation, the supernatant was carefully removed, and the pellets

were resuspended in 50 µL of NIB wash buffer. Approximately 5 µL of nuclei were diluted

tenfold and stained with DAPI (Sigma Cat. D9542). Subsequently, the nuclei's quality and

density were evaluated with a hemocytometer under a microscope. The original nuclei were

further diluted with a DNB buffer to achieve a final concentration of 1,000 nuclei per µL.

Ultimately, a total of 16,000 nuclei were used as input for snRNA-seq library preparation. For

snRNA-seq library preparation, we employed the Chromium Next GEM Single Cell 3'GEM Kit

v3.1 from 10X Genomics (Cat# PN-1000123), following the manufacturer's instructions

(10xGenomics, CG000315_ChromiumNextGEMSingleCell3-

_GeneExpression_v3.1_DualIndex_RevB). The libraries were subsequently sequenced using the

Illumina NovaSeq 6000 in dual-index mode with 10 cycles for the i7 and i5 indices, respectively.

The raw BCL files obtained after sequencing were demultiplexed and converted into FASTQ

format using the default settings of the 10X Genomics tool cellranger mkfastq[101] (v7.0.0). The

raw reads were processed with cellranger count[101] (v7.0.0) using the Japonica rice reference

genome[102] (v7.0). Genes were kept if they were expressed in more than three cells and each cell

having a gene expression level of at least 1,000 but no more than 10,000 expressed genes. Cells

with over 5% mitochondria or chloroplast counts were filtered out. The expression matrix was

normalized to mitigate batch effects based on global-scaling normalization and multicanonical

correlation analysis in Seurat[103] (v4.0). The Scrublet tool[104] was employed to predict doublet

cells in this dataset. SCTransform in Seurat was used to normalize the data and identify variable

genes. The nearest neighbors were computed using FindNeighbors using 30 PCA dimensions.

The clusters were identified using FindClusters with a resolution of 1. The cell types were

annotated based on the marker gene list (Supplementary Table 2). To identify genes exhibiting

higher expression in a particular cell type than in the others, we utilized the 'cpm' function from

edgeR[105] (v3.38.1), for normalizing the expression matrix. Genes within a specific cell type that

displayed more than 1.5-fold change in log2 Counts Per Million (CPM) values compared to the

average log2 CPM across all cell types were determined as specifically expressed genes in that

particular cell type.

Transcriptome similarity between cell types of *O. sativa* and *Z. mays* was assessed using the

MetaNeighbor package[106]. To statistically compare similarities across different cell types, we

randomly divided the cells of each type into five groups. Each group was then used as input for

the MetaNeighbor analysis. The area under the receiver operating characteristic curve (auROC)

score obtained from MetaNeighbor was used as the similarity score in our analysis.

Slide-seq library preparation and data analysis

Root tissues from rice seedlings 14 days after sowing were used for the Slide-seq V2 spatial

transcriptomics. The tissues were embedded in the Optimal Cutting Temperature (OCT)

741 compound, snap-frozen in a cold 2-methylbutane bath, and cryosectioned into 10 um thick slices.

742 The spatial transcriptome library was constructed following a published method[107,108]. In brief,

743 the tissue slices were placed on the Slide-seq V2 puck and underwent the RNA hybridization and

744 reverse transcription process. After tissue clearing and spatial bead collections, the cDNA was

745 synthesized and amplified for a total of 14 cycles. The library was constructed using Nextera XT

746 Library Prep Kit (Illumina, USA) following the manufacturer's instructions.

748 The reads alignment and quantification were conducted following the Slide-seq pipeline

749 (https://github.com/MacoskoLab/slideseq-tools). The data processing is similar to the procedures

750 applied in the analysis of snRNA-seq but setting a resolution of 0.7 for the FindClusters function

751 in Seurat[103] (v4.0). The cell types were annotated based on the histology of cross-sectioned roots.

752 The marker genes of each cluster were identified using the Wilcoxon signed rank test in

753 FindAllMarkers.

755 **RNA *in situ* Hybridization**

756 The rice samples were put into the vacuum tissue processor (HistoCore PEARL, Leica) to fix,

757 dehydrate, clear, and embed, and were subsequently embedded in paraffin (Paraplast Plus,

758 Leica). The samples were sliced into 8 μm sections with a microtome (Leica RM2265). The

759 cDNAs of the genes were amplified with their specific primer pairs in situ_F/in situ_R and

760 subcloned into the pGEM-T vector (Supplementary Table 27). The pGEM-gene vectors were

761 used as the template to generate sense and antisense RNA probes. Digoxigenin-labeled RNA

762 probes were prepared using a DIG Northern Starter Kit (Roche) according to the manufacturer's

763 instructions. Slides were observed under bright fields through a microscope (ZEISS) and

764 photographed with an Axiocam 512 color charge-coupled device (CCD) camera.

766 **Raw reads processing of scATAC-seq**

767 Data processing was executed independently for each tissue and/or replicate. Initially, raw BCL

768 files were demultiplexed and converted into FASTQ format, utilizing the default settings of the

769 10X Genomics tool cellranger-atac make-fastq[109] (v1.2.0). Employing the Japonica rice

770 reference genome[102] (v7.0), the raw reads underwent processing using cellranger-atac count[109]

771 (v1.2.0). These steps encompassed adaptor/quality trimming, mapping, and barcode

attachment/correction. Subsequently to the initial processing, reads that were uniquely mapped

with mapping quality > 10 and correctly paired were subjected to further refinement through

SAMtools view (v1.7; -f 3 -q 10; Li et al. 2009). To mitigate the impact of polymerase chain

reaction (PCR) duplicates, the Picard tool MarkDuplicates[110] (v2.16.0) was applied on a per-

nucleus basis. To elevate data quality, a blacklist of regions was devised to exclude potentially

spurious reads. The methodology involved the exclusion of regions displaying bias in Tn5

integration from Tn5-treated genomic DNA. Specifically, regions characterized by 1-kilobase

windows with coverage exceeding four times the genome-wide median were eliminated. We

further leveraged ChIP-seq input data[6], to filter out collapsed sequences in the reference using

the same criteria. This blacklist also incorporated sequences of low-complexity and

homopolymeric sequences through RepeatMasker[111] (v4.1.2). Moreover, nuclear sequences

exhibiting homology surpassing 80% to mitochondrial and chloroplast genomes[112] (BLAST+;

v2.11.0) were also included within the blacklist. Furthermore, BAM alignments were converted

into BED format, wherein the coordinates of reads mapping to positive and negative strands

were subjected to a shift by +4 and -5, respectively. The unique Tn5 integration sites per barcode

were finally retained for subsequent analyses.

Identifying high-quality nuclei

To ensure the acquisition of high-quality nuclei, we harnessed the capabilities of the Socrates

package for streamlined processing[20]. To gauge the fraction of reads within peaks, we employed

MACS2[113] (v2.2.7.1) with specific parameters (genomesize = 3e8, shift= -75, extsize = 150, fdr

= 0.05) on the bulk Tn5 integration sites. Subsequently, we quantified the number of integration

sites per barcode using the callACRs function. Next, we estimated the proximity of Tn5

integration sites to genes, focusing on a 2 kb window surrounding the TSS. This estimation was

achieved through the buildMetaData function, which culminated in the creation of a meta file.

For further refinement of cell selection, we harnessed the findCells function, implementing

several criteria: 1) A minimum read depth of 1,000 Tn5 integration sites was required. 2) The

total number of cells was capped at 16,000. 3) The proportion of reads mapping to TSS sites was

above 0.2, accompanied by a z-score threshold of 3. 4) Barcode FRiP scores were required to

surpass 0.1, alongside a z-score threshold of 2. 5) We filtered out barcodes exhibiting a

proportion of reads mapping to mitochondrial and chloroplast genomes that exceeded two

standard deviations from the library mean. 6) We finally used the detectDoublets function to
estimate doublet likelihood by creating synthetic doublets and conducting enrichment analysis.
These multiple steps ensured the meticulous identification and selection of individual cells,
facilitating a robust foundation for subsequent analyses.
807

Nuclei clustering
For the nuclei clustering, we leveraged all functions from the Socrates package[20]. We binned the
entire genome into consecutive windows, each spanning 500 bp. We then tabulated the count of
windows featuring Tn5 insertions per cell. Barcodes falling below one standard deviation from
the mean feature counts (with a z-score less than 1) were excluded. Moreover, barcodes with
fewer than 1,000 features were eliminated. We pruned windows that exhibited accessibility in
less than 0.5% or more than 99.5% of all nuclei. To standardize the cleaned matrix, we applied a
term frequency-inverse document frequency normalization function. The dimensionality of the
normalized matrix underwent reduction through the utilization of non-negative matrix
factorization, facilitated by the R package RcppML[114] (v0.3.7). We retained 50 column vectors
from an uncentered matrix. Subsequently, we selected the top 30,000 windows that displayed the
highest residual variance across all cells. This selection was based on fitting a model where the
proportion of cells with accessibility served as an independent variable and variance as the
dependent variable. To further reduce the dimensionality of the nuclei embedding, we employed
the UMAP technique using umap-learn (k = 50, min_dist = 0.1, metrix = 'euclidean') in R[115]
(v0.2.8.0). Furthermore, we clustered nuclei using the callClusters function within the Socrates
framework[20]. Louvain clustering was applied, with a setting of k=50 nearest neighbors at a
resolution of 0.7. This process underwent 100 iterations with 100 random starts. Clusters with an
aggregated read depth of less than 1 million and 50 cells were subsequently eliminated. To filter
outlying cells in the UMAP embedding, we estimated the mean distance for each nucleus using
its 50 nearest neighbors. Nuclei that exceeded 3 standard deviations from the mean distance were
deemed outliers and removed from consideration.
830

Estimation of gene accessibility scores
To estimate the gene accessibility, we employed a strategy wherein the Tn5 insertion was
counted across both the gene body region as well as a 500 bp extension upstream. Subsequently,

we employed the SCTransform algorithm from the Seurat package[103] (v4.0) to normalize the

count matrix that was then transformed into a normalized accessibility score, with all positive

values scaled to 1.

Cell type validation

Upon completing the initial annotation process, which was based on a curated list of marker

genes, we further expanded our marker repertoire by incorporating markers collected from

published bulk RNA-seq data encompassing a diverse array of cell types, which were acquired

via laser capture dissection, as well as a published scRNA-seq data encompassing panicle related

cell types[116]. In brief, we collected the markers from several studies, including three distinct cell

types within rice leaves[117] and ten cell types across rice seed organs[118,119]. From these sources,

we selected the top 100 variably expressed markers for each cell type and employed them to

compute cell identity enrichment scores. We undertook a comprehensive assessment of markers

linked to the different cell types. Subsequently, we randomly drew 100 markers from this pool

and repeated this procedure 1,000 times to construct a null distribution based on marker

chromatin accessibility scores. For each target cell type marker, we compared their accessibility

scores to this null distribution. This facilitated the derivation of an enrichment score per cell,

delineating the marker's significance for each representative cell type. We next employed a

MAGIC algorithm[120] to refine these enrichment scores. These scores were then mapped onto a

UMAP plot, enhancing the cell identity annotation.

Furthermore, we undertook validation of our single-cell chromatin accessibility atlas through

integration with published scRNA-seq from rice root tissue. This validation was achieved

through two distinct approaches (Supplementary Fig. 8). In the first approach, we leveraged the

marked enrichment technique, adapting the above mentioned methodology with the

incorporation of the top 20 markers derived from marker identification using the 'FindMarkers'

function in Seurat[103] (v4.0). Following the acquisition of a smoothed score for each cell type,

individual cells were annotated to specific cell types based on the largest enrichment score within

that cell type. A threshold was further set, requiring the maximum score to exceed 0.5 for

confident labeling; otherwise, the cell was labeled as 'Unknown'. The second approach entailed

employing a k-nearest neighbor (knn) strategy. This strategy commenced with the normalization

of scRNA-seq datasets, mirroring the process applied to scATAC-seq datasets. The top 3,000

most variable genes within the scATAC-seq dataset were then identified using the Seurat

function 'FindVariableFeatures', subsequently filtering to include only genes common to both

datasets. By treating the scRNA-seq cells as a reference, a dimension reduction process was

conducted to generate a loading matrix, which was then utilized to project the scATAC-seq cells

onto the scRNA-seq cell embedding. The integration of these two datasets was achieved through

the Harmony algorithm[121] (v0.1.0). Within the dual embeddings, the 20 nearest neighbors of

each scATAC-seq cell in the scRNA-seq dataset were computed. The most frequent label among

these RNA neighbors (> 10 cells) was subsequently assigned as the label for each scATAC-seq

cell or designated as NA if no label meeting this threshold was identified.

Identification of *de novo* marker genes

For each cell type, we used edgeR[105], to identify cell-type-specific genes that are differentially

accessible. To determine if genes are accessible in a specific cell type, we compared the genes in

the target cell type to those in all other cell types, which served as the reference. We identified

genes with an FDR < 0.05 and log2 (Fold change) > 0 as candidate genes that are specifically

accessible in a particular cell type.

Cell cycle prediction

The prediction of cell cycle stages per nucleus was executed similarly to annotating cell

identities based on the aforementioned enriched scores. In brief, we collected a set of 55 cell-

cycle marker genes from a previous study[122]. For every cell-cycle stage, the cumulative gene

accessibility score for each nucleus was computed. These resultant scores were subsequently

normalized using the mean and standard deviation derived from 1,000 permutations of the 55

random cell-cycle stage genes, with exclusion of the focal stage. Z-scores corresponding to each

cell-cycle stage were transformed into probabilities using the 'pnorm' function in R.

Furthermore, the cell-cycle stage displaying the highest probability was designated as the most

probable cell stage.

ACR identification

Upon segregating the comprehensive single-base resolution Tn5 insertion sites BED dataset into

distinct subsets aligned with annotated cell types, we executed the MACS2 tool[113] (v2.2.7.1) for

precise peak identification per cell type. Notably we employed non-default parameters,

specifically: --extsize 150, --shift -75, -nomodel -keep-dup all. To mitigate potential false

positives, a permutation strategy was applied, generating an equal number of peaks based on

regions that were mappable and non-exonic. This approach encompassed the assessment of Tn5

insertion sites and density within both the original and permuted peak groups. By scrutinizing the

permutation outcomes, we devised empirically derived false discovery rate (FDR) thresholds

specific to each cell type. This entailed determining the minimum Tn5 density score within the

permutation cohort where the FDR remained < 0.05. To further eliminate peaks that exhibited

significant overlap with nucleosomes, we applied the NucleoATAC tool[123] (v0.2.1) to identify

potential nucleosome placements. Peaks that featured over 50% alignment with predicted

nucleosomes were systematically removed. The average fragment size of reads overlapping with

the peaks were calculated and the peaks with the average fragment size > 150 bp were filtered

out. Ultimately, the pool of peaks for each cell type was amalgamated and fine-tuned, yielding

500 bp windows that were centered on the summit of ACR coverage.

Identification of cell-type-specific ACRs

To identify cell-type-specific ACRs across the *O. sativa* atlas, we used a modified entropy metric

in our prior study[24]. Briefly, this method calculates the accessibility of ACRs, normalizes these

values using CPM, and then calculates both the entropy and specificity scores for each ACR in

each cell type. We combined this method with a bootstrapping approach[124]. For each cell type

and species, a sample of 250 cells was taken 5,000 times with replacement, and the specificity

metric was calculated as described above. This specificity metric was then compared against a

series of 5,000 null distributions, each consisting of a random shuffle of 250 cells from mixed

cell populations. This allowed us to set thresholds for what we would expect from non-cell-type-

specific data. A nonparametric test was then used to compare the median real bootstrap

specificity score to the null distributions. ACRs were labeled as cell-type-specific if they had a *p*

value less than 0.001. Cell-type-specific ACRs were those with a significant *p* value in a given

cell type and found to be specific in one or two cell types for leaf ACRs across five examined

species, and in one, two or three cell types for ACRs in the *O. sativa* atlas. This method was used

for all species in this study. Due to the number of ACRs and cell types in the *O. sativa* atlas, each

tissue was analyzed independently, and the results were merged downstream.

Interactions between distal cell-type-specific ACRs and cell-type-specific genes

Raw Hi-C data from *O. sativa* were collected including five replicates[25]. Low-quality reads were

filtered out using Trimmomatic[125]. The clean reads from each replicate were mapped to the

Japonica rice reference genome[102]. These mapped reads were then used to obtain normalized

contact maps through a two-step approach in the HiC-Pro software[126] in parallel mode. The

analysis was run using the default configuration file, with modifications to specify a minimal

mapping quality of 10 and enzyme recognition sites of MboI. The valid pairs files obtained from

all replicates of the same species were combined using the 'merge_persample' step in HiC-Pro

for further analysis. Fit-HiC[127] was used to identify intra-chromatin loops. The input contact

maps for Fit-HiC were generated from valid pairs files using the 'validPairs2FitHiC-fixedSize'

script at a 5 kb resolution. Fragments and bias files were generated using 'creatFitHiCFragments-

fixedsize' and 'HiCKRy' scripts separately. Significant intra-chromosomal interactions were

identified by running FitHiC at a 5 kb resolution. These significant interactions were further

merged and filtered using 'merge-filter.sh' script at a 5 kb resolution and with a FDR of 0.05.

Each bin of the significant intra-chromatin loops was intersected with distal cell-type-specific

ACRs using the 'intersect' function in bedtools[128]. The bins associated with those intersecting the

distal cell-type-specific ACRs were further examined to determine if any intersected with

promoter regions of cell-type-specific genes, defined as regions 2 kb upstream of the genes.

Aligning pseudotime trajectories between rice and maize

Motif deviation scores were computed utilizing ChromVAR[129] (v1.18.0) for both rice and maize

cells originated from the trajectories of root developing xylem and cortex development. To refine

these scores, a diffusion approach based on the MAGIC algorithm[130] was used. We further used

cellAlign tool[131] (v0.1.0), a technique that standardized the imputed deviation scores across a

predefined set of points (n = 200) distributed evenly along the trajectories of rice and maize. This

normalization strategy aimed to mitigate technical biases inherent in the data. For each motif pair

shared between rice and maize, a comprehensive global alignment procedure ensured to align the

imputed deviation scores across pseudotime for both *O. sativa* and *Z. mays*. Subsequently, we

calculated the normalized distance between the two species using the cellAlign tool. Motif

clustering based on these distance scores, using k-means, yielded two distinct groups: Group 1,

characterized by relatively higher distances, and Group 2. A linear regression framework was

subsequently introduced, using distance scores and pseudotime as predictive variables for motifs

within these two groups. In instances where motif pairs within either group exhibited positive or

negative coefficients, we classified them as 'shiftEarlyrice' or 'shiftEarlymaize'. Notably, motif

pairs in Group 1 were designated as 'conserved' if their coefficients bore identical positive or

negative attributes. To enhance the robustness of our findings, p values acquired from the linear

regression analysis underwent adjustment using the Benjamini-Hochberg procedure, effectively

addressing multiple comparisons. If the corrected p value exceeded 0.05, the motif pairs were

categorized as 'Unknown'.

Correlation between chromatin accessibility of TF genes and motif deviation

We sourced rice and *A. thaliana* TFs from PlantTFDB[132] (v4.0) database. To identify rice

orthologs of *A. thaliana* TFs, we employed BLAST[112] (BLAST+; v2.11.0) by utilizing protein

fasta alignments with an e-value threshold of 1e-5 used for significance. Alignments were

restricted to fasta sequences categorized as TFs from either species. To further refine the putative

orthologs, we applied filters based on functional similarity to *A. thaliana* TFs. Alignments with

less than 15% identity were excluded, along with rice TFs associated with distinct families. From

the remaining candidates, we selected the orthologs demonstrating the highest Pearson

correlation coefficient concerning the motif deviation scores. Motif deviation scores of specific

TF motifs within nuclei were computed *via* chromVAR[129] (v1.18.0).

Linear-model based motif enrichment analysis

We employed the FIMO tool from the MEME suite[133] (v5.1.1) with a significance threshold of p

value $< 10^{-5}$ to predict motif locations. The motif frequency matrix used was sourced from the

JASPAR plants motif database[134] (v9). Subsequently, we constructed a binarized peak-by-motif

matrix and a motif-by-cell count matrix. This involved multiplying the peak-by-cell matrix with

the peak-by-motif matrix. To address potential overrepresentation and computational efficiency,

down-sampling was implemented. Specifically, we standardized the cell count by randomly

selecting 412 cells per cell type per species. This count represents the lowest observed cell count

for a given cell type across all species. For each cell type annotation, total motif counts were

predicted through negative binomial regression. This involved two input variables: an indicator

column for the annotation, serving as the primary variable of interest, and a covariate

representing the logarithm of the total number of nonzero entries in the input peak matrix for

each cell. The regression provided coefficients for the annotation indicator column and an

intercept. These coefficients facilitated the estimation of fold changes in motif counts for the

annotation of interest in relation to cells from all other annotations. This iterative process was

conducted for all motifs across all cell types. The obtained p values were adjusted using the

Benjamini-Hochberg procedure to account for multiple comparisons. Finally, enriched motifs

were identified by applying a dual filter criterion: corrected p values < 0.01, fold-change of the

top enriched TF motif in cell type-specific peaks for all cell types should be over 1, and beta

(motif enrichment score) > 0.05 or beta > 0.

Binomial test-based motif enrichment analysis

To assess the enrichment of motifs in a target set of ACRs, we performed analysis for each

specific motif. We randomly selected an equivalent number of ACRs as found in the target set,

repeating this process 100 times. Notably, the randomly selected ACR set did not overlap with

the actual target set of ACRs. Following this, we computed the average ratio of ACRs capturing

the motif within the null distribution.

1006

Subsequently, we executed an exact Binomial test[135], wherein we set this ratio as the

hypothesized probability of success. The number of ACRs overlapping the motif in the target set

was considered the number of successes, while the total number of ACRs in the target set

represented the number of trials. The alternative hypothesis was specified as 'two.sided'. This

meticulous approach allowed us to robustly evaluate and identify significant motif enrichments

within the target set of ACRs.

1013

1014   **Construction of control sets for enrichment tests**

1015   To check if non-CDS QTNs could be significantly captured by ACRs, we generated control sets

1016   by simulating sequences with the same length as non-CDS QTNs 100 times, yielding a mean

1017   proportion for the control sets. The binomial test p value was calculated by comparing the mean

1018   ratio to the observed overlapping ratio of non-CDS QTNs captured by ACRs.

1019

1020   To check if ACRs could significantly capture CNS, we generated control sets by simulating

1021   sequences with the same length as ACRs 100 times, yielding a mean proportion for the control

1022   sets. The binomial test p value was calculated by comparing the mean ratio to the observed

1023   overlapping ratio of ACRs capturing the CNS.

1024

1025   To perform comparative analysis of expression levels and chromatin accessibility of genes

1026   surrounding broad ACRs under and outside of H3K27me3 peaks, we sampled the same number

1027   of ACRs per cell type regarding the broad ACRs not under H3K27me3 peaks. This step is to

1028   make sure that their nearby gene chromatin accessibility exhibited similar values compared to

1029   the broad ACRs under the H3K27me3 peaks.

1030

1031   To check if the H3K27me3-broad-ACRs could significantly capture the known PREs and

1032   capture the EMF2b ChIP-seq peaks, we generated control sets by randomly selecting not-

1033   H3K27me3-broad-ACR instances 100 times, yielding a mean number value for the control sets.

1034   The Binomial test p value was calculated by comparing the mean ratio to the observed number of

1035   H3K27me3-broad ACRs overlapped with the PREs.

1036

To test if H3K27me3-broad ACRs in *O. sativa*, *Z. mays*, and *S. bicolor* significantly capture six

known motifs, we generated control sets by simulating sequences with the same length as ACRs

100 times, yielding a mean proportion for the control sets. The binomial test p value was

calculated by comparing the mean ratio to the observed overlapping ratio of H3K27me3-broad

ACRs capturing the motifs. The same process was conducted to examine whether the EMF2b

peaks significantly capture the six motifs.

1043

Identification of H3K27me3-broad ACRs

We first implemented a series of cutoffs to determine whether the peak is accessible for a

specific cell type. For each cell type, we first normalized the read coverage depth obtained from

the MACS2 tool divided by total count of reads, and ensured that the maximum of normalized

coverage within the peak exceeded a predefined threshold set at 2. Additionally, we calculated

Tn5 integration sites per peak, filtering out peaks with fewer than 20 integration sites.

Subsequently, we constructed a peak by cell type matrix with Tn5 integration site counts. This

matrix underwent normalization using the 'cpm' function wrapped in edgeR[105] (v3.38.1) and

'normalize.quantiles' function wrapped within preprocessCore[136] (v1.57.1) in the R

programming environment. To further refine our selection, a threshold of 2 was set for the counts

per million value per peak per cell type. Peaks that satisfied these distinct cutoff criteria were

deemed accessible in the designated cell types. For analyzing cell types with fewer than 10

samples (n < 10), we established criteria where H3K27me3-broad ACRs must be accessible in at

least n-1 cell types, while cell-type-specific ACRs should be accessible in fewer than 3 of the

examined cell types. For analyses involving more than 10 cell types, we adjusted the criteria:

H3K27me3-broad ACRs must be accessible in at least n−2 cell types, and cell-type-specific

ACRs should be accessible in fewer than 4 of the examined cell types.

1061

De novo **motif analysis**

To identify position weight matrix of six known motifs within 170 *A. thaliana* PREs[74], we

employed the streme function with default settings from the MEME suite[133] (v5.1.1). The control

sequences were built up to match each PRE sequence by excluding exons, PREs, and

unmappable regions, and they possess a similar GC content (< 5% average difference) and same

sequence length compared to the positive set.

1068

Identification of syntenic regions

Identification of syntenic gene blocks was done using the GENESPACE[137] (v1.4). In brief, to

establish orthologous relationships between ACR sequences, ACRs in the *O. sativa* genome were

extended to incorporate the two closest gene models for a 'query block' since GENESPACE

only draws relationships between protein coding sequences. Then the GENESPACE function

'query_hits' was used with the argument 'synOnly = TRUE' to retrieve syntenic blocks. The

resulting syntenic hits were further filtered to allow only a one-to-one relationship between

O.sativa and the corresponding species. The corresponding syntenic blocks were then named and

numbered, and both the genes and genomic coordinates were recorded.

1078

To further identify corresponding ACRs within these blocks we set up a BLASTN pipeline[138]

(v2.13.0). For each comparison of species, using *O. sativa* as the reference the underlying

nucleotide sequences of the syntenic regions were extracted using Seqkit, and used as the blast

reference database[139] (v2.5.1). The sequences underlying the ACRs within the same syntenic

region in a different species were then used as the query. The blast was done using the following

parameters to allow for alignment of shorter sequences '-task blastn-short -evalue 1e-3 -

max_target_seqs 4-word_size 7 -gapopen 5 -gapextend 2 -penalty -1 -reward 1 -outfmt 6'. This

procedure was run for each syntenic region separately for all species comparisons. The resulting

BLASTN files were combined, and then filtered using a custom script. Alignments were only

considered valid if the e-value passed a stringent threshold of 1e-3, and the alignment was

greater than 20 nucleotides with the majority of the shared ACRs (92% to 94%) containing the

alignment regions including TF motif binding sites (Supplementary Fig. 19). The resulting

filtered BLAST files, and the BED files generated from these BLAST files allowed us to draw

our relationships between ACRs in the corresponding syntenic space. For all analyses, ACRs

were considered to have conserved cell-type-specificity if these ACRs would be aligned by

BLAST and had the same cell-type as assigned by the above method.

1095

1096 **Estimation of conservation scores**

1097 Conservation scores were predicted using PhyloP[140] (v1.0), where values are scaled between 0 to

1098 1, with one being highly conserved and 0 being non-conserved. Phylogenies to train PhyloP were

1099 generated using PhyloFit[141] (v1.0), and neutral and conserved sequences were identified using

1100 the whole genome aligner progressive cactus.

1101

ChIP-seq analysis

The clean reads of EMF2b were downloaded from a previous study[83]. The reads were mapped to

the rice reference genome[102] (v7.0) using bowtie2[142] (v2.5.2) with the following parameters: '--

very-sensitive --end-to-end'. Reads with MAPQ > 5 were used for the subsequent analysis.

Aligned reads were sorted and duplicated reads were removed using SAMtools[143] (v1.7). Peak

calling was performed using epic2[144] with the following parameters: '-fdr 0.01 --bin-size 150 --

gaps-allowed 1'. The peak 'BED' and 'BIGWIG' files of H3K27me3 ChIP-seq data for leaf,

root, and panicle rice organs were downloaded from RiceENCODE[145]

(http://glab.hzau.edu.cn/RiceENCODE/).

1111

Genetic variants calling in ZS97 genotype in *O. sativa*

We obtained raw sequencing data for the ZS97 genotype of *O. sativa* from a published study[85].

After quality filtering the raw reads using fastp[146] (v0.23.4), we aligned them to the Japonica *O.*

sativa reference genome (Ouyang et al., 2007) using the BWA-MEM algorithm[147] (v0.7.8). We

then used Picard tool MarkDuplicates[110] (v2.16.0) to remove PCR duplicates. The final genetic

variants file was generated using the HaplotypeCaller function in GATK[148] (GATK 4.2.3.0).

1118

1119 **GO enrichment test**

1120 The GO enrichment tests were performed based on the AgriGO[149] (v2) by setting the Chi-square

1121 statistical test and multi-test adjustment method is Hochberg (FDR).

1122

1123 **Additional resources**

1124 Cell-type resolved data can be viewed through our public Plant Epigenome JBrowse Genome

1125 Browser[150] (http://epigenome.genetics.uga.edu/PlantEpigenome/index.html)

1126

Data Availability

scATAC-seq data encompassing 18 libraries from nine organs were accessible in NCBI

(PRJNA1007577/GSE252040;

https://dataview.ncbi.nlm.nih.gov/object/PRJNA1007577?reviewer=kgarq48dii11vomg44kgr1jq

66; PRJNA1052039;

https://dataview.ncbi.nlm.nih.gov/object/PRJNA1052039?reviewer=flhu9sl84o5m999r1ph8tlm

mbg)

1134

Acknowledgements

This research was funded by the National Science Foundation (IOS-2134912) to SRW and RJS

and the UGA Office of Research to RJS. APM and JPM were supported by the National

Institutes of Health (K99GM144742) and (T32GM142623), respectively. DW was supported by

National Science Foundation MCB-2224729 and IOS-1953279. We thank Changhui Sun (Rice

Research Institute, Sichuan Agricultural University, Chengdu 611130, China) for identifying

genetic variants of ZS97 using NIP as reference genotype in *O. sativa*.

1142

Contributions

R.J.S., H.Y., J.P.M., A.P.M., M.A.A.M., D.W., S.Z., and S.R.W. designed and conceived

experiments and managed the project. X.Z., Y.L., Z.L., X.T., S.Z., Y.W., and H.Y. participated

in material collection and sample processing. H.Y., J.P.M., and T.R. performed the

bioinformatics analyses. H.Y. and J.P.M. wrote the manuscript. Y.L. Y.W. and Z.L. contributed

to marker validation. R.J.S., J.P.M., A.P.M., M.A.A.M., D.W., S.Z., and S.R.W. edited the

manuscript. L.H. contributed to some computing resources.

1150

1151   **Competing interests**

1152   R.J.S. is a co-founder of REquest Genomics, LLC, a company that provides epigenomic services.

1153

1154   **Supplementary information**

1155   **Supplementary Note:**

1156   1. Cell-type annotation and validation

1157   **Supplementary Tables**

1158   Supplementary Tables 1–27.

References

Preissl, S., Gaulton, K. J. & Ren, B. Characterizing cis-regulatory elements using single-cell epigenomics. *Nature Reviews Genetics* **24**, 21-43 (2023).

Wittkopp, P. J. & Kalay, G. Cis-regulatory elements: molecular mechanisms and evolutionary processes underlying divergence. *Nature Reviews Genetics* **13**, 59-69 (2012).

Marand, A. P., Eveland, A. L., Kaufmann, K. & Springer, N. M. cis-Regulatory elements in plant development, adaptation, and evolution. *Annual review of plant biology* **74**, 111-137 (2023).

Oka, R. *et al.* Genome-wide mapping of transcriptional enhancer candidates using DNA and chromatin features in maize. *Genome biology* **18**, 1-24 (2017).

Maher, K. A. *et al.* Profiling of accessible chromatin regions across multiple plant species and cell types reveals common gene regulatory principles and new control modules. *The Plant Cell* **30**, 15-36 (2018).

Lu, Z. *et al.* The prevalence, evolution and chromatin signatures of plant regulatory elements. *Nature Plants* **5**, 1250-1259 (2019).

Reynoso, M. A. *et al.* Evolutionary flexibility in flooding response circuitry in angiosperms. *Science* **365**, 1291-1295 (2019).

Kajala, K. *et al.* Innovation, conservation, and repurposing of gene function in root cell type development. *Cell* **184**, 3333-3348. e3319 (2021).

Ricci, W. A. *et al.* Widespread long-range cis-regulatory elements in the maize genome. *Nature plants* **5**, 1237-1249 (2019).

Cusanovich, D. A. *et al.* A single-cell atlas of in vivo mammalian chromatin accessibility. *Cell* **174**, 1309-1324. e1318 (2018).

Domcke, S. *et al.* A human cell atlas of fetal chromatin accessibility. *Science* **370**, eaba7612 (2020).

Lu, Z. *et al.* Tracking cell-type-specific temporal dynamics in human and mouse brains. *Cell* **186**, 4345-4364. e4324 (2023).

Javelle, M., Vernoud, V., Rogowsky, P. M. & Ingram, G. C. Epidermis: the formation and functions of a fundamental plant tissue. *New Phytologist* **189**, 17-39 (2011).

Kadioglu, A., Terzi, R., Saruhan, N. & Saglam, A. Current advances in the investigation of leaf rolling caused by biotic and abiotic stress factors. *Plant Science* **182**, 42-48 (2012).

Xu, Y. *et al.* Overexpression of OsZHD1, a zinc finger homeodomain class homeobox transcription factor, induces abaxially curled and drooping leaf in rice. *Planta* **239**, 803-816 (2014).

Dorrity, M. W. *et al.* The regulatory landscape of Arabidopsis thaliana roots at single-cell resolution. *Nature communications* **12**, 3334 (2021).

Farmer, A., Thibivilliers, S., Ryu, K. H., Schiefelbein, J. & Libault, M. Single-nucleus RNA and ATAC sequencing reveals the impact of chromatin accessibility on gene expression in Arabidopsis roots at the single-cell level. *Molecular Plant* **14**, 372-383 (2021).

Tu, X., Marand, A. P., Schmitz, R. J. & Zhong, S. A combinatorial indexing strategy for low-cost epigenomic profiling of plant single cells. *Plant Communications* **3** (2022).

Nobori, T. *et al.* Time-resolved single-cell and spatial gene regulatory atlas of plants under pathogen attack. *bioRxiv*, 2023.2004. 2010.536170 (2023).

20   Marand, A. P., Chen, Z., Gallavotti, A. & Schmitz, R. J. A cis-regulatory atlas in maize at
single-cell resolution. *Cell* **184**, 3041-3055. e3021 (2021).

21   Feng, D. *et al.* Chromatin accessibility illuminates single-cell regulatory dynamics of rice
root tips. *BMC biology* **20**, 274 (2022).

22   Zhang, L. *et al.* Asymmetric gene expression and cell-type-specific regulatory networks
in the root of bread wheat revealed by single-cell multiomics analysis. *Genome Biology*
24, 65 (2023).

23   Swift, J. *et al.* Single nuclei sequencing reveals C4 photosynthesis is based on rewiring
of ancestral cell identity networks. *bioRxiv*, 2023.2010. 2026.562893 (2023).

24   Mendieta, J. P. *et al.* Investigating the cis-Regulatory Basis of C3 and C4 Photosynthesis
in Grasses at Single-Cell Resolution. *National Academy of Sciences of the United States*
of America **121**, e2402781121 (2024).

25   Dong, Q. *et al.* Genome-wide Hi-C analysis reveals extensive hierarchical chromatin
interactions in rice. *The Plant Journal* **94**, 1141-1156 (2018).

26   Wei, X. *et al.* A quantitative genomics map of rice provides genetic insights and guides
breeding. *Nature Genetics* **53**, 243-253 (2021).

27   Yang, Y. *et al.* Natural variation of OsGluA2 is involved in grain protein content
regulation in rice. *Nature communications* **10**, 1949 (2019).

28   Zemach, A. *et al.* Local DNA hypomethylation activates genes in rice endosperm.
Proceedings of the National Academy of Sciences **107**, 18729-18734 (2010).

29   Xu, Q. *et al.* DNA demethylation affects imprinted gene expression in maize endosperm.
Genome biology **23**, 77 (2022).

30   Ohashi-Ito, K. & Fukuda, H. HD-Zip III homeobox genes that include a novel member,
ZeHB-13 (Zinnia)/ATHB-15 (Arabidopsis), are involved in procambium and xylem cell
differentiation. *Plant and Cell Physiology* **44**, 1350-1358 (2003).

31   Wu, R. *et al.* CFL1, a WW domain protein, regulates cuticle development by modulating
the function of HDG1, a class IV homeodomain transcription factor, in rice and
Arabidopsis. *The Plant Cell* **23**, 3392-3411 (2011).

32   Zhong, R., Richardson, E. A. & Ye, Z.-H. The MYB46 transcription factor is a direct
target of SND1 and regulates secondary wall biosynthesis in Arabidopsis. *The Plant Cell*
19, 2776-2792 (2007).

33   Wang, Z. *et al.* Salicylic acid promotes quiescent center cell division through ROS
accumulation and down-regulation of PLT1, PLT2, and WOX5. *Journal of Integrative*
Plant Biology **63**, 583-596 (2021).

34   Gontarek, B. C., Neelakandan, A. K., Wu, H. & Becraft, P. W. NKD transcription factors
are central regulators of maize endosperm development. *The Plant Cell* **28**, 2916-2936
(2016).

35   Sun, X. *et al.* Activation of secondary cell wall biosynthesis by miR319-targeted TCP 4
transcription factor. *Plant Biotechnology Journal* **15**, 1284-1294 (2017).

36   Ortiz-Ramírez, C. *et al.* Ground tissue circuitry regulates organ complexity in maize and
Setaria. *Science* **374**, 1247-1252 (2021).

37   Lv, Z., Zhao, W., Kong, S., Li, L. & Lin, S. Overview of molecular mechanisms of plant
leaf development: a systematic review. *Frontiers in Plant Science* **14** (2023).

38   Le Hir, R. & Bellini, C. The plant-specific Dof transcription factors family: new players
involved in vascular system development and functioning in Arabidopsis. *Frontiers in*
Plant Science **4**, 164 (2013).

39 Dai, X. *et al.* Chromatin and regulatory differentiation between bundle sheath and
mesophyll cells in maize. *The Plant Journal* **109**, 675-692 (2022).
40 Yanagisawa, S. Dof1 and Dof2 transcription factors are associated with expression of
multiple genes involved in carbon metabolism in maize. *The Plant Journal* **21**, 281-288
(2000).
41 Yanagisawa, S. & Sheen, J. Involvement of maize Dof zinc finger proteins in tissue-
specific and light-regulated gene expression. *The Plant Cell* **10**, 75-89 (1998).
42 Borba, A. R. *et al.* Synergistic binding of bHLH transcription factors to the promoter of
the maize NADP-ME gene used in C4 photosynthesis is based on an ancient code found
in the ancestral C3 state. *Molecular biology and evolution* **35**, 1690-1705 (2018).
43 Shimadzu, S., Furuya, T. & Kondo, Y. Molecular mechanisms underlying the
establishment and maintenance of vascular stem cells in Arabidopsis thaliana. *Plant and
Cell Physiology* **64**, 274-283 (2023).
44 Otero, S. & Helariutta, Y. Companion cells: a diamond in the rough. *Journal of
experimental botany*, erw392 (2016).
45 Ramachandran, V. *et al.* Plant-specific Dof transcription factors VASCULAR-
RELATED DOF1 and VASCULAR-RELATED DOF2 regulate vascular cell
differentiation and lignin biosynthesis in Arabidopsis. *Plant molecular biology* **104**, 263-
281 (2020).
46 Kubo, H., Kishi, M. & Goto, K. Expression analysis of ANTHOCYANINLESS2 gene in
Arabidopsis. *Plant science* **175**, 853-857 (2008).
47 Amanda, D. *et al.* DEFECTIVE KERNEL1 (DEK1) regulates cell walls in the leaf
epidermis. *Plant Physiology* **172**, 2204-2218 (2016).
48 Ingram, P., Dettmer, J., Helariutta, Y. & Malamy, J. E. Arabidopsis Lateral Root
Development 3 is essential for early phloem development and function, and hence for
normal root system development. *The Plant Journal* **68**, 455-467 (2011).
49 Chen, X. *et al.* SQUAMOSA promoter-binding protein-like transcription factors: Star
players for plant growth and development. *Journal of integrative plant biology* **52**, 946-
951 (2010).
50 Denyer, T. *et al.* Spatiotemporal developmental trajectories in the Arabidopsis root
revealed using high-throughput single-cell RNA sequencing. *Developmental cell* **48**, 840-
852. e845 (2019).
51 Fang, J. *et al.* The URL1–ROC5–TPL2 transcriptional repressor complex represses the
ACL1 gene to modulate leaf rolling in rice. *Plant Physiology* **185**, 1722-1744 (2021).
52 Horstman, A. *et al.* AIL and HDG proteins act antagonistically to control cell
proliferation. *Development* **142**, 454-464 (2015).
53 Rombolá-Caldentey, B., Rueda-Romero, P., Iglesias-Fernández, R., Carbonero, P. &
Oñate-Sánchez, L. Arabidopsis DELLA and two HD-ZIP transcription factors regulate
GA signaling in the epidermis through the L1 box cis-element. *The Plant Cell* **26**, 2905-
2919 (2014).
54 Yu, L. H. *et al.* Arabidopsis EDT 1/HDG 11 improves drought and salt tolerance in
cotton and poplar and increases cotton yield in the field. *Plant biotechnology journal* **14**,
72-84 (2016).
55 Hong, S.-Y., Kim, O.-K., Kim, S.-G., Yang, M.-S. & Park, C.-M. Nuclear import and
DNA binding of the ZHD5 transcription factor is modulated by a competitive peptide
inhibitor in Arabidopsis. *Journal of Biological Chemistry* **286**, 1659-1668 (2011).

56   Rosado, D., Ackermann, A., Spassibojko, O., Rossi, M. & Pedmale, U. V. WRKY
transcription factors and ethylene signaling modify root growth during the shade-
avoidance response. *Plant Physiology* **188**, 1294-1311 (2022).
57   Brockington, S. F. *et al.* Evolutionary analysis of the MIXTA gene family highlights
potential targets for the study of cellular differentiation. *Molecular Biology and Evolution*
30, 526-540 (2013).
58   Wolfe, K. H., Gouy, M., Yang, Y.-W., Sharp, P. M. & Li, W.-H. Date of the monocot-
dicot divergence estimated from chloroplast DNA sequence data. *Proceedings of the*
National Academy of Sciences **86**, 6201-6205 (1989).
59   Woolfe, A. *et al.* Highly conserved non-coding sequences are associated with vertebrate
development. *PLoS biology* **3**, e7 (2005).
60   Babarinde, I. A. & Saitou, N. Genomic locations of conserved noncoding sequences and
their proximal protein-coding genes in mammalian expression dynamics. *Molecular*
biology and evolution **33**, 1807-1817 (2016).
61   Song, B. *et al.* Conserved noncoding sequences provide insights into regulatory sequence
and loss of gene expression in maize. *Genome research* **31**, 1245-1257 (2021).
62   Nelson, A. C. & Wardle, F. C. Conserved non-coding elements and cis regulation:
actions speak louder than words. *Development* **140**, 1385-1395 (2013).
63   Hendelman, A. *et al.* Conserved pleiotropy of an ancient plant homeobox gene uncovered
by cis-regulatory dissection. *Cell* **184**, 1724-1739. e1716 (2021).
64   Pennacchio, L. A. *et al.* In vivo enhancer analysis of human conserved non-coding
sequences. *Nature* **444**, 499-502 (2006).
65   McEwen, G. K. *et al.* Ancient duplicated conserved noncoding elements in vertebrates: a
genomic and functional analysis. *Genome research* **16**, 451-465 (2006).
66   Turner, E. E. & Cox, T. C. Genetic evidence for conserved non-coding element function
across species–the ears have it. *Frontiers in physiology* **5**, 7 (2014).
67   Nolte, M. J. *et al.* Functional analysis of limb transcriptional enhancers in the mouse.
Evolution & development **16**, 207-223 (2014).
68   Salvi, S. *et al.* Conserved noncoding genomic sequences associated with a flowering-time
quantitative trait locus in maize. *Proceedings of the National Academy of Sciences* **104**,
11376-11381 (2007).
69   Vierstra, J. *et al.* Mouse regulatory DNA landscapes reveal global principles of cis-
regulatory evolution. *Science* **346**, 1007-1012 (2014).
70   Leypold, N. A. & Speicher, M. R. Evolutionary conservation in noncoding genomic
regions. *Trends in Genetics* **37**, 903-918 (2021).
71   Ciren, D., Zebell, S. & Lippman, Z. B. Extreme restructuring of cis-regulatory regions
controlling a deeply conserved plant stem cell regulator. *PLoS Genetics* **20**, e1011174
(2024).
72   Liu, L. *et al.* Enhancing grain-yield-related traits by CRISPR–Cas9 promoter editing of
maize CLE genes. *Nature Plants* **7**, 287-294 (2021).
73   Cao, R. *et al.* Role of histone H3 lysine 27 methylation in Polycomb-group silencing.
Science **298**, 1039-1043 (2002).
74   Xiao, J. *et al.* Cis and trans determinants of epigenetic silencing by Polycomb repressive
complex 2 in Arabidopsis. *Nature genetics* **49**, 1546-1552 (2017).
75   Schmitz, R. J., Grotewold, E. & Stam, M. Cis-regulatory sequences in plants: Their
importance, discovery, and future challenges. *The plant cell* **34**, 718-741 (2022).

76   Ouyang, W. *et al.* Haplotype mapping of H3K27me3-associated chromatin interactions
defines topological regulation of gene silencing in rice. *Cell Reports* **42** (2023).
77   Minnoye, L. *et al.* Chromatin accessibility profiling methods. *Nature Reviews Methods*
Primers **1**, 10 (2021).
78   Zhang, T.-Q., Chen, Y., Liu, Y., Lin, W.-H. & Wang, J.-W. Single-cell transcriptome
atlas and chromatin accessibility landscape reveal differentiation trajectories in the rice
root. *Nature communications* **12**, 2053 (2021).
79   Yu, Y., Zhang, H., Long, Y., Shu, Y. & Zhai, J. Plant public RNA-seq database: a
comprehensive online database for expression analysis of~ 45 000 plant public RNA-seq
libraries. *Plant Biotechnology Journal* **20**, 806 (2022).
80   Bai, X. *et al.* Duplication of an upstream silencer of FZP increases grain yield in rice.
Nature Plants **3**, 885-893 (2017).
81   Wang, Z. *et al.* AraENCODE: A comprehensive epigenomic database of Arabidopsis
thaliana. *Molecular Plant* **16**, 1113-1116 (2023).
82   Tonosaki, K. & Kinoshita, T. Possible roles for polycomb repressive complex 2 in cereal
endosperm. *Frontiers in plant science* **6**, 144 (2015).
83   Tan, F.-Q. *et al.* A coiled-coil protein associates Polycomb Repressive Complex 2 with
KNOX/BELL transcription factors to maintain silencing of cell differentiation-promoting
genes in the shoot apex. *The Plant Cell* **34**, 2969-2988 (2022).
84   Zhao, L. *et al.* Integrative analysis of reference epigenomes in 20 rice varieties. *Nature*
communications **11**, 2658 (2020).
85   Zhang, J. *et al.* Building two indica rice reference genomes with PacBio long-read and
Illumina paired-end sequencing data. *Scientific data* **3**, 1-8 (2016).
86   Huang, Y. *et al.* OsNCED5, a 9-cis-epoxycarotenoid dioxygenase gene, regulates salt and
water stress tolerance and leaf senescence in rice. *Plant science* **287**, 110188 (2019).
87   Goel, M. *et al.* The vast majority of somatic mutations in plants are layer-specific.
Genome Biology **25**, 1-18 (2024).
88   Andrews, G. *et al.* Mammalian evolution of human cis-regulatory elements and
transcription factor binding sites. *Science* **380**, eabn7930 (2023).
89   Engelhorn, J. *et al.* Phenotypic variation in maize can be largely explained by genetic
variation at transcription factor binding sites. *bioRxiv*, 2023.2008. 2008.551183 (2023).
90   Zhao, T. & Schranz, M. E. Network-based microsynteny analysis identifies major
differences and genomic outliers in mammalian and angiosperm genomes. *Proceedings*
of the National Academy of Sciences **116**, 2165-2174 (2019).
91   Villar, D. *et al.* Enhancer evolution across 20 mammalian species. *Cell* **160**, 554-566
(2015).
92   Reineke, A. R., Bornberg-Bauer, E. & Gu, J. Evolutionary divergence and limits of
conserved non-coding sequence detection in plant genomes. *Nucleic acids research* **39**,
6029-6043 (2011).
93   Kaplinsky, N. J., Braun, D. M., Penterman, J., Goff, S. A. & Freeling, M. Utility and
distribution of conserved noncoding sequences in the grasses. *Proceedings of the*
National Academy of Sciences **99**, 6147-6151 (2002).
94   Guo, H. & Moose, S. P. Conserved noncoding sequences among cultivated cereal
genomes identify candidate regulatory sequence elements and patterns of promoter
evolution. *The Plant Cell* **15**, 1143-1158 (2003).

95   Inada, D. C. *et al.* Conserved noncoding sequences in the grasses4. *Genome Research* **13**,
2030-2041 (2003).
96   Clark, J. W. & Donoghue, P. C. Whole-genome duplication and plant macroevolution.
Trends in plant science **23**, 933-945 (2018).
97   Del Pozo, J. C. & Ramirez-Parra, E. Whole genome duplications in plants: an overview
from Arabidopsis. *Journal of Experimental Botany* **66**, 6991-7003 (2015).
98   Jump, A. S., Marchant, R. & Peñuelas, J. Environmental change and the option value of
genetic diversity. *Trends in plant science* **14**, 51-58 (2009).
99   Wiles, E. T. & Selker, E. U. H3K27 methylation: a promiscuous repressive chromatin
mark. *Current opinion in genetics & development* **43**, 31-37 (2017).
100 Zhang, X., Marand, A. P., Yan, H. & Schmitz, R. J. scifi-ATAC-seq: massive-scale
single-cell chromatin accessibility sequencing using combinatorial fluidic indexing.
Genome Biology **25**, 90 (2024).
101 Zheng, G. X. *et al.* Massively parallel digital transcriptional profiling of single cells.
Nature communications **8**, 14049 (2017).
102 Ouyang, S. *et al.* The TIGR rice genome annotation resource: improvements and new
features. *Nucleic acids research* **35**, D883-D887 (2007).
103 Satija, R., Farrell, J. A., Gennert, D., Schier, A. F. & Regev, A. Spatial reconstruction of
single-cell gene expression data. *Nature biotechnology* **33**, 495-502 (2015).
104 Wolock, S. L., Lopez, R. & Klein, A. M. Scrublet: computational identification of cell
doublets in single-cell transcriptomic data. *Cell systems* **8**, 281-291. e289 (2019).
105 Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *bioinformatics* **26**, 139-
140 (2010).
106 Crow, M., Paul, A., Ballouz, S., Huang, Z. J. & Gillis, J. Characterizing the replicability
of cell types defined by single cell RNA-sequencing data using MetaNeighbor. *Nature*
communications **9**, 884 (2018).
107 Rodriques, S. G. *et al.* Slide-seq: A scalable technology for measuring genome-wide
expression at high spatial resolution. *Science* **363**, 1463-1467 (2019).
108 Stickels, R. R. *et al.* Highly sensitive spatial transcriptomics at near-cellular resolution
with Slide-seqV2. *Nature biotechnology* **39**, 313-319 (2021).
109 Satpathy, A. T. *et al.* Massively parallel single-cell chromatin landscapes of human
immune cell development and intratumoral T cell exhaustion. *Nature biotechnology* **37**,
925-936 (2019).
110 Institute, B. Picard toolkit. *Broad Institute, GitHub repository* (2019).
111 Smit, A. F. Repeat-Masker Open-3.0. *http://www. repeatmasker. org* (2004).
112 Camacho, C. *et al.* BLAST+: architecture and applications. *BMC bioinformatics* **10**, 1-9
(2009).
113 Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome biology* **9**, 1-9
(2008).
114 DeBruine, Z. J., Melcher, K. & Triche Jr, T. J. Fast and robust non-negative matrix
factorization for single-cell experiments. *bioRxiv*, 2021.2009. 2001.458620 (2021).
115 McInnes, L., Healy, J. & Melville, J. Umap: Uniform manifold approximation and
projection for dimension reduction. *arXiv preprint arXiv:1802.03426* (2018).

1432 116 Zong, J. *et al.* A rice single cell transcriptomic atlas defines the developmental
1433   trajectories of rice floret and inflorescence meristems. *New Phytologist* **234**, 494-512
1434   (2022).

1435 117 Hua, L. *et al.* The bundle sheath of rice is conditioned to play an active role in water
1436   transport as well as sulfur assimilation and jasmonic acid synthesis. *The Plant Journal*
1437   **107**, 268-286 (2021).

1438 118 Itoh, J.-I. *et al.* Genome-wide analysis of spatiotemporal gene expression patterns during
1439   early embryogenesis in rice. *Development* **143**, 1217-1227 (2016).

1440 119 Wu, T.-Y., Müller, M., Gruissem, W. & Bhullar, N. K. Genome wide analysis of the
1441   transcriptional profiles in different regions of the developing rice grains. *Rice* **13**, 1-19
1442   (2020).

1443 120 Van Dijk, D. *et al.* Recovering gene interactions from single-cell data using data
1444   diffusion. *Cell* **174**, 716-729. e727 (2018).

1445 121 Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with
1446   Harmony. *Nature methods* **16**, 1289-1296 (2019).

1447 122 Pettkó-Szandtner, A. *et al.* Core cell cycle regulatory genes in rice and their expression
1448   profiles across the growth zone of the leaf. *Journal of plant research* **128**, 953-974
1449   (2015).

1450 123 Schep, A. N. *et al.* Structured nucleosome fingerprints enable high-resolution mapping of
1451   chromatin architecture within regulatory regions. *Genome research* **25**, 1757-1770
1452   (2015).

1453 124 Pattengale, N. D., Alipour, M., Bininda-Emonds, O. R., Moret, B. M. & Stamatakis, A. in
1454   *Research in Computational Molecular Biology: 13th Annual International Conference,*
1455   *RECOMB 2009, Tucson, AZ, USA, May 18-21, 2009. Proceedings 13.* 184-200
1456   (Springer).

1457 125 Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
1458   sequence data. *Bioinformatics* **30**, 2114-2120 (2014).

1459 126 Servant, N. *et al.* HiC-Pro: an optimized and flexible pipeline for Hi-C data processing.
1460   *Genome biology* **16**, 1-11 (2015).

1461 127 Kaul, A., Bhattacharyya, S. & Ay, F. Identifying statistically significant chromatin
1462   contacts from Hi-C data with FitHiC2. *Nature protocols* **15**, 991-1012 (2020).

1463 128 Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing
1464   genomic features. *Bioinformatics* **26**, 841-842 (2010).

1465 129 Schep, A. N., Wu, B., Buenrostro, J. D. & Greenleaf, W. J. chromVAR: inferring
1466   transcription-factor-associated accessibility from single-cell epigenomic data. *Nature*
1467   *methods* **14**, 975-978 (2017).

1468 130 Dijk, D. v. *et al.* MAGIC: A diffusion-based imputation method reveals gene-gene
1469   interactions in single-cell RNA-sequencing data. *BioRxiv*, 111591 (2017).

1470 131 Alpert, A., Moore, L. S., Dubovik, T. & Shen-Orr, S. S. Alignment of single-cell
1471   trajectories to compare cellular expression dynamics. *Nature methods* **15**, 267-270
1472   (2018).

1473 132 Jin, J. *et al.* PlantTFDB 4.0: toward a central hub for transcription factors and regulatory
1474   interactions in plants. *Nucleic acids research*, gkw982 (2016).

1475 133 Grant, C. E., Bailey, T. L. & Noble, W. S. FIMO: scanning for occurrences of a given
1476   motif. *Bioinformatics* **27**, 1017-1018 (2011).

1477	134	Castro-Mondragon, J. A. *et al.* JASPAR 2022: the 9th release of the open-access database of transcription factor binding profiles. *Nucleic acids research* **50**, D165-D173 (2022).
1479	135	Wagner-Menghin, M. M. Binomial test. *Encyclopedia of statistics in behavioral science* (2005).
1481	136	Bolstad, B. M. & Bolstad, M. B. M. Package 'preprocessCore'. (2013).
1482	137	Lovell, J. T. *et al.* GENESPACE tracks regions of interest and gene copy number variation across multiple genomes. *Elife* **11**, e78526 (2022).
1484	138	Sayers, E. W. *et al.* Database resources of the national center for biotechnology information. *Nucleic acids research* **49**, D10 (2021).
1486	139	Shen, W., Le, S., Li, Y. & Hu, F. SeqKit: a cross-platform and ultrafast toolkit for FASTA/Q file manipulation. *PloS one* **11**, e0163962 (2016).
1488	140	Pollard, K. S., Hubisz, M. J., Rosenbloom, K. R. & Siepel, A. Detection of nonneutral substitution rates on mammalian phylogenies. *Genome research* **20**, 110-121 (2010).
1490	141	Siepel, A. & Haussler, D. Phylogenetic estimation of context-dependent substitution rates by maximum likelihood. *Molecular biology and evolution* **21**, 468-488 (2004).
1492	142	Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nature methods* **9**, 357-359 (2012).
1494	143	Li, H. *et al.* The sequence alignment/map format and SAMtools. *bioinformatics* **25**, 2078-2079 (2009).
1496	144	Stovner, E. B. & Sætrom, P. epic2 efficiently finds diffuse domains in ChIP-seq data. *Bioinformatics* **35**, 4392-4393 (2019).
1498	145	Xie, L. *et al.* RiceENCODE: A comprehensive epigenomic database as a rice Encyclopedia of DNA Elements. *Molecular Plant* **14**, 1604-1606 (2021).
1500	146	Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **34**, i884-i890 (2018).
1502	147	Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler transform. *bioinformatics* **25**, 1754-1760 (2009).
1504	148	McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome research* **20**, 1297-1303 (2010).
1507	149	Tian, T. *et al.* agriGO v2. 0: a GO analysis toolkit for the agricultural community, 2017 update. *Nucleic acids research* **45**, W122-W129 (2017).
1509	150	Hofmeister, B. T. & Schmitz, R. J. Enhanced JBrowse plugins for epigenomics data visualization. *BMC bioinformatics* **19**, 1-6 (2018).

Extended Data Fig. 1. Pipeline of cell identity annotation and quality control of rice ACRs.

a, A pipeline of cell identity annotation corresponding to nine strategies shown in Supplementary Note 1.

b-c, A comparison between ACRs (n = 120,048) and control regions (n = 120,048) in terms of SNP density (**b**) and conservation scores (**c**). The SNP data were downloaded from Rice SNP-Seek Database (Mansueto et al. 2017). The SNP density is the number of SNPs per 1,000 bp within the ACRs or control regions. The ACRs exhibited lower SNP density and higher conservation scores compared to a control composed of non-ACR DNA. The control set was created to match each ACR by excluding exons, ACRs, and unmappable regions, and they possess a similar GC content (< 5% average difference) compared to the positive set.

d, Row normalized chromatin accessibility z-scores for ACRs across cell types.

e, Spearman correlation is shown between the total number of Tn5 insertions and number of cell-type-specific ACRs across all cell types. The identification of cell-type-specific ACRs was independent of sequencing depth, as evidenced by the lack of correlation between Tn5 insertions and the number of cell-type-specific ACRs per cell type.

f, A two-dimensional density plot illustrates the median chromatin accessibility across 126 cell types for 120,048 ACRs, along with the range of chromatin accessibility (calculated as the difference between the maximum and minimum values).

g, Categories of distal cell-type-specific leaf ACRs showing potential interaction with genes based on examining chromatin loops derived from rice leaf Hi-C data.

Extended Data Fig. 2. Characterization of QTNs located within cell-type-specific ACRs and pseudotime trajectories.

a, The ratio of non-CDS QTNs overlapping with ACRs to all non-CDS QTNs.

b, A bar plot reveals the non-CDS QTNs significantly overlapped ACRs.

c, Analysis of cell-type-aggregate chromatin accessibility of ACRs surrounding *OsROS1* across endosperm-related cell types and cell types in early seedling tissue. The gray bar highlights endosperm-specific cytosine methylation depletion over endosperm-specific ACRs.

d, Overview of pseudotime developmental trajectory analysis.

e, Relative chromatin accessibility of ACRs and TF genes associated with pseudotime (x-axis). The left side displays three genes neighboring the enriched ACRs along the trajectory gradient.

f, Distribution of motif-to-motif distances (upper panel) and number of motifs (lower panel) from k-mean groups.

Extended Data Fig. 3. Characterization of ACRs across five grass species.

a, Number of syntenic regions identified in all five species. Note that Os was used as the reference due to the inclusion of the atlas, biasing results to this segment of the tree.

b, Enrichment of cell-type-specific ACRs in non-syntenic regions, compared to syntenic regions. Significance testing was performed using a two-sided Fisher's exact test. 'na' indicates that the broadly-accessible ACRs were not enriched in non-syntenic regions for genic, proximal, and distal categories. The percentage for broad and cell-type-specific ACRs within non-syntenic or syntenic regions collectively sum to 100%.

c, ACR count of broadly-accessible and cell-type-specific ACRs within syntenic and non-syntenic regions.

d, A summary plot displaying the different categories of ACRs:
1) Based on their proximity to the nearest gene, ACRs are divided into three groups: genic ACRs (overlapping a gene), proximal ACRs (within 2 kb of genes), and distal ACRs (more than 2 kb away from genes).
2) According to the statistical approach described in the methods section (Identification of cell-type-specific ACRs in leaf tissue across five species), ACRs are classified as either broad ACRs, which are accessible in most cell types (p value >= 0.0001), or cell-type-specific ACRs, which are accessible in a limited number of cell types (p value < 0.0001).
3) Based on their syntenic status, ACRs are grouped into shared ACRs (matching sequences accessible in both species), variable ACRs (matching sequences accessible in only one species), and species-specific ACRs (sequences exclusive to a single species).

e, The distribution of cell-type-specific ACRs across different cell types was characterized by a similar percentage representation.

f, The distribution of distance of ACRs to their nearest genes in broadly accessible and cell-type-specific ACRs across five species.

g, A summary plot displaying criteria to define H3K27me3-broad and H3K27me3-cell-type-specific ACRs. For the analysis of cell types in leaf tissue across rice, sorghum, and maize, where the number of cell types analyzed is fewer than 10, we set criteria such that H3K27me3-broad ACRs must be accessible in at least n-1 cell types, whereas cell-type-specific ACRs should be accessible in fewer than 3 out of the n examined cell types. For seedling and seminal root tissues in rice, where more than 10 cell types are examined, we adjusted the criteria. Here, H3K27me3-broad ACRs are required to be accessible in at least n-2 cell types, and cell-type-specific ACRs should be accessible in fewer than 4 out of the n examined cell types.

Extended Data Fig. 4. Examination of the chromatin accessibility of motifs enriched in specific cell types across species.

a, Spearman correlation between chromatin accessibility of TF genes and motif deviation. Note that for *U. fusca* species, the analysis was omitted due to the limited number of cell types (four) available for spearman correlation, making it statistically unfeasible.

b, Spearman correlation of motif deviation to chromatin accessibility and expression of TF genes in *O. sativa* seedling organ.

c, Mean motif coverage centered on ACRs (red line) and control regions (gray line) for each species, with 95% confidence intervals shown as a shaded polygon.

d, The UMAP panels highlight one TF motif (HDG1) enriched with ACRs in epidermis, where their cognate TFs are known to accumulate.

e, The UMAP panels highlight one representative TF motif underlying the WRKY family enriched with ACRs in epidermis cells, where their cognate TFs are known to accumulate.

f-g, Presenting box (**e**) and heatmap (**f**) plots illustrating the collapsed motif enrichment patterns into TF motif families across various species for each cell type. Each dot within the box represents a TF motif family. The score for each super TF motif family was calculated by averaging the enrichment scores of all the TF motif members within that super family. The DOF TF motif family was highlighted by a red frame in the heatmap. The reduced motif enrichment cutoff setting reveals additional motif families enriched in companion cells and sieve elements, compared to the higher beta cutoff setting as shown in Fig. 3e. It is worth noting that we still observed lower enrichment scores in *O. sativa* compared to other species.

Extended Data Fig. 5. Characterization of ACRs within syntenic regions.

a, The bar plot illustrates the percentage of *O. sativa* ACRs within syntenic and non-syntenic regions when compared to another species.

b, The pie charts depict percentages of broadly accessible (Broad) and cell-type-specific (CT) ACRs within three classes of ACR conservation in Fig. 3b. The total number of ACR in *O. sativa* is below the pie charts.

c, The percentage of broad and cell-type-specific ACRs using *Z. mays*, *S. bicolor*, *P. miliaceum*, or *U. fusca* as the baseline underlying three classes shown in Fig. 3b. The x-axis labels, highlighted by a red frame, are the inverse to x-axis labels in Fig. 3c by using *O. sativa* as a baseline. The significance test was done by using the Fisher's exact test (alternative = 'two.sided'). '**' denotes a *p* value < 0.01, and '*' denotes a *p* value < 0.05.

d, Characterization of ACRs under 'shared ACR' class and their proximity categorizations. **Left,** the number of ACRs under the 'shared ACR' class, categorized into four combinations based on broadly accessible and cell-type-specific ACRs. **Right,** a heatmap of ACR categories based on their proximity to genes: genic ACRs overlapping genes, proximal ACRs (within 2 kb of genes), and distal ACRs (more than 2 kb away from the TSS).

e, The percentage of shared, variable, and species-specific ACR classes in genic, proximal, and distal manners for each species pair. The percentage for each class within genic, proximal, and distal groups collectively sum to 100%. E.g. Genic, proximal and distal broad shared ACRs sum to 100%.

f, The pie charts illustrate percentage of cell-type-specific *O. sativa* ACRs syntenic to cell-type-specific ACRs in corresponding species.

g, The pie charts illustrate mesophyll and bundle sheath specific *O. sativa* ACRs syntenic to cell-type-specific ACRs in corresponding species.

Extended Data Fig. 6. Epidermal cell specific ACRs are less conserved in sequence than other examined cell types.

a, The analysis demonstrates an enrichment of cell-type-specific ACRs based on Fisher's exact test. This test assesses whether cell-type-specific ACRs are more likely to be situated in the 'species-specific' class, compared to other cell types. Statistically significant differences (*p* value < 0.05) in all pairwise comparisons are denoted by distinct letters, determined using Fisher's exact test with the alternative set to 'two.sided'. Bars sharing the same letter indicate that they are not significantly (*p* value > 0.05) different from each other.

b, The proportion of cell-type-specific ACRs identified collectively across all cell types within each species pair under three syntenic classes. The percentage for each cell type within the three classes collectively sum to 100%.

c, This panel has the same meaning as panel **a**, and corresponds to an enrichment test on panel **b**. *U. fusca*-to-*O. sativa* did not include a bar for protoderm as this cell type has not been identified in *U. fusca* scATAC-seq dataset.

d, The proportion of cell-type-specific ACRs identified collectively across all cell types within each species pair under three syntenic classes (share, variable, and species-specific classes). The percentage for each cell type within the three classes collectively sum to 100%. The asterisk indicates three cell types not examined in the comparisons within leaf tissue presented in Fig. 4e.

e, This test assesses whether cell-type-specific ACRs are more likely to be situated in the 'species-specific' class, compared to other cell types. Statistically significant differences (*p* value < 0.05) in all pairwise comparisons are denoted by distinct letters, determined using a Fisher's exact test with the alternative set to 'two.sided'. Bars sharing the same letter indicate that they are not significantly (*p* value > 0.05) different from each other. The asterisk indicates three cell types not examined in the comparisons within leaf tissue presented in Fig. 4e.

f, The proportion of cell-type-specific ACRs identified collectively across all cell types within each species pair under three syntenic classes (share, variable, and species-specific classes). The percentage for each cell type within the three classes collectively sum to 100%.

g, This test assesses whether cell-type-specific ACRs are more likely to be situated in the 'species-specific' class, compared to other cell types. Statistically significant differences (*p* value < 0.05) in all pairwise comparisons are denoted by distinct letters, determined using Fisher's exact test with the alternative set to 'two.sided'. Bars sharing the same letter indicate that they are not significantly (*p* value > 0.05) different from each other.

h, The MetaNeighbor analysis quantifies transcriptome similarity among cell types in *O. sativa* compared to *Z. mays*. 'n=5' denotes five replicates, which were created by randomly dividing the cells from each cell type into five groups, with each group serving as input for the MetaNeighbor analysis. Statistically significant differences (*p* value < 0.05) in all pairwise comparisons are denoted by distinct letters, determined using Wilcoxon signed rank test with the alternative set to 'two.sided'. The groups sharing the same letter indicate that they are not significantly (*p* value > 0.05) different from each other.

i, The percentage of broad and cell-type-specific ACRs using either *Z. mays* (*Zm*) or *S. bicolor* (*Sb*) as the baseline and using either *P. miliaceum* (*Pm*) or *U. fusca* (*Uf*) as the baseline underlying three classes shown in Fig. 4b. The significance test was done by using the Fisher's exact test (alternative = 'two.sided'). '**' denotes a p value < 0.01, and '*' denotes a *p* value < 0.05.

j, The percentage of cell-type-specific ACRs identified across all cell types within each species pair split into three classes shown in Fig. 4b. The percentage for each cell type within the three classes collectively sum to 100%.

Extended Data Fig. 7. Identification of TF motifs and TEs associating with epidermal cell divergence.

a, TF motif enrichment tests were performed in species-specific ACRs per species pair using *O. sativa* as the baseline (See Methods: Binomial test-based motif enrichment analysis). The TF motif with a *q* value more than 0.05 was indicated by filling them with a white color.

b, This panel utilized the same method as panel a to conduct TF motif enrichment tests in syntenic and non-syntenic ACRs in the *O. sativa*-to-*Z. mays* species pair (The results from other species pairs were shown in Supplementary Table 8). The '-Log10 (*q* value)' of each TF family was calculated by averaging '-Log10 (*q* value)' across all individual TF motifs within this family. TF family names were marked besides each dark dot, while the PLINC, HD-ZIP, SBP, and WRKY family were highlighted using blue color. The x-axis indicates the number of TF motifs detected to be enriched (*q* value < 0.05) in the relative TF family.

c, TF motif enrichment tests were performed in species-specific ACRs neighboring *O. sativa* ortholog exhibiting higher expression levels in epidermis based on Binomial test (See Methods: Binomial test-based motif enrichment analysis). Significance testing in violin plot was performed using the t-test (alternative = 'two.sided').

d, GO enrichment test was performed in *O. sativa* orthologous genes based on agriGO (Tian et al. 2017).

e, A screenshot of *LIP1* accessibility in *O. sativa* and *Z. mays* L1 cells which contains an *O. sativa* epidermal specific and species-specific ACR with two ZHD1 motif sites. No corresponding ZHD1 motifs were found in *ZmLIP1*.

f-g, These two plots have the same meanings as panel **d** and **e**, but focusing on species-specific ACRs in *Z. mays*.

h, Enrichment of TE family on ACRs within non-syntenic regions relative to those in syntenic regions. Significance testing was performed using Fisher's exact test (alternative = 'two.sided'). The TE with a *p* value more than 0.05 was indicated by filling them with a white color.

i, TF motif enrichment tests were performed on the epidermis specific ACRs overlapping with LTR *Gypsy* TEs based on Binomial test (See Methods: Binomial test-based motif enrichment analysis). The TE with a *p* value more than 0.05 was indicated by filling them with a white color.

Extended Data Fig. 8. Examination of *O. sativa* atlas ACRs corresponding to variable *Z. mays* ACRs.

a, Percentage of ACRs overlapping CNS. The significance test was done by using the Binomial test (** indicate p value < 0.01; alternative = 'two.sided'). We generated control sets by simulating sequences with the same length as ACRs 100 times, yielding a mean proportion for the control sets. The binomial test p value was calculated by comparing the mean ratio to the observed overlapping ratio of ACRs capturing the CNS.

b, Three classes depicting variations in ACR conservation between two species. 'Shared CNS ACRs': CNS ACRs with matching sequences that are accessible in both species; 'Variable CNS ACRs': CNS ACRs with matching sequences, but are only accessible in one species; 'Species-specific CNS ACRs': CNS ACRs where the sequence is exclusive to a single species.

c, The pie charts depict percentages of CNS ACRs within three classes of ACR conservation in panel a. The total number of ACR in *O. sativa* is below the pie charts.

d, Comparison of length of alignment regions and motif count within shared CNS ACRs between with same and different cell-type specificity

e, A sketch illustrating whether variable ACRs containing CNS in *Z. mays* capture ACRs derived from the *O. sativa* atlas. The left pie chart panel represents the percentage of *Z. mays* leaf CNS ACRs that were accessible in non-leaf cell types in the *O. sativa* atlas.

f, A density plot illustrating the number of non-leaf *O. sativa* cell-types in which an *O. sativa* ACR syntenic to *Z. mays* variable ACRs are accessible.

g-i, The bar plots represent the count of *O. sativa* atlas cell-type-specific ACRs accessible in non-leaf cell states from seminal root, panicle, and early seed organs. These ACRs were derived from cell-type-specific ACRs from Fig. 4e.

j, The percentage of non-syntenic ACRs capturing CNS. All pairwise comparisons are statistically significant (p value < 0.05) as indicated by different letters based on t-test (alternative = 'two.sided'). If two bars have the same letter, then they are equivalent (p value > 0.05).

Extended Data Fig. 9. Assessment of transcriptional state of genes in proximity to H3K27me3-broad-ACRs.

a, Pie charts showcasing the composition of different categories of ACRs, and bar plots displaying the number of ACRs within the H3K27me3-broad-ACR group.

b, No significant difference in the distance to nearby genes was identified between H3K27me3-broad ACRs to H3K27me3-absent ACRs as shown in Fig. 5c.

c, Comparative analysis of expression levels and chromatin accessibility of genes surrounding broad ACRs under and outside of H3K27me3 peaks. ** indicate *p* value < 0.01, which was performed by the Wilcoxon signed rank test (alternative = 'two.sided').

d, Four categories of gene sets overlapped with H3K27me3 peaks and comparisons of their expression between bulk RNA-seq data libraries from two different organs. The 'n' above the box indicates the number of libraries used in this comparison. Significance testing was performed using the Wilcoxon signed rank test (alternative = 'two.sided').

e, Percentage of H3K27me3-broad ACRs nearby genes exhibiting no expression across any cell type was significantly higher than the other ACRs associated with the not expressed genes. Significance testing was performed using Fisher's exact test (alternative = 'two.sided').

f, Percentage of H3K27me3-broad ACRs nearby genes exhibiting cell-type-specific expression was significantly higher than the other genes not associating with H3K27me3-broad-ACRs. Significance testing was performed using Fisher's exact test (alternative = 'two.sided').

Extended Data Fig. 10. Characterization of H3K27me3-broad ACRs potentially enriched for silencer CREs.

a, A screenshot illustrates a H3K27me3-broad ACR harboring a reported silencer within an H3K27me3 peak in the panicle organ, located approximately 5.3 kb upstream of *FZP* gene. SBM: Secondary Branch Meristem. PBM: Primary Branch Meristem. SM: Spikelet Meristem. FM: Floral Meristem. le: Lemma. lo: Lodicule. pa: Palea. rg: Rudimentary glume. sl: Sterile lemma.

b, The number of H3K27me3-broad-ACR capturing known 53 PREs is determined based on *A. thaliana* scATAC-seq data. The significance test was done by using the Binomial test (alternative = 'two.sided'; See Methods: Construction of control sets for enrichment tests).

c, Percentage of H3K27me3 and H3K27me3 absent in broad ACRs that contain PRE-associated motifs.

d, Pie charts illustrate the different types of PRE-associated motifs within H3K27me3-broad and H3K27me3-absent-broad ACRs.

e, Distribution of PRE motif count per ACR for H3K27me3-broad and H3K27me3-absent-broad ACRs categories.

f, Percentage of EMF2b peaks in *O. sativa* capturing six known motifs enriched in PREs in *A. thaliana*.
** indicate *p* value < 0.01, which was performed by the Binomial test (alternative = 'two.sided'; See Methods: Construction of control sets for enrichment tests).

g, Alignment of H3K27me3 and EMF2b chromatin attributes at summits of distinct ACR groups in *O. sativa*. CT ACRs mean cell-type-specific ACRs, other ACR means ACRs not classified as the broad or CT ACRs according to the classification criteria in Extended Data Fig. 3g.

h, Comparison of expression of genes nearby H3K27me3-broad ACRs that contain PRE motifs with SNPs in 'Zhenshan 97' (ZS97) genotype using 'Nipponbare' (NIP) as the reference. Significant test was performed by the Wilcoxon signed rank test (alternative = 'two.sided'). CPM: Counts Per Million.

i, Alignment of H3K27me3 chromatin attributes at summits of transcription start site (TSS) of genes derived from panel h. The control genes include genes overlapping with H3K27me3, which are shared between both genotypes.

j, A screenshot illustrates an H3K27me3-broad ACR containing a PRE-associated motif with an SNP situated at 1.2 kb upstream of *LOC_Os11g08020* gene, which might be associated with a lower H3K27me3 signal and higher expression of the *LOC_Os11g08020* in the 'Zhenshan 97' genotype.

k, Four TF families, highlighted in red, were significantly enriched in H3K27me3-broad ACRs. The motif data were collected from 568 TFs from *A. thaliana* belonging to 24 families within the JASPAR database (Castro-Mondragon et al. 2022). The *p* value was computed using a hypergeometric test (alternative = 'two.sided').

Extended Data Fig. 11. Evolutionary dynamics of H3K27me3-broad ACRs across different species.

a, An example of a syntenic region containing H3K27me3-broad ACRs that were conserved across *O. sativa*, *Z. mays*, and *S. bicolor*.

b, The pie charts depict percentages of H3K27me3-broad ACRs within three classes of ACR conservation in Fig. 3b.

c, Three categories of *O. sativa* H3K27me3-broad ACRs that are syntenic to regions from another species.

d, Comparison of ACR percentage underlying three classes corresponding to panel **c** associated with H3K27me3 or not. The *p* values displayed in the bottom bar plot were computed using Fisher's exact test (alternative = 'two.sided').