Title:

Identification of KIFC1 as a putative vulnerability in lung cancers with centrosome amplification

Authors:

Christopher Zhang[1,2]*, Benson Wu[1,2]*, Yin Fang Wu[2], Caterina di Ciano-Oliveira[2], Isabel Soria-Bretones[3], Nhu-An Pham[3], Andrew J. Elia[3], Raj Chari[4], Wan L. Lam[5,6], Mark R. Bray[3], Tak W. Mak[3,7,8], Ming-Sound Tsao[1,3,8], David W. Cescon[3,9], Kelsie L. Thu[1,2]
*These authors contributed equally to this work

Affiliations:

[1] Department of Laboratory Medicine and Pathobiology, University of Toronto
[2] Keenan Research Centre for Biomedical Science, St. Michael's Hospital, Unity Health Toronto
[3] Princess Margaret Cancer Centre, University Health Network
[4] Laboratory Animal Sciences Program, Frederick National Laboratory for Cancer Research
[5] Department of Integrative Oncology, BC Cancer Research Institute
[6] Department of Pathology and Laboratory Medicine, University of British Columbia
[7] Departments of Immunology University of Toronto
[8] Departments of Medical Biophysics, University of Toronto
[9] Institute of Medical Science, University of Toronto

Competing Interests Statement:
Christopher Zhang, Benson Wu, Yin Fang Wu, Caterina di Ciano-Oliveira, Nhu-An Pham, Andrew J. Elia, Raj Chari, Wan L. Lam, Ming S. Tsao, Kelsie L. Thu - none
Isabel Soria-Bretones - ISB is currently an employee of Repare Therapeutics
Mark Bray - MRB is currently an employee of Treadwell Therapeutics
Tak Mak - Owns equity in Treadwell Therapeutics Inc. and Agios Pharmaceuticals and is a consultant for AstraZeneca and Tessa Therapeutics
David Cescon - DWC reports consultancy and advisory relationships with AstraZeneca, Daiichi Sankyo, Exact Sciences, Eisai, Gilead, GlaxoSmithKline, Inflex, Inivata/NeoGenomics, Lilly, Merck, Novartis, Pfizer, Roche and Saga; research funding to their institution from AstraZeneca, Guardant Health, Gilead, GlaxoSmithKline, Inivata/NeoGenomics, Knight, Merck, Pfizer, ProteinQure and Roche

Word Count:

4,492

Abstract

Centrosome amplification (CA), an abnormal increase in the number of centrosomes in the cell, is a recurrent phenomenon in lung and other malignancies. Although CA contributes to tumor development and progression by promoting genomic instability (GIN), it also induces mitotic stress that jeopardizes cellular integrity. The presence of extra centrosomes leads to the formation of multipolar mitotic spindles prone to causing lethal chromosome segregation errors during cell division. To sustain the benefits of CA, malignant cells are dependent on adaptive mechanisms to mitigate its detrimental consequences, and these mechanisms represent therapeutic vulnerabilities. We aimed to discover genetic dependencies associated with CA in lung cancer. Combining a CRISPR/Cas9 functional genomics screen with analyses of tumor genomic data, we identified the motor protein KIFC1 as a putative vulnerability specifically in lung cancers with CA. KIFC1 expression was positively correlated with CA in lung adenocarcinoma (LUAD) cell lines and with a gene expression signature predictive of CA in LUAD tumor tissues. High *KIFC1* expression was associated with worse patient outcomes, smoking history, and indicators of GIN. KIFC1 loss-of-function sensitized LUAD cells to potentiation of CA and sensitization was associated with a diminished ability of KIFC1-depleted cells to cluster extra centrosomes into pseudo-bipolar mitotic spindles. Our work suggests that KIFC1 inhibition represents a novel approach for potentiating GIN to lethal levels in LC with CA by forcing cells to divide with multipolar spindles, rationalizing the clinical development of KIFC1 inhibitors and further studies to investigate its therapeutic potential.

INTRODUCTION

Lung cancer (LC) is the leading cause of cancer death worldwide[1] and is broadly divided into small-cell lung cancer (SCLC) and non-small cell lung cancer (NSCLC). Of the various NSCLC subtypes, lung adenocarcinoma (LUAD) is most prevalent, accounting for ~40% of all cases[2]. Although advances in targeted therapies and immunotherapy have expanded the treatment landscape for LC[3], the 5-year net survival rate has not risen above 22%[4]. This highlights the need for new therapeutic approaches that target mechanisms supporting lung tumor growth and progression.

Centrosomes are microtubule-organizing organelles with important roles in cell motility and polarity, and generation of mitotic spindles required to segregate chromosomes during mitosis[5]. Centrosome amplification (CA), an abnormal increase in the number of centrosomes in a cell, is a common feature of many malignancies including LC[5–7]. Several routes to CA have been described including DNA damage, failure of cytokinesis, centriole overduplication, and carcinogen exposure[8–11]. The presence of extra centrosomes has been shown to promote genomic instability (GIN)[12, 13] which fuels tumor adaptation to stress such as that induced by anti-cancer therapy[14]. This fitness advantage may explain the prevalence of CA in cancer; however, having supernumerary centrosomes comes with a cost. Cells with CA are prone to forming multipolar mitotic spindles that if left uncorrected cause severe chromosome segregation errors, mitotic catastrophe, and cell death[15]. Thus, cancer cells must manage extra centrosomes to prevent multipolar divisions and their deleterious effects. Several coping mechanisms have been proposed with clustering of extra centrosomes into pseudo-bipolar spindles being the most well established[5, 12, 16]. As such, the proteins regulating these adaptive mechanisms represent vulnerabilities and potential therapeutic targets in cancer cells with CA.

Multiple reports have demonstrated the occurrence and clinical relevance of CA in LC[6, 7, 17].

However, despite its occurrence, the mechanisms LC use to cope with CA have not been directly

investigated. We sought to identify collateral dependencies associated with the presence of extra

centrosomes in LC models. Integrating a CRISPR/Cas9 screen with tumor genomic analyses, we

identified KIFC1 as a putative survival factor for LUAD with CA. We discovered that KIFC1 inactivation

is lethal specifically in LC cells with CA. Our analyses of cell lines and independent tumor datasets

revealed that KIFC1 expression is positively correlated with CA in LUAD suggesting centrosome

number could indicate LC sensitivity to KIFC1 inhibition. We found that KIFC1 depletion sensitized

LUAD cells to potentiation of CA and that this phenotype was associated with a diminished ability to

cluster extra centrosomes. Collectively, these findings nominate KIFC1 as a therapeutic target in LUAD

with CA, rationalizing the development of KIFC1-specific inhibitors and further preclinical investigation

of its therapeutic potential.

METHODS

Tumor and cell line genomic analyses

RNA-sequencing (RNASeqV2) and associated clinical data for The Cancer Genome Atlas (TCGA) Pan-

Cancer (PANCAN) cohort were accessed from the National Cancer Institute's Genomic Data Commons

portal[18]. Mutation status for the most common oncogenic drivers (*EGFR, KRAS, MET, ROS1, RET,*

ALK, NRAS and BRAF) in TCGA LUAD tumors was obtained from cBioPortal[19]. Gene expression

and essentiality data for the Cancer Cell Line Encyclopedia (CCLE) were obtained from the DepMap

database (https://depmap.org/portal/)[20]. Normalized microarray-derived gene expression data for 4

additional LUAD cohorts were obtained from the Gene Expression Omnibus (GSE50081, GSE31210,

GSE68465, GSE75037). CA20 gene expression signature scores for TCGA lung tumors and CCLE cell

lines were obtained from a published study[21], and were calculated for additional cohorts as described[22].

Immunohistochemistry

Tumor tissues were banked after informed written patient consent following a protocol approved by the University Health Network Research Ethics Board. Formalin-fixed paraffin embedded tissues from 10 NSCLC patient-derived xenograft (PDX) models were processed for immunohistochemistry to detect centrosomes using the anti-pericentrin antibody ab4448 (Abcam, 1:1000 dilution). Mitotic cells were classified as having CA if they contained more than 2 centrosomes and the proportion with CA was calculated for each tumor. At least 30 mitotic cells were counted per tumor tissue section.

Cell culture and lentivirus production

Cell lines were grown at 37°C and 5% CO_2 in DMEM, Hams F12 or RPMI 1640 media supplemented with 10% fetal bovine serum, 2mM glutamine, and 100U/ml of penicillin-streptomycin. NCI-H1299, -H1975, -H1944, -H2291, -H1792, -H2122, -H23, -H2030, -H1568, -H1650, HCC827, HCC2935 were generous gifts from the late Dr. Adi Gazdar. PC9 was gifted from Dr. William Lockwood and A549 was gifted from Dr. Haibo Zhang. Cells were confirmed to be negative for mycoplasma using a PCR-based assay[23]. Platinum-E (PE) cells were obtained from Cell BioLabs and used to produce lentivirus by transfection with psPAX2 and pMD2.G (Addgene #12260, #12259) using Lipofectamine 3000.

CRISPR/Cas9 genetic screen

H1299 cells were engineered with stable Cas9 expression using Lenti-Cas9-blast (Addgene #52962). Efficient nuclease activity was confirmed using a reporter system (Addgene #67979, #67980; **Supp.Fig.1**). We synthesized a custom single guide RNA (sgRNA) knockout (KO) library with 13,243 sgRNA targeting 3,319 genes that are "druggable" with FDA-approved or investigational compounds annotated in the DrugBank and bindingDB databases[24, 25], as well as negative controls (sgLacZ, sgLuciferase, sgGFP) (**Supp. Table 1**). This library was cloned into pRC0162 (Addgene #195319) and sequence verified. H1299-Cas9 cells were infected with the library at a multiplicity of infection of 30% for single integrations per cell to yield 500X library coverage. Following 48hr of puromycin selection, cells were pooled and split into 3 replicates each for treatment with vehicle (DMSO) or 7.5nM of CFI-400945 (945). Ten million cells were collected at time zero (T0) and the remainder were cultured in the presence of DMSO or 945 for 17 cell doublings and then harvested for targeted sgRNA sequencing. Screens hits were identified using the DrugZ algorithm[26].

Competition Assays

H1975, PC9, and A549 were engineered to express functional Cas9 as above. sgRNA targeting AAVS1 (AV, negative control) or KIFC1 (K3, K4) were cloned into LentiGuide-puro-NLS-GFP or LentiGuide-puro-NLS-mCherry (#185473, 185474) and transduced into Cas9+ cells. sgRNA sequences are provided in **Supp. Table 2**. mCherry+ AV (wildtype, WT) were mixed 1:1 with GFP+ K3/K4/AV cells to compare the relative fitness of cells with and without KIFC1 perturbation. Cells were treated with DMSO or 7.5nM 945 and mCherry/GFP+ fractions were measured by flow cytometry over multiple passages.

KIFC1 knockdown, dose response, and viability assays

Human Silencer Select siRNA assays s7906, s7907, and Negative Control No. 1 (Thermo Fisher

Scientific) were used to knockdown (KD) KIFC1. Cells were transfected with siRNA using

Lipofectamine RNAiMAX (50nM in H1299, H1975, A549; 75nM in PC9) and 24 hours later, were

sampled to confirm KD and plated for functional experiments. Dose response, clonogenic survival, and

AZ82 cytotoxicity assays were done using sulforhodamine B staining as previously described[27].

Proliferation was assessed using the ATP-Lite assay.

Immunofluorescence, immunoblotting, and real-time quantitative PCR

Immunofluorescence (IF) was used to detect centrosomes using the following antibodies: CEP192

(1:1000, Bethyl Labs#A302-324A)[28], α-tubulin (1:1000, Sigma-Aldrich #T9026), goat anti-mouse

AlexaFluor-488, and goat anti-rabbit AlexaFluor-555. DNA was stained with DAPI. CA was calculated

as described for IHC above. Western blots were done using the following antibodies: KIFC1 (1:1000,

CST#12313), GAPDH (1:5000, CST#97166), and β-ACTIN (1:5000, Proteintech#66009-1-IG). RT-

qPCR to confirm siRNA-mediated *KIFC1* knockdown was done using *HPRT1* as the endogenous

control. Primer sequences are listed in **Supp. Table 2**.

Statistics

Statistical analyses are described in the figure legends and were performed using R and GraphPad Prism.

RESULTS

Centrosome amplification (CA) is prevalent in NSCLC tissues

Previous studies have reported centrosome abnormalities in 29-53% of NSCLC tissues[6, 7]. We

evaluated CA in 10 patient-derived xenograft (PDX) models of NSCLC (4 LUAD, 3 squamous cell

carcinoma (LUSC), and 3 large cell carcinoma) using IHC to detect the established centrosome marker,

pericentrin[29]. Cases were chosen to represent different NSCLC subtypes. Our assessment of these

tumors confirmed that CA is prevalent in NSCLC **(Fig. 1A)**. CA was detected in mitotic cells from all

10 PDX, with the percentage of cells with CA ranging from 18-66% **(Fig. 1B)**. We observed that

centrosomes in cells with CA were disorganized in multipolar spindles or clustered into pseudo-bipolar

spindles[12, 16] **(Fig. 1A)**. The presence of CA in clinical NSCLC tissues rationalized investigation of

vulnerabilities associated with extra centrosomes in cell models.

CRISPR screen identifies KIFC1 as a putative CA-coping factor in LC

To discover genetic dependencies in LC cells with CA, we performed a CRISPR/Cas9 screen in the

LUAD cell line, H1299. We selected H1299 because of its high Cas9 editing efficiency (**Supp. Fig. 1A**),

its relatively high basal CA which implied adaptation to CA (**Fig.1C; Supp. Fig. 4C**), and its amenability

to potentiation of CA with low doses of the PLK4 inhibitor, CFI-400945 (945)[30] (**Supp. Fig. 1B**).

H1299-Cas9 cells were infected with a custom sgRNA KO library targeting ~3,300 "druggable" genes,

grown in the presence of DMSO or sublethal dose of 945 (7.5nM) that potentiated CA for 17 cell

doublings (**Supp. Fig. 1C**), and then harvested for sgRNA sequencing. We first compared sgRNA

abundance in DMSO-treated cells at end point to cells harvested at screen onset (T0) to confirm the

depletion of sgRNA targeting essential genes. This identified *MYC* as the top hit with 7 other genes

classified by the DepMap as "common essential genes" ranking in the top 10 hits (*CDC25, SRSF1, RPL4,*

PSMA3, RPL14, ATP2A2, and *SCD*)[20] (**Supp. Fig. 1D-E**). Notably, *NRAS*, an established oncogenic

driver in H1299[31] was identified as the 10th ranked hit **(Supp. Fig. 1D-E)**. Detection of these essential

genes confirmed the technical success of our CRISPR/Cas9 screen.

Sampling of H1299 cells during the screen confirmed that 7.5nM 945 potentiated CA to ~65% of cells and increased the incidence of multipolar spindles compared to DMSO (**Fig.1C-E**). Accordingly, growth in the presence of 945 was slower (**Supp. Fig. 1C**). We reasoned that cells containing sgRNA targeting genes essential for survival in the presence of extra centrosomes would be selectively depleted in populations treated with 945 compared to DMSO. Analysis of sgRNA abundance in 945- versus DMSO- treated populations identified 16 putative CA survival genes with an FDR q-value < 0.25 (**Fig.1F; Supp.Table 3**). To prioritize hits from our screen, we examined their correlations with a gene expression signature of CA called CA20 in the TCGA LUAD cohort[32]. CA20 is a composite of 20 genes associated with the presence of CA that was validated in clinical breast tumors[21, 22]. We hypothesized that genes important for coping with extra centrosomes would be highly expressed in tumors with CA and positively correlated with CA20 scores. Of the 16 candidates, none were CA20 signature genes and KIFC1 exhibited the strongest positive correlation with CA20 in LUAD (**Fig.1G**). KIFC1 is a minus-end directed kinesin motor protein involved in mitotic spindle assembly and organization, as well as vesicle and organelle trafficking[33]. It also has a well-described role in centrosome clustering in cancer cells[33]. Thus, our CA sensitization screen suggested that H1299 cells are dependent on KIFC1 to cope with CA, nominating KIFC1 as a therapeutic vulnerability in LC with supernumerary centrosomes.

KIFC1 is associated with clinical factors in LUAD

We next aimed to determine the clinical relevance of KIFC1 in patient-derived tumor tissues by analyzing KIFC1 mRNA expression in the TCGA LUAD cohort. Consistent with previous reports in pan-cancer or LC-focused studies [34–37], KIFC1 was significantly overexpressed in LUAD compared to non-malignant lung tissues (**Fig.2A**), and LUAD patients whose tumors had high KIFC1 expression had significantly worse 5 year survival outcomes than those with low expression in a univariate model (**Fig.2B**). KIFC1 expression remained significantly associated with overall survival in a Cox-proportional hazards model including additional prognostic factors including age, stage, sex and smoking

history **(Table 1)**. Expression was also elevated in tumors of greater stage, and for the first time, we

discovered that expression was significantly higher in smokers relative to non-smokers (**Fig.2C-D**).

Lastly, we found that KIFC1 expression was modestly elevated in males relative to females and in tumors

without oncogenic driver mutations compared to those with oncogenic alterations (**Supp. Fig.2A**). An

ANOVA suggested that smoking history was the clinical factor most strongly associated with KIFC1

expression **(Table 2)**. Although KIFC1 was overexpressed and correlated with CA20 in LUSC, we did

not observe any significant associations between KIFC1 expression and patient characteristics or survival

(**Supp. Fig. 3**). We then investigated the robustness of these associations in 4 independent LUAD cohorts

with gene expression and clinical data available (GSE75037, GSE50081, GSE31210, GSE68465) (**Supp.**

Fig.2B-F)[38–41]. These analyses confirmed that high KIFC1 expression was associated with smoking

(3/4 cohorts), advanced stage (4/4 cohorts), and worse 5-year survival (4/4 cohorts). Together these

findings indicate the clinical and prognostic significance of KIFC1 in LUAD.

CA and GIN correlate with KIFC1 expression in LUAD models and tissues

Since our screen nominated KIFC1 as a genetic dependency of LC cells with CA, we suspected KIFC1

expression would be higher in cells with extra centrosomes. To test this hypothesis, we evaluated KIFC1

protein expression and CA in a panel of LUAD cell lines. Both KIFC1 expression and CA were variable

across models (**Supp. Fig. 4A-C**). In support of our hypothesis, we identified a positive correlation

between KIFC1 protein expression and CA across LUAD cell lines (**Fig.3A**). We next evaluated the

association between CA20 and KIFC1 dependency scores from the DepMap database in a larger panel

of LC cell lines from the Cancer Cell Line Encyclopedia (CCLE) collection[20]. Our analyses revealed

no association between CA20 and KIFC1 essentiality across 52 NSCLC or 13 LUSC lines, but indicated

trends towards a greater dependency on KIFC1 in 15 LUAD and 11 small-cell lung cancer cell lines

predicted to have high levels of CA (**Fig.3B-C; Supp.Fig4D-E**). Thus, KIFC1 may be a LUAD-specific

dependency. As noted above (**Fig.1G**), we observed a strong positive correlation between KIFC1

expression and CA20 in LUAD tumors from the TCGA (**Fig.3D**), and we validated this correlation in 4

independent cohorts (**Supp.Fig.2E**). Because CA drives genomic instability (GIN)[13, 42, 43], we also

investigated associations between KIFC1 expression and measures of GIN in the TCGA LUAD cohort.

This revealed significant positive correlations between KIFC1 and fraction of genome altered and

aneuploidy scores[18] (**Fig.3E-F**). Collectively, these findings demonstrate that KIFC1 expression is

associated with CA and GIN in LUAD models and tissues.

LUAD cells are dependent on KIFC1 for survival upon CA potentiation

We next sought to functionally validate KIFC1 as a dependency of LC cells with CA using *in vitro*

growth competition assays that mimic CRISPR screen conditions. We used models with different

endogenous levels of KIFC1 and CA and examined the effects of KIFC1 loss-of-function (LOF) on

cellular fitness. H1299 and H1975 were selected as representative models with relatively high basal

KIFC1 expression and CA, and A549 and PC9 as models with low KIFC1 and CA (**Supp. Fig.4**).

Treatment with low-dose 945 potentiated CA and increased the proportion of multipolar spindles in each

LUAD line except A549 (**Fig.4A-B**). Cas9+ derivatives of these lines were transduced with a GFP+

lentivector encoding independent sgRNA targeting KIFC1 (K3 or K4), or a GFP+ or mCherry+

lentivector encoding an sgRNA targeting AAVS1 (AV, wildtype (WT) control). CRISPR-mediated

KIFC1 inactivation was evident in all models relative to controls (**Fig.4C**). We competed GFP+ WT or

GFP+ K3/K4 cells with mCherry+ WT cells in the presence of DMSO or 945 and monitored the

GFP:mCherry+ ratio over several passages. These experiments validated KIFC1 as a hit from our screen,

as GFP+ cells were depleted in the 945-treatment condition over time (**Fig.4D**), confirming that KIFC1

LOF sensitized H1299 cells to CA potentiation. We observed the same phenotype in H1975 although the

effect was less prominent, possibly due to H1975's lower endogenous KIFC1 expression and levels of

CA (**Supp.Fig.4**). We also found that KIFC1 LOF reduced the fitness of H1299 cells treated with DMSO

(**Fig.4D**), indicating that KIFC1 is required to compensate for the relatively high basal CA in the H1299

model. KIFC1-perturbation did not reduce the fitness of PC9 or A549 cells with relatively low CA and KIFC1 expression (**Fig.4D**). These results implicate KIFC1 as a specific dependency in LUAD cell lines with high basal CA (H1299, H1975). The lack of sensitization in PC9 suggests this model uses KIFC1-independent mechanisms to cope with CA induction, while the lack of effect in A549 likely reflects their resistance to CA induction.

Depletion of KIFC1 impairs cell viability and suppresses centrosome clustering in LUAD cell lines with CA

Although multiple compounds have been reported to inhibit KIFC1[44–46], their weak potencies require treating cells at high doses that preclude attributing treatment-induced phenotypes specifically to KIFC1 inhibition[47]. We investigated the effects of AZ82, the KIFC1 inhibitor reported to have the best specificity[47], on LUAD viability. We observed cytotoxic effects of AZ82 at 7.5uM but they did not correlate with KIFC1 expression (**Supp.Fig.5**). For example, PC9 exhibited the greatest sensitivity to AZ82 in contrast to the results of our CRISPR screen validation experiments (**Fig.4D; Supp.Fig.5**). As such, we used two independent siRNAs to knockdown (KD) KIFC1 to investigate its therapeutic potential. Both siRNA yielded effective KD in all 4 models (**Fig.5A-B**). Dose response curves indicated that KIFC1 depletion enhanced sensitivity to 945 in H1299 (**Fig.5C**). A similar albeit less pronounced effect was observed in H1975 and PC9, and KIFC1 KD had no effect on the sensitivity of A549 to 945. We also conducted clonogenic survival assays to measure the effects of KIFC1 KD on cell viability. KIFC1 KD significantly reduced colony survival in H1299 and H1975 when CA was potentiated by 945 treatment, but had no effect in 945-treated PC9 and A549 cells (**Fig.5D**). Depletion of KIFC1 alone had no effect on cell proliferation or clonogenic survival (**Supp.Fig.6A-B**).

CA can lead to the formation of multipolar spindles during mitosis (**Fig.1D-E**) and KIFC1 counters their deleterious effects by clustering extra centrosomes to promote viable chromosome segregation and cell

division. We suspected that the dependence of LUAD cells with CA on KIFC1 was due to its role in

centrosome clustering, so we evaluated the effects of KIFC1 KD on spindle polarity (**Fig.5E**). KIFC1

depletion in H1299 led to a decrease in centrosome clustering evident by a reduced proportion of mitotic

cells with pseudo-bipolar spindles and an increased proportion of cells with multipolar spindles. This

effect was more pronounced when CA was potentiated with 945 treatment. KIFC1 KD did not affect

centrosome clustering in H1975, PC9, or A549 cells treated with DMSO. Like H1299, 945 treatment

reduced pseudo-bipolar and increased multipolar spindles in H1975 (**Fig.5E**). KD of KIFC1 in PC9 and

A549 increased the proportion of multipolar spindles in 945-treated cells but this phenotype was not

reproduced with both siRNAs and the magnitude of change was less than that observed for H1299 and

H1975 (**Fig.5E**). These data suggest that KIFC1 promotes survival of LUAD cells with CA by clustering

extra centrosomes into pseudo-bipolar spindles to avoid the lethal consequences of cell divisions with

multipolar spindles.

DISCUSSION

Centrosome amplification (CA) is a recurrent feature of many malignancies that promotes invasiveness

and genomic instability (GIN), and consequently, cancer development and progression[12, 13, 42, 43,

48]. Not surprisingly, CA is associated with poor prognosis in many cancer types[17, 21, 29, 49]. Despite

the prevalence of CA in lung tumors[6, 7, 17], the vulnerabilities imposed by CA in LC and its therapeutic

potential have not been defined. By integrating a CRISPR/Cas9 screen, genomic analyses of LUAD

tumors, and functional studies in multiple cell lines, we identified and validated *KIFC1* as a dependency

of LUAD with supernumerary centrosomes. Kinesin family member C1 (KIFC1 or HSET) belongs to

the kinesin-14 family of minus-end directed motor proteins that facilitate organization of the mitotic

spindle[33]. Both the tail and motor domains of KIFC1 bind to microtubules, allowing it to crosslink

microtubules and slide them apart or together to organize spindle poles[50, 51]. Although KIFC1 is not

essential in non-malignant cells, it is indispensable in some cancer cell lines[52, 53]. Studies of multipolar

divisions in cells with CA have demonstrated that KIFC1 clusters extra centrosomes into spindles with

pseudo-bipolar arrangements[42, 52, 54, 55]. This action is thought to occur by KIFC1 ratcheting

microtubules emanating from different centrosomes to group them together[54].

Our discovery of KIFC1 as a putative vulnerability in LC is consistent with reports in breast, prostate,

ovarian, liver, and renal cancers[45, 52, 56–58]. Liu and colleagues investigated KIFC1 using functional

studies in 3 NSCLC lines and reported that KIFC1 depletion reduced viability and induced G2/M arrest

and p21 expression but they did not evaluate centrosomes[35]. However, they reported that KIFC1 KD

reduced viability to the greatest extent in H1299 cells, which we found to have relatively high basal CA

among the LUAD models we assessed. Advancing on correlative studies that did not consider centrosome

number, our unique CA-focused approach nominated KIFC1 as a dependency specifically in LUAD

models with high basal levels of CA, likely due to its established role in mitigating CA-associated mitotic

stress via centrosome clustering[45, 52, 56–58]. Patel *et al*. identified KIFC1 as a putative vulnerability

in breast cancers based on integrative genomic analyses of tissues and cell lines. Similar to our finding,

this group showed that KIFC1 was preferentially essential in triple-negative breast cancer models with

CA[57]. Although the CA20 gene expression signature is only a surrogate for CA, our assessment of

KIFC1 dependency and CA20 in the DepMap database[20] further suggested that KIFC1 is a CA-specific

dependency in LUAD and possibly SCLC. KIFC1's classification as a strongly selective dependency,

but not a common essential gene, in DepMap likely indicates that a small fraction of cancer models with

high basal CA are dependent on KIFC1.

Liu *et al*. also reported that KIFC1 was overexpressed in NSCLC compared to non-malignant lung tissues

and that high expression was associated with worse prognosis, which is consistent with findings from

other LC studies[35–37, 59, 60] and reports in other malignancies[34, 36, 61–65]. Our analyses of

multiple LC cohorts confirmed associations of KIFC1 with clinical factors including survival and stage

as well as CA20. For the first time, we found that KIFC1 expression was elevated in LUAD from patients

with a smoking history. This observation was reproducible across multiple cohorts, raising the possibility

that tobacco smoke leads to increased KIFC1 expression as a consequence of inducing CA[66]. Celik *et*

al. recently reported that KIFC1 was overexpressed in 59% of lung tumors and that 96% of these

exhibited hypomethylation of KIFC1[59]. This suggests that recurrent overexpression of KIFC1 in LC

is driven by epigenetic mechanisms, but this remains to be validated in additional cohorts. An integrative

pan-cancer study by Wu *et al*. found that KIFC1 was rarely perturbed at the genetic level in LUAD and

other tumor types, suggesting genetic mechanisms are unlikely to account for KIFC1 overexpression[34].

Further studies are needed to decipher the mechanisms driving KIFC1 overexpression but ample

evidence indicates that KIFC1 expression is a robust prognostic biomarker in LC.

Targeting kinesins that regulate spindle assembly in cancer cells is a promising therapeutic approach

particularly in genomically unstable tumors[67–69]. Inhibitors of KIF18A and Eg5 (encoded by *KIF11*)

are being investigated in clinical trials[68, 70]. Our work nominates KIFC1 as a therapeutic target in LC,

with CA as a putative predictive biomarker. Our finding that KIFC1 expression was significantly higher

in LUAD lacking common driver mutations and in smokers compared to former and never smokers

suggests that KIFC1-targeted therapy could be useful in these currently underserved patient populations.

Encouragingly, KIFC1 KD does not affect proliferation or survival in non-malignant cells *in vitro*[52,

57, 71], but studies of KIFC1-targeted therapy in animal models are required to confirm its therapeutic

index. None of the KIFC1 inhibitors described to date (e.g. AZ82, CW069, SR17852, and KAA[44–46,

72]) have progressed to clinical studies, which may be attributable to poor potencies, specificities, and/or

bioavailability[47, 73]. Thus, development of clinical-grade KIFC1 inhibitors is warranted, especially

given their potential utility in diverse cancers. Our analyses of CA and the CA20 gene signatures suggest

that various opportunities exist to measure CA as a potential biomarker. As such, validation of these

candidate response-predictive features could facilitate the translation of KIFC1-targeted therapy by informing rational patient selection.

Our work and that of others indicates that cultured cancer cells exhibit less CA than malignant cells in tumor tissues[74]. Accordingly, we treated LC cells with low dose 945[30] to potentiate CA and better mimic clinical tumors and observed that KIFC1 LOF sensitized cells to CA. This observation was consistent with a study that reported that combining KIFC1 KD with potentiation of CA by cisplatin enhanced cytotoxicity in TNBC cells[57]. Together, these results suggest that KIFC1 inhibition could synergize with chemotherapy and/or irradiation which induce CA and/or multipolar spindles[11, 75, 76]. Supporting this concept, a recent study showed that KIFC1 KD sensitized LC to radiotherapy, though CA was not measured[71]. Further studies to define rational and effective KIFC1 inhibitor combinations are warranted, but our work nominates 945 as a promising partner for KIFC1-targeted therapy. Confirming whether 945 treatment enhances CA in tumor tissues from treated patients would provide additional evidence to support this strategy. Since 945 does not induce CA in non-malignant cells, this combination could be clinically tractable[30, 77, 78].

The recurrence of CA and KIFC1 overexpression in lung tumors, and the dependence of LC with CA on KIFC1 emphasize its therapeutic potential[33, 34]. We propose that KIFC1-targeted therapy could provide a novel strategy for potentiating GIN to lethal levels by forcing LC to divide with multipolar spindles, and that CA and KIFC1 expression could serve as biomarkers to guide effective use. We believe the efficacy of KIFC1-targeted therapy will be greatest when combined with treatments that exacerbate CA in lung tumors, and suggest that developing clinical-grade KIFC1 inhibitors and identifying and evaluating rational therapeutic combinations is warranted.

Acknowledgements:

393 We thank the UHN Therapeutics group for providing CFI-400945 for our studies, TCGA for tumor

394 genomics data, and the late Dr. Adi Gazdar for sharing LUAD cell lines. This work was supported by St.

395 Michael's Hospital, the Canada Research Chairs program, the Canadian Foundation for Innovation, and

396 the Canadian Institutes of Health Research (CIHR). C.Z. was supported by a CIHR scholarship.

Author Contributions:

Data Availability:

408 Datasets analyzed are available from public sources as described, and CRISPR screen data are available

409 from the corresponding author upon reasonable request.

Competing Interests:

ISB is currently an employee of Repare Therapeutics. MRB is currently an employee of Treadwell

Therapeutics. TWM owns equity in Treadwell Therapeutics Inc. and Agios Pharmaceuticals and is a

consultant for AstraZeneca and Tessa Therapeutics. DWC reports consultancy and advisory relationships

with AstraZeneca, Daiichi Sankyo, Exact Sciences, Eisai, Gilead, GlaxoSmithKline, Inflex,

Inivata/NeoGenomics, Lilly, Merck, Novartis, Pfizer, Roche and Saga; research funding to their

institution from AstraZeneca, Guardant Health, Gilead, GlaxoSmithKline, Inivata/NeoGenomics,

Knight, Merck, Pfizer, ProteinQure and Roche.

References

1. Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, Bray F (2021) Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. CA Cancer J Clin 71:209–249

2. Politi K, Herbst RS (2015) Lung cancer in the era of precision medicine. Clin Cancer Res 21:2213–2220

3. Herbst RS, Morgensztern D, Boshoff C (2018) The biology and management of non-small cell lung cancer. Nature 553:446–454

4. Siegel RL, Miller KD, Fuchs HE, Jemal A (2022) Cancer statistics, 2022. CA Cancer J Clin 72:7–33

5. Sabat-Pośpiech D, Fabian-Kolpanowicz K, Prior IA, Coulson JM, Fielding AB (2019) Targeting centrosome amplification, an Achilles' heel of cancer. Biochem Soc Trans 47:1209–1222

6. Koutsami MK, Tsantoulis PK, Kouloukoussa M, et al (2006) Centrosome abnormalities are frequently observed in non-small-cell lung cancer and are associated with aneuploidy and cyclin E overexpression. J Pathol 209:512–521

7. Jung CK, Jung JH, Lee KY, Kang CS, Kim M, Ko YH, Oh CS (2007) Centrosome abnormalities in non-small cell lung cancer: correlations with DNA aneuploidy and expression of cell cycle regulatory proteins. Pathol Res Pract 203:839–847

8. Godinho SA, Pellman D (2014) Causes and consequences of centrosome abnormalities in cancer. Philos Trans R Soc Lond B Biol Sci. https://doi.org/10.1098/rstb.2013.0467

9. Liao W-T, Lin P, Cheng T-S, Yu H-S, Chang LW (2007) Arsenic promotes centrosome abnormalities and cell colony formation in p53 compromised human lung cells. Toxicol Appl Pharmacol 225:162–170

10. Thaiparambil J, Dong L, Jasso D, Huang J-A, El-Zein RA (2020) Mitotic Spindle Apparatus Abnormalities in Chronic Obstructive Pulmonary Disease Cells: A Potential Pathway to Lung

Cancer. Cancer Prev Res . https://doi.org/10.1158/1940-6207.CAPR-19-0557

11. Löffler H, Fechter A, Liu FY, Poppelreuther S, Krämer A (2013) DNA damage-induced

centrosome amplification occurs via excessive formation of centriolar satellites. Oncogene

32:2963–2972

12. Ganem NJ, Godinho SA, Pellman D (2009) A mechanism linking extra centrosomes to

chromosomal instability. Nature 460:278–282

13. Lingle WL, Barrett SL, Negron VC, D'Assoro AB, Boeneman K, Liu W, Whitehead CM,

Reynolds C, Salisbury JL (2002) Centrosome amplification drives chromosomal instability in

breast tumor development. Proc Natl Acad Sci U S A 99:1978–1983

14. Sansregret L, Vanhaesebroeck B, Swanton C (2018) Determinants and clinical implications of

chromosomal instability in cancer. Nat Rev Clin Oncol 15:139–150

15. Holland AJ, Cleveland DW (2012) Losing balance: the origin and impact of aneuploidy in cancer.

EMBO Rep 13:501–514

16. Quintyne NJ, Reing JE, Hoffelder DR, Gollin SM, Saunders WS (2005) Spindle multipolarity is

prevented by centrosomal clustering. Science 307:127–129

17. Chan JY (2011) A clinical overview of centrosome amplification in human cancers. Int J Biol Sci

7:1122–1144

18. Thorsson V, Gibbs DL, Brown SD, et al (2018) The Immune Landscape of Cancer. Immunity

48:812–830.e14

19. Gao J, Aksoy BA, Dogrusoz U, et al (2013) Integrative analysis of complex cancer genomics and

clinical profiles using the cBioPortal. Sci Signal 6:l1

20. Tsherniak A, Vazquez F, Montgomery PG, et al (2017) Defining a Cancer Dependency Map. Cell

170:564–576.e16

21. de Almeida BP, Vieira AF, Paredes J, Bettencourt-Dias M, Barbosa-Morais NL (2019) Pan-cancer

association of a centrosome amplification gene expression signature with genomic alterations and

clinical outcome. PLoS Comput Biol 15:e1006832

22.   Ogden A, Rida PCG, Aneja R (2017) Prognostic value of CA20, a score based on centrosome

amplification-associated genes, in breast tumors. Sci Rep 7:262

23.   Uphoff CC, Drexler HG (2011) Detecting mycoplasma contamination in cell cultures by

polymerase chain reaction. Methods Mol Biol 731:93–103

24.   Wishart DS, Feunang YD, Guo AC, et al (2018) DrugBank 5.0: a major update to the DrugBank

database for 2018. Nucleic Acids Res 46:D1074–D1082

25.   Liu T, Lin Y, Wen X, Jorissen RN, Gilson MK (2007) BindingDB: a web-accessible database of

experimentally determined protein-ligand binding affinities. Nucleic Acids Res 35:D198–201

26.   Colic M, Wang G, Zimmermann M, et al (2019) Identifying chemogenetic interactions from

CRISPR screens with drugZ. Genome Med 11:52

27.   Thu KL, Silvester J, Elliott MJ, et al (2018) Disruption of the anaphase-promoting complex

confers resistance to TTK inhibitors in triple-negative breast cancer. Proc Natl Acad Sci U S A

115:E1570–E1577

28.   Zhu F, Lawo S, Bird A, Pinchev D, Ralph A, Richter C, Müller-Reichert T, Kittler R, Hyman AA,

Pelletier L (2008) The mammalian SPD-2 ortholog Cep192 regulates centrosome biogenesis. Curr

Biol 18:136–141

29.   Denu RA, Zasadil LM, Kanugh C, Laffin J, Weaver BA, Burkard ME (2016) Centrosome

amplification induces high grade features and is prognostic of worse outcomes in breast cancer.

BMC Cancer 16:47

30.   Mason JM, Lin DC-C, Wei X, et al (2014) Functional characterization of CFI-400945, a Polo-like

kinase 4 inhibitor, as a potential anticancer agent. Cancer Cell 26:163–176

31.   Ohashi K, Sequist LV, Arcila ME, et al (2013) Characteristics of lung cancers harboring NRAS

mutations. Clin Cancer Res 19:2584–2591

32.   Cancer Genome Atlas Research Network (2014) Comprehensive molecular profiling of lung

adenocarcinoma. Nature 511:543–550

33. Xiao Y-X, Yang W-X (2016) KIFC1: a promising chemotherapy target for cancer treatment?

Oncotarget 7:48656–48670

34. Wu H, Duan Y, Gong S, Zhu Q, Liu X, Liu Z (2022) An Integrative Pan-Cancer Analysis of

Kinesin Family Member C1 (KIFC1) in Human Tumors. Biomedicines.

https://doi.org/10.3390/biomedicines10030637

35. Liu Y, Zhan P, Zhou Z, et al (2016) The overexpression of KIFC1 was associated with the

proliferation and prognosis of non-small cell lung cancer. J Thorac Dis 8:2911–2923

36. Sun Y, Zhang Y, Lang Z, Huang J, Zou Z (2019) Prognostic and clinicopathological significance

of kinesin family member C1 in various cancers: A meta-analysis. Medicine 98:e17346

37. Grinberg-Rashi H, Ofek E, Perelman M, et al (2009) The Expression of Three Genes in Primary

Non–Small Cell Lung Cancer Is Associated with Metastatic Spread to the Brain. Clin Cancer Res

15:1755–1761

38. Director's Challenge Consortium for the Molecular Classification of Lung Adenocarcinoma,

Shedden K, Taylor JMG, et al (2008) Gene expression-based survival prediction in lung

adenocarcinoma: a multi-site, blinded validation study. Nat Med 14:822–827

39. Der SD, Sykes J, Pintilie M, Zhu C-Q, Strumpf D, Liu N, Jurisica I, Shepherd FA, Tsao M-S

(2014) Validation of a histology-independent prognostic gene signature for early-stage, non-small-

cell lung cancer including stage IA patients. J Thorac Oncol 9:59–64

40. Okayama H, Kohno T, Ishii Y, et al (2012) Identification of genes upregulated in ALK-positive

and EGFR/KRAS/ALK-negative lung adenocarcinomas. Cancer Res 72:100–111

41. Girard L, Rodriguez-Canales J, Behrens C, et al (2016) An Expression Signature as an Aid to the

Histologic Classification of Non-Small Cell Lung Cancer. Clin Cancer Res 22:4880–4889

42. Basto R, Brunk K, Vinadogrova T, Peel N, Franz A, Khodjakov A, Raff JW (2008) Centrosome

amplification can initiate tumorigenesis in flies. Cell 133:1032–1042

43. Levine MS, Bakker B, Boeckx B, et al (2017) Centrosome Amplification Is Sufficient to Promote

Spontaneous Tumorigenesis in Mammals. Dev Cell 40:313–322.e5

44. Wu J, Mikule K, Wang W, et al (2013) Discovery and mechanistic study of a small molecule

inhibitor for motor protein KIFC1. ACS Chem Biol 8:2201–2208

45. Watts CA, Richards FM, Bender A, et al (2013) Design, synthesis, and biological evaluation of an

allosteric inhibitor of HSET that targets cancer cells with supernumerary centrosomes. Chem Biol

20:1399–1410

46. Zhang W, Zhai L, Wang Y, et al (2016) Discovery of a novel inhibitor of kinesin-like protein

KIFC1. Biochem J 473:1027–1035

47. Yukawa M, Yamauchi T, Kurisawa N, Ahmed S, Kimura K-I, Toda T (2018) Fission yeast cells

overproducing HSET/KIFC1 provides a useful tool for identification and evaluation of human

kinesin-14 inhibitors. Fungal Genet Biol 116:33–41

48. Godinho SA, Picone R, Burute M, Dagher R, Su Y, Leung CT, Polyak K, Brugge JS, Théry M,

Pellman D (2014) Oncogene-like induction of cellular invasion from centrosome amplification.

Nature 510:167–171

49. Mittal K, Kaur J, Jaczko M, et al (2020) Centrosome amplification: a quantifiable cancer cell trait

with prognostic value in solid malignancies. Cancer Metastasis Rev.

https://doi.org/10.1007/s10555-020-09937-z

50. Fink G, Hajdo L, Skowronek KJ, Reuther C, Kasprzak AA, Diez S (2009) The mitotic kinesin-14

Ncd drives directional microtubule–microtubule sliding. Nat Cell Biol 11:717–723

51. Guha S, Patil A, Muralidharan H, Baas PW (2021) Mini-review: Microtubule sliding in neurons.

Neurosci Lett 753:135867

52. Kwon M, Godinho SA, Chandhok NS, Ganem NJ, Azioune A, Thery M, Pellman D (2008)

Mechanisms to suppress multipolar divisions in cancer cells with extra centrosomes. Genes Dev

22:2189–2203

53. Choe MH, Kim J, Ahn J, Hwang S-G, Oh JS, Kim J-S (2018) Centrosome Clustering Is a Tumor-

selective Target for the Improvement of Radiotherapy in Breast Cancer Cells. Anticancer Res

38:3393–3400

54. Goshima G, Nédélec F, Vale RD (2005) Mechanisms for focusing mitotic spindle poles by minus

end–directed motor proteins. J Cell Biol 171:229–240

55. Godinho SA, Kwon M, Pellman D (2009) Centrosomes and cancer: how cancer cells divide with

too many centrosomes. Cancer Metastasis Rev 28:85–98

56. Li Y, Lu W, Chen D, Boohaker RJ, Zhai L, Padmalayam I, Wennerberg K, Xu B, Zhang W

(2015) KIFC1 is a novel potential therapeutic target for breast cancer. Cancer Biol Ther 16:1316–

1322

57. Patel N, Weekes D, Drosopoulos K, et al (2018) Integrated genomics and functional validation

identifies malignant cell specific dependencies in triple negative breast cancer. Nat Commun

9:1044

58. Parvin A, Hao S-L, Tan F-Q, Yang W-X (2020) Inhibition of kinesin motor protein KIFC1 by

AZ82 induces multipolar mitosis and apoptosis in prostate cancer cell. Gene 760:144989

59. Celik B, Pasin O, Sen S, et al (2023) DNA methylation of KIFC1 gene in determination of

histological diagnosis, prognosis and metastasis of lung cancer. Pathol Res Pract 249:154742

60. Tabassum G, Singh P, Gurung R, et al (2023) Investigating the role of Kinesin family in lung

adenocarcinoma via integrated bioinformatics approach. Sci Rep 13:9859

61. Fu X, Zhu Y, Zheng B, et al (2018) KIFC1, a novel potential prognostic factor and therapeutic

target in hepatocellular carcinoma. Int J Oncol 52:1912–1922

62. Kostecka LG, Olseen A, Kang K, Torga G, Pienta KJ, Amend SR (2021) High KIFC1 expression

is associated with poor prognosis in prostate cancer. Med Oncol 38:47

63. Ogden A, Garlapati C, Li XB, et al (2017) Multi-institutional study of nuclear KIFC1 as a

biomarker of poor prognosis in African American women with triple-negative breast cancer. Sci

Rep 7:42289

64.   Mittal K, Choi DH, Klimov S, et al (2016) A centrosome clustering protein, KIFC1, predicts

aggressive disease course in serous ovarian adenocarcinomas. J Ovarian Res 9:17

65.   Sabat-Pośpiech D, Fabian-Kolpanowicz K, Kalirai H, Kipling N, Coupland SE, Coulson JM,

Fielding AB (2022) Aggressive uveal melanoma displays a high degree of centrosome

amplification, opening the door to therapeutic intervention. Hip Int 8:383–394

66.   Shinmura K, Iwaizumi M, Igarashi H, Nagura K, Yamada H, Suzuki M, Fukasawa K, Sugimura H

(2008) Induction of centrosome amplification and chromosome instability in p53-deficient lung

cancer cells exposed to benzo[a]pyrene diol epoxide (B[a]PDE). J Pathol 216:365–374

67.   Tischer J, Gergely F (2019) Anti-mitotic therapies in cancer. J Cell Biol 218:10–11

68.   Shahin R, Aljamal S (2022) Kinesin spindle protein inhibitors in cancer: from high throughput

screening to novel therapeutic strategies. Future Sci OA 8:FSO778

69.   Cohen-Sharir Y, McFarland JM, Abdusamad M, et al (2021) Aneuploidy renders cancer cells

vulnerable to mitotic checkpoint inhibition. Nature 590:486–491

70.   Braun J, Möckel MM, Strittmatter T, Marx A, Groth U, Mayer TU (2015) Synthesis and

biological evaluation of optimized inhibitors of the mitotic kinesin Kif18A. ACS Chem Biol

10:554–560

71.   Wang X, Raman N, Lemtiri-Chlieh G, et al (2023) Griseofulvin Radiosensitizes Non-Small Cell

Lung Cancer Cells and Activates cGAS. Mol Cancer Ther 22:519–528

72.   Kurisawa N, Yukawa M, Koshino H, Onodera T, Toda T, Kimura K-I (2020) Kolavenic acid

analog restores growth in HSET-overproducing fission yeast cells and multipolar mitosis in MDA-

MB-231 human cells. Bioorg Med Chem 28:115154

73.   Sharma N, Setiawan D, Hamelberg D, Narayan R, Aneja R (2023) Computational benchmarking

of putative KIFC1 inhibitors. Med Res Rev 43:293–318

74.   Mittal K, Choi DH, Ogden A, et al (2017) Amplified centrosomes and mitotic index display poor

concordance between patient tumors and cultured cancer cells. Sci Rep 7:43984

75. Mullee LI, Morrison CG (2016) Centrosomes in the DNA damage response—the hub outside the

centre. Chromosome Research 24:35–51

76. Scribano CM, Wan J, Esbona K, et al (2021) Chromosomal instability sensitizes patient breast

tumors to multipolar divisions induced by paclitaxel. Sci Transl Med 13:eabd4811

77. Kawakami M, Mustachio LM, Zheng L, et al (2018) Polo-like kinase 4 inhibition produces

polyploidy and apoptotic death of lung cancers. Proc Natl Acad Sci U S A 115:1913–1918

78. Chan CY-K, Yuen VW-H, Chiu DK-C, et al (2022) Polo-like kinase 4 inhibitor CFI-400945

suppresses liver cancer through cell cycle perturbation and eliciting antitumor immunity.

Hepatology. https://doi.org/10.1002/hep.32461

 FIGURE LEGENDS

Figure 1. CRISPR/Cas9 genetic screen identifies KIFC1 as a dependency in lung cancers with

centrosome amplification (CA). (A) Representative IHC images of mitotic cells in NSCLC tissues from

patient-derived xenografts (PDX) stained for the centrosome marker pericentrin. Arrowheads point to

centrosomes. Scale bar indicates 10um. (B) Characterization of CA in 10 NSCLC PDX. Centrosomes

were scored in mitotic cells only and cells were classified as having CA if they contained > 2 pericentrin

foci. (C) Potentiation of CA in H1299-Cas9+ cells by 945 treatment during the CRISPR screen. Cells

were sampled from each replicate (N=3/treatment arm). Bar plots indicate mean with SD, asterisks

indicate t-test p<0.0001. (D) Representative IF images exhibiting bipolar (BP), multipolar (MP), and

pseudo-bipolar (P-BP) mitotic spindles in H1299 cells. Scale bar indicates 5um. (E) Characterization of

mitotic spindles in H1299-Cas9+ cells sampled during the CRISPR screen. Counts were pooled across

the 3 replicates for each treatment condition. Asterisks indicate chi-square test p<0.0001. A minimum of

160 total mitotic cells were counted for each treatment condition for IF analyses in (C) and (E). (F)

Normalized Z-scores for each gene targeted in the sgRNA library calculated with the DrugZ algorithm.

Z-scores represent the relative abundance of sgRNAs in cells treated with 945 relative to DMSO control.

Negative Z-scores indicate genes targeted by sgRNAs that were significantly depleted in cells in which

CA was potentiated by 945. (G) Correlations between gene expression and CA20 scores for the top 16

genes identified by the CRISPR screen in the TCGA LUAD cohort. Correlation coefficients indicate

those for Pearson's correlation analyses.

Figure 2. KIFC1 expression is associated with multiple clinical factors in LUAD patients. KIFC1

expression was analyzed in the TCGA LUAD cohort to investigate associations with malignancy (A), 5-

year survival (B), stage (C), and smoking history (D). Asterisks indicate significance for Mann Whitney

or Kruskal-Wallis tests (*p<0.05, **p<0.01, ****p<0.0001). Error bars indicate mean and SD. NS =

never smokers, CS = current smokers, FS < 15 = former smoker for fewer than 15 years, FS > 15 =

former smoker for over 15 years. The association between KIFC1 expression and outcome was assessed

in patients whose tumors exhibited KIFC1 expression ranking in the top and bottom tertiles of expression.

Asterisks indicate $p < 0.01$ for a log-rank test on the Kaplan-Meier survival curves.

Figure 3. Centrosome amplification and genomic instability are positively correlated with KIFC1

expression in LUAD models and tissues. (A) Correlation between KIFC1 expression and CA in 11

LUAD models. Protein expression was quantified from western blots and CA from IF conducted under

basal conditions (Supp. Fig. 4). (B,C) Correlation between KIFC1 dependency and CA20 in NSCLC (B)

and LUAD (C) cell lines from the CCLE. KIFC1 gene effect scores were accessed from the DepMap

database and negative scores indicate genetic dependency. CA20 scores were obtained from the de

Almeida 2019 study. (D) Association between KIFC1 mRNA expression and CA20 in the TCGA LUAD

cohort. (E,F) Correlation between KIFC1 mRNA expression and fraction of genome altered (FGA) and

aneuploidy score. FGA and aneuploidy scores were obtained from cBioPortal. All correlation

coefficients indicated were calculated using Pearson's correlation analyses.

Figure 4. Validation of KIFC1 as a genetic dependency in LUAD models with CA. (A) Potentiation

of CA by low-dose 945 treatment (7.5nM) in Cas9+ cell models. Centrosomes were detected using IF

for CEP192 and CA was scored as the proportion of cells with greater than two centrosomes. Error bars

indicate the mean and SD (N=3), and asterisks indicate p-value significance for t-tests between

treatments. (B) Classification of mitotic spindles as bipolar (BP), multipolar (MP), or pseudo-bipolar (P-

BP) for each treatment condition. Asterisks indicate p-value significance for a chi-square test. For (A)

and (B), data for 3 replicates are pooled and at least 150 total mitotic cells were scored for each treatment

condition. (C) Confirmation of CRISPR-mediated loss of KIFC1 expression in LUAD cell lines. Cas9+

cells were transduced with mCherry (mCH) or GFP lentivectors to express sgRNA targeting AAVS1

(AV) or KIFC1 (K3 targeting exon 3 and K4 targeting exon 4). (D) Multicolour competition assays

comparing the relative fitness of cells with wildtype (AV) and genetically perturbed *KIFC1* (K3, K4) in

DMSO- and 945-treated conditions. mCH-AV cells were mixed 1:1 with GFP-AV, GFP-K3, or GFP-K4

cells and grown over multiple passages. The GFP+ fraction was monitored over time by flow cytometry.

Data are representative of two independent experiments (*p<0.05, **p<0.01, ***p<0.001,

****p<0.0001).

Figure 5. Depletion of KIFC1 sensitizes LUAD models with high basal CA to potentiation of CA.

(A,B) Confirmation of siRNA-mediated KIFC1 knockdown by western blotting (A) and RT-qPCR (B).

Relative quantification (RQ; $2^{-\Delta\Delta Ct}$) values were calculated using the $\Delta\Delta Ct$ method and error bars indicate

the minimum and maximum RQ for 3 technical replicates. RNA and protein were collected 24hr post

transfection at the time that cells were seeded for functional assays. Representative knockdowns are

shown. NTC = non-targeting control. K1, K2 indicate independent siRNA targeting KIFC1. (C) Dose

response assays for 945 treatment in cells with and without KIFC1 knockdown. Cell viability at each

dose was normalized to that for DMSO-treated controls. Each curve represents mean viability data for 3-

5 biological replicates. Error bars indicate SEM. Numbers indicate area under the curve. (D) Clonogenic

survival assays. Data are representative of 2-3 biological replicates per model. Colony survival was

normalized to that observed in DMSO-treated conditions for each siRNA. Error bars indicate SEM of 6

technical replicates. (E) Classification of mitotic spindles as bipolar (BP), multipolar (MP), or pseudo-

bipolar (P-BP). Data are pooled for 3-4 replicate experiments and at least 150 total mitotic cells were

scored for each treatment condition. Asterisks indicate p-value significance for chi-square tests between

NTC and K1 or K2 siRNA for each treatment condition (*p<0.05, **p<0.01, ***p<0.001,

****p<0.0001).

Supplementary Figure 1. **Technical validation of H1299 CRISPR/Cas9 screen.** (A) Confirmation of Cas9 editing efficiency in H1299-Cas9 cells. Cas9+ cells were transduced with either a BFP-GFP-empty vector or a BFP-GFP-sgGFP vector, and GFP expression was quantified by flow cytometry 48hr later. The percentage of BFP+GFP+ cells is indicated in Q2. (B) Potentiation of centrosome amplification in H1299-Cas9 cells by treatment with low doses of 945. Centrosomes were scored based on immunofluorescence for CEP192. Error bars indicate mean and SD for two replicates. (C) Growth curves of DMSO and 945-treated cell populations during the CRISPR screen. Error bars indicate the mean and SD for the 3 technical replicates. (D) Identification of essential genes in H1299 cells from the CRISPR screen. Normalized Z-scores for target genes in DMSO-treated cells at screen end point relative to screen onset (T0) are shown. Z-scores represent the relative abundance of sgRNAs targeting each gene in the cell populations. Negative Z-scores indicate genes targeted by sgRNAs that were significantly depleted in DMSO-treated cells at the end of the screen. (E) Abundance of sgRNA targeting *MYC* and *NRAS* at screen onset (T0) compared to endpoint (after 17 doublings).

Supplementary Figure 2. **Association of KIFC1 expression with clinical features in additional LUAD cohorts.** A) KIFC1 expression in LUAD from males and females in the TCGA dataset. B) KIFC1 expression in the TCGA cohort segregated by mutation status for common oncogenic drivers in LUAD: *EGFR, KRAS, MET, ROS1, RET, ALK, NRAS* and *BRAF*. In 4 additional LUAD datasets from the gene expression omnibus, KIFC1 expression was investigated in relation to smoking history (B), sex (C), stage (D), CA20 score (E), and 5-year survival (F). Asterisks indicate significance for Mann Whitney or Kruskal-Wallis tests. Error bars indicate mean and SD. NS = never smokers, CS = current smokers, ES = ever smokers, FS = former smoker. Pearson's correlation coefficients are indicated for associations between KIFC1 expression and CA20 scores. *Note that only 19/20 genes in the CA20 signature were evaluable in the GSE68465. Association between KIFC1 expression and 5-year survival was assessed in patients whose tumors exhibited KIFC1 expression ranking in the top and bottom tertiles of expression.

Asterisks indicate significance for log-rank tests on the Kaplan-Meier survival curves (*p<0.05,

**p<0.01, ***p<0.001, ****p<0.0001).

Supplementary Figure 3. **Association of KIFC1 expression with clinical features in lung squamous**

cell carcinoma (LUSC) patients. KIFC1 expression was analyzed in the TCGA LUSC cohort to

investigate associations with malignancy (A), smoking history (B), sex (C), and stage (D). Asterisks

indicate significance for Mann Whitney or Kruskal-Wallis tests (****p<0.0001). Error bars indicate

mean and SD. NS = never smokers, CS = current smokers, FS < 15 = former smoker for fewer than 15

years, FS > 15 = former smoker for over 15 years. (E) Association between KIFC1 expression and 5-

year survival was assessed in patients whose tumors exhibited KIFC1 expression ranking in the top and

bottom tertiles of expression. The survival association was investigated using a log-rank test on the

Kaplan-Meier survival curves. (F) Correlation between KIFC1 mRNA expression and CA20 scores.

Correlation coefficient is that for Pearson's correlation.

Supplementary Figure 4. **Centrosome amplification and KIFC1 expression in LUAD cell lines.** (A)

Western blot for KIFC1 expression in 14 LUAD lines. (B) Quantification of protein expression in (A).

KIFC1 levels were normalized to GAPDH. (C) Proportion of cells with centrosome amplification in 11

LUAD lines. Centrosomes were identified by immunofluorescence staining for CEP192. (D) Correlation

between KIFC1 dependency and CA20 in LUSC (D) and SCLC (E) cell lines from the CCLE. KIFC1

gene effect scores were accessed from the DepMap database and negative scores indicate genetic

dependency. CA20 scores were obtained from the de Almeida 2019 study. The correlation coefficients

indicated were calculated using Pearson's correlation analyses.

Supplementary Figure 5. **Cytotoxicity of AZ82 in LUAD cell lines.** LUAD cells were treated for 5

days with DMSO or AZ82 at two doses, 3.5uM (A) and 7.5uM (B). Surviving cells were fixed, stained

with SRB, and solubilized for colorimetric readouts on a spectrophotometer. Values indicate viability in

AZ82-treated cells relative to DMSO controls. Error bars indicate SD for replicate experiments.

Supplementary Figure 6. Effect of KIFC1 knockdown on proliferation and clonogenic survival. A) Cell

proliferation was measured using the ATP-Lite assay over 5 days. Values indicate luminescence signals

normalized to Day 0. Growth curves represent data pooled from 4-6 replicate experiments and error bars

indicate SEM. B) Raw colony numbers for each siRNA condition are plotted for the DMSO-treated cells

from Fig.5. Colonies were grown for 11-14 days. Colony numbers in cells transfected with NTC and K1

or K2 siRNA were compared using t-tests. Data represent 3-5 experimental replicates. Error bars indicate

SEM.

Fig.1

Fig.2

Fig.3

Fig.4

Fig.5