1 **De novo annotation of the wheat pan-genome reveals complexity and diversity within the**
2 **hexaploid wheat pan-transcriptome**
3
Authors

Benjamen White[1†], Thomas Lux[2†], Rachel L Rusholme-Pilcher[1†], Angéla Juhász[6†], Gemy
Kaithakottil[1], Susan Duncan[1,3], James Simmonds[3], Hannah Rees[1], Jonathan Wright[1], Joshua
Colmer[1], Sabrina Ward[1], Ryan Joynson[1,4], Benedict Coombes[1], Naomi Irish[1], Suzanne
Henderson[1], Tom Barker[1], Helen Chapman[1], Leah Catchpole[1], Karim Gharbi[1], Moeko
Okada[5,16,17], Hirokazu Handa[18], Shuhei Nasuda[19], Kentaro K. Shimizu[5,16], Heidrun Gundlach[2],
Daniel Lang[2], Guy Naamati[7], Erik J. Legg[8], Arvind K. Bharti[8], Michelle L. Colgrave[6,9], Wilfried
Haerty[1], Cristobal Uauy[3], David Swarbreck[1], Philippa Borrill[3], Jesse A. Poland[10], Simon
Krattinger[10], Nils Stein[11,15], Klaus F.X. Mayer[2,12], Curtis Pozniak[13], 10+ Wheat Genome Project,
Manuel Spannagl^[2], Anthony Hall^[1,14]
14
[1]Earlham Institute, Norwich Research Park, Norwich, NR4 7UH, UK.
[2]PGSB Plant Genome and Systems Biology, Helmholtz Center Munich, German Research Center
for Environmental Health, Neuherberg, Germany.
[3]John Innes Centre, Norwich Research Park, Norwich, NR4 7UH, UK.
[4]Limagrain Europe, Clermont-Ferrand, Auvergne-Rhône-Alpes, France.
[5]Department of Evolutionary Biology and Environmental Studies, University of Zurich, Zurich,
Switzerland.
[6]Australian Research Council Centre of Excellence for Innovations in Peptide and Protein
Science, School of Science, Edith Cowan University, Joondalup, WA, 6027, Australia.
[7]EMBL-EBI, Wellcome Genome Campus, Hinxton, Cambridgeshire, UK.
[8]Syngenta Crop Protection, Research Triangle Park, NC, USA.
[9]CSIRO Agriculture and Food, St Lucia, QLD 4067, Australia.
[10]Plant Science Program, Biological and Environmental Science and Engineering Division, King
Abdullah University of Science and Technology (KAUST), Thuwal 23955-6900, Kingdom of Saudi
Arabia
[11]Center of Integrated Breeding Research (CiBreed), Georg-August-University, Göttingen,
Germany
[12]School of Life Sciences, Technical University Munich, Freising, Germany.
[13]Crop Development Centre, The University of Saskatchewan, Saskatoon, Canada.
[14]School of Biological Sciences, University of East Anglia, Norwich, UK
[15]Leibniz Institute of Plant Genetics and Crop Plant Research (IPK) Gatersleben, Seeland,
Germany
[16]Kihara Institute for Biological Research, Yokohama City University, Yokohama, Japan
[17]Graduate School of Science and Technology, Niigata University, Niigata, Japan
[18]Graduate School of Life and Environmental Sciences, Kyoto Prefectural University, Kyoto,
Japan
[19]Graduate School of Agriculture, Kyoto University, Kyoto, Japan
42
43 †These authors contributed equally to this work.

45 *Correspondence: manuel.spannagl@helmholtz-muenchen.de, Anthony.Hall@earlham.ac.uk

Abstract

Wheat is the most widely cultivated crop in the world with over 215 million hectares grown annually. However, to meet the demands of a growing global population, breeders face the challenge of increasing wheat production by approximately 60% within the next 40 years. The 10+ Wheat Genomes Project recently sequenced and assembled the genomes of 15 wheat cultivars to develop our understanding of genetic diversity and selection within the pan-genome of wheat. Here, we provide a wheat pan-transcriptome with *de novo* annotation and differential expression analysis for nine of these wheat cultivars, across multiple different tissues and whole seedlings sampled at dusk/dawn. Analysis of these *de novo* annotations facilitated the discovery of genes absent from the Chinese Spring reference, identified genes specific to particular cultivars and defined the core and dispensable genomes. Expression analysis across cultivars and tissues revealed conservation in expression between a large core set of homoeologous genes, but also widespread changes in sub-genome homoeolog expression bias between cultivars. Co-expression network analysis revealed the impact of divergence of sub-genome homoeolog expression and identified tissue-associated cultivar-specific expression profiles. In summary, this work provides both a valuable resource for the wider wheat community and reveals diversity in gene content and expression patterns between global wheat cultivars.

Introduction

Wheat (*Triticum aestivum*) is the most widely grown crop and is cultivated in 12 mega-environments across the world[1], with 777.7 metric tonnes harvested globally in 2021/22 (www.fao.org). Pressures of climate change, political instability, a move to more sustainable farming and a reduction in agricultural land are putting increasing demand on international wheat harvests[2]. Efforts to overcome these pressures can be accelerated by understanding the genetic diversity of global wheat cultivars and their pan-transcriptional variation.

Wheat has a large (15Gb) allohexaploid (BBAADD) genome, derived from a series of relatively recent hybridisation events[3]. Its size, evolutionary history, and high repeat content, despite hindering genome assembly, make wheat an interesting model for the evolution of large polyploid genomes. Step changes in technology have enabled the chromosome-level assembly of nine high-quality wheat genomes by a global consortium. These genomes revealed evidence of widespread structural rearrangements, introgression from wild relatives and the impacts of parallel international breeding programmes[4]. To date, these genomes were annotated only by projecting Chinese Spring gene models across the new assemblies. The generation of *de novo* annotations for these genomes provides a key insight into gene gain and loss, reveals novel gene models across wheat cultivars and facilitates comparative gene expression analysis between cultivars.

Previous analyses of the wheat transcriptional landscape described tissue-specific changes in gene expression in two cultivars, using a common Chinese Spring reference genome[5].

Polyploidy leads to complex effects on gene expression resulting from structural variation, gene
duplication, deletion and neofunctionalization, ultimately increasing variation in gene
expression and the plasticity of the species. To date, studies of plant pan-transcriptomes either
rely on read alignment to a single reference genome which can result in reference bias, or
generate *de novo* transcript assemblies from short read data that can accumulate errors and
technical artefacts[6].
94
Here, we generate *de novo* gene annotations, incorporating long reads for the nine assembled
wheat cultivars, providing a valuable resource for wheat researchers and breeders. We identify
evidence of widespread gene duplication and deletion, revealing the population structures
imposed by repeated hybridisations from wild relatives and different breeding programmes.
We define the hexaploid wheat core and dispensable transcriptome and our analysis of gene
expression and gene networks across different tissues and between cultivars reveals
conservation and divergence in expression balance across homoeologous sub-genomes. We
exemplify the value of these analyses through an in-depth investigation of the pan-genome
variability of prolamin gene content and expression; a key trait for quality and health aspects in
wheat.
105
Results
***De-novo* gene annotations of the pan-cultivars define the core and accessory gene sets**
To precisely assess the gene content and differences in gene expression, copy number and the
presence/absence of genes between the wheat cultivars, we generated a *de novo* gene
annotation for each of the nine pan-cultivars. We used an established automated annotation
pipeline which built evidence-based gene model predictions using a comprehensive
transcriptomic dataset. This dataset was made up of Iso-Seq data from roots and shoots (390-
700 K reads per sample), and RNA-seq data (150 bp paired-end read, 56-85 M pairs of reads per
sample) obtained for each cultivar from five distinct tissue types and whole aerial organs
sampled at dawn and dusk (**Figure 1A,** see methods for a full description and **Extended Data
Figure 1** for details of quality control). In addition to the transcriptomic dataset, the gene
annotation pipeline also used protein homology and *ab initio* prediction. Finally, a gene
consolidation procedure (**Extended Data Figure 2A**) was developed to identify and correct for
missed gene models. This step ensures the best possible comparability between the wheat
genomes and gene repertoire[7].
121
The number of high-confidence gene models identified ranges from 140,178 for CDC Landmark
to 145,065 for Norin 61 (**Figure 1B**). Low-confidence genes, primarily representing gene
fragments, pseudogenes and gene models with only weak support, are in the range of 315,390
(Mace) to 405,664 (SY Mattis). With a maximal difference of 3.5%, the number of high-
confidence (HC) genes appears to be similar across cultivars, whereas most of the differences in
gene number observed can be attributed to the low-confidence gene set. For around 70% of
the HC genes we obtained evidence for transcription in at least one condition.
129
130 We benchmarked the quality of the *de novo* gene predictions against BUSCO v5.1.2 with the
131 poales_odb10 lineage dataset, representing 4,896 Poales near-universal single-copy orthologs.

On average, we found more than 99.8% of the BUSCO genes represented at least one complete copy and 86% by three complete copies (**Figure 1B**). This is an improvement in complete BUSCO genes over the gene projections from Chinese Spring used in the first study of the wheat pan-genome[4] and can be explained by the *de novo* gene annotation strategy applied here, which included comparable RNA-seq and Iso-Seq datasets and *ab initio* prediction, as well as the final consolidation step. The *de novo* annotations are available in Ensembl Plants release 52.

Figure 1. Study design, de-novo gene annotations and orthologous framework. **A**. Overview of transcriptome data generated for this study of the wheat pan-cultivars. 1 & 2: whole aerial organs sampled at dawn and dusk, 3: root, 4: complete spike at heading (GS59), 5: flag leaf 7 days post anthesis (GS71), 6: whole grains 15 days post anthesis (GS77). **B.** *De novo* gene prediction results for each pan-cultivar (left side, "genes", separated for A, B and D sub-genome) as well as a summary of the BUSCO completeness assessment of gene models (right

side, "BUSCO"). **C**. GENESPACE construction and visualisation of orthologous genes within the
wheat pan-genome, using *de novo* predicted gene models.

Genes and gene families exclusive or amplified in a specific cultivar are of major interest in a
pan-genomic context[8]. While genes present in all compared cultivars are referred to as the core
genome, cloud and shell genes are found only in one (cloud) or shared in a subset of cultivars
(shell). The improved gene annotation enabled the construction of a high-density orthologous
framework for wheat. GENESPACE[9] was used to derive syntenic relationships between all
chromosomes and sub-genomes, allowing in-detail investigation of macro- and micro-synteny
(**Figure 1C**) and gene copy number variations. While previously identified rearrangements such
as the chromosome 5B/7B translocation in SY Mattis and Arina*LrFor* were confirmed, additional
frequent small-scale structural variations can now be examined in the context of their gene
content. We found a 16 Mb inversion, split into three segments of around 5Mb each, on
chromosome 3D between Canadian cultivars CDC Stanley and CDC Landmark which coincides
with the locations of QTLs related to biomass and grain weight[10].

We identified groups of orthologous genes (referred to as orthogroups) among the wheat high-
confidence gene models of all cultivars. A total of 54,865 orthogroups contained 99.7% of all
genes, with 173 orthogroups identified as cultivar-specific and 3,756 genes not clustered in any
orthogroup - defining the cloud genome. Cloud and shell genes have previously been found to
be associated with disease resistance[11], adaptation to new environments[12], or important
agricultural traits[13]. Within the shell genome, our analysis identified orthogroups that are
shared only between specific cultivars. Examples include CDC Stanley and CDC Landmark from
Canada, Mace and LongReach Lancer from Australia or Arina*LrFor*, SY Mattis and Julius from
Europe (highlighted in yellow in **Figure 2A**) which all share exclusive sets of genes. These
observed patterns likely reflect the complex breeding history of the selected pan-cultivars
which represent wheat lines from different regions, growth habits and breeding programs.
Inspection of the chromosomal location of these gene groups identified multiple clusters
(**Figure 2B and Extended Data Figure 3**) that are likely associated with crosses to distinct
material or hybridisations with wild or domesticated relatives; events common in wheat[14].

Proportions of core (genes present in all cultivars), shell (genes present in 2-8 cultivars) and
cloud (genes found in only one cultivar and unclustered genes) genes were found to be similar
across the pan-cultivars (**Figure 2C**). On average 76.34% of genes were classified as core,
23.32% as shell, and 0.33% as cloud (**Extended Data Figure 2B**). Amongst the core gene set, we
found biological functions associated with basic metabolic, catabolic and DNA repair/replication
processes enriched (**Supp. Table 1**), while stress response and regulation of gene expression
were overrepresented in the shell genes (**Supp. Table 2**). In the set of cloud genes, functions
related to chromatin organisation and reproductive processes were found to be enriched
(**Supp. Table 3**). Expression patterns of core, shell and cloud genes revealed pronounced
differences globally, but not between the sub-genomes (**Figure 2D**). As observed in other pan-
genomes[15], core genes tend to be higher expressed in all sub-genomes and tissues, as
compared to both shell and cloud genes.

188
Figure 2. **The wheat core-, shell-, and cloud- genome and homoelogous expression patterns.**
A. UpSet plot showing intersects of orthogroup conservation between pan-cultivars and the
relation to their breeding programs and sowing season. **B.** A representation of CDC Stanley
chromosome 3B showing the positions of Canadian-specific genes (top bar), heatmaps showing
coverage scores between genes in CDC Stanley and CDC Landmark (middle bar) and coverage
scores between CDC Stanley and Norin 61 (bottom bar). Coverage scores are calculated using
kmers from each CDC Stanley gene to search the genomes of the other cultivar and range from
0 to 1 with values closer to 1 indicating greater similarity. Regions of greater difference are
shown as darker bands. The plot shows the 0-50 Mb region of chromosome 3B (indicated by a
red box). The mean of the coverage score between CDC Stanley genes in this region and genes
in the non-Canadian lines is plotted. A cluster of four Canadian-specific genes (marked by a red
dashed line) lies in a region which is noticeably different between CDC Stanley and the non-
Canadian lines potentially representing an introgression. **C.** Number of genes belonging to core,
shell and cloud ortholog groups across cultivars. **D.** Violin plots of core, cloud and shell log
average gene expression across all combined cultivars and tissues, in each sub-genome. Higher
mean expression was observed in core genes across all sub-genomes. **E.** Ternary plots, of stable
(left) and dynamic (right) 30-let expression, where there is a homoeolog present on each sub-
genome, of all tissue in all cultivars, combined, showing more overall balanced expression in
stable 30-lets and unbalanced expression in dynamic 30-lets.
208

Duplication of genes has been identified as a major driver of gene function evolution and
adaptation in plants[12]. In wheat, a large number of tandem duplications was previously found
both in the Chinese Spring (IWGSC v1.1) reference genome and the pan-genome assemblies[16,4].
Our full *de novo* gene annotation of the wheat pan-genome, in combination with the extensive
gene expression data presented in this study, allowed for an in-depth assessment of gene
duplication dynamics across cultivars in hexaploid wheat.
215
We identified on average 5,040 tandem arrays (HC genes only) in each cultivar, with the lowest
in CDC Landmark (4,914) and Arina*LrFor* as the highest (5,172). In addition, we tested whether
there is a bias in expression towards one member of the array. We found that for 2,520 arrays
one of the two members was biased in its expression with respect to the other member,
whereas for 1,800 arrays both copies were expressed at similar levels or varied depending on
the tissue (**Extended Data Figure 2C**). 719 tandem arrays were not expressed under the
investigated conditions. Amongst all tandemly duplicated genes in wheat, biological functions
associated with phosphorylation, response to stimulus and stress and reproductive processes
were enriched (**Supp. Table 4**). We also investigated the conservation of tandem arrays across
all pan-cultivars. Around 69% of the tandem arrays identified in a specific cultivar were found to
be shared with all other pan-cultivars, with the remaining 31% tandems showing varying
conservation (**Extended Data Figure 2D**). These results highlight the impact of tandemly
duplicated genes as a potential key driver of evolution and adaptation. Besides functional
redundancy of homoeologous genes in hexaploid wheat, tandem genes and their expression
(bias) are therefore an important target for breeding applications.
231
Conservation of Global Expression in the Wheat Pan-Transcriptome
To investigate changes in global gene expression across cultivars, biological replicates from
whole aerial organs at dusk and dawn, and from flag leaf, root, spike and grain, were used to
generate normalised gene expression counts. We observed from principal component analysis
of the normalised counts that most of the variance is represented by the first principal
component, representing the different developmental stages, and also similar grouping of
expression overall (**Extended Data Figure 1A, B**). We then used these normalised counts from
the nine cultivars together with complete *de novo* annotations for the core, shell, and cloud
group genes, to explore differences in expression between tissues across all cultivars. The
patterns of expression observed in each individual orthologous class were consistent across
tissues, and between sub-genomes, with core genes also showing an overall higher mean
expression than either shell or cloud (**Figure 2D, Extended Data Figure 4A**). Indicating a global
conservation of expression, irrespective of tissue type or sub-genome biases.
245
The tissue-specific gene index (tau) was employed to assess the degree of gene expression
specificity to flag leaf, root, spike or grain tissues across all cultivars (**Extended Data Figure 4B**).
We observed the least number of tau genes in flag leaf (1,005 - 3,202 specific genes), that were
significantly less (t-test; $p < 0.001$) overall compared to either root (4,736 - 8,974 specific
genes), spike (5,453 - 9,323 specific genes) or grain (3,955 - 12,157 specific genes), that showed
no significant difference between each other. However, the number of specific genes showed
the least cultivar variability for flag leaf tissues, compared to the wide range in the number of

grain-specific genes observed between cultivars. This could be the result of contrasting
transcriptomic complexity between flag leaf and grain tissues, representing different
developmental stages of maturity and metabolic activity. In polyploid crops agricultural traits
are often modulated by an interaction of homoeologous copies of genes[5]. In wheat, previous
studies have focused on tissue-specific expression across homologous triads, identifying sets of
triads that are either balanced or unbalanced in their sub-genome expression. Here, we
compared variation in triad expression across cultivars using all 13,521 identified sets of 30-let
genes with a homoeolog present on each sub-genome of the *de novo* annotated cultivars. Using
previously reported cut-off values[5], we observed similar sub-genome expression in these 30-
lets, in each of the cultivars, to that reported previously in Chinese Spring, with 102 also being
classed as not expressed (**Extended Data Figure 4C**)[5]. However, when comparing the bias of
sub-genome expression, we observed 8,028 (59.37%) of these 30-lets to have a conserved,
'stable', balanced expression between the three homoeologous copies across all cultivars
(**Extended Data Figure 4D**). Whereby 'stable' expression relates to a conserved sub-genome
expression bias between cultivars, as opposed to a 'dynamic' expression where a change in sub-
genome expression bias can be observed in one or more cultivars.
269
As well as conservation of the balanced state we also see conservation in dominance or
suppression within triad groups with 276 showing stable suppressed expression and 63 stable
dominant expression. Stably expressed 30-lets showed GO term enrichment for essential
biological processes associated with photosynthesis, translation, DNA replication, exocytosis,
glycolytic process and cell redox homeostasis.(**Extended Data Figure 4E**). Whilst the 5,052
37.36% 'dynamically' expressed 30-lets that showed a change in the bias of sub-genome
expression in at least one cultivar were found to be significantly enriched for transmembrane
transport, response to stress, response to oxidative stress, defence response and
photosynthesis. These dynamic 30-lets were observed to be less fixed to a specific sub-genome
expression pattern compared to stably expressed 30-lets, showing a further Euclidean distance
from a, b, c or centroid points (**Figure 2E**). Across these dynamically expressed 30-lets, 4,467
showed balanced expression in at least one cultivar, with B sub-genome suppression being the
next most represented balance of expression occurring in 1,972 of the dynamic 30-let sets
(**Extended Data Figure 4F**). Overall, more suppression of expression was seen than dominance.
The Kruskal-Wallis test, applied to assess differences in the mean values of the dynamic 30-let
bias across the cultivars, revealed no significant differences when examining the total
percentage of each expression bias ($p > 0.05$). This suggests that the bias of dynamic
expression, whilst different for individual 30-lets, has been proportionally conserved across
these cultivars.

289
290
Figure 3. Components of the wheat consensus network with functional annotation and UK
cultivar Claire-specific differences A. Number of genes in each module from the seven cultivar
consensus network. **B.** Hierarchical clustering of 23 consensus module eigengenes identifying
six metamodules. **C.** GO-slim terms of biological processes associated with genes in 6
metamodules. Only terms with p<0.05 and >10 significant genes are shown. Bubble colour
indicates the -log$_2$p-value significance from Fisher's exact test and size indicates the frequency
of the GO-slim term in the underlying EBI Gene Ontology Annotation database (larger bubbles
indicate more general terms). **D**. Claire-specific network of 18 genes with divergent expression
patterns compared to all other cultivars. Node size is scaled to the log$_2$ average expression +1 of
each gene in spikelet tissues and edge width reflects the weight of the connection between
nodes. **E.** Log$_2$ fold change in expression across all tissues, between Claire and remaining
cultivars of 18 Claire-specific genes.

Conserved patterns of co-expression
In addition to the nine cultivars for which we generated *de novo* annotations, we also collected
expression data from the same tissues and conditions for the remaining five cultivars included
in the wheat pan-genome. To exploit these additional transcriptomic datasets in the absence of
de novo annotations, we employed the consensus network function of WGCNA[17] which uses
expression counts from a single reference (CSv1.1) to enable the identification of gene
expression modules that represent biological pathways conserved across all cultivars.

To explore how regulatory networks are conserved across tissues and cultivars of the pan-
transcriptome, we constructed our consensus co-expression network spanning seven cultivars;

314   four with chromosome pseudomolecule assemblies (Arina*LrFor*, Julius, Jagger, Norin 61) and
315   three with scaffold assemblies; two UK derived (Cadenza, Claire) and one key CIMMYT (Weebill)
316   cultivar. Together these seven cultivars represent a wide range of the genetic variation
317   observed in the pan-genome[4].
318
To identify sets of genes with highly correlated expression patterns across cultivars we used
high-confidence genes expressed at greater than five normalised counts in at least three
samples, with less than two standard deviations in normalised counts, between sample
replicates. The resulting 48,337 genes from each cultivar (338,359 genes in total) were used to
construct the consensus network, with modules ranging in size from 84 to 9,183 genes (**Figure
3A**). This consensus network accounted for 70% (34,069 genes) of the genes from each cultivar
in our original dataset. These modules comprised co-expressed genes that exhibited highly
similar expression patterns across all six developmental stages in all seven cultivars. A
comparison of the consensus module eigengenes (MEs) for these 23 modules demonstrated
the conservation of expression patterns between cultivars (**Extended Data Figure 5**). We used
hierarchical clustering to further collapse the 23 consensus modules into six consensus
metamodules (**Figure 3B**); comprising 32,936 genes from each cultivar. GO-slim term analysis
(**Figure 3C**) and distinct transcription factor superfamily membership (**Extended Data Figure 6**)
indicated that each metamodule could be associated with distinct biological processes (**Supp.
Table 5**).
334
We defined a threshold for classifying inter-consensus module relationships and used this to
make pairwise comparisons of the ME for each consensus module and identify modules with
divergent or similar patterns of expression. We then used these module relationships to
compare how the 30-let triads were split across our consensus network. Within the genes used
to build our consensus network, we identified 6,867 of the 10,521 complete triad sets (50.8%).
3,640 (53.0%) of these triads were assigned to modules within the network, with the remaining
3,227 (47%) triads having at least one member present in the unclustered set of genes. This set
of genes that could not be fitted into the consensus network will contain genes with low
variance or low expression across tissues, and genes that do not show the same pattern of
expression across all seven cultivars. Of the 3,640 triads within our network 3,548 (96%)
belonged to either the same or similar expression modules, reflecting the conservation of
expression between the A, B and D sub-genomes. 2,431 of these triads (66.8%) belonged to the
stable category of 30-lets defined previously through comparison of individual triad expression
balance across all tissues and 9 cultivars. The identification of co-expression modules containing
similarly expressed triad members reveals additionally conserved genes, tightly connected to
these stable triads. Using the consensus network, we also observed 146 of the 3,640 triads (4%)
where sub-genome members were split across divergent modules. These triads were
significantly enriched for GO terms associated with the regulation of signal transduction and
DNA metabolic process (**Supp. Table 6**).
354
355
Using a consensus network approach to identify cultivar-specific co-expression patterns

Whilst our consensus network enables robust biological inferences through the identification of
conserved gene sets across all cultivar members of the network, we also report a set of 14,268
unclustered genes that cannot be fitted to the consensus modules. These 14,268 genes may
have an expression profile that correlates with a consensus ME, but as this pattern of
expression is not conserved across all seven cultivars, these genes will not be placed within the
consensus network. Within our unclustered gene set we identified 1,753 such genes where the
pattern of expression for a single cultivar was closely correlated to a consensus ME displaying
an expression profile divergent to the module containing the same gene in the remaining six
cultivars. This enabled us to identify sets of co-expressed cultivar-specific genes (**Supp. Table 7**).
We visualised 12 of these cultivar-specific network fragments using igraph[18] including a set of
18 linked genes with increased expression in spike tissue in Claire compared to other cultivars
(**Figure 3D & 3E**). The most highly connected gene in this subnetwork (TraesCS7A02G446400) is
a transducin/WD40 repeat protein. These proteins are key regulators of both plant
developmental and stress processes, and are known to participate in histone modification,
transcriptional regulation and signal transduction[19]. Additional genes in this cluster, are
annotated as protein phosphatases, an eRF1 transcription factor, a calcium-binding EF-hand
domain-containing protein and a polyadenylation specificity factor. We hypothesise that the
cultivar-specific expression pattern observed in Claire linked to increased expression in spike
tissues could be the result of cultivar-specific regulation of a developmental or stress response.
376
Co-expression network analysis using *de novo* gene models
Our consensus network approach, using a common reference, enabled us to identify high
confidence, conserved expression modules and identify cultivar-specific co-expressed gene
sets. However, the use of a common reference meant that we were unable to assess the *de*
novo contribution of each genome to the consensus network. Of the seven cultivars within
the consensus network, Arina*LrFor*, Jagger, Julius and Norin 61 each have corresponding *de*
novo annotations. We used these *de novo* gene models to identify a total of 4,682 *de novo*
annotated genes without a corresponding CSv1.1 orthologue and used expression counts from
these *de novo* gene models to build a *de novo* co-expression network of 13 modules (3,975
genes, **Supp. Table 8**). Each of these 13 modules could be closely correlated with at least one
consensus module from the consensus network (**Supp. Table 8**), indicating that our *de novo*
modules were not exhibiting patterns of expression distinct from those previously identified in
the seven-cultivar consensus network. One of these *de novo* derived modules was significantly
enriched ($p<0.000003$) for Jagger *de novo* gene models with increased expression in flag leaf,
spike and root tissues (**Extended Data Figure 7**). The three most significant GO terms enriched
within this module of 50 genes indicated a role in transcriptional regulation (GO:0065007,
GO:0031323, GO:0050789). 15 of these 50 genes were also annotated as transcription factors
with FHY3/FAR1 DNA binding domains (**Supp. Table 8**). These domains are known to be
involved in phytochrome signalling in *Arabidopsis*[20] and in wheat are hypothesised to
contribute to the regulation of *Ppd-B1a* and *PhyC* known to control photoperiodic sensitivity to
flowering[21].
398
Our work demonstrates the strength of a consensus network approach in identifying potentially
biologically conserved pathways between cultivars where *de novo* annotations are not available

for each member of the network. Using this method we were able to reveal cultivar-specific co-
expressed genes for several cultivars including Claire and Weebill, for which we do not currently
have *de novo* gene models. In addition, further extending our co-expression analysis to include
the *de novo* gene models of four of the chromosome level assemblies revealed additional *de*
novo co-expressed modules exhibiting cultivar-specificity, such as the *de novo* module enriched
for Jagger genes, that would not have been captured in the consensus network.
407
408   Developing *de novo* annotations for all 14 of the cultivars within the wheat pan-genome[17] will
409   be invaluable in uncovering the complete regulatory network landscape of the wheat pan-
410   transcriptome. Associating these co-expression profiles with the core, shell and cloud
411   components of the wheat pan-genome will enable us to explore how structural rearrangements
412   and introgressions across the wheat genome perturb these regulatory networks.
413

A case study: Uncovering variation in the prolamin super-family and immune reactive
proteins across the pan-cultivars

Prolamins represent a large superfamily in wheat involved in stress responses, cell growth and
plant development, as well as end-use quality and protein content. Along with HMW-glutenins
they are also potential triggers for various immune reactions in a subset of the human
population. As a case study, we investigated both the qualitative and quantitative differences in
the 687 genes from the prolamin superfamily and HMW-glutenins across the newly generated
wheat pan-genome and pan-transcriptome data. We observed clear expression differences
both for individual developmental stages and also between wheat cultivars for many genes
from the prolamin superfamily highlighting spatiotemporal variation in expression profile
(**Figure 4A**).
425

Comparison of reference grain allergens identified in the Chinese Spring reference genome
(IWGSC v1.1) and across the pan-genome cultivars[22,23,24] with the expression patterns of
potentially immune reactive gene products indicated differences in the major allergens and
antigens (glutenins and gliadins). SY Mattis and LongReach Lancer showed lower gene
expression levels in alpha and gamma gliadins with gene set enrichment analysis of gene
families highlighting gamma gliadins are primarily enriched in the downregulated genes
(**Extended Data Figure 8, Supp. Table 9&10**).
433

Detailed analysis of celiac disease (CD) related epitopes encoded in the gliadin and glutenin
genes in the pan-genome revealed variability in their expression patterns. We found lower
expression of HLA-DQ epitope containing genes in SY Mattis and LongReach Lancer and higher
values in Cadenza and Jagger. Cultivar-specific analysis showed that Arina*LrFor* and SY Mattis
contained lower alpha gliadin DQ epitope expressions due to significant differences in the
expression activities of the three sub-genomes which might be affected by differences in the
related transcription factor gene expression profiles (**Figure 4B, 4C, Supp. Table 11&12,**
Extended Data Figure 9). While sub-genome specific expression patterns of gamma gliadin DQ
epitopes did not reveal significant variation, the expression of alpha-gliadin genes with DQ
epitopes originating from the A genome was lower in SY Mattis and LongReach Lancer, while
the highly immunogenic D genome alpha gliadin epitope expression levels were lower in the

cultivar Arina*LrFor* (**Extended Data Figure 10A**). Our results indicate that fine-tuned sub-
genome-specific balance in the expression profiles may be associated with differences in the
regulatory transcription factor profiles (**Figure 4B, 4C**).
448

449
Figure 4. **Gene and expression variation in the prolamin family across the wheat pan-**
cultivars. A. Prolamin superfamily gene expression across cultivars; **B**. Celiac disease epitope
expression across cultivars. Epitope expression profiles were calculated as sum of gene
expression profiles with the highlighted HLA-DQ epitopes for each sub-genome. **C**. Relative
proportion of cumulative expression profiles of transcription factor families showing strong co-
expression pattern (Pearson correlation values > 0.8) with the epitope-coding prolamin genes.
Results show significant differences in the NAC, AP2/EREBP and MYB transcription factor gene
expressions, major regulators of storage protein gene expression. **D**. Representation of the
variation graph for the region of 6D containing the alpha gliadin locus (**Extended Data Figure**
10B). Horizontal coloured lines depict paths through the graph for each cultivar; Norin 61 (6D:
26,703,647-27,222,360 bp), CDC Stanley (6D: 28,164,601-28,660,350 bp) and Mace (6D:
26,808,846-27,298,593bp), with SY Mattis (6D: 26,645,382-27,096,594 bp) and Julius (6D:
26,983,100-27,437,565 bp) sharing a single path. Rectangular blocks (a-p) represent individual

genes at corresponding locations across cultivars (blue: in common to all 4 cultivars, orange:
occurring in one cultivar and purple: in common to 2 cultivars). Gene d is present as a single
copy in Norin 61, and duplicated in CDC Stanley, SY Mattis, Julius and Mace. This duplication is
represented as a loop in the path through the graph for these cultivars (**Extended Data Figure**
11).
468
Gliadin and glutenin loci were found to be highly conserved in all cultivars, with some variation
due to the presence of pseudogenes and gene duplications (**Extended Data Figure 10B**).
Reverse translated consensus sequences of the known CD-specific T-cell epitopes were mapped
to the genomes of all cultivars to determine the number and location of gliadin and glutenin
genes containing CD-related immune reactive peptide regions (**Extended Data Figure 10B,**
Supp. Table 13). The number or combination of epitopes in the loci was not significantly
different between the pan-cultivars. However, the gamma-gliadin and alpha-gliadin gene
models with a high number of epitopes were found in cultivars Arina*LrFor*, Norin 61 and Mace,
respectively (**Extended Data Figure 10B, Supp. Table 13**).
478
Although highly conserved in their locus structure on chromosome 6D, alpha-gliadin genes
encoding highly immunogenic proteins showed copy number variation within the wheat pan-
genome. We constructed a localised pan-genome graph from five cultivars (Norin 61, CDC
Stanley, SY Mattis, Julius, Mace) and extracted the subgraph of the alpha gliadin-containing
locus (**Figure 4D, Extended Data Figure 11**). Inspection of the subgraph helped to resolve the
complex structure of the locus, with copy number variation observed as a loop in the paths of
SY Mattis, Julius, CDC Stanley, and Mace (2 copies of alpha-gliadin genes) but not within the
Norin 61 path (single alpha gliadin copy). While in total 4 to 6 epitopes were identified in the
alpha-gliadins of the wheat pan-genome cultivars, 8 epitopes were detected in cultivars Mace
and Norin 61 (**Extended Data Figure 10B**). These results indicate that gene copy number
expansion primarily affected the centre of the locus and resulted in the increase of gene
variants with high epitope counts. Comparison of promoter profiles indicates differences in the
expression regulation when epitope-poor and epitope-rich gene copy variants of the same
chromosome 6D locus are compared. While genome-wide construction and interpretation of
pan-genome graphs remains a daunting task for complex genomes such as wheat, we found
localised subgraphs, augmented by our *de novo* annotations, particularly helpful in resolving
complex loci, and uncovering structural variation as also demonstrated in the current draft
human pan-genome[25].
497
498
Discussion
We have built *de novo* gene annotations for nine wheat assemblies representative of global
breeding programs[4]. Our consolidated gene annotation approach generated a robust set of
core, high-confidence genes shared across the pan-cultivars. It also identified genes and gene
families that are found exclusively in or amplified in cultivars derived from specific breeding
programmes. It is likely that some of this variation has come through widespread introgression
events[26], often associated with adaptation to biotic or abiotic stress[13]. Our annotations also
identified cultivar-specific variation in tandem gene duplication. Novel gene content, gene

duplication and neo-functionalisation together with gene expression patterns will have an
impact on researchers and breeders as they identify genes underlying traits, manipulate gene
expression or incorporate and track new genetic variation.
510
Our analysis of global gene expression identified sets of genes with stable homoeologous
expression patterns between cultivars, demonstrating tightly regulated key biological
processes. We also identified homoeologous triads diverging in their expression patterns
between cultivars, revealing genes enriched for processes associated with biotic and abiotic
stress. Understanding the regulatory networks driving these altered patterns will provide
important targets for manipulating these processes. Using network analysis, we identified
widespread conservation of expression patterns across tissues and cultivars before focusing on
cultivar-specific gene sets, to reveal networks of genes involved in stress responses in the
developing grain and the photoperiodic control of flowering. These cultivar-specific network
changes may be the result of wheat breeding programmes targeted to local environments. We
also demonstrated the utility of our new resources by investigating genomic variation in the
prolamin superfamily, focusing on immunogenic potential.
523
524 In conclusion, this study reveals layers of hidden diversity spanning our modern wheat cultivars.
525 Previously overlooked, this diversity is likely to underpin the agronomic success of wheat over a
526 wide range of global mega-environments.
527
Materials and Methods

Plant Materials and Growth Conditions

The 14 cultivars were grown in a Controlled Environment Room (CER) (Conviron BDW80;
Conviron, Winnipeg, Canada) set at 16 h day/8 h night photoperiod (300 µmol m^{-2} s^{-1}, lights on
at 05:00, lights off at 21:00), temperatures of 20/16 °C, respectively, and 60% relative
humidity. Plants were sampled in triplicate at the 3-leaf stage (Zadoks GS13), harvesting whole
roots and whole aerial organs separately, four hours after dawn (09:00). Whole aerial organs
were also sampled two hours after dusk (23:00). Plants for subsequent adult plant sampling
were treated according to their vernalisation requirements. In the case of spring wheat
cultivars (CDC Landmark, CDC Stanley, Paragon, Cadenza, Mace and LongReach Lancer),
seedlings were grown as described above. At 3-leaf stage, seedlings were transferred to 1 L
pots containing Petersfield Cereal Mix (Petersfield, Leicester, UK) and maintained under the
same CER conditions as described previously. For winter wheat cultivars (Norin 61, Julius,
Jagger, Arina*LrFor*, Robigus, Claire and SY Mattis), seedlings were transferred in 40-well trays (7
days after sowing) to a vernalisation CER running at 6 °C with 8 h day/16 h night photoperiod
for 61 days. After this period the plants were transferred to 1 L pots containing Petersfield
Cereal Mix (Petersfield, Leicester, UK) and moved to the same CER and settings as described for
the spring wheat cultivars. For both spring and winter wheat cultivars, three additional samples
were harvested: complete spike at heading (GS59), flag leaf 7 days post-anthesis (GS71) and
whole grains 15 days post-anthesis (GS77). All samples were harvested four hours after dawn
(09:00), and a single plant was used per each of the three biological replicates.

Sample Preparation and Sequencing

Total RNA was extracted using Qiagen RNeasy Plant Mini Kit (cat. no. 74904) and DNAse treated using an Invitrogen TURBO DNase kit (cat. no. AM2238) according to the manufacturer's protocol. Bead purification of the RNA was conducted using the Agencourt RNAClean XP beads.system (cat. no. A63987). Final sample concentrations were verified using a Qubit 4 Fluorometer, and the integrity of the RNA was checked on the Agilent 2100 Bioanalyzer, using the RNA 6000 nano kit (Agilent, 5067-1511), running the plant total RNA assay. The directional RNA-seq libraries were constructed using the NEBNext Ultra II Directional RNA Library prep for Illumina kit (NEB, E7760L) utilising the NEBNext Poly(A) mRNA Magnetic Isolation Module (NEB, E7490L) and NEBNext Multiplex Oligos for Illumina (96 Unique Dual Index Primer Pairs) (cat. no. E6440S/L) at a concentration of 10 µM. The final libraries were equimolar pooled, a q-PCR was performed and the pool was sequenced on a Illumina NovaSeq 6000 with 150 bp paired-end reads.

The Iso-Seq libraries were constructed from 1 µg of total RNA per sample and full-length cDNA were then generated using the SMARTer PCR cDNA synthesis kit (Takara Bio Inc, 639506). The libraries were sequenced on the Sequel Instrument v1, using 1 SMRTcell v2 per library. All libraries had 600-minute movies, 120 minutes of immobilisation time, and 120 minutes of pre-extension time.

Data Quality Control and Sample Validation

We used a set of cultivar specific SNPs to confirm the cultivar origin of each replicate and the developmental stage of each sample was validated through a machine learning approach trained using the pooled RNA-seq samples and then run on the entire set of biological replicates. Principal component analysis of the pooled samples shows them to cluster by developmental stage as expected.

Alignment and Gene Expression Analysis

Samples were aligned to the IWGSC Chinese Spring RefSeq 1.1 reference genome, using HISAT2 v2.0.4, acting as a common reference to allow inclusion of UK cultivars and comparison with *de novo* annotations, and normalised counts were generated using DESeq2 with the RefSeq 1.1 gene set[27,16,28].

GO term analysis

Functional enrichment of differentially expressed genes for biological processes was performed using the gene ontology enrichment analysis package, topGO[29] in R (v3.6.0, with the following parameters: nodeSize = 10, algorithm = "parentchild", classicFisher test p < 0.05). GO terms refer to ontology terms for biological processes unless otherwise specified and were obtained from Ensembl Plants 51, using the BioMart tool. Bubble plots were plotted using ggplot in R, adapting code from[30].

Tissue-Specific Index

The specificity of gene expression to developmental stages was determined using the tissue-specific index[31]. Where N is the number of developmental stages (condition), and x_i is the

593 expression profile component for a given gene in each condition, normalised by the maximal
594 expression value of the given gene from all conditions considered. This allowed us to classify
595 genes as being highly specific to one condition (tau => 0.8). Assignment of tau values was
596 performed in R using code adapted from previous work[32].
597
Sub-genome Expression Bias
Analysis of sub-genome expression focused on 30-let homoeologs with a 1:1:1 relationship
across all three sub-genomes. Of these, 13,521 were determined to be macrosyntenic,
belonging to the same sub-genome in all cultivars (excluding UK cultivars which are not
assembled), and 10,653 as microsyntenic, belonging to the same chromosome and sub-genome
in all cultivars (excluding UK cultivars). From these 66 30-lets were not taken forward in the
analysis due to low expression and/or quality filtering determined by DESeq2 (R package v
4.0.3) of at least one homoeolog in each set. Relative expression of 30-lets across homoeologs
and associated sub-genome expression biases were calculated as previously reported, through
use of our triad.expression R package (https://github.com/AHallLab/triad.expression).
608
Co-expression Analysis
Network construction
The WGCNA R package[17] (R version 3.6.0) was used to build co-expression networks for seven
cultivars (Arina*LrFor*, Cadenza, Claire, Jagger, Julius, Norin 61 and Weebill) for which we had
triplicate biological replicates for each developmental stage. These cultivars also span the range
of genetic variation observed in the previously published pan-genome[4]. The expression matrix
for the seven selected cultivars containing DESeq2 normalised counts aligned to 102,443 high
confidence CSv1.1 genes was filtered and genes, where the sum of counts across all samples
was greater than 5 in at least 3 samples, were retained (92,976 genes). To reduce background
noise we removed genes where the expression of any replicate was >2σ from the mean
expression of that sample set. The resulting 48,337 genes were submitted to WGCNA to
construct a signed hybrid consensus network using the blockwiseConsensusModules ()
function. A soft power threshold of 18 was used, together with the following parameters;
minModuleSize = 30, corType = bicor, maxPOutliers = 0.05, mergeCutHeight = 0.3,
minKMEtoStay = 0.2, maxBlockSize = 46,000. Eigengenes were then extracted for each module,
per cultivar from the resulting consensus network.
625
Defining thresholds for classifying inter-module relationships
To classify inter-module relationships and identify modules with divergent or similar patterns of
expression we defined a threshold of module similarity. Initially we calculated the distance
between each pairwise consensus module comparison, using the Pearson correlation distance.
We used the maximum distance of each of these pairwise comparisons, for each module and
calculated the median of these maxima. Next, we investigated the proportion of 1,663 triads
(from 30-lets) identified as split across the 23 previously defined consensus modules, that
would be classed as divergent using a module similarity threshold of 0-100%. From these results
we selected a module similarity threshold of 75% the median of maximum distances, with
distances above this classed as divergent and distances below, classed as similar.
636

Identifying metamodules
We used the R package clValid[33] to determine the optimal number of clusters for the 7 cultivar
23 consensus module eigengenes. The resulting Dunn index[34] and silhouette width[35] indicated
that the optimum number of clusters for our ME dataset was 6. We calculated the pairwise
Pearson correlation coefficients for all our 161 cultivar consensus ME (cor()) and converted this
to a dissimilarity matrix (as.dist()). We used hierarchical clustering of this dissimilarity matrix
(hclust()) to define consensus metamodules. As the JAG magenta consensus module fell into a
different metamodule to the remaining 6 cultivars we omitted the magenta consensus module
from our metamodule construction and downstream enrichment analysis. We used the
moduleEigengenes() function to compute the ME of each metamodule and carried out GOterm
and TF superfamily enrichment analysis.
648
649 *Transcription factor superfamily enrichment*
650 Genes annotated as members of transcription factor (TF) superfamilies[5] were identified in each
651 metamodule and the frequency of each TF superfamily compared to the frequency observed in
652 the 32,936 genes used to construct metamodules. TF families were classed as either
653 significantly under or overrepresented in each module using Fisher's exact test (p <= 0.05).
654
Identifying cultivar-specific expression patterns
We used the consensuskME() function in WGCNA to determine the maximum eigengene-based
connectivity (kME) of each gene within the unclustered gene set of 14,268 genes, to the
consensus ME for each cultivar. Those genes with a positive association greater than 0.7 to a
consensus ME were retained. To reveal putative networks specific to a single cultivar we
identified genes from the 13,708 genes that demonstrated associations greater than 0.7 kME
for at least one cultivar, per gene, where a minimum of 4 out of the 7 cultivars exhibited >0.7
kME and that in pairwise comparison to all other cultivars, for the specific cultivar being
assessed, the gene was assigned to a divergent module.
664
Connectivity within each set of genes demonstrating single cultivar-specificity was determined
using the R package igraph[18]. Using the graph adjacency() function, graph adjacencies were
created for each specific cultivar set based on the Pearson correlation distances between genes
in a pairwise fashion. These directed graphs were simplified to remove multiple edges and
loops, and filtered to retain only those connections with an absolute Pearson correlation > 0.8.
The mst() function using the prim algorithm was used to create a minimum spanning tree and
the resulting subgraphs were visualised using plot() with isolated nodes excluded.
672
Gene annotation
For the structural gene annotation of the chromosome-scale assembled pan-cultivars, we
combined de novo gene calling and homology-based approaches with RNAseq, Isoseq, and
protein datasets. The RNAseq data were mapped using STAR[36] (v2.7.8a) and further assembled
into transcripts by StringTie[37] (v2.1.5, parameters -m 150-t -f 0.3). PacBio Iso-Seq transcripts
were derived from the raw reads using PacBio SMRT Link software (v5.1.0.26412rev2,
pbsmrtpipe.pipelines.sa3_ds_isoseq2, default parameters). The Iso-Seq transcripts were
aligned to the genome assemblies using GMAP[38] (v2018-07-04). To assist the homology-based

annotation approach, Triticeae protein sequences from publicly available datasets (UniProt,
https://www.uniprot.org, 05/10/2016) were aligned against the genome sequence assemblies
of all pan-cultivars using GenomeThreader[39] (v1.7.1; arguments -startcodon -finalstopcodon -
species rice -gcmincoverage 70 -prseedlength 7 -prhdist 4). All transcripts derived from RNAseq,
IsoSeq, and aligned protein sequences were combined using Cuffcompare [40] (v2.2.1). Stringtie
(version 2.1.5, parameters --merge -m150) was employed to merge all sequences into a pool of
candidate transcripts. To identify potential open reading frames and to predict protein
sequences within the candidate transcript set, TransDecoder (version 5.5.0;
http://transdecoder.github.io) was used.
690
691 We used Augustus[41] (v3.3.3) for the *ab initio* gene prediction. Guiding hints based on the
692 RNAseq, protein, IsoSeq and TE datasets described above were used to counteract potential
693 over-prediction (details in[42]). Augustus was run using the wheat model.
694
695 A consolidated set of gene models was selected using Mikado[43], as implemented in the Minos
696 pipeline (https://github.com/EI-CoreBioinformatics/minos), with models scored and selected
697 based on a combination of intrinsic qualities and support from transcriptome and protein
698 alignments.
699
BLASTP[44] (ncbi-blast v2.3.0+, parameters -max_target_seqs 1 -evalue 1e-05) was used to
compare potential protein sequences with a trusted set of reference proteins (Uniprot
Magnoliophyta, reviewed/Swissprot, downloaded on 3 Aug 2016; https://www.uniprot.org).
This approach was employed to differentiate gene candidates into complete and valid genes,
non-coding transcripts, pseudogenes, and transposable elements. This step was assisted by
PTREP (Release 19; http://botserv2.uzh.ch/kelldata/trep-db/index.html), a database of
hypothetical proteins containing deduced amino acid sequences in which internal frameshifts
have been removed in many cases. We selected best hits for each predicted protein from each
of the three databases used. Only hits with an e-value below 10e-10 were considered.
Functional annotation of all protein sequences predicted in our pipeline was performed with
the AHRD pipeline (https://github.com/groupschoof/AHRD).
711
We classified predicted proteins into two confidence classes: high and low confidence. Hits with
subject coverage (for protein references) or query coverage (transposon database) greater than
80% were considered significant and protein sequences were classified as high-confidence
based on following criteria: protein sequence was complete and had a subject and query
coverage above the threshold in the UniMag database or no BLAST hit in UniMag but in UniPoa
and not PTREP; a low-confidence protein sequence was incomplete and had a hit in the UniMag
or UniPoa database but not in PTREP. Alternatively, it had no hit in UniMag, UniPoa, or PTREP,
but the protein sequence was complete. In a second refinement step, low-confidence proteins
with an AHRD-score of 3* were promoted to high-confidence.
721
722 BUSCO[45] (v5.1.2.) software was used to evaluate the completeness and accuracy of structural
723 gene predictions with the 'poales_odb10' database containing a total of 4896 single-copy

724 genes. The evidence-based part of the annotation pipeline is deposited at
725 https://github.com/PGSB-HMGU/plant.annot.
726

Consolidation
Pairwise whole genome alignments were generated using lastz[46]. The resulting alignments were
stitched together into a single whole genome alignment using TBA/multiz[47]. The MAF output
was converted into HAL format using maf2hal[48].
De novo gene annotation from one cultivar was lifted over to all other cultivars using the whole
genome alignment and the halLiftover tool, whereas only full-length gene models were kept.
Missing gene models in one cultivar were identified using bedtools[49].
734

Tandem array detection
Tandem arrays were identified using the tandem discovery model from the JCVI package[50].
Expression bias was calculated using a modified method described previously[5]. Here we used
normalised read counts instead of TPM values and a cut-off of 0.8. The following categories
were assigned: only1 for tandems with only one gene expressed and no expression data for
second gene; expressed1 for tandems in which only one gene is expressed under all RNASeq
conditions; variable where expression can shift between array members depending on the
condition; balanced, where both array members are equally expressed. noExpr states that no
expression data was available.
744

Orthogroup analysis
The longest isoforms from high-confidence genes were used as input for Orthofinder[51].
Orthofinder was run using standard parameters. We used the UpSetR in the R package
(http://gehlenborglab.org/research/projects/upsetr/) to analyse and visualise how many
orthogroups are shared between the cultivars or are unique to a single species. GENESPACE[9]
was used to derive and visualise syntenic relationships between all chromosomes and sub-
genomes.
752

Analysis of Canadian-specific genes
Taking each genome in turn as a reference, kmers of length 51 were identified from genic
regions using the annotation for that reference. These kmers were used to search the genomes
of the other cultivars and a coverage score was computed[52] between each gene in the
reference and every other genome. The coverage score (a value between 0 and 1) can be used
as a proxy for sequence similarity/difference between genes in different cultivars where values
closer to 0 indicate greater difference and values closer to 1 indicate similarity. Coverage scores
for genes along chromosomes were plotted using the seaborn visualisation library[53] in Jupyter
notebook. Coverage scores were also visualised as heatmaps with coverage scores close to 0
represented as dark bands.
763

Comparative analysis of immune reactive regions in the wheat pan-genome
Reference allergen identification and chromosome 6D comparison
Reference allergens in the wheat pan-genome were filtered using blastn algorithm against the
identified sequences in the IWGSC v1 gene annotation v1.1[22]. To identify unannotated gliadin

and glutenin gene models and to compare the potential immune reactivity of the wheat
cultivars, known coeliac diseases associated HLA-DQ T-cell epitopes were reverse translated
and the consensus nucleotide sequences were used for a motif search with 100% sequence
identity. The mapped epitope-rich regions were used for the detailed comparison of the alpha
gliadin locus in chromosome 6D. Additional gene models representing complete gene models
with DQ epitopes were manually annotated. The locus organisation was compared to the
Chinese Spring chromosome 6D alpha gliadin locus in the IWGSC v1 reference genome
assembly[22].
776
777 *Promoter motif enrichment analysis*
778 1000 bp 5'-end non-coding sequences were extracted from the chromosome 6D loci and used
779 for motif enrichment analysis in MEME-SEA[54]. The JASPAR core plant 2022 motif collection was
780 used as a background database.
781
782 *Epitope expression analysis*
783 Epitope expression values were calculated using the TPM gene expression values of genes
784 where the reverse translated consensus epitope sequence was detected multiplied by the
785 number of epitopes in each sequence. The obtained values were summed for each epitope type
786 as well as summed for epitope types at genome levels.
787
788 *Gene co-expression analysis*
789 TPM>1 log2 transformed TPM gene expression data were used to create a grain co-expression
790 network using co-expression cut-off value of 0.8. The resulting network was annotated with the
791 reference allergen-specific information for disease relatedness and gene family. The first
792 neighbour network was visualised in Cytoscape.
793
Pan-genome graph construction of 10Mb 6D region
We extracted a 10Mb region (20-30Mb) encompassing the alpha gliadin locus from the top of
chromosome 6D for the cultivars Norin 61, CDC Stanley, SY Mattis, Julius, and Mace. To
estimate the divergence of the input sequences, we used mash-2.2[55], specifically the mash
triangle command to calculate a maximum sequence divergence of 0.039. To account for
possible underestimation of sequence divergence and localised structural variants we specified
a minimum mapping identity value (-p 90) for pan-graph construction using PGGB[56] together
with segment size (-s 30kb), number of mappings (-n 6), minimum length of exact matches (-k
311), target sequence length for POA (-G 13117, 13219), mean length of each sequence pad for
POA (-O 0.03) and k-mer size for mapping (-K 111). Default settings were used for all other
parameters.
805
Extracting the alpha-gliadin locus sub pan-graph
Using ODGI toolkit[57] we extracted the subgraph of the alpha-gliadin locus from our 6D graph
build. We used the odgi extract command together with coordinates of the Norin 61 gene
models described in **Supp. Table 14** extracts the 520.7kb region encompassing the locus (6D:
26,703,647-27,222,360bp) and the corresponding paths intersecting with this region in CDC
Stanley (6D: 28,164,601-28,660,350 bp), SY Mattis (6D: 26,645,382-27,096,594 bp), Julius (JUL

6D: 26,983,100-27,437,565 bp), and Mace (6D: 26,808,846–27,298,593 bp). We used odgi sort
to sort the resulting subgraph and odgi procbed to adjust the coordinates of the gene models
for each cultivar to fit the resulting subgraph. odgi inject allowed us to visualise the placement
of these gene models across the graph and identify cultivar-specific haplotypes. We generated
a graphical fragment assembly (gfa) of this sub pan-graph using odgi view (**Supp. Table 15).**

Data Availability
The genome sequence and gene annotations of all wheat cultivars can be viewed and
downloaded in Ensembl Plants (https://plants.ensembl.org/index.html). This includes the *de*
novo genes for the chromosome-level cultivars generated within this study and projected genes
for all assemblies from the IWGSC RefSeq v1.1 annotation. All raw data used in this study is
available at the European Nucleotide Archive under accession PRJEB51827.

Code Availability
Relevant code repositories are referenced throughout the Methods sections.

Supplementary Tables
All supporting tables and associated materials are available at
https://opendata.earlham.ac.uk/wheat/under_license/toronto/Hall_2024-01-
01_wheat_pantranscriptome.

Acknowledgements
We would like to acknowledge BBS/E/T/000PR9816 (NC1 - Supporting EI's ISPs and the UK
Community with Genomics and Single Cell Analysis) for data generation and BB/CCG1720/1 for
the physical HPC infrastructure and data centre delivered via the NBI Research Computing
group. Earlham Institute Strategic Programme Grant Decoding Biodiversity BBX011089/1
(BBS/E/ER/230002B). Delivering Sustainable Wheat BB/X011003/1 (BBS/E/ER/230003C).
Norwich Research Park Biosciences Doctoral Training Partnership grant BB/M011216/1. We
also would like to thank the Australian Research Council Centre of Excellence for Innovations in
Peptide and Protein Science (CE200100012) and Coeliac Australia (G1005443) as well as MEXT
JSPS KAKENHI 22H05179, 22H02316, 22K21352, Swiss National Science Foundation
310030_212551, Japan Science and Technology Agency JPMJCR16O3 and URPP Evolution in
Action. Helmholtz Munich would like to acknowledge support by BMBF project number
031B0190 (de.NBI). We would also like to thank the 10+ Wheat Genome Project for their
feedback and guidance.

Author Contributions
CP arranged supplying seeds. JS & CU managed growing plants and supplying materials. SD &
HR prepared samples and performed extractions for sequencing. NI, SH, TB, HC, LC and KG
generated and managed the iso-seq and RNA-seq data production. BW, RJ & JC performed
quality control of biological replicate data. SW & BW reworked the existing package for sub-
genome expression bias characterisation. BW carried out gene expression analysis, tissue-
specific indexing and investigation of sub-genome biases. TL, GK & DS performed *de novo*
annotations. TL, DL & HG consolidated *de novo* annotations, generated orthogroups, and

investigated introgressions and copy number variation. RRP performed co-expression analysis, constructed regulatory networks and gliadin pan-graph, and investigated cultivar-specific expression. JW performed genome kmer comparison. AJ performed an analysis of allergen and prolamin diversity, MG supervised the prolamin analysis. KS investigated Norin-specific gene expression patterns. GN imported and managed data for EnsemblPlants. BW, TL, RRP, JW, CU, MS & AH contributed to preparing the manuscript. PB, WH, BC, CU, NS, SK, JP, KM & CP provided intellectual input. Syngenta provided funding for data generation. The 10+ Wheat Genome Project, EL, AB, AH and MS for conceptualisation of the project.

Ethics declarations

The authors declare no competing interests.

References

1. Braun, H.-J., Rajaram, S. & Ginkel, M. van. CIMMYT's approach to breeding for wide adaptation. *Euphytica* **92**, 175–183 (1996).

2. Erenstein, O. *et al.* Wheat Improvement, Food Security in a Changing Climate. 47–66 (2022) doi:10.1007/978-3-030-90673-3_4.

3. Levy, A. A. & Feldman, M. Evolution and origin of bread wheat. *Plant Cell* **34**, 2549–2567 (2022).

4. Walkowiak, S. *et al.* Multiple wheat genomes reveal global variation in modern breeding. *Nature* **588**, 277–283 (2020).

5. Ramirez-Gonzalez, R. H. *et al.* The transcriptional landscape of polyploid wheat. *Science (New York, NY)* **361**, eaar6089 (2018).

6. Jin, M. *et al.* Maize pan-transcriptome provides novel insights into genome complexity and quantitative trait variation. *Scientific Reports* **6**, 18936 (2016).

7. Thind, A. K. *et al.* Chromosome-scale comparative sequence analysis unravels molecular mechanisms of genome dynamics between two wheat cultivars. *Genome Biol.* **19**, 104 (2018).

8. Khan, A. W. *et al.* Super-Pangenome by Integrating the Wild Side of a Species for Accelerated Crop Improvement. *Trends Plant Sci.* **25**, 148–158 (2020).

9. Lovell, J. T. *et al.* GENESPACE tracks regions of interest and gene copy number variation across multiple genomes. *eLife* **11**, e78526 (2022).

10. He, F. *et al.* Genomic variants affecting homoeologous gene expression dosage contribute to agronomic trait variation in allopolyploid wheat. *Nat. Commun.* **13**, 826 (2022).

11. Leister, D. Tandem and segmental gene duplication and recombination in the evolution of plant disease resistance genes. *Trends Genet.* **20**, 116–122 (2004).

12. Hanada, K., Zou, C., Lehti-Shiu, M. D., Shinozaki, K. & Shiu, S.-H. Importance of Lineage-Specific Expansion of Plant Tandem Duplicates in the Adaptive Response to Environmental Stimuli . *Plant Physiol.* **148**, 993–1003 (2008).

13. Bayer, P. E., Golicz, A. A., Scheben, A., Batley, J. & Edwards, D. Plant pan-genomes are the new reference. *Nat. Plants* **6**, 914–920 (2020).

14. He, F. *et al.* Exome sequencing highlights the role of wild-relative introgression in shaping
the adaptive landscape of the wheat genome. *Nat. Genet.* **51**, 896–904 (2019).

15. Gordon, S. P. *et al.* Extensive gene content variation in the Brachypodium distachyon pan-
genome correlates with population structure. *Nat. Commun.* **8**, 2184 (2017).

16. IWGSC, T. I. W. G. S. C. *et al.* Shifting the limits in wheat research and breeding using a fully
annotated reference genome. *Science (New York, NY)* **361**, eaar7191 (2018).

17. Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network
analysis. *BMC Bioinform.* **9**, 559 (2008).

18. Cśardi, G. & Nepusz, T. The igraph software package for complex network research. in.

19. Hu, R. *et al.* Genome-wide identification and analysis of WD40 proteins in wheat (Triticum
aestivum L.). *BMC Genom.* **19**, 803 (2018).

20. Li, J., Li, G., Wang, H. & Deng, X. W. Phytochrome Signaling Mechanisms. *Arab. Book* **2011**,
e0148 (2011).

21. Kiseleva, A. A., Potokina, E. K. & Salina, E. A. Features of Ppd-B1 expression regulation and
their impact on the flowering time of wheat near-isogenic lines. *BMC Plant Biol.* **17**, 172 (2017).

22. Juhász, A. *et al.* Genome mapping of seed-borne allergens and immunoresponsive proteins
in wheat. *Sci. Adv.* **4**, eaar8602 (2018).

23. Halstead-Nussloch, G. *et al.* Multiple Wheat Genomes Reveal Novel Gli-2 Sublocus Location
and Variation of Celiac Disease Epitopes in Duplicated α-Gliadin Genes. *Front. Plant Sci.* **12**,
715985 (2021).

24. Shimizu, K. K. *et al.* De Novo Genome Assembly of the Japanese Wheat Cultivar Norin 61
Highlights Functional Variation in Flowering Time and Fusarium-Resistant Genes in East Asian
Genotypes. *Plant Cell Physiol.* **62**, 8–27 (2020).

25. Liao, W.-W. *et al.* A draft human pangenome reference. *Nature* **617**, 312–324 (2023).

26. Cheng, H. *et al.* Frequent intra- and inter-species introgression shapes the landscape of
genetic variation in bread wheat. *Genome Biol.* **20**, 136 (2019).

27. Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome alignment
and genotyping with HISAT2 and HISAT-genotype. *Nat. Biotechnol.* **37**, 907–915 (2019).

28. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

29. Rahnenfuhrer, A. A. topGO: Enrichment Analysis for Gene Ontology. R package version
2.46.0. (2021).

30. Vega, J. J. D. *et al.* Differential expression of starch and sucrose metabolic genes linked to
varying biomass yield in Miscanthus hybrids. *Biotechnol. Biofuels* **14**, 98 (2021).

31. Yanai, I. *et al.* Genome-wide midrange transcription profiles reveal expression level
relationships in human tissue specification. *Bioinformatics* **21**, 650–659 (2005).

32. Kryuchkova-Mostacci, N. & Robinson-Rechavi, M. A benchmark of gene expression tissue-
specificity metrics. *Brief. Bioinform.* **18**, 205–214 (2017).

33. Brock, G., Pihur, V., Datta, S. & Datta, S. clValid: An R Package for Cluster Validation. *Journal*
of Statistical Software **25**, 1–22 (2008).

34. Dunn†, J. C. Well-Separated Clusters and Optimal Fuzzy Partitions. *J. Cybern.* **4**, 95–104
(1974).

35. Rousseeuw, P. J. Silhouettes: A graphical aid to the interpretation and validation of cluster
analysis. *J. Comput. Appl. Math.* **20**, 53–65 (1987).

36. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

37. Kovaka, S. *et al.* Transcriptome assembly from long-read RNA-seq alignments with
StringTie2. *Genome Biol.* **20**, 278 (2019).

38. Wu, T. D. & Watanabe, C. K. GMAP: a genomic mapping and alignment program for mRNA
and EST sequences. *Bioinformatics* **21**, 1859–1875 (2005).

39. Gremme, G., Brendel, V., Sparks, M. E. & Kurtz, S. Engineering a software tool for gene
structure prediction in higher organisms. *Inf. Softw. Technol.* **47**, 965–978 (2005).

40. Ghosh, S. & Chan, C.-K. K. Plant Bioinformatics, Methods and Protocols. *Methods Mol. Biol.*
1374, 339–361 (2015).

41. Hoff, K. J. & Stanke, M. Predicting Genes in Single Genomes with AUGUSTUS. *Curr. Protoc.*
Bioinform. **65**, e57 (2019).

42. Nachtweide, S. & Stanke, M. Gene Prediction, Methods and Protocols. *Methods Mol. Biol.*
1962, 139–160 (2019).

43. Venturini, L., Caim, S., Kaithakottil, G. G., Mapleson, D. L. & Swarbreck, D. Leveraging
multiple transcriptome assembly methods for improved gene structure annotation. *GigaScience*
7, giy093- (2018).

44. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment search
tool. *Journal of molecular biology* **215**, 403–410 (1990).

45. Seppey, M., Manni, M. & Zdobnov, E. M. Gene Prediction, Methods and Protocols. *Methods*
Mol. Biol. **1962**, 227–245 (2019).

46. Kiełbasa, S. M., Wan, R., Sato, K., Horton, P. & Frith, M. C. Adaptive seeds tame genomic
sequence comparison. *Genome Res.* **21**, 487–493 (2011).

47. Blanchette, M. *et al.* Aligning Multiple Genomic Sequences With the Threaded Blockset
Aligner. *Genome Res.* **14**, 708–715 (2004).

48. Hickey, G., Paten, B., Earl, D., Zerbino, D. & Haussler, D. HAL: a hierarchical format for
storing and analyzing multiple genome alignments. *Bioinformatics* **29**, 1341–1342 (2013).

49. Quinlan, A. R. BEDTools: The Swiss-Army Tool for Genome Feature Analysis. *Curr. Protoc.*
Bioinform. **47**, 11.12.1-11.12.34 (2014).

50. Tang, H. *et al.* Synteny and Collinearity in Plant Genomes. *Science* **320**, 486–488 (2008).

51. Emms, D. M. & Kelly, S. OrthoFinder: solving fundamental biases in whole genome
comparisons dramatically improves orthogroup inference accuracy. *Genome Biol.* **16**, 157
(2015).

52. Clavijo, B. J., Accinelli, G. G., Yanes, L., Barr, K. & Wright, J. Skip-mers: increasing entropy
and sensitivity to detect conserved genic regions with simple cyclic q-grams. *bioRxiv* (2017)
doi:10.1101/179960.

53. Waskom, M. seaborn: statistical data visualization. *J. Open Source Softw.* **6**, 3021 (2021).

54. Bailey, T. L. & Grant, C. E. SEA: Simple Enrichment Analysis of motifs. *bioRxiv*
2021.08.23.457422 (2021) doi:10.1101/2021.08.23.457422.

55. Ondov, B. D. *et al.* Mash: fast genome and metagenome distance estimation using MinHash.
Genome Biol. **17**, 132 (2016).

56. Garrison, E. *et al.* Building pangenome graphs. *Biorxiv Prepr Serv Biology* (2023)
doi:10.1101/2023.04.05.535718.

57. Guarracino, A., Heumos, S., Nahnsen, S., Prins, P. & Garrison, E. ODGI: understanding
pangenome graphs. *Bioinformatics* **38**, btac308- (2022).

984
985