<u>**Title:**</u> **Annexin- and calcium-regulated priming of legume root cells for**

endosymbiotic infection

Authors:

Ambre Guillory[1][§], Joëlle Fournier[1][§], Audrey Kelner[1][§], Karen Hobecker[2], Marie-Christine

Auriac[1], Lisa Frances[1], Anaïs Delers[1], Léa Pedinotti[4], Aurélie Le Ru[3], Jean Keller[4], Pierre-Marc

Delaux[4], Caroline Gutjahr[2], Nicolas Frei Dit Frey[4] and Fernanda de Carvalho-Niebel[1][*]

Affiliations :

[1]LIPME, INRAE, CNRS, Université de Toulouse, Castanet-Tolosan, France.

[2]Max-Planck-Institute of Molecular Plant Physiology, Potsdam-Golm, Germany.

[3]FRAIB-TRI imaging platform, Université de Toulouse, CNRS, UPS, Castanet-Tolosan, France.

[4]LRSV, Université de Toulouse, CNRS, UPS, Toulouse INP, Castanet-Tolosan, France.

[§]these authors contributed equally to this work

[*]Correspondence to: Fernanda de Carvalho-Niebel (fernanda.de-carvalho-niebel@inrae.fr)

Abstract

Legumes establish endosymbioses with arbuscular mycorrhizal (AM) fungi or rhizobia bacteria
to improve mineral nutrition. Symbionts are hosted in privileged habitats, root cortex (for AM
fungi) or nodules (for rhizobia) for efficient nutrient exchange. To reach these habitats, plants
form cytoplasmic bridges, which are key to predicting and guiding the cellular route of entry
of fungal hyphae or rhizobia-filled infection threads (ITs). However, the underlying
mechanisms are poorly studied. Here we show that unique ultrastructural changes and Ca^{2+}
spiking signatures, closely linked to MtAnn1 annexin accumulation, accompany rhizobia-
associated bridge formation. Loss of *MtAnn1* function in *M. truncatula* affects Ca^{2+} spike
amplitude, cytoplasmic configuration and rhizobia infection efficiency, consistent with a role
of MtAnn1 in regulating infection priming. *MtAnn1,* which evolved in species establishing
intracellular symbioses, is also AM-symbiosis-induced and required for proper arbuscle
formation. Together, we propose that MtAnn1 is part of an ancient Ca^{2+}-regulatory module
for transcellular endosymbiotic infection.

Introduction

Plants benefit from associations with microbes in the soil which improve the acquisition of
essential nutrients for growth. These associations include the ancestral symbiosis with
arbuscular mycorrhizal (AM) fungi providing phosphorous, or the more recently evolved
interactions with endosymbiotic bacteria for nitrogen acquisition[1-3]. Microbial endosymbionts
are hosted in privileged intracellular niches, where optimal conditions are met for efficient
nutrient exchange (e.g. the root cortex hosting arbuscules formed by AM fungi, or root
nodules, hosting nitrogen-fixing bacteria). Extensive studies on model legumes such as
Medicago truncatula, combined with evolution-based studies across a broad species range
revealed that nodule-forming nitrogen-fixing endosymbiosis (RNS) evolved ~100 million years
ago in species from related angiosperm plant lineages, by co-opting signalling components
from the ancient AM symbiosis and recruiting new key genes through neo-functionalization
and/or rewiring of expression[2,4,5]. These endosymbioses share a common symbiotic pathway
that uses calcium (Ca^{2+}) as a key secondary messenger to trigger downstream signalling[3,6].
Upon perception of mycorrhizal Myc and rhizobial Nod factor signals, Ca^{2+} oscillations (spiking)
are triggered in and around the root hair nucleus through the concerted action of nuclear
envelope channel/pump complexes[3,6,7]. While Ca^{2+} spiking is critical in the nucleus (after its
decoding by Ca^{2+}/calmodulin-dependent protein kinases CCaMK or Does not Make Infection,
DMI3)[8,9] to regulate host transcriptional reprogramming[10-12], the functional relevance of
sustained Ca^{2+} spiking in the cytoplasmic compartment has not yet been established. The
rhizobia- and AM-induced *MtAnn1*[13] and *MtAnn2*[14] genes, which encode cytosolic Ca^{2+}-
binding annexins that can bind to membranes to modulate a variety of Ca^{2+} -regulated
processes (e.g. ion conductance, exocytosis/endocytosis)[15,16], are potential players in this
process.

Following signal exchange to reach compatibility, infection of root tissues by AM fungi or
rhizobia bacteria occurs in most legumes transcellularly, through the *de novo* construction of
apoplastic compartments, delimited by host cell wall/ membrane interfaces, which physically
separate microsymbionts from the host cytoplasm[17-19]. The hyphae of AM fungi progress
through epidermal and cortical root tissues until the fungus reaches the inner cortex to form

highly branched arbuscules inside plant cells. Arbuscules comprise a massive network of membrane tubules at the symbiotic host-fungus interface[20,21], which likely ensures the release of nutrients, particularly phosphate, to the plant. In RNS, rhizobia enter the roots of most legume species *via* apoplastic tubular structures called infection threads (ITs)[18,19] that grow from root hairs to the developing nodule underneath[22] under the influence of key plant hormones, intercellular signalling and long-distance regulatory pathways[23-25].

ITs are formed in root hairs after rhizobia are entrapped and enclosed in a globular infection chamber[26]. This plant-driven process requires targeted cell wall remodelling at the IT interface[26,27] for polar growth, under the control of the infectosome protein complex[28-30]. Transcellular IT infection also imposes tight regulation of the plant's cell cycle status through the recruitment of mitotic regulators in root hairs and acquisition of specific pre-mitotic states before and during IT passage in the cortex[31-34]. Finally, the route of IT entry is also fine-tuned by the host, which creates a broad cytoplasmic column filled with endoplasmic reticulum (ER) and surrounded by microtubule arrays to guide IT progression[35,36]. In root hairs, this bridge connects the tip of the IT to the nucleus to direct IT elongation. In the adjacent cortex, differentiated root cells with a large central vacuole enter an activated state (hereafter termed pre-infection priming), with characteristic transvacuolar cytoplasmic strands emanating from the nucleus before a transvacuolar cytoplasmic bridge, the pre-infection thread (PIT[37]), forms in anticipation of future IT passage. PIT bridges have been reported to occur in nodules of several legume and actinorhizal species[37-39]. Moreover, PIT bridges closely resemble Pre-Penetration Apparatus (PPA)[17,40] cytoplasmic columns that precede AM fungal hyphae transcellular passage. The fact that PIT/PPA-like remodelling is not seen in pathogen interactions[41], suggests their specific and probably ancient recruitment for endosymbiosis. Parallels have been drawn between PIT/PPA and premitotic phragmosome transvacuolar bridges formed in cells preparing for mitosis, as these processes share similarities in tightly controlling plant cell division status, membrane trafficking and microtubule dynamics[33,34,38,42].

The development of fluorescent protein fusions labelling the endoplasmic reticulum network (ER) has made it possible to visualize the dynamics of ER-enriched bridge remodelling *in vivo* during PPA/PIT formation[35,40,43]. Complementary *in vivo* studies using Cameleon Ca^{2+} sensors revealed that primed cortical cells exhibit a low-frequency Ca^{2+} spiking signature that switches to high frequency at the onset of infection by AM fungi or rhizobia[43]. CCaMK/DMI3 is critical for PPA formation in *Medicago*[41]. Moreover, a gain-of-function variant in *Lotus* can trigger cytoplasmic remodelling in the cortex in the absence of symbionts[44]. Thus, CCaMK/DMI3-mediated signalling can bypass early symbiotic activation to trigger the priming response in the cortex. These studies suggest that cytoplasmic bridge reprogramming involves fine-tuning of Ca^{2+} spiking signatures, but the underlying genetic pathways or molecular players have not been identified.

In this study, using resolutive microscopy methods in different *M. truncatula* genetic backgrounds, we provide novel insights on ultrastructural rearrangements and *in vivo* Ca^{2+} signalling dynamics in plant cells preparing for rhizobia infection. We show that an intimate link between specific Ca^{2+} spiking signatures and the accumulation of annexin MtAnn1 characterises this priming response. MtAnn1 is a clear indicator of cytoplasmic reprograming for infection, and its expression in this context is induced by the master symbiotic transcription factor Nodule Inception (NIN)[4]. Through phylogenetics and mutant phenotyping, we provide evidence that *MtAnn1*, recruited during evolution in plants establishing root endosymbioses, likely plays an ancestral role for successful rhizobia and AM fungi infection. We propose that

MtAnn1 was recruited to shape calcium-regulated transvacuolar cytoplasmic rearrangements
for efficient endosymbiotic infection.

Results

Cell-specific ultrastructural changes in plant cells prior to IT entry

PIT bridge formation is a widespread process in RNS nodules[37,39], but detailed studies of
this process are lacking. Previous work in *M. sativa* and *M. tuncatula skl* mutant roots showed
that microtubule rearrangements accompany PIT formation[38], however overall cellular
features underlying this process remain unknown. Here, we set up an adapted methodology
to gain access to ultrastructural changes underlying rhizobia pre-infection priming in *M.*
truncatula roots. Segments of early infected roots, harvested 5-6 dpi with rhizobia, were
prepared for light and transmission electron microscopy (TEM) using an Epon-embedding
procedure. These analyses revealed specific cytoplasmic rearrangements in primed outer
cortical cells (C1) next to an infected root hair site (Fig. 1A-C) compared to mock control (Fig.
1H-J) root samples. TEM analyses showed that the thick cytoplasmic bridge was highly
enriched in endoplasmic reticulum (ER) (white arrows, Fig. 1E) and large quantities of small
vacuole-like structures with sizes ranging from 0.5 to 5 μm (v in Fig. 1E). These activated cells
also featured significant and unusually high mitochondrial clustering around the nucleus (Fig.
1E and Fig. S1). This mitochondria-nuclei association was consistently observed in independent
pre-infection (PI) primed cells (Fig. 1E and Fig. S1B-G) compared to actively dividing cells in the
nodule primordia (NP) (Fig. 1D, Fig. S1B-C and S1H-K). At a later stage, when the IT had
progressed across this small vacuole- and ER-rich cytoplasmic bridge, the nucleus was often
found in close proximity to the IT (Fig. 1F-G). Thus, the plant nucleus not only guides the
progression of ITs but also establishes close physical interactions with them. Overall, drastic
ultrastructural changes take place specifically in outer cortical cells (C1 or C2) primed for
rhizobia infection, namely a unique transcellular cytoplasmic organization, enriched with ER,
small vacuole-like structures and a significant increase in mitochondria-nucleus association.

MtAnn1 and Ca^{2+} spiking dynamics mark symbiotically engaged and infected plant cells

Ca^{2+} spiking responses are transiently regulated in cortical cells preparing for infection[43], but
how these responses relate to cytoplasmic remodelling is unknown. To study this, we used an
in vivo approach based on co-imaging a fluorescent Ca^{2+}sensor and a cytoplasmic marker to
simultaneously monitor Ca^{2+} spiking and cytoplasmic remodelling in individual cells. We used
M. truncatula A17 (wild-type) and the supernodulating mutant *sunn*, which shows wild-type
infection but at higher levels, facilitating live imaging studies of early rhizobia infection
events[26,28,45]. These *in vivo* analyses cannot provide as much data from the wild-type
background in which infection thread events are more rare than in *sunn*. However, results
were confirmed in the two genetic backgrounds (*A17, sunn*) by independent experiments.
Furthermore, as sites of interest were individually imaged multiple times in living roots,
responses were dynamically monitored and confirmed overtime by follow-up observations.

We used the red fluorescent nuclear NR-GECO1 sensor for Ca^{2+}imaging because of its
better compatibility with the green fluorescence cytoplasmic marker and higher sensitivity to
detect symbiotic Ca^{2+} spiking, as compared to a Förster resonance energy transfer (FRET)
cameleon sensor[46]. To validate this sensor in our system, we first monitored Ca^{2+} spiking in *S.*

meliloti-responsive root hairs of the nodulation susceptible zone, just below the infection
zone, in both *A17* and *sunn,* 1-4 dpi with rhizobia. Using NR-GECO1, we confirmed that *S.*
meliloti triggers periodic, high-frequency Ca^{2+} spiking in these root hairs with profiles
indistinguishable from those induced by purified Nod factors, as previously reported[46,47] (Fig.
S2 and Movies 1-2). Subsequent analysis of rhizobia-infected sites in *sunn* at 2-7 dpi, revealed
specific low frequency Ca^{2+} spiking profiles in the outer cortex underlying rhizobia infection
sites (Fig. S3). Cortical cells that spiked (Fig. S3B, D, F and H), included those that later became
infected (Fig. S3C and G). Furthermore, only cortical cells in close contact with the infected
epidermal cell spiked, while other distant cortical cells did not. This suggests that potential
short distance signalling from the infected cell is required for pre-infection priming of cortical
cells. Our results are consistent with previous data using a FRET-cameleon Ca^{2+} probe[43], and
confirm that low-frequency Ca^{2+} spiking is associated with rhizobial pre-infection priming of
cortical cells.

MtAnn1, encoding a cytosolic Ca^{2+} and membrane-binding annexin protein, is induced by
rhizobia both in root hairs and in outer cortical cells preparing for infection[13,31], where Ca^{2+}
spiking responses are coincidently transiently regulated (Fig. S3)[43]. Thus, the use of a cytosolic
MtAnn1-GFP fusion expressed under its native rhizobia-induced promoter seemed a suitable
tool to monitor cytoplasmic remodelling during rhizobia infection. Comparative localisation
analyses were performed in *M. truncatula* A17 and/or *sunn* genotypes using a GFP-ER marker,
which labels the cytoplasmic ER endomembrane network, and the MtAnn1-GFP fusion. This
revealed similar labelling of root hair cytoplasmic bridges by both markers (Fig S4). Co-
localization studies confirmed that MtAnn1-GFP and a red mcherry-ER fusion co-labelled
cytoplasmic bridges in both root hair (Fig. S5) and cortical cells (Fig. S6). MtAnn1-GFP fusion,
expressed under its native promoter, specifically labels cytoplasmic bridges of pre-infection
primed cells (Fig. S6A-B) and no labelling was observed in non-activated cells in the vicinity of
the red mCherry-ER marker (driven by the p35S) labelling the cytoplasm. Combined, these
data support the conclusion that the MtAnn1-GFP fusion specifically marks cytoplasmic
bridges, where it partially co-localizes with the ER network.

Co-expression of red NR-GECO1 Ca^{2+} sensor and green MtAnn1-GFP fusion was then used
to live track the spatio-temporal dynamics of Ca^{2+} spiking and MtAnn1 accumulation
simultaneously in *M. truncatula* A17. The MtAnn1/NR-GECO1 double-labelling greatly
facilitated detection of root hairs in early stages of bacterial entrapment (RHE) and IT polar
growth (IT). RHE- or IT- root hairs consistently showed both MtAnn1-GFP fusion signal and
high frequency Ca^{2+} spiking profiles (Fig. 2A-B, F). However, the amplitude of Ca^{2+} spikes
dropped significantly from RHE to IT stage root hairs (Fig. 2C-E). Conversely, relative levels of
MtAnn1-GFP fluorescence tended to increase in IT versus RHE root hairs (Fig. 2A-B and Fig.
S7), consistent with root hair transcriptome data[31]. Overall, our data suggest that decreased
Ca^{2+} spiking amplitude and increased MtAnn1 levels are associated with polar IT growth.

MtAnn1-GFP also labelled cytoplasmic bridges of cortical cells primed for infection in both
wild-type A17 and *sunn* (Fig. 3A-C, Fig. S6 and S8). MtAnn1-GFP-labelled cells were those in
close contact with the infected root hair, which consistently showed low frequency Ca^{2+}
spiking. Furthermore, this low-frequency Ca^{2+} spiking profile switches to a high-frequency
pattern when the cortical cell is infected (Fig. S8), consistent with previously published data[43].
Strikingly, nearby cortical cells not labelled with MtAnn1 did not show any spike in both A17
and *sunn* (Fig. 3 and Fig. S8). Collectively, our data establish that IT development and pre-

infection priming involve stage- and cell-type-specific Ca^{2+} spiking signatures closely linked to
MtAnn1-GFP bridge accumulation.

Rhizobia root hair infection is required for pre-infection priming of adjacent cortical cells

MtAnn1-GFP labelling and Ca^{2+} spiking mark primed outer cortical cells (Fig 3 and Fig S8),
but it is unclear if IT formation is required for this process. Co-expression of NR-GECO1 and
MtAnn1-GFP in infection-defective *ern1* (*ERF Required for Nodulation 1*)[48-50], revealed that
Ca^{2+} spiking and cytoplasmic-bridge formation are still seen in root hairs with arrested
chambers (Fig. S9A-C). MtAnn1-GFP labelling and Ca^{2+} spiking were also seen in the outer
cortex adjacent to root hairs with arrested-infections in *ern1* (Fig. S9D-F), but the pattern
appeared to be deregulated (more cells labelled with MtAnn1-GFP or Ca^{2+} spiking sometimes
in *MtAnn1* unlabelled cells, never seen in the wild-type control A17 or *sunn*, Fig. 3 and Fig. S8).
This suggests that ERN1 is not required to promote pre-infection priming, but in its absence
the priming response is deregulated.

CCaMK/DMI3 is critical for PPA formation in *Medicago*[41], however, it is unknown if it is also
required for PIT formation. Since *dmi3*[8] is impaired in root hair infection, it is not possible to
access PIT formation in this genetic background. We thus took advantage of stable transgenic
lines complemented with DMI3 only in root hairs[51], to investigate if Ca^{2+} spiking and MtAnn1-
GFP could still label cortical cells in this line compared to wildtype A17 control (Fig. 3).
Complemented *dmi3* pEXT-DMI3 lines showed Ca^{2+} spiking, rescued IT development and
MtAnn1 labelling in root hairs (Fig. S9G and Movie 3). However, no Ca^{2+} spiking or MtAnn1-
GFP labelling was seen in the cortex, compared to the control A17 or *sunn* (Fig. 3 and Fig. S8).
No cytoplasmic remodelling in the cortex of *dmi3* pEXT-DMI3 could be detected by further
analysis of a constitutively expressed GFP (Fig. S9I-J). Thus, MtAnn1-labelled cytoplasmic
bridges were clearly visible only in infected root hairs of *dmi3* pEXT-DMI3 and not in adjacent
cortical cells (no MtAnn1 labelling or GFP-labelled cytoplasm remodelling, Fig S9). This
suggests that PIT infection priming uses a DMI3-dependent pathway that must be active in the
cortex.

Taken together, these analyses indicate that root hair infection chamber formation is
sufficient in *ern1* to trigger MtAnn1-GFP-labelled bridge in the root hair itself and adjacent
cortex. The absence of cortical priming in *dmi3* is consistent with the conclusion that, like in
PPA, DMI3 is required for PIT reprogramming.

NIN controls pre-infection priming and the associated expression of *MtAnn1*

The master symbiotic transcription factor Nodule INception (NIN) is genetically
downstream of DMI3 and its mutation abolishes rhizobia-induced *MtAnn1* expression[31,52]. To
test if NIN is required for *MtAnn1* expression, we compared the spatio-temporal expression
profile of a *pMtAnn1:GUS* fusion in *M. truncatula* A17 wild-type and *nin* roots. In control
conditions, the *pMtAnn1:GUS* fusion was active in root tips, lateral roots and some
endodermal cells (Fig. 4A). Upon *S. meliloti* inoculation, strong promoter activity was detected
at rhizobial infection sites, both in root hairs and outer cortex (Fig. 4B-C), and later in other
nodule primordium cortical cells[13] but not in *nin* (Fig. 4D). This confirms the NIN-dependent
control of *MtAnn1* expression. Conversely, epidermal expression of NIN (*pEXPA:NIN*[53])
resulted in *pMtAnn1:GUS* fusion activity in the absence of rhizobia in *nin* (Fig. 4E-F). This

indicates that NIN can bypass early symbiotic signalling to promote *pMtAnn1:GUS* activity.
Transactivation assays in *Nicotiana benthamiana* (Fig. 4G and Fig. S10) showed that NIN can
transcriptionally activate *pMtAnn1:GUS*. This activation was abolished when a NIN mutated
version was used (Fig. 4G and Fig. S10). These data show that the expression of *MtAnn1* in
rhizobia infection sites is under NIN dependence and suggests that NIN is a possible direct
MtAnn1 regulator.

nin fails to form infection chambers and ITs but can still trap rhizobia in a curled root hair[26].
The question remains if a cytoplasmic bridge can still form in *nin*. To assess this, we examined
cytoplasmic bridge formation in *nin* transformed with *pEXPA:NIN* or a *pEXPA:GUS* control (Fig.
4H-M). Since *MtAnn1* is transcriptionally dependent on NIN (Fig. 4A-G), we could not use the
pMtAnn1:MtAnn1-GFP fusion for these studies, but instead used a *pENOD11:GFP-ER* fusion
co-expressed with *pEXPA:NIN* or *pEXPA:GUS* (control) in *nin*. Green GFP-ER fluorescence
marked the perinuclear ER network and the tip of growing root hairs in control and NIN-
complemented samples (Fig. 4H-J). However, thick cytoplasmic bridges marked by GFP-ER
were only detected in NIN-complemented root hairs (Fig. 4I-J). No GFP-ER signal was observed
in the root cortex of either control or NIN-complemented roots (Fig. 4K-M). Moreover, the
constitutive Ds-Red marker was also unable to detect cytoplasmic remodelling in *nin* (Fig. 4K-
M), whereas this marker allowed visualization of remodelling in control *sunn* roots (as
illustrated in Fig. S11). This suggests that a functionally active NIN in the cortex is required to
trigger cytoplasmic remodelling.

Together, these data infer that NIN acts downstream of DMI3 to control *MtAnn1* gene
transcription (possibly directly) and pre-infection priming.

MtAnn1 mutation affects efficiency of rhizobial infection and nodule differentiation

Since *MtAnn1* expression is closely linked to early stages of rhizobial entry, we investigated
the impact of its mutation on rhizobial infection by analysing *M. truncatula Tnt1* R108
insertion mutant lines (Fig. S12). Promoter (*ann1-1*) or exonic mutant lines (*ann1-2* and *ann1-*
3) showed reduced (*ann1-1*) or nearly abolished (*ann1-2* and *ann1-3*) *MtAnn1* expression
compared with R108 and R108S, a wild-type sibling from the same mutant population (Fig.
S12A-B). Rhizobia infection was assessed in *ann1-2* and *ann1-3,* where *MtAnn1* expression was
most reduced, by monitoring bacterial β-galactosidase reporter activity in root and nodule
tissues. ITs formed in root hairs and progressed into the cortex in mutant roots, but with lower
efficiency (Fig. 5A-B). This did not affect nodule numbers, but their infection levels and relative
VPY expression (Fig. 5C-P and Fig. S12-S13). At 4 dpi, there was a consistent decrease in the
number of fully colonized NPs in mutants (40-50 %, compared to 70-90 % in wild-type lines)
(Fig. 5C). These NPs were often infected by multiple ITs compared with controls (Fig. 5D-J),
suggesting an inefficient process. A mixture of well and poorly colonized nodules was still
observed in mutant root systems at 10 dpi (Fig. 6M-P). Although not statistically significant,
there was a trend towards higher *MtAnn2* expression in rhizobia-inoculated roots of *ann1-2*
and *ann1-3* mutants than in wild-type lines, suggesting a possible compensatory effect (Fig.
S12).

Consistent with the R108 *ann1* mutant analyses, RNAi *MtAnn1* knockdown in *M. truncatula*
A17 resulted in reduced NP/nodule infection at 5 dpi (Fig. 5Q-T). Infected nodules in RNAi roots
show the same multiple ITs above as in annexin mutants (compare RNAi Fig. 5T, V with control
Fig. S, U, W). RNAi roots also contain abnormal NPs with poor or no colonization, with several

associated root hair ITs (Figure 5X-Z). These are not seen in control NPs (Fig. W, arrowheads).
Additional nodule number reduction was seen in RNAi roots, possibly due to the double
knockdown of *MtAnn1* and *MtAnn2* (Fig. 5Q and Fig. S12). Overall, impaired *MtAnn1* function
in mutant and RNAi contexts affected rhizobial root and nodule infection, in R108 and A17 *M.*
truncatula backgrounds.

While *ann1* mutants showed no major changes in root architecture or nodule number,
striking variations in shape (less elongated overall) and colour (paler pink) were seen at 21 dpi
(Fig. 6A-C and Fig. S13), reminiscent of potential differentiation and/or nitrogen fixation
defects[54]. To better understand why *ann1* nodules were smaller and less elongated, we
analysed thin and ultrathin sections of wild-type and mutant nodules. As the two mutant
alleles led to similar nodulation defects and the wild-type sibling (R108S) behaved reliably as
R108 (see Fig. 5 and S12-S13), we chose to perform these delicate thin/ultrathin sections on
one representative of each genotype (R108S and *ann1-3*). Meristematic (I), infection (II),
amyloplast-rich interzone (IZ) and nitrogen fixation (III) zones were clearly distinguished in
R108S nodules (Fig. 6D-H, N and Fig. S14). However, *ann1-3* nodules showed variable zonation
patterns, either wild-type-like or with significantly abnormally large IZ regions, sometimes
accompanied by senescing cells (arrows) (Fig. 6I-M, O-P and Fig. S14). Despite this IZ defect
(Fig. 6P), *ann1-3* nodules formed bacteroids. However, they did not appear to complete their
differentiation in Zone III-like bacteroids and instead remained interzone-type bacteroids[55],
imbedded in a white vesicle-enriched region (black arrows) (Fig. 6R-X). As a result, *ann1-3*
mutant nodules had reduced nitrogen-fixing capacity (Fig. 5F), which ultimately led to reduced
plant growth (Fig. S4D). Taken together, these results indicate that *Mtann1* mutant nodules
are affected in their differentiation and nitrogen fixing abilities.

MtAnn1-dependent modulation of Ca^{2+} spiking and ultrastructural organization of the
cytoplasmic bridge

As Ca^{2+} spiking and MtAnn1 dynamics were closely linked in the peri-nuclear regions of
primed cells (Figs. 2-3), we assessed if *MtAnn1* mutation impacts symbiotic Ca^{2+} spiking.
Although more challenging than previous live analysis in the hyper infectious *sunn* context (Fig.
2), we succeeded in monitoring Ca^{2+} spiking responses at infection sites in both wild-type
R108S and *ann1-3* transgenic roots expressing the NR-GECO1 Ca^{2+} sensor. Similarly, as seen in
sunn (Fig. 2), a drop in Ca^{2+} spiking amplitude was observed in R108S in IT-containing RHs
compared with RHE root hairs (Fig. 7A). Conversely, this drop was not clearly observed in *ann1-*
3 (Fig. 7B). Likewise, a global deregulated Ca^{2+} spiking amplitude pattern was observed in many
S. *meliloti*-responsive root hairs of *ann1-3* compared with wild-type R108S, while Ca^{2+} spiking
frequency was not significantly changed in any root hair categories (Fig. 7C and S15). Together,
these data indicate a global deregulation of Ca^{2+} spiking amplitude in *ann1-3*, suggestive of
the need of *MtAnn1* for modulating symbiotic Ca^{2+} spiking responses.

Ca^{2+} spiking and MtAnn1 dynamics accompany cytoplasmic bridge formation (Fig. 2-3), so
we wondered how the deregulated Ca^{2+} spiking response in *ann1-3* (Fig. 7) could impact
cytoplasmic bridge formation. *MtAnn1* is expressed during root infection (Fig. 4), but also in
the nodule pre-infection and infection zones[56]. Thus, the defective nodule differentiation
phenotype of *ann-1-3* could be related to a possible deregulated infection in the nodule. We
thus analysed the cytoplasmic composition of infected cells in R108S and *ann1-3* apical nodule
zones by TEM, which can reveal ultrastructural organelle features and distribution that

confocal imaging cannot. In wild-type R108S, ITs progressed through a small vacuole-like
enriched cytoplasmic bridge containing a dense ER network (Fig. 7D-E, H-I and H'-I'), as shown
similarly in A17 (Fig. 1). In *ann-1-3*, the small vacuole-like and ER-enriched cytoplasmic
environment around the IT was strongly attenuated (Fig. 7F-G, J-K, J'-K'). Thus, mutation in
MtAnn1 resulted in a deregulated Ca^{2+} spiking response and a change in the ER and vesicle
composition of the cytoplasmic bridge.

MtAnn1 is required for efficient mycorrhizal root colonization

Recently developed genomic and transcriptomic databases from around 300 plant species
establishing different types of endosymbioses enabled to address key endosymbiosis-related
ancestral gene functions[57]. We have thus exploited these extensive resources to trace the
evolutionary history of *MtAnn1*. Previous studies of the monophyletic plant annexin group,
mainly composed of angiosperm sequences, suggested their emergence through duplications
from a common ancestor prior to the monocot/eudicot split[58]. Consistent with these data, we
found that the *MtAnn1* annexin cluster likely arose from sequential pre-Angiosperms, Eudicots
and Papilionoideae-specific duplications (Fig S8). *MtAnn1* belongs to one of the Papilionoideae
duplicated clades, which mainly includes species that establish intracellular endosymbioses
(Fig. S16), except for *Caryophyallae* and *Cephalotus* (status uncertain). *Ann1* orthologs were
also detected in Papilionoideae species that lost RNS but maintained AMS. This indicates that
the function of *Ann1* is not restricted to RNS but could expand to AMS. Strikingly, *Ann1* from
different species show consistent transcriptional activation during both root nodule and AM
symbioses[5], suggesting an ancestral role in root endosymbiotic infection. To determine if
MtAnn1 indeed plays a role during AMS, we first investigated if *MtAnn1* promoter activity was
spatio-temporally regulated in *M. truncatula* roots colonized by the AM fungus
Rhizophagus irregularis. Strong activation of the *pMtAnn1:GUS* fusion was only observed in
root segments colonized by *R. irregularis* at 4 wpi, especially in cortical cells containing
arbuscules (Fig. 8A-D). *R. irregularis* colonization rates were then monitored in *ann1-3*
compared to R108S wild-type roots at 6 dpi.

While R108S roots had dense colonization by *R. irregularis*, with cortex cells fully filled with
arbuscules, mutant roots had a lower proportion of colonized root sectors and arbuscule
density (Fig. 8E-H and K-L). In addition, arbuscules formed in the mutant showed a slightly but
significant reduction in length and this was associated with a trend increase in the expression
of the arbuscule senescence-related marker *MYB1*[59] (Fig. 8F-J and M-N). These data highlight
the importance of *MtAnn1* for the efficient establishment of the ancient AM symbiosis.

Discussion

Rhizobia root infection occurs transcellularly in most legumes via apoplastic IT structures
that grow polarly to reach the developing nodule. Plant cells engaged in this process form
cytoplasmic bridges that precede and accompany the route for ITs to progress. This process,
analogous to PPA bridges formed during AM symbiosis, is likely governed by common
ancestral cellular mechanisms, though knowledge remains limited. Here we report cell-
specific Ca^{2+} responses and cytoplasmic ultrastructural rearrangements that prime *M.*

truncatula plant cells for rhizobia infection. We show that this priming state, under DMI3-NIN
genetic control, is marked by an intimate cellular link between Ca^{2+} spiking and annexin
MtAnn1 accumulation. Moreover, MtAnn1 is needed for modulation of Ca^{2+} spiking responses
and creation of an infection-permissive, dense, ER- and small vacuole-rich cytoplasmic
environment. Ultimately, *MtAnn1* is required for successful root nodule differentiation and
efficient AM root colonization, implying an ancient recruitment of *MtAnn1* in endosymbiosis.
Our work supports the need to fine-tune cell-specific Ca^{2+}-spiking responses by annexin Ca^{2+}
binding proteins for the transcellular passage of endosymbionts.

Organelle dynamics shaping rhizobia pre-infection priming

Ultrastructure dissection of plant cells preparing for transcellular rhizobia infection
revealed marked enrichment and distribution of organelles during this process (Fig. 1). This
priming state is seen only in a few outer cortical cells in close contact with the infected root
hair cell (Fig. 1, 3 and S8), highlighting the likely need for signals from the infected cell to
promote it. A marked clustering of mitochondria around the nucleus is observed in primed
cells (Fig. 1 and S1). This may reflect the energy-demanding nature of this process. However,
dividing nodule primordia cells with presumably high metabolic activity show no such
enrichment. Alternatively, perinuclear mitochondria may mark a specific regulated cell cycle
status, likely to occur in these cells[33,34,60] or contribute to modulating cytosolic Ca^{2+}
oscillations, as shown in other systems[61].

Consistent with previous findings[35], TEM analysis revealed that a dense ER-rich network
occupy the transcellular cytoplasmic bridge in primed or rhizobia infected *M. truncatula* cells
(Fig. 1 and 7). The bridge is also rich in small vacuole-like structures[62], but we cannot exclude
that multivesicular bodies (MVBs), seen in the PPA bridge, are part of this organelle
population. Future use of molecular markers[62] may help discriminate between them. Dynamic
changes in plant vacuole size have been reported to occur in different cell types, and in early
meristematic cells this is thought to increase cytosolic area for subsequent daughter cell
formation events[62,63]. An analogous need for increased cytosolic area is possibly required for
IT passage. The concomitant reduction of small vacuoles and ER (involved in vacuole
biogenesis[64]) in *MtAnn1* mutants (Fig. 7) raises the question of their functional
interconnection in this process.

Role of MtAnn1 in modulating cell- and stage-specific symbiotic Ca^{2+} responses

High frequency Ca^{2+} spiking, triggered by Nod factors in root hairs is key for early rhizobia
recognition [3,6,46]. Here, we showed that a second Ca^{2+} spiking phase occurs in root hairs
undergoing rhizobia infection and involves a decrease in Ca^{2+} peak amplitude as ITs extend
into root hairs (Fig. 2). This drop was also reported in root outer cortical cells crossed by an
IT[43], consistent with this switch being key for IT progression. Ca^{2+} amplitude regulation fine-
tunes different biological processes ranging from gene expression, stomatal closure,
gravitropic responses to synaptic plasticity in animals[65-67]. The deregulated Ca^{2+} amplitude
profile and defective infection caused by *MtAnn1* mutation (Fig. 5 and 7) points to a functional
link between MtAnn1 and regulation of Ca^{2+} spiking amplitude during rhizobia infection.

We determined that Ca^{2+} spiking and strong MtAnn1-GFP labelling are hallmarks of rhizobia
infection and pre-infection priming (Fig. 3 and S8) and rely on the cell-autonomous activity of
DMI3 and NIN (Fig. 4 and S9). Thus, spatially distinct Ca^{2+} spiking responses are first triggered
in non-infected root hairs independently of DMI3-NIN[68,69], before DMI3-NIN-dependent Ca^{2+}
spiking is induced in cells engaged for infection. Using the NR-GECO1 sensor, we observe a
unique low frequency Ca^{2+} spiking only in primed cells of the cortex in both A17 and *sunn*
backgrounds (Fig. 3 and S8), in agreement with previous findings using a FRET-cameleon
sensor[43]. Although root hairs show similar remodelling and MtAnn1 labelling, they never show
such low frequency Ca^{2+} spiking, a signature likely reflecting a commitment state to infection.

In *Medicago*, symbiotic Ca^{2+} spiking is generated in the nucleoplasm and perinuclear
cytoplasm by the concerted action of nuclear envelope ion channels (DMI1, CNGC15), a Ca^{2+}
pump (MCA8), and under negative feedback regulation by a nuclear calmodulin (CaM2)[6,70].
MtAnn1 is a cytosolic phospholipid- and Ca^{2+}-interacting protein of the annexin family[13],
viewed as dynamic regulators that associate with membranes when intracellular Ca^{2+} levels
change[15,16]. Annexin family members are involved in the regulation of ion transport,
modulation of cytoplasmic Ca^{2+} signals and ROS signalling in diverse abiotic or biotic
responses[71-75]. Although predominantly cytoplasmic, annexins can also topologically associate
with the ER[76,77] and modulate the activity of Ca^{2+}-release channels[78]. As MtAnn1 partially
localizes to the ER network in cytoplasmic bridges (Fig. S4-S6), it may exert such a regulatory
function. The radical change in the cytosolic distribution of the ER and small vacuoles by the
MtAnn1 mutation (Fig. 7) also establishes a functional link between MtAnn1 and the creation
of this specific cytoplasmic environment. Ca^{2+} orchestrates membrane trafficking and fusion,
while vacuoles, emitters and receivers of Ca^{2+} signals, can be directly affected by altered
cytosolic Ca^{2+} levels[79,80]. Annexins are Ca^{2+} sensing proteins involved in endomembrane
trafficking in plants[81], either in conjunction with the membrane fusion machinery or by
regulating exocytosis. Thus, MtAnn1 may directly or indirectly impact Ca^{2+}-regulated vesicle
trafficking and/or vacuole morphology. However, the biological relevance of such a dense
cytoplasmic strand for IT passage remains to be determined.

Evolutionary recruitment of *MtAnn1* for root endosymbioses

In *Mtann1* mutants, root infection by rhizobia is reduced (Fig. 5) and this culminates in
mature nodules with reduced size, altered shape, abnormal zonation and reduced nitrogen
fixation ability. These phenotypes are reminiscent of the reduced infection levels and nodule
size observed after *PvAnn1* RNAi knock-down in common bean[82]. Such phenotypes probably
stem from the early deregulated Ca^{2+} spiking, impaired cytoplasmic configuration and
consequently defective infection caused by the *MtAnn1* mutation. Furthermore, as *MtAnn1* is
expressed in nodule pre-infection and infection zones[56], the defective nodule differentiation
phenotype is likely a consequence of maintained deregulated infection in the nodule. Finally,
the close paralogue *MtAnn2*[14] may act redundantly with *MtAnn1* in early root infection (Fig.
5) where they are both co-expressed[31], but it is unlikely to compensate for *MtAnn1* loss of
function in nodules as it has a distinct spatial expression profile.

We showed that the subclade containing *MtAnn1* emerged mainly in plant species
establishing intracellular symbioses. Induced expression of *MtAnn1* in rhizobia and AM
symbioses and its requirement for proper formation of nitrogen-fixing nodules and cortical
arbuscules (Figs. 5 and 8) suggest ancestral and shared roles for *MtAnn1* in endosymbiotic

infection. We propose that *MtAnn1* fine-tunes oscillatory Ca^{2+} responses to create an optimal
dense cytoplasmic environment for rhizobia IT and AM fungi hyphae to progress. As *MtAnn1*
is transcriptionally regulated by NIN, even in the absence of symbiotic signalling (Fig. 4), we
hypothesize that *MtAnn1* is a direct target of NIN and that its evolutionary recruitment for
RNS involved the acquisition of putative NIN *cis*-regulatory motifs, which are indeed present
in the *MtAnn1* promoter. Since NIN is a major regulator in RNS[83], the AM-associated
expression of *MtAnn1* is possibly under the control of another, so far unidentified regulator.

In conclusion, we showed that cell specific Ca^{2+} spiking signals shape rhizobia pre-infection
priming to prepare the optimal dense cytoplasmic environment for IT passage in *Medicago*,
and MtAnn1 emerges as a modulator of this process. MtAnn1, necessary for efficient rhizobia
and AM colonization, may have been recruited during evolution to integrate an ancestral Ca^{2+}-
regulatory module for guiding endosymbiotic infection. MtAnn1, with a conserved K/H/RGD
protein-interacting motif[15], may work in partnership with other players, and discovering them
could help identify other critical components in this process.

Methods

Plant materials and microbial strains

M. truncatula Jemalong A17 and R108 wild-type ecotypes, a regenerated transgenic line (referred
to as A2) expressing the *p35S-GFP-HDEL* construct[35] as well as *sunn* (*sunn-2* allele)[84], *dmi3* (*dmi3-*
1 allele)[8], *nin* (*nin-1* allele)[69] and *ern1* (*bit1* allele) & *ern1 ern2* (*bit1 ern2-1* alleles)[48,50] mutants
were used in this work. The *dmi3* line carrying the *pEXT:DMI3* construct[51], hereafter called
dmi3 + pEXT:DMI3, was kindly provided by C. Remblière (LIPME, Toulouse). The three R108
Tnt1 mutants carrying insertions in *MtAnn1* were obtained from Oklahoma State University
and are designated *ann1-1* (NF0830), *ann1-2* (NF17737) and *ann1-3* (NF4963). These lines
were self-crossed and genotyped by PCR and sequencing at each generation using primers -
344 pMtAnn1-Fw, Mtr14183-Rev, -914 pMtAnn1-Fw, LTR6- and *LTR4-* (Supplementary Table
S1). Homozygous mutants and sibling wild-type lines (referred to as WT) were selected for
seed multiplication and phenotyping. R108S is a wild-type sibling derived from line NF4963.
Seeds were scarified and surface-sterilized prior to germination on inverted soft agar plates[45]
and used for *A. rhizogenes* transformation or nodulation experiments. *N. benthamiana* seeds
were germinated in potting soil in glass-covered trays before transfer to pots (7 cm x 7 cm x
6.5 cm) at 21°C in a growth chamber under a 16h photoperiod and 70 µE/m/s light intensity
for 3 weeks until bacterial infiltration. The strains *A. rhizogenes* ARquA1, *A. tumefaciens*
GV3103 & GV3101 and *S. meliloti* 2011 (*Sm*2011-lac), constitutively expressing a *hemA-lacZ*
fusion, have been previously described[45]. The *S. meliloti* 2011 strain *Sm*2011-cCFP,
constitutively expressing a cyan fluorescent protein was kindly provided by P. Smit,
Wageningen (The Netherlands). *R. irregularis* spores (DAOM197198) were purchased from
Agronutrition (Labège, France).

DNA constructs

pMtAnn1:MtAnn1-GFP[13], *pENOD11:GFP-ER*[35] and *p35S-GFP-ER*[85] in pBIN121and *p2x35S-mCherry-*
ER in pBIN20[86] were used to study live pre-infection priming responses in *M. truncatula*.
p2x35S:NR-GECO1 in pCAMBIA-CR1ΔDs-Red[46] was used to monitor Ca^{2+} spiking in *M.*
truncatula. *pMtAnn1:GUS* in pLP100[13], *pEXPA:NIN* in pK7WG2-R*, p35S:3xHA-NIN* and

p35S:3xHA-NIN in PAM-PAT35S-3xHA-GTW[53], were used in transcription activation studies in
M. truncatula or *N. benthamiana*. *p35S:MtAnn1-GFP* and *p35S:GFP* in pBIN121[13], were used
in mutant complementation studies. The *pMtAnn1:RNAi-MtAnn1* and *pMtAnn1:GUS* (control)
constructs were generated here by Golden-Gate cloning into the pCambia-CR1 vector[46].
Briefly, DNA fragments corresponding to 2,2 kb of the genomic *MtAnn1* sequence upstream of the
ATG, 942 bp of the *MtAnn1* coding sequence in sense and antisense orientations, and the GUS coding
sequence, were generated by PCR amplification using the primer pairs *pMtAnn1-GG-A-Fw/pMtAnn1-*
GG-B-Rev, MtAnn1-ATG-GG-B-Fw/MtAnn1-STOP-GG-C-Rev, MtAnn1 antisens-STOP-GG-X-Fw/MtAnn1
antisens-ATG-GG-D-Rev and *GUS-GG-B-Fw/GUS-GG-D-Rev*, respectively (Supplementary Table S1).
DNA fragments were then cloned into a pBluescript SK vector and validated by DNA
sequencing prior to Golden Gate assembly. Golden-Gate reactions with plasmids comprising
(i) the *MtAnn1* promoter, *MtAnn1* coding sequences (sense and antisense) and a 1,3 kb intron
spacer[23] or (ii) the *MtAnn1* promoter and the *GUS* coding sequence, were used to generate
MtAnn1 RNAi (*pMtAnn1:MtAnn1 sens-intron-MtAnn1 antisens*) or GUS control
(*pMtAnn1:GUS*) constructs in the binary pCambia-CR vector. All binary vectors used for *A.*
rhizogenes transformation of *M. truncatula* (pLP100, pCambia, pK7WG2-R, pBIN121) include
a kanamycin resistance cassette in the T-DNA region. pK7G2-R and pCambia-CR1 vectors also
include a Ds-Red cassette.

Generation of *M. truncatula* transgenic material

A. rhizogenes-mediated transformation of *M. truncatula* A17 (WT, *sunn-2*, *ern1*, *ern1 ern2* and
dmi3) or R108 (WT sibling and *ann1-3*) was performed as described[45]. Briefly, germinated
seedlings had their root tips removed with a scalpel and were placed on Fahraeus medium
plates (12 cm x 12 cm) supplemented with 0.5 mM NH4NO3 and 20 mg/L kanamycin, before
inoculation with a drop (~3 µL) of *A. rhizogenes* aqueous suspension (OD600nm 0.5). For co-
transformation experiments with strains *pMtAnn1:MtAnn1-GFP + p2x35S:NR-GECO1*,
p2x35S:NR-GECO1 + pENOD11:GFP-ER, *pEXPA:NIN + pENOD11:GFP-ER* or *pEXPA:NIN +*
pMtAnn1:GUS (in pLP100), equal volumes of bacterial suspensions were mixed prior to
seedling inoculation. The lower ¾ part of the plates were sealed with parafilm and placed
vertically in plastic boxes under controlled 16h light/ 8h dark photoperiod conditions, first at
20°C for 1 week, then at 25°C for two to three weeks. Composite *M. truncatula* plants with
kanamycin resistant roots were selected. In some experiments, kanamycin-resistant
composite roots were re-selected on the basis of constitutive fluorescence of the Ds-Red
marker (for pK7WG2-R *pEXPA:NIN* and pCAMBIA-CR1 *pMtAnn1:RNAi-MtAnn1* and
pMtAnn1:GUS constructs), fusion fluorescence in lateral roots (for *pMtAnn1:MtAnn1-GFP* and
pENOD11:GFP-ER constructs) or in the nucleus (for *p2x35S:NR-GECO1*). Selected composite
plants were then transferred to appropriate nitrogen- and antibiotic-free plates or pots for
subsequent inoculation with rhizobia.

Plant growth and microbial inoculation procedures

For *in vivo* imaging experiments, kanamycin-resistant and fluorescent-positive composite
plants of A17 (WT, *sunn-2*, *ern1*, *ern1 ern2* and *dmi3*) or R108 (R108S and *ann1-3*) carrying
pMtAnn1:MtAnn1-GFP + p2x35S:NR-GECO1, pEXPA:NIN + pENOD11:GFP-ER or *p2x35S:NR-*
GECO1 + pENOD11:GFP-ER constructs were transferred to nitrogen-free Fahraeus plates
supplemented with Amoxycillin sodium/Clavulanate potassium 5:1 (200 mg/L) for 3-7 d, then

transferred to nitrogen-free 0.5% [w/v] phytagel Fahraeus plates supplemented with 50nM 2-
amino ethoxyvinyl glycine (AVG) for 3 days, as described previously[24]. Whole root systems
were then inoculated with 0.5 to 1 mL of the *Sm*2011-cCFP suspension (OD$_{600nm}$ 0.001), applied
between the medium and the LUMOX film. The inoculated root systems were kept protected
from light, by wrapping ¾ of the plates with dark plastic bags in a culture room at 20°C or 25°C
with a 16-h photoperiod and a light intensity of 70 mE/s/m^2, until microscopy observations.

For high-resolution microscopy analyses of pre-infection priming in A17 and nodule
differentiation in WT/*ann-3*, germinated seedlings of *M. truncatula* A17 and R108 (R108S and
ann1-3) were grown for 3 days on nitrogen-free paper/Fahraeus Kalys HP696 agar plates[46]
before roots were either spot or flood-inoculated with water (control) or a *Sm*2011-LacZ
(OD$_{600nm}$ 0.01) suspension, as described in[29]. For expression analyses of *pMtAnn1:GUS* (in
pLP100) in A17 and *nin* (co-expressing or not *pEXPA:NIN*), kanamycin-resistant and/or Ds-Red
fluorescence-positive composite plants were transferred 3 weeks after *A. rhizogenes*
transformation to the same nitrogen-free paper/Fahraeus Kalys HP696 agar plates and grown
for 3 days prior to flood inoculation (for 1h) of the entire root system with *Sm*2011-LacZ
(OD$_{600nm}$ 0.01). Plants grown *in vitro* on plates were all cultivated at 25°C under a 16-h
photoperiod and a light intensity of 70 mE/s/m^2 with their root systems protected from light
by wrapping ¾ of the plates with dark plastic bags.

For phenotyping experiments, germinated seedlings of *M. truncatula* R108 lines (R108, R108S
and *ann1-1 to ann1-3* mutants), and selected composite plants (2-3 weeks after
transformation) of A17-transformed with MtAnn1-RNAi/control constructs or R108 (R108S or
ann1-3)-transformed with *p35S:GFP/p35S:MtAnn1-GFP* constructs, were transferred to 8 x 8
x 7cm pots (3 plants/pot for germinated seedlings and 2 plants/pot for composite plants) filled
with inert attapulgite substrate (Oil Dri US Special; http://www.oildri.com/), supplemented
with 10mL nitrogen-free Fahraeus medium. For longer growth periods (> 3 wpi), plants were
grown in 9 x 9 x 8 cm pots (3 plants/pot) supplemented with 14mL medium. Pots were placed
in small greenhouses at 25°C, with a 16 h photoperiod and a light intensity of 100 mE/s/m^2,
and inoculated with a suspension of *Sm*2011-LacZ (OD$_{600nm}$ 0.1), after 3 days of nitrogen
starvation.

To assess mycorrhizal root colonization of WT and *ann1-3* mutant lines, germinated seedlings
were transferred to 8 x 8 x 7cm pots (1 plant per pot), filled with a 1:1 mix of Zeolite substrate
fractions 1.0-2.5 mm and 0.5-1.0 mm (Symbiom LTD, Lanskroun, Czech Republic), and
inoculated with *R. irregularis* spores (strain DAOM197198, 150 spores per pot). Pots were
placed in 60 x 40 x 12 cm trays in a 16 h photoperiod chamber (light intensity: 300 µmol/s/m^2)
at a day-time temperature of 22 °C and a night-time temperature of 20 °C, and 70% humidity.
Plants were watered weekly with a modified low-phosphate and low-nitrogen Long Ashton
solution (7.5 µM Na$_2$HPO$_4$, 750 µM KNO$_3$, 400 µM Ca(NO$_3$)$_2$, 200 mg/L MES buffer, pH 6.5).

β-glucuronidase (GUS) and β-galactosidase enzymatic assays

Root segments from *M. truncatula* composite plants expressing *pMtAnn1:GUS* (in pLP100) or
pEXPA:NIN + pMtAnn1:GUS constructs were collected from control (water) or rhizobia-
inoculated roots and incubated in 0.5 % paraformaldehyde/0.1 M potassium phosphate buffer
pH 7.0, for 1 h, prior to histochemical (blue) staining for GUS activity for 2-5 hours at 37 °C
using 1 mM of the substrate X-Gluc (5-bromo-4-chloro-3-indoxyl-b-D-GlcA,
cyclohexylammonium salt, B7300; Biosynth, Staad, Switzerland) as described[50]. For

mycorrhizal roots expressing *pMtAnn1:GUS*, composite plants were harvested 4 weeks after inoculation with *R. irregularis*. The root system was vacuum-infiltrated five times for 5 min each with GUS staining solution (0.5 mg/ml X-Gluc, 100 mM phosphate buffer pH 7.0, 100 mM EDTA, 0.5 mM $K_4[Fe(CN)_6]$, 0.5 mM $K_3[Fe(CN)_6]$ and 0.1% Triton X-100), and then incubated in this solution for 1 h at 37 °C in the dark. Histochemical GUS staining of *N. benthamiana* leaf discs was carried out in the same X-Gluc substrate but supplemented with 0,1 % triton, first under vacuum for 20 min at room temperature before incubation at 37 °C for 3 h. Enzymatic GUS fluorimetric assays of *N. benthamiana* leaf discs were done according to [45]. Briefly, tissues were ground using a MM400 grinder (Retsch) and homogenized in GUS extraction buffer before 1 µg of total protein extracts were used for enzymatic reactions at 37°C using 1 mM of the 4-MUG substrate (4-Methylumbelliferyl-β-D-glucuronide hydrate, Biosynth M-5700). GUS activity was measured using a FLUOstar Omega 96 microplate reader (BMG LABTECH, France) through quantification of the fluorescence of 4-MU (4-Methylumbelliferone, Sigma) reaction product. To reveal the constitutive β-galactosidase activity of the *S. meliloti* strain *Sm2011-lacZ* in ITs, root samples (sometimes pre-stained for GUS activity) were rinsed and fixed for 1 h in 1.25 % glutaraldehyde/Z buffer (10 mM KCl, 1 mM $MgCl_2$, and 0.1 M phosphate buffer, pH 7.0) as described[45]. After rinsing in Z-buffer, root samples were incubated overnight in the dark at 28 °C in Z-buffer containing 2 mM Magenta-Gal (5-bromo-6-chloro-3-indoxyl-b-D-galactopyranoside; B7200; Biosynth) or X-Gal (5-bromo-4-chloro-3-indolyl-b-D-galactopyranoside, W5376C; Thermo Fisher Scientific, Guilford, CT). GUS and/or LacZ-stained tissues were cleared for 30 s with 12 % sodium hypochlorite solution before microscopy observations.

Tissue harvesting and microscopy methods

Roots of *M. truncatula* composite plants expressing different fluorescent fusions and grown under LUMOX film were observed using a Leica TCS SP8 AOBS laser scanning confocal microscope equipped with a 40x long distance water immersion objective (HCX APO L U-V-I 40x/0.80 WATER). Confocal images were recorded using Leica LAS-X software, before and after rhizobia inoculation (from 1 to 7 dpi). For GFP, CFP and DsRed or mCherry fluorescent proteins, 458 nm and 488 nm Argon laser lines and a 561 nm diode were used for excitation, respectively, with emission windows set at 465-495 nm, 500-525 nm and 570-600 nm (DsRed) or 600-630 nm (mCherry) (hybrid detector), respectively. To avoid potential interference, sequential mode was used to acquire CFP and red fluorescent protein fluorescence separately from GFP fluorescence and bright-field images. For confocal imaging of calcium responses, the 561 nm diode was used to excite the mApple red fluorescent protein from the NR-GECO1 sensor, and the emitted fluorescence was recovered using a hybrid detector in the 600 to 643 nm emission window. Time series (t-series) were acquired at 5 s intervals for 10 to 15 min, with pinhole diameter set to 3 Airy units, as described[46].

Roots or nodulated roots of *M. truncatula* A17 and R108 (R108S and *ann1-3*) or composite plants were selected based on their fluorescence and/or analyzed before or after GUS and/or β-galactosidase staining using stereomicroscopy (Leica S6E) or light microscope (AxioPlan II Imaging; Carl Zeiss, Oberkochen, Germany). For *pMtAnn1:GUS* (in pLP100) expression analysis, samples were harvested 0 and 4 dpi with rhizobia. For comparative phenotyping of wild-type and *MtAnn1* mutant or RNAi roots, β-galactosidase stained nodulated root samples were harvested 4-5 or 8-10 dpi with rhizobia. Roots expressing *pEXPA:NIN + pMtAnn1:GUS*

were dehydrated in ethanol series and embedded in glycol methacrylate (Technovit 7100; Haereus-Kulzer) according to the manufacturer's instructions. Sections of 10 μm were counterstained with aqueous basic fushin solution (0.007%) for observations. To quantify nodule size and rhizobia infection levels , β-galactosidase-stained nodulated root systems (10 dpi) from WT and *ann1-3* plants were scanned (Objectscan 1600, Microtek) and the acquired images (TIFF) were used for image quantification (see details in the following part of the methods).

For high-resolution microscopy analyses of pre-infection priming responses in A17, control roots and rhizobia-infected root regions or nodule primordia were carefully isolated from spot-inoculated or flood-inoculated root systems 5-6 dpi, after histochemical revelation of rhizobial β-galactosidase activity. Harvested root sections or nodule primordia were fixed in 2% glutaraldehyde, diluted in 0.2 M cacodylate buffer pH=7.3 for a few hours at room temperature or 1-3 days at 4°C, then rinsed with buffer, before progressive dehydration in an ethanol series and final inclusion in LR White resin following manufacturer's instructions (EMS). For comparative ultrastructural analysis of isolated R108S and *ann1-3* nodules, nodule samples were carefully isolated from rhizobia-inoculated root systems 14-16 dpi and fixed under vacuum in 2 successive batches of 2.5% glutaraldehyde diluted in 0.2 M cacodylate buffer pH=7.3, the first batch containing saponin or triton (0.1 % final) for 3-5 days at 4°C. After rinsing in buffer, post-fixation in 2% osmium tetroxyde (diluted in 0.2 M sodium cacodylate), rinsing again in buffer, progressive dehydration in an ethanol series and incubation in propylene oxide (2 x 1h), samples were finally embedded in Epon 812 resin (EMS) following the manufacturer's instructions. For all samples, semi-thin (1 μm) and ultra-thin (80 nm) sections were generated using an Ultracut E ultramicrotome (Reichert Jung). Semi-thin sections of root/nodule primordia (5-6 dpi) were stained with basic fuchsin (0.07 % in water) to help visualize ITs, while mature nodule sections (14-16 dpi) were stained in an aqueous solution with methylene blue (0.2 %), toluidine blue (1 %), borax (1 %) and basic fuchsin (0.07 %) before observation by bright field microscopy (Axioplan II Imaging; Carl Zeiss). Ultrathin sections (80 nm) were contrasted with Uranyless and lead citrate (Delta microscopy) and observed using a Hitachi 7700 electron microscope (Hitachi High-Tech).

For histochemical GUS staining of mycorrhizal roots, selected composite plants transformed with *pMtAnn1:GUS* were transferred to 7 x 7 x 8 cm pots (5 plants/pot) filled with washed quartz sand (grain size 0.7-1.2 mm) pre-watered with modified half-strength Hoagland solution containing 20 μM phosphate[79]. Each plant was inoculated with 500 spores of *R. irregularis* DAOM197198 (C-grade, Agronutrition, Toulouse, France). Plants were grown in a Polyklima cabinet at 22°C constant temperature, 60 % air humidity and 16-h-light/8-h-dark cycles, at 200 μE/s/m2 light intensity and a combination of warm and cold LED 'True DayLight' at a 40:25 ratio. Pots were watered twice per week with 30-40 ml of autoclaved de-ionized tap water and fertilized once per week with 30-40 ml of modified half-strength Hoagland solution containing 20 μM phosphate. To visualize the fungal structures, the GUS staining solution was exchanged for 10 % KOH and the roots incubated for 15 min at 95°C. After KOH removal, the roots were washed 3 times with de-ionized water. After addition of 0.1 M HCl, the roots were further incubated for 2 h in the dark at room temperature. HCl was removed and the roots were washed 3 times with de-ionized water and once with PBS. This was followed by an overnight incubation at room temperature in the dark with 200 ng/μL WGA Alexa Fluor 488 (Invitrogen W11261) in PBS. Roots were imaged with a LEICA DM6B epifluorescence microscope. To compare the extent of mycorrhizal colonization of roots

between WT and *ann1-3* lines, whole root systems were collected 6 weeks post-inoculation
with *R. irregularis* and cleared by boiling (95°C) in 10% KOH for 5 minutes, rinsed with water,
then stained by boiling in an acidic ink solution (5% acetic acid, 5% Sheaffer black ink #94321,
in water) to reveal fungal structures. Overall intraradical colonization rates and arbuscular
density were estimated in ink-stained roots under a stereomicroscope (Leica S6E) using the
gridline intersect method.

Microscopy image analyses

Analysis of confocal t-series was performed using the Fiji software[80]. Intensity data were
calculated from selected regions of interest (ROIs), corresponding to single nuclei, imaged in
independent roots, in 2-3 independent experiments. Maximal z-projections of stacks and
merged confocal images for illustration purpose were prepared using Leica confocal software
or Fiji. To quantify relative amplitudes of Ca^{2+} spiking, the mean fluorescence of each ROI was
measured as a function of time and used to determine the Signal-to-Noise Ratio (SNR). The
SNR measures the difference between the fluorescence at each time-point (F_t) and the
fluorescence baseline, calculated as the average of 4 fluorescence values taken 10 and 15 sec
before and after t (F_{t-15}; F_{t-10}; F_{t+10}; F_{t+15}). A threshold of 0.3 was applied to the SNR to define
individual peaks, and the amplitude of Ca^{2+} spiking was calculated for each nucleus as the mean
of peak SNR values. Spiking frequency, defined as the number of peaks observed in 10 min,
was also calculated. In each experiment where calcium spiking was monitored, nuclei
displaying no spiking were also included, resulting in the sample size (n) being higher in the
frequency graph than in the associated amplitude graph.

For quantification of MtAnn1-GFP fluorescence levels, root hairs were imaged at either the
RHE or IT stages with identical acquisition settings in 3 independent experiments. In each
acquired confocal z-stack, 4 to 6 successive confocal sections encompassing the cytoplasmic
aggregation around the site of rhizobia entrapment (RHE) or the cytoplasmic bridge
characteristic of the IT stage were selected. In ImageJ, the threshold function was used to
generate a mask based on a maximal z-projection of the selected sub-stack, from which the
contours of the cytoplasmic zone (ROI) were defined. The mean grey value was then measured
in the defined ROI on an average z-projection of the same sub-stack.

Scanned images of β-galactosidase-stained root systems from R108S and *ann1* mutants (10
dpi) were imported to Ilastik software[87] to enable machine-learning recognition of blue β-
galactosidase-stained nodules (pixel-based classification method). Objects identified as
nodules were exported in a new single segmentation .tiff file for analysis in ImageJ to measure
specific nodule features (size and infection level, based on the blue intensity of rhizobial β-
galactosidase activity).

Transient expression assays in *N. benthamiana* leaves

An *A. tumefaciens* GV3101 strain carrying the *pMtAnn1:GUS* fusion construct (in pLP100) was
used for infiltration studies in *N. benthamiana* alone or with *A. tumefaciens* GV3103 strains
carrying *p35S:NIN* or *p35S:NINΔ* constructs[52], designed to constitutively express NIN or NINΔ
under the 35S promoter in transactivation experiments. Briefly, *A. tumefaciens* strains grown
overnight at 28°C in LB medium with appropriate antibiotics were harvested and resuspended
in Agromix (10 mM $MgCl_2$, 10 mM MES/KOH pH 5.6 and Acetosyringone in DMSO Sigma-

Aldrich 150 µM) and kept in the dark for at least 2h, at room temperature. Equal volumes of *A. tumefaciens* cultures (OD_{600}=0.25) were used to co-infiltrate leaf discs of 3-week-old *N. benthamiana* plants. Infiltrated plants were kept at 21°C in a growth chamber (16-h photoperiod and a light intensity of 70 mE/s/m2) for 36 h prior to histochemical GUS assays or storage at -70°C prior to protein extraction for Western-blot or fluorimetric GUS assays.

Western blot analyses

N. benthamiana leaf discs, previously stored at -80°C, were ground using a MM400 (Retsch) crusher and resuspended in Laemmli 2X sample buffer for SDS-PAGE (Bio-rad). Samples were then placed at 95°C for 3 min and centrifuged at 13200 rpm for 1 min. For each sample, 10 µL of the supernatant was loaded in a polyacrylamide 4-15% Mini-PROTEAN precast gel (Bio-Rad) along with 5 µL of pre-stained protein ladder marker (ThermoScientific, Lithuania). The Mini-Protean tank Electrophoresis System (Biorad, USA) was used for gel migration in 1 X Tris/Glycine/SDS Buffer (Bio-Rad). Protein transfer to nitrocellulose membranes (Bio-Rad) was performed using a Trans-Blot Turbo semi-dry transfer system (Bio-Rad). Membranes were stained with Ponceau Red before incubation for 1 h in blocking 1 x TBS solution (Tris base, NaCl and H_2O), 0.1 % Tween, 5 % milk at room temperature, with mild agitation, before rinsing 3 x in TBS-Tween 0.1 % for 10 min, and incubated overnight in the same solution at 4°C with a 6000 X dilution of anti-HA-peroxidase antibodies (Sigma-Aldrich) in 0.5 % milk in TBS-Tween 0.1 % to reveal HA-tagged NIN proteins following chemiluminescence revelation using the Clarity Western ECL Substrate (Bio-Rad) and the ChemiDoc Touch imaging system (Bio-Rad).

Acetylene reduction assay

To assess nitrogenase activity, grouped nodules were isolated from root systems of plants grown in pots 3 weeks after inoculation with *Sm*2011-LacZ and tested for acetylene reduction. Nodulated root fragments were incubated at 25°C in sealed 60 mL vials containing 0.5 mL of nitrogen-free Fahraeus liquid medium in the presence of 10 % (v/v) acetylene for 3 h. 400 µL of gaz was then collected from each vial and ethylene production was quantified by gas chromatography (model no. GC 7280A; Agilent Technologies, Lexington, MA).

RNA extraction and quantitative RT-PCR analysis

Total RNA was extracted from *M. truncatula* control and rhizobia-inoculated roots (4-5 dpi) using the Macherey-Nagel total RNA isolation kit according to the manufacturer's instructions. DNA-free RNA samples were quantified, and RNA integrity was verified by Agilent RNA Nano Chip (Agilent Technologies). First-strand complementary DNA synthesis was performed using 1 µg of total RNA using an anchored oligo (dT) and Transcriptor Reverse Transcriptase (Roche) following the manufacturers' protocol. Quantitative RT-PCR was performed on 384-well plates, with Light Cycler 480 systems fromRoche or Biorad and using the SYBR Green I Master mixes (Roche or Takyon, Eurogentec), according to manufacturer's instructions. Reactions and cycling conditions were performed as described before[29] using Primer pairs listed in Supplementary Table S1. The specificity of the PCR amplification was verified by analysing the PCR dissociation curve and sequencing of the PCR product. Transcript levels were normalized to the endogenous Ubiquitin reference.

Phylogenetic analysis

Protein sequence of *MtAnn1* (MtrunA17_Chr8g0352611) was used as query to search against a database containing 227 plant genomes covering the main lineages of green plants and five SAR genomes as outgroups using the BLASTp v2.14.0+ with an e-value threshold of 1e-10[55]. Homologous proteins were then aligned using Muscle v5.1 with the "super5" option[82] and trimmed to remove positions with more than 60% of gaps using trimAl v1.4.rev22[83]. The trimmed alignment served as matrix for phylogenetic reconstruction using FastTree v2.1.11-OpenMP[84]. To gain insights about the evolution of Ann1 in relation with the nitrogen-fixing symbiosis, the flowering clade corresponding containing orthologs of MtAnn1 has been extracted and proteins re-aligned using Muscle with default parameters before trimming as described above. Few spurious sequences with obvious mispositioning and abnormal short sequences were removed. Then, maximum likelihood tree has been reconstructed using IQ-TREE v2.1.2[85] after testing the best evolution model using ModelFinder[86] as implemented in IQ-TREE2 and according to the Bayesian Information Criteria. Branch supports have been tested with 10,000 replicates of both SH-aLRT[87] and UltraFast Bootstraps with UFBoot2[88]. Finally, trees were visualized and annotated in the iTOL platform v6.8.1[89].

Graph generation and statistical analyses

Graph preparation and statistical analyses were performed using R (http://r-project.org), except for graphs in Fig. 5C and D, which were generated using Microsoft Excel and for which contingency analyses were performed using GraphPad Prism 10. For all other analyses, data were presented as boxplots, normal distribution of the data was evaluated using the Shapiro-Wilk test, and homogeneity of variance was assessed using Fisher or Bartlett tests. When applicable, a transformation was performed to normalize data distribution (Log10 or BoxCox). Parametric statistical tests (t-test, ANOVA) were used to analyze data with a normal distribution, while non-parametric equivalents (Mann-Whitney, Kruskal-Wallis) were used for data with a non-normal distribution. Number of individually analyzed samples (n), replicates and *p* significance levels are indicated in Figure legends. In detail, data was analyzed as follows: Fig. 2: values in Fig. 2E follow a normal distribution and display homogeneous variance, so a two-tailed Student t-test was performed (t=-3.0927, df=39, p=0.0037); values in Fig. 2F do not follow a normal distribution and were thus analyzed using a two-tailed Mann-Whitney test (W=315, p=0.1531). Fig. 4: Log10-transformed values in Fig. 4G follow a normal distribution and variance homogeneity and were thus analyzed using one-way ANOVA followed by Tukey honest significant difference (HSD) tests (F=149.5, df=2, p<2e-16). Fig. 5: Values in Fig. 5A and 5B do not follow a normal distribution, hence Kruskal-Wallis tests were carried out (K=6.499, p= 0.0388 for A, and K= 6.204, p=0.0449 for B). Fig. 5C and 5D were built from contingency tables reporting numbers of nodule primordia belonging to different categories: fully infected (+) or poorly infected or not infected (+/- or -) in C, or NPs with a sole IT extended to the cortex (1 IT) or several ITs (≥ 2 ITs) in D. Relative proportions of these categories were compared one-on-one between R108S and each other genotypes (R108, *ann1-2* and *ann1-3*), by running separate Fisher's exact tests. Results of said tests are (C) R108S vs R108 p=0. 0,2738, R108S vs

ann1-2 p< 0,0077, R108S vs *ann1-3* p= 0,0428, and (D) R108S vs R108 p=0.4432, R108S vs *ann1-2* p=0.0128, R108S vs *ann1-3* p=0.0007. Values in Fig. 5O follow a normal distribution and variance homogeneity and were thus analyzed using one-way ANOVA followed by Tukey honest significant difference (HSD) tests (F=180,81, df=3, p<0,076), and values in 5P do not follow a normal distribution and were thus analyzed by a Kruskal-Wallis test (K=55.44 and p<0.0001). In Fig. 5Q, values follow a normal distribution and variance homogeneity, hence a two-tailed Student t-test was carried out (t=2.4153, df=99, p=0.0176), while values in Fig. 5R do not follow a normal distribution, thus two-tailed Mann-Whitney tests were applied (W=1724, p=0.0023). Fig. 6: Values in Fig. 6P follow a normal distribution but not homogeneity of variances, thus a Welsh two-sample t-test was carried out (t=4.8433, df=8.2905, p=0.0012). Log10-transformed values in Fig. 6Q show a normal distribution and variance homogeneity and were thus analyzed using a two-tailed Student t-test (t=-3.066, df=27, p=0.0049). Fig. 7: Values in Fig. 7A-B, which show a normal distribution and variance homogeneity were analyzed using one-tailed Student t-tests, due to the low number of values (7A: t=2.1821, df=9, p=0.0285; 7B: t=0.9612, df=22, p=0.1735). In Fig. 7C, values do not follow a normal distribution, thus a two-tailed Mann-Whitney test was performed (W=9944, p=0.0018). Fig. 8: In Fig. 8, values in 8I and BoxCox-transformed values in 8J (λ=1.353535) show a normal distribution and homogeneity of variance and were thus analyzed using two-tailed Student t-test (8I: t=3.6816, df=72, p=0.00045; 8J: t=3.8831, df=72, p=0.0002). Distribution of values is not normal in Fig. 8K, so data was analyzed via a Mann Whitney test (W=9103, p=0.0046.) Log-transformed values of Fig. 8L follow a normal distribution so a two-tailed Student t-test was applied (t=2.758, df=17, p=0.0134). Supplementary Fig. S1: In Fig. S1B, values do not follow a normal distribution, thus a Mann Whitney test was carried out (W=35, p=0.005349), while in Fig. S1C values do show a normal distribution, so a two-tailed Student t-test was performed (t=3.9724, df=10, p=0.0026). In Supplementary Fig. S2E-H, values show a non-normal distribution and were thus analyzed using two-tailed Mann-Whitney tests (S2E: W=853, p=1.811e-07; S2F: W=205, p=2.893e-05, S2G: W=348, p=3.377e-09, S2H: W=237, p=1.926e-09). In Supplementary Fig. S7C, values follow a normal distribution, hence a two-tailed Student t-test was performed (t=1.9307, df=6, p=0.1017). In Supplementary Fig. S12B and S12C, Log-transformed data follow a normal distribution, so one-way ANOVAs followed by Tukey honest significant difference (HSD) tests were carried out (respectively F=321.1, df=9, p<2e-16 for B, and F=1.701, df=9, p=0.106 for C). Log-transformed data in Fig. S12D, and non-transformed data in Fig. S12E follow a normal distribution, hence two-tailed Student t-tests were employed (respectively t=4.4053, df=22, p=0.0002 for D, and t=2.0775, df=22, p=0.0496 for E). In Fig. S12F, values do not have a normal distribution, so a Kruskal-Wallis test was used (K=43.0318, p=4.5265e-10). In Fig. S12G, data follows a normal distribution, so a one-way ANOVA followed by Tukey honest significant difference (HSD) tests was carried out (F=4.593, df=2, p=0.0143). In Supplementary Fig. S13C and S13E, data follow a normal distribution (after a Log-transformation for S13C), hence one-way ANOVAs followed by Tukey honest significant difference (HSD) tests were used for comparisons (respectively F=2.223, df=4, p=0.0765 for S13C and F=4.398, df=4, p=0.0034 for S13E). Values in Fig. S13D do not follow a normal

distribution, so statistical analysis was done using a Kruskal-Wallis test (K=9.15, p=0.0575). In Supplementary Fig. S15C and S15E, values do not follow a normal distribution and were thus analyzed using two-tailed Mann-Whitney tests (Respectively W=18215, p=0.3726 for S15C, and W=86, p=0.57 for S15E), while values in S15D do follow a normal distribution and variance homogeneity, and were analyzed using a two-tailed Student t-test (t=-1.0271, df=10, p=0.3286).

Reporting summary. Further information on research design is available in the Nature Research Reporting Summary linked to this article.

Data availability

The authors declare that all data supporting the results of this study are available within the article and its Supplementary Information Files. Materials generated in this study are available from the corresponding authors upon request.

References

Lanfranco, L., Fiorilli, V. & Gutjahr, C. Partner communication and role of nutrients in the arbuscular mycorrhizal symbiosis. *New Phytol* **220**, 1031-1046 (2018). https://doi.org/10.1111/nph.15230

Huisman, R. & Geurts, R. A Roadmap toward Engineered Nitrogen-Fixing Nodule Symbiosis. *Plant Commun* **1**, 100019 (2020). https://doi.org/10.1016/j.xplc.2019.100019

Genre, A. & Russo, G. Does a Common Pathway Transduce Symbiotic Signals in Plant-Microbe Interactions? *Front Plant Sci* **7**, 96 (2016). https://doi.org/10.3389/fpls.2016.00096

Griesmann, M. *et al.* Phylogenomics reveals multiple losses of nitrogen-fixing root nodule symbiosis. *Science* **361** (2018). https://doi.org/10.1126/science.aat1743

Libourel, C. *et al.* Comparative phylotranscriptomics reveals ancestral and derived root nodule symbiosis programmes. *Nat Plants* **9**, 1067-1080 (2023). https://doi.org/10.1038/s41477-023-01441-w

Charpentier, M. Calcium Signals in the Plant Nucleus: Origin and Function. *J Exp Bot* (2018). https://doi.org/10.1093/jxb/ery160

Krönauer, C. & Radutoiu, S. Understanding Nod factor signalling paves the way for targeted engineering in legumes and non-legumes. *Curr Opin Plant Biol* **62**, 102026 (2021). https://doi.org/10.1016/j.pbi.2021.102026

Lévy, J. *et al.* A putative Ca^{2+} and calmodulin-dependent protein kinase required for bacterial and fungal symbioses. *Science* **303**, 1361-1364 (2004). https://doi.org/10.1126/science.1093038

Tirichine, L. *et al.* Deregulation of a Ca^{2+}/calmodulin-dependent kinase leads to spontaneous nodule development. *Nature* **441**, 1153-1156 (2006). https://doi.org/10.1038/nature04862

Singh, S., Katzer, K., Lambert, J., Cerri, M. & Parniske, M. CYCLOPS, a DNA-binding transcriptional activator, orchestrates symbiotic root nodule development. *Cell Host Microbe* **15**, 139-152 (2014). https://doi.org/10.1016/j.chom.2014.01.011

Cerri, M. R. *et al.* The ERN 1 transcription factor gene is a target of the CC a MK/CYCLOPS complex and controls rhizobial infection in Lotus japonicus. *New Phytologist* **215**, 323-337 (2017).

Pimprikar, P. & Gutjahr, C. Transcriptional Regulation of Arbuscular Mycorrhiza Development. *Plant and Cell Physiology* **59**, 678-695 (2018). https://doi.org/10.1093/pcp/pcy024

13 De Carvalho-Niebel, F., Timmers, A. C., Chabaud, M., Defaux-Petras, A. & Barker, D. G. The Nod
factor-elicited annexin MtAnn1 is preferentially localised at the nuclear periphery in
symbiotically activated root tissues of Medicago truncatula. *Plant J* **32**, 343-352 (2002).

14 Manthey, K. *et al.* Transcriptome profiling in root nodules and arbuscular mycorrhiza identifies
a collection of novel genes induced during Medicago truncatula root endosymbioses. *Mol*
Plant Microbe Interact **17**, 1063-1077 (2004). https://doi.org/10.1094/MPMI.2004.17.10.1063

15 Davies, J. M. Annexin-Mediated Calcium Signalling in Plants. *Plants (Basel)* **3**, 128-140 (2014).
https://doi.org/10.3390/plants3010128

16 Gerke, V. *et al.* Annexins-a family of proteins with distinctive tastes for cell signaling and
membrane dynamics. *Nat Commun* **15**, 1574 (2024). https://doi.org/10.1038/s41467-024-
45954-0

17 Gutjahr, C. & Parniske, M. Cell and developmental biology of arbuscular mycorrhiza symbiosis.
Annu Rev Cell Dev Biol **29**, 593-617 (2013). https://doi.org/10.1146/annurev-cellbio-101512-
122413

18 de Carvalho-Niebel, F., Fournier, J., Becker, A. & Marín Arancibia, M. Cellular insights into
legume root infection by rhizobia. *Curr Opin Plant Biol* **81**, 102597 (2024).
https://doi.org/10.1016/j.pbi.2024.102597

19 Gao, J. P., Liang, W., Liu, C. W., Xie, F. & Murray, J. D. Unraveling the rhizobial infection thread.
J Exp Bot (2024). https://doi.org/10.1093/jxb/erae017

20 Roth, R. *et al.* Arbuscular cell invasion coincides with extracellular vesicles and membrane
tubules. *Nat Plants* **5**, 204-211 (2019). https://doi.org/10.1038/s41477-019-0365-4

21 Ivanov, S., Austin, J., Berg, R. H. & Harrison, M. J. Extensive membrane systems at the host-
arbuscular mycorrhizal fungus interface. *Nat Plants* **5**, 194-203 (2019).
https://doi.org/10.1038/s41477-019-0364-5

22 Xiao, T. T. *et al.* Fate map of Medicago truncatula root nodules. *Development* **141**, 3517-3528
(2014). https://doi.org/10.1242/dev.110775

23 Gaudioso-Pedraza, R. *et al.* Callose-Regulated Symplastic Communication Coordinates
Symbiotic Root Nodule Development. *Curr Biol* **28**, 3562-3577.e3566 (2018).
https://doi.org/10.1016/j.cub.2018.09.031

24 Lin, J., Frank, M. & Reid, D. No Home without Hormones: How Plant Hormones Control Legume
Nodule Organogenesis. *Plant Commun* **1**, 100104 (2020).
https://doi.org/10.1016/j.xplc.2020.100104

25 Chaulagain, D. & Frugoli, J. The Regulation of Nodule Number in Legumes Is a Balance of Three
Signal Transduction Pathways. *Int J Mol Sci* **22** (2021). https://doi.org/10.3390/ijms22031117

26 Fournier, J. *et al.* Remodeling of the infection chamber before infection thread formation
reveals a two-step mechanism for rhizobial entry into the host legume root hair. *Plant*
physiology **167**, 1233--1242 (2015). https://doi.org/10.1104/pp.114.253302

27 Su, C. *et al.* Transcellular progression of infection threads in Medicago truncatula roots is
associated with locally confined cell wall modifications. *Curr Biol* **33**, 533-542.e535 (2023).
https://doi.org/10.1016/j.cub.2022.12.051

28 Liu, C. W. *et al.* A protein complex required for polar growth of rhizobial infection threads. *Nat*
Commun **10**, 2848 (2019). https://doi.org/10.1038/s41467-019-10029-y

29 Lace, B. *et al.* RPG acts as a central determinant for infectosome formation and cellular
polarization during intracellular rhizobial infections. *Elife* **12** (2023).
https://doi.org/10.7554/eLife.80741

30 Li, X. *et al.* RPG interacts with E3-ligase CERBERUS to mediate rhizobial infection in Lotus
japonicus. *PLoS Genet* **19**, e1010621 (2023). https://doi.org/10.1371/journal.pgen.1010621

31 Breakspear, A. *et al.* The root hair "infectome" of Medicago truncatula uncovers changes in
cell cycle genes and reveals a requirement for Auxin signaling in rhizobial infection. *Plant Cell*
26, 4680-4701 (2014). https://doi.org/10.1105/tpc.114.133496

32   Gao, J. P. *et al.* Intracellular infection by symbiotic bacteria requires the mitotic kinase
AURORA1. *Proc Natl Acad Sci U S A* **119**, e2202606119 (2022).
https://doi.org/10.1073/pnas.2202606119

33   Yang, W. C. *et al.* Rhizobium nod factors reactivate the cell cycle during infection and nodule
primordium formation, but the cycle is only completed in primordium formation. *Plant Cell* **6**,
1415-1426 (1994). https://doi.org/10.1105/tpc.6.10.1415

34   Batzenschlager Morgane1, Lace Beatrice1, Zhang Ning1,2, Su Chao1, Egli Sabrina1, Krohn
Pascal1,3,4 & Salfeld Jule1, D. F. A., 4, Laux Thomas1,2, and Ott Thomas1,2,*. (bioRxiv, 2023).

35   Fournier, J. *et al.* Mechanism of infection thread elongation in root hairs of Medicago
truncatula and dynamic interplay with associated rhizobial colonization. *Plant Physiol* **148**,
1985-1995 (2008). https://doi.org/10.1104/pp.108.125674

36   Perrine-Walker, F. M., Lartaud, M., Kouchi, H. & Ridge, R. W. Microtubule array formation
during root hair infection thread initiation and elongation in the Mesorhizobium-Lotus
symbiosis. *Protoplasma* **251**, 1099-1111 (2014). https://doi.org/10.1007/s00709-014-0618-z

37   van Brussel, A. A. *et al.* Induction of pre-infection thread structures in the leguminous host
plant by mitogenic lipo-oligosaccharides of Rhizobium. *Science* **257**, 70-72 (1992).
https://doi.org/10.1126/science.257.5066.70

38   Timmers, A. C., Auriac, M. C. & Truchet, G. Refined analysis of early symbiotic steps of the
Rhizobium-Medicago interaction in relationship with microtubular cytoskeleton
rearrangements. *Development* **126**, 3617-3628 (1999).

39   Berg, R. W. Vol. 77 1351–1357 (Can. J. Bot., 1999).

40   Genre, A., Chabaud, M., Faccio, A., Barker, D. G. & Bonfante, P. Prepenetration apparatus
assembly precedes and predicts the colonization patterns of arbuscular mycorrhizal fungi
within the root cortex of both Medicago truncatula and Daucus carota. *Plant Cell* **20**, 1407-
1420 (2008). https://doi.org/10.1105/tpc.108.059014

41   Genre, A., Ortu, G., Bertoldo, C., Martino, E. & Bonfante, P. Biotic and abiotic stimulation of
root epidermal cells reveals common and specific responses to arbuscular mycorrhizal fungi.
Plant Physiol **149**, 1424-1434 (2009). https://doi.org/10.1104/pp.108.132225

42   Russo, G. & Genre, A. Divide and Be Conquered-Cell Cycle Reactivation in Arbuscular
Mycorrhizal Symbiosis. *Front Plant Sci* **12**, 753265 (2021).
https://doi.org/10.3389/fpls.2021.753265

43   Sieberer, B. J., Chabaud, M., Fournier, J., Timmers, A. C. & Barker, D. G. A switch in Ca2+ spiking
signature is concomitant with endosymbiotic microbe entry into cortical root cells of Medicago
truncatula. *Plant J* **69**, 822-830 (2012). https://doi.org/10.1111/j.1365-313X.2011.04834.x

44   Takeda, N., Maekawa, T. & Hayashi, M. Nuclear-localized and deregulated calcium- and
calmodulin-dependent protein kinase activates rhizobial and mycorrhizal responses in Lotus
japonicus. *Plant Cell* **24**, 810-822 (2012). https://doi.org/10.1105/tpc.111.091827

45   Cerri, M. R. *et al.* Medicago truncatula ERN transcription factors: regulatory interplay with
NSP1/NSP2 GRAS factors and expression dynamics throughout rhizobial infection. *Plant
Physiol* **160**, 2155-2172 (2012). https://doi.org/10.1104/pp.112.203190

46   Kelner, A., Leitão, N., Chabaud, M., Charpentier, M. & de Carvalho-Niebel, F. Dual Color Sensors
for Simultaneous Analysis of Calcium Signal Dynamics in the Nuclear and Cytoplasmic
Compartments of Plant Cells. *Front Plant Sci* **9**, 245 (2018).
https://doi.org/10.3389/fpls.2018.00245

47   Wais, R. J., Keating, D. H. & Long, S. R. Structure-function analysis of nod factor-induced root
hair calcium spiking in Rhizobium-legume symbiosis. *Plant Physiol* **129**, 211-224 (2002).
https://doi.org/10.1104/pp.010690

48   Middleton, P. H. *et al.* An ERF transcription factor in Medicago truncatula that is essential for
Nod factor signal transduction. *Plant Cell* **19**, 1221-1234 (2007).
https://doi.org/10.1105/tpc.106.048264

49 Andriankaja, A. *et al.* AP2-ERF transcription factors mediate Nod factor--dependent Mt
ENOD11 activation in root hairs via a novel cis-regulatory motif. *The Plant Cell* **19**, 2866--2885
(2007).

50 Cerri, M. R. *et al.* The Symbiosis-Related ERN Transcription Factors Act in Concert to Coordinate
Rhizobial Host Root Infection. *Plant Physiol* **171**, 1037-1054 (2016).
https://doi.org/10.1104/pp.16.00230

51 Rival, P. *et al.* Epidermal and cortical roles of NFP and DMI3 in coordinating early steps of
nodulation in Medicago truncatula. *Development* **139**, 3383-3391 (2012).
https://doi.org/10.1242/dev.081620

52 Schiessl, K. *et al.* NODULE INCEPTION Recruits the Lateral Root Developmental Program for
Symbiotic Nodule Organogenesis in Medicago truncatula. *Curr Biol* **29**, 3657-3668.e3655
(2019). https://doi.org/10.1016/j.cub.2019.09.005

53 Vernie, T. *et al.* The NIN Transcription Factor Coordinates Diverse Nodulation Programs in
Different Tissues of the Medicago truncatula Root. *Plant Cell* **27**, 3410-3424 (2015).
https://doi.org/10.1105/tpc.15.00461

54 Liu, H. *et al.* A genetic screen for plant mutants with altered nodulation phenotypes in
response to rhizobial glycan mutants. *New Phytol* **220**, 526-538 (2018).
https://doi.org/10.1111/nph.15293

55 Vasse, J., de Billy, F., Camut, S. & Truchet, G. Correlation between ultrastructural
differentiation of bacteroids and nitrogen fixation in alfalfa nodules. *J Bacteriol* **172**, 4295-4306
(1990). https://doi.org/10.1128/jb.172.8.4295-4306.1990

56 de Carvalho Niebel, F., Lescure, N., Cullimore, J. V. & Gamas, P. The Medicago truncatula
MtAnn1 gene encoding an annexin is induced by Nod factors and during the symbiotic
interaction with Rhizobium meliloti. *Molecular plant-microbe interactions* **11**, 504--513 (1998).

57 Radhakrishnan, G. V. *et al.* An ancestral signalling pathway is conserved in intracellular
symbioses-forming plant lineages. *Nat Plants* **6**, 280-289 (2020).
https://doi.org/10.1038/s41477-020-0613-7

58 Clark, G. B., Morgan, R. O., Fernandez, M. P. & Roux, S. J. Evolutionary adaptation of plant
annexins has diversified their molecular structures, interactions and functional roles. *New
Phytol* **196**, 695-712 (2012). https://doi.org/10.1111/j.1469-8137.2012.04308.x

59 Floss, D. S. *et al.* A Transcriptional Program for Arbuscule Degeneration during AM Symbiosis
Is Regulated by MYB1. *Curr Biol* **27**, 1206-1212 (2017).
https://doi.org/10.1016/j.cub.2017.03.003

60 Seguí-Simarro, J. M., Coronado, M. J. & Staehelin, L. A. The mitochondrial cycle of Arabidopsis
shoot apical meristem and leaf primordium meristematic cells is defined by a perinuclear
tentaculate/cage-like mitochondrion. *Plant Physiol* **148**, 1380-1393 (2008).
https://doi.org/10.1104/pp.108.126953

61 Costa, A., Navazio, L. & Szabo, I. The contribution of organelles to plant intracellular Calcium
signalling. *J Exp Bot* (2018). https://doi.org/10.1093/jxb/ery185

62 Cui, Y. *et al.* A whole-cell electron tomography model of vacuole biogenesis in Arabidopsis root
cells. *Nat Plants* **5**, 95-105 (2019). https://doi.org/10.1038/s41477-018-0328-1

63 Cui, Y., Zhao, Q., Hu, S. & Jiang, L. Vacuole Biogenesis in Plants: How Many Vacuoles, How Many
Models? *Trends Plant Sci* **25**, 538-548 (2020). https://doi.org/10.1016/j.tplants.2020.01.008

64 Viotti, C. ER and vacuoles: never been closer. *Front Plant Sci* **5**, 20 (2014).
https://doi.org/10.3389/fpls.2014.00020

65 Evans, R. C. & Blackwell, K. T. Calcium: amplitude, duration, or location? *Biol Bull* **228**, 75-83
(2015). https://doi.org/10.1086/BBLv228n1p75

66 Whalley, H. J. & Knight, M. R. Calcium signatures are decoded by plants to give specific gene
responses. *New Phytol* **197**, 690-693 (2013). https://doi.org/10.1111/nph.12087

67 Xu, T., Niu, J. & Jiang, Z. Sensing Mechanisms: Calcium Signaling Mediated Abiotic Stress in
Plants. *Front Plant Sci* **13**, 925863 (2022). https://doi.org/10.3389/fpls.2022.925863

1055	68	Wais, R. J. *et al.* Genetic analysis of calcium spiking responses in nodulation mutants of
1056		Medicago truncatula. *Proc Natl Acad Sci U S A* **97**, 13407-13412 (2000).
1057		https://doi.org/10.1073/pnas.230439797

69 Marsh, J. F. *et al.* Medicago truncatula NIN is essential for rhizobial-independent nodule
organogenesis induced by autoactive calcium/calmodulin-dependent protein kinase. *Plant*
Physiol **144**, 324-335 (2007). https://doi.org/10.1104/pp.106.093021

70 Del Cerro, P. *et al.* Engineered CaM2 modulates nuclear calcium oscillation and enhances
legume root nodule symbiosis. *Proc Natl Acad Sci U S A* **119**, e2200099119 (2022).
https://doi.org/10.1073/pnas.2200099119

71 Malabarba, J. *et al.* ANNEXIN1 mediates calcium-dependent systemic defense in Arabidopsis
plants upon herbivory and wounding. *New Phytol* **231**, 243-254 (2021).
https://doi.org/10.1111/nph.17277

72 Kodavali, P. K. *et al.* Structural and functional characterization of annexin 1 from Medicago
truncatula. *Plant Physiol Biochem* **73**, 56-62 (2013).
https://doi.org/10.1016/j.plaphy.2013.08.010

73 Liao, C., Zheng, Y. & Guo, Y. MYB30 transcription factor regulates oxidative and heat stress
responses through ANNEXIN-mediated cytosolic calcium signaling in Arabidopsis. *New Phytol*
216, 163-177 (2017). https://doi.org/10.1111/nph.14679

74 Richards, S. L. *et al.* Annexin 1 regulates the H2O2-induced calcium signature in Arabidopsis
thaliana roots. *Plant J* **77**, 136-145 (2014). https://doi.org/10.1111/tpj.12372

75 Laohavisit, A. *et al.* Arabidopsis annexin1 mediates the radical-activated plasma membrane
$Ca^{2}+$- and K+-permeable conductance in root cells. *Plant Cell* **24**, 1522-1533 (2012).
https://doi.org/10.1105/tpc.112.097881

76 Tichá, M. *et al.* Advanced Microscopy Reveals Complex Developmental and Subcellular
Localization Patterns of ANNEXIN 1 in. *Front Plant Sci* **11**, 1153 (2020).
https://doi.org/10.3389/fpls.2020.01153

77 Zhao, Z. X. *et al.* ANNEXIN 8 negatively regulates RPW8.1-mediated cell death and disease
resistance in Arabidopsis. *J Integr Plant Biol* **63**, 378-392 (2021).
https://doi.org/10.1111/jipb.13025

78 Vais, H. *et al.* ER-luminal [Ca. *Elife* **9** (2020). https://doi.org/10.7554/eLife.53531

79 Peiter, E. The plant vacuole: emitter and receiver of calcium signals. *Cell Calcium* **50**, 120-128
(2011). https://doi.org/10.1016/j.ceca.2011.02.002

80 Himschoot, E., Pleskot, R., Van Damme, D. & Vanneste, S. The ins and outs of Ca. *Curr Opin*
Plant Biol **40**, 131-137 (2017). https://doi.org/10.1016/j.pbi.2017.09.003

81 Konopka-Postupolska, D. & Clark, G. Annexins as Overlooked Regulators of Membrane
Trafficking in Plant Cells. *Int J Mol Sci* **18** (2017). https://doi.org/10.3390/ijms18040863

82 Carrasco-Castilla, J., Ortega-Ortega, Yolanda, Jáuregui-Zúñiga, David, Juárez-Verdayes, Marco
A, Arthikala, Manoj-Kumar, Monroy-Morales, Elisabeth, Nava, Noreide, Santana, Olivia,
Sánchez-López, Rosana, Quinto, Carmen. 108-119 (Environmental and Experimental Botany,
2018).

83 Liu, J. & Bisseling, T. Evolution of NIN and NIN-like Genes in Relation to Nodule Symbiosis.
Genes (Basel) **11** (2020). https://doi.org/10.3390/genes11070777

84 Schnabel, E., Journet, E. P., de Carvalho-Niebel, F., Duc, G. & Frugoli, J. The Medicago
truncatula SUNN gene encodes a CLV1-like leucine-rich repeat receptor kinase that regulates
nodule number and root length. *Plant Mol Biol* **58**, 809-822 (2005).
https://doi.org/10.1007/s11103-005-8102-y

85 Haseloff, J., Siemering, K. R., Prasher, D. C. & Hodge, S. Removal of a cryptic intron and
subcellular localization of green fluorescent protein are required to mark transgenic
Arabidopsis plants brightly. *Proc Natl Acad Sci U S A* **94**, 2122-2127 (1997).
https://doi.org/10.1073/pnas.94.6.2122

86   Nelson, B. K., Cai, X. & Nebenführ, A. A multicolored set of in vivo organelle markers for co-
localization studies in Arabidopsis and other plants. *Plant J* **51**, 1126-1136 (2007).
https://doi.org/10.1111/j.1365-313X.2007.03212.x
87   Berg, S. *et al.* ilastik: interactive machine learning for (bio)image analysis. *Nat Methods* **16**,
1226-1232 (2019). https://doi.org/10.1038/s41592-019-0582-9

1110

Acknowledgements

The *M. truncatula* R108 mutant plants utilized in this research project, which are jointly owned
by the Centre National De La Recherche Scientifique, were obtained from Noble Research
Institute, LLC and were created through research funded, in part, by grants from the National
Science Foundation, NSF #DBI-0703285 and NSF #IOS-1127155. We thank S. Bensmihen for
providing the *pEXT:DMI3* construct[51] and Celine Remblière for providing seeds of the *M.
truncatula dmi3* line carrying the *pEXT:DMI3* construct. This work was supported by French
National Research grants TULIP ANR-10-LABX-41, COME-IN ANR-14-CE35-0007-01 and LIVE-
SWITCH ANR-19-CE20-0026-01 to FdCN and by the European Research Council (ERC) under
the European Union's Horizon 2020 research and innovation program (grant No. 759731) to
CG. A. G. was funded by an ANR postdoctoral grant (ANR-19-CE20-0026-01), A. K. and A. D.
were funded by PhD grants from the French Ministry of National Education and Research. We
acknowledge the TRI-FRAIB imaging facility, member of the national infrastructure France-
BioImaging supported by the French National Research Agency (ANR-10-INBS-04).

1125

Author contributions

1127 Coordinated the work: FdCN; designed experiments: FdCN, AG, JF, AK, KH, MCA, LF, NFDF, CG;
1128 performed experiments: AG, JF, AK, KH, MCA, LF, AD, NFDF; Machine learning methods
1129 development: AG, ALR; phylogenetic analysis: JK, PMD; analysed data: AG, JF, AK, KH, MCA,
1130 LF, AD, LP, JK, PMD, CG, NFDF, FdCN; wrote the manuscript: FdCN, AG and JF.

1131

Fig. 1. Cell-specific structural changes in *M. truncatula* cells primed for infection.

(**A-B**) Representative images of longitudinal sections of *M. truncatula* A17 roots 5 days after spot-inoculation with *S. meliloti.* These are consecutive 1 μm sections stained with basic fuchsin to reveal cell outlines and content. Arrows indicate ITs in root hair epidermal cells and arrowheads indicate a C1 cortical cell exhibiting pre-infection priming. The region of interest in **B** (highlighted by a white square) is shown in **C-E** by transmission electron microscopy (TEM) of 80 nm sections. Note that the primed C1 outer cortical cell (arrowhead in **C**, and enlarged image in **E** exhibits a unique structural organization compared with neighbouring cells. Primed cells comprise a cytoplasmic bridge enriched in endoplasmic reticulum (white arrows in **E**), and numerous vesicles or small vacuole-like (SV-like) structures (SVs) (v). They also have a large nucleus (n) with enriched mitochondria (asterisks) around it, as quantified in Fig. S1. TEM

analysis of primed cells were performed in n=8 sections derived from 5 independent infection sites from 2 independent experiments. (**F**-**G**) At a later stage, when the IT (arrows) has progressed across the cytoplasmic bridge towards inner tissues, the nucleus of the crossed cell is frequently found closely associated to the IT (**G**). (**H**-**J**) Representative images of *M. truncatula* A17 root sections after mock (water) treatment for 5 days. (**H**) A 1 µm longitudinal section of a mock control root stained with basic fuchsin. (**I**-**J**) TEM of 80 nm sections of a mock control root. The region of interest in **I** (highlighted by a white square) is shown in **J**. Note that non-symbiotically activated cells have a large vacuole and nuclei near the cell periphery. Other abbreviations: cortical cells (C1, C2, C3), nucleolus (nuc). Scale bars: **A**-**B**, **H** = 50 µm, **C**, **I** = 20 µm, **D**-**G**, **J** = 5 µm. See additional primed cells and mitochondria quantification in Fig. S1.

Fig. 2. Ca^{2+} spiking amplitude drops along root hair IT development.

(**A**-**B**) Representative bright-field and corresponding confocal images of (**A**) a root hair with entrapped CFP expressing (magenta)-*S. meliloti* (RHE) and (**B**) a root hair with a growing infection thread (IT) in *sunn*. Nuclei expressing the NR-GECO1 Ca^{2+} sensor appear in red, and the MtAnn1-GFP fluorescence (green) labels the cytoplasmic zone around the IT and the cytoplasmic bridge connecting and surrounding the nucleus. (**C**-**D**) Representative Ca^{2+} spiking traces of RHE (**C**) and growing IT stages (**D**) in *sunn*. The relative concentration of Ca^{2+} ions in the nucleus is reflected by the intensity of NR-GECO1 fluorescence, expressed as signal-to-noise ratio (SNR, cf. Methods section). The number of root hair with nuclear spiking/total number of root hairs are indicated between parentheses. Nuclei are counted as spiking when showing more than 2 peaks in 10 min. (**E**-**F**) Quantification of nuclear Ca^{2+} spiking: Spiking amplitude (**E**), expressed as average SNR of spikes per nucleus, and spiking frequency (**F**),

expressed as number of spikes in 10 minutes per nucleus. Box plots represent the distribution of individual values (indicated by open circles) from root hairs with entrapped rhizobia RHE, n=23 in **E** and n=24 in **F**) or with an IT (n=18 in **E** and n=21 in **F**) in *sunn,* 2-4 dpi with *S. meliloti* from three independent biological experiments. Median (central line), mean (solid black circle) and outliers (cross) are indicated. Two-tailed T-test (**E**) and Mann-Whitney test (**F**) of values were performed in R (asterisks indicate statistical difference; **p < 0.01). Scale bars: **A-B** = 10 µm. See also Fig. S2 and Movies 1-2 for Ca^{2+} spiking responses in A17 and/or *sunn* root hairs, and Figs. S4-S5 and S7 for MtAnn1-GFP localization and fluorescence quantification in root hair bridges.

Fig. 3. Strong MtAnn1-GFP fusion fluorescence and low frequency Ca^{2+} spikes are hallmarks of pre-infection priming in the cortex.

Representative images of rhizobia infection sites in roots co-expressing MtAnn1-GFP (green) and NR-GECO1 (red) in *M. truncatula* A17. Confocal images from **A** to **C** illustrate an infected root hair with CFP-labelled rhizobia in IT (in magenta, arrow) and the nucleus (n) in front, guiding the growth of the IT in a cytoplasmic bridge (arrowhead) (**A**), the base of the root hair cell (**B**, dotted line) and the neighbour C1a-c outer cortical cells (**C**, dotted lines). C1a-C1c are also shown in **D**. (**D-E**) Representative traces of Ca^{2+} spiking from outer cortical cells from two

independent sites (C1a is also shown in **C-D**, and C1g in **E**, is from another site), adjacent to an infected root hair (shown in **A** for **D**, and not shown for the second site **E**). Traces from other neighbouring cortical cells (C1b-c, C2d-f in **D** and C1h-l in **E**) not in direct contact with the infection sites are shown. Cells are indicated in the upper panels. Images were obtained from A17 roots 4 dpi (**A-D**) or 3dpi (**E**) with *S. meliloti*. Note that only cortical cells co-expressing MtAnn1-GFP show detectable low frequency Ca^{2+} spiking (compare C1a-b and C1c, C1g and C1h-l, respectively). Traces are expressed as signal-to-noise ratio (SNR). Images in **A-C** are maximal z-projections of sub-stacks and images in **D-E** are maximal projections of whole time series. Scale bars: **A-E** = 20 µm. See also Figs. S3, S6 and S8 for Ca^{2+} spiking responses and MtAnn1-GFP dynamics in primed cells in the cortex in *sunn*. See also Fig. S3-S6 and Fig. S8-S9 and Movie 3.

Fig. 4 NIN is required for cytoplasmic pre-infection priming and *MtAnn1* expression.

(**A-F**) Expression of the *pMtAnn1:GUS* fusion in *M. truncatula* A17 (**A-C**) or *nin* (**D-F**) roots non-inoculated (**A, E-F**) or inoculated with *S. meliloti* at 4 dpi (**B-D**). (**A-B**) *MtAnn1* exhibit constitute

promoter activity during root development (in root tips and developing lateral roots, arrowheads) and endodermal cells (asterisks). (**B**-**C**) Upon rhizobia inoculation, GUS activity is strongly detected at rhizobia infection sites (ITs, arrows in **B**-**C**). (**C**) A close view of an infection site (black square in **B**) enables to visualize *LacZ*-expressing rhizobia in magenta (arrow). (**D**-**F**) The infection-related expression profile of *MtAnn1* is abolished in rhizobia-inoculated *nin* roots (**D**). (**E**-**F**) Expression of NIN under the epidermis-specific promoter *pEXPA* in *nin* roots promotes *pMtAnn1:GUS* fusion activity in the epidermis, even in the absence of rhizobia (**E**-**F**). A longitudinal 10 μm root section counterstained with basic fushin show *pMtAnn1:GUS* activity (blue) restricted to the epidermal cell layer (**F**). Data in **A**-**F** are from two independent experiments (n=41 in A17, n=31 in *nin*, n=37 in *nin + pEXPA:NIN*). (**G**) Transactivation of the *pMtAnn1:GUS* fusion by NIN in *A. tumefaciens*-infiltrated leaf discs. Fluorimetric GUS assays were performed using 1 μg of total protein extracts. Box plots represent the distribution of values (indicated as open circles) of individual plants (n=25 per sample) from 4 independent experiments. Median (central line), mean (solid black circle) and outliers (cross) are indicated. Different letters above the box blots indicate statistically significant difference (One-way ANOVA, α= 5%). Images of representative leaf discs are included. (**H**-**M**) *pENOD11:*GFP-ER expression in *nin* transformed with *pEXPA:GUS* control (**H**, **K**) or complemented with *pEXPA:NIN* (**I**-**J**, **L**-**M**). Control or complemented *nin* roots all co-express the fluorescent marker *pUbiquitin*-DsRed, which labels cytosol and nucleoplasm in red. Green GFP-ER fluorescence labels the perinuclear ER network of epidermal cells and the tip of growing root hairs in **H**-**J**. (**I**-**J**)In complemented *nin*, GFP-ER also labels cytoplasmic columns (arrowheads) in rhizobia-enclosed (arrow in **J**) RHE stage root hairs. (**K**-**M**) In both control (**K**) and complemented *nin* roots (in **M**, boxed area in **L**), outer cortical cells (co) show no GFP-ER labelling. These Ds-Red labelled cells also show no signs of cytoplasmic reorganisation, whereas using the same Ds-Red marker we can detect cytoplasmic rearrangements in *sunn* (See Fig S11). **J** and **M** are higher magnification of boxed areas of **I** and **L**, respectively. Data were obtained from 2 independent experiments (n=7 for *nin + pEXPA:GUS* and n=19 for *nin + pEXPA:NIN*). Asterisks in **J** and **M** mark the xy positions of the infected root hair base. Abbreviations: epi=epidermis; co=cortex. Scale bars: **A**-**F** = 100 μm, **H**-**M** = 40 μm. See also Fig. S10-S11.

Fig. 5. Root Infection is negatively impacted in *ann1* mutants and RNAi transgenic roots.

Relative rhizobia infection levels were monitored in *ann1* mutants and *MtAnn1* RNAi transgenic roots with reduced *MtAnn1* or *MtAnn1/MtAnn2* expression levels (shown in Fig.

S12). Histochemical X-gal staining of mutant or RNAi *Medicago* roots (and corresponding controls) infected with a *lacZ*-expressing *S. meliloti* strain was used to quantify (**A**-**D**, **O**-**R**) and visualize (**E**-**N** and **S**-**Z**) rhizobia infection events in blue. (**A**-**J**) Early root infection by rhizobia was analyzed in root systems of R108 (n=17), R108S (n=21), *ann1-2* (n=36) and *ann1-3* (n=38) plants from 2 independent experiments at 4 dpi with rhizobia. The number of ITs in-root hairs (**A**) and ITs that had reached the outer cortex (**B**) are indicated. Different letters above the box blots indicate statistically significant difference (Kruskal-Wallis tests, α= 5%). (**C**-**D**) Graphs illustrate the proportion of plants having low to none (- or +/-) or colonized (+) nodule primordia (NP) (**C**), and of nodule primordia colonized by 1 or several ITs in *ann1* mutants compared to R108 and/or R108S lines (**D**). Asterisks in **C**-**D** indicate statistical differences relative to the R108S control (Fisher's exact tests, * p < 0.05, **p < 0.01, ***p < 0.001). (**E**-**J**) Representative images of R108 (**E** and **F**), R108S (**G**), *ann1-2* (**H**) and *ann1-3* (**I** and **J**) infected nodule primordia at 4 dpi. (**K**-**N**) Representative images of R108, R108S and *ann1* nodules at 10 dpi with rhizobia. Note the higher proportions of less infected (arrows) nodules in mutants compared to WT lines. (**O**-**P**) R108, R108S, *ann1-2* and/or *ann1-3* nodule number (**O**) or infection levels (**P**) were analyzed 10 dpi with rhizobia. Infection levels (**P**) were quantified by automated ImageJ measurement of the intensity of bacterial X-gal blue staining of 10 dpi nodules on R108S (196 nodules from 21 plants), *ann1-2* (295 nodules from 30 plants) and *ann1-3* (310 nodules from 21 plants) roots, from 2 independent experiments. Different letters above the box blots indicate statistically significant differences (One-way ANOVA in 0 and Kruskal-Wallis in P tests, α = 5%). (**Q**-**Z**) RNAi knockdown of *MtAnn1* and *MtAnn2 in M. truncatula A17* roots. Total nodule/NP number (**Q**) and infected nodule/NP number (**R**) were quantified in in *M. truncatula* A17 roots expressing *pMtAnn1:GUS* (control) or *pMtAnn1:RNAi-MtAnn1* (*RNAi-MtAnn1*) at 5 dpi with *S. meliloti-LacZ* from 3 independent experiments (control, n=51 and *RNAi-MtAnn1,* n=50 plants). (**S**-**Z**) Representative images of nodules or NP formed in control (**S**, **U**, **W**) and *RNAi-MtAnn1* (**T**, **V**, **X**-**Z**). Fully colonized MtAnn1-RNAi nodules/NP with several ITs (arrows) are shown in **T**-**V**. Abnormal RNAi NPs are shown in **X**-**Z** compared to control NP (arrowheads in **W**). A two-tailed t-test (in **Q**) and a two-tailed Mann-Whitney test (in **R**) were performed in R (asterisks indicate statistical difference, * p<0.05, ** p<0.01). Box plots (**A**, **B**, **O**-**R**) represent the distribution of individual values (indicated as open circles in **A**-**B**, **O** and **Q**-**R,** and as black dots in **P**. Median (central line), mean (solid black circle) and outliers (crosses) are indicated. Scale bars: **E**-**J**, **S**-**Z** = 100 µm, **K**-**N** = 1mm**.** See also Fig. S12-S13.

Fig. 6. *MtAnn1* **mutants show impaired nodule differentiation and function.**

(**A**-**C**) Representative images of entire nodules of wild-type R108S (**A**) and *ann1-3* mutants (**B**-**C**) collected at 15 dpi with *S. meliloti*. (**D**-**M**) Representative images of 1 μm longitudinal sections of 21 dpi nodules of R108S (**D**-**H** and **N**) and *ann1-3* (**I**-**M** and **O**) after staining with toluidine blue and/basic fuchsin. Meristematic (ZI), infection (ZII), interzone (IZ) and nitrogen fixing (ZIII) zones are labelled (**D**-**O**). The amyloplast-rich IZ is indicated between brackets in **D**-**O**. (**P**) Nodule IZ area was measured in images of 1 μm nodule sections from R108S (n=7) and *ann1-3* (n=8) and IZ relative proportion per nodule are shown for each genotype (see detailed

methodology in Fig. S14). (**Q**) Nitrogen fixation capacity was measured in nodulated root segments of R108S (n=14) and *ann1-3* (n=15) collected 21 dpi with *S. meliloti,* from 3 independent experiments (and comprising respectively 365 and 417 nodules). Box plots in **P**-**Q** represent the distribution of individual values (indicated as open circles). Median (central line), mean (solid black circle) and outliers (crosses) are indicated. Two-tailed Welsh (in **P**) or t-tests (in **Q**) of values were performed in R (asterisks indicate significant differences, **p<>0.01). (**R**-**Z**) Ultrastructural analyses by TEM of IZ or ZIII zones in R108S and *ann1-3* nodules. Close up of IZ (**R**-**U**) and IZ/ZIII (**V**-**Z**) of R108S or *ann1-3* nodules with uninvaded cells (UC) and invaded cells (IC) filled with bacteroids. Black arrowheads indicate amyloplasts located near intercellular spaces of IC and within UC cells in the IZ of R108S and *ann1-3,* and also in the IZ/ZIII of *ann1-3*. Differentiated bacteroids (white asterisks) are seen in both R108S and *ann1-3* (**T**, **U**, **Y** and **Z**). However, in *ann1-3* nodules they mostly appear as interzone-type bacteroids[55] imbedded in white vesicle-enriched region (black arrows). Scale bars: **A**-**C** = 1 mm, **D**-**O** = 50 µm, **R**-**Z** = 5 µm. See also Fig. S12-14 for complementary molecular and phenotypic description of *MtAnn1* mutants.

Fig. 7. Modified Ca²⁺ spiking and IT cytoplasmic environment in the absence of *MtAnn1*.

(**A-C**) Transgenic roots expressing the NR-GECO1 Ca²⁺ sensor were generated to monitor nuclear Ca²⁺ oscillations in in *ann1-3* mutant or wild-type (R108S) plants. Variations in Ca²⁺ ion concentration, through changes in the relative intensity of NR-GECO1 fluorescence, is expressed as signal-to-noise ratio (SNR, cf. Methods section). Box plots represent average amplitude of spikes (spike SNR), calculated separately for each nucleus of root hairs with

entrapped rhizobia (RHE) or with a growing IT (IT) in (**A**) R108S (R108S RHE, n=7; R108S IT, n=4) or in (**B**) *ann1-3* (*ann1-3* RHE, n=15; *ann1-3* IT, n=9) or in *S. meliloti*-responsive root hairs in the close competence zone (**C**) (R108S, n=160 and *ann1-3*, n=156) at 1-4 dpi. Data was obtained from three independent experiments. One-tailed t-tests (in **A**-**B**) and a two-tailed Mann-Whitney test (in **C**) were performed in R (asterisks indicate significant differences, *p<0.05; **p<0.01). Median (central line), mean (solid black circle) and outliers (cross) are indicated. (**D**-**K**) Transmission electron microscopy analyses of infected cells from apical zone 2 of WT R108S (**D**-**E**, **H**-**I**, and **H'**-**I'**) or *ann1-3* (**F**-**G**, **J**-**K**, and **J'**-**K'**) 15 dpi nodules. ITs (arrows) in WT are embedded in vesicles (yellow arrowheads) and ER-rich (blue arrowheads) cytoplasmic bridges, which are less visible in *ann1-3*. **H'**-**K'** are complementary inverted LUT images of **H**-**K** that were generated by ImageJ. Scale bars: **D**-**G** = 5 µm, **H**-**K** = 2 µm. See also Fig. S15.

Fig. 8. *MtAnn1* **is required for efficient mycorrhization** *of M. truncatula* **roots.**

(**A-D**) *MtAnn1* promoter activity in independent *M. truncatula* roots after inoculation with *R. irregularis* at 4 wpi (**A-C**) or with water (mock) (**D**). Representative images of root sectors colonized (**A-B**) and non-colonized (**C**) by fungi are shown. Blue GUS staining in the root cortex

(arrows) correlates with *R. irregularis* colonized areas, visualized in yellow by WGA-Alexa Fluor 488 fluorescence staining (WGA). (**E**-**J**) Representative images of ink-stained AM-colonized roots (**E**-**H**) or WGA-stained arbuscules (**F**-**G**, **I**-**J**) formed in R108S or *ann1-3 M. truncatula* roots. (**K**-**L**) Determination of intraradical colonization by *R. irregularis* in WT and *ann1-3* plants by the gridline intersect method. The relative surface of the root system displaying vesicles, arbuscules and/or intraradical hyphae is given in (**K**), while the relative surface corresponding specifically to arbuscules is given in (**L**). Box plots represent values from R108S (WT) (n=36) and *ann1-3* (n=38) root systems at 6 weeks post-inoculation (wpi), from two independent biological experiments. (**M**) Relative arbuscule lengths were compared in R108S (WT) and *ann1-3* roots. (**N**) Q-RT-PCR expression analysis of *MYB1* in total RNA samples from independent R108S (WT) (n=10) or *ann1-3* (=9) plants inoculated with *R. irregularis* (6 wpi, 2 independent experiments). Individual values (open circles or black dots), median (central line) and mean (solid black circle) are indicated in **K**-**N**. Two-tailed Student (**K**-**L**, **N**) and Mann Whitney (**M**) tests were performed in R (asterisks indicate significant differences, **p<0.01, ***p<0.001). Scale bars: **A**-**D** = 100 μm, **E**-**H** = 50 μm, **F**-**J** = 20 μm. See also Fig. S16.