A Haplotype-resolved, Chromosome-scale Genome for *Malus domestica* Borkh. 'WA 38'

Running Title: A high-quality genome for the 'WA 38' apple.

Huiting Zhang[1,2,*], Itsuhiro Ko[3,4,*], Abigail Eaker[3,4,*], Sabrina Haney[5], Ninh Khuu[3], Kara Ryan[6], Aaron B. Appleby[7], Brendan Hoffmann[8], Henry Landis[6], Kenneth Pierro[8], Noah Willsea[9], Heidi Hargarten[2], Alan Yocca[2,10], Alex Harkess[10], Loren Honaas[2,**], Stephen Ficklin[1,**]

[1] Department of Horticulture, Washington State University, Pullman, WA, 99164

[2] USDA Agricultural Research Service, Wenatchee, WA, USA 98801

[3] Department of Plant Pathology, Washington State University. Pullman, WA, 99164

[4] Program of Molecular Plant Sciences, Washington State University, Pullman, WA, 99164

[5] Department of Animal Science, Washington State University. Pullman, WA, 99164

[7] Department of Crop and Soil Science, Washington State University. Pullman, WA, 99164

[8] Integrated Plant Sciences Program, Washington State University. Pullman, WA, 99164

[9] Department of Horticulture, WSU Tree Fruit Research and Extension Center, Wenatchee, WA, 98801

[10] HudsonAlpha Institute for Biotechnology, Huntsville, AL, 35806

* Co-first author.

** Corresponding authors.

Loren Honaas: loren.honaas@usda.gov

Stephen Ficklin: stephen.ficklin@wsu.edu

Abstract

Genome sequencing for agriculturally important Rosaceous crops has made rapid progress both in completeness and annotation quality. Whole genome sequence and annotation gives breeders, researchers, and growers information about cultivar specific traits such as fruit quality, disease resistance, and informs strategies to enhance postharvest storage. Here we present a haplotype-phased, chromosomal level genome of *Malus domestica*, 'WA 38', a new apple cultivar released to market in 2017 as Cosmic Crisp ®. Using both short and long read sequencing data with a k-mer based approach, chromosomes originating from each parent were assembled and segregated. This is the ***first*** pome fruit genome fully phased into parental haplotypes in which chromosomes from each parent are identified and separated into their unique, respective haplomes. The two haplome assemblies, 'Honeycrisp' originated HapA and 'Enterprise' originated HapB, are about 650 Megabases each, and both have a BUSCO score of 98.7% complete. A total of 53,028 and 54,235 genes were annotated from HapA and HapB, respectively. Additionally, we provide genome-scale comparisons to 'Gala', 'Honeycrisp', and other relevant cultivars highlighting major differences in genome structure and gene family circumscription. This assembly and annotation was done in collaboration with the American Campus Tree Genomes project that includes 'WA 38' (Washington State University), 'd'Anjou' pear (Auburn University), and many more. To ensure transparency, reproducibility, and applicability for any genome project, our genome assembly and annotation workflow is recorded in detail and shared under a public GitLab repository. All software is containerized, offering a simple implementation of the workflow.

Keywords

Apple genomics, *Malus domestica* 'WA 38', genome sequence, comparative genomics, plant genomics, haplotype-resolved assembly, genome annotation

Introduction

For economically important crop species, having full-resolution reference genomes aids in the understanding of traits associated with commodity quality, disease resistance, long-term storage, and shelf life. Apple (*Malus domestica*) is the number one consumed fruit in the United States, with a Farm-Gate Revenue of $3.2 billion in the U.S. (USApple, 2024), and $78 billion globally (FGN, 2020). There are over 7,000 apple varieties grown world wide (Washington Apple Commission, 2024), each with unique colors, flavors, and textures (N.C. Cooperative Extension, n.d.). Therefore, a single genome is unlikely to capture the complexity of all cultivars within this highly heterozygous species (Li *et al.* 2022b; Zhang *et al.* 2022). One such cultivar is 'WA 38', commercially released as Cosmic Crisp® in 2017 by the Pome Fruit Breeding Program at Washington State University's (WSU) Tree Fruit Research and Extension Center (Figure 1 A, B) and has reached the top 10 best selling apple cultivars in the United States (Truscott, 2023). 'WA 38' is a cross between 'Honeycrisp' and 'Enterprise', made using classical breeding methods in 1997. One parent, 'Honeycrisp', is well-known for its crisp texture, firmness retention in storage, disease resistance, and cold hardiness, but is highly susceptible to production and postharvest issues (Khan *et al.* 2022). The other parent, 'Enterprise', is an easy-to-grow cultivar that has extended postharvest storage capabilities, however it is not widely cultivated commercially due to its less desirable eating quality (Crosby *et al.* 1994). Their resulting cross has been met with favorable reviews for its appealing color, texture, flavor, cold hardiness, long-term storage capabilities (>1 year), and scab resistance (Evans *et al.* 2012). However, it inherited undesirable traits as well, such as a propensity for physiological symptoms that may be related to mineral imbalances (Sallato *et al.* 2021; Sheick *et al.* 2023), maturity at harvest (Serra et al 2023), and an 'off' flavor that has been brought up by consumers that may be the result of improper picking times, crop load management, handling/packing practices or other post harvest processes (Mendoza, M., Hanrahan, I., & Bolaños, G., 2020). Most concerning is green spot (Figure 1 C, D, and F), a corking disorder that seems to be unique to 'WA 38', but with etiology similar to disorders associated with mineral imbalances such as bitter pit and drought spot (Sheick *et al.* 2022, 2023). The propensity for and cause of physiological disorders often differs on a cultivar-by-cultivar basis (Pareek 2019), and a genetic basis for such predispositions is likely (Liebhard *et al.* 2003; Johnston and Brookfield 2012; Di Guardo *et al.* 2013; Lum *et al.* 2016). Thus, improved resolution of cultivar-specific genomic differences is critical for advancing our understanding of how economically important traits, such as physiological disorders, are inherited and how they can be managed more efficiently.

Figure 1. 'WA 38', a cultivar of apple developed by the Washington State University Apple
Breeding Program (a cross between 'Honeycrisp' and 'Enterprise'), marketed as Cosmic
*Crisp®. **A)** 'WA 38' apples ready for harvest on the mother tree, located at the WSU and USDA-*
*ARS Columbia View Research Orchard near Orondo, WA, USA. **B)** The 'WA 38' mother tree. **C***
*& **D**) Green spot, a corking disorder which results in green blemishes on the fruit peel and*
brown, corky cortex tissue. Symptom severity generally increases during fruit maturation and
*time in storage, resulting in cullage. **E**) Natural peel greasiness as a result of more advanced*
maturity at harvest can interfere with artificial waxes applied in the packinghouse after removal
*from postharvest storage, creating unappealing, dull spots. **F**) Green Spot symptoms can begin*
to appear while fruit is still developing on the tree. Photo Credits: A&B: Heidi Hargarten/USDA-
ARS; C&D: Bernardita Sallato/WSU; E: Carolina Torres/WSU; F: Ross Courtney/Good Fruit
Grower.

To develop full resolution reference genomes of superior quality, having skilled bioinformaticians
is required. To train the next generation of bioinformaticians for agricultural genomic research, a
national effort spearheaded by Auburn University, HudsonAlpha Institute for Biotechnology, and
Washington State University was started in 2021 - The American Campus Tree Genomes
project (ACTG). ACTG aims to break through institutional barriers that have traditionally
prevented many students from accessing valuable, hands-on research projects and experience
in bioinformatics (Sharman, S, n.d.). To accomplish this goal, a course has been developed to
involve students in genome projects from inception, through analysis, to publication (Harkess,
2022). During the course, students learn genome assembly and annotation workflows using the
raw sequence data from genomes of beloved trees (e.g., Toomer's oak and 'd'Anjou' pear
(Yocca *et al.* 2024) at Auburn University, Sabal palm at University of South Carolina) and are

listed as authors on the final publication. The 'WA 38' genome introduced here was developed

through ACTG by students from Washington State University, presenting three major outcomes:

1) a fully annotated, chromosomal level, haplotype-resolved genome of 'WA 38' utilizing PacBio

HiFi, Dovetail Omni-C, and Illumina DNA and RNA sequencing data, 2) identification of unique

regions of interest using a comparative genomics approach with other economically important

M. domestica cultivars including 'Gala', 'Fuji', and 'Honeycrisp', and 3) establishment of a

containerized, reproducible, flexible, high performance computing workflow for complete

genome assembly and annotation (Figure 2, Supplemental Figure 1).

Figure 2. Schematic chart of 'WA 38' genome project.

Methods

Workflows developed for each stage of the project and the summary workflow of the whole

project are available in Supplemental Figure S1. Scripts with parameters for each computation

step and methods in markdown format are available in GitLab at: https://gitlab.com/ficklinlab-

public/wa-38-genome.

Sample Collection

Approximately 20 grams of young leaf material was harvested from the *Malus domestica* 'WA

38' mother tree at the Washington State University and USDA-ARS Columbia View Research

Orchard near Orondo, WA, USA and flash-frozen in liquid nitrogen. Tissue was sent to the

HudsonAlpha Institute for Biotechnology in Huntsville, AL, USA for DNA extraction, sequencing

library preparation, and sequencing, following the same protocol (detailed below) used to

generate the 'd'Anjou' pear genome (Yocca *et al.* 2024).

To assess heterozygosity and genome size of 'WA 38', DNA was extracted using a standard

CTAB isolation method (Doyle and Doyle 1987). Illumina TruSeq DNA PCR-free libraries were

constructed from 3 ug of input DNA following the manufacturer's instruction, and sequenced on

an Illumina NovaSeq6000.

For PacBio HiFi sequencing, high molecular weight DNA was isolated using a Nanobind Plant

Nuclei Big DNA kit (Circulomics-PacBio, Menlo Park, CA), with 4 g of input tissue and a 2-hour

lysis. DNA purity, quantity, and fragment sizes were measured via spectrophotometry, Qubit™

dsDNA Broad Range assay (Invitrogen™), and Femto Pulse system (Agilent, Santa Clara, CA),

respectively. DNA that passed quality control was sheared with a Megaruptor (Diagenode,

Denville, NJ) and size-selected to roughly 25 kb on a BluePippin. The SMRTbell Express

Template Prep Kit 2.0 (PacBio, Menlo Park, CA) was used to construct the PacBio sequencing

library, and HiFi reads were produced using circular consensus sequencing (CCS) mode with

two 8M flow cells on a PacBio Sequel II long-read system.

To scaffold PacBio HiFi contigs into chromosome pseudomolecules, a Dovetail Genomics

Omni-C library was generated using 1 g of flash-frozen young leaf material as input following

the manufacturer's instruction (Dovetail Genomics, Scotts Valley, CA), and sequenced on an

Illumina NovaSeq6000 S4 PE150 flow cell.

Sequence quality assessment and genome complexity analysis

Adapter sequences were trimmed from the raw Illumina shot-gun DNA reads using fastp

(v0.23.2) (Chen *et al.* 2018) with all the other trimming functions disabled. Both the raw and

trimmed Illumina reads, PacBio HiFi reads, and Omni-C Illumina reads were assessed for

quality with FastQC (v0.11.9) (Andrews, 2010). Genome complexity, *i.e.* nuclear genome size

and ploidy, was estimated using Jellyfish (v2.2.10) (Marçais and Kingsford 2011). With trimmed

paired-end Illumina reads as input and a k-mer size set to 21, a k-mer count file was generated

by Jellyfish. The k-mer histogram, also created by Jellyfish, was visualized in GenomeScope

(v1.0) (Vurture *et al.* 2017) with the following parameters: k-mer size = 21, Read length = 151,

and Max k-mer coverage = 1000. A summary statistic report of the sequence quality and

complexity analysis was generated with MultiQC (v1.13a).

Genome Assembly

Genome assembly and scaffolding

Phased haplomes were assembled by Hifiasm (v0.16.1) (Cheng *et al.* 2021) with default

parameters, using both the Omni-C data and the PacBio HiFi long reads. The statistical

summary of the assembly was produced using the assimilation_stats Perl scripts described in

(Earl *et al.* 2011). Both hifiasm-assembled haplotype unitigs were then sorted by MUMmer

(v3.23) (Kurtz *et al.* 2004) using the 'nucmer' function with flag -maxmatch. The resulting files

were uploaded to the Assemblytics web server (http://assemblytics.com; Nattestad and Schatz

2016) to visualize structural variations in two haplotype unitigs with the default settings.

Following the initial assembly step, bwa (Li and Durbin 2009) was used to index the draft

contigs, and subsequently to align the Hi-C reads to the indexed contigs. The sorted files were

input into Phase Genomics hic_qc (https://github.com/phasegenomics/hic_qc;

phasegenomics, n.d.) to validate the overall quality of the library. Both assembled haplomes

were scaffolded into chromosomes with YaHS (Danecek *et al.* 2021; Zhou *et al.* 2022), using

default parameters.

Assembly curation, completeness assessment, and telomere identification

Hi-C files were generated using YaHS Juicer Pre (v1.2a.2-0) with flag -a allowing manual

curation. The resulting files were used as input for Juicer Tools Pre (v 1.22.01) to generate Hi-C

contact maps (Durand *et al.* 2016; Zhou *et al.* 2022). Juicebox Assembly Tools (v1.11.08) was

used to explore the Hi-C maps for miss-assemblies (Robinson *et al.* 2018). After manual

examination of the Hi-C maps, the final genome assembly was generated by linking remaining

files from YaHS Juicer Pre and original HiFi scaffold, using YaHS Juicer Post (v 1.2a.2-0,

(Durand *et al.* 2016)).

For consistency and reproducibility, 'WA 38' chromosomes were renamed and reorientated to

match published genomes. First, MUMmer (v3.23) was used to align the 'WA 38' assembly to

the 'Gala' v1 HapA assembly using the –maxmatch parameter for unique matches (Kurtz *et al.*

2004; Sun *et al.* 2020). Next, Assemblytics dotplot was used to identify 'WA 38' scaffolds that

aligned with the 'Gala' v1 chromosomes and 'WA 38' scaffolds were renamed accordingly. To

determine orientation, each 'WA 38' chromosome was aligned to the corresponding 'Gala' v1

chromosomes using LASTZ (v 1.02.00) implemented in Geneious (v9.0.5; Harris. 2007) with the

'search both strands' option. The chromosomes on the reverse stand were reoriented with the

Reverse Complement (RC) function in Geneious (Supplemental Figure S2). The resulting

assembly was searched against NCBI's RefSeq Plastid database (NCBI Organelle genome

resources, n.d.) using megablast and a custom virus and bacteria database using Kraken

(v2.1.3; Wood and Salzberg, 2014) to identify contaminants. Scaffolds identified as plastid or

microbe contaminants were removed in the assembly.

The cleaned assembly was compared to the 'Honeycrisp' genome assembly with a kmer

approach using meryl (v1.4.1, Rhie et al., 2020). Chromosomes with a 'Honeycrisp' origin were

placed in HapA, whereas the others were placed in HapB.

The two final haplome assemblies were compared to each other using MUMmer and

Assemblytics as described above to identify structural variants. Benchmarking universal single-

copy gene orthologs (BUSCO, v5.4.3_cv1) analysis was performed in genome mode with the

eudictos_odb10 database to assess completeness (Manni *et al.* 2021).

Structural and functional annotation

Repeat annotation

Repetitive elements from both haplotypes were annotated using EDTA (v2.0.0; Ou *et al.* 2019),

with flags 'sensitive=1' and 'anno=1'. The full coding sequence from 'Gala' HapA, obtained from

the Genome Database for Rosaceae (GDR; Jung *et al.* 2019), was used as reference to aid

repeat finding. The custom transposable element library generated by EDTA was then imported

to RepeatMasker (Smit *et al.* 2013-2015) to further identify potentially overlooked repetitive

elements and create masked versions of the genome. Three masked versions were generated:

softmasked, N masked, and X masked.

Telomeres were identified by tidk (v0.2.41; Brown *et al.* 2023) with the following parameters:

explore --minimum 2 --maximum 20 and the default database provided by the software.

Gene Annotation

To annotate gene space, a combination of *ab initio* prediction and evidence-based prediction

were performed on the softmasked assemblies with two rounds of BRAKER using transcriptome

and homologous protein evidence. PASA (v2.5.2; Haas *et al.* 2003) was then used to refine

gene models and add UTR annotation. Lastly, a custom script was used for filtering. The

detailed methods are described below.

BRAKER1 - annotation with transcriptome evidence

To perform transcriptome guided annotation, same RNA-seq data from Khan et al. (2022) (eight

tissue types from six pome fruit cultivars including 'WA 38', BioProject: PRJNA791346) were

first aligned to the 'WA 38' haplomes using the STAR aligner implemented in GEMmaker

(v2.1.0) Nextflow workflow (Hadish *et al.* 2022). The resulting read alignments were used as

extrinsic evidence in BRAKER1 (Hoff *et al.* 2016) to predict gene models in each softmasked

haplome with the following parameters: --softmasking, --UTR=off, --species=malus_domestica.

BRAKER2 - annotation with homologous protein evidence

To provide protein evidence data for BRAKER2 (Brůna *et al.* 2021), protein sequences from

three sources were used: 1) Predicted protein sequences of 13 Rosaceae genomes retrieved

from GDR (*Fragaria vesca* v4a2 (Li *et al.* 2019), *Malus baccata* v1.0 (Chen *et al.* 2019), *M.*

domestica var.Gala v1 (Sun *et al.* 2020), *M. domestica* var.GDDH13 v1.1 (Daccord *et al.* 2017),

M. domestica 'Honeycrisp' v1.0 (Khan *et al.* 2022), *M. sieversii* v1 (Sun *et al.* 2020), *M.*

sylvestris v1 (Sun *et al.* 2020), *Prunus persica* v2.0.a1 (Verde *et al.* 2017), *Pyrus betulifolia v1.0*

(Dong et al. 2020), *P. communis* 'd'Anjou' v2.3 (Yocca *et al.* 2023), *P. pyrifolia* 'Nijisseikiv' v1.0

(Shirasawa *et al.* 2021)*, Rosa chinensis* 'Old Blush' v2.0.a1 (Raymond *et al.* 2018), and *Rubus*

occidentalis v3 (VanBuren *et al.* 2018)); 2) Peptide sequences predicted from *de novo*

transcriptome assemblies used in the 'Honeycrisp' genome annotation (Khan *et al.* 2022); and

3) Viridiplantae OrthoDBv11 protein sequences (Kuznetsov *et al.* 2022). In the same manner as

BRAKER1, the softmasked haplome assemblies were used as input.

TSEBRA - transcript selection

The gene annotation results from BRAKER1 and BRAKER2 were merged and filtered based on

the supporting evidence using TSEBRA (v.1.0.3; Gabriel *et al.* 2021) with the default

configuration (file obtained in August 2022) provided by TSEBRA developers.

PASA - gene model curation and UTR annotation

Two sources of transcriptome assembly evidence were obtained to facilitate PASA annotation:

1) Transcript sequences predicted from *de novo* transcriptome assemblies used by 'Honeycrisp'

genome annotation; and 2) Reference guided assemblies created with read alignment files from

GEMmaker (see the BRAKER1 section for details) using Trinity (Grabherr et al. 2011) with max

intron size set to 10,000. Four rounds of PASA (v2.5.2) curation were performed using the

aforementioned evidence and a starting annotation. The first round of PASA curation used

TSEBRA annotation as the starting annotation, and annotations from the previous round were

used as the starting annotation for rounds two through four. The curation results from each

round were manually inspected using the PASA web portal. No significant improvement was

observed after the fourth round of curation, therefore no further rounds were performed.

Gene model filtering and gene renaming

Repeat and gene model annotations were loaded to IGV (v2.15.1; Robinson *et al.* 2011) for

manual inspection. Three types of erroneous gene models were observed consistently

throughout the annotations. Type 1: Genes overlapping with repeat regions (e.g. transposon

was wrongly annotated as a gene), Type 2: Gene models overlapping with each other on the

same strand (e.g. single gene was wrongly annotated with multiple gene models), and Type 3:

Gene models with splice variants that had no overlap (e.g. different genes were wrongly

annotated as the single gene's splice variants). A custom script was used to address these

errors. The Type 1 error was resolved by removing genes with 90% of its coding region

overlapping with repeat regions. The Type 2 error was resolved by removing the shorter gene of

a pair that overlaps on the same strand. The Type 3 error was resolved by splitting splice

variant models with no overlap into two separate gene models. Finally, custom scripts were

used to generate the final annotation files (gene, mRNA, cds, protein, gff3) and rename genes

to match the naming convention proposed by GDR

(https://www.rosaceae.org/nomenclature/genome). The longest isoforms of each transcript were

needed for some downstream analysis and were extracted using a modified version of the

get_longest_isoform_seq_per_trinity_gene.pl script provided by Trinity (Grabherr *et al.* 2011).

Functional Annotation

The final gene sets from both 'WA 38' haplomes were annotated using EnTAPnf (Hart *et al.*

2020) with Interproscan, Panther, RefSeq, and uniprot_sprot databases that are automatically

downloaded using the download.py script provided by EnTAPnf.

Comparative Analysis

Synteny analysis

276 A synteny comparison was performed using GENESPACE (Lovell *et al.* 2022) with five *Malus*

277 *domestica* assemblies and annotations (GDDH13 from Daccord *et al.* 2017), both haplomes of

278 'Honeycrisp' from Khan *et al.* 2022, and both haplomes from 'WA 38'). Default parameters were

279 used. Only the longest isoforms were used for 'WA 38'.

Gene family analysis

Gene family, or orthogroup, analyses were carried out to identify shared and unique gene

families in 'WA 38' and other pome fruit genomes (i.e., *Malus* sp. and *Pyrus* sp. A full list of

genomes analyzed can be found in Supplemental Table S1) following the method described by

(Khan *et al.* 2022). Briefly, predicted protein sequences from the selected pome fruit genomes

were classified into a pre-computed orthogroup database (26Gv2.0) using the 'both HMMscan

and BLASTp' option implemented in the GeneFamilyClassifier tool from PlantTribes2 (Wafula *et*

al. 2022). Overlapping orthogroups among *M. domestica* genomes were calculated and

visualized with the UpSet plot function implemented in TBtools v2.030 (Chen *et al*., 2023).

A Core OrthoGroup (CROG) - Rosaceae gene count analysis was carried out following the

method described by (Wafula *et al.* 2022). First, a CROG gene count matrix was created by

counting genes classified into CROGs from each pome fruit genome. Next, the matrix was

visualized as a clustermap using the Seaborn clustermap package (CROGs with standard

deviation of 0 were removed prior to plotting) with rows normalized by *z*-score. Finally, the

derived *z*-score of CROGs in each genome was summarized into a boxplot to illustrate z-score

distribution using the boxplot function in Seaborn.

Gene evidence source mapping

Each gene was screened against the following evidence source: Transcriptome evidence

covering the entire gene (Full support); Transcriptome evidence covering part of the gene (Any

support); Homologous protein evidence covering the entire gene (Full support); Homologous

protein evidence covering part of the gene (Any support); Has a EnTAP functional annotation

from any database; Assignment to a PlantTribes2 Orthogroup. Transcriptome and homologous

protein evidence were mapped to genes by using "selectSupportedSubsets.py" script provided

by BRAKER (Brůna *et al.* 2021) and BEDtools (Quinlan and Hall 2010). Summaries of evidence

source mapping are available in Supplemental Table S2 and S3. The following subsets of genes

were extracted and were subject to BUSCO completeness analysis and CROG gene count

analysis: Subset 1, Genes with **full** support from either RNAseq or homologous protein

evidence; Subset 2, Genes with **any** support from either RNAseq or homologous protein

evidence; Subset 3, Genes from Subset 1 plus gene with both EnTAP and PlantTribes2

annotation; Subset 4, Genes from Subset 1 plus genes with either EnTAP or PlantTribes2

annotation.

Chloroplast & Mitochondria Assembly and Annotation

The chloroplast genome was assembled from trimmed Illumina shot-gun DNA reads using

NOVOplasty (v4.3.1; Dierckxsens *et al.* 2017) with the *Malus sierversii* chloroplast genome

(NCBI accession ID: MH890570.1; Naizaier *et al.* 2019) as the reference sequence and the

NOVOplasty *Zea mays RUBP* gene as the seed sequence. The assembled chloroplast was

annotated using GeSeq Web Server (website accessed on Dec. 19th, 2023; Tillich *et al.* 2017)

with settings for 'circular plastid genomes for land plants' and the following parameters:

annotating plastid inverted repeats and plastid trans-spliced *rps12*. Additionally, annotations

from third party softwares Chloë (v0.1.0) and ARAGORN (v1.2.38), as well as a BLAT (v.35×1)

search against all land plant chloroplast reference sequences (CDS and rRNA), were integrated

with the GeSeq results. Genes identified by multiple tools were manually reviewed to produce

the final, curated annotation. The curated chloroplast annotation was visualized by OGDRAW

(v1.3.1; Greiner *et al.* 2019).

The mitochondrial genome sequence was isolated from the Hifiasm assembled contigs using

MitoHifi (v3.2; Uliano-Silva *et al.* 2023). The *M. domestica* mitochondria sequencing from NCBI

(NC_018554.1; Goremykin *et al.* 2012), which contained 57 genes consisting of 4 rRNAs, 20

tRNAS, and 33 protein-coding genes, was used as the closely related reference sequence.

Briefly, MitoHifi compares the assembled contigs to the reference mitogenome using the BLAST

algorithm. The resulting contigs were manually filtered by size and redundancy and then are

circulated. To increase the annotation quality, GeSeq was deployed in mitochondrial mode with

the *M. domestica* NCBI RefsSeq sequence to annotate the 'WA 38' mitochondria assembly.

Fragmented genes from the annotation were manually removed prior to visualization in

OGDRAW (v1.3.1; Greiner *et al.* 2019).

Results

A Complete, Reproducible, Publicly-available Workflow

Results

A Complete, Reproducible, Publicly-available Workflow

To ensure transparency and reproducibility, the 'WA 38' Whole Genome Assembly and

Annotation (WA 38 WGAA) project workflow was made publicly accessible through a GitLab

repository (https://gitlab.com/ficklinlab-public/wa-38-genome). This repository contains the

complete manual workflow for assembly and annotation of the genome. It organizes each step

in order of execution, using ordered, numeric directory prefixes where each directory includes

detailed method documentation and scripts that were executed for each analysis. All parameter

settings, as well as any command line manipulation of the files generated are noted in the

scripts or methods. Summary diagrams for the manually executed workflow are available in

Supplemental Figure S1. All software utilized in the project has been containerized and shared

on Docker Hub (https://hub.docker.com/u/systemsgenetics). Any user that follows the workflow

can retrieve the public data and repeat the steps to reproduce the results. Leveraging these

resources from the 'WA 38' WGAA project, and as part of our commitment to knowledge

sharing, we have initiated an American Campus Tree Genome (ACTG) course GitHub

organization (https://github.com/actg-course/). This organization comprises three main

repositories: 1) wgaa-compute: a generic whole genome assembly and annotation workflow

template, derived from the 'WA 38' WGAA project, that can be adapted for other species; 2)

wgaa-docker: the Docker recipes for all the software employed in the project; and 3) wgaa-doc:

an open-source and editable documentation repository containing teaching materials for current

and future ACTG instructors, providing a collaborative space for instructors to learn from and

contribute to the enhancement of the course materials.

Nuclear Genome Assembly

Nuclear Genome Assembly

Sequence Quality Assessment

Raw sequencing data (Table 1) was assessed for read quality. The Illumina shotgun short read

data consisted of 807.2 million total reads with a mean length of 151bp for a total of 121.9

Gigabases (Gb) of data after adapter trimming. Filtered Illumina data GC content is 38% and

has 91.8% Q20 bases and 83.4% Q30 bases. Duplication rates ranged from 23.3% to 27.8%.

PacBio long read raw data consisted of 3.9 million reads from 85-49,566bp in length for a total

of 60.0 GB. Sequence duplication rates ranged from 2.2% to 2.4%. PacBio sequence GC

content is 38%, same as the Illumina data. In addition, a 402x coverage (201x for each

haplome) of Omni-C data was generated to facilitate the assembly and phasing.

Table 1. Yield of Illumina DNA short reads (Shotgun and Omni-C) and PacBio HiFi sequencing

reads from young leaf tissues of 'WA 38'.

	Long Read	Short Read	
	PacBio HiFi	Shotgun DNA seq	OmniC-Seq
Total read number	3,870,263	807,220,896	1,730,268,360
Number of bases (Gb)	60.0	121.9	261.3
Coverage*	92x	188x	402x
Average length (bp)	15,495	151	151

* calculated with the size of a haploid genome (650 Mb).

Genome complexity

Using a *k*-mer frequency approach, genome characteristics such as heterozygosity and genome

size were estimated (Figure 3). Analysis of both short and long reads resulted in an estimated

heterozygosity of ~1.35%, similar to estimates from the 'Honeycrisp' cultivar (1.27%; Khan *et al.*

2022). Estimate genome size was 467Mb from the short reads and 606Mb from the long reads.

These estimates are lower than expected from other apple genomes ('Honeycrisp': 660-674 Mb;

Khan *et al.* 2022 and 'Golden Delicious': ~701 Mb; Li *et al.* 2016) and the final assembly (Table

2). Additionally, the percent of unique sequence was estimated at 69.5% for the short reads and

53.4% for the long reads, with the long read estimate being more consistent with what is

expected from the 'Honeycrisp' (51.7%; Khan *et al.* 2022) and of wild apple species *Malus*

baccata (58.6%; Chen *et al.* 2019).

Genome assembly, scaffolding, and curation

For initial assembly, scaffolding, and curation, two unsorted, phased haplomes, called Hap1 and

Hap2, were assembled and scaffolded using both PacBio long reads and Omni-C short reads.

Hi-C maps of the haplome assemblies show no mis-assemblies (Supplemental Figure S3). For

Hap1 and Hap2, a total of 22 joins and 20 joins, respectively, were made in the scaffolding step

to build the final assemblies into 17 chromosomes each, with the remaining scaffolds

representing unincorporated contigs. Unincorporated contigs were investigated and found to be

bacterial or other contamination and were removed. After removing contaminants, Hap1 is

645.41 Mb in length with an N50 of 36.1 Mb, while Hap2 is 651.07 Mb in length with an N50 of

37.2 Mb. Additional assembly statistics for both haplomes are included in Supplemental Table

S4. 'WA 38' has a comparable genome size to other previously sequenced apple cultivars,

including its parent 'Honeycrisp' (Khan *et al.* 2022). Notably, the 'WA 38' scaffold N50 is among

the longest across all published apple genomes, indicating high levels of assembly contiguity

(Supplemental Table S5).

Figure 3. Genome complexity of 'WA 38' genome using PacBio long read data (A) and Illumina

short read (B). The output figure was generated by GenomeScope (k=21).

Haplotype-binning, Structural Comparison, and Completeness Assessment

The K-mer based binning method identified the origin of chromosomes in each haplome

assembly. Ten out of the 17 chromosomes in Hap1 originated from 'Honeycrisp', while the other

seven were from 'Enterprise'. After reorganizing the chromosomes based on parent

contribution, the haplome containing all the 'Honeycrisp' origin chromosomes is designated as

HapA, whereas the 'Enterprise' originated haplome is designated as HapB. HapA and HapB are

structurally similar; a total of ~44 Mb are affected by structural variants and are mainly

contributed by indels and repeat expansion and contractions (Supplemental Table S6 and

Supplemental Figure S4). Additionally, three large inversions are observed on chromosomes 1,

11, and 13 (Supplemental Figure S5). Based on the BUSCO analysis, both the HapA and HapB

assemblies were 98.7% complete, with only 19 BUSCOs missing and 12 partially detected

(Supplemental Table S7). This BUSCO score suggests high genome completeness for both

haplomes, comparable to the 'Fuji' apple genome assemblies, which is most contiguous of all

apple genomes to date (Table 2 and Supplemental Table S5; Li et al., 2024).

Table 2. Comparison of genomic features and assembly statistics of the 'WA 38' genome and

previously published apple genomes.

	'WA 38'		'Honeycrisp'		'Antonovka'	'Gala'	GDDH13	'Fuji'
	HapA	HapB	HapA	HapB				
Number of Scaffold	17	17	473	215	168	812	1,081	1,358
Haploid genome size (Mb)	645.41	651.07	674	660	643.5	652.4	709.6	736.9
N50 (Mb)	36.1	37.2	31.6	32.8	35.85	23.9	5.5	36.8
L50	8	8	8	8	8	8	NA	9
Number of protein-coding genes	53,028	54,235	47,563	48,655	45,085	45,352	45,116	49,972
Complete BUSCO (%) Assembly	98.7	98.7	98.6	98.7	97.6	97.9	98	98.8
Complete BUSCO (%) Annotation	98.5	98.4	96.8	97.4	97.25	95.5	96.1	97.2
Number of orthogroups in 26Gv2.0	10,494	10,511	10,350	10,366	10,293	10,095	10,117	10,243
Reference	This paper		Khan et al. 2022		Švara et al., 2023	Sun et al. 2021	Daccord et al. 2017	Li et al., 2024

NA: Data not available. For consistency, genome statistics and BUSCO analyses were

performed on the publicly available genomes using the same methods used for 'WA 38', except

for N50 and L50 of GDDH13 as the scaffold assembly is not publicly available. 'Antonovka' data

is the average of the two haplomes. The unphased version of 'Fuji' was used. A more in-depth

comparison is available in Supplemental Table S5.

Nuclear Genome Structural Annotation

Repeat annotation

In both haplomes, approximately 58.7% of the assembly was predicted to be repetitive regions

by EDTA (Ou *et al.* 2019; Table 3). RepeatMasker identified an additional 4% repeat elements,

resulting in a total of 62.7% repeat regions in both HapA and HapB, comparable to the

'Honeycrisp' genome (Khan *et al.* 2022). In both haplomes, the most dominant type of repeat

element is long terminal repeat (LTR), followed by Terminal Inverted Repeat (TIR) (Table 3,

Supplemental Table S8), consistent with that in 'Honeycrisp'. We also compared the repeat

landscape of 'WA 38' with 'd'Anjou' pear which was annotated with the same methodology.

While they share the major repeat classes, 'd'Anjou' pear has a much lower percentage of

repeat elements (Table 3).

Through telomere search in each haplotype, we discover that telomere repeat regions are

present in almost every chromosome of each haplome. The most enriched telomere repeat unit

is a 7-mer "AAACCCT" and its reverse complement "AGGGTTT", which has been reported as

overrepresented in the *Arabidopsis thaliana* genome (Choi *et al.* 2021), opposed to "CCCATTT"

and "TTTTAGGG" reported in the most recent T2T 'Golden Delicious' apple genome (Su *et al.*

2024). A list of telomere repeat regions and units for both haplotypes were deposited in

Supplemental Table S9.

Table 3. Summary of repetitive element annotation in the 'WA 38' and other apple genomes.

Class		'WA 38' (%)		'Honeycrisp' (%)		'd'Anjou' pear (%)	
		HapA	HapB	HapA	HapB	Hap1	Hap2
LTR	Copia	9.37	10.22	9.73	9.6	5.6	5.73
	Gypsy	17.19	18.32	20.29	17.8	12.32	12.88
	unknown	16.37	14.52	14.89	16.86	8.46	10
TIR	CACTA	1.94	2.15	2.21	1.95	1.4	1.4
	Mutator	3.96	4.18	4.16	4.25	3.47	3.41

		This paper		Khan et al. 2022		Yocca et al., 2023	
	PIF Harbinger	2.4	2.52	2.43	2.6	1.81	1.81
	Tc1_Mariner	0.16	0.24	0.15	0.27	0.13	0.11
	hAT	2.15	2.37	2.3	2.31	0.58	0.84
	polinton	0	0	0	0.01	0	0
nonLTR	LINE_element	0.14	0.16	0.18	0.17	0.14	0.14
	unknown	0.11	0.09	0.09	0.18	0.06	0.06
nonTIR	helitron	3.41	2.20	2.95	3.18	1.56	1.92
Other repeat region		1.52	1.74	2.91	2.78	3.98	4.22
RM*		3.98	3.99	NA	NA	NA	NA
Total		62.71	62.71	62.43	61.97	39.78	42.52
Reference		This paper		Khan et al. 2022		Yocca et al., 2023	

repeat regions annotated by RepeatMakser

Gene space annotation

To annotate the gene space, we utilized a combination of *ab initio* prediction and evidence-

based prediction with transcriptome and homologous protein, functions implemented in

BRAKER2 (Brůna *et al.* 2021). However, BRAKER2 was unable to annotate UTR regions and

yielded erroneous gene models and splice variants (Supplemental Figure S6). Therefore, the

gene models were further processed with PASA (Haas *et al.* 2003) and a custom script. A total

of 53,028 and 54,235 genes were annotated from HapA and HapB, respectively, more than

most published apple genomes (Table 2, Supplemental Table S10). The complete BUSCO

scores for HapA and HapB annotations are 98.5% and 98.4%, respectively, the highest score

among all *M. domestica* genomes sequenced to date (Supplemental Table S5). The average

protein annotated from HapA and HapB contains 361.3 and 356.4 amino acids, respectively,

similar to that of other *M. domestica* annotations (Supplemental Table S11). On average, 1.3

splice variants were identified for each gene in both HapA and HapB annotations. The only

other apple genome with splice variant annotation is 'Honeycrisp', and on average, 1.05 splice

variants were annotated per gene (Supplemental Table S11). Additionally, 53.5% and 52.2% of

the annotated transcripts from HapA and HapB, respectively, contain untranslated regions

(UTRs). Notably, 'WA 38' is the only other apple genome besides 'GDDH13' and 'Fuji' that has

more than half of the genes annotated with UTRs.

The 'WA 38' genes were named in accordance with the convention following guidance from the

Genome Database for Rosaceae (GDR). This convention was first proposed by our group for

the 'Honeycrisp' genome and was later adopted with modification by GDR (Gene name

example: *drMalDome.wa38.v1a1.ch10A.g00001.t1*). This convention meets recommendations

proposed by the AgBioData consortium to reduce gene ID ambiguity and improve

reproducibility.

Nuclear Genome Functional Annotation

EnTAP (Hart *et al.* 2020) functional annotation assigned functional terms to 89.5% and 88.8% of

proteins annotated from HapA and HapB, respectively. Specifically, an average of 83% and

55% of all proteins (including both HapA and HapB) have strongly supported hits in the NCBI

RefSeq (O'Leary *et al.* 2016) and UniProt database, respectively, 75% were annotated with an

InterPro term, and 88% have functional annotations from at least one of the databases included

in InterProScan. EggNOG (O'Leary *et al.* 2016; Huerta-Cepas *et al.* 2019) search provided

additional function information: 90% of the annotated proteins were assigned into EggNOG

orthogroups, 84% were annotated with protein domains, 21% were classified into KEGG

pathways, and 63%, 53%, and 61% proteins were annotated with GO biological process,

cellular component, and molecular function terms, respectively (Supplemental Table S12).

Comparative Analyses

Synteny and gene family analyses were performed to investigate the similarity and unique

features of 'WA 38' genome to other closely related species and cultivars.

Synteny analysis was performed to compare the genomes of 'WA 38', one of its parents,

'Honeycrisp', and the most referenced apple genome, 'GDDH13', using GeneSpace. The two

'WA 38' haplomes are highly collinear with each other and with the other apples, especially the

two 'Honeycrisp' haplomes. Although inversions at various scales were observed between the

two 'WA 38' haplomes e.g. large inversions on chromosomes 1, 11, 13 (Supplemental Figure

S5 and S7), they have minor effects on gene order (Figure 4), likely due to the small number of

genes annotated from those inverted regions.

Figure 4. Riparian plot comparing 'WA 38' Haplotype A and B with 'Honeycrisp' Haplotype A

and B and 'Golden Delicious' (GDDH13) genomes by gene rank order.

Gene family analysis is performed using PlantTribes2 and the pre-constructed 26Gv2.0 scaffold

orthogroup database (Wafula *et al.* 2022). Out of the 18,110 pre-constructed orthogroups,

proteins from all apple annotations (including 6 published scion cultivar genomes, 2 rootstock

genomes, and the 'WA 38' genome from this work) are found in 11,698 orthogroups. 'Golden

Delicious' Genome v1.0 (Velasco *et al.* 2010) was omitted from this analysis due to poor

annotation quality. Proteins from HapA and HapB of 'WA 38' were classified into 10,494 and

10,511 orthogroups, respectively, similar or slightly higher in number compared to previously

published *M. domestica* genomes, including 'Honeycrisp', 'Gala', and 'GDDH13' (Table 2,

Figure 5). An investigation into shared and unique orthogroups across all the scion genomes

showed that most orthogroups (8,800 or 75%) are shared by all six apple genomes considered.

Additionally, 824 orthogroups are shared by both 'WA 38' haplomes and the seven other

annotations (each of the two haplomes from 'Honeycrisp' and 'Antonovka 172670-B' are

counted as unique annotations). 'Honeycrisp' shared the largest number of orthogroups with

'WA 38', as expected due to being a parent of 'WA 38' (Supplemental Table S13). These results

indicate that the 'WA 38' annotation captures genes in virtually all *M. domestica* orthogroups.

Additionally, 39 orthogroups were unique to 'WA 38' (*i.e.* present only in the two 'WA 38'

haplomes) and each haplome of 'WA 38' contains 44 unique orthogroups (Figure 5).

Figure 5. Upset plot of shared and unique orthogroups among Malus domestica genomes.

Rows in the bottom of the figure are genomes used for the comparison. Columns (categories, x-

axis of the bar graph) are annotated with black or gray dots where black is present and gray is

absent. The height of the black bars (y-axis of the bar graph) is scaled to match the number of

orthogroup in each category, which are also printed above the bars.

In addition to identifying the shared and unique orthogroup, a CoRe OrthoGroup (CROG) -

Rosaceae analysis was performed to further investigate orthogroup contents. As expected, in

the CROG gene count clustermap (Figure 6), 'WA 38' clustered closely with 'Honeycrisp'. The

'WA 38' + 'Honeycrisp' group is clustered with 'GDDH13', as expected based on pedigree

(Howard *et al.* 2017). Interestingly, a strong 'publication bias', first mentioned by Wafula et al.,

2022, is observed: genomes released in the same publication or annotated by the same

researcher clustered together. Such groups are: 'Gala', *Malus sieversii*, and *M. sylvestris* (Sun

et al., 2021*)*; 'Fuji', 'M9', and 'MM106' (Li *et al*., 2024); *M. fusca* (Mansfeld *et al.* 2023) and

Pyrus communis 'd'Anjou' (Yocca *et al.* 2024); 'Honeycrisp' (Khan *et al.* 2022) and 'WA 38'. The

CROG gene count *z*-score box plot shows (Figure 7) that the average *z*-score of 'WA 38' gene

counts are slightly higher than expected (with 0 as the perfect score), indicating that there are a

number of CROGs containing more genes from the 'WA 38' annotations compared to other

apples.

Figure 6. CoRe OrthoGroup (CROG) - Rosaceae gene count clustermap. Each row represents a CROG and each column represents a genomes. Color indicates the number of genes in each cell relative to the row average (z-score). Warmer color indicates more genes. Cooler color indicates fewer genes. The darker a color, the closer the value is to the row average. Genome and annotation abbreviations can be found in Supplemental Table 1.

532　*Figure 7. Boxplot summarizing z-score distribution of CROG gene counts in selected pome fruit*

533　*genomes. Genome and annotation abbreviations can be found in Supplemental Table 1.*

Gene model evidence source mapping

The final gene model annotation contains *ab initio* prediction and genes with transcript evidence

and/or homologous protein support. Although high BUSCO completeness scores are obtained

from both haplome annotations, their gene numbers are greater than expected (45,000-49,000

based on previous publications). Therefore, we explored evidence supporting a gene model to

be a true positive, including extrinsic evidence (from transcript and homologous protein) used in

gene model annotation and comparative genomic evidence (EnTAP functional annotation and

gene family circumscription), and assessed completeness via a BUSCO analysis (Table 4). The

most stringent filter, the same strategy deployed in the 'Honeycrisp' genome annotation, was to

remove genes without full support from both transcript and homologous protein evidence

(Subset 1 in Table 4). This strategy removed ~10,000 genes from both haplomes and left

~43,000 genes in each annotation. Complete BUSCO score for this gene set decreased by ~1%

compared to the original full gene set. In the other three subsets (2-4) of genes, where less

stringent criteria were applied, ~3,000-4,000 genes were removed and complete BUSCO scores

maintained above 98%. In two of the subsets where the genes with functional and gene family

were taken into consideration (Subset 3 & 4), complete BUSCO scores remained the same as

the original gene set even after removing thousands of genes. CROG gene count analyses

were performed on the original full set, Subset 1 and Subset 3. The CROG gene count

clustermaps from the three gene sets showed highly similar clustering patterns (Figure 6 and

Supplemental Figure S8), indicating that removing genes did not alter the overall gene family

circumscription. The average CROG gene count z-score decreased from 0.330 in the original

full set, to 0.297 in Subset 3, and to 0.008 in Subset 1, indicating values closer to expectation as

more rigorous evidence categories are applied.

Table 4. Summary of genes mapped with various evidence source and completeness

assessments of those gene subsets.

	Number of genes		Complete BUSCO (%)	
	HapA	HapB	HapA	HapB
Original full set	53,028	54,235	98.5	98.4
Subset 1. Genes with full support*	43,079	43,590	97.5	97.6
Subset 2. Genes with any support*	49,829	50,861	98.2	98.2
Subset 3. Genes with full support + EnTAP & PT2	49,417	50,005	98.5	98.4
Subset 4. Genes with full support + EnTAP or PT2	50,087	50,743	98.5	98.4

* Full or any support from either RNA transcriptome or homologous protein evidence.

Plastid Genomes Assembly and Annotation

The chloroplast genome of the 'WA 38' apple is 159,915 bp in length, which is smaller than

most assembled *Malus* chloroplast genomes (Naizaier *et al.* 2019; Yan *et al.* 2019; Zhao *et al.*

2019; Ha *et al.* 2020; Miao *et al.* 2022; Li *et al.* 2022a). The plastome consisted of a typical

quadripartite structure with a pair of inverted repeat regions (IR) of the same length (26,352 bp)

separated by a long single copy region (LSC) (88,052 bp) and a short single copy region (SSC)

(19,159 bp). The IR regions and the SSC regions were all similar in length to that of other *Malus*

chloroplasts (Naizaier *et al.* 2019; Yan *et al.* 2019; Zhao *et al.* 2019; Ha *et al.* 2020; Miao *et al.*

2022; Li *et al.* 2022a)*. A total of 134 unique genes were annotated, including 86 protein-coding

genes, 42 tRNA genes, and 7 rRNA genes. Moreover, eight protein-coding genes (*ycf1, ycf2,*

rpl2, rpl23, ndhB, rps7, rps12, rps19-fragment), ten tRNA genes (*trnE-UUC, trnI-GAU, trnA-*

UGC, trnL-CAA, trnM-CAU, trnN-GUU, trnR-ACG, trnI-CAU, trnN-GUU, trnV-GAC), all four

rRNA genes (*rrn16, rrn23, rrn4.5, rrn5*) were located wholly within the IR regions (Figure 8).

Twelve protein-coding genes, eight tRNA genes, and one rRNA gene (*rrn16*) contain introns.

The majority of which contained one intron (19 genes), with only two genes (*pafl* and *clpP1*)

containing two introns.

The mitochondrial genome of the 'WA 38' apple is 451,423bp long and contains 64 annotated

genes. This annotation includes 4 rRNA genes (two copies of 26S, and one copy of both 18S

and 5S), 20 tRNA genes (including two copies of *trnaA-FME* and three copies of *trnaF-GAA*),

and 40 protein-coding genes (including two copies of *atp1, apt8, cox3, nad6, nad7, rnaseH,*

rps12, *rps3*, and *sdh4*). (Figure 9)

Figure 8. Chloroplast genome map of 'WA 38' with annotation. The outer circle shows the

locations of genes, colored according to their function and biological pathways as shown in the

figure legend. Forward-encoded genes are drawn on the outside of the circle, while reverse-

encoded genes are on the inside of the circle. The middle circle shows locations of the four

major sections of the chloroplast: LSC (long single copy), SSC (short single copy), IRA (inverted

repeat A), and IRB (inverted repeat B). The inner gray circle shows GC content across the

chloroplast genome.

Figure 9. Mitochondrial genome map of 'WA 38' with annotation. The outer circle shows the locations of genes, colored according to their function and biological pathways as shown in the figure legend. Forward-encoded genes are drawn on the outside of the circle, while reverse-encoded genes are on the inside of the circle. The inner gray circle shows GC content across the mitochondria genome.

Discussion

Genomes are essential resources for research communities. In order to provide accessible, hands-on training to the next generation of plant genome scientists, we engaged students in the construction of a genome for the 'WA 38' (Cosmic Crisp®) apple. Our guiding philosophy is

'inclusion and novelty', where we aim to build a high-quality reference genome that is useful to a

wide range of current and future research communities.

We emphasized assembly quality by leveraging our recent 'Honeycrisp' genome (Khan et al.

2022) to fully resolve haplotypes, *i.e.* the specific genetic contributions of each parent are known

and are represented in each respective haplome. As the first pome fruit genome to achieve this

level of resolution, the 'WA 38' genome provides a unique resource for researchers across

various fields to explore genome-scale genomic signatures that were previously unattainable for

pome fruit research. Examples include a more in-depth understanding of genetic variation and

inheritance, identification of alleles associated with specific traits (paving the way for allele

specific expression experiments), and opportunities to perform trait association analyses with

higher resolution (useful for breeding programs to identify new genetic markers linked to

desirable traits) (Talbot *et al.* 2024).

We also emphasized genome annotation quality, aiming to provide a hierarchy of hypothesized

gene models, where we compile a more complete list of putative genes, with increasingly

stringent evidence categories allowing users to access and use the appropriate set of

annotations for their application. By breaking from convention where a single stringency for

genome annotation has historically been set in published genomes, our approach provides an

annotation matrix that allows users to explore gene space as a function of annotation support.

Our original, full gene set contains ~54,000 putative gene models, almost 9,000 more than most

other *Malus* genomes (Supplemental Table S5). Subsequent filtering using various evidence

sources successfully adjusted the gene number closer to expected, although this resulted in

reduced completeness in some cases (Table 4). Subset 1, where only genes with full support

were selected, is the most stringent criteria we used for gene selection. Although the BUSCO

completeness score dropped by ~1%, it's still among the highest in *Malus* annotations and the

average CROG gene count z-score indicates that the overall number of genes in CROG are

very close to expectation (Supplemental Figure S8). However, a collection of 'cold' orthogroups

(containing fewer than expected number of genes compared to the rest annotations) emerged in

the 'Honeycrisp' plus 'WA 38' cluster from the CROG analysis (highlighted with a box in

Supplemental Figure S8). Since these cold spots were not observed in the original full gene set

nor the less rigorously filtered Subset 3, and are unique to the genomes annotated with the

same method and same filtering strategy, they are likely the result of a methodological bias.

This subset, Subset 1, is expected to contain fewer false positives at the cost of also dropping a

small amount of true positives; suitable for analysis that requires high-confidence gene models,

such as reconstructing species or pedigree relationship. Subset 3, which contains all genes

from Subset 1 and genes with both EnTAP and PlantTribes2 evidence, has a similar gene

number to the most recently published apple genomes, namely 'Honeycrisp', 'Fuji', 'M9', and

'MM106'. Subset 3 maintained the same BUSCO completeness score and did not have the

'cold' orthogroup observed in Subset 1. Thus, Subset 3 may contain more false positive genes,

but it also retains the most true positives; suitable for most analyses that can tolerate a small

amount of false positive gene models. Furthermore, similar to the 'Honeycrisp' plus 'WA 38'

cluster with shared unique 'cold' orthogroup zones in the Subset 1 CROG analysis, genomes

annotated by the same research group tend to exhibit similar gene count patterns (CROG

analysis - Figure 6), suggesting that methodological bias in a seemingly subjective analysis may

lead to a more similar gene landscape within those annotations. The most surprising examples

are the cluster of 'Gala' with the two wild *Malus* progenitors (*i.e.* different species), and the

cluster of *Malus fusca* with *Pyrus communis* 'd'Anjou' (*i.e.* different genera). In addition,

although most of the published *Malus* genome annotations have a similar number of genes

(~45,000, Supplemental Table S5), the CROG analysis identified different collections of

orthogroups with higher (warm color) or lower (cool color) than average gene counts across

clusters. These 'warm' and 'cool' orthogroup spots are not necessarily indicative of gene family

expansions or contractions (a separate analysis would be required), but does provide valuable

insight into the gene space within the context of lineage-specific genome annotations and

highlights potential areas for genome resource improvement. We believe the methodological

bias revealed by the CROG analysis should be addressed or acknowledged before further

analyses of gene family expansions and contractions in *Malus* is performed.

Throughout this project, we emphasized community engagement and enforce standardization of

genome resources. The AgBioData Genome Nomenclature working group is dedicated to

providing recommendations for consistent genome and gene model nomenclature that meets

the FAIR data principle (Wilkinson *et al.* 2016). We worked together with this working group and

the Rosaceae community genome database (Genome Database for Rosaceae, GDR, (Jung *et*

al. 2019)) to improve the existing nomenclature for Rosaceae genomes. The adoption of

standardized nomenclature for plant genomes represents a significant advancement in the field

of plant genomics as it helps reduce confusion and potential errors, thereby enhancing the

reliability and reproducibility of genomic research. In addition, we followed a previously-

established gene family classification protocol (Wafula *et al.* 2022; Khan *et al.* 2022) that

circumscribed genes into pre-computed orthogroups. Such a practice not only reduces

computational resource requirements, but also allows researchers to more easily compare

findings across studies. The uniformity, achieved by taking advantage of the already-existing

community resource, facilitates clearer communication, ensuring that discoveries are accurately

attributed and understood in the context of existing knowledge.

Our work emphasized the "reproducibility" of FAIR (Findable, Accessible, Interoperable and

Reproducible) data. All bioinformatics analyses follow some workflow whether it is manually

developed as work progresses by the researcher or is the product of an automated workflow

managed by software tools like Galaxy (The Galaxy Community, 2022) (graphical interface),

Nextflow (di Tommaso *et al*. 2017) or Snakemake (Mölder et al. 2021) (command-line interface).

Automated workflows create reproducible analyses because the version and parameters are

easily documented and software is commonly dockerized. For manually developed workflows,

the process is prone to being haphazard and disorganized and difficult to share. Thus, many

workflows are simply reduced to a brief description of software tools in Methods sections of

journal articles with software versions and important parameters often missing. As introduced in

the Results section, we provide a complete set of scripts and dockeried software to completely

recreate every analysis in the assembly and annotation of the WA 38 genome. The

organizational structure of the repository follows the Bioinformatics Notebook protocol

developed by our team (https://gitlab.com/ficklinlab-public/bioinformatics-notebook/). The goal of

this protocol is to ensure that complex manually executed workflows can be shared for

reproducibility, the format is readable by others and backups of critical data are supported.

Briefly, the directories are ordered using a numeric prefix indicating the order that analyses

should be performed. Inside each directory are sub-directories with smaller tasks. For each

task all relevant scripts and instructions are provided. All software used by the project is

dockerized and scripts contain the full parameter set used for every step. While there are areas

for improvement, the protocol, when followed, allows for easy sharing of the workflow via a Git

repository. In our view, this approach is a novel contribution towards FAIR data by ensuring that

non-automated workflows can be shared and are fully reproducible.

In addition to providing a fully reproducible workflow for the assembly of the 'WA 38' genome.

We generalized the scripts for any genome assembly and shared those as part of the three

ACTG GtiHub repositories mentioned in the Results setion. The new ACTG general workflow is

designed to provide training that is applicable for a wide range of species. The ACTG

repositories are a work in progress as we seek to create a generic, species-agnostic workflow

that will serve the broader American Campus Tree Genome (ACTG) community.

Availability of source code and requirements

Project name: 'WA 38' whole genome assembly and annotation

Project home page: https://gitlab.com/ficklinlab-public/wa-38-genome

Operating system(s): Platform independent

Programming language: bash, python, awk, perl

Other requirements: singularity, nextflow, java, python

License: Not applicable

Any restrictions to use by non-academics: No restrictions

RRID: Not applicable

Data Availability

711 Raw reads generated for this project are publicly available at NCBI under BioProject:

712 PRJNA1072127. Genome assembly and annotation are available on GDR:

713 https://www.rosaceae.org/Analysis/20220983

Competing interests

715 The author(s) declare that they have no competing interests.

Funding

717 This work was supported by the Washington Tree Fruit Research Commission (WTFRC) project

718 #AP-19-103 and USDA ARS internal appropriation funds.

Authors' contributions

720 L.H and S.F acquired funding for this project. All authors contributed to data analysis, data

721 interpretation, and manuscript writing.

References

Andrews. S, 2010 FastQC: A Quality Control Tool for High Throughput Sequence Data [Online]. Available online at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc/

NC State Extension, n.d. Apples - *Malus domestica*, North Carolina Extension Gardener Plant Toolbox. Retrieved May 23, 2024 [online] https://plants.ces.ncsu.edu/plants/malus-domestica/common-name/apples/

Brůna, T., K. J. Hoff, A. Lomsadze, M. Stanke, and M. Borodovsky, 2021 BRAKER2: automatic eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS supported by a protein database. NAR Genom Bioinform 3: lqaa108.

Chen C., Wu Y., Li J., Wang X., Zeng Z., Xu J., Liu Y., Feng J., Chen H., He Y., and Xia R., 2023 TBtools-II: A "one for all, all for one" bioinformatics platform for biological big-data mining. Mol. Plant 16: 1733–1742.

Cheng, H., G. T. Concepcion, X. Feng, H. Zhang, and H. Li, 2021 Haplotype-resolved de novo assembly using phased assembly graphs with hifiasm. Nat. Methods 18: 170–175.

Chen, X., S. Li, D. Zhang, M. Han, X. Jin *et al.*, 2019 Sequencing of a Wild Apple (*Malus baccata*) Genome Unravels the Differences Between Cultivated and Wild Apple Species Regarding Disease Resistance and Cold Tolerance. G3 9: 2051–2060.

Chen, S., Y. Zhou, Y. Chen, and J. Gu, 2018 fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics 34: i884–i890.

Choi, J. Y., L. R. Abdulkina, J. Yin, I. B. Chastukhina, J. T. Lovell *et al.*, 2021 Natural variation in plant telomere length is associated with flowering time. Plant Cell 33: 1118–1134.

Brown, M., P. M. González De la Rosa, and M. Blaxter, (2023). A Telomere Identification Toolkit (v0.2.41). Zenodo. https://doi.org/10.5281/zenodo.10091385

Crosby, J. A., J. Janick, P. C. Pecknold, J. C. Goffreda, and S. S. Korban, 1994 'Enterprise' Apple. HortScience 29: 825–826.

Daccord, N., J.-M. Celton, G. Linsmith, C. Becker, N. Choisne *et al.*, 2017 High-quality de novo assembly of the apple genome and methylome dynamics of early fruit development. Nat. Genet. 49: 1099–1106.

Danecek, P., J. K. Bonfield, J. Liddle, J. Marshall, V. Ohan *et al.*, 2021 Twelve years of SAMtools and BCFtools. Gigascience 10: giab008.

di Tommaso, P., M. Chatzou, E. W. Floden, P. P. Barja, E. Palumbo, and C. Notredame, 2017. Nextflow enables reproducible computational workflows. Nature Biotechnology, 35(4), 316–319.

Dierckxsens, N., P. Mardulyn, and G. Smits, 2017 NOVOPlasty: de novo assembly of organelle
genomes from whole genome data. Nucleic Acids Res. 45: e18.

Di Guardo, M., A. Tadiello, B. Farneti, G. Lorenz, D. Masuero *et al.*, 2013 A multidisciplinary
approach providing new insight into fruit flesh browning physiology in apple (*Malus* x
domestica Borkh.). PLoS One 8: e78004.

Dong, X., Z. Wang, L. Tian, Y. Zhang, D. Qi *et al.*, 2020 De novo assembly of a wild pear (*Pyrus*
betuleafolia) genome. Plant Biotechnol. J. 18: 581–595.

Doyle, J. J., and J. L. Doyle, 1987 A rapid DNA isolation procedure for small quantities of fresh
leaf tissue: RESEARCH.

Durand, N. C., M. S. Shamim, I. Machol, S. S. P. Rao, M. H. Huntley *et al.*, 2016 Juicer
Provides a One-Click System for Analyzing Loop-Resolution Hi-C Experiments. Cell Syst 3:
95–98.

Earl, D., K. Bradnam, J. St John, A. Darling, D. Lin *et al.*, 2011 Assemblathon 1: a competitive
assessment of de novo short read assembly methods. Genome Res. 21: 2224–2241.

Evans, K. M., B. H. Barritt, B. S. Konishi, L. J. Brutcher, and C. F. Ross, 2012 "WA 38" Apple.
HortScience 47: 1177–1179.

FGN, 2020 Global apple market reached $78B: set to continue moderate growth. Fruit
Growers News. Retrieved May 23, 2024 [online]
https://fruitgrowersnews.com/news/global-apple-market-reached-78m-set-to-continue-
moderate-growth/

Gabriel, L., K. J. Hoff, T. Brůna, M. Borodovsky, and M. Stanke, 2021 TSEBRA: transcript
selector for BRAKER. BMC Bioinformatics 22: 566.

Goremykin, V. V., P. J. Lockhart, R. Viola, and R. Velasco, 2012 The mitochondrial genome of
Malus domestica and the import-driven hypothesis of mitochondrial genome expansion in
seed plants. Plant J. 71: 615–626.

Grabherr, M. G., B. J. Haas, M. Yassour, J. Z. Levin, D. A. Thompson *et al.*, 2011 Full-length
transcriptome assembly from RNA-Seq data without a reference genome. Nat. Biotechnol.
29: 644–652.

Greiner, S., P. Lehwark, and R. Bock, 2019 OrganellarGenomeDRAW (OGDRAW) version
1.3.1: expanded toolkit for the graphical visualization of organellar genomes. Nucleic Acids
Res. 47: W59–W64.

Haas, B. J., A. L. Delcher, S. M. Mount, J. R. Wortman, R. K. Smith Jr *et al.*, 2003 Improving the
Arabidopsis genome annotation using maximal transcript alignment assemblies. Nucleic
Acids Res. 31: 5654–5666.

Hadish, J. A., T. D. Biggs, B. T. Shealy, M. R. Bender, C. B. McKnight *et al.*, 2022 GEMmaker:
process massive RNA-seq datasets on heterogeneous computational infrastructure. BMC
Bioinformatics 23: 1–11.
Ha, Y.-H., B. Maisupova, K. Choi, H.-J. Kim, D. Dosmanvetov *et al.*, 2020 Report on a complete
chloroplast genome sequence of wild apple tree, *Malus sieversii* (Lebed.) M. Roem.
Mitochondrial DNA B Resour. 5: 1504–1505.
Harkess, A., 2022 The American Campus Tree Genomes Documentation. Retrieved
May 23, 2024 [online] https://actg-wgaa.readthedocs.io/en/latest/index.html
Harris, R.S., 2007 Improved pairwise alignment of genomic DNA. Ph.D. Thesis, The
Pennsylvania State University. Retrieved May 23, 2024 [online]
https://www.bx.psu.edu/~rsharris/rsharris_phd_thesis_2007.pdf
Hart, A. J., S. Ginzburg, M. (sam) Xu, C. R. Fisher, N. Rahmatpour *et al.*, 2020 EnTAP: Bringing
faster and smarter functional annotation to non-model eukaryotic transcriptomes. Mol. Ecol.
Resour. 20: 591–604.
Hoff, K. J., S. Lange, A. Lomsadze, M. Borodovsky, and M. Stanke, 2016 BRAKER1:
Unsupervised RNA-Seq-Based Genome Annotation with GeneMark-ET and AUGUSTUS.
Bioinformatics 32: 767–769.
Howard, N. P., E. van de Weg, D. S. Bedford, C. P. Peace, S. Vanderzande *et al.*, 2017
Elucidation of the 'Honeycrisp' pedigree through haplotype analysis with a multi-family
integrated SNP linkage map and a large apple (*Malus×domestica*) pedigree-connected
SNP data set. Horticulture Research 4: 17003.
Huerta-Cepas, J., D. Szklarczyk, D. Heller, A. Hernández-Plaza, S. K. Forslund *et al.*, 2019
eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated orthology resource
based on 5090 organisms and 2502 viruses. Nucleic Acids Res. 47: D309–D314.
Johnston, J. W., and P. Brookfield, 2012 Delivering postharvest handling protocols for apples
and pears faster: Integrating "omics" and physiology approaches. Acta Hortic. 23–28.
Jung, S., T. Lee, C.-H. Cheng, K. Buble, P. Zheng *et al.*, 2019 15 years of GDR: New data and
functionality in the Genome Database for Rosaceae. Nucleic Acids Res. 47: D1137–D1145.
Khan, A., S. B. Carey, A. Serrano, H. Zhang, H. Hargarten *et al.*, 2022 A phased, chromosome-
scale genome of "Honeycrisp" apple (*Malus domestica*). GigaByte 2022: gigabyte69.
Kurtz, S., A. Phillippy, A. L. Delcher, M. Smoot, M. Shumway *et al.*, 2004 Versatile and open
software for comparing large genomes. Genome Biol. 5: R12.
Kuznetsov, D., F. Tegenfeldt, M. Manni, M. Seppey, M. Berkeley *et al.*, 2022 OrthoDB v11:
annotation of orthologs in the widest sampling of organismal diversity. Nucleic Acids Res.

51: D445–D451.

Li, X., Z. Ding, H. Miao, J. Bao, and X. Tian, 2022a Complete chloroplast genome studies of
different apple varieties indicated the origin of modern cultivated apples from and. PeerJ
10: e13107.

Li, H., and R. Durbin, 2009 Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics 25: 1754–1760.

Liebhard, R., M. Kellerhals, W. Pfammatter, M. Jertmini, and C. Gessler, 2003 Mapping
quantitative physiological traits in apple (*Malus* x *domestica* Borkh.). Plant Mol. Biol. 52:
511–526.

Li, X., L. Kui, J. Zhang, Y. Xie, L. Wang *et al.*, 2016 Improved hybrid de novo genome assembly
of domesticated apple (*Malus* x *domestica*). Gigascience 5: 35.

Li, W., C .Chu, H. Li et al., 2024 Near-gapless and haplotype-resolved apple genomes
provide insights into the genetic basis of rootstock-induced dwarfing. Nat Genet 56, 505–
516.

Li, W., J. Liu, H. Zhang, Z. Liu, Y. Wang *et al.*, 2022b Plant pan-genomics: recent advances,
new challenges, and roads ahead. J. Genet. Genomics 49: 833–846.

Li, Y., M. Pi, Q. Gao, Z. Liu, and C. Kang, 2019 Updated annotation of the wild strawberry
Fragaria vesca V4 genome. Horticulture Research 6: 1–9.

Lovell, J. T., A. Sreedasyam, M. Eric Schranz, M. Wilson, J. W. Carlson *et al.*, 2022
GENESPACE tracks regions of interest and gene copy number variation across multiple
genomes.

Lum, G. B., B. J. Shelp, J. R. DeEll, and G. G. Bozzo, 2016 Oxidative metabolism is associated
with physiological disorders in fruits stored under multiple environmental stresses. Plant
Sci. 245: 143–152.

Manni, M., M. R. Berkeley, M. Seppey, F. A. Simão, and E. M. Zdobnov, 2021 BUSCO Update:
Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage
for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. Mol. Biol. Evol. 38: 4647–4654.

Mansfeld, B. N., A. Yocca, S. Ou, A. Harkess, E. Burchard *et al.*, 2023 A haplotype resolved
chromosome-scale assembly of North American wild apple *Malus fusca* and comparative
genomics of the fire blight Mfu10 locus. Plant J. 116: 989–1002.

Marçais, G., and C. Kingsford, 2011 A fast, lock-free approach for efficient parallel counting of
occurrences of k-mers. Bioinformatics 27: 764–770.

Mendoza, M., Hanrahan, I., & Bolaños, G., 2022 2022 Update: Additional WA 38
harvest and storage considerations. Retrieved May 23, 2024 [online]

https://treefruit.wsu.edu/article/2022-update-additional-wa-38-harvest-and-storage-

consideration

Miao, H., J. Bao, X. Li, Z. Ding, and X. Tian, 2022 Comparative analyses of chloroplast

genomes in "Red Fuji" apples: low rate of chloroplast genome mutations. PeerJ 10:

e12927.

Mölder, F., K. P. Jablonski, B. Letcher, M. B. Hall, C. H. Tomkins-Tinch *et al.* 2021

Sustainable data analysis with Snakemake [version 2; peer review: 2 approved].

F1000Research, 10:33

Naizaier, R., Z. Qu, S. Wu, and X. Tian, 2019 The complete chloroplast genome of Malus

sieversii (Rosaceae), a wild apple tree in Xinjiang, China. Mitochondrial DNA B Resour. 4:

983–984.

Nattestad, M., and M. C. Schatz, 2016 Assemblytics: a web analytics tool for the detection of

variants from an assembly. Bioinformatics 32: 3021–3023.

NCBI Organelle genome resources, n.d. Retrieved May 23, 2024 [online]

https://www.ncbi.nlm.nih.gov/genome/organelle/

O'Leary, N. A., M. W. Wright, J. R. Brister, S. Ciufo, D. Haddad *et al.*, 2016 Reference

sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional

annotation. Nucleic Acids Res. 44: D733–45.

Ou, S., W. Su, Y. Liao, K. Chougule, J. R. A. Agda *et al.*, 2019 Benchmarking transposable

element annotation methods for creation of a streamlined, comprehensive pipeline.

Genome Biol. 20: 1–18.

Pareek, S. T. de F. S., 2019 Postharvest physiological disorders in fruits and vegetables (S.

Tonetto de Freitas & S. Pareek, Eds.). CRC Press, Boca Raton : Taylor & Francis, 2018.

phasegenomics, n.d. Hic_QC: a (very) simple script to QC Hi-C data. GitHub.

https://github.com/phasegenomics/hic_qc

Quinlan, A. R., and I. M. Hall, 2010 BEDTools: a flexible suite of utilities for comparing genomic

features. Bioinformatics 26: 841–842.

Raymond, O., J. Gouzy, J. Just, H. Badouin, M. Verdenaud *et al.*, 2018 The *Rosa* genome

provides new insights into the domestication of modern roses. Nat. Genet. 50: 772–777.

Rhie, A., Walenz, B.P., Koren, S. et al., 2020 Merqury: reference-free quality,

completeness, and phasing assessment for genome assemblies. Genome Biol 21, 245.

Robinson, J. T., H. Thorvaldsdóttir, W. Winckler, M. Guttman, E. S. Lander *et al.*, 2011

Integrative genomics viewer. Nat. Biotechnol. 29: 24–26.

Robinson, J. T., D. Turner, N. C. Durand, H. Thorvaldsdóttir, J. P. Mesirov *et al.*, 2018

Juicebox.js Provides a Cloud-Based Visualization System for Hi-C Data. Cell Syst 6: 256–258.e1.

Sallato, B., M. D. Whiting, and J. Munguia, 2021 Rootstock and nutrient imbalance leads to "Green Spot" development in 'WA 38' apples. HortScience 56: 1542–1548.

Serra, S., Goke, A., Sheick, R., Mendoza, M., Schmidt, T., Hanrahan, I., Ross, C., & Musacchi, S., 2023 Effects of harvest timing on maturity, fruit quality, and consumer acceptance of 'WA 38' apples. *Acta Horticulturae*, *1366*, 61–68.

Sharman, S., n.d. Learning from the trees: American Campus Tree Genomes Project pushes equality in genomic science. Retrieved May 23, 2024 [online] https://hudsonalpha.shorthandstories.com/learning-from-the-trees/

Sheick, R., S. Serra, S. Musacchi, and D. Rudell, 2023 Metabolic fingerprint of 'WA 38' green spot symptoms reveals increased production of epicuticular metabolites by parenchyma. Sci. Hortic. 321: 112257.

Sheick, R., S. Serra, D. Rudell, and S. Musacchi, 2022 Investigations of multiple approaches to reduce Green Spot incidence in 'WA 38' apple. Agronomy (Basel) 12: 2822.

Shirasawa, K., A. Itai, and S. Isobe, 2021 Chromosome-scale genome assembly of Japanese pear (*Pyrus pyrifolia*) variety 'Nijisseiki' DNA Res. 28:dsab001

Smit, AFA, Hubley, R & Green, P., n.d. RepeatMasker Open-4.0.2013-2015 Retrieved May 23, 2024 [online] http://www.repeatmasker.org

Sun, X., C. Jiao, H. Schwaninger, C. T. Chao, Y. Ma *et al.*, 2020 Phased diploid genome assemblies and pan-genomes provide insights into the genetic history of apple domestication. Nat. Genet. 52: 1423–1432.

Su, Y., X. Yang, Y. Wang, J. Li, Q. Long *et al.*, 2024 Phased Telomere-to-Telomere Reference Genome and Pangenome Reveal an Expansion of Resistance Genes during Apple Domestication. Plant Physiol. kiae258

Talbot, S. C., K. J. Vining, J. W. Snelling, J. Clevenger, and S. A. Mehlenbacher, 2024 A haplotype-resolved chromosome-level assembly and annotation of European hazelnut (C. avellana cv. Jefferson) provides insight into mechanisms of eastern filbert blight resistance. G3. jkae021

The Galaxy Community, The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2022 update. Nucleic Acids Research. 50: W345–W351.

Tillich, M., P. Lehwark, T. Pellizzer, E. S. Ulbricht-Jones, A. Fischer *et al.*, 2017 GeSeq - versatile and accurate annotation of organelle genomes. Nucleic Acids Res. 45: W6–W11.

trinityrnaseq, n.d. get_longest_isoform_seq_per_trinity_gene.pl. GitHub. Retrieved May

23, 2024 [online]

https://github.com/trinityrnaseq/trinityrnaseq/blob/master/util/misc/get_longest_isoform_s

eq_per_trinity_gene.pl

Truscott, S., 2023 WSU's Cosmic Crisp® joins top 10 bestselling U.S. apple varieties.

CAHNRS News Washington State University. Retrieved May 23, 2024 [online]

https://news.cahnrs.wsu.edu/article/wsus-cosmic-crisp-joins-top-10-bestselling-u-s-

apple-varieties/

Uliano-Silva, M., J. G. R. N. Ferreira, K. Krasheninnikova, G. Formenti, L. Abueg *et al.*, 2023

MitoHiFi: a python pipeline for mitochondrial genome assembly from PacBio high fidelity

reads. BMC Bioinformatics 24: 1–13.

USApple, 2024 Industry at a glance Retrieved May 23, 2024 [online],

https://usapple.org/industry-at-a-glance

VanBuren, R., C. M. Wai, M. Colle, J. Wang, S. Sullivan *et al.*, 2018 A near complete,

chromosome-scale assembly of the black raspberry (*Rubus occidentalis*) genome.

Gigascience 7.:

Velasco, R., A. Zharkikh, J. Affourtit, A. Dhingra, A. Cestaro *et al.*, 2010 The genome of the

domesticated apple (*Malus × domestica* Borkh.). Nat. Genet. 42: 833–839.

Verde, I., J. Jenkins, L. Dondini, S. Micali, G. Pagliarani *et al.*, 2017 The Peach v2.0 release:

high-resolution linkage mapping and deep resequencing improve chromosome-scale

assembly and contiguity. BMC Genomics 18: 225.

Vurture, G. W., F. J. Sedlazeck, M. Nattestad, C. J. Underwood, H. Fang *et al.*, 2017

GenomeScope: fast reference-free genome profiling from short reads. Bioinformatics 33:

2202–2204.

Wafula, E. K., H. Zhang, G. Von Kuster, J. H. Leebens-Mack, L. A. Honaas *et al.*, 2022

PlantTribes2: Tools for comparative gene family analysis in plant genomics. Front. Plant

Sci. 13: 1011199.

Washington Apple Commission, 2021 Did you know? Apple Facts. Retrieved May 23,

2024 [online] https://waapple.org/did-you-know/

Wilkinson, M. D., M. Dumontier, I. J. J. Aalbersberg, G. Appleton, M. Axton *et al.*, 2016 The

FAIR Guiding Principles for scientific data management and stewardship. Sci Data 3:

160018.

Wood, D. E., and S. L. Salzberg., 2014 Kraken: ultrafast metagenomic sequence classification

using exact alignments. Genome Biol 15, R46

Yan, M., X. Zhao, J. Zhou, Y. Huo, Y. Ding *et al.*, 2019 The complete chloroplast genome of

959  cultivated apple (*Malus domestica* Cv. "Yantai Fuji 8"). Mitochondrial DNA B Resour. 4:

960   1213–1216.

961 Yocca, A., M. Akinyuwa, N. Bailey, B. Cliver, H. Estes *et al.*, 2024 A chromosome-scale

962   assembly for 'd'Anjou' pear. G3 14.3, jkae003

963 Zhang, H., E. K. Wafula, J. Eilers, A. E. Harkess, P. E. Ralph *et al.*, 2022 Building a foundation

964   for gene family analysis in Rosaceae genomes with a novel workflow: A case study in

965   Pyrus architecture genes. Front. Plant Sci. 13: 975942.

966 Zhao, X., M. Yan, Y. Ding, X. Chen, and Z. Yuan, 2019 The complete chloroplast genome of

967   apple rootstock 'M9.' Mitochondrial DNA B Resour 4: 2187–2188.

968 Zhou, C., S. A. McCarthy, and R. Durbin, 2022 YaHS: yet another Hi-C scaffolding tool.

969   Bioinformatics 39: btac808.