1

Feedforward circuits enable modality-specific control of cortical inhibition during

behavioral state transitions.

Marcel de Brito Van Velze[1], Dhanasak Dhanasobhon[1], Marie Martinez[1], Annunziato Morabito[1],

Emmanuelle Berthaux[1], Cibele Martins Pinho[1], Yann Zerlaut[1#] and Nelson Rebola[1#]

[1]ICM, Paris Brain Institute, Hôpital de la Pitié-Salpêtrière, Sorbonne Université, Inserm, CNRS,

Paris, 75013, France.

#These authors contributed equally.

Number of figures: 7; Supplementary figures: 4; Supplementary tables: 0

Keywords: Neocortical interneurons, Somatostatin positive Interneurons, Somatosensory cortex,

Spontaneous behaviours, Locomotion, Whisking.

Acknowledgements:

This study was supported by the Centre National de la Recherche Scientifique, Agence Nationale

de la Recherche (grants ANR-EXCIGLY and ANR-DecoSensoMol1) the European Research

Council (ERC-STG- 678250 to N.R.), the European Union's Horizon 2020 research and

innovation programme (Marie Skłodowska-Curie Grant 892175 *"InProsMod"* to Y.Z.); by the Paris

Brain Institute, and the Fondation pour la Recherche Médicale (FRM; fellowship

ARF201909009117 to Y.Z.). We thank the Paris Brain Institute core facilities, namely iVector and

ICMice PHENOPARC. We would also like to thank Dr. Alberto Bacci (ICM, Paris), Dr. Joana

Lourenco (ICM, Paris) and Dr. Nicolas Chenouard (ICM, Paris) for the valuable discussions and

comments on the manuscript.

34
35

SUMMARY (236 words)

Active behavior strongly influences sensory information processing in primary sensory cortices. Local interneurons, especially those involved in disinhibitory circuits like somatostatin (SST) and vasoactive intestinal peptide (VIP)-positive interneurons, play a crucial role in regulating cortical function during behavior. While numerous studies have focused on the recruitment of pyramidal neurons, the intricate network mechanisms governing the behavior-state-dependent modulation of inhibitory neurons remain unclear. Using intravital two-photon calcium imaging we have studied the neuronal pathways that regulate SST- and VIP-IN activity in sensory cortices during spontaneous behaviours. We now reveal that active behavioral states associated with increased locomotor activity strongly recruit L2/3 SST-INs in primary somatosensory (S1) but not visual (V1) cortex indicating modality-specific mechanisms control activity of cortical interneurons during active states. Such a difference between V1 and S1 disappears in the presence of visual sensory drive (ambient light) suggesting a critical involvement of feedforward sensory pathways in the positive modulation of SST-INs during spontaneous behaviors. In agreement, we now unveil that inactivating the somatosensory thalamus significantly reduces the recruitment of SST-INs in S1 by locomotion. In contrast, interfering with disinhibitory circuits via chemogenetic inactivation of VIP-INs further increased the sensitivity of SST-INs to variations in behavioural states. Our work thus reveals a previous unknown role for thalamic inputs in driving SST-INs, suggesting that by integrating feedforward activity with neuromodulation, SST-INs play a central role in adapting sensory processing to behavioral states in a sensory modality-specific manner.

Introduction:

In the mammalian brain, the cerebral cortex is responsible for complex cognitive functions, including a coherent representation of the external world that is crucial for an effective interaction with a constantly changing environment. Importantly, sensation can occur under drastically different behavior conditions, which have been reported to result in various perceptual outcomes (Busse et al., 2017; McGinley et al., 2015a). A hallmark of sensory systems is thus to provide a flexible strategy to process incoming information in a context-dependent manner (McCormick et al., 2020).

Local neocortical interneurons, in particular those involved in disinhibitory circuits, like vasoactive intestinal peptide (VIP) and somatostatin (SST) expressing interneurons (INs), have been shown to be particularly sensitive to behavior state transitions and to be key contributors to the state-dependent regulation of cortical function (Fu et al., 2015, 2014; Gasselin et al., 2021; Gentet et al., 2012; Muñoz et al., 2017; Polack et al., 2013; Reimer et al., 2014). In V1, recruitment of VIP-INs and consequent decrease in SST-IN mediated inhibition has been suggested to underlie the gain modulation of visual-evoked responses observed during locomotion (Fu et al., 2014). Yet, the mechanisms controlling the engagement of SST-INs across active behavioral states are unclear. In addition, behavioral modulations have been shown to differentially impact pyramidal cell activity across sensory cortices (Ayaz et al., 2013; Niell and Stryker, 2010; Polack et al., 2013; Saleem et al., 2013; Schneider et al., 2014; Shimaoka et al., 2018; Vinck et al., 2015; Zhou et al., 2014). However, whether inhibitory circuits display similar or variable sensitivity to behavioral state transitions across sensory cortical areas remains unexplored.

A recent study revealed that modulation of INs activity in V1 by brain states can be largely predicted by the first transcriptomic principal component (Bugeon et al., 2022). This transcriptomic component suggested that the increase in SST-INs activity, observed in alert states, is associated with cholinergic signalling (Bugeon et al., 2022). However, this finding contrasts with a previous report that SST-INs in V1 lose cholinergic modulation during development (Yaeger et al., 2019).

4

Paradoxically, VIP-mediated disinhibition has been proposed to explain the increase in SST-INs
activity during active behavioral states but this hypothesis lacks experimental evidence (Garcia
Del Molino et al., 2017). Both hypotheses point to neuromodulatory inputs, in particular cholinergic
axonal projections to be at the origin of the sensitivity of VIP- and SST-INs to behavioral state
transitions. Yet, recent results highlighted that thalamic activity is essential for the large and
sustained depolarizations associated with arousal/movement-related signals in L2/3 PNs in V1
(Nestvogel and McCormick, 2022). These results suggest that behavior states can modulate
neocortical activity not only through neuromodulatory projections such as cholinergic and
noradrenergic pathways, but also via glutamatergic thalamic inputs. As SST-INs receive
pronounced inputs from local L2/3 PNs it is therefore possible that feedforward activity transmitted
via the thalamus participates in the behavioral-dependent modulation of SST-INs in primary
sensory regions. Still, as glutamatergic synapses from L2/3 PNs into SST-INs have relatively low
release probabilities it is unclear whether an increase in L2/3 PNs activity during active states is
sufficient to drive SST-INs. Altogether, the specific neuronal mechanisms governing the activity
of SST-INs during spontaneous behaviors are still not fully understood.

The modulation of disinhibitory circuits by active behavioral states has been the focus of
considerable scrutiny in V1 (Bugeon et al., 2022; Dipoppa et al., 2018; Fu et al., 2014; Pakan et
al., 2016; Polack et al., 2013; Reimer et al., 2014). Locomotion, in particular, has been found to
influence the activity of VIP- and SST-INs in a context-dependent manner, with the modulation of
SST-INs being contingent on the presence or absence of ambient light (Dipoppa et al., 2018;
Pakan et al., 2016). Whether such modulation by locomotion extends to other sensory cortices
remains unclear. In S1, behavioral periods associated with increased whisking (in stationary
animals) have been reported to increase L2/3 VIP- activity with concomitant decrease in L2/3
SST-INs firing in agreement with the presence of disinhibitory circuits (Gasselin et al., 2021;
Gentet et al., 2012; Lee et al., 2013; Muñoz et al., 2017; Yu et al., 2019). Nevertheless, in S1,
and in stark contrast to V1, our knowledge regarding the impact of locomotion on the operation of

SST- and VIP-INs remains limited. This becomes especially relevant considering that, during

navigation, mice typically employ locomotion in conjunction with whisking to explore their

surroundings.

Here we used intravital two-photon calcium imaging in S1 and V1 of awake, head-fixed

mice, to investigate how transitions in behavioral states (namely whisking and locomotion)

modulate the activity of SST- and VIP-INs, key disinhibitory elements of cortical networks. We

found that in darkness, the recruitment of SST-INs by locomotion was significantly higher in S1

than V1. We also observed that the increase of SST-INs activity in S1 occurred despite the strong

recruitment of VIP-INs, a known inhibitor of SST-INs. In addition, neither the strength of inhibition

provided by VIP-INs nor the sensitivity of SST-INs to cholinergic signaling was different across

sensory regions. Importantly, inactivation of thalamic function using muscimol microinjections

limited the increase in activity of SST-INs by locomotion. These results reveal that behavior state-

transitions, often associated with changes in arousal, induce sensory modality specific changes

to the activity of inhibitory circuits, most likely through inputs coming from the sensory thalamus.

By integrating feedforward thalamic activity transmitted indirectly via L2/3 PNs as well as

neuromodulatory information coming from VIP-mediated inhibition and cholinergic innervation,

SST-INs occupy a pivotal position in cortical networks by adapting sensory processing to

behavioral states in a sensory modality-specific manner.

Results:

Activity of L2/3 SST-INs in S1 and V1 during spontaneous behavior

We initially focused our attention on the *vivo* response of SST-INs to behavioral state transitions in the mouse primary somatosensory cortex (S1). In fact, multiple studies have studied the activity of disinhibitory circuits during active behavioral states in V1. However, it remains unclear whether inhibitory circuits display similar or variable sensitivity to behavior state transitions in other sensory cortical areas including S1. To monitor alterations in firing dynamics of SST-INs we expressed the genetically encoded calcium indicator GCaMP6s selectively in SST-IN using transgenic mice (SST-cre) (Figure 1A) and changes in somatic fluorescence of GCaMP6s were used as a proxy for spiking activity. Fluorescence signals were recorded using two-photon imaging in awake head fixed mice that spontaneously transitioned between rest and locomoting periods on a circular treadmill (Figure 1A). Whisking activity was simultaneously acquired using a video camera focused on the mouse snout (Figure 1A). This experimental setup allowed us to study how SST-INs activity varied across active behavioral states namely during whisking with and without associated running.

In our initial analysis, we categorized recording periods based on whether animals were engaged in whisking, locomotion, or neither. Periods during which whisking or locomotion were not detected were classified as resting (see Methods). Through the quantification of average GCaMP6s fluorescence levels we found that in S1, SST-INs displayed markedly higher activity levels during periods of locomotion compared to resting ($\Delta F/F_{0\ rest}$ = 0.80 ± 0.02; $\Delta F/F_{0\ run}$ = 1.84 ± 0.06; n = 2030 cells in 10 mice, p<0.0001, Mann-Whitney U test, Figure 1A-C). This engagement of SST-INs during locomotion in S1 was also observed when we investigated the variability in the response to behavior state transitions at the single cell level. For that we quantified the correlation between baseline activity and running speed for individual SST-INs (Figure 1D). We observed a predominantly positive correlation coefficient across SST-INs in S1 (Pearson correlation, ρ = 0.22 ± 0.01, n = 2030, Figure 1D). Specifically, 59% of SST-INs exhibited a statistically significant (see

methods) positive modulation in response to locomotion, while only a smaller but not negligible

fraction of SST-INs displayed a significant negative modulation (20%) (Figure 1E)

These results contrasted with previous reports in V1, where in darkness, locomotion had

on average, no impact in SST-INs activity (Dipoppa et al., 2018; Pakan et al., 2016). In order to

test if this functional difference was linked to our experimental conditions or if in fact reflected a

previously unknown region-specific modulation of SST-INs by locomotion we repeated the

experiments listed above but now monitoring SST-INs activity in V1 (Figure 1E). In agreement

with previous studies we observed that the modulation of SST-INs by locomotion in the dark in

V1 was quite modest ($\Delta F/F_{0\ rest}$ = 0.70 ± 0.03; $\Delta F/F_{0\ run}$ = 0.84 ± 0.03; n = 881 cells in 10 mice,

p<0.0001, Mann-Whitney U; (Dipoppa et al., 2018; Pakan et al., 2016) (Figure 1E,F) contrasting

the results obtained in S1. This difference in activity of SST-INs during locomotion between S1

and V1 was also observed at the single cell level. In contrast to S1, our observations in V1

revealed a comparable ratio of SST-INs exhibiting positive (33%) and negative (25%) correlations

between running speed and spontaneous activity (Figure 1G).

In order to better estimate the rate of change in calcium levels in SST-INs between S1

and V1 we generated population averages of all running or whisking bouts aligned to locomotion

or whisking onset respectively (Figure 1H, I). In these population averages it was possible to

observe that both locomotion as well as whisking induced changes in $\Delta F/F0$ in SST-INs that were

significantly larger in S1 than in V1 (Figure 1J, I). The differential sensitivity of SST-INs between

the two sensory areas to behavioral state transitions was particularly evident when comparing the

cross-correlation curves between spontaneous activity of SST-INs and the locomotion or whisking

traces (Pearson correlation: locomotion, S1, ρ = 0.20 ± 0.01, n=2030; V1, ρ = 0.05 ± 0.01, n=633;

whisking, S1, ρ = 0.15 ± 0.01, n=853; V1, ρ = 0.04 ± 0.01, n=633, Figure 1K). Altogether these

results suggest that in the neocortex, behavior-dependent modulation of SST-INs activity is not

uniform across sensory regions.

Contribution of whisking and locomotion to SST-INs modulation in S1

Our results indicate that in S1 active behaviors like whisking and locomotion, result in

increased activity of SST-INs. However locomotion in mice is closely linked to the motion of their

whiskers (Mitchinson et al., 2007). In contrast, whisking can take place in both run and rest

conditions. We thus decided to further investigate how the two different active states (whisking

and locomotion) impacted the activity of SST-INs. In our experimental conditions mice were

observed to spend an average of 32% of their time running, 47% whisking, and within this

whisking period, 16% did not involve associated ambulatory activity (see Methods) (Figure 2A-B).

To facilitate a more straightforward comparison on how whisking and locomotion shaped activity

in SST-INs in S1 we specifically isolated recording periods in which mice were engaged in active

whisking without simultaneous running (Figure 2C-E). Under these conditions, which we defined

as "whisking only," the activity of SST-INs exhibited distinct differences compared to whisking

episodes that were accompanied by locomotion. Specifically, SST-INs were not positively

modulated during active whisking showing in contrast a tendency towards inhibition in agreement

with previous works (Gasselin et al., 2021; Gentet et al., 2012; Lee et al., 2013; Muñoz et al.,

2017) (Figure 2C-E). To further quantify how the different behavioral states influenced SST-INs

activity we separated calcium activity and we defined whisking only modulation index (WOMI) and

whisking modulation index (WMI) similar to our previous LMI calculation (see Methods). We

observed that both WMI and LMI were clearly positive and rather similar in agreement with the

observation that in mice most whisking bouts are actually associated with locomotion (Sofroniew

et al., 2014). In contrast WOMI, was only slightly positive with a large percentage of cells

displaying negative modulation (Figure 2F).

During our imaging sessions, we observed significant variation among mice in terms of

the duration of whisking without concurrent locomotion (% time WO = 15.6 ± 1.7%, CV = 105%,

n = 86 sessions). In light of this observation, we embarked on investigating whether the response

profile of SST-INs differed between sessions characterized by high or low percentages of WO.

Our rationale was that sessions with a higher proportion of WO activity would provide a better

opportunity to sample and visualize the specific impact of whisking with and without associated

locomotion. To achieve this, we employed an arbitrary threshold of 20% to categorize imaging

sessions into those with high or low percentages of WO (Figure 2G, H). We subsequently

identified all whisking bouts and averaged corresponding variations in GCaMP6s fluorescence

across the two different groups of sessions aligned to whisking onset. Upon examination of the

population's average response, distinct variations in the response of SST-INs to whisking became

evident between the two types of sessions (Figure 2G, H). In sessions with a high percentage of

whisking only, whisking onset was associated with a clear decrease in the activity of SST-INs

immediately after whisking onset. In contrast, in sessions where mice spent only a small fraction

of their time whisking without locomotion, whisking onset was clearly associated with an increase

in SST-INs firing (Figure 2G-H). Overall, these results indicate that in S1, whisking at rest induces

a decrease in SST-INs activity, in agreement with previous reports (Gentet et al., 2012; Muñoz et

al., 2017; Yu et al., 2019). On the other hand, when mice engage in active behavioral states

involving locomotion, activity of somatostatin interneurons (SST-INs) is increased, specifically in

the primary somatosensory cortex (S1), but not in the primary visual cortex (V1; Fig. 1). These

findings suggest that distinct active behavioral states exert varying effects on the activity of S1

SST-INs.

Morphoelectric and transcriptomic analysis have highlighted that SST-INs, even within

L2/3, are a highly heterogeneous neuronal population (Gouwens et al., 2020; Wu et al., 2023).

We probed if sensitivity to locomotion was also heterogeneous across individual SST-INs in S1.

To evaluate this, we separated SST-INs according to their response to WOMI. We observed that

SST-INs that displayed a negative WOMI, were on average less sensitive to locomotion (Figure

2I). In contrast, SST-INs with positive WOMI mostly increase their activity with locomotion (Figure

2I). Variable modulation of SST-INs for locomotion was observed even within single fields of view,

suggesting that variability in SST-IN responses cannot be attributed to distinct behavioral states

(Figure S1). In summary, these findings indicate the presence of various subtypes of SST-INs in

Layer 2/3, exhibiting varying degrees of sensitivity to transitions in behavioral states.

VIP-INs are equally engaged by locomotion in V1 and S1

Having observed that in S1, SST-INs are recruited during active behavioral states,

specifically during locomotion, we proceeded to investigate the underlying neuronal mechanisms

involved in such modulation. Taking into account previous works we identified three main potential

pathways that could participate in controlling L2/3 SST-INs activity during spontaneous behaviors:

(i) inhibitory top-down modulation via VIP-INs (Fu et al., 2014; Lee et al., 2013; Pfeffer et al., 2013;

Pi et al., 2013) (ii) increased local recurrent excitatory L2/3 PNs activity (Nestvogel and

McCormick, 2022; Pakan et al., 2016) and (iii) direct excitatory neuromodulation (*e.g.* via direct

cholinergic inputs) (Yaeger et al., 2019). We initially focused on VIP-INs, which are known to be

important inhibitors of SST-IN function (Fu et al., 2014; Lee et al., 2013; Pfeffer et al., 2013; Pi et

al., 2013) and are strong recipients of neuromodulatory inputs (*e.g.* cholinergic) associated with

active behavioral states (Alitto and Dan, 2013; Gasselin et al., 2021). Additionally, theoretical work

suggested that VIP-INs activation in certain conditions can result in unexpected increase in SST-

INs function through disinhibition of local PNs (Garcia Del Molino et al., 2017). Therefore, we

studied the contribution of VIP-INs to the activity patterns of SST-INs during behavioral state

transitions. We also explored the possibility that differential activity/function of VIP-INs could be

at the origin of the distinct sensitivity of SST-INs in V1 and S1 to locomotion.

We started by examining how activity of VIP-INs varied in S1 during the same behavioral

transitions as we described for SST-INs (Figure 3A-C). In line with previous findings (Fu et al.,

2015, 2014), we observed an increase in activity of VIP-INs during ambulatory periods in S1

($\Delta F/F_{0\ rest}$ = 0.79 ± 0.02; $\Delta F/F_{0\ run}$ = 2.23 ± 0.05; n = 1915 cells; in 10 mice, p<0.0001, Mann-

Whitney U, Figure 3A, B, D) with the majority of VIP-INs displaying a positive modulation by

locomotion (LMI = 0.38 ± 0.01; Figure 3E). A similar observation was obtained while estimating

Pearson correlation between GCaMP6f or GCaMP6s fluorescence and running speed. VIP-INs

displayed on average a positive correlation (ρ = 0.30 ± 0.01, Figure 3F) with only a very small

fraction (6%) of VIP-INs displaying a statistically significant decrease in activity (Figure 3F) during

locomotion. It is worth mentioning that the percentage of VIP-INs displaying negative modulation

by locomotion was significantly less than SST-INs (see Figure 1). Despite the similar average

positive modulation by locomotion between VIP- and SST-INs in S1, we observe that at the

population level, VIP-INs displayed a higher sensitivity to behavioral state transitions than SST-

INs (Figure S2). This was reflected into a higher correlation coefficient between spontaneous

activity and running speed (VIP: ρ = 0.30 ± 0.01 vs SST: 0.22± 0.01, p < 0.001, Mann-Whitney U)

(Figure S1A) and different response profile between running speed and the variation in ΔF/F when

compared with SST-INs (Figure S2B). Similar findings were observed when analyzing whisking

only periods. In contrast to SST-INs, we observed that whisking only was associated with a clear

positive increase in the activity of VIP-INs in S1 (Figure S2C,D). This increased sensitivity of VIP-

INs to active states is likely explained by the enrichment of VIP-INs in nicotinic cholinergic

receptors when compared to SST-INs (Figure S2E, F and (Gasselin et al., 2021)).

Importantly, and in contrast to SST-INs we observed comparable modulation of VIP-INs

by locomotion in V1 and S1 (Figure 3C, G-I) (Pakan et al., 2016). Analogous findings were

obtained while studying whisking periods. We further investigated a potential difference in the

strength of inhibition from VIP-INs to SST-INs between the two primary sensory regions using ex-

vivo whole-cell patch-clamp recording. For that we used animals that expressed the Cre

Recombinase in VIP-INs while the FLP recombinase was expressed in SST-INs (Figure 3J). Such

transgenic mice allowed us to selectively express channelrhodopsin (ChR2) in VIP-INs and to

record light-evoked IPSCs from fluorescently labeled SST-INs (Figure 3J). We normalized

responses through comparison to simultaneously patched neighboring PNs. As previously

reported, we observed that VIP-mediated inhibition was higher in SST-INs when compared to

local PNs, but that this ratio was similar between S1 and V1 (Figure 3K). Taken together, our

findings indicate that changes in VIP-IN-mediated inhibition are likely not responsible for the

differential impact of locomotion and whisking in SST-INs between V1 and S1. In contrast, VIP-

INs are active during similar behavior periods as SST-INs in S1 suggesting possible functional

interactions during active states.

VIP-INs inhibit SST-INs during active states in S1.

In agreement with the classical "disinhibitory" scenario (Pfeffer et al., 2013), the activation

of VIP-INs is expected to decrease the activity of SST-INs. Our observation that during rest/run

transitions both VIP- and SST-INs in S1 increase their activity is not easy to reconcile with such

a hypothesis. Modeling studies raised the hypothesis that such as disinhibition of local PNs by

VIP-INs activation can be responsible for a simultaneous increase in VIP- and SST-INs firing

during active states (Garcia Del Molino et al., 2017). Hence, we performed experiments to assess

the influence of manipulating VIP-INs activity on the recruitment of SST-INs during active

behaviors in S1.

To achieve this, we used VIP-Cre::SST-Flp transgenic mice to express the inhibitory

DREADD hM4Di in VIP-INs while GCaMP6s was selectively expressed in SST-INs (Figure 4A-

B). Mice were initially imaged under control conditions, and then re-imaged 30 minutes after the

injection of CNO (1-10mg/Kg) to monitor potential changes in the activity of SST-INs (Figure 4B).

In both control conditions and after CNO injection, we observed an increase in average activity of

SST-INs during locomotion (control: $\Delta F/F_{0\ rest} = 0.71 \pm 0.04$; $\Delta F/F_{0\ run} = 1.52 \pm 0.07$; n = 393; Run:

$\Delta F/F_{0\ rest} = 0.61 \pm 0.04$; $\Delta F/F_{0\ run} = 1.81 \pm 0.08$; n = 308 cells in 5 mice, p<0.0001, Mann-Whitney

U) (Figure 4C). Interestingly, when investigating the modulation at the single cell level, we found

that SST-INs exhibited greater sensitivity to locomotion, as evidenced by elevated LMI and

Pearson correlation values (LMI: control = 0.30 ± 0.02, n = 393, CNO = 0.40 ± 0.02, n = 308 cells,

p<0.001, Mann-Whitney U test; Pearson: control = 0.30 ± 0.02, n = 393 , CNO = 0.44 ± 0.02, n =

308 cells, p<0.01, Mann-Whitney U test) (Figure 4D). In control conditions the proportion of

negatively modulated cells was also reduced in CNO treated mice (Figure 4D). We subsequently averaged activity over all whisking periods aligning for whisking onset (Figure 4E). In control condition we observed that immediately after whisking there was a decrease in SST-INs activity that progressively increased as locomotion speed increased (Figure 4E-F). In CNO treated conditions, the inhibitory effect after whisking was absent and overall, there was a larger increase in positive modulation of SST-INs by locomotion (Figure 4F-E). Detailed analysis in the variation of SST-INs activity with running speed also revealed that after CNO treatment a similar speed produced a larger increase in $\Delta F/F_0$ signal in SST-INs (Figure 4G, H). These results suggest that during active states VIP-INs have an overall inhibitory action in SST-INs and raises the question about the underlying mechanism responsible for the active recruitment of L2/3 SST-IN during locomotor activity.

Visual stimulation alters SST-INs activity during active states in V1 but not in S1.

Based on our results, the elevation in SST-INs activity during locomotion is not primarily driven by VIP-mediated disinhibition of PNs. Instead, our results suggest the involvement of alternative mechanisms. Whisking as well as locomotion are known to increase activity of neuromodulatory inputs, namely cholinergic axons in neocortical regions (e.g. (Reimer et al., 2014)). Recent work in mice suggested that L2/3 SST-INs in V1 loose their responsiveness to cholinergic signaling, which could potentially account for their decreased sensitivity to active states (Yaeger et al., 2019) but see (Bugeon et al., 2022). We tested if cholinergic signaling in SST-INs was different between V1 and S1, which could potentially explain our results. Using carbachol, a general agonist of acetylcholine receptors, we observed both in V1 as well as S1 a depolarization of SST-INs that was blocked by the muscarinic receptor antagonist atropine (Figure S3). This finding suggests equal equipment in cholinergic signaling in SST-INs between the two sensory areas.

We thus looked for alternative pathways that could explain the positive modulation of SST-

INs during active states in S1. Recent work revealed that thalamic function is essential for the

arousal/movement-related activation of PNs in V1 (Nestvogel and McCormick, 2022). Inspired by

those findings we tested if feedforward signals transmitted to the cortex are also at the origin of

the observed modulation of SST-INs by active behavioral states. Such a possibility could also

eventually explain the difference in the modulation of SST-INs in V1 and S1 in the dark. Previous

works have reported that SST-INs in V1, which display modest to no modulation by locomotion in

the dark (Dipoppa et al., 2018; Fu et al., 2014; Pakan et al., 2016), show increased activity during

active states when mice are facing a grey screen (Dipoppa et al., 2018; Pakan et al., 2016). These

results are compatible with the idea that increased activity of thalamic neurons contribute to

defining the magnitude or direction of modulation of cortical SST-INs to behavioral transitions in

a sensory modality specific manner. We thus investigated whether in our working conditions visual

stimulation which is expected to increase the activity of the feedforward pathways into V1 altered

the sensitivity of SST-INs to behavioral transitions from rest to run in both S1 and V1. To achieve

this, we monitored the activity of SST-INs in both V1 and S1 under two conditions: in the dark and

in front of a grey screen (Figure 5A). We assessed changes in activity for both VIP- and SST-INs

from rest/run transitions by measuring variations of LMI as well as the Pearson correlation

between spontaneous activity and running speed (Figure 5). Our observations revealed that VIP-

INs activity remains relatively constant between S1 and V1, regardless of the dark or grey screen

conditions (Figure 5A-D). In contrast, SST-INs in V1 displayed a significantly greater sensitivity to

locomotion when facing a grey screen. Interestingly, the modulation of SST-INs in S1 by

locomotion appeared to be relatively unaffected by the dark versus grey screen conditions (Figure

5A-D). Altogether our results thus suggest that activity of feedforward pathways are important for

sensitivity of SST-INs to behavior state transitions.

Thalamic activity is required for the modulation of SST-INs by spontaneous behaviors.

As visual stimuli altered sensitivity of SST-INs to locomotion in V1 but not S1 we reasoned

that most likely thalamic activity is required for the modulation of SST-INs by active behavioral

states. We raised the hypothesis that in the dark, whisking leads to an increase in the S1

associated thalamic nuclei function (Petty et al., 2021) that is subsequently relayed to S1 and

explains the sensitivity of SST-INs to locomotion in the dark but not in V1. To directly test this

hypothesis, we inactivated the VPM and POM with microinjections of the GABA(A) agonist

muscimol (0.5 mM) and monitored the effect in SST-INs during resting and active states. We first

validated that we were able to effectively inactivate the somatosensory thalamus. For that we

conducted *in vivo* calcium imaging experiments in S1 while monitoring activity of PNs as well as

SST-INs in response to whisker stimulation in control conditions and after muscimol injections in

the somatosensory thalamus (Figure 6). We observed that in control conditions whisker

stimulation induced a clear increase in the activity of both SST-INs and PNs (Figure 6A-C and

Figure S4). In contrast, after muscimol injection this increase was absent in both cell types (Figure

6A-C and Figure S4). These results confirmed the effectiveness of muscimol in reducing thalamic

activity. Subsequently, we monitored the modulation of SST-INs in S1 by whisking and locomotion

in control conditions and after thalamus inactivation (Figure D-E). Interestingly when we evaluated

response of SST-INs to both whisking and locomotion there was a clear modification of the

response. In muscimol conditions there was a significant decrease in the response of SST-INs

during locomotion (control: $\Delta F/F_{0\ rest}$ = 0.53 ± 0.01; $\Delta F/F_{0\ run}$ = 1.02 ± 0.03; n = 1657; Run: $\Delta F/F_{0}$

$_{rest}$ = 0.53 ± 0.02; $\Delta F/F_{0\ run}$ = 0.76 ± 0.04; n = 541 cells in 8 mice, p<0.0001, Mann-Whitney U).

This effect translated into a decreased LMI both at the single cell level as well as when comparing

average values across the difference tested mice (LMI single cells: control = 0.18 ± 0.01, n =

1657, muscimol = 0.09 ± 0.01, n =541, <0.0001, Mann-Whitney U; Figure 6H, I). The Pearson

correlation between spontaneous activity and running speed was also significantly reduced by

thalamus inactivation (Pearson single cells: control = 0.18 ± 0.01, n = 1657, muscimol = 0.14 ±

0.01, n =541, <0.01, Mann-Whitney U; Figure 6I). Plotting variation in ΔF/F vs running speed

revealed once more that in control conditions SST-INs activity increased with running speed and

that this increase is significantly reduced after muscimol injections (Figure 6J). Overall, our results

suggest that in S1, modulation of SST-INs by spontaneous behaviors requires thalamic activity.

In S1 locomotion is associated with increased L2/3 pyramidal cell activity.

SST-INs are most likely not contacted directly by thalamic inputs (Sermet et al., 2019) and

receive information indirectly via their important connectivity with L2/3 PNs. We thus monitored

the *in vivo* spiking activity of L2/3 in S1 during behavior state transitions namely running and

whisking in control conditions and after muscimol injections in the somatosensory thalamus

(Figure 7). Our observations revealed a significant rise in spiking activity among L2/3 principal

neurons (PNs) in S1 during the transition from rest to running (control: $\Delta F/F_{0\,rest}$ = 0.22 ± 0.01; $\Delta F/F_{0}$

$_{run}$= 0.80 ± 0.03; n = 1918; p <0.0001, Mann-Whitney U; Figure 7B, C)). At the population level, we

noted that 44% of the cells exhibited a statistically significant increase in activity, while only 20%

displayed a decrease (Figure 7A-C). In contrast, after muscimol injection in the thalamus this

average increase in activity was completely absent (Figure 7E-G). In fact, positive correlation

between locomotion and GCaMP6s fluorescence traces was lost upon thalamus inactivation

(control: Pearson: 0.11 ± 0.01: n = 1918; run= -0.07 ± 0.0`; n = 371; p <0.0001, Mann-Whitney U;

Figure 7F, G). These results are in agreement with what has recently been reported using

electrophysiological approaches in the primary visual cortex (Nestvogel and McCormick,

2022). Furthermore, the increase in average ΔF/F values during running periods in pyramidal

cells was also not present in cells where somatosensory thalamus was inactivated by muscimol

injection (control: $\Delta F/F_{0\,rest}$ = 0.27 ± 0.01; $\Delta F/F_{0\,run}$= 0.22 ± 0.02; n = 371; p <0.0001, Mann-Whitney

U; Figure 7F, G). Hence, the convergence of locomotion-induced modulation and susceptibility

to thalamic inactivation in L2/3 PNs, similar to what we observed for SST-INs, further supports

the hypothesis that SST-INs in S1 inherit the increased activity during active behavioral states

from L2/3 PNs.

Strongly heterogeneous glutamatergic inputs to SST enable PN-drive despite an overall

weak connection strength

Glutamatergic inputs from L2/3 PNs to SST-INs are extremely weak, displaying low initial

release probably that progressively increases with continuous activation due to robust synaptic

facilitation (Kapfer et al., 2007; Pala and Petersen, 2015; Rozov et al., 2001). How can such weak

synapses transmit the increase in activity regimes in L2/3 PNs associated with active behavioral

states to L2/3 SST-INs? To address this, we performed paired recording between L2/3 PNs and

SST-INs in S1 and evaluated the strength of the excitatory connections between pairs of

connected neurons (Figure 7H). We observed that the synaptic strength (amplitude of first EPSC)

was highly variable among connections with an overall reduced mean amplitude and displaying

pronounced short facilitation (mean = 32 ± 6 pA, median = 12.8 pA, CV = 137%, n = 50) (Figure

7I, J). Interestingly, we observed that a small fraction (32 %) of cells displayed large initial current

amplitudes (amplitude of first EPSC >30 pA). In a subset of connected pairs, we managed to

successfully establish a third connection and we compared the synaptic properties between the

two different presynaptic neurons into the same postsynaptic SST-IN (Figure 7K,

L). Unexpectedly we observed that there was a positive correlation between the amplitude of the

synaptic currents between the two presynaptic neurons. These results suggest that the variability

in synaptic strength is not randomly distributed across glutamatergic synaptic inputs innervating

SST-INs but is determined by the postsynaptic subtype of SST-IN (Figure 7M). Overall, these

results indicate that a subpopulation of SST-INs is expected to be particularly sensitive to

variations in the activity level of L2/3 PNs that occur during the behavioral transitions from rest

active states in S1 in line with our initial *in vivo* observations (see Figure 1).

462
463

Discussion:

Using in vivo two-photon calcium imaging in awake mice we observed that behavioral state transitions do not equally affect the activity of L2/3 SST-INs across primary sensory regions. Specifically, we noted that in dark conditions, locomotion led to an increased activity of SST-INs in the S1 but had no effect in V1. Interestingly, when visual stimulation was present, this difference was lost and transitions from rest to running now positively modulated SST-INs in both primary sensory regions. Variable recruitment of VIP-INs does not seem to be at the origin of such differential behavior-state dependent modulation of SST-INs between the two sensory regions. In addition, using a chemogenetic approach we revealed that VIP-INs mainly inhibit SST-INs during rest/run transitions and are not the source of positive modulation of SST-INs by locomotion. In contrast, we observed that inactivation of somatosensory thalamus strongly reduces the modulation of SST-INs in S1 by spontaneous behaviors. Altogether our results provide a detailed description of the neuronal mechanisms at the origin of the differential modulation of VIP- and SST-INs by behavior states in S1.

The responses of neurons in primary sensory neocortical regions to sensory stimuli are influenced by behavioral states including, arousal level, whisking, attention, and locomotion (Bennett et al., 2013; Iriki et al., 1996; McGinley et al., 2015b; Petersen and Crochet, 2013). Disinhibitory circuits have been recognized as key players in mediating the neuromodulatory effects of such behavioral state transitions in neocortical networks. Specifically, the activity of VIP- and SST-INs is believed to allow gain control by behavioral state in the neocortex (Fu et al., 2015, 2014). In fact, previous studies in V1 revealed that, in the dark, VIP-INs are recruited during locomotion while the activity of SST-INs remains unchanged (Dipoppa et al., 2018; Fu et al., 2014; Pakan et al., 2016). Interestingly, our findings indicate that in S1, SST-INs are strongly recruited by locomotion even in the absence of light. This result is unexpected given the inhibitory connection between VIP-INs and SST-INs, as well as the established role of VIP-INs in inhibiting

SST-INs during whisking in stationary animals (Gasselin et al., 2021; Gentet et al., 2012; Lee et

al., 2013; Pfeffer et al., 2013; Pi et al., 2013). Interestingly, when we selected from our recording

sessions periods of whisking with no associated locomotion, we reproduced previous findings

observed in non-locomoting mice. Namely, whisking resulted in an increase of VIP-INs activity

and a decrease in recruitment of SST-INs (Gentet et al., 2012; Muñoz et al., 2017). Hence, our

observations indicate that whisking during rest or during locomotion are associated with distinct

cortical states, with the latter resulting in significantly stronger recruitment of SST-INs in S1.

Notwithstanding the multiple evidence that SST-INs are sensitive to behavioral state

transitions the neuronal mechanisms at the origin of such modulations are not obvious. In the S1,

whisking in stationary animals mainly inhibits SST-INs in layer 2/3 (Gentet et al., 2012; Muñoz et

al., 2017). Through the manipulation of VIP-INs function, Munoz and colleagues provided

compelling evidence for VIP-INs to be the source of SST-INs inhibition during whisking in awake

mice (Muñoz et al., 2017). Surprisingly, during such experiments, the authors observed a reversal

in the response of SST-INs. Inhibiting VIP-INs resulted in a transformation of the modulation into

a notable positive recruitment of SST-INs by whisking (Muñoz et al., 2017). This finding highlights

the pivotal role of VIP-INs in governing the activity of SST-INs during whisking. However, the

question that emerges concerns the source of excitation by whisking in L2/3 SST-INs when VIP-

INs are silenced. We hypothesized that the mechanisms underlying such positive modulation are

probably also at the origin of positive modulation by locomotion that we now observed in S1.

Interestingly, in L5 SST-INs, pharmacological block or genetic removal of muscarinic receptors

abolishes positive increase in SST-INs by whisking indicating that cholinergic inputs are likely to

participate in increased activity of SST-INs by whisking (Muñoz et al., 2017). However, if a similar

mechanism is present in L2/3 SST-INs is unknown. At odds with a possible cholinergic modulation

of L2/3 SST-INs during spontaneous behaviours, Yaeger and colleagues (Yaeger et al., 2019),

reported that in the V1 of adult mice (P>56), SST-INs lose their sensitivity to cholinergic

modulation. However, it is worth noting that Yaeger et al. mainly reported changes at the

suprathreshold level upon cholinergic receptor activation. Consistent with the notion that

cholinergic modulation of SST-INs is absent in adult mice, theoretical analysis yielded a

hypothesis proposing that VIP-INs, rather paradoxically, may be responsible for the increase in

SST-INs, a phenomenon termed "response reversal," achieved through the disinhibition of local

PNs (Garcia Del Molino et al., 2017).

Our observations reveal that inhibiting VIP-INs does not abolish the positive modulation

of SST-INs by locomotion; instead, it has the opposite effect, increasing the sensitivity of SST-

INs to transitions from rest to running. This suggests that positive recruitment of SST-INs involves

mechanisms independent of VIP-INs. Furthermore, we noted that despite relatively similar

recruitment of VIP-INs across sensory regions, SST-INs in S1 and V1 displayed different

response profiles to locomotion particularly in dark conditions. This result further argues that VIP-

INs are likely not the cause of SST-INs increase during locomotion in the neocortex. In contrast,

our experimental results illustrate that in the presence of visual stimulation modulation of SST-

INs by locomotion is altered in V1 but not in S1 suggesting that sensory region-specific

mechanisms are involved in such modulation.

Based on our current findings, we propose a new working model in which the positive

modulation of SST-INs activity across behavioral states relies on the interaction between thalamic

activity and neuromodulatory pathways, specifically cholinergic inputs. Our hypothesis is strongly

supported by our data, which demonstrates that inactivation of the thalamus significantly

diminishes the modulation of SST-INs in S1 by locomotion. We propose that the involvement of

thalamic activity partially explains the region-specific effects of spontaneous behaviors on SST-

INs due to different levels of ongoing sensory drive. Activity of thalamocortical VPM neurons, that

relay somatosensory information to S1, increases with whisking (Gutnisky et al., 2017; Poulet et

al., 2012; Urbain et al., 2015). While comparatively less explored, recent findings suggest a

notable upsurge in VPM activity also during locomotion (Huo et al., 2020). For the moment is

unclear if this representation of self-motion in VPM neurons is due to cortical inputs (Poulet et al.,

2012), neuromodulation (Petty et al., 2021) or proprioception (Severson et al., 2017).

Nonetheless, we anticipate that the combined effects of whisking and locomotion lead to an

overall increase in thalamic activity, which is subsequently transmitted to the neocortex.

Interestingly, our findings align with recent research indicating that depolarization of L2/3 principal

neurons (PNs) in V1 during locomotion is dependent on thalamic function (Nestvogel and

McCormick, 2022). In fact, we obtained similar results in our study (Figure 6). Under normal

conditions, locomotion leads to an overall increase in activity (positive LMI) in L2/3 PNs (Figure

7), and this increase is entirely abolished when thalamic activity is disrupted. As SST-INs are

highly contacted by L2/3 PNs we suggest that the increase in SST-INs is inherited from L2/3 PNs.

Worth noting that in V1 the differential response of SST-INs to locomotion is also observed in L2/3

PN whose LMI is significantly larger in the present of visual stimulation (Pakan et al., 2016),

compatible with the idea that SST-INs display behavior dependent modulation in part transmitted

from local L2/3 PNs.

We also observed significant variability at the single-cell level regarding the sensitivity of

S1 L2/3 SST-INs to behavioral state transitions (see Figure 1 and 2). This variability was followed

by a pronounced variance in the strength of unitary connections between PNs and SST-INs

(Figure 7). Previous proposals have suggested that skewed distributions of excitatory

postsynaptic potential (EPSP) amplitudes in SST-INs, with a relatively small mean but a skew

towards larger amplitudes, ensured individual PNs to excite a subset of SST-INs (Kapfer et al.,

2007). However, our findings indicate that these distributions are not entirely random among SST-

INs and that the strength of these connections is influenced by the postsynaptic cell. Recent

transcriptomic data further suggest the presence of at least three distinct subtypes of SST-INs in

S1 L2/3 (Wu et al., 2023). Remarkably the expression of the Elfn1, which is known to retrogradely

regulate release dynamics at glutamatergic inputs, exhibits differential expression patterns among

SST-INs subtypes (Lim et al., 2018; Paul et al., 2017). This leads us to speculate that the

properties of glutamatergic inputs are not uniform across the different SST-IN subtypes in L2/3,

potentially conferring unique computational properties to neocortical circuits. In the future, it would

be interesting to directly investigate whether *in vivo* SST-INs exhibiting greater sensitivity to

locomotion correspond to those with stronger unitary connections between PNs and SST-INs that

we recorded *in vitro*. Subpopulations of SST-INs, with respect to their response during rest-run

transitions, were previously documented in V1 (Reimer et al., 2014). However, it remained

uncertain whether these responses genuinely corresponded to distinct subpopulations of SSTs

or if they were merely a result of transgenic mouse models used to label SST-INs (Hu et al.,

2013). In a more recent development, a study that employed a genetic approach to identify

subpopulations of SST-INs unveiled distinct functional specializations within these discrete

subsets of SST-INs (Green et al., 2023), further supporting the idea of the existence of SST-IN

subgroups.

During our thalamic muscimol injections, we observed a complete block of the positive

modulation of PNs by locomotion while it was only reduced in SST-INs. Since the experiments on

PNs and SST-INs were not conducted simultaneously, it remains possible that experimental

differences, such as incomplete thalamic inactivation during SST-INs imaging sessions, could

account for the disparate results despite ensuring successful targeting and delivery of muscimol

to the somatosensory thalamus. However, our in vitro results indicate that SST-INs are equipped

with cholinergic receptors. Thus, it is still possible that the remaining modulation of SST-INs is

mediated by cholinergic neuromodulation that remained active during our thalamus inactivation

experiment. Such a possible scenario would indicate that both thalamic and cholinergic pathways

are essential in shaping the responses of SST-INs in S1 across the different behavioral states.

Nonetheless, future work involving the selective removal of muscarinic receptors from SST-INs is

needed to allow definite claims about the in vivo contribution of cholinergic pathways in the control

of L2/3 SST-INs during spontaneous behaviors.

Also evident was the observation that SST- and VIP-INs are equipped with different types

of cholinergic receptors. VIP-INs were particularly rich in nicotinic receptors which were almost

absent in SST-INs that seemed to express mainly muscarinic receptors. This differential

equipment of cholinergic receptors is expected to result also in differential sensitivity to behavioral

state transitions. In particular the high levels of nicotinic levels in VIP-INs are compatible with their

fast activation to even small behavioral state variations like fast whisking bouts (Gasselin et al.,

2021). In contrast for SST-INs, due to mainly muscarinic receptors, longer and more sustained

activation of cholinergic inputs activation is required to mediate depolarizations in SST-INs.

Mechanistically it is now clear that both thalamus as well as neuromodulatory inputs like

cholinergic projections originating from the basal forebrain participate in the modulation of cortical

networks by behavioral state. Less obvious is the computational role of the two different pathways.

The cholinergic pathway, via VIP-INs activation, has been proposed to underlie the gain control

of L2/3 PNs to sensory evoked activity across behavioral states (Fu et al., 2014). Thalamic activity

is required for the important depolarizations of L2/3 neurons (Nestvogel and McCormick, 2022)

during locomotion but their precise computational role is still ill defined. In any case by integrating

feedforward thalamic activity transmitted indirectly via L2/3 PNs as well as neuromodulatory

information both via VIP-mediated inhibition and via direct cholinergic innervation, SST-INs

occupy a key position in cortical networks to adjust activity of the two pathways and adjust

neuronal activity in function of behavior state.

References:

Alitto, H., Dan, Y., 2013. Cell-type-specific modulation of neocortical activity by basal forebrain input. Front. Syst. Neurosci. 6.

Ayaz, A., Saleem, A.B., Schölvinck, M.L., Carandini, M., 2013. Locomotion controls spatial integration in mouse visual cortex. Curr. Biol. CB 23, 890–894. https://doi.org/10.1016/j.cub.2013.04.012

Bennett, C., Arroyo, S., Hestrin, S., 2013. Subthreshold Mechanisms Underlying State-Dependent Modulation of Visual Responses. Neuron 80, 350–357. https://doi.org/10.1016/j.neuron.2013.08.007

Bugeon, S., Duffield, J., Dipoppa, M., Ritoux, A., Prankerd, I., Nicoloutsopoulos, D., Orme, D., Shinn, M., Peng, H., Forrest, H., Viduolyte, A., Reddy, C.B., Isogai, Y., Carandini, M., Harris, K.D., 2022. A transcriptomic axis predicts state modulation of cortical interneurons. Nature 607, 330–338. https://doi.org/10.1038/s41586-022-04915-7

Busse, L., Cardin, J.A., Chiappe, M.E., Halassa, M.M., McGinley, M.J., Yamashita, T., Saleem, A.B., 2017. Sensation during Active Behaviors. J. Neurosci. 37, 10826–10834. https://doi.org/10.1523/JNEUROSCI.1828-17.2017

Dipoppa, M., Ranson, A., Krumin, M., Pachitariu, M., Carandini, M., Harris, K.D., 2018. Vision and Locomotion Shape the Interactions between Neuron Types in Mouse Visual Cortex. Neuron 98, 602-615.e8. https://doi.org/10.1016/j.neuron.2018.03.037

Fu, Y., Kaneko, M., Tang, Y., Alvarez-Buylla, A., Stryker, M.P., 2015. A cortical disinhibitory circuit for enhancing adult plasticity. eLife 4, e05558. https://doi.org/10.7554/eLife.05558

Fu, Y., Tucciarone, J.M., Espinosa, J.S., Sheng, N., Darcy, D.P., Nicoll, R.A., Huang, Z.J., Stryker, M.P., 2014. A cortical circuit for gain control by behavioral state. Cell 156, 1139–1152. https://doi.org/10.1016/j.cell.2014.01.050

Garcia Del Molino, L.C., Yang, G.R., Mejias, J.F., Wang, X.-J., 2017. Paradoxical response reversal of top-down modulation in cortical circuits with three interneuron types. eLife 6, e29742. https://doi.org/10.7554/eLife.29742

Gasselin, C., Hohl, B., Vernet, A., Crochet, S., Petersen, C.C.H., 2021. Cell-type-specific nicotinic input disinhibits mouse barrel cortex during active sensing. Neuron 109, 778-787.e3. https://doi.org/10.1016/j.neuron.2020.12.018

Gentet, L.J., Kremer, Y., Taniguchi, H., Huang, Z.J., Staiger, J.F., Petersen, C.C.H., 2012. Unique functional properties of somatostatin-expressing GABAergic neurons in mouse barrel cortex. Nat. Neurosci. 15, 607–612. https://doi.org/10.1038/nn.3051

Gouwens, N.W., Sorensen, S.A., Baftizadeh, F., Budzillo, A., Lee, B.R., Jarsky, T., Alfiler, L., Baker, K., Barkan, E., Berry, K., Bertagnolli, D., Bickley, K., Bomben, J., Braun, T., Brouner, K., Casper, T., Crichton, K., Daigle, T.L., Dalley, R., Frates, R.A. de, Dee, N., Desta, T., Lee, S.D., Dotson, N., Egdorf, T., Ellingwood, L., Enstrom, R., Esposito, L., Farrell, C., Feng, D., Fong, O., Gala, R., Gamlin, C., Gary, A., Glandon, A., Goldy, J., Gorham, M., Graybuck, L., Gu, H., Hadley, K., Hawrylycz, M.J., Henry, A.M., Hill, D., Hupp, M., Kebede, S., Kim, T.K., Kim, L., Kroll, M., Lee, C., Link, K.E., Mallory, M., Mann, R., Maxwell, M., McGraw, M., McMillen, D., Mukora, A., Ng, Lindsay, Ng, Lydia, Ngo, K., Nicovich, P.R., Oldre, A., Park, D., Peng, H., Penn, O., Pham, T., Pom, A., Popović, Z., Potekhina, L., Rajanbabu, R., Ransford, S., Reid, D., Rimorin, C., Robertson, M., Ronellenfitch, K., Ruiz, A., Sandman, D., Smith, K., Sulc, J., Sunkin, S.M., Szafer, A., Tieu, M., Torkelson, A., Trinh, J., Tung, H., Wakeman, W., Ward, K., Williams, G., Zhou, Z., Ting, J.T., Arkhipov, A., Sümbül, U., Lein, E.S., Koch, C., Yao, Z., Tasic, B., Berg, J., Murphy, G.J., Zeng, H., 2020. Integrated Morphoelectric and Transcriptomic Classification of Cortical GABAergic Cells. Cell 183, 935-953.e19. https://doi.org/10.1016/j.cell.2020.09.057

Green, J., Bruno, C.A., Traunmüller, L., Ding, J., Hrvatin, S., Wilson, D.E., Khodadad, T.,
Samuels, J., Greenberg, M.E., Harvey, C.D., 2023. A cell-type-specific error-correction
signal in the posterior parietal cortex. Nature 620, 366–373.
https://doi.org/10.1038/s41586-023-06357-1
Gutnisky, D.A., Yu, J., Hires, S.A., To, M.-S., Bale, M.R., Svoboda, K., Golomb, D., 2017.
Mechanisms underlying a thalamocortical transformation during active tactile sensation.
PLOS Comput. Biol. 13, e1005576. https://doi.org/10.1371/journal.pcbi.1005576
Hu, H., Cavendish, J., Agmon, A., 2013. Not all that glitters is gold: off-target recombination in
the somatostatin–IRES-Cre mouse line labels a subset of fast-spiking interneurons.
Front. Neural Circuits 7.
Huo, Y., Chen, H., Guo, Z.V., 2020. Mapping Functional Connectivity from the Dorsal Cortex to
the Thalamus. Neuron 107, 1080-1094.e5. https://doi.org/10.1016/j.neuron.2020.06.038
Iriki, A., Tanaka, M., Iwamura, Y., 1996. Attention-induced neuronal activity in the monkey
somatosensory cortex revealed by pupillometrics. Neurosci. Res. 25, 173–181.
https://doi.org/10.1016/0168-0102(96)01043-7
Kapfer, C., Glickfeld, L.L., Atallah, B.V., Scanziani, M., 2007. Supralinear increase of recurrent
inhibition during sparse activity in the somatosensory cortex. Nat. Neurosci. 10, 743–
753. https://doi.org/10.1038/nn1909
Lee, S., Kruglikov, I., Huang, Z.J., Fishell, G., Rudy, B., 2013. A disinhibitory circuit mediates
motor integration in the somatosensory cortex. Nat. Neurosci. 16, 1662–1670.
https://doi.org/10.1038/nn.3544
Lim, L., Pakan, J.M.P., Selten, M.M., Marques-Smith, A., Llorca, A., Bae, S.E., Rochefort, N.L.,
Marín, O., 2018. Optimization of interneuron function by direct coupling of cell migration
and axonal targeting. Nat. Neurosci. 21, 920–931. https://doi.org/10.1038/s41593-018-
0162-9
McCormick, D.A., Nestvogel, D.B., He, B.J., 2020. Neuromodulation of Brain State and
Behavior. Annu. Rev. Neurosci. 43, 391–415. https://doi.org/10.1146/annurev-neuro-
100219-105424
McGinley, M.J., David, S.V., McCormick, D.A., 2015a. Cortical Membrane Potential Signature of
Optimal States for Sensory Signal Detection. Neuron 87, 179–192.
https://doi.org/10.1016/j.neuron.2015.05.038
McGinley, M.J., Vinck, M., Reimer, J., Batista-Brito, R., Zagha, E., Cadwell, C.R., Tolias, A.S.,
Cardin, J.A., McCormick, D.A., 2015b. Waking State: Rapid Variations Modulate Neural
and Behavioral Responses. Neuron 87, 1143–1161.
https://doi.org/10.1016/j.neuron.2015.09.012
Mitchinson, B., Martin, C.J., Grant, R.A., Prescott, T.J., 2007. Feedback control in active
sensing: rat exploratory whisking is modulated by environmental contact. Proc. R. Soc. B
Biol. Sci. https://doi.org/10.1098/rspb.2006.0347
Muñoz, W., Tremblay, R., Levenstein, D., Rudy, B., 2017. Layer-specific modulation of
neocortical dendritic inhibition during active wakefulness. Science 355, 954–959.
https://doi.org/10.1126/science.aag2599
Nestvogel, D.B., McCormick, D.A., 2022. Visual thalamocortical mechanisms of waking state-
dependent activity and alpha oscillations. Neuron 110, 120-138.e4.
https://doi.org/10.1016/j.neuron.2021.10.005
Niell, C.M., Stryker, M.P., 2010. Modulation of visual responses by behavioral state in mouse
visual cortex. Neuron 65, 472–479. https://doi.org/10.1016/j.neuron.2010.01.033
Pakan, J.M., Lowe, S.C., Dylda, E., Keemink, S.W., Currie, S.P., Coutts, C.A., Rochefort, N.L.,
2016. Behavioral-state modulation of inhibition is context-dependent and cell type
specific in mouse visual cortex. eLife 5, e14985. https://doi.org/10.7554/eLife.14985

Pala, A., Petersen, C.C.H., 2015. In Vivo Measurement of Cell-Type-Specific Synaptic
       Connectivity and Synaptic Transmission in Layer 2/3 Mouse Barrel Cortex. Neuron 85,
       68–75. https://doi.org/10.1016/j.neuron.2014.11.025

Paul, A., Crow, M., Raudales, R., He, M., Gillis, J., Huang, Z.J., 2017. Transcriptional
       Architecture of Synaptic Communication Delineates GABAergic Neuron Identity. Cell
       171, 522-539.e20. https://doi.org/10.1016/j.cell.2017.08.032

Petersen, C.C.H., Crochet, S., 2013. Synaptic Computation and Sensory Processing in
       Neocortical Layer 2/3. Neuron 78, 28–48. https://doi.org/10.1016/j.neuron.2013.03.020

Petty, G.H., Kinnischtzke, A.K., Hong, Y.K., Bruno, R.M., 2021. Effects of arousal and
       movement on secondary somatosensory and visual thalamus. eLife 10, e67611.
       https://doi.org/10.7554/eLife.67611

Pfeffer, C.K., Xue, M., He, M., Huang, Z.J., Scanziani, M., 2013. Inhibition of inhibition in visual
       cortex: the logic of connections between molecularly distinct interneurons. Nat. Neurosci.
       16, 1068–1076. https://doi.org/10.1038/nn.3446

Pi, H.-J., Hangya, B., Kvitsiani, D., Sanders, J.I., Huang, Z.J., Kepecs, A., 2013. Cortical
       interneurons that specialize in disinhibitory control. Nature 503, 521–524.
       https://doi.org/10.1038/nature12676

Polack, P.-O., Friedman, J., Golshani, P., 2013. Cellular mechanisms of brain state–dependent
       gain modulation in visual cortex. Nat. Neurosci. 16, 1331–1339.
       https://doi.org/10.1038/nn.3464

Poulet, J.F.A., Fernandez, L.M.J., Crochet, S., Petersen, C.C.H., 2012. Thalamic control of
       cortical states. Nat. Neurosci. 15, 370–372. https://doi.org/10.1038/nn.3035

Reimer, J., Froudarakis, E., Cadwell, C.R., Yatsenko, D., Denfield, G.H., Tolias, A.S., 2014.
       Pupil Fluctuations Track Fast Switching of Cortical States during Quiet Wakefulness.
       Neuron 84, 355–362. https://doi.org/10.1016/j.neuron.2014.09.033

Rozov, A., Burnashev, N., Sakmann, B., Neher, E., 2001. Transmitter release modulation by
       intracellular Ca2+ buffers in facilitating and depressing nerve terminals of pyramidal cells
       in layer 2/3 of the rat neocortex indicates a target cell-specific difference in presynaptic
       calcium dynamics. J. Physiol. 531, 807–826. https://doi.org/10.1111/j.1469-
       7793.2001.0807h.x

Saleem, A.B., Ayaz, A., Jeffery, K.J., Harris, K.D., Carandini, M., 2013. Integration of visual
       motion and locomotion in mouse visual cortex. Nat. Neurosci. 16, 1864–1869.
       https://doi.org/10.1038/nn.3567

Schneider, D.M., Nelson, A., Mooney, R., 2014. A synaptic and circuit basis for corollary
       discharge in the auditory cortex. Nature 513, 189–194.
       https://doi.org/10.1038/nature13724

Sermet, B.S., Truschow, P., Feyerabend, M., Mayrhofer, J.M., Oram, T.B., Yizhar, O., Staiger,
       J.F., Petersen, C.C., 2019. Pathway-, layer- and cell-type-specific thalamic input to
       mouse barrel cortex. eLife 8, e52665. https://doi.org/10.7554/eLife.52665

Severson, K.S., Xu, D., Van de Loo, M., Bai, L., Ginty, D.D., O'Connor, D.H., 2017. Active touch
       and self-motion encoding by Merkel cell-associated afferents. Neuron 94, 666-676.e9.
       https://doi.org/10.1016/j.neuron.2017.03.045

Shimaoka, D., Harris, K.D., Carandini, M., 2018. Effects of Arousal on Mouse Sensory Cortex
       Depend on Modality. Cell Rep. 22, 3160–3167.
       https://doi.org/10.1016/j.celrep.2018.02.092

Sofroniew, N.J., Cohen, J.D., Lee, A.K., Svoboda, K., 2014. Natural Whisker-Guided Behavior
       by Head-Fixed Mice in Tactile Virtual Reality. J. Neurosci. 34, 9537–9550.
       https://doi.org/10.1523/JNEUROSCI.0712-14.2014

Syeda, A., Zhong, L., Tung, R., Long, W., Pachitariu, M., Stringer, C., 2022. Facemap: a
       framework for modeling neural activity based on orofacial tracking.
       https://doi.org/10.1101/2022.11.03.515121

Urbain, N., Salin, P.A., Libourel, P.-A., Comte, J.-C., Gentet, L.J., Petersen, C.C.H., 2015.
Whisking-Related Changes in Neuronal Firing and Membrane Potential Dynamics in the
Somatosensory Thalamus of Awake Mice. Cell Rep. 13, 647–656.
https://doi.org/10.1016/j.celrep.2015.09.029
Vinck, M., Batista-Brito, R., Knoblich, U., Cardin, J.A., 2015. Arousal and Locomotion Make
Distinct Contributions to Cortical Activity Patterns and Visual Encoding. Neuron 86, 740–
754. https://doi.org/10.1016/j.neuron.2015.03.028
Wu, S.J., Sevier, E., Dwivedi, D., Saldi, G.-A., Hairston, A., Yu, S., Abbott, L., Choi, D.H.,
Sherer, M., Qiu, Y., Shinde, A., Lenahan, M., Rizzo, D., Xu, Q., Barrera, I., Kumar, V.,
Marrero, G., Prönneke, A., Huang, S., Kullander, K., Stafford, D.A., Macosko, E., Chen,
F., Rudy, B., Fishell, G., 2023. Cortical somatostatin interneuron subtypes form cell-type-
specific circuits. Neuron 0. https://doi.org/10.1016/j.neuron.2023.05.032
Yaeger, C.E., Ringach, D.L., Trachtenberg, J.T., 2019. Neuromodulatory control of localized
dendritic spiking in critical period cortex. Nature 567, 100–104.
https://doi.org/10.1038/s41586-019-0963-3
Yu, J., Hu, H., Agmon, A., Svoboda, K., 2019. Recruitment of GABAergic Interneurons in the
Barrel Cortex during Active Tactile Behavior. Neuron 104, 412-427.e4.
https://doi.org/10.1016/j.neuron.2019.07.027
Zhou, M., Liang, F., Xiong, X.R., Li, L., Li, H., Xiao, Z., Tao, H.W., Zhang, L.I., 2014. Scaling
down of balanced excitation and inhibition by active behavioral states in auditory cortex.
Nat. Neurosci. 17, 841–850. https://doi.org/10.1038/nn.3701
783
784

Figure Legends

Figure 1: SST-INs in V1 and S1 are differentially modulated by behavioral state transitions.
(A) Two-photon (2P) calcium imaging in S1 of awake mice. Mice were head fixed and were free
to run on a circular treadmill. Whisking activity was tracked via a video camera detecting infra-red
light focused on mouse snout illuminated with 850 nm LED. Bottom: Two-photon laser scanning
microscopy (2PLSM) image (maximum-intensity projection, MIP) of SST-Cre positive neurons
labelled with GCaMP6s in S1. Scale bar is 100 μm **(B)** Representative GCaMP6s fluorescence
traces (black) shown as $\Delta F/F_0$ (up) as well as normalized $\Delta F/F_0$ (bottom) for SST-INs displayed
in A. Each row represents a neuron sorted by weight on the first principal component (PC) of their
activity. Locomotion trace is shown on top (red). **(C)** Mean $\Delta F/F_0$ during periods of rest and run
($\Delta F/F_{0\ rest}$ = 0.80 ± 0.02; $\Delta F/F_{0\ run}$ = 1.84 ± 0.06; n = 2030 cells in 16 mice, p<0.0001, Mann-
Whitney U test). **(D)** Summary plot of Pearson correlation coefficients between calcium activity
and running speed. SST-INs showing statistically significant (see methods) modulation by
locomotion are represented by colored bars, with light orange indicating negative modulation and
positive modulation denoted by brown. The pie chart provides an overview of the overall
percentage of SST-INs showing positive, negative, or no effect in response to locomotion. **(E)**
Same as B but for SST-INs in the primary visual cortex. (**F, G**) Same as C and D but for V1. $\Delta F/F_0$

rest = 0.70 ± 0.03; $\Delta F/F_{0\ run}$ = 0.84 ± 0.03; n = 881 cells in 10 mice, p<0.0001, Mann-Whitney U.
(H) Event-triggered average traces of SST-INs GCaMP6s signal in S1 aligned to the start of
running (left) or whisking (right) along with running speed (red) and whisking (blue). Normalized
$\Delta F/F$ is displayed for all analyzed cells (bottom). Traces represent mean ± SEM values across all
recorded cells. **(I)** Same as G but for V1. **(J)** Summary plot of variation in $\Delta F/F$ between rest and
run and rest and whisking for the event-triggered average traces shown in G and H. Values were
obtained by averaging $\Delta F/F$ trace for each individual SST-INs in a 2 s window before and after
event onset (see Methods). $\Delta F/F_{0\ Loco\ S1}$ = 0.40 ± 0.03, n = 2030; $\Delta F/F_{0\ Loco\ V1}$ = 0.03 ± 0.02, n=
633, p<0.0001, Mann-Whitney U; $\Delta F/F_{0\ Whisk\ S1}$ = 0.08 ± 0.01, n = 853, n = 2030, $\Delta F/F_{0\ Whisk\ V1}$ =
0.002 ± 0.007; n = 853 cells in 10 mice, p<0.0001, Mann-Whitney U. **(K)** Cross-correlation
between in vivo calcium activity and running speed for SST-INs in S1 and V1 (left). (right) Average
correlation coefficient values at zero-time for locomotion and whisking for SST-INs recorded in
V1 and S1 (ρloco S1 = 0.22 ± 0.01, n = 2030, ρloco V1 = 0.06 ± 0.01, n = 883, p<0.0001, Mann-
Whitney U; ρwhisk S1 = 0.15 ± 0.01, n = 2030, ρloco V1 = 0.05 ± 0.01, n = 883, p<0.0001, Mann-
Whitney U).
818

819
820
Figure 2: Recruitment of SST-INs in S1 is primarily associated with behavioral state
transitions characterized by increased locomotor activity. (A) Example session illustrating
simultaneous recording traces for whisking (blue) and running speed (red). Whisking only periods
(no locomotor activity) are highlighted in light green. **(B)** Quantification of the fraction of time mice
spent whisking, wishing only and running during imaging sessions. **(C)** Event-triggered average
traces of SST-INs GCaMP6s fluorescence signal in S1 aligned to whisking onset. Only whisking
bouts not associated with locomotion were used (whisking only). Normalized ΔF/F is displayed
for all analyzed cells (bottom). Traces represent mean ± SEM values across all recorded cells.
(D) Enlarged view of average traces shown in C. **(E)** Summary plot of variation in ΔF/F between
rest and whisking for the event-triggered average traces shown in C and panel H figure1. Values
were obtained by averaging ΔF/F traces for each individual SST-INs in a 2 s window before and
after event onset. Note that during whisking only, there is either no effect or a slight tendency to
inhibit SST-INs. However, when whisking is associated with running periods, it clearly increases
the activity of SST-INs. $\Delta F/F_{0\ Whisk} = 0.08 \pm 0.01$, n = 853; $\Delta F/F_{0\ Whisk\ only} = -0.01 \pm 0.01$; n = 853

cells in 10 mice, p<0.001, Mann-Whitney U. **(F)** Summary plot of average mean locomotion,
whisking and whisking only modulation indices (LMI = 0.26 ± 0.01, n=2030; WMI = 0.21 ± 0.01,
n=856; WOMI = 0.06 ± 0.01, n=830) **(G)** Event-triggered average traces of SST-INs GCaMP6s
signal (ΔF/F) in S1 aligned to whisking onset. Results were separated according to percentage
time mice spent whisking only during the imaging session. Average of GCaMP6s fluorescence
from sessions with whisking only periods longer than 20% of the recording time is shown on the
left and less than 20% is illustrated on the right. Traces represent mean ± SEM values across all
recorded cells. **(H)** Summary plot of variation in ΔF/F between rest and whisking for the event-
triggered average traces shown in G. Average values were obtained 2 s before and 2 s after event
whisking onset (see Methods). Values were calculated for each individual SST-INs. ($\Delta F/F_0$
$_{Whisk<20\%}$ = 0.11 ± 0.02, n = 602; $\Delta F/F_{0\ Whisk<20\%}$ = -0.02 ± 0.01, n = 251, p<0.0001, Mann-Whitney
U). **(I)** Histograms of the distribution of locomotion modulation indices (LMI = $(R_L - R_S)/(R_L + R_S)$,
where R_L and R_S are the mean $\Delta F/F0$ during locomotion and stationary periods, respectively), for
SST-INs that displayed negative or positive WOMI.
849

850
Figure 3: VIP interneurons are strongly modulated by locomotion in S1. A) *(top)* Illustration
of a mouse brain depicting the location of S1 and V1. *(bottom)* 2PLSM image (maximum-intensity
projection, MIP) of VIP-Cre positive neurons labeled with GCaMP6f in S1. **B)** Representative
traces of in vivo calcium (GCaMP6f) activity in VIP-INs. Locomotion trace is shown on top (red)
followed by the calcium activity (green). Deconvolved activity (in black) is shown under each trace
C) Same as B but for VIP-INs in V1. **D)** Mean $\Delta F/F_0$ during periods of rest and run, ($\Delta F/F_{0\ rest}$ =
0.80 ± 0.02; $\Delta F/F_{0\ run}$ = 2.23 ± 0.05; n = 1915 cells in 11 mice, p<0.0001, Mann-Whitney U test).
E) Histogram of locomotion modulation indices for VIP-INs in S1. Inset depicts the average LMI
value (LMI = 0.38 ± 0.01, n= 1915). **F)** Distribution of Pearson correlation coefficients between

calcium activity and locomotion trace for VIP-INs in S1. Colored bars indicate VIP-INs that are
statistically significantly modulated by locomotion, either in a negative (orange) or positive
manner. The pie chart provides an overview of the overall percentage of VIP-INs showing positive,
negative, or no effect in response to locomotion. **G, H , I)** Same as D, E, F but for VIP-INs in V1.
J) Illustration of the double transgenic mouse line used for targeted protein expression in VIP-
and SST-INs respectively. Adeno-associated virus (AAV) coding Cre-dependent expression of
Channelrhodopsin2-EGFP along with a second AAV coding FLP-dependent expression of
mCherry were infected in primary visual (V1) and primary sensory (S1) cortices of VIP-Cre::SST-
Flp mice. Schematic diagram of experimental approach where VIP-INs were photo-activated via
ChR2 while a PN and neighboring SST-IN were patched in L2/3. **K)** *(left)* Example traces of IPSCs
simultaneously recorded in PNs and SST in V1 and S. *(middle)* Scatter plot of IPSCs amplitudes
recorded in SST-INs and PNs in S1 and V1 following VIP-INs photostimulation. (*right*) Ratio of
light-evoked IPSCs in V1 and S1 (V1: n = 7 cells, 2 mice ; S1: n = 10 cells, 2 mice). Mann Whitney
test p=0.8125. Error bars represent SEM.
874

875
Figure 4:VIP-INs mainly inhibit SST-INs during spontaneous behaviors. A) Simplified
illustration of the double transgenic mouse line used to target protein expression in VIP- and SST-
INs respectively based on the Cre/Lox and FLP/FRT system. Adeno-associated viruses (AAV)
were used to express in a Cre-dependent manner G_i-coupled inhibitory (hM4Di) designer
receptors exclusively activated by designer drugs (DREADD) in VIP-INs. A second AAV was used
to drive in a FLP-dependent manner the expression of GCaMP6s in SST-INs. **B)** Schematic
illustration of the canonical cellular organization in S1 and the expected negative modulation of
VIP-INs via activation of hM4Di by CNO injection. **C)** Mean $\Delta F/F_0$ during periods of rest and run
in control conditions and 30 after CNO injection (left), ($\Delta F/F_{0\ rest,\ control}$ = 0.71 ± 0.02; $\Delta F/F_{0\ run,\ control}$
= 1.5 ± 0.07; n = 393 cells in 5 mice, p<0.0001, Mann-Whitney U test; $\Delta F/F_{0\ rest,\ CNO}$ = 0.61 ± 0.04;
$\Delta F/F_{0\ run,\ CNO}$ = 1.81 ± 0.08; n = 308 cells in 5 mice, p<0.0001, Mann-Whitney U test). Summary
plot of LMI values obtained for SST-INs in control and after CNO injection (right), (LMI control =
0.30 ± 0.02, n = 393; LMI CNO = 0.4 ± 0.02, n = 308; p<0.001, Mann-Whitney U test). **D)** Variation
in Pearson correlation values between running speed and GCaMP6s fluorescence signals for
SST-INs recorded in control and CNO treated mice (ρloco control = 0.30 ± 0.02, n = 393, ρloco
CNO = 0.44 ± 0.02, n = 308, p<0.0001, Mann-Whitney U). Percentage of SST-INs exhibiting
statistically significant positive, negative, or non-significant Pearson correlation values is shown

on the right. **E)** Event-triggered average traces of SST-INs GCaMP6s signals in S1 aligned to the
onset of whisking along with running speed (red) and whisking (blue). Normalized ΔF/F is
displayed for all analyzed cells (bottom). Results are displayed for control conditions (left) and
after CNO injection (right). Traces represent mean ± SEM values across all recorded cells. **F)**
Enlarged view of average traces shown in E. **G)** Summary plot of variation in ΔF/F between rest
and whisking for the event-triggered average traces shown in F. Average values were obtained 2
s before and 2 s after event whisking onset (see Methods). Values were calculated for each
individual SST-INs. Note that in control conditions whisking induces a brief decrease in GCaMP6s
fluorescence that is absent in CNO treated mice. **H and F)** Variation in ΔF/F values for SST-INs
with running speed for control and CNO conditions.
903

904
905
Figure 5: Visual stimulation modulates the recruitment of SST-INs by locomotion in V1 but
not in S1. A) Schematic of the experimental setup used to probe the effect of visual stimulation
in the behaviour state-dependent modulation of the activity of VIP and SST-INs in V1 and S1. **B)**
Summary plots of LMI and Pearson correlation between running speed and GCaMP6
fluorescence signal for VIP- and SST-INs in the dark and facing a grey screen. LMI = 0.23 ± 0.01,
n =505, for VIP in the dark versus 0.21 ± 0.02, n = 222, in light; p > 0.9999, and 0.08 ± 0.01, n =
881 for SST in the dark versus 0.31 ± 0.02, n = 116 in light, p < 0.0001. Pearson Correlation: 0.21
± 0.01, n = 505, for VIP in the dark versus 0.20 ± 0.02, n = 222, in light; 0.06 ± 0.01, n = 881, for
SST in the dark versus versus 0.31 ± 0.02, n = 116 for SST-INs in light, p < 0.0001, Mann-Whitney
U test. **C)** Same as B but for S1. Values represent mean ± SEM.
916

917
Figure 6: Thalamic activity is required for the modulation of SST-INs by locomotion in S1.
A) Schematic illustration of experimental approach used to monitor effectiveness of thalamus
inactivation using muscimol injection in somatosensory thalamus. Whisker deflection was
achieved using localized air puff while cortical activity in S1 was monitored using *in vivo* two-
photon calcium imaging. **B)** Event-triggered average traces of SST-INs GCaMP6s signals in S1
aligned to whisker puff along with running speed (red). Normalized ΔF/F is displayed for all
analyzed cells (bottom). Results are displayed for control conditions (left) and for mice previously
injected with muscimol in the thalamus (*right*). Traces represent mean ± SEM values across all
recorded cells. **C)** Summary plot of variation in ΔF/F between rest and after whisker stimulation
for the event-triggered average traces shown in B. Average values were obtained 2 s before and
2 s after whisker stimulation onset (see Methods). **D)** Representative GCaMP6s fluorescence

38

traces (black) shown as $\Delta F/F_0$ (up) as well as normalized $\Delta F/F_0$ (bottom) for SST-INs in S1 for
mice spontaneously transitioning between rest and active behavior periods. Each row represents
a neuron sorted by weight on the first principal component (PC) of their activity. Locomotion trace
is shown on top (red). **E)** Same as A but for mice treated with muscimol injection in the
somatosensory thalamus (VPM and POM). **F)** Event-triggered average traces of SST-INs
GCaMP6s signals in S1 aligned to the onset of locomotion along with running speed (red) and
whisking (blue). Normalized $\Delta F/F$ is displayed for all analyzed cells (bottom). Results are
displayed for mice previously injected with muscimol in the thalamus. Traces represent mean ±
SEM values across all recorded cells. **G)** Mean $\Delta F/F_0$ during periods of rest and run in control
conditions and 30 min microinjection of muscimol in the somatosensory thalamus ($\Delta F/F_{0 \text{ rest control}}$
= 0.53 ± 0.01; $\Delta F/F_{0 \text{ run control}}$ = 1.01 ± 0.03; n = 1657 cells in 8 mice, p<0.0001, Mann-Whitney U
test; $\Delta F/F_{0 \text{ rest muscimol}}$ = 0.53 ± 0.02; $\Delta F/F_{0 \text{ run muscimol}}$ = 0.76 ± 0.04; n = 541 cells in 8 mice, p<0.0001,
Mann-Whitney U test). **H)** Summary plot of average LMI obtained in control and after muscimol
injection per the different individual mice tested ($LMI_{control}$ = 0.23 ± 0.06, $LMI_{muscimol}$ = 0.08 ± 0.06,
n = 8 mice, p = 0.0391, Wilcoxon test. **I)** Summary plots for LMI (left) and Pearson correlation
values (middle) between running speed and calcium activity for SST-INs in S1 obtained in control
and after muscimol injection in the thalamus ($LMI_{control}$ = 0.18 ± 0.01, n = 1657, $LMI_{muscimol}$ = 0.10
± 0.01, n = 541, p < 0.0001 Mann-Whitney U test; Pearson, $\rho_{control}$ = 0.18 ± 0.01, n = 1657, $\rho_{muscimol}$
= 0.14 ± 0.01, n = 541, p < 0.05 Mann-Whitney U test) . Percentage of SST-INs exhibiting
statistically significant positive, negative, or non-significant Pearson correlation values is shown
on the right. **J)** Variation in $\Delta F/F$ values for SST-INs with running speed for control and muscimol
injection conditions.
951

952
Figure 7: Behavioral state transitions from rest to active states increase activity of
pyramidal cells in S1. A) Top: schematic illustration of layer 2/3 microcircuits in S1; bottom: in
vivo 2PSLM images of S1 PNs labeled with GCaMP6s. **B)** Raster plot of the pyramidal cell activity
in the S1 from awake mice spontaneously transitioning between resting periods and periods of
locomotion. Each row represents a neuron sorted by weight on the first principal component (PC)
of their activity. **C)** Mean $\Delta F/F_0$ during periods of rest and run (Control: $\Delta F/F_{0\ rest}$ = 0.21 ± 0.01;
$\Delta F/F_{0\ run}$ = 0.80 ± 0.04; n = 1918 cells in 3 mice, p<0.0001, Mann-Whitney U test). **E)** Cross-
correlation between in vivo calcium activity and running speed for Pyramidal neurons in S1 in
control condition and after muscimol injections in somatosensory thalamus. **F-G)** Same as B and
C but mice previously microinjected with muscimol in thalamus **(**Muscimol: $\Delta F/F_{0\ rest}$ = 0.18 ± 0.01;
$\Delta F/F_{0\ run}$ = 0.17 ± 0.16; n = 371 cells in 2 mice, p<0.0001, Mann-Whitney U test). **H)** Schematic
illustration of dual whole-cell patch clamp configuration used to record synaptic properties of
unitary connections between individual L2/3 PNs and local SST-INs. **I)** Example traces of a train
of EPSCs recorded from two different SST-INs in response to presynaptic action potential
induction in two different L2/3 PNs. Note the pronounced different in the amplitude of the first
EPSC between the two SST-INs. **J)** Histogram of amplitudes of first EPSC obtained for 50
synaptic unitary connections. Inset depicts the relationship between amplitude and failure rate for
the 50 connections. **K)** Same as for **H** but illustrating that the properties of two different
presynaptic L2/3 PNs were recorded for the same postsynaptic SST-INs. **L)** Example traces of
EPSCs recorded in two different SST-INs for four presynaptic PNs. **M)** Summary plot indicating

the relationship between the amplitude obtained when stimulating PN1 or PN2 for the same
postsynaptic SST-INs.
975
976

977
Supplementary figure 1: Variability of SST-INs response within individual fields of view in
S1 to behavioral transitions from rest to run. A) Representative GCaMP6s fluorescence traces
(black) shown as $\Delta F/F_0$ (up) as well as normalized $\Delta F/F_0$ (bottom) for SST-INs for individual fields
recorded in awake mice spontaneously transitioning between rest and run periods. Each row
represents a neuron sorted by weight on the first principal component (PC) of their activity.
Locomotion (red) and whisking (blues) traces are shown on top. **B)** Scatter plots of the mean
amplitude of fluorescence changes (DF/F_0) of each neuron illustrated in A for locomotion periods
versus stationary periods.
986

987
988
Supplementary figure 2: Response of SST- and VIP-INs in S1 to behavioral transitions from
rest to run. **A)** Cross-correlation curves between running speed and calcium activity for VIP and
SST-INs. Values at zero time are displayed on the right. **B)** Variation in ΔF/F values for SST- and
VIP-INs with average running speed. **B)** Event-triggered average traces of VIP-INs GCaMP6s
fluorescence signal in S1 aligned to whisking onset. Only whisking bouts not associated with
locomotion were used (whisking only). Normalized ΔF/F is displayed for all analyzed cells
(bottom). Traces represent mean ± SEM values across all recorded cells. **(C)** Enlarged view of
average traces shown in A. **B)** Schematic illustration of whole-cell current clamp recording in L2/3
of primary somatosensory (S1) cortex while puff applying nicotine 100 ⌈M in SST or VIP-INs (left).
Example traces of puff-evoked responses in SST- and VIP-INs (right). **C)** Summary plot of
variations in membrane potential induced by nicotine applications in both SST- and VIP-INs (SST
[n = 11] vs VIP [n = 9], p < 0.0001, Mann-Whitney test).
1001

1002
Supplementary figure 3: SST-INs are equipped with cholinergic receptors in both V1 and
S1.
A) Schematic illustration of whole-cell current clamp recording in L2/3 of primary somatosensory
(S1) cortex while puff applying carbachol (2mM) in SST (up). Example traces of puff-evoked
responses in SST-INs (down). **B)** Example traces of current clamp recordings illustrating the
variation in membrane potential in SST-INs to puff application of carbachol in both V1 (P>56: top
right) and S1 (P<56 middle right; P>56 bottom right). **C)** Summary plot of the variation of
membrane potential induced by carbachol (CCH) in SST-INs in V1 and S1 (V1, SST [n = 7]; S1,
SST with CCH puff alone [n = 20] versus SST with CCH puff in the presence of atropine 10⌠M [n
= 7], p = 0.007, Mann-Whitney test)**.** Error bars are SEM.
1013

1014
Supplementary figure 4: Somatosensory thalamus inactivation with muscimol injection
decreases whisker-evoked pyramidal cell activity in S1. A) Event-triggered average traces of
L2/3 PNs GCaMP6s signals in S1 aligned to whisker puff along with running speed (red).
Normalized ΔF/F is displayed for all analyzed cells (bottom). Results are displayed for control
conditions (left) and for mice previously injected with muscimol in the thalamus (*right*). Traces
represent mean ± SEM values across all recorded cells. **B)** Summary plot of variation in ΔF/F
between rest and after whisker stimulation for the event-triggered average traces shown in B.
Average values were obtained 2 s before and 2 s after whisker stimulation onset (see Methods).
1023

Experimental Procedures:

Mice:

Animals were housed in the Paris Brain Institute accredited by the French Ministry of Agriculture. All experiments were performed in compliance with French and European regulations on care and protection of laboratory animals (EU Directive 2010/63, French Law 2013-118, February 6th, 2013), and were approved by local ethics committees and by the French Ministry of Research and Innovation (authorization numbers #11199). All animal work was performed at the ICM PHENO-ICMice facility. Both male and female mice aged 4 weeks or older were used with the following genotypes: SST-IRES-Cre (SSTtm2.1(cre)Zjh/J; JAX 013044); SST-IRES-Cre X Ai9 (Gt(ROSA)26Sortm9(CAG-tdTomato); JAX 007909); VIP-IRES-Cre (Viptm1(cre)Zjh/AreckJ; JAX 031628) X Ai9 (Gt(ROSA)26Sortm9(CAG-tdTomato); JAX 007909); VIP-cre::SST-FLP ($Sst^{tm3.1(flpo)Zjh}$/J; JAX 028579). Animals were maintained on a 12-hour light/dark cycles with food and water provided ad libitum. Animals were housed in groups of 2-6 animals per cage, as often as possible, to favor social interactions, a condition that prevents the emergence of depressive and anxious-like behavior (Berry et al., 2012; Ieraci et al., 2016; Kappel et al., 2017).

Slice preparation

Acute parasagittal slices (320 μm) were prepared from adult mice, starting from postnatal days 25–70. Mice were deeply anesthetized with a mix of ketamine/xylazine (mix of i.p. ketamine 16 mg/kg and xylazine 13 mg/kg) and perfused transcardially with ice-cold cutting solution containing the following (in mM): 220 Sucrose, 11 Glucose, 2.5 KCl,1.25 NaH2PO4, 25 NaHCO3, 7 MgSO4, 0.5 CaCl2. After the perfusion the brain was quickly removed and slices prepared using a vibratome (Leica VT1200S). Slices containing S1 barrel field or V1 were transferred to ACSF solution at 34 ^{o}C containing (in mM): 125 NaCl, 2.5 KCl, 2 CaCl2, 1 MgCl2, 1.25 NaH2PO4, 25

1051 NaHCO3, 15 Glucose for 15–20 min. After the period of recovery, slices were kept at room

1052 temperature for a period of 5-6 h.

1053

Electrophysiology:

Whole-cell patch-clamp recordings were performed close to physiological temperature

(32–34 °C) using a Multiclamp 700B amplifier (Molecular Devices) and fire-polished thick-walled

glass patch electrodes (1.85 mm OD, 0.84 mm ID, World Precision Instruments); 3.5–5 MOhm

tip resistance. For voltage-clamp recordings, cells were whole-cell patched using following

intracellular solution (in mM): 90 Cs-MeSO3, 10 EGTA, 40 HEPES, 4 MgCl2, 5 QX-314, 2.5

CaCl2,10 Na2Phosphocreatine, 0.3 MgGTP, 4 Na2ATP (300 mOsm pH adjusted to 7.3 using

CsOH). Extracellular synaptic stimulation was achieved by applying voltage pulses (20 us, 5–50

V; Digitimer Ltd, UK) via a second patch pipette filled with ACSF and placed 20–40 mm from

soma. For current clamp experiments patch pipettes were filled with the following intracellular

solution (in mM): 135 K-gluconate, 5 KCl, 10 HEPES, 0.01 EGTA, 10 Na2phosphocreatine, 4

MgATP, 0.3 NaGTP (295 mOsm, pH adjusted to 7.3 using KOH). The membrane potential (Vm)

was held at -60 mV, if necessary, using small current injection (typically in a range between -50

pA and 200 pA). Recordings were not corrected for liquid junction potential. Series resistance was

compensated online by balancing the bridge and compensating pipette capacitance. APs were

initiated by brief current injection ranging from 1000–2000 pA and 1–5 ms duration. For all

experiments data were discarded if series resistance, measured with a 10 mV pulse in voltage

clamp configuration, was >20 MOhm or changed by more than 20% across the course of an

experiment. For current clamp experiment cells were excluded if input resistance varied by more

than 25% of the initial value. Recordings of puff evoked cholinergic responses in SST- and VIP-

INs were performed at the resting membrane potential in the presence of 10 μM CNQX and 10

μM bicuculine.

All recordings were low-pass filtered at 10 kHz and digitized at 100 kHz using an analog-

to-digital converter (model NI USB 6259, National Instruments, Austin, TX, USA) and acquired

with Nclamp software (Rothman and Silver, 2018) running in Igor PRO (Wavemetrics, Lake

Oswego, OR, USA). The intracellular solution used for the experiments in which a presynaptic

cell was depolarized by a train of action potentials to induce neurotransmitter release on a post-

synaptic cell/element was (in mM): 130 K-MeSO3, 4 MgCl2,10 HEPES, 0.01 EGTA, 4 Na2ATP,

0.3 NaGTP (300 mOsm, pH 7.3 adjusted with NaOH). Paired recordings connections were probed

using a train of 5/9 action potentials at frequency of 20 Hz.

1084

In vivo two-photon calcium imaging

Surgery

Mice were initially anesthetized with Isoflurane. Induction of anaesthesia was performed

at 4% isoflurane, Iso-Vet, and an air flow of 250ml/min. Mice were then placed in a stereotactic

device (Kopf) kept at 37 °C using a regulated heating blanket and a thermal probe and maintained

under anaesthesia using 1-2% Isoflurane and an air flow of 250 ml/min until the end of the

surgery. For calcium imaging in VIP- and SST-INs, cranial window implantation and AAV viral

stereotaxic injection was performed in the same surgical procedure. In order to prevent brain

swelling and post-operative pain, anti-inflammatory (Dexamethasone, 2 µg/g Dexadreson, MSD)

and analgesic drugs (Buprenorphine, 0.1 mg/kg Buprecare) were injected subcutaneously. After

shaving the scalp, the skin was cleaned by wiping multiple times with an antiseptic solution and

70% alcohol. After injecting a local anaesthetic under the scalp (lidocaine, 10mg/kg - Xylovet or

Laocaine) a section of the skin (a circular section with 1.5 cm diameter) was removed using

surgical scissors. The periosteum was then carefully removed and the skull was scrapped using

a dental drill, in order to remove all residues. The surface of the skull was subsequently thoroughly

cleaned and dried using cottons swabs and a physiological saline solution. Using the stereotactic

device, the center of the cranial window was marked and a circular piece of bone was removed

using a 3 mm biopsy punch (LCH-PUK-30 Kai Medical). This step of the surgical procedure was

considered critical since damage to the dura would result in rapidly opacifying cranial windows.

Once the brain was exposed, the AAV viral vectors carrying the genetically encoded calcium

indicators (GCaMP6 f and s) were stereotactically injected. For calcium imaging of VIP-INs, SST-

INs and pyramidal neurons, AAV9 Syn-Flex-GCaMP6f-WPRE-SV40 ($2.1x10^{13}$ AddGene-

100833), AAV9 Syn-Flex-GCaMP6s-WPRE-SV40 ($2.5x10^{13}$ AddGene-100845) or AAV9 ENN-

AAV-CaMKII.GCaMP6f-WPRE-SV40 ($2.5x10^{13}$ AddGene-100834) was used respectively. Using

a Nanojet III (Drummond Scientific), 200nl of virus were injected in the barrel cortex (from Bregma,

1.1 RC; 3.3 ML) or visual cortex (from bregma: 3.0 RC; 2.4 ML) at a rate of 1nl/s followed by 5

minutes of wait to prevent backflow. A round of 3mm glass coverslip (CS-3R Warner Instruments)

was then placed over the craniotomy and glued in place using dental cement (Superbond). A

stainless-steel head post (Luigs & Neumann) was then attached to the skull also using dental

cement. For 3 days following the cranial windows surgery, mice were injected once a day with a

mix of anti-inflammatory (Dexamethasone, 2µg g^{-1} Dexadreson, MSD) and analgesic drugs

(Buprenorphine, 0.1 mg/kg Buprecare). Two weeks after surgery, mice were first habituated to

the experimenter by handling. The following days, mice were head-fixed in the acquisition setup

for increasing periods of time (5 – 45 minutes).

Thalamus inactivation

In order to inactivate the somatosensory-associated thalamus, following the acquisition of

several control imaging sessions, mice were placed in a stereotactic device, and put under light

isoflurane anesthesia. A hole was drilled over the ipsilateral thalamus and using a cannula, a

bolus of fluorescent Muscimol (0.5 mM, Muscimol, BODIPY™ TMR-X Conjugate Muscimol,

BODIPY™ TMR-X Conjugate, ThermoFischer Scientific) was slowly injected into the thalamus

(Coordinates: -1.82 RC, 1.5 ML, 3.5 depth). After injection, the craniotomy was covered with

silicone (Kwik-Cast, WPI) and the mouse was allowed to recover for an additional 30 minutes.

After recovery, two-photon imaging was performed for a maximum of one hour post muscimol

injection. The effective inactivation of the thalamus was confirmed functionally by recording the

response of pyramidal cells as well as local cortical interneurons to whisker stimulation as well as

by performing histology just after the imaging sessions.

Two-Photon Imaging

In vivo two-photon calcium imaging was performed with an Ultima IV two-photon laser-

scanning microscope system (Bruker), using a 20×, 1.0 N.A. water immersion objective (Olympus)

with the femtosecond laser (MaiTai DeepSee, Spectra Physics) tuned to 920 nm for imaging of

cells expressing GCaMP6s/f and to 1020 nm for imaging of cells expressing mCherry.

Fluorescence light was separated from the excitation path through a long pass dichroic (660dcxr;

Chroma, USA), split into green and red channels with a second long pass dichroic (575dcxr;

Chroma, USA), and cleaned up with band pass filters (hq525/70 and hq607/45; Chroma, USA).

Fluorescence was detected using both proximal epifluorescence photomultiplier tubes (gallium

arsenide phosphide, H7422PA-40 SEL, Hamamatsu). Time-series movies of neuronal

populations expressing GCaMP6f were acquired at the frame rate of 30Hz (512 × 512 pixels field

of view; 1.13 µm/pixel). The duration of each focal plane movie was 300s (9000 frames) to track

spontaneous neuronal activity. During the recording periods animals were free to run on a circular

treadmill covered with a soft foam. This foam softened the movement of the animals and provided

better traction (this was important since mice adapted more rapidly to the experimental setup and

movements were more fluid during the recordings). Locomotion was monitored using a rotary

encoder (H5-1000-NE-S, US Digital) and digitized with a NIDAQ analog-to-digital converter

(model NI USB 6363, National Instruments, Austin, TX, USA) at 2000 Hz. Whisker movement

was recorded using a FLIR camera (BFS-U3-17S7M-C) at a frame rate of 30Hz and using the

Spinnaker SDK software (FLIR Systems Inc, Orlando, USA) equipped with a cleanup filter

(FB850-40, Thor labs) to block laser shining from during two-photon imaging. The camera was

focused on the whisker pad of the mouse using a 10X (13 - 130mm FL) C-Mount, Close Focus

Zoom lens and each frame acquisition was controlled via Neuromatic running in IGORPRO

(Wavemetrics) environment using a national instruments digital-to-analog board (NI-6393). The

mouse snout was illuminated via an 850 nm LED (M850L3, Thor labs).

Whisker Stimulation

For whisker stimulation experiments, a NPI PDES-02DX (NPI Electronic GmbH) puffer

was used. Tubing was connected to the puffer and a short puff (1 sec) of air was aimed at the

mouse whiskers. The tip of the puffer was kept out of reach of exploratory whisking range. The

whisker puff was randomly triggered during imaging sessions and the time of puff was recorded

via the digitization of voltage trigger using a analog-to-digital converter (model NI USB 6259,

National Instruments, Austin, TX, USA)

CNO Experiments:

For experiments involving DREADDs, mice were injected with clozapine-N-oxide (CNO).

A water-soluble salt preparation of CNO, CNO dihydrochloride (Tocris, Bio-Techne LTD,

Abingdon, UK, catalog no.: 6329), which was dissolved in sterile saline. Several concentrations

of CNO were tested (1, 5 and 10mg/kg) and no significant changes were found so data was all

data was pulled together.

Visual stimulation

To study the effect of light on the recruitment of VIP- and SST-INs in the cortex, an

isoluminant grey screen (Lilliput 669GL 7") was placed at 45° from the center line of the mouse

and at approximately 20cm from the left eye (contralateral to the craniotomy) of the mouse. Visual

stimulation trials were interleaved with control trials. In order to reduce light contamination, a

custom funnel was used to shield the objective from light coming from the screen.

Calcium Imaging Analysis

Motion correction, Image segmentation and calcium signal extraction

Motion correction, image segmentation and calcium signal extraction were performed using the python 3.6 package Suite2P (Pachitariu et al., 2016). This package analyses data in four stages: image registration, region of interest (ROI) detection, quality control, and activity and neuropil extraction. The first stage, image registration, is responsible for motion correction and works by performing a mean image of a few frames and then shifting (in the x and y axis) each frame so that it optimally fits the mean image. The second stage, ROI detection, detects the cells in the frame. This is done by detecting the pixels with the highest variability across time and then expanding the region to the surrounding pixels until this variability drops below a defined threshold. Although this method works quite well, the disadvantage of this method is that since only active cells are detected, it is difficult to determine the number of silent cells in field of view, which could be important for functional tasks. In a third stage, quality control, the program separates structures that correspond to cells from the remaining structures by assigning them a score. The results of this stage were always manually curated. In the last stage of the analysis, activity and neuropil extraction, is performed by averaging the pixel intensity in the ROI for each time point. To determine the neuropil, a donut-like surrounding the ROI is generated and the activity within this area is also averaged for each time point. Neuropil plays a critical role in calcium imaging since it represents the activity of nearby structures surrounding the cell of interest that might be excited by the 2P excitation PSF. Since these structures might have a different behavioral profile than the cell of interest their activity might contaminate the signals detected in the cell soma (Dipoppa et al., 2018).

Analysis of fluorescence signals

Raw fluorescence traces were corrected for neuropil contamination by subtracting

Suite2p-neuropil traces using a fixed scaling coefficient of 0.7 (Dipoppa et al., 2018). To be able

to compare activity across cells and mouse lines, fluorescence signals were normalized. To do

this we used the $\Delta F/F_0$ method, calculated using the following formula: $(F-F_0)/F_0$, where F is the

fluorescence and F_0 is the baseline fluorescence. When studying the activity of interneurons (VIP-

and SST-INs), the baseline fluorescence was calculated using the lower 5^{th} percentile. Results

were systematically evaluated using a different baseline method to estimate F0 (MinMax baseline

method with a sliding window of 60s) and observations reported in the present manuscript were

insensitive to the baseline method used. For the analysis of pyramidal cell activity, a sliding

average window (60 s) defined as MinMax method was used.

Locomotion Analysis

Changes in positioning of the circular treadmill (sampled at 2 kHz) were interpolated onto

a down-sampled rate of 30Hz in order to match the sampling rate of the two-photon imaging.

Locomotion periods, were defined as the periods where speed was higher than 0.1 cm/s. All

points 0.5 seconds before and 1.5 seconds after a locomotion period were considered stationary

periods. Events spaced by less than 0.5s were joined. Sessions in which the percentage of either

locomotion or stationary periods were lower than 5%, were excluded. The locomotion modulation

index (LMI) was defined as the difference between the mean $\Delta F/F_0$ during locomotion (F_{run}) and

stationary (F_{rest}) periods, normalized by the sum of both values: LMI = $(F_{run} - F_{rest})/(F_{run} + F_{rest})$

(Pakan et al., 2016). To calculate statistically significant Pearson Correlations between

fluorescence and running speed, comparisons were made with 1000 shuffled (circular shuffling)

locomotion traces. Pearson correlation coefficient was considered statistically significant if value

of original data was above a 95% confidence interval generated from the 1000 shuffled traces.

Whisking Analysis

Whisker pad movement was tracked using the python framework Facemap (Syeda et al.,

2022) (https://github.com/MouseLand/facemap). Whisker-pad movement was measured by

selecting an ROI over the whisker pad and calculating the motion energy index (MEI) across the

video. MEI was defined as the sum of the absolute change in pixel intensity within the ROI

between adjacent video frames. Using custom python code, whisking pad motion energy was

filtered with a gaussian filter (sigma of 3), normalized with a Min Max method and resampled to

30Hz. Whisking periods were defined as periods where normalized MEI exceeded 10 %. Whisking

only periods were defined as periods in which there was whisking but not locomotion. Whisking

modulation index (WMI) and whisking only modulation index (WOMI) were calculated in a similar

way to LMI but using the variations in GCaMP6s fluorescence during resting or whisking periods.

Event-triggered averages were computed by automatically identifying frames with event

onsets, namely those related to whisking and/or locomotion. Due to the slow time course of

GCaMP6s and to minimize cross-talk between events, we specifically chose locomotion events

that were not preceded by another locomotion period within a 5-second interval. For whisking only

analysis, whisking episodes not associated with locomotion were included in the average (5se

before and 5 sec after). In Figure 3I, locomotion events were further categorized in agreement

with locomotion duration. Once multiple locomotion and/or whisking events were detected, the

results were combined and averaged to calculate an average value for each cell. Subsequently,

these individual average cell values were averaged together and presented with their respective

average and SEM.

1242

Statistics

All statistical analysis were performed in Prism (GraphPad Software, Inc.). The error bars

in all graphs indicate standard error of the mean (S.E.M.) and statistics were performed with two-

tailed tests. For statistical tests comparing the average $\Delta F/F_0$ of neurons between two behavioral

states (stationary versus locomotion periods) in the same subject, we used Wilcoxon signed-rank

tests. For statistical tests comparing the distribution of LMI, Pearson Correlation or behavior

properties between brain regions or interneuron populations, Mann-Whitney U tests were used.

When comparing two conditions across behavioral states, Kruskal-Wallis test was used.

1251
1252
1253