Feedforward circuits enable modality-specific control of cortical inhibition during behavioral state transitions.

Marcel de Brito Van Velze[1], Dhanasak Dhanasobhon[1], Marie Martinez[1], Annunziato Morabito[1], Emmanuelle Berthaux[1], Cibele Martins Pinho[1], Yann Zerlaut[1#] and Nelson Rebola[1#]

[1]ICM, Paris Brain Institute, Hôpital de la Pitié-Salpêtrière, Sorbonne Université, Inserm, CNRS, Paris, 75013, France.

#These authors contributed equally.

Number of figures: 7; Supplementary figures: 4; Supplementary tables: 0

Keywords: Neocortical interneurons, Somatostatin positive Interneurons, Somatosensory cortex, Spontaneous behaviours, Locomotion, Whisking.

1

Acknowledgements:

This study was supported by the Centre National de la Recherche Scientifique, Agence Nationale de la Recherche (grants ANR-EXCIGLY and ANR-DecoSensoMol1) the European Research Council (ERC-STG- 678250 to N.R.), the European Union's Horizon 2020 research and innovation programme (Marie Skłodowska-Curie Grant 892175 "*InProsMod*" to Y.Z.); by the Paris Brain Institute, and the Fondation pour la Recherche Médicale (FRM; fellowship ARF201909009117 to Y.Z.). We thank the Paris Brain Institute core facilities, namely iVector and ICMice PHENOPARC. We would also like to thank Dr. Alberto Bacci (ICM, Paris) and Dr. Joana Lourenco (ICM, Paris) for the valuable discussions and comments on the manuscript.

SUMMARY (236 words)

Active behavior strongly influences sensory information processing in primary sensory cortices. Local interneurons, especially those involved in disinhibitory circuits like somatostatin (SST) and vasoactive intestinal peptide (VIP)-positive interneurons, play a crucial role in regulating cortical function during behavior. While numerous studies have focused on the recruitment of pyramidal neurons, the intricate network mechanisms governing the behavior-state-dependent modulation of inhibitory neurons remain unclear. Using intravital two-photon calcium imaging we have studied the neuronal pathways that regulate SST- and VIP-IN activity in sensory cortices during spontaneous behaviours. We now reveal that active behavioral states associated with increased locomotor activity strongly recruit L2/3 SST-INs in primary somatosensory (S1) but not visual (V1) cortex indicating modality-specific mechanisms control activity of cortical interneurons during active states. Such a difference between V1 and S1 disappears in the presence of visual sensory drive (ambient light) suggesting a critical involvement of feedforward sensory pathways in the positive modulation of SST-INs during spontaneous behaviors. In agreement, we now unveil that inactivating the somatosensory thalamus significantly reduces the recruitment of SST-INs in S1 by locomotion. In contrast, interfering with disinhibitory circuits via chemogenetic inactivation of VIP-INs further increased the sensitivity of SST-INs to variations in behavioural states. Our work thus reveals a previous unknown role for thalamic inputs in driving SST-INs, suggesting that by integrating feedforward activity with neuromodulation, SST-INs play a central role in adapting sensory processing to behavioral states in a sensory modality-specific manner.

Introduction:

In the mammalian brain, the cerebral cortex is responsible for complex cognitive functions, including a coherent representation of the external world that is crucial for an effective interaction with a constantly changing environment. Importantly, sensation can occur under drastically different behavior conditions, which have been reported to result in various perceptual outcomes (Busse et al., 2017; McGinley et al., 2015a). A hallmark of sensory systems is thus to provide a flexible strategy to process incoming information in a context-dependent manner (McCormick et al., 2020).

Local neocortical interneurons, in particular those involved in disinhibitory circuits, like vasoactive intestinal peptide (VIP) and somatostatin (SST) expressing interneurons (INs), have been shown to be particularly sensitive to behavior state transitions and to be key contributors to the state-dependent regulation of cortical function (Fu et al., 2015, 2014; Gasselin et al., 2021; Gentet et al., 2012; Muñoz et al., 2017; Polack et al., 2013; Reimer et al., 2014). In V1, recruitment of VIP-INs and consequent decrease in SST-IN mediated inhibition has been suggested to underlie the gain modulation of visual-evoked responses observed during locomotion (Fu et al., 2014). Yet, the mechanisms controlling the engagement of SST-INs across active behavioral states are unclear. In addition, behavioral modulations have been shown to differentially impact pyramidal cell activity across sensory cortices (Ayaz et al., 2013; Niell and Stryker, 2010; Polack et al., 2013; Saleem et al., 2013; Schneider et al., 2014; Shimaoka et al., 2018; Vinck et al., 2015; Zhou et al., 2014). However, whether inhibitory circuits display similar or variable sensitivity to behavioral state transitions across sensory cortical areas remains unexplored.

A recent study revealed that modulation of INs activity in V1 by brain states can be largely predicted by the first transcriptomic principal component (Bugeon et al., 2022). This transcriptomic component suggested that the increase in SST-INs activity, observed in alert states, is associated with cholinergic signalling (Bugeon et al., 2022). However, this finding

4

contrasts with a previous report that SST-INs in V1 lose cholinergic modulation during

development (Yaeger et al., 2019). Paradoxically, VIP-mediated disinhibition has been

proposed to explain the increase in SST-INs activity during active behavioral states but this

hypothesis lacks experimental evidence (Garcia Del Molino et al., 2017). Both hypotheses point

to neuromodulatory inputs, in particular cholinergic axonal projections to be at the origin of the

sensitivity of VIP- and SST-INs to behavioral state transitions. Yet, recent results highlighted

that thalamic activity is essential for the large and sustained depolarizations associated with

arousal/movement-related signals in L2/3 PNs in V1 (Nestvogel and McCormick, 2022). These

results suggest that behavior states can modulate neocortical activity not only through

neuromodulatory projections such as cholinergic and noradrenergic pathways, but also via

glutamatergic thalamic inputs. As SST-INs receive pronounced inputs from local L2/3 PNs it is

therefore possible that feedforward activity transmitted via the thalamus participates in the

behavioral-dependent modulation of SST-INs in primary sensory regions. Still, as glutamatergic

synapses from L2/3 PNs into SST-INs have relatively low release probabilities it is unclear

whether an increase in L2/3 PNs activity during active states is sufficient to drive SST-INs.

Altogether, the specific neuronal mechanisms governing the activity of SST-INs during

spontaneous behaviors are still not fully understood.

The modulation of disinhibitory circuits by active behavioral states has been the focus of

considerable scrutiny in V1 (Bugeon et al., 2022; Dipoppa et al., 2018; Fu et al., 2014; Pakan et

al., 2016; Polack et al., 2013; Reimer et al., 2014). Locomotion, in particular, has been found to

influence the activity of VIP- and SST-INs in a context-dependent manner, with the modulation

of SST-INs being contingent on the presence or absence of ambient light (Dipoppa et al., 2018;

Pakan et al., 2016). Whether such modulation by locomotion extends to other sensory cortices

remains unclear. In S1, behavioral periods associated with increased whisking (in stationary

animals) have been reported to increase L2/3 VIP- activity with concomitant decrease in L2/3

SST-INs firing in agreement with the presence of disinhibitory circuits (Gasselin et al., 2021;

Gentet et al., 2012; Lee et al., 2013; Muñoz et al., 2017; Yu et al., 2019). Nevertheless, in S1,

and in stark contrast to V1, our knowledge regarding the impact of locomotion on the operation

of SST- and VIP-INs remains limited. This becomes especially relevant considering that, during

navigation, mice typically employ locomotion in conjunction with whisking to explore their

surroundings.

Here we used intravital two-photon calcium imaging in S1 and V1 of awake, head-fixed

mice, to investigate how transitions in behavioral states (namely whisking and locomotion)

modulate the activity of SST- and VIP-INs, key disinhibitory elements of cortical networks. We

found that in darkness, the recruitment of SST-INs by locomotion was significantly higher in S1

than V1. We also observed that the increase of SST-INs activity in S1 occurred despite the

strong recruitment of VIP-INs, a known inhibitor of SST-INs. In addition, neither the strength of

inhibition provided by VIP-INs nor the sensitivity of SST-INs to cholinergic signaling was

different across sensory regions. Importantly, inactivation of thalamic function using muscimol

microinjections limited the increase in activity of SST-INs by locomotion. These results reveal

that behavior state-transitions, often associated with changes in arousal, induce sensory

modality specific changes to the activity of inhibitory circuits, most likely through inputs coming

from the sensory thalamus. By integrating feedforward thalamic activity transmitted indirectly via

L2/3 PNs as well as neuromodulatory information coming from VIP-mediated inhibition and

cholinergic innervation, SST-INs occupy a pivotal position in cortical networks by adapting

sensory processing to behavioral states in a sensory modality-specific manner.

Results:

Activity of L2/3 SST-INs in S1 and V1 during spontaneous behavior

We initially focused our attention on the *vivo* response of SST-INs to behavioral state transitions in the mouse primary somatosensory cortex (S1). In fact, multiple studies have studied the activity of disinhibitory circuits during active behavioral states in V1. However, it remains unclear whether inhibitory circuits display similar or variable sensitivity to behavior state transitions in other sensory cortical areas including S1. To monitor alterations in firing dynamics of SST-INs we expressed the genetically encoded calcium indicator GCaMP6s selectively in SST-IN using transgenic mice (SST-cre) (Figure 1A) and changes in somatic fluorescence of GCaMP6s were used as a proxy for spiking activity. Fluorescence signals were recorded using two-photon imaging in awake head fixed mice that spontaneously transitioned between rest and locomoting periods on a circular treadmill (Figure 1A). Whisking activity was simultaneously acquired using a video camera focused on the mouse snout (Figure 1A). This experimental setup allowed us to study how SST-INs activity varied across active behavioral states namely during whisking with and without associated running.

In our initial analysis, we categorized recording periods based on whether animals were engaged in whisking, locomotion, or neither. Periods during which whisking or locomotion were not detected were classified as resting (see Methods). Through the quantification of average GCaMP6s fluorescence levels we found that in S1, SST-INs displayed markedly higher activity levels during periods of locomotion compared to resting ($\Delta F/F_{0\ rest}$ = 0.80 ± 0.02; $\Delta F/F_{0\ run}$ = 1.84 ± 0.06; n = 2030 cells in 10 mice, p<0.0001, Mann-Whitney U test, Figure 1A-C). This engagement of SST-INs during locomotion in S1 was also observed when we investigated the variability in the response to behavior state transitions at the single cell level. For that we quantified the correlation between baseline activity and running speed for individual SST-INs (Figure 1D). We observed a predominantly positive correlation coefficient across SST-INs in S1 (Pearson correlation, ρ = 0.22 ± 0.01, n = 2030, Figure 1D). Specifically, 59% of SST-INs

exhibited a statistically significant (see methods) positive modulation in response to locomotion,

while only a smaller but not negligible fraction of SST-INs displayed a significant negative

modulation (20%) (Figure 1E)

These results contrasted with previous reports in V1, where in darkness, locomotion had

on average, no impact in SST-INs activity (Dipoppa et al., 2018; Pakan et al., 2016). In order to

test if this functional difference was linked to our experimental conditions or if in fact reflected a

previously unknown region-specific modulation of SST-INs by locomotion we repeated the

experiments listed above but now monitoring SST-INs activity in V1 (Figure 1E). In agreement

with previous studies we observed that the modulation of SST-INs by locomotion in the dark in

V1 was quite modest ($\Delta F/F_{0\ rest}$ = 0.70 ± 0.03; $\Delta F/F_{0\ run}$ = 0.84 ± 0.03; n = 881 cells in 10 mice,

p<0.0001, Mann-Whitney U; (Dipoppa et al., 2018; Pakan et al., 2016) (Figure 1E,F) contrasting

the results obtained in S1. This difference in activity of SST-INs during locomotion between S1

and V1 was also observed at the single cell level. In contrast to S1, our observations in V1

revealed a comparable ratio of SST-INs exhibiting positive (33%) and negative (25%)

correlations between running speed and spontaneous activity (Figure 1G).

In order to better estimate the rate of change in calcium levels in SST-INs between S1

and V1 we generated population averages of all running or whisking bouts aligned to locomotion

or whisking onset respectively (Figure 1H, I). In these population averages it was possible to

observe that both locomotion as well as whisking induced changes in $\Delta F/F0$ in SST-INs that

were significantly larger in S1 than in V1 (Figure 1J, I). The differential sensitivity of SST-INs

between the two sensory areas to behavioral state transitions was particularly evident when

comparing the cross-correlation curves between spontaneous activity of SST-INs and the

locomotion or whisking traces (Pearson correlation: locomotion, S1, ρ = 0.20 ± 0.01, n=2030;

V1, ρ = 0.05 ± 0.01, n=633; whisking, S1, ρ = 0.15 ± 0.01, n=853; V1, ρ = 0.04 ± 0.01, n=633,

Figure 1K). Altogether these results suggest that in the neocortex, behavior-dependent

modulation of SST-INs activity is not uniform across sensory regions.

Contribution of whisking and locomotion to SST-INs modulation in S1

Our results indicate that in S1 active behaviors like whisking and locomotion, result in increased activity of SST-INs. However locomotion in mice is closely linked to the motion of their whiskers (Mitchinson et al., 2007). In contrast, whisking can take place in both run and rest conditions. We thus decided to further investigate how the two different active states (whisking and locomotion) impacted the activity of SST-INs. In our experimental conditions mice were observed to spend an average of 32% of their time running, 47% whisking, and within this whisking period, 16% did not involve associated ambulatory activity (see Methods) (Figure 2A-B). To facilitate a more straightforward comparison on how whisking and locomotion shaped activity in SST-INs in S1 we specifically isolated recording periods in which mice were engaged in active whisking without simultaneous running (Figure 2C-E). Under these conditions, which we defined as "whisking only," the activity of SST-INs exhibited distinct differences compared to whisking episodes that were accompanied by locomotion. Specifically, SST-INs were not positively modulated during active whisking showing in contrast a tendency towards inhibition in agreement with previous works (Gasselin et al., 2021; Gentet et al., 2012; Lee et al., 2013; Muñoz et al., 2017) (Figure 2C-E). To further quantify how the different behavioral states influenced SST-INs activity we separated calcium activity and we defined whisking only modulation index (WOMI) and whisking modulation index (WMI) similar to our previous LMI calculation (see Methods). We observed that both WMI and LMI were clearly positive and rather similar in agreement with the observation that in mice most whisking bouts are actually associated with locomotion (Sofroniew et al., 2014). In contrast WOMI, was only slightly positive with a large percentage of cells displaying negative modulation (Figure 2F).

During our imaging sessions, we observed significant variation among mice in terms of the duration of whisking without concurrent locomotion (% time WO = 15.6 ± 1.7%, CV = 105%, n = 86 sessions). In light of this observation, we embarked on investigating whether the

response profile of SST-INs differed between sessions characterized by high or low percentages of WO. Our rationale was that sessions with a higher proportion of WO activity would provide a better opportunity to sample and visualize the specific impact of whisking with and without associated locomotion. To achieve this, we employed an arbitrary threshold of 20% to categorize imaging sessions into those with high or low percentages of WO (Figure 2G, H). We subsequently identified all whisking bouts and averaged corresponding variations in GCaMP6s fluorescence across the two different groups of sessions aligned to whisking onset. Upon examination of the population's average response, distinct variations in the response of SST-INs to whisking became evident between the two types of sessions (Figure 2G, H). In sessions with a high percentage of whisking only, whisking onset was associated with a clear decrease in the activity of SST-INs immediately after whisking onset. In contrast, in sessions where mice spent only a small fraction of their time whisking without locomotion, whisking onset was clearly associated with an increase in SST-INs firing (Figure 2G-H). Overall, these results indicate that in S1, whisking at rest induces a decrease in SST-INs activity, in agreement with previous reports (Gentet et al., 2012; Muñoz et al., 2017; Yu et al., 2019). On the other hand, when mice engage in active behavioral states involving locomotion, activity of somatostatin interneurons (SST-INs) is increased, specifically in the primary somatosensory cortex (S1), but not in the primary visual cortex (V1; Fig. 1). These findings suggest that distinct active behavioral states exert varying effects on the activity of S1 SST-INs.

Morphoelectric and transcriptomic analysis have highlighted that SST-INs, even within L2/3, are a highly heterogeneous neuronal population (Gouwens et al., 2020; Wu et al., 2023). We probed if sensitivity to locomotion was also heterogeneous across individual SST-INs in S1. To evaluate this, we separated SST-INs according to their response to WOMI. We observed that SST-INs that displayed a negative WOMI, were on average less sensitive to locomotion (Figure 2I). In contrast, SST-INs with positive WOMI mostly increase their activity with locomotion (Figure 2I). Variable modulation of SST-INs for locomotion was observed even

within single fields of view, suggesting that variability in SST-IN responses cannot be attributed

to distinct behavioral states (Figure S1). In summary, these findings indicate the presence of

various subtypes of SST-INs in Layer 2/3, exhibiting varying degrees of sensitivity to transitions

in behavioral states.

VIP-INs are equally engaged by locomotion in V1 and S1

Having observed that in S1, SST-INs are recruited during active behavioral states,

specifically during locomotion, we proceeded to investigate the underlying neuronal

mechanisms involved in such modulation. Taking into account previous works we identified

three main potential pathways that could participate in controlling L2/3 SST-INs activity during

spontaneous behaviors: (i) inhibitory top-down modulation via VIP-INs (Fu et al., 2014; Lee et

al., 2013; Pfeffer et al., 2013; Pi et al., 2013) (ii) increased local recurrent excitatory L2/3 PNs

activity (Nestvogel and McCormick, 2022; Pakan et al., 2016) and (iii) direct excitatory

neuromodulation (*e.g.* via direct cholinergic inputs) (Yaeger et al., 2019). We initially focused on

VIP-INs, which are known to be important inhibitors of SST-IN function (Fu et al., 2014; Lee et

al., 2013; Pfeffer et al., 2013; Pi et al., 2013) and are strong recipients of neuromodulatory

inputs (*e.g.* cholinergic) associated with active behavioral states (Alitto and Dan, 2013; Gasselin

et al., 2021). Additionally, theoretical work suggested that VIP-INs activation in certain

conditions can result in unexpected increase in SST-INs function through disinhibition of local

PNs (Garcia Del Molino et al., 2017). Therefore, we studied the contribution of VIP-INs to the

activity patterns of SST-INs during behavioral state transitions. We also explored the possibility

that differential activity/function of VIP-INs could be at the origin of the distinct sensitivity of SST-

INs in V1 and S1 to locomotion.

We started by examining how activity of VIP-INs varied in S1 during the same behavioral

transitions as we described for SST-INs (Figure 3A-C). In line with previous findings (Fu et al.,

2015, 2014), we observed an increase in activity of VIP-INs during ambulatory periods in S1

($\Delta F/F_{0\ rest}$ = 0.79 ± 0.02; $\Delta F/F_{0\ run}$ = 2.23 ± 0.05; n = 1915 cells; in 10 mice, p<0.0001, Mann-

Whitney U, Figure 3A, B, D) with the majority of VIP-INs displaying a positive modulation by

locomotion (LMI = 0.38 ± 0.01; Figure 3E). A similar observation was obtained while estimating

Pearson correlation between GCaMP6f or GCaMP6s fluorescence and running speed. VIP-INs

displayed on average a positive correlation (ρ = 0.30 ± 0.01, Figure 3F) with only a very small

fraction (6%) of VIP-INs displaying a statistically significant decrease in activity (Figure 3F)

during locomotion. It is worth mentioning that the percentage of VIP-INs displaying negative

modulation by locomotion was significantly less than SST-INs (see Figure 1). Despite the similar

average positive modulation by locomotion between VIP- and SST-INs in S1, we observe that at

the population level, VIP-INs displayed a higher sensitivity to behavioral state transitions than

SST-INs (Figure S2). This was reflected into a higher correlation coefficient between

spontaneous activity and running speed (VIP: ρ = 0.30 ± 0.01 vs SST: 0.22± 0.01, p < 0.001,

Mann-Whitney U) (Figure S1A) and different response profile between running speed and the

variation in $\Delta F/F$ when compared with SST-INs (Figure S2B). Similar findings were observed

when analyzing whisking only periods. In contrast to SST-INs, we observed that whisking only

was associated with a clear positive increase in the activity of VIP-INs in S1 (Figure S2C,D).

This increased sensitivity of VIP-INs to active states is likely explained by the enrichment of VIP-

INs in nicotinic cholinergic receptors when compared to SST-INs (Figure S2E, F and (Gasselin

et al., 2021)).

Importantly, and in contrast to SST-INs we observed comparable modulation of VIP-INs

by locomotion in V1 and S1 (Figure 3C, G-I) (Pakan et al., 2016). Analogous findings were

obtained while studying whisking periods. We further investigated a potential difference in the

strength of inhibition from VIP-INs to SST-INs between the two primary sensory regions using

ex-vivo whole-cell patch-clamp recording. For that we used animals that expressed the Cre

Recombinase in VIP-INs while the FLP recombinase was expressed in SST-INs (Figure 3J).

Such transgenic mice allowed us to selectively express channelrhodopsin (ChR2) in VIP-INs

and to record light-evoked IPSCs from fluorescently labeled SST-INs (Figure 3J). We normalized responses through comparison to simultaneously patched neighboring PNs. As previously reported, we observed that VIP-mediated inhibition was higher in SST-INs when compared to local PNs, but that this ratio was similar between S1 and V1 (Figure 3K). Taken together, our findings indicate that changes in VIP-IN-mediated inhibition are likely not responsible for the differential impact of locomotion and whisking in SST-INs between V1 and S1. In contrast, VIP-INs are active during similar behavior periods as SST-INs in S1 suggesting possible functional interactions during active states.

VIP-INs inhibit SST-INs during active states in S1.

In agreement with the classical "disinhibitory" scenario (Pfeffer et al., 2013), the activation of VIP-INs is expected to decrease the activity of SST-INs. Our observation that during rest/run transitions both VIP- and SST-INs in S1 increase their activity is not easy to reconcile with such a hypothesis. Modeling studies raised the hypothesis that such as disinhibition of local PNs by VIP-INs activation can be responsible for a simultaneous increase in VIP- and SST-INs firing during active states (Garcia Del Molino et al., 2017). Hence, we performed experiments to assess the influence of manipulating VIP-INs activity on the recruitment of SST-INs during active behaviors in S1.

To achieve this, we used VIP-Cre::SST-Flp transgenic mice to express the inhibitory DREADD hM4Di in VIP-INs while GCaMP6s was selectively expressed in SST-INs (Figure 4A-B). Mice were initially imaged under control conditions, and then re-imaged 30 minutes after the injection of CNO (1-10mg/Kg) to monitor potential changes in the activity of SST-INs (Figure 4B). In both control conditions and after CNO injection, we observed an increase in average activity of SST-INs during locomotion (control: $\Delta F/F_{0\ rest}$ = 0.71 ± 0.04; $\Delta F/F_{0\ run}$ = 1.52 ± 0.07; n = 393; Run: $\Delta F/F_{0\ rest}$ = 0.61 ± 0.04; $\Delta F/F_{0\ run}$ = 1.81 ± 0.08; n = 308 cells in 5 mice, p<0.0001, Mann-Whitney U) (Figure 4C). Interestingly, when investigating the modulation at the single cell

level, we found that SST-INs exhibited greater sensitivity to locomotion, as evidenced by elevated LMI and Pearson correlation values (LMI: control = 0.30 ± 0.02, n = 393, CNO = 0.40 ± 0.02, n = 308 cells, p<0.001, Mann-Whitney U test; Pearson: control = 0.30 ± 0.02, n = 393 , CNO = 0.44 ± 0.02, n = 308 cells, p<0.01, Mann-Whitney U test) (Figure 4D). In control conditions the proportion of negatively modulated cells was also reduced in CNO treated mice (Figure 4D). We subsequently averaged activity over all whisking periods aligning for whisking onset (Figure 4E). In control condition we observed that immediately after whisking there was a decrease in SST-INs activity that progressively increased as locomotion speed increased (Figure 4E-F). In CNO treated conditions, the inhibitory effect after whisking was absent and overall, there was a larger increase in positive modulation of SST-INs by locomotion (Figure 4F-E). Detailed analysis in the variation of SST-INs activity with running speed also revealed that after CNO treatment a similar speed produced a larger increase in $\Delta F/F_0$ signal in SST-INs (Figure 4G, H). These results suggest that during active states VIP-INs have an overall inhibitory action in SST-INs and raises the question about the underlying mechanism responsible for the active recruitment of L2/3 SST-IN during locomotor activity.

Visual stimulation alters SST-INs activity during active states in V1 but not in S1.

Based on our results, the elevation in SST-INs activity during locomotion is not primarily driven by VIP-mediated disinhibition of PNs. Instead, our results suggest the involvement of alternative mechanisms. Whisking as well as locomotion are known to increase activity of neuromodulatory inputs, namely cholinergic axons in neocortical regions (e.g. (Reimer et al., 2014)). Recent work in mice suggested that L2/3 SST-INs in V1 loose their responsiveness to cholinergic signaling, which could potentially account for their decreased sensitivity to active states (Yaeger et al., 2019) but see (Bugeon et al., 2022). We tested if cholinergic signaling in SST-INs was different between V1 and S1, which could potentially explain our results. Using carbachol, a general agonist of acetylcholine receptors, we observed both in V1 as well as S1 a

depolarization of SST-INs that was blocked by the muscarinic receptor antagonist atropine (Figure S3). This finding suggests equal equipment in cholinergic signaling in SST-INs between the two sensory areas.

We thus looked for alternative pathways that could explain the positive modulation of SST-INs during active states in S1. Recent work revealed that thalamic function is essential for the arousal/movement-related activation of PNs in V1 (Nestvogel and McCormick, 2022). Inspired by those findings we tested if feedforward signals transmitted to the cortex are also at the origin of the observed modulation of SST-INs by active behavioral states. Such a possibility could also eventually explain the difference in the modulation of SST-INs in V1 and S1 in the dark. Previous works have reported that SST-INs in V1, which display modest to no modulation by locomotion in the dark (Dipoppa et al., 2018; Fu et al., 2014; Pakan et al., 2016), show increased activity during active states when mice are facing a grey screen (Dipoppa et al., 2018; Pakan et al., 2016). These results are compatible with the idea that increased activity of thalamic neurons contribute to defining the magnitude or direction of modulation of cortical SST-INs to behavioral transitions in a sensory modality specific manner. We thus investigated whether in our working conditions visual stimulation which is expected to increase the activity of the feedforward pathways into V1 altered the sensitivity of SST-INs to behavioral transitions from rest to run in both S1 and V1. To achieve this, we monitored the activity of SST-INs in both V1 and S1 under two conditions: in the dark and in front of a grey screen (Figure 5A). We assessed changes in activity for both VIP- and SST-INs from rest/run transitions by measuring variations of LMI as well as the Pearson correlation between spontaneous activity and running speed (Figure 5). Our observations revealed that VIP-INs activity remains relatively constant between S1 and V1, regardless of the dark or grey screen conditions (Figure 5A-D). In contrast, SST-INs in V1 displayed a significantly greater sensitivity to locomotion when facing a grey screen. Interestingly, the modulation of SST-INs in S1 by locomotion appeared to be relatively unaffected by the dark versus grey screen conditions (Figure 5A-D). Altogether our results thus

suggest that activity of feedforward pathways are important for sensitivity of SST-INs to

behavior state transitions.

Thalamic activity is required for the modulation of SST-INs by spontaneous behaviors.

As visual stimuli altered sensitivity of SST-INs to locomotion in V1 but not S1 we

reasoned that most likely thalamic activity is required for the modulation of SST-INs by active

behavioral states. We raised the hypothesis that in the dark, whisking leads to an increase in

the S1 associated thalamic nuclei function (Petty et al., 2021) that is subsequently relayed to S1

and explains the sensitivity of SST-INs to locomotion in the dark but not in V1. To directly test

this hypothesis, we inactivated the VPM and POM with microinjections of the GABA(A) agonist

muscimol (0.5 mM) and monitored the effect in SST-INs during resting and active states. We

first validated that we were able to effectively inactivate the somatosensory thalamus. For that

we conducted *in vivo* calcium imaging experiments in S1 while monitoring activity of PNs as well

as SST-INs in response to whisker stimulation in control conditions and after muscimol

injections in the somatosensory thalamus (Figure 6). We observed that in control conditions

whisker stimulation induced a clear increase in the activity of both SST-INs and PNs (Figure 6A-

C and Figure S4). In contrast, after muscimol injection this increase was absent in both cell

types (Figure 6A-C and Figure S4). These results confirmed the effectiveness of muscimol in

reducing thalamic activity. Subsequently, we monitored the modulation of SST-INs in S1 by

whisking and locomotion in control conditions and after thalamus inactivation (Figure D-E).

Interestingly when we evaluated response of SST-INs to both whisking and locomotion there

was a clear modification of the response. In muscimol conditions there was a significant

decrease in the response of SST-INs during locomotion (control: $\Delta F/F_{0\ rest} = 0.53 \pm 0.01$; $\Delta F/F_0$

$_{run} = 1.02 \pm 0.03$; n = 1657; Run: $\Delta F/F_{0\ rest} = 0.53 \pm 0.02$; $\Delta F/F_{0\ run} = 0.76 \pm 0.04$; n = 541 cells in

8 mice, p<0.0001, Mann-Whitney U). This effect translated into a decreased LMI both at the

single cell level as well as when comparing average values across the difference tested mice

(LMI single cells: control = 0.18 ± 0.01, n = 1657, muscimol = 0.09 ± 0.01, n =541, <0.0001,

Mann-Whitney U; Figure 6H, I). The Pearson correlation between spontaneous activity and

running speed was also significantly reduced by thalamus inactivation (Pearson single cells:

control = 0.18 ± 0.01, n = 1657, muscimol = 0.14 ± 0.01, n =541, <0.01, Mann-Whitney U;

Figure 6I). Plotting variation in ΔF/F vs running speed revealed once more that in control

conditions SST-INs activity increased with running speed and that this increase is significantly

reduced after muscimol injections (Figure 6J). Overall, our results suggest that in S1,

modulation of SST-INs by spontaneous behaviors requires thalamic activity.

In S1 locomotion is associated with increased L2/3 pyramidal cell activity.

SST-INs are most likely not contacted directly by thalamic inputs (Sermet et al., 2019)

and receive information indirectly via their important connectivity with L2/3 PNs. We thus

monitored the *in vivo* spiking activity of L2/3 in S1 during behavior state transitions namely

running and whisking in control conditions and after muscimol injections in the somatosensory

thalamus (Figure 7). Our observations revealed a significant rise in spiking activity among L2/3

principal neurons (PNs) in S1 during the transition from rest to running (control: $\Delta F/F_{0\,rest}$ = 0.22 ±

0.01; $\Delta F/F_{0\,run}$ = 0.80 ± 0.03; n = 1918; p <0.0001, Mann-Whitney U; Figure 7B, C)). At the

population level, we noted that 44% of the cells exhibited a statistically significant increase in

activity, while only 20% displayed a decrease (Figure 7A-C). In contrast, after muscimol

injection in the thalamus this average increase in activity was completely absent (Figure 7E-G).

In fact, positive correlation between locomotion and GCaMP6s fluorescence traces was lost

upon thalamus inactivation (control: Pearson: 0.11 ± 0.01: n = 1918; run= -0.07 ± 0.0`; n = 371;

p <0.0001, Mann-Whitney U; Figure 7F, G). These results are in agreement with what has

recently been reported using electrophysiological approaches in the primary visual cortex

(Nestvogel and McCormick, 2022). Furthermore, the increase in average ΔF/F values during

running periods in pyramidal cells was also not present in cells where somatosensory thalamus

was inactivated by muscimol injection (control: $\Delta F/F_{0\,rest}$ = 0.27 ± 0.01; $\Delta F/F_{0\,run}$ = 0.22 ± 0.02; n =

371; p <0.0001, Mann-Whitney U; Figure 7F, G). Hence, the convergence of locomotion-

induced modulation and susceptibility to thalamic inactivation in L2/3 PNs, similar to what we

observed for SST-INs, further supports the hypothesis that SST-INs in S1 inherit the increased

activity during active behavioral states from L2/3 PNs.

Strongly heterogeneous glutamatergic inputs to SST enable PN-drive despite an overall

weak connection strength

Glutamatergic inputs from L2/3 PNs to SST-INs are extremely weak, displaying low initial

release probably that progressively increases with continuous activation due to robust synaptic

facilitation (Kapfer et al., 2007; Pala and Petersen, 2015; Rozov et al., 2001). How can such

weak synapses transmit the increase in activity regimes in L2/3 PNs associated with active

behavioral states to L2/3 SST-INs? To address this, we performed paired recording between

L2/3 PNs and SST-INs in S1 and evaluated the strength of the excitatory connections between

pairs of connected neurons (Figure 7H). We observed that the synaptic strength (amplitude of

first EPSC) was highly variable among connections with an overall reduced mean amplitude and

displaying pronounced short facilitation (mean = 32 ± 6 pA, median = 12.8 pA, CV = 137%, n =

50) (Figure 7I, J). Interestingly, we observed that a small fraction (32 %) of cells displayed large

initial current amplitudes (amplitude of first EPSC >30 pA). In a subset of connected pairs, we

managed to successfully establish a third connection and we compared the synaptic properties

between the two different presynaptic neurons into the same postsynaptic SST-IN (Figure 7K,

L). Unexpectedly we observed that there was a positive correlation between the amplitude of

the synaptic currents between the two presynaptic neurons. These results suggest that the

variability in synaptic strength is not randomly distributed across glutamatergic synaptic inputs

innervating SST-INs but is determined by the postsynaptic subtype of SST-IN (Figure 7M).

Overall, these results indicate that a subpopulation of SST-INs is expected to be particularly

sensitive to variations in the activity level of L2/3 PNs that occur during the behavioral

transitions from rest active states in S1 in line with our initial *in vivo* observations (see Figure 1).

469
470

Discussion:

Using in vivo two-photon calcium imaging in awake mice we observed that behavioral state transitions do not equally affect the activity of L2/3 SST-INs across primary sensory regions. Specifically, we noted that in dark conditions, locomotion led to an increased activity of SST-INs in the S1 but had no effect in V1. Interestingly, when visual stimulation was present, this difference was lost and transitions from rest to running now positively modulated SST-INs in both primary sensory regions. Variable recruitment of VIP-INs does not seem to be at the origin of such differential behavior-state dependent modulation of SST-INs between the two sensory regions. In addition, using a chemogenetic approach we revealed that VIP-INs mainly inhibit SST-INs during rest/run transitions and are not the source of positive modulation of SST-INs by locomotion. In contrast, we observed that inactivation of somatosensory thalamus strongly reduces the modulation of SST-INs in S1 by spontaneous behaviors. Altogether our results provide a detailed description of the neuronal mechanisms at the origin of the differential modulation of VIP- and SST-INs by behavior states in S1.

The responses of neurons in primary sensory neocortical regions to sensory stimuli are influenced by behavioral states including, arousal level, whisking, attention, and locomotion (Bennett et al., 2013; Iriki et al., 1996; McGinley et al., 2015b; Petersen and Crochet, 2013). Disinhibitory circuits have been recognized as key players in mediating the neuromodulatory effects of such behavioral state transitions in neocortical networks. Specifically, the activity of VIP- and SST-INs is believed to allow gain control by behavioral state in the neocortex (Fu et al., 2015, 2014). In fact, previous studies in V1 revealed that, in the dark, VIP-INs are recruited during locomotion while the activity of SST-INs remains unchanged (Dipoppa et al., 2018; Fu et al., 2014; Pakan et al., 2016). Interestingly, our findings indicate that in S1, SST-INs are strongly recruited by locomotion even in the absence of light. This result is unexpected given the inhibitory connection between VIP-INs and SST-INs, as well as the established role of VIP-INs

in inhibiting SST-INs during whisking in stationary animals (Gasselin et al., 2021; Gentet et al.,

2012; Lee et al., 2013; Pfeffer et al., 2013; Pi et al., 2013). Interestingly, when we selected from

our recording sessions periods of whisking with no associated locomotion, we reproduced

previous findings observed in non-locomoting mice. Namely, whisking resulted in an increase of

VIP-INs activity and a decrease in recruitment of SST-INs (Gentet et al., 2012; Muñoz et al.,

2017). Hence, our observations indicate that whisking during rest or during locomotion are

associated with distinct cortical states, with the latter resulting in significantly stronger

recruitment of SST-INs in S1.

Notwithstanding the multiple evidence that SST-INs are sensitive to behavioral state

transitions the neuronal mechanisms at the origin of such modulations are not obvious. In the

S1, whisking in stationary animals mainly inhibits SST-INs in layer 2/3 (Gentet et al., 2012;

Muñoz et al., 2017). Through the manipulation of VIP-INs function, Munoz and colleagues

provided compelling evidence for VIP-INs to be the source of SST-INs inhibition during whisking

in awake mice (Muñoz et al., 2017). Surprisingly, during such experiments, the authors

observed a reversal in the response of SST-INs. Inhibiting VIP-INs resulted in a transformation

of the modulation into a notable positive recruitment of SST-INs by whisking (Muñoz et al.,

2017). This finding highlights the pivotal role of VIP-INs in governing the activity of SST-INs

during whisking. However, the question that emerges concerns the source of excitation by

whisking in L2/3 SST-INs when VIP-INs are silenced. We hypothesized that the mechanisms

underlying such positive modulation are probably also at the origin of positive modulation by

locomotion that we now observed in S1. Interestingly, in L5 SST-INs, pharmacological block or

genetic removal of muscarinic receptors abolishes positive increase in SST-INs by whisking

indicating that cholinergic inputs are likely to participate in increased activity of SST-INs by

whisking (Muñoz et al., 2017). However, if a similar mechanism is present in L2/3 SST-INs is

unknown. At odds with a possible cholinergic modulation of L2/3 SST-INs during spontaneous

behaviours, Yaeger and colleagues (Yaeger et al., 2019), reported that in the V1 of adult mice

(P>56), SST-INs lose their sensitivity to cholinergic modulation. However, it is worth noting that

Yaeger et al. mainly reported changes at the suprathreshold level upon cholinergic receptor

activation. Consistent with the notion that cholinergic modulation of SST-INs is absent in adult

mice, theoretical analysis yielded a hypothesis proposing that VIP-INs, rather paradoxically,

may be responsible for the increase in SST-INs, a phenomenon termed "response reversal,"

achieved through the disinhibition of local PNs (Garcia Del Molino et al., 2017).

Our observations reveal that inhibiting VIP-INs does not abolish the positive modulation

of SST-INs by locomotion; instead, it has the opposite effect, increasing the sensitivity of SST-

INs to transitions from rest to running. This suggests that positive recruitment of SST-INs

involves mechanisms independent of VIP-INs. Furthermore, we noted that despite relatively

similar recruitment of VIP-INs across sensory regions, SST-INs in S1 and V1 displayed different

response profiles to locomotion particularly in dark conditions. This result further argues that

VIP-INs are likely not the cause of SST-INs increase during locomotion in the neocortex. In

contrast, our experimental results illustrate that in the presence of visual stimulation modulation

of SST-INs by locomotion is altered in V1 but not in S1 suggesting that sensory region-specific

mechanisms are involved in such modulation.

Based on our current findings, we propose a new working model in which the positive

modulation of SST-INs activity across behavioral states relies on the interaction between

thalamic activity and neuromodulatory pathways, specifically cholinergic inputs. Our hypothesis

is strongly supported by our data, which demonstrates that inactivation of the thalamus

significantly diminishes the modulation of SST-INs in S1 by locomotion. We propose that the

involvement of thalamic activity partially explains the region-specific effects of spontaneous

behaviors on SST-INs due to different levels of ongoing sensory drive. Activity of

thalamocortical VPM neurons, that relay somatosensory information to S1, increases with

whisking (Gutnisky et al., 2017; Poulet et al., 2012; Urbain et al., 2015). While comparatively

less explored, recent findings suggest a notable upsurge in VPM activity also during locomotion

(Huo et al., 2020). For the moment is unclear if this representation of self-motion in VPM

neurons is due to cortical inputs (Poulet et al., 2012), neuromodulation (Petty et al., 2021) or

proprioception (Severson et al., 2017).   Nonetheless, we anticipate that the combined effects of

whisking and locomotion lead to an overall increase in thalamic activity, which is subsequently

transmitted to the neocortex. Interestingly, our findings align with recent research indicating that

depolarization of L2/3 principal neurons (PNs) in V1 during locomotion is dependent on thalamic

function (Nestvogel and McCormick, 2022). In fact, we obtained similar results in our study

(Figure 6). Under normal conditions, locomotion leads to an overall increase in activity (positive

LMI) in L2/3 PNs (Figure 7), and this increase is entirely abolished when thalamic activity is

disrupted. As SST-INs are highly contacted by L2/3 PNs we suggest that the increase in SST-

INs is inherited from L2/3 PNs. Worth noting that in V1 the differential response of SST-INs to

locomotion is also observed in L2/3 PN whose LMI is significantly larger in the present of visual

stimulation (Pakan et al., 2016), compatible with the idea that SST-INs display behavior

dependent modulation in part transmitted from local L2/3 PNs.

We also observed significant variability at the single-cell level regarding the sensitivity of

S1 L2/3 SST-INs to behavioral state transitions (see Figure 1 and 2). This variability was

followed by a pronounced variance in the strength of unitary connections between PNs and

SST-INs (Figure 7). Previous proposals have suggested that skewed distributions of excitatory

postsynaptic potential (EPSP) amplitudes in SST-INs, with a relatively small mean but a skew

towards larger amplitudes, ensured individual PNs to excite a subset of SST-INs (Kapfer et al.,

2007). However, our findings indicate that these distributions are not entirely random among

SST-INs and that the strength of these connections is influenced by the postsynaptic cell.

Recent transcriptomic data further suggest the presence of at least three distinct subtypes of

SST-INs in S1 L2/3 (Wu et al., 2023). Remarkably the expression of the Elfn1, which is known

to retrogradely regulate release dynamics at glutamatergic inputs, exhibits differential

expression patterns among SST-INs subtypes (Lim et al., 2018; Paul et al., 2017). This leads us

to speculate that the properties of glutamatergic inputs are not uniform across the different SST-

IN subtypes in L2/3, potentially conferring unique computational properties to neocortical

circuits. In the future, it would be interesting to directly investigate whether *in vivo* SST-INs

exhibiting greater sensitivity to locomotion correspond to those with stronger unitary connections

between PNs and SST-INs that we recorded *in vitro*. Subpopulations of SST-INs, with respect to

their response during rest-run transitions, were previously documented in V1 (Reimer et al.,

2014). However, it remained uncertain whether these responses genuinely corresponded to

distinct subpopulations of SSTs or if they were merely a result of transgenic mouse models

used to label SST-INs (Hu et al., 2013). In a more recent development, a study that employed a

genetic approach to identify subpopulations of SST-INs unveiled distinct functional

specializations within these discrete subsets of SST-INs (Green et al., 2023), further supporting

the idea of the existence of SST-IN subgroups.

During our thalamic muscimol injections, we observed a complete block of the positive

modulation of PNs by locomotion while it was only reduced in SST-INs. Since the experiments

on PNs and SST-INs were not conducted simultaneously, it remains possible that experimental

differences, such as incomplete thalamic inactivation during SST-INs imaging sessions, could

account for the disparate results despite ensuring successful targeting and delivery of muscimol

to the somatosensory thalamus. However, our in vitro results indicate that SST-INs are

equipped with cholinergic receptors. Thus, it is still possible that the remaining modulation of

SST-INs is mediated by cholinergic neuromodulation that remained active during our thalamus

inactivation experiment. Such a possible scenario would indicate that both thalamic and

cholinergic pathways are essential in shaping the responses of SST-INs in S1 across the

24

different behavioral states. Nonetheless, future work involving the selective removal of muscarinic receptors from SST-INs is needed to allow definite claims about the in vivo contribution of cholinergic pathways in the control of L2/3 SST-INs during spontaneous behaviors.

Also evident was the observation that SST- and VIP-INs are equipped with different types of cholinergic receptors. VIP-INs were particularly rich in nicotinic receptors which were almost absent in SST-INs that seemed to express mainly muscarinic receptors. This differential equipment of cholinergic receptors is expected to result also in differential sensitivity to behavioral state transitions. In particular the high levels of nicotinic levels in VIP-INs are compatible with their fast activation to even small behavioral state variations like fast whisking bouts (Gasselin et al., 2021). In contrast for SST-INs, due to mainly muscarinic receptors, longer and more sustained activation of cholinergic inputs activation is required to mediate depolarizations in SST-INs.

Mechanistically it is now clear that both thalamus as well as neuromodulatory inputs like cholinergic projections originating from the basal forebrain participate in the modulation of cortical networks by behavioral state. Less obvious is the computational role of the two different pathways. The cholinergic pathway, via VIP-INs activation, has been proposed to underlie the gain control of L2/3 PNs to sensory evoked activity across behavioral states (Fu et al., 2014). Thalamic activity is required for the important depolarizations of L2/3 neurons (Nestvogel and McCormick, 2022) during locomotion but their precise computational role is still ill defined. In any case by integrating feedforward thalamic activity transmitted indirectly via L2/3 PNs as well as neuromodulatory information both via VIP-mediated inhibition and via direct cholinergic innervation, SST-INs occupy a key position in cortical networks to adjust activity of the two pathways and adjust neuronal activity in function of behavior state.

References:

Alitto, H., Dan, Y., 2013. Cell-type-specific modulation of neocortical activity by basal forebrain input. Front. Syst. Neurosci. 6.

Ayaz, A., Saleem, A.B., Schölvinck, M.L., Carandini, M., 2013. Locomotion controls spatial integration in mouse visual cortex. Curr. Biol. CB 23, 890–894. https://doi.org/10.1016/j.cub.2013.04.012

Bennett, C., Arroyo, S., Hestrin, S., 2013. Subthreshold Mechanisms Underlying State-Dependent Modulation of Visual Responses. Neuron 80, 350–357. https://doi.org/10.1016/j.neuron.2013.08.007

Bugeon, S., Duffield, J., Dipoppa, M., Ritoux, A., Prankerd, I., Nicoloutsopoulos, D., Orme, D., Shinn, M., Peng, H., Forrest, H., Viduolyte, A., Reddy, C.B., Isogai, Y., Carandini, M., Harris, K.D., 2022. A transcriptomic axis predicts state modulation of cortical interneurons. Nature 607, 330–338. https://doi.org/10.1038/s41586-022-04915-7

Busse, L., Cardin, J.A., Chiappe, M.E., Halassa, M.M., McGinley, M.J., Yamashita, T., Saleem, A.B., 2017. Sensation during Active Behaviors. J. Neurosci. 37, 10826–10834. https://doi.org/10.1523/JNEUROSCI.1828-17.2017

Dipoppa, M., Ranson, A., Krumin, M., Pachitariu, M., Carandini, M., Harris, K.D., 2018. Vision and Locomotion Shape the Interactions between Neuron Types in Mouse Visual Cortex. Neuron 98, 602-615.e8. https://doi.org/10.1016/j.neuron.2018.03.037

Fu, Y., Kaneko, M., Tang, Y., Alvarez-Buylla, A., Stryker, M.P., 2015. A cortical disinhibitory circuit for enhancing adult plasticity. eLife 4, e05558. https://doi.org/10.7554/eLife.05558

Fu, Y., Tucciarone, J.M., Espinosa, J.S., Sheng, N., Darcy, D.P., Nicoll, R.A., Huang, Z.J., Stryker, M.P., 2014. A cortical circuit for gain control by behavioral state. Cell 156, 1139–1152. https://doi.org/10.1016/j.cell.2014.01.050

Garcia Del Molino, L.C., Yang, G.R., Mejias, J.F., Wang, X.-J., 2017. Paradoxical response reversal of top-down modulation in cortical circuits with three interneuron types. eLife 6, e29742. https://doi.org/10.7554/eLife.29742

Gasselin, C., Hohl, B., Vernet, A., Crochet, S., Petersen, C.C.H., 2021. Cell-type-specific nicotinic input disinhibits mouse barrel cortex during active sensing. Neuron 109, 778-787.e3. https://doi.org/10.1016/j.neuron.2020.12.018

Gentet, L.J., Kremer, Y., Taniguchi, H., Huang, Z.J., Staiger, J.F., Petersen, C.C.H., 2012. Unique functional properties of somatostatin-expressing GABAergic neurons in mouse barrel cortex. Nat. Neurosci. 15, 607–612. https://doi.org/10.1038/nn.3051

Gouwens, N.W., Sorensen, S.A., Baftizadeh, F., Budzillo, A., Lee, B.R., Jarsky, T., Alfiler, L., Baker, K., Barkan, E., Berry, K., Bertagnolli, D., Bickley, K., Bomben, J., Braun, T., Brouner, K., Casper, T., Crichton, K., Daigle, T.L., Dalley, R., Frates, R.A. de, Dee, N., Desta, T., Lee, S.D., Dotson, N., Egdorf, T., Ellingwood, L., Enstrom, R., Esposito, L., Farrell, C., Feng, D., Fong, O., Gala, R., Gamlin, C., Gary, A., Glandon, A., Goldy, J., Gorham, M., Graybuck, L., Gu, H., Hadley, K., Hawrylycz, M.J., Henry, A.M., Hill, D., Hupp, M., Kebede, S., Kim, T.K., Kim, L., Kroll, M., Lee, C., Link, K.E., Mallory, M., Mann, R., Maxwell, M., McGraw, M., McMillen, D., Mukora, A., Ng, Lindsay, Ng, Lydia, Ngo, K., Nicovich, P.R., Oldre, A., Park, D., Peng, H., Penn, O., Pham, T., Pom, A., Popović, Z., Potekhina, L., Rajanbabu, R., Ransford, S., Reid, D., Rimorin, C., Robertson, M., Ronellenfitch, K., Ruiz, A., Sandman, D., Smith, K., Sulc, J., Sunkin, S.M., Szafer, A., Tieu, M., Torkelson, A., Trinh, J., Tung, H., Wakeman, W., Ward, K., Williams, G., Zhou, Z., Ting, J.T., Arkhipov, A., Sümbül, U., Lein, E.S., Koch, C., Yao, Z., Tasic, B., Berg, J., Murphy, G.J., Zeng, H., 2020. Integrated Morphoelectric and Transcriptomic Classification of Cortical GABAergic Cells. Cell 183, 935-953.e19. https://doi.org/10.1016/j.cell.2020.09.057

Green, J., Bruno, C.A., Traunmüller, L., Ding, J., Hrvatin, S., Wilson, D.E., Khodadad, T., Samuels, J., Greenberg, M.E., Harvey, C.D., 2023. A cell-type-specific error-correction signal in the posterior parietal cortex. Nature 620, 366–373. https://doi.org/10.1038/s41586-023-06357-1

Gutnisky, D.A., Yu, J., Hires, S.A., To, M.-S., Bale, M.R., Svoboda, K., Golomb, D., 2017. Mechanisms underlying a thalamocortical transformation during active tactile sensation. PLOS Comput. Biol. 13, e1005576. https://doi.org/10.1371/journal.pcbi.1005576

Hu, H., Cavendish, J., Agmon, A., 2013. Not all that glitters is gold: off-target recombination in the somatostatin–IRES-Cre mouse line labels a subset of fast-spiking interneurons. Front. Neural Circuits 7.

Huo, Y., Chen, H., Guo, Z.V., 2020. Mapping Functional Connectivity from the Dorsal Cortex to the Thalamus. Neuron 107, 1080-1094.e5. https://doi.org/10.1016/j.neuron.2020.06.038

Iriki, A., Tanaka, M., Iwamura, Y., 1996. Attention-induced neuronal activity in the monkey somatosensory cortex revealed by pupillometrics. Neurosci. Res. 25, 173–181. https://doi.org/10.1016/0168-0102(96)01043-7

Kapfer, C., Glickfeld, L.L., Atallah, B.V., Scanziani, M., 2007. Supralinear increase of recurrent inhibition during sparse activity in the somatosensory cortex. Nat. Neurosci. 10, 743–753. https://doi.org/10.1038/nn1909

Lee, S., Kruglikov, I., Huang, Z.J., Fishell, G., Rudy, B., 2013. A disinhibitory circuit mediates motor integration in the somatosensory cortex. Nat. Neurosci. 16, 1662–1670. https://doi.org/10.1038/nn.3544

Lim, L., Pakan, J.M.P., Selten, M.M., Marques-Smith, A., Llorca, A., Bae, S.E., Rochefort, N.L., Marín, O., 2018. Optimization of interneuron function by direct coupling of cell migration and axonal targeting. Nat. Neurosci. 21, 920–931. https://doi.org/10.1038/s41593-018-0162-9

McCormick, D.A., Nestvogel, D.B., He, B.J., 2020. Neuromodulation of Brain State and Behavior. Annu. Rev. Neurosci. 43, 391–415. https://doi.org/10.1146/annurev-neuro-100219-105424

McGinley, M.J., David, S.V., McCormick, D.A., 2015a. Cortical Membrane Potential Signature of Optimal States for Sensory Signal Detection. Neuron 87, 179–192. https://doi.org/10.1016/j.neuron.2015.05.038

McGinley, M.J., Vinck, M., Reimer, J., Batista-Brito, R., Zagha, E., Cadwell, C.R., Tolias, A.S., Cardin, J.A., McCormick, D.A., 2015b. Waking State: Rapid Variations Modulate Neural and Behavioral Responses. Neuron 87, 1143–1161. https://doi.org/10.1016/j.neuron.2015.09.012

Mitchinson, B., Martin, C.J., Grant, R.A., Prescott, T.J., 2007. Feedback control in active sensing: rat exploratory whisking is modulated by environmental contact. Proc. R. Soc. B Biol. Sci. https://doi.org/10.1098/rspb.2006.0347

Muñoz, W., Tremblay, R., Levenstein, D., Rudy, B., 2017. Layer-specific modulation of neocortical dendritic inhibition during active wakefulness. Science 355, 954–959. https://doi.org/10.1126/science.aag2599

Nestvogel, D.B., McCormick, D.A., 2022. Visual thalamocortical mechanisms of waking state-dependent activity and alpha oscillations. Neuron 110, 120-138.e4. https://doi.org/10.1016/j.neuron.2021.10.005

Niell, C.M., Stryker, M.P., 2010. Modulation of visual responses by behavioral state in mouse visual cortex. Neuron 65, 472–479. https://doi.org/10.1016/j.neuron.2010.01.033

Pakan, J.M., Lowe, S.C., Dylda, E., Keemink, S.W., Currie, S.P., Coutts, C.A., Rochefort, N.L., 2016. Behavioral-state modulation of inhibition is context-dependent and cell type specific in mouse visual cortex. eLife 5, e14985. https://doi.org/10.7554/eLife.14985

Pala, A., Petersen, C.C.H., 2015. In Vivo Measurement of Cell-Type-Specific Synaptic
Connectivity and Synaptic Transmission in Layer 2/3 Mouse Barrel Cortex. Neuron 85,
68–75. https://doi.org/10.1016/j.neuron.2014.11.025
Paul, A., Crow, M., Raudales, R., He, M., Gillis, J., Huang, Z.J., 2017. Transcriptional
Architecture of Synaptic Communication Delineates GABAergic Neuron Identity. Cell
171, 522-539.e20. https://doi.org/10.1016/j.cell.2017.08.032
Petersen, C.C.H., Crochet, S., 2013. Synaptic Computation and Sensory Processing in
Neocortical Layer 2/3. Neuron 78, 28–48. https://doi.org/10.1016/j.neuron.2013.03.020
Petty, G.H., Kinnischtzke, A.K., Hong, Y.K., Bruno, R.M., 2021. Effects of arousal and
movement on secondary somatosensory and visual thalamus. eLife 10, e67611.
https://doi.org/10.7554/eLife.67611
Pfeffer, C.K., Xue, M., He, M., Huang, Z.J., Scanziani, M., 2013. Inhibition of inhibition in visual
cortex: the logic of connections between molecularly distinct interneurons. Nat. Neurosci.
16, 1068–1076. https://doi.org/10.1038/nn.3446
Pi, H.-J., Hangya, B., Kvitsiani, D., Sanders, J.I., Huang, Z.J., Kepecs, A., 2013. Cortical
interneurons that specialize in disinhibitory control. Nature 503, 521–524.
https://doi.org/10.1038/nature12676
Polack, P.-O., Friedman, J., Golshani, P., 2013. Cellular mechanisms of brain state–dependent
gain modulation in visual cortex. Nat. Neurosci. 16, 1331–1339.
https://doi.org/10.1038/nn.3464
Poulet, J.F.A., Fernandez, L.M.J., Crochet, S., Petersen, C.C.H., 2012. Thalamic control of
cortical states. Nat. Neurosci. 15, 370–372. https://doi.org/10.1038/nn.3035
Reimer, J., Froudarakis, E., Cadwell, C.R., Yatsenko, D., Denfield, G.H., Tolias, A.S., 2014.
Pupil Fluctuations Track Fast Switching of Cortical States during Quiet Wakefulness.
Neuron 84, 355–362. https://doi.org/10.1016/j.neuron.2014.09.033
Rozov, A., Burnashev, N., Sakmann, B., Neher, E., 2001. Transmitter release modulation by
intracellular Ca2+ buffers in facilitating and depressing nerve terminals of pyramidal cells
in layer 2/3 of the rat neocortex indicates a target cell-specific difference in presynaptic
calcium dynamics. J. Physiol. 531, 807–826. https://doi.org/10.1111/j.1469-
7793.2001.0807h.x
Saleem, A.B., Ayaz, A., Jeffery, K.J., Harris, K.D., Carandini, M., 2013. Integration of visual
motion and locomotion in mouse visual cortex. Nat. Neurosci. 16, 1864–1869.
https://doi.org/10.1038/nn.3567
Schneider, D.M., Nelson, A., Mooney, R., 2014. A synaptic and circuit basis for corollary
discharge in the auditory cortex. Nature 513, 189–194.
https://doi.org/10.1038/nature13724
Sermet, B.S., Truschow, P., Feyerabend, M., Mayrhofer, J.M., Oram, T.B., Yizhar, O., Staiger,
J.F., Petersen, C.C., 2019. Pathway-, layer- and cell-type-specific thalamic input to
mouse barrel cortex. eLife 8, e52665. https://doi.org/10.7554/eLife.52665
Severson, K.S., Xu, D., Van de Loo, M., Bai, L., Ginty, D.D., O'Connor, D.H., 2017. Active touch
and self-motion encoding by Merkel cell-associated afferents. Neuron 94, 666-676.e9.
https://doi.org/10.1016/j.neuron.2017.03.045
Shimaoka, D., Harris, K.D., Carandini, M., 2018. Effects of Arousal on Mouse Sensory Cortex
Depend on Modality. Cell Rep. 22, 3160–3167.
https://doi.org/10.1016/j.celrep.2018.02.092
Sofroniew, N.J., Cohen, J.D., Lee, A.K., Svoboda, K., 2014. Natural Whisker-Guided Behavior
by Head-Fixed Mice in Tactile Virtual Reality. J. Neurosci. 34, 9537–9550.
https://doi.org/10.1523/JNEUROSCI.0712-14.2014
Syeda, A., Zhong, L., Tung, R., Long, W., Pachitariu, M., Stringer, C., 2022. Facemap: a
framework for modeling neural activity based on orofacial tracking.
https://doi.org/10.1101/2022.11.03.515121

Urbain, N., Salin, P.A., Libourel, P.-A., Comte, J.-C., Gentet, L.J., Petersen, C.C.H., 2015.
Whisking-Related Changes in Neuronal Firing and Membrane Potential Dynamics in the
Somatosensory Thalamus of Awake Mice. Cell Rep. 13, 647–656.
https://doi.org/10.1016/j.celrep.2015.09.029
Vinck, M., Batista-Brito, R., Knoblich, U., Cardin, J.A., 2015. Arousal and Locomotion Make
Distinct Contributions to Cortical Activity Patterns and Visual Encoding. Neuron 86, 740–
754. https://doi.org/10.1016/j.neuron.2015.03.028
Wu, S.J., Sevier, E., Dwivedi, D., Saldi, G.-A., Hairston, A., Yu, S., Abbott, L., Choi, D.H.,
Sherer, M., Qiu, Y., Shinde, A., Lenahan, M., Rizzo, D., Xu, Q., Barrera, I., Kumar, V.,
Marrero, G., Prönneke, A., Huang, S., Kullander, K., Stafford, D.A., Macosko, E., Chen,
F., Rudy, B., Fishell, G., 2023. Cortical somatostatin interneuron subtypes form cell-type-
specific circuits. Neuron 0. https://doi.org/10.1016/j.neuron.2023.05.032
Yaeger, C.E., Ringach, D.L., Trachtenberg, J.T., 2019. Neuromodulatory control of localized
dendritic spiking in critical period cortex. Nature 567, 100–104.
https://doi.org/10.1038/s41586-019-0963-3
Yu, J., Hu, H., Agmon, A., Svoboda, K., 2019. Recruitment of GABAergic Interneurons in the
Barrel Cortex during Active Tactile Behavior. Neuron 104, 412-427.e4.
https://doi.org/10.1016/j.neuron.2019.07.027
Zhou, M., Liang, F., Xiong, X.R., Li, L., Li, H., Xiao, Z., Tao, H.W., Zhang, L.I., 2014. Scaling
down of balanced excitation and inhibition by active behavioral states in auditory cortex.
Nat. Neurosci. 17, 841–850. https://doi.org/10.1038/nn.3701
793
794

Figure Legends

Figure 1:   SST-INs in V1 and S1 are differentially modulated by behavioral state
transitions. (A) Two-photon (2P) calcium imaging in S1 of awake mice. Mice were head fixed
and were free to run on a circular treadmill. Whisking activity was tracked via a video camera
detecting infra-red light focused on mouse snout illuminated with 850 nm LED. Bottom: Two-
photon laser scanning microscopy (2PLSM) image (maximum-intensity projection, MIP) of SST-
Cre positive neurons labelled with GCaMP6s in S1. Scale bar is 100 µm **(B)** Representative
GCaMP6s fluorescence traces (black) shown as $\Delta F/F_0$ (up) as well as normalized $\Delta F/F_0$
(bottom) for SST-INs displayed in A. Each row represents a neuron sorted by weight on the first
principal component (PC) of their activity. Locomotion trace is shown on top (red). **(C)** Mean
$\Delta F/F_0$ during periods of rest and run ($\Delta F/F_{0\ rest} = 0.80 \pm 0.02$; $\Delta F/F_{0\ run} = 1.84 \pm 0.06$; n = 2030
cells in 16 mice, p<0.0001, Mann-Whitney U test). **(D)** Summary plot of Pearson correlation
coefficients between calcium activity and running speed. SST-INs showing statistically
significant (see methods) modulation by locomotion are represented by colored bars, with light
orange indicating negative modulation and positive modulation denoted by brown. The pie chart
provides an overview of the overall percentage of SST-INs showing positive, negative, or no
effect in response to locomotion. **(E)** Same as B but for SST-INs in the primary visual cortex. (**F,**

G) Same as C and D but for V1. $\Delta F/F_{0\ rest}$ = 0.70 ± 0.03; $\Delta F/F_{0\ run}$ = 0.84 ± 0.03; n = 881 cells in
10 mice, p<0.0001, Mann-Whitney U. **(H)** Event-triggered average traces of SST-INs GCaMP6s
signal in S1 aligned to the start of running (left) or whisking (right) along with running speed
(red) and whisking (blue). Normalized $\Delta F/F$ is displayed for all analyzed cells (bottom). Traces
represent mean ± SEM values across all recorded cells. **(I)** Same as G but for V1. **(J)** Summary
plot of variation in $\Delta F/F$ between rest and run and rest and whisking for the event-triggered
average traces shown in G and H. Values were obtained by averaging $\Delta F/F$ trace for each
individual SST-INs in a 2 s window before and after event onset (see Methods). $\Delta F/F_{0\ Loco\ S1}$ =
0.40 ± 0.03, n = 2030; $\Delta F/F_{0\ Loco\ V1}$ = 0.03 ± 0.02, n= 633, p<0.0001, Mann-Whitney U; $\Delta F/F_{0}$
$_{Whisk\ S1}$ = 0.08 ± 0.01, n = 853, n = 2030, $\Delta F/F_{0\ Whisk\ V1}$ = 0.002 ± 0.007; n = 853 cells in 10 mice,
p<0.0001, Mann-Whitney U. **(K)** Cross-correlation between in vivo calcium activity and running
speed for SST-INs in S1 and V1 (left). (right) Average correlation coefficient values at zero-time
for locomotion and whisking for SST-INs recorded in V1 and S1 (ρloco S1 = 0.22 ± 0.01, n =
2030, ρloco V1 = 0.06 ± 0.01, n = 883, p<0.0001, Mann-Whitney U; ρwhisk S1 = 0.15 ± 0.01, n
= 2030, ρloco V1 = 0.05 ± 0.01, n = 883, p<0.0001, Mann-Whitney U).
828

829
830
Figure 2: Recruitment of SST-INs in S1 is primarily associated with behavioral state
transitions characterized by increased locomotor activity. (A) Example session illustrating
simultaneous recording traces for whisking (blue) and running speed (red). Whisking only
periods (no locomotor activity) are highlighted in light green. **(B)** Quantification of the fraction of
time mice spent whisking, wishing only and running during imaging sessions. **(C)** Event-
triggered average traces of SST-INs GCaMP6s fluorescence signal in S1 aligned to whisking
onset. Only whisking bouts not associated with locomotion were used (whisking only).
Normalized $\Delta F/F$ is displayed for all analyzed cells (bottom). Traces represent mean ± SEM
values across all recorded cells. **(D)** Enlarged view of average traces shown in C. **(E)** Summary
plot of variation in $\Delta F/F$ between rest and whisking for the event-triggered average traces shown
in C and panel H figure1. Values were obtained by averaging $\Delta F/F$ traces for each individual
SST-INs in a 2 s window before and after event onset. Note that during whisking only, there is
either no effect or a slight tendency to inhibit SST-INs. However, when whisking is associated
with running periods, it clearly increases the activity of SST-INs. $\Delta F/F_{0\ Whisk} = 0.08 \pm 0.01$, n =

853; $\Delta F/F_{0 \text{ Whisk only}}$ = -0.01 ± 0.01; n = 853 cells in 10 mice, p<0.001, Mann-Whitney U. **(F)**
Summary plot of average mean locomotion, whisking and whisking only modulation indices (LMI
= 0.26 ± 0.01, n=2030; WMI = 0.21 ± 0.01, n=856; WOMI = 0.06 ± 0.01, n=830) **(G)** Event-
triggered average traces of SST-INs GCaMP6s signal ($\Delta F/F$) in S1 aligned to whisking onset.
Results were separated according to percentage time mice spent whisking only during the
imaging session. Average of GCaMP6s fluorescence from sessions with whisking only periods
longer than 20% of the recording time is shown on the left and less than 20% is illustrated on
the right. Traces represent mean ± SEM values across all recorded cells. **(H)** Summary plot of
variation in $\Delta F/F$ between rest and whisking for the event-triggered average traces shown in G.
Average values were obtained 2 s before and 2 s after event whisking onset (see Methods).
Values were calculated for each individual SST-INs. ($\Delta F/F_{0 \text{ Whisk<20%}}$ = 0.11 ± 0.02, n = 602;
$\Delta F/F_{0 \text{ Whisk<20%}}$ = -0.02 ± 0.01, n = 251, p<0.0001, Mann-Whitney U). **(I)** Histograms of the
distribution of locomotion modulation indices (LMI = $(R_L - R_S)/(R_L + R_S)$, where R_L and R_S are
the mean $\Delta F/F0$ during locomotion and stationary periods, respectively), for SST-INs that
displayed negative or positive WOMI.
860

Figure 3: VIP interneurons are strongly modulated by locomotion in S1. A) *(top)* Illustration
of a mouse brain depicting the location of S1 and V1. *(bottom)* 2PLSM image (maximum-
intensity projection, MIP) of VIP-Cre positive neurons labeled with GCaMP6f in S1. **B)**

Representative traces of in vivo calcium (GCaMP6f) activity in VIP-INs. Locomotion trace is
shown on top (red) followed by the calcium activity (green). Deconvolved activity (in black) is
shown under each trace **C)** Same as B but for VIP-INs in V1. **D)** Mean $\Delta F/F_0$ during periods of
rest and run, ($\Delta F/F_{0\ rest}$ = 0.80 ± 0.02; $\Delta F/F_{0\ run}$ = 2.23 ± 0.05; n = 1915 cells in 11 mice,
p<0.0001, Mann-Whitney U test). **E)** Histogram of locomotion modulation indices for VIP-INs in
S1. Inset depicts the average LMI value (LMI = 0.38 ± 0.01, n= 1915). **F)** Distribution of
Pearson correlation coefficients between calcium activity and locomotion trace for VIP-INs in
S1. Colored bars indicate VIP-INs that are statistically significantly modulated by locomotion,
either in a negative (orange) or positive manner. The pie chart provides an overview of the
overall percentage of VIP-INs showing positive, negative, or no effect in response to locomotion**.**
G, H , I) Same as D, E, F but for VIP-INs in V1. **J)** Illustration of the double transgenic mouse
line used for targeted protein expression in VIP- and SST-INs respectively. Adeno-associated
virus (AAV) coding Cre-dependent expression of Channelrhodopsin2-EGFP along with a
second AAV coding FLP-dependent expression of mCherry were infected in primary visual (V1)
and primary sensory (S1) cortices of VIP-Cre::SST-Flp mice. Schematic diagram of
experimental approach where VIP-INs were photo-activated via ChR2 while a PN and
neighboring SST-IN were patched in L2/3. **K)** *(left)* Example traces of IPSCs simultaneously
recorded in PNs and SST in V1 and S. *(middle)* Scatter plot of IPSCs amplitudes recorded in
SST-INs and PNs in S1 and V1 following VIP-INs photostimulation. (*right*) Ratio of light-evoked
IPSCs in V1 and S1 (V1: n = 7 cells, 2 mice ; S1: n = 10 cells, 2 mice). Mann Whitney test
p=0.8125. Error bars represent SEM.
886

887
Figure 4:VIP-INs mainly inhibit SST-INs during spontaneous behaviors. A) Simplified
illustration of the double transgenic mouse line used to target protein expression in VIP- and
SST-INs respectively based on the Cre/Lox and FLP/FRT system. Adeno-associated viruses
(AAV) were used to express in a Cre-dependent manner G_i-coupled inhibitory (hM4Di) designer
receptors exclusively activated by designer drugs (DREADD) in VIP-INs. A second AAV was
used to drive in a FLP-dependent manner the expression of GCaMP6s in SST-INs. **B)**
Schematic illustration of the canonical cellular organization in S1 and the expected negative
modulation of VIP-INs via activation of hM4Di by CNO injection. **C)** Mean $\Delta F/F_0$ during periods
of rest and run in control conditions and 30 after CNO injection (left), ($\Delta F/F_{0\ rest,\ control}$ = 0.71 ±
0.02; $\Delta F/F_{0\ run,\ control}$ = 1.5 ± 0.07; n = 393 cells in 5 mice, p<0.0001, Mann-Whitney U test; $\Delta F/F_0$
$_{rest,\ CNO}$ = 0.61 ± 0.04; $\Delta F/F_{0\ run,\ CNO}$ = 1.81 ± 0.08; n = 308 cells in 5 mice, p<0.0001, Mann-
Whitney U test).   Summary plot of LMI values obtained for SST-INs in control and after CNO
injection (right), (LMI control = 0.30 ± 0.02, n = 393; LMI CNO = 0.4 ± 0.02, n = 308; p<0.001,
Mann-Whitney U test). **D)** Variation in Pearson correlation values between running speed and
GCaMP6s fluorescence signals for SST-INs recorded in control and CNO treated mice (ρloco
control = 0.30 ± 0.02, n = 393, ρloco CNO = 0.44 ± 0.02, n = 308, p<0.0001, Mann-Whitney U).
Percentage of SST-INs exhibiting statistically significant positive, negative, or non-significant

Pearson correlation values is shown on the right. **E)** Event-triggered average traces of SST-INs
GCaMP6s signals in S1 aligned to the onset of whisking along with running speed (red) and
whisking (blue). Normalized ∆F/F is displayed for all analyzed cells (bottom). Results are
displayed for control conditions (left) and after CNO injection (right). Traces represent mean ±
SEM values across all recorded cells. **F)** Enlarged view of average traces shown in E. **G)**
Summary plot of variation in ∆F/F between rest and whisking for the event-triggered average
traces shown in F. Average values were obtained 2 s before and 2 s after event whisking onset
(see Methods). Values were calculated for each individual SST-INs. Note that in control
conditions whisking induces a brief decrease in GCaMP6s fluorescence that is absent in CNO
treated mice. **H and F)** Variation in ∆F/F values for SST-INs with running speed for control and
CNO conditions.
916

917
918
Figure 5: Visual stimulation modulates the recruitment of SST-INs by locomotion in V1
but not in S1. **A)** Schematic of the experimental setup used to probe the effect of visual
stimulation in the behaviour state-dependent modulation of the activity of VIP and SST-INs in V1
and S1. **B)** Summary plots of LMI and Pearson correlation between running speed and
GCaMP6 fluorescence signal for VIP- and SST-INs in the dark and facing a grey screen. LMI =
0.23 ± 0.01, n =505, for VIP in the dark versus 0.21 ± 0.02, n = 222, in light; p > 0.9999, and
0.08 ± 0.01, n = 881 for SST in the dark versus 0.31 ± 0.02, n = 116 in light, p < 0.0001.
Pearson Correlation: 0.21 ± 0.01, n = 505, for VIP in the dark versus 0.20 ± 0.02, n = 222, in
light; 0.06 ± 0.01, n = 881, for SST in the dark versus versus 0.31 ± 0.02, n = 116 for SST-INs in
light, p < 0.0001, Mann-Whitney U test. **C)** Same as B but for S1. Values represent mean ±
SEM.
930

931
Figure 6: Thalamic activity is required for the modulation of SST-INs by locomotion in S1.
A) Schematic illustration of experimental approach used to monitor effectiveness of thalamus
inactivation using muscimol injection in somatosensory thalamus. Whisker deflection was
achieved using localized air puff while cortical activity in S1 was monitored using *in vivo* two-
photon calcium imaging. **B)** Event-triggered average traces of SST-INs GCaMP6s signals in S1
aligned to whisker puff along with running speed (red). Normalized ΔF/F is displayed for all
analyzed cells (bottom). Results are displayed for control conditions (left) and for mice
previously injected with muscimol in the thalamus (*right*). Traces represent mean ± SEM values
across all recorded cells. **C)** Summary plot of variation in ΔF/F between rest and after whisker
stimulation for the event-triggered average traces shown in B. Average values were obtained 2
s before and 2 s after whisker stimulation onset (see Methods). **D)** Representative GCaMP6s

fluorescence traces (black) shown as $\Delta F/F_0$ (up) as well as normalized $\Delta F/F_0$ (bottom) for SST-
INs in S1 for mice spontaneously transitioning between rest and active behavior periods. Each
row represents a neuron sorted by weight on the first principal component (PC) of their activity.
Locomotion trace is shown on top (red). **E)** Same as A but for mice treated with muscimol
injection in the somatosensory thalamus (VPM and POM). **F)** Event-triggered average traces of
SST-INs GCaMP6s signals in S1 aligned to the onset of locomotion along with running speed
(red) and whisking (blue). Normalized $\Delta F/F$ is displayed for all analyzed cells (bottom). Results
are displayed for mice previously injected with muscimol in the thalamus. Traces represent
mean ± SEM values across all recorded cells. **G)** Mean $\Delta F/F_0$ during periods of rest and run in
control conditions and 30 min microinjection of muscimol in the somatosensory thalamus ($\Delta F/F_0$
$_{rest\ control}$ = 0.53 ± 0.01; $\Delta F/F_{0\ run\ control}$ = 1.01 ± 0.03; n = 1657 cells in 8 mice, p<0.0001, Mann-
Whitney U test; $\Delta F/F_{0\ rest\ muscimol}$ = 0.53 ± 0.02; $\Delta F/F_{0\ run\ muscimol}$ = 0.76 ± 0.04; n = 541 cells in 8
mice, p<0.0001, Mann-Whitney U test). **H)** Summary plot of average LMI obtained in control and
after muscimol injection per the different individual mice tested ($LMI_{control}$ = 0.23 ± 0.06,
$LMI_{muscimol}$ = 0.08 ± 0.06, n = 8 mice, p = 0.0391, Wilcoxon test. **I)** Summary plots for LMI (left)
and Pearson correlation values (middle) between running speed and calcium activity for SST-
INs in S1 obtained in control and after muscimol injection in the thalamus ($LMI_{control}$ = 0.18 ±
0.01, n = 1657, $LMI_{muscimol}$ = 0.10 ± 0.01, n = 541, p < 0.0001 Mann-Whitney U test; Pearson,
$\rho_{control}$ = 0.18 ± 0.01, n = 1657, $\rho_{muscimol}$ = 0.14 ± 0.01, n = 541, p < 0.05 Mann-Whitney U test) .
Percentage of SST-INs exhibiting statistically significant positive, negative, or non-significant
Pearson correlation values is shown on the right. **J)** Variation in $\Delta F/F$ values for SST-INs with
running speed for control and muscimol injection conditions.
965

966
Figure 7: Behavioral state transitions from rest to active states increase activity of
pyramidal cells in S1. A) Top: schematic illustration of layer 2/3 microcircuits in S1; bottom: in
vivo 2PSLM images of S1 PNs labeled with GCaMP6s. **B)** Raster plot of the pyramidal cell
activity in the S1 from awake mice spontaneously transitioning between resting periods and
periods of locomotion. Each row represents a neuron sorted by weight on the first principal
component (PC) of their activity. **C)** Mean $\Delta F/F_0$ during periods of rest and run (Control: $\Delta F/F_{0\ rest}$
= 0.21 ± 0.01; $\Delta F/F_{0\ run}$ = 0.80 ± 0.04; n = 1918 cells in 3 mice, p<0.0001, Mann-Whitney U test).
E) Cross-correlation between in vivo calcium activity and running speed for Pyramidal neurons
in S1 in control condition and after muscimol injections in somatosensory thalamus. **F-G)** Same
as B and C but mice previously microinjected with muscimol in thalamus **(**Muscimol: $\Delta F/F_{0\ rest}$ =
0.18 ± 0.01; $\Delta F/F_{0\ run}$ = 0.17 ± 0.16; n = 371 cells in 2 mice, p<0.0001, Mann-Whitney U test). **H)**
Schematic illustration of dual whole-cell patch clamp configuration used to record synaptic
properties of unitary connections between individual L2/3 PNs and local SST-INs. **I)** Example
traces of a train of EPSCs recorded from two different SST-INs in response to presynaptic
action potential induction in two different L2/3 PNs. Note the pronounced different in the
amplitude of the first EPSC between the two SST-INs. **J)** Histogram of amplitudes of first EPSC
obtained for 50 synaptic unitary connections. Inset depicts the relationship between amplitude
and failure rate for the 50 connections. **K)** Same as for **H** but illustrating that the properties of
two different presynaptic L2/3 PNs were recorded for the same postsynaptic SST-INs. **L)**
Example traces of EPSCs recorded in two different SST-INs for four presynaptic PNs. **M)**

Summary plot indicating the relationship between the amplitude obtained when stimulating PN1
or PN2 for the same postsynaptic SST-INs.
989
990

A

991
Supplementary figure 1: Variability of SST-INs response within individual fields of view in
S1 to behavioral transitions from rest to run. A) Representative GCaMP6s fluorescence
traces (black) shown as $\Delta F/F_0$ (up) as well as normalized $\Delta F/F_0$ (bottom) for SST-INs for
individual fields recorded in awake mice spontaneously transitioning between rest and run
periods. Each row represents a neuron sorted by weight on the first principal component (PC)
of their activity. Locomotion (red) and whisking (blues) traces are shown on top. **B)** Scatter plots
of the mean amplitude of fluorescence changes (DF/F_0) of each neuron illustrated in A for
locomotion periods versus stationary periods.
1000

43

1001
1002
Supplementary figure 2: Response of SST- and VIP-INs in S1 to behavioral transitions
from rest to run. A) Cross-correlation curves between running speed and calcium activity for
VIP and SST-INs. Values at zero time are displayed on the right. **B)** Variation in ΔF/F values for
SST- and VIP-INs with average running speed. **B)** Event-triggered average traces of VIP-INs
GCaMP6s fluorescence signal in S1 aligned to whisking onset. Only whisking bouts not
associated with locomotion were used (whisking only). Normalized ΔF/F is displayed for all
analyzed cells (bottom). Traces represent mean ± SEM values across all recorded cells. **(C)**
Enlarged view of average traces shown in A. **B)** Schematic illustration of whole-cell current
clamp recording in L2/3 of primary somatosensory (S1) cortex while puff applying nicotine 100
⌈M in SST or VIP-INₛ (left). Example traces of puff-evoked responses in SST- and VIP-INs
(right). **C)** Summary plot of variations in membrane potential induced by nicotine applications in
both SST- and VIP-INs (SST [n = 11] vs VIP [n = 9], p < 0.0001, Mann-Whitney test).
1015

1016
Supplementary figure 3: SST-INs are equipped with cholinergic receptors in both V1 and
S1.
A) Schematic illustration of whole-cell current clamp recording in L2/3 of primary somatosensory
(S1) cortex while puff applying carbachol (2mM) in SST (up). Example traces of puff-evoked
responses in SST-INs (down). **B)** Example traces of current clamp recordings illustrating the
variation in membrane potential in SST-INs to puff application of carbachol in both V1 (P>56:
top right) and S1 (P<56 middle right; P>56 bottom right). **C)** Summary plot of the variation of
membrane potential induced by carbachol (CCH) in SST-INs in V1 and S1 (V1, SST [n = 7]; S1,
SST with CCH puff alone [n = 20] versus SST with CCH puff in the presence of atropine 10⎾M
[n = 7], p = 0.007, Mann-Whitney test)**.** Error bars are SEM.
1027

1028
Supplementary figure 4: Somatosensory thalamus inactivation with muscimol injection
decreases whisker-evoked pyramidal cell activity in S1. A) Event-triggered average traces
of L2/3 PNs GCaMP6s signals in S1 aligned to whisker puff along with running speed (red).
Normalized ∆F/F is displayed for all analyzed cells (bottom). Results are displayed for control
conditions (left) and for mice previously injected with muscimol in the thalamus (*right*). Traces
represent mean ± SEM values across all recorded cells. **B)** Summary plot of variation in ∆F/F
between rest and after whisker stimulation for the event-triggered average traces shown in B.
Average values were obtained 2 s before and 2 s after whisker stimulation onset (see Methods).
1037

Experimental Procedures:

Mice:

Animals were housed in the Paris Brain Institute accredited by the French Ministry of Agriculture. All experiments were performed in compliance with French and European regulations on care and protection of laboratory animals (EU Directive 2010/63, French Law 2013-118, February 6th, 2013), and were approved by local ethics committees and by the French Ministry of Research and Innovation (authorization numbers #11199). All animal work was performed at the ICM PHENO-ICMice facility. Both male and female mice aged 4 weeks or older were used with the following genotypes: SST-IRES-Cre (SSTtm2.1(cre)Zjh/J; JAX 013044); SST-IRES-Cre X Ai9 (Gt(ROSA)26Sortm9(CAG-tdTomato); JAX 007909); VIP-IRES-Cre (Viptm1(cre)Zjh/AreckJ; JAX 031628) X Ai9 (Gt(ROSA)26Sortm9(CAG-tdTomato); JAX 007909); VIP-cre::SST-FLP ($Sst^{tm3.1(flpo)Zjh}$/J; JAX 028579). Animals were maintained on a 12-hour light/dark cycles with food and water provided ad libitum. Animals were housed in groups of 2-6 animals per cage, as often as possible, to favor social interactions, a condition that prevents the emergence of depressive and anxious-like behavior (Berry et al., 2012; Ieraci et al., 2016; Kappel et al., 2017).

Slice preparation

Acute parasagittal slices (320 µm) were prepared from adult mice, starting from postnatal days 25–70. Mice were deeply anesthetized with a mix of ketamine/xylazine (mix of i.p. ketamine 16 mg/kg and xylazine 13 mg/kg) and perfused transcardially with ice-cold cutting solution containing the following (in mM): 220 Sucrose, 11 Glucose, 2.5 KCl,1.25 NaH2PO4, 25 NaHCO3, 7 MgSO4, 0.5 CaCl2. After the perfusion the brain was quickly removed and slices prepared using a vibratome (Leica VT1200S). Slices containing S1 barrel field or V1 were transferred to ACSF solution at 34 °C containing (in mM): 125 NaCl, 2.5 KCl, 2 CaCl2, 1 MgCl2,

1065   1.25 NaH2PO4, 25 NaHCO3, 15 Glucose for 15–20 min. After the period of recovery, slices

1066   were kept at room temperature for a period of 5-6 h.

1067

Electrophysiology:

Whole-cell patch-clamp recordings were performed close to physiological temperature

(32–34 ^{o}C) using a Multiclamp 700B amplifier (Molecular Devices) and fire-polished thick-walled

glass patch electrodes (1.85 mm OD, 0.84 mm ID, World Precision Instruments); 3.5–5 MOhm

tip resistance. For voltage-clamp recordings, cells were whole-cell patched using following

intracellular solution (in mM): 90 Cs-MeSO3, 10 EGTA, 40 HEPES, 4 MgCl2, 5 QX-314, 2.5

CaCl2,10 Na2Phosphocreatine, 0.3 MgGTP, 4 Na2ATP (300 mOsm pH adjusted to 7.3 using

CsOH). Extracellular synaptic stimulation was achieved by applying voltage pulses (20 us, 5–50

V; Digitimer Ltd, UK) via a second patch pipette filled with ACSF and placed 20–40 mm from

soma. For current clamp experiments patch pipettes were filled with the following intracellular

solution (in mM): 135 K-gluconate, 5 KCl, 10 HEPES, 0.01 EGTA, 10 Na2phosphocreatine, 4

MgATP, 0.3 NaGTP (295 mOsm, pH adjusted to 7.3 using KOH). The membrane potential (Vm)

was held at -60 mV, if necessary, using small current injection (typically in a range between -50

pA and 200 pA). Recordings were not corrected for liquid junction potential. Series resistance

was compensated online by balancing the bridge and compensating pipette capacitance. APs

were initiated by brief current injection ranging from 1000–2000 pA and 1–5 ms duration. For all

experiments data were discarded if series resistance, measured with a 10 mV pulse in voltage

clamp configuration, was >20 MOhm or changed by more than 20% across the course of an

experiment. For current clamp experiment cells were excluded if input resistance varied by more

than 25% of the initial value. Recordings of puff evoked cholinergic responses in SST- and VIP-

INs were performed at the resting membrane potential in the presence of 10 μM CNQX and 10

μM bicuculine.

All recordings were low-pass filtered at 10 kHz and digitized at 100 kHz using an analog-

to-digital converter (model NI USB 6259, National Instruments, Austin, TX, USA) and acquired

with Nclamp software (Rothman and Silver, 2018) running in Igor PRO (Wavemetrics, Lake

Oswego, OR, USA). The intracellular solution used for the experiments in which a presynaptic

cell was depolarized by a train of action potentials to induce neurotransmitter release on a post-

synaptic cell/element was (in mM): 130 K-MeSO3, 4 MgCl2,10 HEPES, 0.01 EGTA, 4 Na2ATP,

0.3 NaGTP (300 mOsm, pH 7.3 adjusted with NaOH). Paired recordings connections were

probed using a train of 5/9 action potentials at frequency of 20 Hz.

1098

In vivo two-photon calcium imaging

Surgery

Mice were initially anesthetized with Isoflurane. Induction of anaesthesia was performed

at 4% isoflurane, Iso-Vet, and an air flow of 250ml/min. Mice were then placed in a stereotactic

device (Kopf) kept at 37 °C using a regulated heating blanket and a thermal probe and

maintained under anaesthesia using 1-2% Isoflurane and an air flow of 250 ml/min until the end

of the surgery. For calcium imaging in VIP- and SST-INs, cranial window implantation and AAV

viral stereotaxic injection was performed in the same surgical procedure. In order to prevent

brain swelling and post-operative pain, anti-inflammatory (Dexamethasone, 2 µg/g Dexadreson,

MSD) and analgesic drugs (Buprenorphine, 0.1 mg/kg Buprecare) were injected

subcutaneously. After shaving the scalp, the skin was cleaned by wiping multiple times with an

antiseptic solution and 70% alcohol. After injecting a local anaesthetic under the scalp

(lidocaine, 10mg/kg - Xylovet or Laocaine) a section of the skin (a circular section with 1.5 cm

diameter) was removed using surgical scissors. The periosteum was then carefully removed

and the skull was scrapped using a dental drill, in order to remove all residues. The surface of

the skull was subsequently thoroughly cleaned and dried using cottons swabs and a

physiological saline solution. Using the stereotactic device, the center of the cranial window was marked and a circular piece of bone was removed using a 3 mm biopsy punch (LCH-PUK-30 Kai Medical). This step of the surgical procedure was considered critical since damage to the dura would result in rapidly opacifying cranial windows. Once the brain was exposed, the AAV viral vectors carrying the genetically encoded calcium indicators (GCaMP6 f and s) were stereotactically injected. For calcium imaging of VIP-INs, SST-INs and pyramidal neurons, AAV9 Syn-Flex-GCaMP6f-WPRE-SV40 (2.1×10^{13} AddGene-100833), AAV9 Syn-Flex-GCaMP6s-WPRE-SV40 (2.5×10^{13} AddGene-100845) or AAV9 ENN-AAV-CaMKII.GCaMP6f-WPRE-SV40 (2.5×10^{13} AddGene-100834) was used respectively. Using a Nanojet III (Drummond Scientific), 200nl of virus were injected in the barrel cortex (from Bregma, 1.1 RC; 3.3 ML) or visual cortex (from bregma: 3.0 RC; 2.4 ML) at a rate of 1nl/s followed by 5 minutes of wait to prevent backflow. A round of 3mm glass coverslip (CS-3R Warner Instruments) was then placed over the craniotomy and glued in place using dental cement (Superbond). A stainless-steel head post (Luigs & Neumann) was then attached to the skull also using dental cement. For 3 days following the cranial windows surgery, mice were injected once a day with a mix of anti-inflammatory (Dexamethasone, 2μg g^{-1} Dexadreson, MSD) and analgesic drugs (Buprenorphine, 0.1 mg/kg Buprecare). Two weeks after surgery, mice were first habituated to the experimenter by handling. The following days, mice were head-fixed in the acquisition setup for increasing periods of time (5 – 45 minutes).

Thalamus inactivation

In order to inactivate the somatosensory-associated thalamus, following the acquisition of several control imaging sessions, mice were placed in a stereotactic device, and put under light isoflurane anesthesia. A hole was drilled over the ipsilateral thalamus and using a cannula, a bolus of fluorescent Muscimol (0.5 mM, Muscimol, BODIPY™ TMR-X Conjugate Muscimol, BODIPY™ TMR-X Conjugate, ThermoFischer Scientific) was slowly injected into the

thalamus (Coordinates: -1.82 RC, 1.5 ML, 3.5 depth). After injection, the craniotomy was

covered with silicone (Kwik-Cast, WPI) and the mouse was allowed to recover for an additional

30 minutes. After recovery, two-photon imaging was performed for a maximum of one hour post

muscimol injection. The effective inactivation of the thalamus was confirmed functionally by

recording the response of pyramidal cells as well as local cortical interneurons to whisker

stimulation as well as by performing histology just after the imaging sessions.

Two-Photon Imaging

In vivo two-photon calcium imaging was performed with an Ultima IV two-photon laser-

scanning microscope system (Bruker), using a 20×, 1.0 N.A. water immersion objective

(Olympus) with the femtosecond laser (MaiTai DeepSee, Spectra Physics) tuned to 920 nm for

imaging of cells expressing GCaMP6s/f and to 1020 nm for imaging of cells expressing

mCherry. Fluorescence light was separated from the excitation path through a long pass

dichroic (660dcxr; Chroma, USA), split into green and red channels with a second long pass

dichroic (575dcxr; Chroma, USA), and cleaned up with band pass filters (hq525/70 and

hq607/45; Chroma, USA). Fluorescence was detected using both proximal epifluorescence

photomultiplier tubes (gallium arsenide phosphide, H7422PA-40 SEL, Hamamatsu). Time-series

movies of neuronal populations expressing GCaMP6f were acquired at the frame rate of 30Hz

(512 × 512 pixels field of view; 1.13 µm/pixel). The duration of each focal plane movie was 300s

(9000 frames) to track spontaneous neuronal activity. During the recording periods animals

were free to run on a circular treadmill covered with a soft foam. This foam softened the

movement of the animals and provided better traction (this was important since mice adapted

more rapidly to the experimental setup and movements were more fluid during the recordings).

Locomotion was monitored using a rotary encoder (H5-1000-NE-S, US Digital) and digitized

with a NIDAQ analog-to-digital converter (model NI USB 6363, National Instruments, Austin, TX,

USA) at 2000 Hz. Whisker movement was recorded using a FLIR camera (BFS-U3-17S7M-C)

at a frame rate of 30Hz and using the Spinnaker SDK software (FLIR Systems Inc, Orlando, USA) equipped with a cleanup filter (FB850-40, Thor labs) to block laser shining from during two-photon imaging. The camera was focused on the whisker pad of the mouse using a 10X (13 - 130mm FL) C-Mount, Close Focus Zoom lens and each frame acquisition was controlled via Neuromatic running in IGORPRO (Wavemetrics) environment using a national instruments digital-to-analog board (NI-6393). The mouse snout was illuminated via an 850 nm LED (M850L3, Thor labs).

Whisker Stimulation

For whisker stimulation experiments, a NPI PDES-02DX (NPI Electronic GmbH) puffer was used. Tubing was connected to the puffer and a short puff (1 sec) of air was aimed at the mouse whiskers. The tip of the puffer was kept out of reach of exploratory whisking range. The whisker puff was randomly triggered during imaging sessions and the time of puff was recorded via the digitization of voltage trigger using a analog-to-digital converter (model NI USB 6259, National Instruments, Austin, TX, USA)

CNO Experiments:

For experiments involving DREADDs, mice were injected with clozapine-N-oxide (CNO). A water-soluble salt preparation of CNO, CNO dihydrochloride (Tocris, Bio-Techne LTD, Abingdon, UK, catalog no.: 6329), which was dissolved in sterile saline. Several concentrations of CNO were tested (1, 5 and 10mg/kg) and no significant changes were found so data was all data was pulled together.

Visual stimulation

To study the effect of light on the recruitment of VIP- and SST-INs in the cortex, an isoluminant grey screen (Lilliput 669GL 7") was placed at 45° from the center line of the mouse and at approximately 20cm from the left eye (contralateral to the craniotomy) of the mouse.

Visual stimulation trials were interleaved with control trials. In order to reduce light
contamination, a custom funnel was used to shield the objective from light coming from the
screen.

Calcium Imaging Analysis

Motion correction, Image segmentation and calcium signal extraction

Motion correction, image segmentation and calcium signal extraction were performed
using the python 3.6 package Suite2P (Pachitariu et al., 2016). This package analyses data in
four stages: image registration, region of interest (ROI) detection, quality control, and activity
and neuropil extraction. The first stage, image registration, is responsible for motion correction
and works by performing a mean image of a few frames and then shifting (in the x and y axis)
each frame so that it optimally fits the mean image. The second stage, ROI detection, detects
the cells in the frame. This is done by detecting the pixels with the highest variability across time
and then expanding the region to the surrounding pixels until this variability drops below a
defined threshold. Although this method works quite well, the disadvantage of this method is
that since only active cells are detected, it is difficult to determine the number of silent cells in
field of view, which could be important for functional tasks. In a third stage, quality control, the
program separates structures that correspond to cells from the remaining structures by
assigning them a score. The results of this stage were always manually curated. In the last
stage of the analysis, activity and neuropil extraction, is performed by averaging the pixel
intensity in the ROI for each time point. To determine the neuropil, a donut-like surrounding the
ROI is generated and the activity within this area is also averaged for each time point. Neuropil
plays a critical role in calcium imaging since it represents the activity of nearby structures
surrounding the cell of interest that might be excited by the 2P excitation PSF. Since these

structures might have a different behavioral profile than the cell of interest their activity might

contaminate the signals detected in the cell soma (Dipoppa et al., 2018).

Analysis of fluorescence signals

Raw fluorescence traces were corrected for neuropil contamination by subtracting

Suite2p-neuropil traces using a fixed scaling coefficient of 0.7 (Dipoppa et al., 2018). To be able

to compare activity across cells and mouse lines, fluorescence signals were normalized. To do

this we used the $\Delta F/F_0$ method, calculated using the following formula: $(F-F_0)/F_0$, where F is the

fluorescence and F_0 is the baseline fluorescence. When studying the activity of interneurons

(VIP- and SST-INs), the baseline fluorescence was calculated using the lower 5^{th} percentile.

Results were systematically evaluated using a different baseline method to estimate F0

(MinMax baseline method with a sliding window of 60s) and observations reported in the

present manuscript were insensitive to the baseline method used. For the analysis of pyramidal

cell activity, a sliding average window (60 s) defined as MinMax method was used.

Locomotion Analysis

Changes in positioning of the circular treadmill (sampled at 2 kHz) were interpolated onto

a down-sampled rate of 30Hz in order to match the sampling rate of the two-photon imaging.

Locomotion periods, were defined as the periods where speed was higher than 0.1 cm/s. All

points 0.5 seconds before and 1.5 seconds after a locomotion period were considered

stationary periods. Events spaced by less than 0.5s were joined. Sessions in which the

percentage of either locomotion or stationary periods were lower than 5%, were excluded. The

locomotion modulation index (LMI) was defined as the difference between the mean $\Delta F/F_0$

during locomotion (F_{run}) and stationary (F_{rest}) periods, normalized by the sum of both values: LMI

= $(F_{run} - F_{rest})/(F_{run} + F_{rest})$ (Pakan et al., 2016). To calculate statistically significant Pearson

Correlations between fluorescence and running speed, comparisons were made with 1000

shuffled (circular shuffling) locomotion traces. Pearson correlation coefficient was considered

statistically significant if value of original data was above a 95% confidence interval generated

from the 1000 shuffled traces.

Whisking Analysis

Whisker pad movement was tracked using the python framework Facemap (Syeda et al.,

2022) (https://github.com/MouseLand/facemap). Whisker-pad movement was measured by

selecting an ROI over the whisker pad and calculating the motion energy index (MEI) across the

video. MEI was defined as the sum of the absolute change in pixel intensity within the ROI

between adjacent video frames. Using custom python code, whisking pad motion energy was

filtered with a gaussian filter (sigma of 3), normalized with a Min Max method and resampled to

30Hz. Whisking periods were defined as periods where normalized MEI exceeded 10 %.

Whisking only periods were defined as periods in which there was whisking but not locomotion.

Whisking modulation index (WMI) and whisking only modulation index (WOMI) were calculated

in a similar way to LMI but using the variations in GCaMP6s fluorescence during resting or

whisking periods.

Event-triggered averages were computed by automatically identifying frames with event

onsets, namely those related to whisking and/or locomotion. Due to the slow time course of

GCaMP6s and to minimize cross-talk between events, we specifically chose locomotion events

that were not preceded by another locomotion period within a 5-second interval. For whisking

only analysis, whisking episodes not associated with locomotion were included in the average

(5se before and 5 sec after). In Figure 3I, locomotion events were further categorized in

agreement with locomotion duration. Once multiple locomotion and/or whisking events were

detected, the results were combined and averaged to calculate an average value for each cell.

Subsequently, these individual average cell values were averaged together and presented with

their respective average and SEM.

1261

Statistics

All statistical analysis were performed in Prism (GraphPad Software, Inc.). The error

bars in all graphs indicate standard error of the mean (S.E.M.) and statistics were performed

with two-tailed tests. For statistical tests comparing the average $\Delta F/F_0$ of neurons between two

behavioral states (stationary versus locomotion periods) in the same subject, we used Wilcoxon

signed-rank tests. For statistical tests comparing the distribution of LMI, Pearson Correlation or

behavior properties between brain regions or interneuron populations, Mann-Whitney U tests

were used. When comparing two conditions across behavioral states, Kruskal-Wallis test was

used.

1271
1272
1273