How smart was *T. rex*? Testing claims of exceptional cognition in dinosaurs and the

application of neuron count estimates in palaeontological research

Caspar, Kai R.[1,2*–], Gutiérrez-Ibáñez, Cristián[3*–], Bertrand, Ornella C.[4], Carr, Thomas[5]

Colbourne, Jennifer A. D.[6]; Erb, Arthur[7,8], George, Hady[9], Holtz, Thomas R., Jr.[10,11], Naish,

Darren[12]; Wylie, Douglas R.[3]; Hurlburt, Grant R.[13*–]

*These authors contributed equally; □ corresponding authors

[1]Institute of Cell Biology, Heinrich Heine University Düsseldorf, Düsseldorf, Germany;[2]Department of

Game Management and Wildlife Biology, Faculty of Forestry and Wood Sciences, Czech University of

Life Sciences, Prague, Czech Republic;[3]Department of Biological Sciences, University of Alberta,

Edmonton, Canada;[4]Institut Català de Paleontologia Miquel Crusafont, Universitat Autònoma de

Barcelona, Barcelona, Spain [5]Department of Biology, Carthage College, Kenosha, Wisconsin, USA

[6]Comparative Cognition Unit, Messerli Research Institute, University of Veterinary Medicine Vienna,

Vienna, Austria;[7] School of GeoSciences, Grant Institute, University of Edinburgh, Edinburgh,

Scotland, UK;[8]Center for Science, Teaching, and Learning, Rockville Centre, New York, USA;[9]School

of Earth Sciences, University of Bristol, Bristol, UK; [10]Department of Geology, University of Maryland,

College Park, Maryland, USA;[11]Department of Paleobiology, National Museum of Natural History,

Washington, DC, USA;[12] School of Biological Sciences, Faculty of Environment and Life Sciences,

University of Southampton, Southampton, United Kingdom;[13]Department of Natural History, Royal

Ontario Museum, Toronto, Ontario, Canada

Abstract

Recent years have seen increasing scientific interest in whether neuron counts can act as

correlates of diverse biological phenomena. Lately, Herculano-Houzel (2023) argued that

fossil endocasts and comparative neurological data from extant sauropsids allow to

reconstruct telencephalic neuron counts in Mesozoic dinosaurs and pterosaurs, which might

act as proxies for behaviors and life history traits in these animals. According to this analysis,

large theropods such as *Tyrannosaurus rex* were long-lived, exceptionally intelligent animals

equipped with "macaque- or baboon-like cognition" whereas sauropods as well as most

ornithischian dinosaurs would have displayed significantly smaller brains and an ectothermic

physiology. Besides challenging established views on Mesozoic dinosaur biology, these

claims raise questions on whether neuron count estimates could benefit research on fossil

animals in general. Here, we address these findings by revisiting Herculano-Houzel's (2023)

work, identifying several crucial shortcomings regarding analysis and interpretation. We

present revised estimates of encephalization and telencephalic neuron counts in dinosaurs,

which we derive from phylogenetically informed modeling and an amended dataset of

endocranial measurements. For large-bodied theropods in particular, we recover significantly

lower neuron counts than previously proposed. Furthermore, we review the suitability of

neurological variables such as neuron numbers and relative brain size to predict cognitive

complexity, metabolic rate and life history traits in dinosaurs, coming to the conclusion that

they are flawed proxies of these biological phenomena. Instead of relying on such

neurological estimates when reconstructing Mesozoic dinosaur biology, we argue that

integrative studies are needed to approach this complex subject.

49 Introduction

The Late Cretaceous North American theropod dinosaur *Tyrannosaurus rex* is a superlative

predator, being among the largest, heaviest, and most powerful (in terms of bite force)

terrestrial carnivores of all time (Gignac and Erickson 2017; Sakamoto 2022; Henderson,

2023). Recently, Herculano-Houzel (2023) proposed that anthropoid primate-level

intelligence should be added to *T. rex*'s already impressive predatory resume based on high

estimated numbers of telencephalic neuron counts in large-bodied theropod taxa. This

conclusion emerged from a paradigm whereby neurological variables estimated from

endocasts can, so it is claimed, be used to infer metabolic parameters, social behaviors, and

longevity in fossil species. Here, we test whether this approach and its remarkable prospects

withstand scrutiny.

The hypothesis of exceptional intelligence in dinosaurs such as *T. rex* challenges the

consensus of crocodile-like cognition in these animals, a position informed by comparative

anatomical data (Rogers, 1998; Witmer & Ridgely, 2009; Hurlburt et al., 2013). Moreover,

this claim bears ramifications that extend beyond specialized biological disciplines due to its

potential to create long-lasting impacts on the public's perspective on dinosaurs, evolution,

and the scientific process. Given the extreme contrast between Herculano-Houzel's (2023)

proposal and more traditional perspectives on dinosaur biology, we revisit the claim of

exceptional intelligence in these animals through an assessment of her methodology and a

reanalysis of the underlying data. By integrating perspectives from both paleontology and

neontology, we evaluate the potential benefits and limitations of neuron count estimation in

research on the behavior and physiology of fossil species. We begin with a brief review of

dinosaur paleoneurology and a discussion of how Herculano-Houzel's (2023) approach aims

to expand the field's methodological tool kit.

Dinosaur paleoneurology and the prospects of neuron count estimates for the field

Paleoneurology is a subfield of paleontology dedicated to research on the nervous systems

of extinct animals. Because soft tissues are not readily preserved in the fossil record,

paleobiologists rely on endocasts when studying the brains of extinct species (Paulina-

Carabajal et al. 2023). An endocast can be a natural (infilling), artificial (mold) or virtual

(digitally reconstructed) cast of the endocranial cavity that is formed by the bones of the

braincase.

The study of extinct species' endocasts, including those of dinosaurs, can be traced back to

the 1800s (e.g., Cuvier, 1812; Marsh, 1879). However, the field was truly defined by Edinger

(1929) who effectively introduced the concept of geological time to neurobiological studies.

Before her, anatomists made comparisons between endocasts and fresh brains, but without

considering the stratigraphic context (Buchholtz & Seyfarth, 2001). Jerison (1973) built on

Edinger's work by studying brain evolution in a quantitative manner and developed the

encephalization quotient (EQ) as an estimate of relative brain size, applicable to both extant

and extinct species. Later, the advent of X-ray computed tomography at the end of the

1990's transformed the field and provided novel ways in which the neurosensory systems of

extinct species could be studied (e.g., Knoll et al., 1999; Witmer et al., 2008). Despite these

crucial innovations, however, paleoneurology has so far remained largely restricted to the

measurement and comparison of gross morphology, limiting our understanding of how the

brains of Mesozoic dinosaurs and other extinct animals worked.

Figure 1: Simplified phylogeny of the Sauropsida (= total group Reptilia) with a focus on the taxon

Ornithodira (the least inclusive clade containing pterosaurs and dinosaurs, see revised definition of

Nesbitt, 2011) and representative color-coded brain morphologies, excluding the pituitary (not to

scale). Blue: olfactory bulb and tracts, Green: pallium (homologous to the mammalian cerebral

cortex), Orange: cerebellum, Yellow: diencephalon and optic tectum; Violet: brain stem. Olfactory

structures, pallium and subpallium comprise the telencephalon. The gray overlay indicates extinct

taxa, the brain morphologies of which are approximated. Note that the brain morphology in *T. rex* and

its relatives (Tyrannosauroidea) is conspicuously plesiomorphic when compared to the other

ornithodirans pictured here (see e.g., Giffin, 1989). Definitions of notable dinosaur clades: Ornithischia

- a large group of primarily herbivorous dinosaurs, excluding the long-necked sauropodomorph

dinosaurs, defined as the most inclusive clade including *Triceratops* but not *Diplodocus* nor

Tyrannosaurus. Most popular representatives of this group include horned or otherwise heavily

armored forms such as *Triceratops*, *Stegosaurus* and *Ankylosaurus* as well as the hadrosaurs,

colloquially known as duck-billed dinosaurs. Sauropodomorpha - the long-necked and often

particularly large-bodied herbivorous dinosaurs, defined as the most inclusive clade including

Diplodocus but not *Triceratops* nor *Tyrannosaurus;* Theropoda - the bipedal, mostly carnivorous

dinosaurs, the most inclusive clade including *Tyrannosaurus* but not *Diplodocus* or *Triceratops.* The

birds are part of this clade (see Baron et al*.,* 2017 for definitions of Ornithischia, Sauropodomorpha

and Theropoda)*;* Tyrannosauroidea, the most inclusive clade of theropods containing *Tyrannosaurus*

but not more bird-like taxa such as *Velociraptor* and *Ornithomimus* (Sereno *et al.,* 2009);

Maniraptoriformes, the least inclusive clade containing *Velociraptor* and *Ornithomimus* but not earlier-

diverging theropods like *Tyrannosaurus* (Holtz, 1996)*.* Silhouettes were taken from PhyloPic (listed

from top to bottom): *Morunasaurus* (in public domain), *Crocodylus* (in public domain),

Rhamphorhynchus (by Scott Hartman), *Olorotitan* (by ДиБгд, vectorized by T. Michael Keesey),

Tyrannosaurus (by Matt Dempsey), *Dromaeosaurus* (by Pranav Iyer), *Dromaius* (by Darren Naish),

Corvus (in public domain).

Pterosaurs and dinosaurs (the latter including birds) form the clade Ornithodira (Fig. 1), the

closest extant relatives of which are crocodilians (Fig. 1). Together, both lineages, which

separated about 250 million years ago, comprise the clade Archosauria (e.g., Legendre et

al., 2016). Next to birds, crocodilians therefore represent a critical reference point in

reconstructing the nervous systems of extinct ornithodirans.

Interestingly, highly disparate patterns of endocranial tissue organization are realized in

these two extant clades. One fundamental difference relates to the portion of the endocranial

cavity which is occupied by the brain rather than by the associated meningeal tissues

(including the dura mater and arachnoid mater) and cerebrospinal fluid (Fig. 2). In

crocodilians, nervous tissue only fills a fraction of the braincase (Hopson, 1979; Jirak &

Janacek, 2017; Watanabe et al. 2019). Longitudinal venous sinuses course along the dorsal

and ventral aspect of the brain, obscuring its true shape in casts of the braincase.

Furthermore, the size of the brain relative to both the endocranial volume and total body

size, decreases during crocodilian ontogeny, even over the course of adulthood (Hurlburt et

al., 2013; but note that absolute brain volume increases with body size, even in adults -

Ngwenya et al., 2013).

Endocast morphology indicates that the endocranial cavity in most non-avian dinosaurs was

organized in crocodilian-like fashion and comparative studies suggest that this configuration

was indeed ancestral for the clade Archosauria (Witmer et al., 2008; Hurlburt et al., 2013;

Fabbri & Bhullar, 2022). For tyrannosauroids specifically, which are among the best-studied

dinosaurs when it comes to palaeoneurology, endocasts representing different ontogenetic

stages suggest that brain size (relative to endocranial volume) decreased with age (Brusatte

et al., 2009; Witmer & Ridgely, 2009; Bever et al., 2013), as is the case in modern

crocodilians. As in, crocodilians, most dinosaurian endocasts do not faithfully capture the

volume and anatomy of the brain, particularly its posterior regions such as the cerebellum

(Watanabe et al., 2019). This contrasts with the situation in birds and most mammals for

which endocasts represent excellent brain size proxies (e.g., Iwaniuk and Nelson, 2002;

Bertrand et al., 2022).

The avian pattern probably evolved at the root of the theropod dinosaur clade

Maniraptoriformes, which includes ornithomimosaurs ('ostrich-mimic' dinosaurs) and

maniraptorans (the bird-like oviraptorosaurs, dromaeosaurids and kin, and birds themselves)

(Osmólska, 2004; Balanoff et al., 2013; Fig. 1). Maniraptoriform brains have enlarged

cerebral and cerebellar regions that almost fully occupy the endocranial cavity, as evidenced

by brain contours faithfully captured by the endocranium and extensive vascular imprints.

There is no evidence that the brains of other dinosaurs similarly contacted the endocranial

surface (pachycephalosaurs pose an exception to this pattern but are not covered in this

article, their endocranial anatomy is described in Hopson, 1979, Giffin, 1989, and Evans,

2005; we discuss other suggested cases of secondarily increased endocranial fills in

dinosaurs in Supplementary File 1). Pterosaurs are similar to maniraptoriforms in also

possessing brains that fit tightly into the endocranial cavity (Witmer et al., 2003).

Aside from general endocranial tissue organization, the neuroarchitecture and circuitry of

the forebrain in birds and crocodilians differs notably from one another (Ulinski & Margoliash,

1990; Briscoe et al., 2018; Briscoe & Ragsdale, 2018). Comparisons with other sauropsids

demonstrate that again the crocodilian condition is more plesiomorphic (Briscoe & Ragsdale,

2018). To which extent non-avian dinosaurs and pterosaurs resembled the two extant

archosaur groups in these regards cannot be reliably reconstructed, since they lack

osteological correlates.

The inferred brain anatomy of various dinosaur groups has been discussed elsewhere

(Paulina-Carabajal et al. 2023) and reviewing it here is beyond the scope of this article. We

aim instead to focus on what endocast-based methods potentially reveal about the behavior

and cognition of extinct species. While considering the aforementioned limitations,

endocasts from fossil ornithodirans allow us to reasonably estimate basic neuroanatomical

measures such as EQ, as well as to deduce specific sensory specializations (e.g., Witmer et

al., 2003; Witmer & Ridgely, 2009; Zelenitsky et al., 2011). Nonetheless, it is generally

assumed that the predictive power of these data in elucidating the cognitive capacities of a

fossil species is low (Paulina-Carabajal et al., 2023). Researchers have long sought to

identify robust morphological correlates of cognition but have found traditional proxies such

as EQ and absolute brain size to be limited and problematic regarding their conceptual

justifications (Van Schaik et al., 2021). Current debates focus on whether refined

neuroanatomical measures such as "cognitive brain size" (Van Schaik et al., 2021) and brain

region-specific neuron counts (Herculano-Houzel, 2011; Kabadayi et al., 2016; Logan et al.,

2018; Sol et al., 2022) might be able to overcome these issues. The quantification of these,

however, seemed out of reach for vertebrate palaeontology.

With this in mind, the approach proposed by Herculano-Houzel (2023) is of great potential

significance: It entails that endocasts of extinct taxa can be used to model neuron counts if

neurological data from related extant species can be taken into account. If valid, this

technique would potentially allow researchers to elucidate aspects of brain physiology that

cannot be inferred from endocast morphology alone. Herculano-Houzel & Kaas (2011) and

Herculano-Houzel et al. (2011) pioneered this approach for fossil hominins and extinct giant

rodents, but Herculano-Houzel (2023) was first in applying this methodology to fossil

sauropsid groups separated from their extant relatives by hundreds of millions of years of

evolution, namely pterosaurs and Mesozoic dinosaurs.

Figure 2: The endocast and endocranial tissue organization in the American alligator
(*Alligator mississippiensis*), illustrating the plesiomorphic condition within the clade Archosauria. Scale
bar = 2 cm in all cases. **a:** Endocast of a wild *A. mississippiensis* (Fla.F&G.HarvestTag 937095),
Dorsal cranial length (DCL), 342.90 mm, right lateral view. Reduced in size to match proportions of
brain in **b** & **c**. **b**: Dura mater around the brain of *A. mississippiensis*, specimen CITES FLM 12-
29409, DCL, 380 mm, left lateral view (reversed). **c**: Brain within arachnoid of FLM 12-29409. Brown-
red material is dried blood filling the subarachnoid space (**SaS**), right lateral view. **Abbreviations:**
Ach, arachnoid mater (covering the cerebellum); **Art**, artery on external wall of dura mater over the

lateral pole of the cerebrum; **CbII**, cerebellum; **CCA**, caudal cerebral artery; **Ch L** or **R**, left or right

cerebral hemisphere; **DSS**, dorsal sagittal sinus; **EthA**, common ethmoid artery; **I C**, internal carotid

artery; **InHA**, interhemispheric artery; **MO**, medulla oblongata; **N.II**, optic nerve; **N.V,** (cast of)

trigeminal nerve; **Nn R**, roots of nerves IX-XI; **OC**, occipital condyle; **OcS**, occipital sinus; **OlBu** & **Tr**,

olfactory bulb & tract; **OtC F,** fossa of otic capsule; **Pineal Loc**, pineal gland location; **Pit**, pituitary

gland; **SaSMe**, mesencephalic subarachnoid space; **SaSR**, rostral SaS; **SaSVe**, ventral SaS; **SN.I**,

first spinal nerve. The rostral end of the cerebrum is below the arrow for **SaSR** in **B**. Both specimens

are housed in the private collection of G. R. Hurlburt.

Indeed, Herculano-Houzel (2011, 2017, 2023) has argued emphatically that neuron counts

represent reliable estimates for cognitive abilities in extant vertebrates, markedly

outperforming other measures such as relative or absolute brain size. If we accept this

premise, accurate modeling of neuron counts in dinosaurs based on endocast volumes and

comparative neurological data might appear as a promising new method to elucidate the

behavior and cognitive capacities of these animals.

The methodology and rationale of Herculano-Houzel (2023)

Herculano-Houzel (2023) reconstructed relative brain size and neuron counts for 29

dinosaur and pterosaur species based on comparative data from extant non-avian and avian

sauropsids ("reptiles" and birds respectively; Olkowicz et al., 2016; Kverková et al., 2022).

Although we want to avoid lengthy discussions about taxonomy, it is worth noting that some

of these are no longer considered valid taxonomic entities (see below; an updated

nomenclature for relevant dinosaurs is included in Tab. 1). For instance, *Rhamphorhynchus*

muensteri and *R. gemmingi* have long been synonymized (Bennett, 1995). Surprisingly,

Herculano-Houzel (2023) inferred an ectothermic metabolism for one, and endothermy for

the other based on assumptions about relative brain size.

Neuron count estimates for fossil taxa only concerned the telencephalon, a major brain

region which is critically involved in cognitive and motor functions as well as the processing

of sensory information. It encompasses the pallium (which is homologous to the cerebral

cortex in humans and other mammals) and subpallium as well as the olfactory bulbs and

tracts (Fig. 1). To understand the rationale behind the approach of reconstructing neuron

counts in fossil species, two important matters must be pointed out: firstly, among jawed

vertebrates, body size and brain size are highly correlated, exhibiting a constant allometric

relationship overall (Tsuboi et al., 2018). It should be noted however, that scaling

relationships can vary to some extent between major taxa as well as between early- and

late-diverging members of a clade (Ksepka et al., 2020; Bertrand et al., 2022). Secondly,

neuronal densities (the number of neurons in a given volume of nervous tissue) can differ

profoundly between different vertebrate taxa. Based on current evidence, the highest neuron

densities among land vertebrates are found in the bird clade Telluraves, consisting of birds

of prey, rollers, parrots, songbirds and kin, while the lowest occur among crocodilians and

turtles (Kverková et al., 2022). For instance, the goldcrest (*Regulus regulus*), short-tailed

shrew (*Blarina* sp.) and painted turtle (*Chrysemys picta*) have brains of equal mass (ca.

0.37 g), but there is remarkable disparity in their whole brain neuron numbers, which range

from 14.3 M in the turtle over 58.8 M in the shrew to 164 M in the passerine bird (Olkowicz et

al., 2016; Kverková et al., 2022). This example illustrates that brain size alone is not a

reliable predictor of neuron counts across distantly related clades (compare Herculano-

Houzel et al., 2014; Olkowicz et al., 2016), which makes their inference in fossil groups

inherently difficult.

To decide which neuronal density patterns apply to specific groups of dinosaurs and

pterosaurs, Herculano-Houzel (2023) relied on brain x body mass regressions. The brain

and body mass datasets used were taken from various literature sources and, as we attempt

to show here, both are problematic. In the resulting regression plot, she identified theropods

clustering distinctly from most other species. On average, they appeared to exhibit larger

brains for a given body size than the remaining dinosaur or pterosaur taxa. When comparing

the regression lines for extinct groups with those of living birds on the one hand and

squamates and turtles on the other, Herculano-Houzel (2023) noted that the theropod

regression fits with the avian one, while the remaining ornithodiran groups aligned more with

the non-avian sauropsid regression line.

Based on these analyses, the author made two critical assumptions: First, since only

theropod brain-body data aligned with those of endothermic extant sauropsids, namely birds,

the other groups (aside from specific pterosaurs and ornithischians that cluster with

theropods) should be considered ectothermic. Second, telencephalic neuron densities in

theropod brain tissue should have been comparable to those found in certain extant bird

taxa (that is, to those found in a polyphyletic assemblage denoted as "pre-K-Pg birds" that

includes Palaeognathae, Galloanserae and Columbiformes and which is considered to form

a neurological grade: Kverková et al., 2022), whereas those of the other groups should

resemble densities encountered in squamates and turtles. No further justification for these

suggestions is provided.

Applying the avian scaling regime, Herculano-Houzel (2023) estimated remarkably high

telencephalic neuron counts in large-bodied theropods such as *Acrocanthosaurus atokensis*

(2.1 billion) and *T. rex* (3.3 billion) which would exceed those of any extant bird and be

comparable to large-bodied Old World monkeys such as baboons (*Papio anubis* - Olkowicz

et al., 2016). Based on this apparent similarity to anthropoid primates, she further speculated

that these giant theropods would have crafted and used tools and exhibited cultural

behaviors (Herculano-Houzel, 2023).

We regard the methodology of Herculano-Houzel (2023) as problematic and disagree with

her physiological and behavioral interpretations. Before we attempt to replicate her findings

with a more refined analytical approach, we want to enumerate the most important flaws of

the article and how they affect the inferences made.

Issues with Herculano-Houzel's method and rationale

A key problem for paleoneurology lies in the fact that an endocast does not necessarily reflect the morphology of an animal's brain. As discussed in previous sections, the endocasts of most non-avian dinosaurs differ markedly in size and shape from the actual brain, as is the case in crocodilians (Fig. 1). Unfortunately, not all studies from which Herculano-Houzel (2023) derived her raw data considered this issue (see below). In addition, the percentage of endocranial space filled by the brain, as well as its proportions may be further influenced by ontogeny (Bever et al., 2013; Hurlburt et al., 2013; Ngwenya et al., 2013; Jirak & Janacek, 2017; Hu et al., 2021). The latter point is relevant because Herculano-Houzel (2023) included several specimens which corresponded to juveniles rather than adults, and thus might have introduced biases to the dataset (see below). Interestingly, at least in crocodilians, neuronal densities in the brain are also affected by ontogenetic stage (Ngwenya et al., 2016).

To arrive at the estimated telencephalic neuron count of > 3 billion for *T. rex*, Herculano-Houzel (2023) assumed a brain mass of 343 g. However, this presupposed that endocast volume equaled brain volume in this species. While it has indeed been claimed that in theropods such as *T. rex*, the brain filled the entire endocranial cavity (Balanoff et al., 2013), this hypothesis is, as previously discussed, contradicted by multiple lines of evidence. More conservative inferences suggest a brain mass of approximately 200 g (Hurlburt, 1996; Hurlburt, et al, 2013; Morhardt, 2016) or possibly even lower (this study; Tab. 1) for *T. rex*. Herculano-Houzel (2023) acknowledged these lower estimates but chose to rely on the inflated values for large theropod brain masses in accompanying figures and the article's Discussion section.

Moreover, while the literature-derived brain mass estimates used for their analyses did in some cases include the olfactory tracts and bulbs (Franzosa & Rowe, 2005, Balanoff et al.,

2013), these were omitted in others (Hurlburt 1996; Hurlburt et al., 2013). This incongruence

created critical biases, affecting both the inference of telencephalic neuron counts and

relative brain size estimates. The latter are additionally skewed by the fact that body masses

used by Herculano-Houzel (2023) were not determined via a uniform methodology but

compiled from sources applying various approaches. There are several ways to estimate

body mass in extinct animals and they can differ greatly regarding their outcomes and

precision (Campione & Evans, 2020). When compared to body mass estimates derived from

stylopodial circumference, a well-established and robust method (Campione & Evans, 2020),

some striking differences become apparent (Tab. 1; Herculano-Houzel, 2023).

Another flaw of Herculano-Houzel's (2023) approach is the neglect of brain morphology to

inform its analyses. To estimate telencephalic neuron numbers in fossil species, the mass of

the telencephalon needs to be approximated first. For theropods, Herculano-Houzel (2023:6)

extrapolated this variable from extant bird data while stating that "within a clade, brain mass

has strongly predictive power to arrive at estimates of numbers of telencephalic neurons in a

brain of known mass, once the neuronal scaling rules that presumably apply are known."

However, this statement can only hold true if the general proportions of the telencephalon

compared to the remaining brain are roughly constant in the group of concern, which is a

precondition that Herculano-Houzel (2023) did not test for in the fossil sample. Indeed, avian

brains only poorly reflect the brain morphologies found in the majority of Mesozoic dinosaurs

(reviewed by Paulina-Carabajal et al. 2023) and their general proportions are only

comparable to those found among maniraptoriform theropods (Balanoff et al., 2013; Fig. 1).

An important difference concerns the pallium, which crucially contributes to higher cognitive

functions, and greatly increased in size within the maniraptoriform radiation (Balanoff et al.,

2013). The same is true for the cerebellum, a part of the brain which is not encompassed by

the telencephalon but is also involved in various aspects of cognition in amniotes (Spence et

al., 2009). Thus, the telencephalic mass and proportions of non-maniraptoriform theropods,

such as *T. rex*, cannot be adequately modeled based on extant birds. Similar limitations

need to be considered when reconstructing traits of, for instance, the pterosaur or
sauropodomorph telencephalon based on extant non-avian sauropsids and they also apply
to our own empirical approach.

Herculano-Houzel (2023) hypothesized that the inferred incongruence in relative brain size
between theropods and other dinosaurs reflects differences in thermobiology, which would
justify applying avian neuronal scaling schemes to the former and non-avian sauropsid
scaling to the latter. Sauropodomorphs as well as selected ornithischians and pterosaurs are
instead suggested to be ectothermic due to their relatively smaller brains. Both of these
assumptions are problematic: First, multiple lines of evidence suggest that ornithodiran
endothermy evolved long before theropods emerged and was likely already present in the
Early Triassic common ancestor of dinosaurs and pterosaurs (e.g., Benton, 2012; Grigg et
al., 2022). We will revisit this evidence and how it challenges the brain size-derived
hypothesis in more detail in the Discussion section of this paper. Herculano-Houzel (2023)
only referenced a single article on dinosaur thermobiology, that of Wiemann et al. (2022), to
defend her standpoint on the matter. The study in question applies a promising but novel
technique to infer endothermy based on lipoxidation end products in fossil bone that still has
to prove itself. While it indeed suggests lowered metabolic rates in some ornithischians, it
also infers an endothermic metabolism for pterosaurs and sauropodomorphs (Wiemann et
al., 2022). Thus, its findings do not align with Herculano-Houzel's (2023) assumptions.

Second comparisons between groups of extant vertebrates, especially birds and mammals,
strongly suggest that there is no uniform relationship between neuron density and relative
brain size or elevated metabolic rates (Kverkova et al., 2022; see also Estienne et al., 2024).
We will elaborate on this aspect in the Discussion section but would like to state at this point
already that it is not straight-forward to assume avian neuron densities in Mesozoic
theropods simply because they exhibited endothermy or a potential increase in relative brain
size. On the other hand, the extensive evidence for endothermy in other dinosaurs and

pterosaurs does not entail that these groups could not have had neuron densities similar to

those found in extant ectothermic sauropsids.

Other issues relate to the statistical methods employed by Herculano-Houzel (2023). Despite

dealing with a large multi-species dataset, their analyses did not take phylogeny into

account, which can produce mathematical artifacts. Phylogenetic relationships among taxa

need to be statistically addressed because shared ancestry can result in non-independence

of species-specific data points (Revell et al., 2008). Such non-independence is known as the

phylogenetic signal, and it has been prominently recovered for relative brain size in extant

sauropsids (Font et al., 2019). Hence, phylogenetically-informed modeling is necessary for

adequately analyzing such datasets (Font et al., 2019).

In light of these substantial shortcomings, we attempt to replicate the findings of Herculano-

Houzel (2023) with phylogenetically informed models of telencephalic neuron counts in fossil

dinosaurs that acknowledge the issues lined out above. Different from her, we do not include

pterosaurs into our analysis due to difficulties with estimating their body mass (especially for

taxa with incomplete postcrania such as *Scaphognathus*) and because of the unclear

taxonomic and ontogenetic status of some of the few available endocasts.

Empirical part: modeling neurological variables in dinosaurs

Endocast sample composition, with notes on endocranial volumes provided by

Hurlburt (1996)

We estimated the mass of the brain (MBr, g; excluding the olfactory tracts and bulbs) as well

as its size relative to body mass (MBd, g) in 31 Mesozoic dinosaur taxa for which data on

endocranial volume (EV, ml) have been published (Tab. 1; Suppl. File 1). Note that this

study does not aim to provide a comprehensive dataset of dinosaur brain sizes. Given the

questions we want to address, we focus on large-bodied theropods and taxa covered in

previous comparative analyses. We included one endocast per species, except for *T. rex*, for

which three adult endocasts (AMNH 5029, AMNH 5117, FMNH PR 2081) were considered.

We only considered species for which we could calculate body mass based on stylopodial

circumference (see below) to reliably infer encephalization. Due to this, our analysis does

not cover all dinosaur species for which complete endocasts are available, nor all species

that Herculano-Houzel (2023) included in her analyses (namely *Conchoraptor gracilis*,

Tsaagan mangas, *Zanabazar junior*, and unnamed troodontid IGM 100/1126). Juvenile

specimens considered by that study (*Alioramus altai* IGM 100/1844, *Gorgosaurus libratus*

ROM 1247, and *Tyrannosaurus rex* CMN 7541 = *"Nanotyrannus lancensis")* were omitted in

this analysis to eliminate the confounding variable of ontogeny.

The only juvenile we include is *Bambiraptor feinbergi* KUVP 129737, which is one of the few

maniraptoriform theropods that we can take into account. For this species, an adult femur

(FIP 002) is available, allowing us to estimate body mass in fully grown individuals. Our

method of body mass inference suggests that KUVP 129737 had attained about 45% of

adult body mass when it died. Data from similar-sized extant rheas (*Rhea americana*),

palaeognath birds which are close neuroanatomical analogs to highly derived theropods

such as *Bambiraptor* (Balanoff et al., 2013), suggest that adult brain mass is already

approached at that point of somatic development (Picasso et al., 2011; Picasso, 2012;). We

therefore combine the juvenile endocranial measurement of *Bambiraptor* with adult body

mass estimates.

Just as Herculano-Houzel (2023) did, we derive a significant portion of our EV values from

Hurlburt (1996). However, many EV figures communicated in this reference must be

considered outdated or otherwise flawed and were carefully bypassed here. We give

detailed reasons for discarding or modifying data from Hurlburt (1996) and the references

provided therein (Jerison, 1973; Hopson, 1979) in Suppl. File 1 Part A. Given that EVs from

this problematic dataset are still widely used (e.g., Müller et al. 2021; Button & Zanno, 2023),

we consider their revision an important aspect of this study. In cases where EVs appeared

doubtful but appropriate illustrations or photographs of specimens were available, we

recalculated EV using manual graphic double integration (GDI; see below for methodology).

This was done for four species (*Allosaurus fragilis, Euoplocephalus tutus, Kentrosaurus*

aethiopicus & Ornithomimus edmontonicus; see Suppl. File 1 for details on specimens).

Brain mass estimates

We estimated the brain mass (MBr, g) of fossil dinosaurs from endocast volume (EV, mL).

Because the specific gravity (density) of living amniote brain tissue approximates one (1.036

g/mL - Iwaniuk & Nelson, 2002), we used brain volume and mass interchangeably (compare

Hurlburt et al., 2013; Herculano-Houzel, 2023). For maniraptoriform species, because their

endocasts preserve brain contours similar to those of avians, we assumed a brain:endocast

ratio of 100%. This is consistent with empirical data on the relationship between MBr and EV

in modern birds, which suggest contributions of meningeal tissue to endocast volume to be

negligible (Iwaniuk & Nelson, 2002). For other dinosaurs, we assumed MBr:EV ratios of 31%

and 42%. Many previous studies have assumed a 50% ratio in these groups (reviewed in

Morhardt, 2016) while some even assumed 100% (Balanoff et al., 2013) or advocated for

intermediate values (e.g., Evans et al., 2008; Knoll & Schwarz-Wings, 2009; Knoll et al.,

2021). The widely adopted 50% ratio was originally proposed by Jerison (1973) and based

on measurements from a likely subadult green iguana (*Iguana iguana*) and a mere visual

estimate of endocranial filling in the tuatara (*Sphenodon punctatus*; the endocranial fill in this

species is now known to be 30% in adults - Roese-Miron et al., 2023). We abandon the

problematic 50% estimate and replace it here by the two aforementioned ratios that are

based on the morphology of extant crocodilians, the closest extant analogs to most non-

avian dinosaurs in regards to endocranial tissue organization, body size and braincase

ossification.   Excluding one anomalous value, the lowest MBr:EV ratio among the three

longest American alligators (*Alligator mississippiensis*) studied by Hurlburt et al (2013) was

found to be 31% (n_{total} = 12, note that this figure excludes the olfactory bulb and tract

portions of the endocranial cavity). This is consistent with observations on the largest Nile

crocodile (*Crocodylus niloticus;* a 16-year-old female) studied by Jirak & Janacek (2017)

when excluding the olfactory tracts and bulb portion of the endocast. The 42% ratio is

derived solely from American alligators. An endocranial fill of 42% was found in an adult

female with a total length of 2.87 m, which roughly approximates both (*a*) the maximum

length for a female American alligator and (*b*) the midpoint length within the size spectrum of

sexually mature alligators (Woodward et al, 1991; Hurlburt & Waldorf, 2002; Hurlburt et al,

2013).

The majority of EV data for Mesozoic dinosaurs were taken and modified from the literature

(detailed out in Tab. 1). In many cases, original sources communicated measurements that

correspond to total EV. This is the volume of the entire endocast, often including the region

of the olfactory tract and bulbs as well as portions of the cranial spinal cord among other

structures. For our analysis, we exclusively relied on the so-called "brain" endocast volume

instead (BrEV; Fig. 3), which was popularized by Jerison (1973) and has been commonly

used since then (e.g., Hurlburt, 1996; Larsson et al., 2000; Paulina- Carabajal & Canale,

2010; Hurlburt et al., 2013). It excludes the spinal cord portion of the endocast caudal to

cranial nerve XII, the volume of nerve trunks from infillings of respective foramina and blood

vessel casts, the labyrinth of the inner ear, the infundibulum, the pituitary fossa, and

especially the volume of the olfactory bulbs and tracts (Fig. 4). The latter are often only

poorly preserved in fossil endocasts, so that relying on specimens with intact casts of

olfactory structures would have reduced our sample size.

In some dinosaurs, there is an obvious constriction and/or a change in surface morphology

at the junction of the cerebrum and olfactory tract (e.g., *Euoplocephalus tutus* AMNH 5337 -

Hopson, 1979; *Stegosaurus ungulatus* CM 106 - Galton, 2001; *Diplodocus longus* CM

11161- Witmer et al., 2008). If present, this was used as a landmark to delineate these brain

regions from one another. In less obvious cases, the junction between the cerebrum and

olfactory tract portion was assumed to be where the ventral curve of the rostral cerebrum

flattens out to approach a horizontal orientation. When selecting the boundary, we erred

towards a more rostral location, to assign as much of the endocast as part of the BrEV as

seemed reasonable. In American alligators, the rostral termination of the cerebrum within the

rostral subarachnoid space is clearly visible (Fig. 2C; SaSR) and consistent with the change

in curvature referred to above.

We used manual GDI to extract BrEV from total EV (relevant details for each specimen are

provided in Supplementary File 1 Part B). The method involves drawing an outline around

two scaled orthogonal two-dimensional views of an endocast, and adding equally spaced

lines perpendicular to the endocast midline (Fig. 3). The mean length (cm) of these lines in

each view (i.e., dorsal, lateral) provides diameters D1 and D2. The volume (mL) of the

desired region is calculated using these two diameters and the length (L, cm) in the formula

for the volume (mL or cm^3) of a cylinder where (all lengths in cm):

$$Volume\ [mL]\ =\ (\pi)(0.25)(D1)(D2)(L)$$

GDI has been demonstrated to produce reasonable estimates of endocast volumes (Fig. 3).

For instance, Jerison (1973) used GDI to calculate a total endocast volume of 536 ml for a *T.*

rex specimen (AMNH 5029), which was 101.13% of the 530 ml volume determined for it by

means of water displacement (Osborn, 1912; Fig. 3). For the same specimen, Jerison (1973)

calculated a 404 ml volume for the "brain region" of the endocast (extending from cranial

nerve XII to the rostral cerebral limit), which was 106.04 % of the 381 ml obtained by CT

volumetry (Hurlburt et al., 2013).

Body mass estimates

We calculated the body mass (MBd, g) of the selected dinosaur taxa (and its mean absolute

percent prediction error - PPE or %PE - Campione & Evans, 2020) in a standardized manner

based on the minimum femoral circumference (as well as humeral circumference in case of

quadrupedal taxa) by aid of the *QE()* and *cQE()* functions from the MASSTIMATE package

(Campione, 2020) in R (R Core Team, 2023). Data on relevant long bone dimensions were

primarily obtained from Benson et al. (2017). Corresponding specimens as well as additional

sources and information on stylopodium circumference measurements are listed in Tab. 1.

To the best of our knowledge, all data correspond to adult specimens.

Figure 3. Exemplary graphic double integration (GDI) of the endocast of *Tyrannosaurus rex* AMNH

5029. Equally spaced lines are drawn across the right lateral and dorsal views respectively. Mean

lengths of the lines drawn across the "brain" portion (BrEV) were 4.8 cm and 6.6 cm for dorsal and

lateral views respectively. BrEV = π x 0.25 x 4.8 x 6.6 x 16.2 = 404 mL (the volume of the entire

endocast was 536 mL). Abbreviations: "Bulb": Olfactory tract and bulbs; "Cord": spinal cord; V:

Trigeminal nerve with its three branches (V1, V2,& V3); X: vagus nerve; XII: hypoglossal nerve.

(Adapted from Fig. 2.7 in Jerison,1973, p. 51).

Table 1: Estimates of brain (MBr, g; derived from endocranial volume, EV) and body mass (MBd, kg) in Mesozoic dinosaurs. For maniraptoriform theropods, we assumed that brain endocast volume equals brain volume. For other dinosaurs, we assumed a brain:endocast ratio of 31 - 42 %. Body mass was calculated based on stylopodial circumference (femoral circumference (FC) for bipedal and femoral as well as humeral (HC) circumference for quadrupedal species). **Footnotes: [1]** *Bambiraptor feinbergi*. MBd from femur circumference, measured on a cast of the right femur of an adult specimen now in the collection of the Vertebrate Paleontology Division, ROM (FIP 002). Original elements of this specimen are now in the collection of the AMNH. **[2]** *Archaeopteryx lithographica*. MBd from FC (14.93 mm) estimated from femur length (60.5mm) of BMNH 37001 (Gatesy, 1991). FC was calculated using the equation from Benson et al. (2017): log10(Femur circumference estimate from femur length) = 1.132 * log10(Femur length) - 0.8429. **[3]** *Carcharodontosaurus saharicus.* MBd from estimated femur circumference (FC = 417.52 mm) from Benson et al. (2017) for BSP 1922 X46. The specimen has been destroyed and is no longer accessible. **[4]** *Sinraptor dongi*. MBd based on FC (283 mm) from Campione & Evans (2020), measurement taken from TMP 93.115.1 (cast of IVPP 10600). Benson et al. (2017) consider IVPP 10600 a subadult based on reported incomplete fusion of cervical vertebrae; However, Paulina-Carabajal & Currie (2012) noted that the degree of cranial suture fusion indicates that the specimen is an adult or at least a large subadult. We consider it an adult here. **[5]** *Tyrannosaurus rex* specimens. MBd based on FC values listed by Persons et al. (2020). For FMNH PR 2081 ("Sue"), but not the other individuals considered here, both FC and EV are available. We associated the EV (313.636 mL) of AMNH 5117 with the MBd (5515 kg) of BHI 3033 ("Stan"), as both specimens have been considered proxies for each other (G. M. Erickson, pers comm. to G.R. Hurlburt, 2005). The EV (381.8 mL) of AMNH 5029 is here linked to the MBd (6430 kg) of CM 9380 (holotype specimen) because it fell between MBd's associated with EVs of FMNH PR 3081 and AMNH 5117. Besides our brain:endocast ratios, we also apply the 57% ratio proposed by Morhardt (2016) to this species.

Species	Group	Body mass (MBd) (kg)	Lower 25% PPE (kg)	Upper 25% PPE (kg)	FC (mm)	HC (mm)	MBd Specimen	MBd Source for FC & HC	Brain Mass (MBr) (g)	MBr/EV	Br EV (mL/cm3)	EV Specimen	EV Meth	Original EV Source
Archaeopteryx lithographica[2]	Non-avian maniraptoriform	3.446	2.563	4.329	14.93	NA	BMNH 37001	Gatesy, 1991	1.52	1	1.52	BMNH 37001	CT	Dominguez Alonso et al. (2004)
Bambiraptor feinbergi[1]	Non-avian maniraptoriform	8.0632	5.9966	10.1298	47	NA	FIP 002	Cast of right femur	14	1	14	KUVP 129737	Water displacement	Burnham (2004)
Citipati osmolskae	Non-avian maniraptoriform	123.993	92.214	155.772	127	NA	IGM 100/978	Benson et al. (2017), #321	22.05	1	22.05	IGM 100/978	CT	Balanoff et al. (2013)
Khaan mckennai	Non-avian maniraptoriform	219.198	163.018	275.378	67.62	NA	IGM 100/1127	Benson et al. (2017), #452	8.8	1	8.8	IGM 100/973	CT	Balanoff et al. (2013)
Ornithomimus edmontonicus	Non-avian maniraptoriform	835.238	621.167	1.049.309	110	NA	ROM 851	Benson et al. (2017), #522	49.89	1	49.89	NMC 12228	GDI (this study)	Photos of endocast by GRH
Shuvuuia deserti	Non-avian maniraptoriform	3.0497	2.2681	3.8313	33	NA	IGM 100/1304	Benson et al. (2017), #574	1.52	1	1.52	IGM 100/977	CT	Balanoff et al. (2013), Balanoff et al. (2024)
Stenonychosaurus inequalis	Non-avian maniraptoriform	473.759	352.335	59.52	89.5	NA	MOR 748 (MTC)	Benson et al. (2017), #621	38.65	1	38.65	RTMP 86.36.457 & 79.8.1	CT	Morhardt (2016)
Acrocanthosaurus atokensis	Non-maniraptoriform theropod	3.454.954	2.569.449	4340.46	426	NA	NCSM 14345	Benson et al. (2017), #242	51.55	0.42	122.74	OMNH 10146	CT	Franzosa & Rowe (2005)
Acrocanthosaurus atokensis	Non-maniraptoriform theropod	3.454.954	2.569.449	4340.46	426	NA	NCSM 14345	Benson et al. (2017), #242	38.05	0.31	122.74	OMNH 10146	CT	Franzosa & Rowe (2005)
Allosaurus fragilis	Non-	2.541.814	1.890.347	3193.28	381	NA	AMNH 680	Benson et al.	41.37	0.42	98.5	UUVP 294	GDI (this study)	Photo of cast by GRH

	maniraptoriform theropod						(2017), #253						
Allosaurus fragilis	Non-maniraptoriform theropod	2.541.814	1.890.347	3193.28	381	NA	AMNH 680	Benson et al. (2017), #253	30.54	0.31	98.5	UUVP 294	GDI (this study)	Photo of cast by GRH
Carcharodontosaurus saharicus	Non-maniraptoriform theropod	3.269.147	2.431.265	4107.03	417.52	NA	BSP 1922 X46	Benson et al. (2017), #307	69.44	0.31	224	SGM-Din 1	CT	Larsson et al. (2000)
Carcharodontosaurus saharicus[3]	Non-maniraptoriform theropod	3.269.147	2.431.265	4107.03	417.52	NA	BSP 1922 X46	Benson et al. (2017), #307	94.08	0.42	224	SGM-Din 1	CT	Larsson et al. (2000)
Carnotaurus sastrei	Non-maniraptoriform theropod	1.641.829	1.221.028	2062.63	325	NA	MACN CH 894	Benson et al. (2017), #308	45.49	0.42	108.3	MACN CH-894	CT	Cerroni & Paulina-Carabajal (2019)
Carnotaurus sastrei	Non-maniraptoriform theropod	1.641.829	1.221.028	2062.63	325	NA	MACN CH 894	Benson et al. (2017), #308	33.57	0.31	108.3	MACN CH-894	CT	Cerroni & Paulina-Carabajal (2019)
Giganotosaurus carolinii	Non-maniraptoriform theropod	6.136.771	4.563.916	7.709.625	525	NA	MUCPv-Ch1	Benson et al. (2017), #399	94.5	0.42	225	MUCPv-CH 1	CT	Paulina-Carabajal & Canale (2010)
Giganotosaurus carolinii	Non-maniraptoriform theropod	6.136.771	4.563.916	7.709.625	525	NA	MUCPv-Ch1	Benson et al. (2017), #399	69.75	0.31	225	MUCPv-CH 1	CT	Paulina-Carabajal & Canale (2010)
Majungasaurus crenatissimus	Non-maniraptoriform theropod	1.614.201	1.200.481	2027.92	323	NA	FMNH PR 2278	Benson et al. (2017), #482	37.51	0.42	89.32	FMNH PR 2100	CT	Sampson & Witmer (2007)

Majungasaurus crenatissimus	Non-maniraptoriform theropod	1.614.201	1.200.481	2027.92	323	NA	FMNH PR 2278	Benson et al. (2017), #482	27.69	0.31	89.32	FMNH PR 2100	CT	Sampson & Witmer (2007)
Sinraptor dongi	Non-maniraptoriform theropod	1.122.287	834.645	1.409.929	283	NA	TMP 93.115.1	Campione & Evans, 2020	29.45	0.31	95	IVPP 10600	CT	Paulina-Carabajal & Currie (2012)
Sinraptor dongi[4]	Non-maniraptoriform theropod	1.122.287	834.645	1.409.929	283	NA	TMP 93.115.1	Campione & Evans, 2020	39.9	0.42	95	IVPP 10600	CT	Paulina-Carabajal & Currie (2012)
Tarbosaurus bataar	Non-maniraptoriform theropod	2.345.113	1744.06	2.946.165	370	NA	MPC-D 552/1	Benson et al. (2017), #610	66.86	0.42	159.2	PIN, no. 553-3/1	Estimated from latex half-cast	Saveliev & Alifanov (2007)
Tarbosaurus bataar	Non-maniraptoriform theropod	2.345.113	1744.06	2.946.165	370	NA	MPC-D 552/1	Benson et al. (2017), #610	49.35	0.31	159.2	PIN, no. 553-3/1	Estimated from latex half-cast	Saveliev & Alifanov (2007)
Tyrannosaurus rex	Non-maniraptoriform theropod	8070.46	6.002.001	10.138.919	580	NA	FMNH PR 2081	Persons et al., 2020	128.4	0.31	414.19	FMNH PR 2081	CT	Hurlburt et al (2013)
Tyrannosaurus rex	Non-maniraptoriform theropod	6.430.357	4.782.256	8.078.457	534	NA	CM 9380	Persons et al., 2020	160.34	0.42	381.76	AMNH 5029	CT	Hurlburt et al (2013)
Tyrannosaurus rex	Non-maniraptoriform theropod	6.430.357	4.782.256	8.078.457	534	NA	CM 9380	Persons et al., 2020	118.35	0.31	381.76	AMNH 5029	CT	Hurlburt et al (2013)
Tyrannosaurus rex	Non-maniraptoriform theropod	5.515.247	4101.69	6.928.805	505	NA	BHI 3033	Persons et al., 2020	178.77	0.57	313.64	AMNH 5117	CT	Morhardt (2016)

Tyrannosaurus rex	Non-maniraptoriform theropod	5.515.247	4101.69	6.928.805	505	NA	BHI 3033	Persons et al., 2020	131.73	0.42	313.64	AMNH 5117	CT	Hurlburt et al (2013)
Tyrannosaurus rex	Non-maniraptoriform theropod	5.515.247	4101.69	6.928.805	505	NA	BHI 3033	Persons et al., 2020	97.23	0.31	313.64	AMNH 5117	CT	Hurlburt et al (2013)
Tyrannosaurus rex [5]	Non-maniraptoriform theropod	8070.46	6.002.001	10.138.919	580	NA	FMNH PR 2081	Persons et al., 2020	173.96	0.42	414.19	FMNH PR 2081	CT	Hurlburt et al (2013)
Amargasaurus cazaui	Non-theropod dinosaur	10194.61	7581.73	12807.49	505	388	MACN-N 15	Benson et al. (2017), #20	35.28	0.42	84	MACN-N 15	CT	Paulina Carabajal et al. (2014)
Amargasaurus cazaui	Non-theropod dinosaur	10194.61	7581.73	12807.49	505	388	MACN-N 15	Benson et al. (2017), #20	26.04	0.31	84	MACN-N 15	CT	Paulina Carabajal et al. (2014)
Apatosaurus sp.	Non-theropod dinosaur	41.268.719	30.691.546	51.845.891	845	640	CM 3018	Benson et al. (2017), #33	43.04	0.42	102.48	BYU 17096	CT	Balanoff et al. (2010)
Apatosaurus sp.	Non-theropod dinosaur	41.268.719	30.691.546	51.845.891	845	640	CM 3018	Benson et al. (2017), #33	31.77	0.31	102.48	BYU 17096	CT	Balanoff et al. (2010)
Buriolestes schultzi	Non-theropod dinosaur	6.424	4.777	08.07	43.27	NA	CAPPA/UFSM 0035	Müller et al. (2021)	1.021	0.42	2.43	CAPPA/UFSM 0035	CT	Müller et al. (2021)
Buriolestes schultzi	Non-theropod dinosaur	6.424	4.777	08.07	43.27	NA	CAPPA/UFSM 0035	Müller et al. (2021)	0.753	0.31	2.43	CAPPA/UFSM 0035	CT	Müller et al. (2021)
Diplodocus sp.	Non-theropod dinosaur	14.813.081	11.016.488	18.609.673	563	460	USNM 10865	Benson et al. (2017), #86	42	0.42	100	CM 11161	CT	L. M. Witmer, pers. comm. (2023)
Diplodocus sp.	Non-theropod dinosaur	14.813.081	11.016.488	18.609.673	563	460	USNM 10865	Benson et al. (2017), #86	31	0.31	100	CM 11161	CT	L. M. Witmer, pers. comm. (2023)

Species	Type						Specimen	Reference				Specimen	Method	Reference
Edmontosaurus annectens	Non-theropod dinosaur	6.610.079	4.915.916	8.304.243	512.3	250.5	AMNH 5730	Benson et al. (2017), #757	126	0.42	300	YPM 618	GDI (Jerison, 1973)	Lull & Wright (1942)
Edmontosaurus annectens	Non-theropod dinosaur	6.610.079	4.915.916	8.304.243	512.3	250.5	AMNH 5730	Benson et al. (2017), #757	93	0.31	300	YPM 618	GDI (Jerison, 1973)	Lull & Wright (1942)
Euoplocephalus tutus	Non-theropod dinosaur	2.329.632	1.732.548	2.926.717	278	244	AMNH 5404	Benson et al. (2017), #766	34.73	0.42	82.7	AMNH 5337	GDI (this study)	Hopson (1979)
Euoplocephalus tutus	Non-theropod dinosaur	2.329.632	1.732.548	2.926.717	278	244	AMNH 5404	Benson et al. (2017), #766	25.64	0.31	82.7	AMNH 5337	GDI (this study)	Hopson (1979)
Giraffatitan brancai	Non-theropod dinosaur	34.003.143	25.288.137	42.718.148	730	654	HMN SII	Benson et al. (2017), #107	130.2	0.42	310	MB.R.2223.1	Plasticine cast	Janensch (1935-36)
Giraffatitan brancai	Non-theropod dinosaur	34.003.143	25.288.137	42.718.148	730	654	HMN SII	Benson et al. (2017), #107	96.1	0.31	310	MB.R.2223.1	Plasticine cast	Janensch (1935-36)
Hypacrosaurus altispinus	Non-theropod dinosaur	3.689.151	2.743.622	4.634.681	395	222	CMN 8501	Benson et al. (2017), #800	85.53	0.31	275.9	ROM 702	CT	Evans et al (2009)
Hypacrosaurus altispinus	Non-theropod dinosaur	3.689.151	2.743.622	4.634.681	395	222	CMN 8501	Benson et al. (2017), #800	115.88	0.42	275.9	ROM 702	CT	Evans et al. (2009)
Iguanodon bernissartensis	Non-theropod dinosaur	8.268.265	6.149.108	10.387.421	490	337.5	RBINS R51	Benson et al. (2017), #805	149.94	0.42	357	RBINS R51	CT	Lauters et al. (2012)
Iguanodon bernissartensis	Non-theropod dinosaur	8.268.265	6.149.108	10.387.421	490	337.5	RBINS R51	Benson et al. (2017), #805	110.67	0.31	357	RBINS R51	CT	Lauters et al. (2012)
Kentrosaurus aethiopicus	Non-theropod dinosaur	1.596.86	1.187.585	2.006.136	245	210	HMN composite specimen	Benson et al. (2017), #813	22.092	0.42	52.6	HMN Ki 124	GDI (this study)	Galton (1988)
Kentrosaurus aethiopicus	Non-theropod dinosaur	1.596.86	1.187.585	2.006.136	245	210	HMN composite	Benson et al. (2017), #813	16.306	0.31	52.6	HMN Ki 124	GDI (this study)	Galton (1988)

						specimen							
Protoceratops andrewsi	Non-theropod dinosaur	82.695	61.5	103.889	93	62	AMNH 6424	Benson et al. (2017), #892	12.6	0.42	30	AMNH 6466	GDI (Jerison, 1973)	Brown & Schlaikjer (1940)
Protoceratops andrewsi	Non-theropod dinosaur	82.695	61.5	103.889	93	62	AMNH 6424	Benson et al. (2017), #892	9.3	0.31	30	AMNH 6466	GDI (Jerison, 1973)	Brown & Schlaikjer (1940)
Psittacosaurus lujiatunensis	Non-theropod dinosaur	28.61	21.278	35.944	74.5	NA	AMNH 6541	Benson et al. (2017), #898	6.006	0.42	14.3	PKUP V1060	CT	Zhou et al. (2007)
Psittacosaurus lujiatunensis	Non-theropod dinosaur	28.61	21.278	35.944	74.5	NA	AMNH 6541	Benson et al. (2017), #898	4.433	0.31	14.3	PKUP V1060	CT	Zhou et al. (2007)
Stegosaurus ungulatus	Non-theropod dinosaur	6.953.916	5.171.627	8.736.205	425	352	YPM 1853	Benson et al. (2017), #927	26.964	0.42	64.2	CM 106	GDI (this study)	Galton (2001)
Stegosaurus ungulatus	Non-theropod dinosaur	6.953.916	5.171.627	8.736.205	425	352	YPM 1853	Benson et al. (2017), #927	19.902	0.31	64.2	CM 106	GDI (this study)	Galton (2001)
Thescelosaurus neglectus	Non-theropod dinosaur	338.505	251.746	425.263	183	NA	AMNH 5891	Benson et al. (2017), #946	11.614	0.42	27.653	NCSM 15728	CT	Button & Zanno (2023)
Thescelosaurus neglectus	Non-theropod dinosaur	338.505	251.746	425.263	183	NA	AMNH 5891	Benson et al. (2017), #946	8.572	0.31	27.653	NCSM 15728	CT	Button & Zanno (2023)
Triceratops sp.	Non-theropod dinosaur	13.274.61	9.872.328	16.676.893	493	490	AMNH 5033	Benson et al. (2017), #950	96.075	0.42	228.75	MOR 1194	CT	Morhardt (2016), L. M. Witmer, pers. comm. (2024)
Triceratops sp.	Non-theropod dinosaur	13.274.61	9.872.328	16.676.893	493	490	AMNH 5033	Benson et al. (2017), #950	70.913	0.31	228.75	MOR 1194	CT	Morhardt (2016), L. M. Witmer, pers. comm. (2024)

Table 2: Estimates of telencephalic neuron counts (*N*; excluding the olfactory system) in Mesozoic dinosaurs. Our inferences are compared with those presented by Herculano-Houzel (2023; HH) if respected species were included in both studies (see text for the rationale of our sample composition). Minimum and maximum estimates based on both avian and non-avian sauropsid regressions are provided.

Species	Group	MBr range (g)	*N* (non-avian_min)	*N* (non-avian_max)	*N* (avian_min)	*N* (avian_max)	MBr (g), HH	*N* (non-avian), HH	*N* (avian), HH
Archaeopteryx lithographica	Non-avian maniraptoriform	1.52-1.52	16,2M	16,2M	56,5M	56,5M	1.47-1.76	15.8M-17.7M	54.2M-62.1M
Bambiraptor feinbergi	Non-avian maniraptoriform	14.00	69,2M	69,2M	277,7M	277,7M	14	62.9M	295.8M
Citipati osmolskae	Non-avian maniraptoriform	22.05	93,2M	93,2M	384,6M	384,6M	22.62	84.4M	424.5M
Khaan mckennai	Non-avian maniraptoriform	8.80	51,1M	51,1M	199,1M	199,1M	8.83	47.4M	209.1M
Ornithomimus edmontonicus	Non-avian maniraptoriform	49.89-49.89	159,1M	159,1M	690,7M	690,7M	87.85	193.6M	1,179M
Shuvuuia deserti	Non-avian maniraptoriform	1.52	16.1M	16.1M	56.5M	56.5M	0.83	11.2M	35.2M
Stenonychosaurus inequalis	Non-avian maniraptoriform	38.65	134.1M	134.1M	575.2M	575.2M	41	121.4M	664.4M
Acrocanthosaurus atokensis	Non-maniraptoriform theropod	38.05-51.55	133,2M	162,5M	568,8M	707,2M	191	311.3M	2,116M
Allosaurus fragilis	Non-maniraptoriform theropod	30.54-41.37	115,3M	140,7M	485,8M	603,9M	168	287.8M	1,921M
Carcharodontosaurus saharicus	Non-maniraptoriform theropod	69.44-94.08	197,5M	241M	875,6M	1,088M			
Carnotaurus sastrei	Non-maniraptoriform theropod	33.57-45.49	122,7M	149,7M	519,9M	646,5M			
Giganotosaurus carolinii	Non-maniraptoriform theropod	69.75-94.50	198,1M	241,7M	878,4M	1,092M			
Majungasaurus crenatissimus	Non-maniraptoriform theropod	27.69-37.51	108,2M	131,9M	452,9M	563,1M			

Sinraptor dongi	Non-maniraptoriform theropod	29.45-39.90	112,6M	137,4M	473,3M	588,5M			
Tarbosaurus bataar	Non-maniraptoriform theropod	49.35-66.86	157,9M	192,7M	685,4M	852,1M			
Tyrannosaurus rex AMNH 5029	Non-maniraptoriform theropod	118.35-160.34	280,1M	341,7M	1,283M	1,595M	343	445.5M	3,289M
Tyrannosaurus rex AMNH 5117 (Morhardt, 2016)	Non-maniraptoriform theropod	178.37	365M	365M	1722.1M	1722.1M			
Tyrannosaurus rex AMNH 5117	Non-maniraptoriform theropod	97.23-131.73	246,2M	300,4M	1,114M	1,385M			
Tyrannosaurus rex FMNH PR 2081	Non-maniraptoriform theropod	128.40-173.96	295,4M	360,5M	1,360M	1,691M	202	322.2M	2,207M
Amargasaurus cazaui	Non-theropod dinosaur	26.04-35.28	103,9M	126,8M	433,4M	538,8M			
Apatosaurus sp.	Non-theropod dinosaur	31.77-43.04	118,4M	144,4M	499,8M	621,4M			
Buriolestes schultzi	Non-theropod dinosaur	0.75-1.02	10.2M	12.4M	34.1M	42.5M			
Diplodocus sp.	Non-theropod dinosaur	31.00-42.00	116,4M	142,1M	491,1M	610,6M	57	148.5M	851.4M
Edmontosaurus annectens	Non-theropod dinosaur	93.00-126.00	239,2M	291,9M	1,079M	1,342M	150	268.5M	1,764M
Euoplocephalus tutus	Non-theropod dinosaur	25.64-34.73	102,9M	125,5M	428,6M	532,8	41	121.4M	664.4M
Giraffatitan brancai	Non-theropod dinosaur	96.10-130.20	244,4M	298,2M	1,105M	1,374M	186	306.3M	2,075M
Hypacrosaurus altispinus	Non-theropod dinosaur	85.53-115.88	226,4M	276,2M	1,016M	1,264M			
Iguanodon bernissartensis	Non-theropod dinosaur	110.67-149.94	268,1M	327,1M	1,223M	1,520M	125	240.2 M	1.538 M
Kentrosaurus aethiopicus	Non-theropod dinosaur	16.31-22.09	76,5M	93,3M	309,8M	385,2M	24	87.5M	443.9M
Protoceratops andrewsi	Non-theropod dinosaur	9.30-12.60	52,9M	64,5M	207,1M	257,5M	28	96.1M	498.5M
Psittacosaurus lujiatunensis	Non-theropod dinosaur	4.43-6.01	32.5M	39.6M	121.8M	151.4M			

Stegosaurus ungulatus	Non-theropod dinosaur	19.90-26.96	87,1M	106,3M	357,4M	444,3M	22.5	84.1M	422.8M
Thescelosaurus neglectus	Non-theropod dinosaur	8.57-11.61	49.9M	60.9M	195.4M	242.9M			
Triceratops sp.	Non-theropod dinosaur	70.91-96.08	199.5M	243.4M	888.9M	1105.1M	72.2	171.7M	1,017M

Phylogenetic modeling of neurological variables

We used data from extant sauropsids to place brain size variables for Mesozoic dinosaurs into their phylogenetic context. To examine variations in the relative size of the brain in fossil taxa and to calculate potential neuronal scaling regimes in extinct dinosaurs, we relied on log-transformed published data on brain mass, telencephalon mass and telencephalic neuronal numbers in extant groups (see below). Allometric equations were calculated with least squares linear regressions using phylogenetic generalized least squares (PGLS) to account for phylogenetic relatedness (Garland and Ives, 2000). PGLS allows the covariance matrix to be modified to accommodate the degree to which trait evolution deviates from Brownian motion, through a measure of phylogenetic correlation, Pagel's λ (Pagel, 1999). PGLS and maximum likelihood estimates of λ were performed using the *ape* (Paradis & Schliep, 2019) and *nlme* (Pinheiro et al., 2017) packages in R.

To compare differences in relative brain size across groups, phylogenetically corrected ANCOVA with Tukey post hoc comparisons were performed using a modified version of the *multcomp* package (Hothorn et al., 2015; modification allowed outputs of the *nlme* package (*gls* objects) to be processed). Because of the uncertainty in estimating both brain mass and body mass in Mesozoic dinosaurs, we opted to test for inter-group differences in two datasets: One with the greatest possible relative brain size i.e., the lowest body mass estimate (lower PPE) for each species and the brain mass estimated from the highest assumed endocranial fill (42%), and one with the lowest relative brain size i.e. the highest body mass estimates (upper PPE) and the lowest assumed endocranial fill (31%). Since we assumed that the brain filled 100% of the endocranial cavity in maniraptoriform theropods, inferred relative brain mass for these species was only affected by differences in the applied body mass estimates.

As mentioned above, an important assumption of Herculano-Houzel (2023) is that theropods in general had relative brain sizes similar to birds. However, there is notable discontinuity in

relative brain size and brain morphology between maniraptoriforms and more basal non-maniraptoriform theropods (Fig. 1, Balanoff et al., 2013). Because of this, we divided our sample of Mesozoic theropods into these two groups (Tabs. 1 & 2). For *T. rex*, mean values for the three available adult brain mass and corresponding body mass estimates were used. We grouped Sauropodomorpha and Ornithischia together as non-theropod dinosaurs and compared relative brain size in this group with that in the two theropod samples. PGLS models of brain mass vs. body mass with clade as a covariate were used to test if relative brain size in these groups differs significantly between them and from extant birds and/or non-avian sauropsids. Relative brain size data for 63 extant non-avian sauropsids (including lepidosaurs, crocodilians, and turtles) and 84 bird species (not including members of the large-brained clade Telluraves) were derived from Hurlburt (1996), Chentanez et al. (1983) and Roese-Miron et al. (2023) and are listed in Supplementary File 2. Importantly, these sources provide brain mass estimates excluding the olfactory tracts and bulbs and thus fit our brain size estimates for Mesozoic dinosaurs.

To test for differences in relative brain size we built a phylogenetic tree for all 175 fossil and extant species. To construct the phylogeny of bird species, we extracted 1,000 fully resolved trees from birdtree.org (Jetz et al., 2012) using the Hackett et al. (2008) backbone, and built a maximum clade credibility (MCC) tree using *phangorn* (Schliep, 2011). For lepidodaurs, we followed Kverková et al. (2022) by using a species level time-calibrated phylogeny (Tonini et al., 2016) and built a MCC tree the same way as we did for birds. For phylogenetic information on turtles and crocodilians, we relied on the Timetree of Life (Kumar et al., 2017). We then stitched the trees together manually, using the divergence times from the Timetree of Life. For Mesozoic dinosaurs (31 species) we used an updated version of the composite phylogeny of Benson et al. (2014, 2018). Phylogenies for fossil dinosaurs, extant non-avian sauropsids, and birds were stitched together manually using Mesquite (Maddison & Maddison, 2023). We opted to set all branch lengths to 1. This was done because clade-specific trees were obtained from various sources applying different phylogenetic methods,

which, together with issues related to the precise dating of some of the fossils covered,

makes it difficult to have well calibrated branch lengths. Importantly, simulation studies have

found that independent contrasts and PGLS are robust to errors in both phylogenetic

topology and branch lengths, so that we do not expect uniform branch lengths to

compromise our analyses (Díaz-Uriarte & Garland 1998; Symonds 2002; Martins &

Housworth 2002; Stone 2011).

Tree building procedures were the same for telencephalic neuron count regressions, but

trees used here included branch lengths. We calculate regression lines between brain mass

and telencephalic number of neurons for extant non-telluravian birds and non-avian

sauropsids. Analogous to Herculano-Houzel (2023), these regressions were then used to

estimate telencephalic neuron counts in dinosaurs, applying either an avian or a reptilian

scaling regime. Since our estimates are based on brain portion endocasts that exclude the

olfactory system, our telencephalic neuron counts correspond to the pallium and subpallium.

Data on whole brain and telencephalic neuron counts as well as on total telencephalic and

brain mass (including olfactory tract and bulbs, since neuron count data excluding these

structures are currently unavailable for the taxa considered here) for birds ($n = 112$) were

derived from Kverková et al. (2022) and Sol et al. (2022), for non-avian sauropsids ($n = 108$)

from Kverková et al. (2022) and for mammals ($n = 39$) from Herculano-Houzel et al. (2015).

The dataset is included in Supplementary File 3.

To get a more precise estimate of the possible number of telencephalic neurons in *T. rex*, we

also modeled scaling regimes for telencephalon mass vs. telencephalic neuron numbers in

extant sauropsids (non-avian sauropsids and non-telluravian birds), using the same

references listed above. We then calculated telencephalic neuron numbers in *T. rex* using

the obtained scaling regimes and applying the telencephalic volumes estimated with a

comparative 3D landmark approach by Morhardt (2016) (referred to as "cerebral

hemispheres" therein, excluding olfactory bulbs and tracts) for specimen AMNH FR 5117.

Based on the estimates of Morhardt (2016), we also comparatively assessed the mass of the

telencephalon and cerebellum in *T. rex*.

We want to note that our neuron count estimates might be biased by the fact that we predict

neuron numbers in the pallium and subpallium (telencephalon excluding the olfactory

system) based on total telencephalic neuron counts (including the olfactory system) here.

This is an issue that in parts also applies to Herculano-Houzel (2023) and which we cannot

circumvent due to limitations of the available raw data.

Results

Relative brain size.

We did not recover notably large relative brain sizes in large-bodied theropods like *T. rex*.

Instead, our analyses suggest that these animals had relative brain dimensions comparable

to extant non-avian sauropsids such as lizards and crocodilians, as did Mesozoic dinosaurs

outside of the clade Theropoda (Table 3). Relative brain sizes similar to those of extant birds

appear to only have emerged among the maniraptoriform theropods: PGLS models showed

a significant difference in relative brain size (intercept) between non-maniraptoriform

theropods, such as *T. rex*, and the more bird-like Maniraptoriformes, which tended to have

larger brains than other dinosaurs (PGLS, max: $F_{4,172}$= 9.49, $p \leq$ 0.0001, λ = 0.707); min:

$F_{4,172}$= 14.03, $p \leq$ 0.0001, λ = 0.707; Fig. 4a-b). Post-hoc analysis shows that both the

maximum and minimum relative brain size estimates for non-maniraptoriform theropods like

T. rex are not significantly different from what would be expected from extant non-avian

sauropsids (Tab. 3). However, both minimum and maximum relative brain size estimates for

these animals are significantly smaller than what would be expected for extant birds (Tab. 3;

Fig. 4). On the other hand, we found that maniraptoriforms show no significant differences in

relative brain size compared to birds (Tab. 3) regardless of whether maximum or minimum

relative brain size was assumed (Tab. 3; Fig 4; note that some maniraptoriforms such as
Shuvuuia deserti cluster with non-avian sauropsids rather than with birds, though). In
contrast to maniraptoriforms, other theropods did not exhibit significantly larger brains than
the non-theropod dinosaurs of the clades Sauropodomorpha and Ornithischia (Tab. 3), data
for which we pool here. Relative brain sizes in these dinosaurs were not recovered to differ
notably from those of non-avian sauropsids. However, if minimum figures are assumed, their
relative brain sizes would have been notably small, approaching a significant difference to
extant non-avian sauropsids (Tab. 3).

Table 3: Tukey post hoc comparisons for a phylogenetic ANCOVA testing for differences in relative
brain size between groups of fossil dinosaurs and extant sauropsids. Significant *p*-values are shown
in bold.

Minimum relative brain size				
	Aves	Non-avian Maniraptoriformes	Non-maniraptoriform Theropoda	Non-theropod Dinosauria
Non-avian Maniraptoriformes	0.57			
Non-maniraptoriform Theropoda	**> 0.00001**	**> 0.00001**		
Non-theropod Dinosauria	**> 0.00001**	**> 0.00001**	0.103	
Non-avian Sauropsida	**> 0.00001**	**0.033**	0.96	0.051
Maximum relative brain size				
	Aves	Non-avian Maniraptoriformes	Non-maniraptoriform Theropoda	Non-theropod Dinosauria
Non-Avian	0.61			

Maniraptoriformes				
Non-Maniraptoriform Theropoda	**0.000106**	**0.0015**		
Non-theropod Dinosauria	**> 0.00001**	**> 0.00001**	0.103	
Non-avian Sauropsida	**> 0.00001**	**0.0033**	0.91	0.79

643 **Table 4**: Regression parameters for different models describing the scaling of neurological traits in

644 extant non-avian sauropsids ("reptiles") and non-telluravian birds. Pagel's λ (ranging between 0 - 1)

645 was used to quantify the phylogenetic signal. See methods for details. SE = standard error.

	Model	Slope (SE)	Intercept (SE)	λ
Avian GLS	log(Tel Neurons N) ~ log(Brain Mass)	0.821 (0.043)	17.5 (0.063)	0
Avian PGLS	Log(Tel Neurons N) ~ log(Brain Mass)	0.717 (0.05)	17.55 (0.12)	0.50
Reptilian GLS	Log(Tel Neurons N) ~ log(Brain Mass)	0.615 (0.03)	16.347 (0.06)	0
Reptilian PGLS	Log(Tel Neurons N) ~ log(Brain Mass)	0.655 (0.03)	16.324 (0.17)	0.82
Avian PGLS	log(Brain Mass) ~ log(Body Mass)	0.584 (0.02)	-2.584 (0.25)	0.96
Reptilian PGLS	log(Brain Mass) ~ log(Body Mass)	0.56 (0.03)	-4.077 (0.26)	0.70

Numbers of neurons

We re-calculated estimates for the number of forebrain neurons in Mesozoic dinosaurs

based on PGLS-derived regressions of brain size vs. number of telencephalic neurons in

extant non-avian sauropsids and birds. Our neuron count estimates are listed in Table 2 and

are compared to those of Herculano-Houzel (2023), whereas regression parameters are

provided in Table 4. While many of the estimates do not differ notably from one another, the

differences for some taxa, especially large theropods, are striking. For *T. rex*, Herculano-

Houzel (2023) provided an estimate of 300-450 M forebrain neurons if modeled based on

extant non-avian sauropsids, and 2-3B based on an avian regression. In contrast, we

estimated a range of 245-360M neurons with a reptilian regression and ~1-2 B with an avian

one (Tab. 2, Fig. 5). Using the forebrain volumes estimated for *T. rex* by Morhardt (2016), we

predict 133-166 M telencephalic neurons in this species if applying a reptilian scaling and

0.989 to 1.25 B based on an avian scaling (Fig. 5).

Figure 4: Relative brain size and forebrain neuronal numbers in Mesozoic dinosaurs and other

amniotes. a. The log-transformed mass of the brain is plotted as a function of the mass of the body for

extant and fossil sauropsids. In the case of fossil species, the mean body and/or brain size is shown

along with standard deviations. The orange dotted line represents the regression line for avian

species (excluding the large-brained clade Telluraves) obtained from PGLS while the pink one

represents the same for extant non-avian sauropsids ("reptiles" in the colloquial sense). **b.** A detail of

the plot shown in **a**. to illustrate the range of relative brain size in *Tyrannosaurus rex* and other

Mesozoic dinosaurs that we consider plausible. Besides our own brain size estimates, the plot

contains those from Morhardt (2016) (specimen AMNH FR 5117, endocranial fill = 57%) and Balanoff

et al. (2013) (specimen AMNH 5029, endocranial fill = 100%, assumed MBd = 5840 kg) **c.** Plot

showing log-transformed brain mass for different groups of extant amniotes plotted against body

mass. **d.** Plot showing log-transformed numbers of telencephalic neurons as a function of the mass of

the forebrain, illustrating neuronal density. Note that non-avian sauropsids and non-primate mammals

differ only moderately from one another here, although mammals have markedly larger brains relative

to body size, as shown in **c**. See methods for data sources. Silhouettes were taken from PhyloPic

(listed clockwise from top left): *Anas* (in public domain) *Morunasaurus* (in public domain),

Dromaeosaurus (in public domain), *Stegosaurus* (by Matt Dempsey), *Allosaurus* (by Tasman Dixon),

Tyrannosaurus (by Matt Dempsey), *Corvus* (in public domain), *Hylobates* (by Kai Caspar), *Antidorcas*

(by Sarah Wernig).

Figure 5: Predicted numbers of neurons in telencephalon (excluding olfactory tracts and

bulbs) of *Tyrannosaurus rex*. Points represent the estimated number of neurons in three adult

specimens of *T. rex* using different inference methods. Estimates from this study using a regression

that takes phylogenetic relationships into account (PGLS, see methods, filled circle, triangle, square

and cross) are plotted in cyan. The estimates from Herculano-Houzel (2023) based on a non-

phylogenetic regression are shown in red (crossed square, asterisk). Different underlying ratios of

brain volume:endocranial volume are annotated. On the left predicted numbers of forebrain neurons

(based on either the avian or extant non-avian sauropsid scaling regime) based on the estimated

volume of the brain portion of the endocast are shown. On the right, analogous to that, the predicted

numbers of telencephalic neurons based on forebrain volumetric estimates by Morhardt (2016) is

plotted.

Figure 6: Relative size of telencephalon (excluding olfactory bulb and tracts) and cerebellum in

T. rex. **a:** The log-transformed mass of the telencephalon in extant non-telluravian birds and non-

avian sauropsids is plotted as a function of the mass of the rest of the brain (total brain -

telencephalon - cerebellum volume). Green dots show the maximum, mid and minimum estimates for

the mass of the *T. rex* telencephalon as estimated from digital endocasts of AMNH 5117 by Morhardt

(2016). The orange dotted line represents the regression line for non-telluravian bird species obtained

from PGLS while the pink represents the same for non-avian sauropsids. **B**: Analogous plot to **a**, but

for the cerebellum. Note that telencephalic mass in extant species includes the olfactory bulb and

tracts.

Discussion of empirical results

We want to emphasize two aspects of our empirical findings that contrast with those of

Herculano-Houzel (2023). Firstly, we did not find relative brain size to separate non-

maniraptoriform theropods such as *T. rex* from extant non-avian sauropsids like crocodilians

and lizards or other large-bodied dinosaurs outside the clade Theropoda; rather, our data

support a grade shift in this trait between maniraptoriform and non-maniraptoriform

theropods, which at least in parts relates to an increase in endocranial fill. As we have

argued beforehand, we see no support for the brains of non-maniraptoriform theropods,

sauropodomorphs, and most ornithischians to have filled the endocranial cavity in a bird-like

fashion. However, we are aware that such a condition, or one that is at least intermediate

between modern birds and crocodilians has been proposed for all of these groups at one

point (e.g., Knoll & Schwarz-Wings, 2009; Morhardt, 2016; Balanoff et al., 2013; Knoll et al.,

2021; see Supplementary File 1 Part C for further comments on that topic). Obviously, future

research might significantly change our understanding of endocranial tissue organization in

Mesozoic dinosaurs and thus challenge the assumptions that we make here. For the time

being, however, we consider our crocodilian-based inferences more plausible and

parsimonious than the alternative suggestions proposed so far.

Our approach suggests that relative brain size in all dinosaurs, except for the majority of

maniraptoriform theropods, does not differ significantly from values present in extant non-

avian reptiles. These results agree with previous conclusions (Hurlburt et al., 2013;

Morhardt, 2016). Nonetheless, we want to stress that it remains unclear how meaningful the

transfer of brain size scaling rules established for the given extant bird (32g – 120kg) and

non-avian sauropsid (1g – 71kg) datasets to large-bodied dinosaurs actually is. Brain-body

size ratios in extant cetaceans drop dramatically in taxa that evolved multi-ton body masses

(Tartarelli & Bisconti, 2006), suggesting that allometric trajectories need to be accounted for.

However, the restricted body mass spectrum of extant birds and reptiles as well as the

limited availability of large-bodied crocodilians and turtles for neurological research hinders

the compilation of such datasets for sauropsids. Furthermore, the scarcity of complete and

adult dinosaur endocasts from taxa that also preserve stylopodial elements to derive body

mass estimates from, limits our understanding of differences in brain size scaling between

taxa. Different clades of mammals and birds have been shown to have distinct allometric

relationships for relative brain size (Ksepka et al., 2020; Smaers et al., 2021). The same

might have been the case in non-avian dinosaurs, biasing comparisons between the

groupings we selected here. In addition to that, there might also be temporal effects on

relative brain size. Such a phenomenon appears to be rampant in mammalian evolution

during the Cenozoic (Bertrand et al., 2022). To our knowledge, this pattern has not been

properly described yet in other vertebrate groups but should be considered in future studies

on brain evolution in long-lived clades such as dinosaurs.

Secondly, our empirical findings do not support Herculano-Houzel's (2023) claim of

exceptionally high telencephalic neuron counts in dinosaurs, particularly in *T. rex* and other

large theropods. Instead, *T. rex* likely did not exhibit more than approximately 1.5 B (or at a

maximum 2 B) telencephalic neurons, even when an avian neuronal density is assumed. If

we assume reptilian neuronal densities, it might even have exhibited neuron numbers an

order of magnitude lower than the 3.3 B suggested by Herculano-Houzel (2023). Apart from

the difficulty of estimating brain mass from a dinosaurian endocast, there is one additional

caveat to our neuron count estimates that needs to be acknowledged and that also applies

to Herculano-Houzel's (2023) study: Telencephalic neuron numbers can only be reliably

derived from total brain mass when the proportions of the studied brains are comparable.

Since brain morphology in many dinosaurian lineages differs significantly from both extant

birds and non-avian sauropsids (Fig. 1, Paulina-Carabajal et al., 2023), biases are thus

ingrained into such estimates. Volumetric modeling of brain regions from endocasts, on

which we relied here for *T. rex* exclusively, could potentially ameliorate this problem to some

degree (Morhardt, 2016) but it is challenging and not yet widely used. For *T. rex*, such

inferences yield lower telencephalic neuron numbers than would be hypothesized based on

our total brain volume estimates, if reptilian scaling rules are applied (Fig. 5). They overlap if

an avian neuron count scaling is assumed (Fig. 5).

We want to emphasize that there is little reason to assume that the brains of non-

maniraptoriform theropods such as *T. rex* had a telencephalic neuronal density similar to that

of extant birds. In living sauropsids, relative brain size is positively correlated with neural

density (Kverková et al., 2022). We show that this measure likely did not differ significantly

between those theropods and extant non-avian sauropsids. Consequently, relative brain size

cannot be used as an argument to defend elevated neuron densities in these animals. The

presence of endothermy in dinosaurs (see below) does also not entail avian neuronal

density (contra Herculano-Houzel, 2023): Similar to birds, mammals have evolved

endothermy and exhibit large relative brain sizes (Fig. 4C; Tsuboi et al., 2018). Furthermore,

they display a unique multilayered cerebral neuroarchitecture (Briscoe & Ragsdale, 2018).

Yet their average forebrain neuronal density is only moderately elevated compared to extant

non-avian sauropsids (at least if anthropoid primates are not considered) and there is a

broad overlap in neuronal density between the groups (Fig. 4D; Kverková et al., 2022),

indicating remarkable conservatism in this trait. With respect to birds, however, the typical

mammalian telencephalic neuron density is remarkably low (Fig. 4D). Interestingly, brain cell

densities (suggestive of high neuron counts but including endothelial and glia cells) on par

with or even higher than those of telluravian birds have recently been identified among

ectotherm teleost fish, with comparatively small relative brain sizes (Estienne et al., 2024).

All of this suggests that metabolic rate and neuronal density are not tightly coupled and that

endothermy cannot be used as a proxy for the latter. Finally, the shape of the endocast and

volumetric estimates of its forebrain and cerebellar portions (compare Fig. 6) suggest that

the brains of *T. rex* and other large non-maniraptoriform theropods were not dissimilar to

those of extant crocodilians (Rogers, 1998; Hurlburt et al., 2013; Morhardt, 2016), which

reflect the plesiomorphic archosaurian condition (Fabbri & Bhullar, 2022). Given this

morphological conservatism and the rather static neuron densities of non-avian amniotes, it

appears appropriate to assume reptilian neuronal densities for these animals.

It is tempting to speculate that the increased neuronal density that sets extant birds apart

from other sauropsids and mammals coevolved with the marked changes in brain

morphology and size that occurred in maniraptoriform theropods. If we indeed assume that

an avian-like brain organization and high neuronal density emerged early within this clade's

history, it is plausible that the Mesozoic dinosaurs with the highest neuron counts, perhaps

above the extant avian range, are represented by the largest-bodied taxa within this group

(for which no complete endocasts are currently available). These include bizarre animals

such as the immense ornithomimosaur *Deinocheirus mirificus* (~ 7 t), the scythe-clawed

Therizinosaurus cheloniformis (~ 5 t) and the giant oviraptorosaur *Gigantoraptor erlianensis*

(~ 2 t). Alternatively, the emergence of volancy in small maniraptoriforms similar to

Archaeopteryx might have driven the evolution of elevated neuron densities, since active

flight likely imposes constraints on skull and brain size (Olkowicz et al., 2016; Shatkovska &

Ghazali, 2021). However, the lack of reliable morphological markers to infer neuron density

renders all these notions speculative. Such a vagueness is inherent to predictions about the

biology of extinct taxa without close living relatives and obviously needs to be

acknowledged. The main argument for assuming avian neuronal densities in any group of

Mesozoic dinosaurs is that the emergence of this trait within the avian stem-lineage cannot

be reliably dated and thus might have significantly preceded the origins of crown birds.

Hence, both a non-avian sauropsid and an avian neuron density (as well as intermediate

conditions) could principally be justified for dinosaurs, although we advocate for the former if

taxa outside the Maniraptoriformes are concerned. Importantly, however, even if we had

robust evidence for high neuron counts in Mesozoic dinosaurs, this would by no means

automatically suggest exceptional cognitive capacities.

General discussion - implications for neuron count and brain size estimates for

vertebrate paleontology

1) Are neuron counts good predictors of cognitive performance?

To infer cognitive abilities in extinct animals from neuron count estimates for the whole brain

or pallium, we first need to be assured that this measure can give us meaningful insight into

behaviors of extant ones. However, while there is some evidence for effects of pallial neuron

counts on species-level cognitive performance in primates (Deaner 2007 - but see below)

and birds (impulse inhibition - Herculano-Houzel, 2017; but see Kabadayi et al. 2017 for

conflicting evidence; foraging-related innovativeness – Sol et al., 2022; but consider

limitations on how innovativeness is measured – Logan et al., 2018), the available data do

not provide consistent support for the hypothesis that more neurons per se enhance

cognition (Barron & Mourmourakis, 2023). As an example, Güntürkün et al. (2017) reviewed

the performance of domestic pigeons (*Columba livia*), corvids and anthropoid primates in a

number of cognitive tasks with the aim to determine if a "cognitive hierarchy" between the

three groups exists. They note that pallial neuron counts in corvids are about 2–6 times

lower than in large-bodied monkeys and apes but 6–17 times higher than in pigeons. Thus,

one would predict major increases in cognitive capacities from pigeons to corvids to

anthropoids. Yet, corvids typically perform on par with anthropoid primates (see also

Kabadayi et al. 2016 and Pika et al. 2020), and pigeons do so as well in some cognitive

dimensions, such as numerical competence and short-term memory (Güntürkün et al.,

2017). In addition, standardized testing of various primate species suggests that small-

brained lemurs with comparatively low neuronal densities (Kverková et al., 2022), monkeys

and great apes rival each other in a number of cognitive dimensions (Schmitt et al., 2012;

Fichtel et al., 2020). In fact, findings that report the influence of absolute brain size (and thus

neuron number) on cognitive performance in primates (Deaner et al. 2007) have failed

replication so far (Fichtel et al. 2020). As a final example, we want to point out large-bodied

dolphin species, which have remarkably high neocortical neuron counts (*Globicephala melas*

- 37 B, *Orcinus orca* - 43 B; Ridgway et al., 2019). Although neuron numbers in these

animals vastly exceed those of humans (15-20 B), there is no evidence that cetacean

cognition is on par or even superior to that of our species (e.g., Manger, 2013; Güntürkün,

2014). Hence, even immense differences in telencephalic neuron numbers do not

necessarily create cognitive divides and their value in predicting cognitive performance is
remarkably limited.

The case becomes even more untenable when we take more complex cognitive phenomena
into consideration, such as habitual tool use. Remarkably, Herculano-Houzel (2023)
suggested that this might be within the realm of possibility for large theropods such as *T. rex*,
as it is for primates and telluravian birds. However, tool use even within these groups is rare,
especially if the more rigorous definition of "tooling" (requiring the deliberate management of
a mechanical interface, see Fragaszy & Mangalam, 2018) is employed: this occurs in only 9
avian and 20 primate genera (Colbourne et al., 2021). While it is true that telencephalon size
in birds has an association with tool use (Lefebvre et al., 2002), this correlation does not hold
any predictive power in the sense that all birds with a certain-sized telencephalon exhibit tool
use. Even within corvids, which telencephalic neuron counts and sophisticated cognitive
abilities overlap with those of anthropoid primates (Olkowicz et al., 2016; Ströckens et al.,
2022), New Caledonian crows (*Corvus moneduloides*), and Hawaiian ʻalalā crows (*C.
hawaiiensis*) are the only species known to employ and manufacture tools in the wild.
Notably, both species inhabit remote islands, and they share unusually straight beaks and
greater binocular overlap than other crows, which are thought to be specific morphological
adaptations to enable tool use (Troscianko et al., 2012; Rutz et al., 2016). A similar situation
can be observed in parrots. These birds display the highest avian telencephalic neuron
counts (Olkowicz et al., 2016; Kverková et al., 2022; Ströckens et al., 2022), and a greatly
enlarged medial spiriform nucleus, which acts as an interface between the pallium and the
cerebellum, enabling enhanced motor cognition (Gutiérrez-Ibáñez et al., 2018). However,
the Tanimbar corella (*Cacatua goffiniana*) is the only parrot known to be a sophisticated tool
user in the wild (O'Hara et al., 2021); tellingly, the Tanimbar corella also inhabits an isolated
Indonesian archipelago. Cases like these indicate that while there might be a chance that a
gross neuron count threshold must be met for such sophisticated vertebrate tool use to
emerge (a notion we would reject since ants evolved remarkable tool use skills with brains

that are small and few in neurons even for the standard of hymenopteran insects - Godfrey

et al., 2021), it is highly unlikely to happen without sufficient ecological pressure, and the

differences between tool using and non-tool using species are likely too subtle to detect via

measurement of neuronal quantities.

Considering these findings, it is unsurprising that taxa converging in neuronal counts often

differ markedly in cognition and behavior. Herculano-Houzel (2023) ranked her neuronal

count estimates for large theropods against those of anthropoid primates, but she might as

well have done so for giraffes (1.7 B neurons), which exceed tool-proficient capuchins (1.1

B) and corvids (0.4-1.2 B) in telencephalic neuron numbers, rivaling macaques (0.8 - 1.7 B)

(Olkowicz et al., 2016). We know little about giraffes' cognitive abilities (Caicoya et al.,

2018), but it would be appropriate to be skeptical of any claim that they might exhibit

"macaque-like" cognition based simply on that measure. Too many other biological traits

divide these taxa, perhaps most strikingly body size. While we agree with many

contemporary authors that relative brain size per se is a flawed measure of cognitive

complexity (e.g., Van Schaik et al., 2021), it must not be ignored. This is especially true if

comparisons between primates and Mesozoic dinosaurs are drawn, since the species

concerned may differ in body mass by several orders of magnitude. Contrary to the

assumptions of Herculano-Houzel (2023), the size of the telencephalon and number of its

neurons must be related to the dimensions of the body, because it processes sensory,

visceral, and motoric information, which scale with body size (Chittka & Niven, 2009; Van

Schaik et al., 2021). This fact is clearly reflected by the pronounced intra- as well as

interspecific body size-dependent scaling of brain size in vertebrates (Tsuboi et al., 2018;

Ksepka et al., 2020; Van Schaik et al., 2021; Bertrand et al., 2022), which can hardly be

explained otherwise. Relative brain size and body size are thus not negligible variables in

comparative cognition and need to be considered in paleoneurology.

The confounding factor of body size on neurological measures might be mitigated by

calculating clade-specific portions of telencephalic mass dedicated to somatic functions (the

regulation of visceral, sensory and motor processes unrelated to cognition) based on

intraspecific variation (Triki et al., 2021; Van Schaik et al., 2021) or by focusing on neuron

counts in brain regions that are evidently not involved in somatic processing (Herculano-

Houzel, 2017; Logan et al., 2018). In fact, a number of studies, particularly in birds, were

able to associate intraspecific differences in certain cognitive dimensions to localized

neurological variation, making this approach a promising one (discussed by Logan et al.,

2018). At the same time however, the great intra- and interspecific heterogeneity in brain

tissue architecture and neurochemistry enormously complicates any interspecific

extrapolations (Logan et al., 2018; Barron & Mourmourakis, 2023). Thus, researchers cannot

translate these findings to extinct species with any tolerable degree of certainty. This issue is

of special relevance when comparing sauropsids with mammals. The mammalian forebrain

exhibits a layered cortex but the pallium of extant sauropsids (and thus likely Mesozoic

dinosaurs) is largely nuclear in organization. As the forebrain increases in size and neuron

counts, a cortical organization can reduce axon length (and therefore processing time and

energetic demands) by bringing adjacent areas closer together through folding, something

that is impossible in a nuclear organization (see Reiner, 2023 for an extensive review).

Neuron counts corresponding to major brain regions, whether empirically determined or

estimated, dramatically simplify neuronal tissue complexity, as do measures such as

absolute brain size or EQ. Based on current evidence, they also represent flawed cognitive

proxies that need to be viewed in the broader context of an animal's ecology, neuroanatomy,

connectomics, and neurochemistry (Fields & Stevens-Graham, 2002; Eyal et al., 2016;

Logan et al., 2018; Barron & Mourmourakis, 2023, Reiner, 2023). All in all, we want to

discourage attempts to predict cognitive performance in extinct species based on endocast-

derived neuron count estimates.

2) Inferring metabolic rate

Apart from inferences about cognition, Herculano-Houzel (2023) claims that relative brain

size should be established as a new thermobiological indicator in vertebrate palaeontology:

Relatively large brains, as inferred for theropods, should be viewed as indicators of

endothermy, while smaller ones, as are found in pterosaurs, sauropodomorphs and many

ornithischians, would indicate ectothermy. Whereas overall brain size in vertebrates is

indeed correlated with metabolic rate (e.g., Yu et al., 2014 - but also note the extreme

variability within ecto- as well as endothermic groups), Herculano-Houzel's (2023) approach

simplifies the matter and ignores a vast body of already available evidence on dinosaur

thermobiology. First, as we have extensively discussed here, relative brain size in large

theropods was probably markedly smaller than suggested by Herculano-Houzel (2023) and

more similar to the condition in extant crocodilians rather than in birds. Second, it is

important to point out that there is a spectrum of metabolic rates in vertebrates (Legendre &

Davesne, 2020) rather than a dichotomy, as suggested by Herculano-Houzel (2023).

Where exactly certain ornithodiran taxa align within this spectrum continues to be debated,

but there is consensus that dinosaurs and pterosaurs, despite their in parts iconically small

brains, had metabolic rates well above the range of extant ectothermic sauropsids (see

references below). Rather than emerging with theropods, contemporary evidence suggests

that endothermy evolved in the ornithodiran stem-lineage or even earlier (Legendre et al.,

2016; Benton, 2021; Grigg et al., 2022) and hence was inherited by pterosaurs and

dinosaurs. The extensive data supporting the presence of endothermy across Ornithodira

has recently been reviewed by Grigg et al. (2022) and includes the presence of hair-like,

sometimes branched, integumentary structures (Benton et al., 2019; Campione et al., 2020),

the efficiency of the ornithodiran respiratory system (Wedel, 2006; Butler et al., 2009;

Aureliano et al., 2022; Wang et al., 2023), bone histology and high skeletal growth rates (de

Ricqlès et al., 2000; Padian et al. 2004; Prondvai et al., 2012; Redelstroff et al. 2013;

Legendre et al., 2016), paleoenvironmental data (Druckenmiller et al., 2021), models of

locomotor costs (Pontzer et al., 2009) and geochemically-derived thermometric findings

(Barrick et al., 1996; Dawson et al. 2020; Wiemann et al., 2022). Nevertheless,

osteohistological evidence suggests that both theropod and non-theropod ornithodiran taxa

varied in their growth and associated metabolic rates (Jenkins et al., 2001; Erickson et al.,

2009; Redelstroff et al. 2013; D'Emic et al., 2023) and a secondary reduction of metabolic

rate in some ornithischian groups appears plausible (Padian et al., 2004; Redelstroff and

Sander, 2009; Wiemann et al., 2022), albeit still compatible with an endothermic physiology

(Grigg et al., 2022).

Overall, we want to emphasize the need for a nuanced perspective on this trait. The

assumption that relative brain size alone (even if inferred correctly) can outperform all the

aforementioned thermophysiological predictors to infer endothermy appears at best

improbable. Its utility to gauge metabolic rate across ornithodiran groups therefore remains

highly doubtful and must be viewed in the framework of other, more robust lines of evidence.

3) Inferring life history traits

Finally, Herculano-Houzel (2023) suggests that neuron count estimates can be used to

model life history traits in Mesozoic ornithodiran taxa. This notion is based on previous

empirical work that showed an association between pallial neuron counts and selected

ontogenetic variables in extant mammals and birds (Herculano-Houzel, 2019). Applied to *T.*

rex, the respective equations predict that females reached sexual maturity at an age of 4–5

years and that the longevity of the species was 42-49 years (Herculano-Houzel, 2023).

These calculations rest on the assumption that *T. rex* had to have 2.2 – 3.3 billion pallial

neurons. As we have shown, this premise appears exceedingly unlikely. Furthermore, the

aforementioned life history predictions are contradicted by the fossil evidence: Sexual

maturity in extinct nonavian dinosaurs can be estimated histologically by the presence of

medullary bone, a tissue that forms as a calcium reservoir for egg shell production and which

is also seen in female birds (Schweitzer et al., 2005; Woodward et al., 2020). The earliest

estimate of sexual maturity in *T. rex*, as estimated by the presence of medullary bone, is 15

years (Woodward et al., 2020; Carr, 2020). If the life history of *T. rex* was similar to extant

American alligators where sexual maturity occurs in animals that attain half of adult size

(which would be in line with the available fossil data - Carr, 2020), then the earliest onset of

sexual maturity in *T. rex* happened in its 12th year of life. Based on these lines of evidence,

Herculano-Houzel's (2023) method greatly underestimates the onset of sexual maturity by 8

to 11 years. Based on the number of lines of arrested growth in its long bones, which are

thought to indicate annual cessations of growth, the chronologically oldest *T. rex* sampled so

far lived up to 33 years (Cullen et al., 2020). Although it is not unreasonable to assume that

T. rex lived longer than three decades, there is yet no histological evidence to support that

hypothesis. Given that Herculano-Houzel's (2023) longevity estimate is based on

problematic premises, it should not be considered a plausible alternative.

In fact, if applied to species other than *T. rex*, the limitations of the aforementioned approach

become even more visible. For instance, if the life history of the sauropod *Apatosaurus*, a

gigantic dinosaur with an adult weight exceeding 30 tonnes, is modeled based on our own

neuron count estimates derived from an avian regression and an endocranial fill of 42%, the

equations suggest a longevity of only 24.5 years and an onset of sexual maturity at 2.2 years

(note that assuming a non-avian sauropsid regression or smaller brain size would result in

an even more fast-paced life history prediction). These figures are obviously unfeasible. We

are aware that Herculano-Houzel (2023) assumes that sauropods such as *Apatosaurus*

were ectothermic animals so that the given equations could not be applied to them.

However, since this notion defies essentially all available evidence on the biology of

sauropods (see above), we choose to ignore it here. To conclude, relationships between life

history and neurology that were established from a selection of extant mammals and birds

cannot be used to reliably infer ontogenetic parameters across non-avian dinosaurs (and

potentially other fossil groups). We strongly discourage relying on them in palaeontological

practice.

1003

Beyond endocasts: What are the limits of inference on dinosaur cognition?

If neuron count estimates and other endocast-derived variables do not allow reliable

predictions about the cognitive abilities of non-avian dinosaurs to be made, what other

methods are available? First of all, trace fossils can provide direct evidence on how

dinosaurs exploited their environment and interacted with both hetero- and conspecifics

(e.g., Carpenter et al. 2005; Varricchio et al. 2007; Lockley et al. 2016; Brown et al. 2021).

While such fossils can provide precise and diverse insights into dinosaur behavior, obvious

limitations render perspectives gained from them extremely patchy, nonetheless.

One further way of inferring cognitive traits in dinosaurs is by comparatively studying

relevant behavioral phenomena in living crocodilians and birds, the groups that form their

extant phylogenetic bracket. While such approaches are starting to gain pace (Zeiträg et al.,

2023), we are not aware that ethological research could so far identify shared physical or

social cognitive skills in crocodilians and birds that have not also been found in turtles and

squamates (in case such comparative data is indeed available - Zeiträg et al., 2022; Font et

al., 2023). Thus, the behavioral resolution of such approaches appears limited thus far.

Cognitive traits identified exclusively in birds or crocodiles cannot simply be extrapolated to

Mesozoic dinosaurs with any degree of certainty since they might represent crown group

apomorphies. Whereas it might be appealing to hypothesize that cognitive patterns found

among modern palaeognaths are representative for their maniraptoriform forerunners

(Jensen et al., 2023; Zeiträg et al., 2023), this idea is (in most cases) not testable and should

hence not be disseminated uncritically.

Inferences on dinosaur cognition are hindered by the fact that both extant crocodilians and

birds are highly derived in their own ways: Convergently to mammals, birds have not only

evolved an enlarged forebrain and cerebellum, but also extensive connections between

these two brain regions (Gutierrez-Ibanez et al., 2018) and descending projections from the

pallium to the brainstem and/or spinal cord (Ulinski and & Margoliash, 1990; Reiner &

Medina, 2000). These circuits are likely essential for enabling various avian behaviors but

are not present in extant non avian sauropsids (Ulinski and & Margoliash 1990; Gutierrez-

Ibanez et al., 2023). It remains obscure when they evolved. Crown-group birds also possess

an apomorphic dorsal projection of the telencephalon, the eminentia sagittalis or wulst,

which appears to be absent even from endocasts of derived non-avian maniraptoriforms

such as *Archaeopteryx* and *Stenonychosaurus* and is prominently involved in visual

cognition (Walsh & Milner, 2011; Iwaniuk & Wylie, 2020). Crocodilians on the other hand

conserve a plesiomorphic brain morphology and cerebral tissue organization (Briscoe et al.,

2018; Briscoe & Ragsdale, 2018). They are unusual in being secondary ectotherms (e.g.,

Seymour et al., 2004; Legendre et al., 2016; Botha et al., 2023) and it is unclear how this

might have affected their neurology and cognition. Thus, the extant archosaurian groups

leave us in a rather suboptimal position to infer cognitive traits in non-avian dinosaurs.

Obviously, even the absence of a given cognitive trait in both crocodilians and basal extant

birds like palaeognaths does not refute its existence in Mesozoic dinosaurs, considering the

diversity and long evolutionary history of this group. In fact, a species' ecology is typically

more indicative of certain behaviors and associated cognitive phenomena than its

phylogenetic affinities. For instance, habitual tool use is an adaptation typically found in

omnivorous extractive foragers (Parker & Gibson, 1977; Parker, 2015) and is only rarely

reported in predators (Shumaker et al., 2011). This is reflected by the fact that the most

common types of tooling actions that have evolved comprise reaching, probing or pounding,

usually in order to access food (Colbourne et al., 2021). It has long been observed that tool

use appears when a species is found in an uncharacteristic niche, for which it lacks the

appropriate morphological adaptations, and thus compensates by using tools to generate a

functionally equivalent behavior (Alcock, 1972; Parker & Gibson, 1977). This is likely why a

number of birds that use tools are found on islands, yet the ability appears absent in their

close mainland relatives (Rutz et al., 2016). Simply put, in order for tool use to evolve, there

needs to be a reason for it to evolve, and there are very few ecological contexts where tool

use is a superior adaptation to its morphological equivalent (Hansell & Ruxton, 2008).

Unfortunately, this type of extremely specific contextual information is nearly absent in long

extinct species. From its iconic tooth and jaw morphology, one can confidently predict that a

hypercarnivorous species like *T. rex* would have no need for tools, but the problem remains

that few assumptions about extinct animal cognition are falsifiable.

In sum, reconstructing cognition in dinosaurs and other fossil taxa without close living

analogs is a challenging endeavor that requires integrative approaches if we are to provide

compelling inferences (de Sousa et al., 2023). Bare neuronal count estimates might be

considered a rather minor contribution to this effort and need to be aligned with data from

comparative anatomy and neurology, ecology, trace fossils, and comparative behavioral

studies on extant animals to offer a plausible picture of cognition in extinct lineages. While

communicating such findings, researchers should acknowledge the limitations of the

presented inferences to allow their audience to delineate between reasoned conclusions and

speculation. In a field such as dinosaur research - avidly followed by popular media and the

public eye - a nuanced view appears especially warranted.

1072

Conclusions

1073

The dinosaurian neuronal count and relative brain size estimates presented by Herculano-

Houzel (2023) are inaccurate due to methodological shortcomings, in particular for *T. rex*.

Accordingly, the biological inferences drawn from them are implausible. As we show here,

there is no compelling evidence that relative brain size in large-bodied theropods differed

significantly from that of extant non-avian sauropsids, and their telencephalic neuron counts

were likely not exceptional, especially for animals of their size. Furthermore, we highlight

issues associated with neuron count estimates in vertebrate paleontology and argue against

their use in reconstructing behavioral and life history variables, especially in animals such as

non-avian dinosaurs, for which disparate neuron densities might be hypothesized based on

different phylogenetic and morphological arguments.

For obvious reasons, many inferences we might make about Mesozoic dinosaur behavior

will remain limited. Nevertheless, we can justify certain predictions - to a degree - within

integrative empirical frameworks to which neuron count estimates might well be added in the

future. Before such steps can be taken, however, a substantially improved understanding of

the relationship between neuron counts and other biological variables, especially cognitive

performance, in extant animals is required.

1090

Institutional abbreviations

AMNH = American Museum of Natural History, New York City, New York, United States;

BMNH / NHMUK = Natural History Museum, London, UK; BSP = Bayerische

Staatssammlung für Paläontologie und historische Geologie, Munich, Germany; BYU =

Brigham Young University, Earth Science Museum, Provo, Utah, United States;

CAPPA/UFSM, Centro de Apoio à Pesquisa Paleontológica da Quarta Colônia /

Universidade Federal de Santa Maria, São João do Polêsine, Rio Grande do Sul, Brazil.

CM = Carnegie Museum of Natural History, Pittsburgh, Pennsylvania. CMN = Canadian

Museum of Nature, Ottawa, Ontario, Canada. DINO = Dinosaur National Monument,

Jensen, Utah, United States. FIP - Florida Institute of Paleontology, Palm Beach, Florida,

United States; FMNH = Field Museum of Natural History, Chicago, Illinois, United States;

FPDM = Fukui Prefectural Dinosaur Museum, Fukui, Japan. HMN / MB.R. = Museum für

Naturkunde, Berlin, Germany; IGM = Mongolian Institute of Geology, Ulaan Bator,

Mongolia; IRSNB / RBINS = Institut Royal des Sciences Naturelles de Belgique, Brussels,

Belgium; IVPP = Institute of Vertebrate Paleontology and Paleoanthropology, Beijing,

China; KUVP = Kansas University Natural History Museum, Lawrence, Kansas, United

States; MACN = Museo Argentino de Ciencias Naturales "Bernardino Rivadavia", Buenos

Aires, Argentina; MPC-D = Institute of Paleontology and Geology, Mongolian Academy of

Sciences, Ulaan Bator, Mongolia; MUCPv-CH = Museo de la Universidad Nacional del

Comahue, colección del Museo Ernesto Bachmann, Villa El Chocón, Argentina; MOR =

Museum of the Rockies, Bozeman, Montana, United States; NMC = Canadian Museum of

Nature, Ottawa, Canada; NCSM = North Carolina Museum of Natural Sciences, Raleigh,

North Carolina, United States; OMNH = Sam Noble Museum at the University of

Oklahoma, Norman, Oklahoma, United States; PIN = Paleontological Institute, Russian

Academy of Sciences, Moscow, Russia; PKUP: Peking University Paleontological

Collections, Beijing, China. ROM = Royal Ontario Museum, Toronto, Ontario, Canada;

RTMP/TMP = Royal Tyrrell Museum of Palaeontology, Drumheller, Alberta, Canada; SGM

= Ministere de l'Energie et des Mines, Rabat, Morocco; USNM = Smithsonian National

Museum of Natural History, Washington, D.C., United States; UUVP = University of Utah,

Salt Lake City, Utah, United States; YPM = Yale Peabody Museum, New Haven

Connecticut, United States.

1122

Acknowledgements

We want to thank David Burnham, Gregory M. Erickson, Ariana Paulina-Carabajal, and

Lawrence M. Witmer for sharing valuable information on fossil specimens and Nicolas E.

Campione for recommendations on body mass calculations. Andrew N. Iwaniuk is

acknowledged for helpful discussions on the methodology and structure of the study and

Jonathan Stone for allowing GRH to perform alligator dissections in his lab. Finally, we thank

Stig Walsh and two anonymous reviewers for their thoughtful and constructive comments on

earlier drafts of this manuscript.

1131

1132   **References**

1133   Alcock, J. (1972). The evolution of the use of tools by feeding animals. *Evolution*, *26*(3), 464-
1134   473.

1135

1136   Alonso, P. D., Milner, A. C., Ketcham, R. A., Cookson, M. J., & Rowe, T. B. (2004). The
1137   avian nature of the brain and inner ear of *Archaeopteryx*. *Nature*, *430*(7000), 666-669.

1138

1139   Balanoff, A. M., Bever, G. S., & Ikejiri, T. (2010). The braincase of *Apatosaurus* (Dinosauria:

1140   Sauropoda) based on computed tomography of a new specimen with comments on variation

1141   and evolution in sauropod neuroanatomy. *American Museum Novitates*, *2010*(3677), 1-32.

1142

1143   Balanoff, A. M., Bever, G. S., Rowe, T. B., & Norell, M. A. (2013). Evolutionary origins of the

1144   avian brain. *Nature*, *501*(7465), 93-96.

1145

1146   Balanoff, A., Ferrer, E., Saleh, L., Gignac, P. M., Gold, M. E. L., Marugán-Lobón, J., ... &

1147   Vaska, P. (2024). Quantitative functional imaging of the pigeon brain: implications for the

1148   evolution of avian powered flight. *Proceedings of the Royal Society B*, *291*(2015), 20232172.

1149

1150   Barrick, R. E., Showers, W. J., & Fischer, A. G. (1996). Comparison of thermoregulation of

1151   four ornithischian dinosaurs and a varanid lizard from the Cretaceous Two Medicine

1152   Formation: evidence from oxygen isotopes. *Palaios*, *11*, 295-305.

1153

1154   Baron, M.G., Norman, D.B., & Barrett, P.M. (2017). A new hypothesis of dinosaur

1155   relationships and early dinosaur evolution. *Nature, 543,* 501-506.

1156

1157   Barron, A. B., & Mourmourakis, F. (2023). The relationship between cognition and brain size

1158   or neuron number. *Brain, Behavior and Evolution*. https://doi.org/10.1159/000532013

1159

1160   Bennett, S. C. (1995). A statistical study of *Rhamphorhynchus* from the Solnhofen

1161   Limestone of Germany: year-classes of a single large species. *Journal of Paleontology*,

1162   *69*(3), 569-580.

1163

1164   Benton, M. J. (2021). The origin of endothermy in synapsids and archosaurs and arms races

1165   in the Triassic. *Gondwana Research*, *100*, 261-289.

1166

1167   Benton, M.J., Dhouailly, D., Jiang, B., & McNamara, M. (2019). The early origin of feathers.

1168   *Trends in Ecology and Evolution. 34*(9), 856-869.

1169

1170   Benson, R. B. J., Hunt, G., Carrano, M. T. & Campione, N. (2017). Data from: Cope's rule

1171   and the adaptive landscape of dinosaur body size evolution. *Dryad Digital Repository.*

1172   *https://doi.org/10.5061/dryad.1t3r4*

1173

1174   Benson, R.B., Campione, N. E., Carrano, M. T., Mannion, P. D., Sullivan, C., Upchurch, P. &

1175   Evans, D. C. (2014). Rates of dinosaur body mass evolution indicate 170 million years of

1176   sustained ecological innovation on the avian stem lineage. *PLoS Biology*, 12(5),
1177   p.e1001853.

1178

1179   Benson, R.B., Hunt, G., Carrano, M. T. & Campione, N. (2018). Cope's rule and the adaptive
1180   landscape of dinosaur body size evolution. *Palaeontology*, *61*(1), 13-48.

1181

1182   Bertrand, O. C., Shelley, S. L., Williamson, T. E., Wible, J. R., Chester, S. G., Flynn, J. J., ...
1183   & Brusatte, S. L. (2022). Brawn before brains in placental mammals after the end-
1184   Cretaceous extinction. *Science*, *376*(6588), 80-85.

1185

1186   Bever, G. S., Brusatte, S. L., Carr, T. D., Xu, X., Balanoff, A. M., & Norell, M. A. (2013). The
1187   braincase anatomy of the late Cretaceous dinosaur *Alioramus* (Theropoda:
1188   Tyrannosauroidea). *Bulletin of the American Museum of Natural History*, *2013*(376)**,** 1-72.

1189

1190   Botha, J., Weiss, B. M., Dollman, K., Barrett, P. M., Benson, R. B., & Choiniere, J. N. (2023).
1191   Origins of slow growth on the crocodilian stem lineage. *Current Biology*, *33*(19), 4261-4268.

1192

1193   Briscoe, S. D., & Ragsdale, C. W. (2018). Homology, neocortex, and the evolution of
1194   developmental mechanisms. *Science*, *362*(6411), 190-193.

1195

1196   Briscoe, S. D., Albertin, C. B., Rowell, J. J., & Ragsdale, C. W. (2018). Neocortical
1197   association cell types in the forebrain of birds and alligators. *Current Biology*, *28*(5), 686-696.
1198

1199

1200   Brown, C.M., Currie, P.J., & Therrien, F. (2021) Intraspecific facial bite marks in
1201   tyrannosaurids provide insight into sexual maturity and evolution of bird-like intersexual
1202   display. *Paleobiology* 48(1), 12-43.

1203

1204   Brusatte, S. L., Carr, T. D., Erickson, G. M., Bever, G. S., & Norell, M. A. (2009). A long-
1205   snouted, multihorned tyrannosaurid from the Late Cretaceous of Mongolia. *Proceedings of*
1206   *the National Academy of Sciences,* 106(41), 17261-17266.

1207

1208   Buchholtz, E. A., & Seyfarth, E.-A. (2001.) The study of "fossil brains": Tilly Edinger (1897–
1209   1967) and the beginnings of paleoneurology. *BioScience*, 51(8), 674-682.

1210

1211   Burnham, D. A. (2004). New information on *Bambiraptor feinbergi* (Theropoda:
1212   Dromaeosauridae) from the Late Cretaceous of Montana. In P. J. Currie, E. B. Koppelhus,
1213   M. A. Shugar & J. L. Wright (eds.), *Feathered dragons: studies on the transition from*
1214   *dinosaurs to birds* (pp. 67-111). Indiana University Press, Bloomington.

1215

1216   Butler, R. J., Barrett, P. M., & Gower, D. J. (2009). Postcranial skeletal pneumaticity and air-
1217   sacs in the earliest pterosaurs. *Biology Letters*, *5*(4), 557-560.

1218

1219   Button, D. J., & Zanno, L. E. (2023). Neuroanatomy of the late Cretaceous *Thescelosaurus*
1220   *neglectus* (Neornithischia: Thescelosauridae) reveals novel ecological specialisations within
1221   Dinosauria. *Scientific Reports*, *13*(1), 19224.

1222

1223   Caicoya, Á. L., Amici, F., Ensenyat, C., & Colell, M. (2019). Object permanence in *Giraffa*
1224   *camelopardalis*: First steps in giraffes' physical cognition. *Journal of Comparative*
1225   *Psychology*, *133*(2), 207.

1226

1227   Campione, N. E. (2020) MASSTIMATE: Body Mass Estimation Equations for Vertebrates.
1228   Available at: https://cran.r-project.org/web/packages/MASSTIMATE/MASSTIMATE.pdf

1229

1230   Campione, N. E., & Evans, D. C. (2020). The accuracy and precision of body mass
1231   estimation in non-avian dinosaurs. *Biological Reviews*, *95*(6), 1759-1797.

1232

1233   Campione, N. E., Barrett, P. M., & Evans, D. C. (2020). On the ancestry of feathers in
1234   Mesozoic dinosaurs. In C. Foth & O. W. M. Rauhut (eds.), *The Evolution of Feathers: From*
1235   *Their Origin to the Present* (pp. 213-243). Springer International Publishing.

1236

1237   Carpenter, K., Sanders, F., McWhinney, L.A., & Wood, L. (2005). Evidence for predator-prey
1238   relationships: examples for *Allosaurus* and *Stegosaurus*. In K. Carpenter (ed.), *The*
1239   *Carnivorous Dinosaurs* (pp. 325-350). Indiana University Press, Bloomington.

1240

1241   Carr, T. D. (2020). A high-resolution growth series of *Tyrannosaurus rex* obtained from
1242   multiple lines of evidence. *PeerJ*, *8*, e9192.

1243

1244   Cerroni, M. A., & Paulina-Carabajal, A. (2019). Novel information on the endocranial
1245   morphology of the abelisaurid theropod *Carnotaurus sastrei*. *Comptes Rendus Palevol*,
1246   *18*(8), 985-995.

1247

Chentanez, T., Huggins, S. E., & Chentanez, V. (1983). Allometric relationships for the
Siamese crocodile, *Crocodylus siamensis. Journal of The Science Society of Thailand*, *9*, 5–
26.

Chittka, L., & Niven, J. (2009). Are bigger brains better?. *Current Biology*, *19*(21), R995-
R1008.

Colbourne, J. A. D., Auersperg, A. M. I., Lambert, M. L., Huber, L., & Völter, C. J. (2021).
Extending the reach of tooling theory: a neurocognitive and phylogenetic perspective. *Topics*
in Cognitive Science, *13*(4), 548–572.

Cullen, T. M., Canale, J. I., Apesteguía, S., Smith, N. D., Hu, D., & Makovicky, P. J. (2020).
Osteohistological analyses reveal diverse strategies of theropod dinosaur body-size
evolution. *Proceedings of the Royal Society B*, *287*(1939), 20202258.

Cuvier, G. (1812). Recherches sur les ossemens fossiles de quadrupèdes, Vol. 3. Deterville,
Paris.

Dawson, R. R., Field, D. J., Hull, P. M., Zelenitsky, D. K., Therrien, F., & Affek, H. P. (2020).
Eggshell geochemistry reveals ancestral metabolic thermoregulation in Dinosauria. *Science*
Advances, *6*(7), eaax9361.

Deaner, R. O., Isler, K., Burkart, J., & Van Schaik, C. (2007). Overall brain size, and not
encephalization quotient, best predicts cognitive ability across non-human primates. *Brain,*
Behavior and Evolution, *70*(2), 115-124.

D'Emic, M. D., O'Connor, P. M., Sombathy, R. S., Cerda, I., Pascucci, T. R., Varricchio, D.,
... & Curry Rogers, K. A. (2023). Developmental strategies underlying gigantism and
miniaturization in non-avialan theropod dinosaurs. *Science*, *379*(6634), 811-814.

Diaz-Uriarte R, Garland T (1998) Effects of branch length errors on the performance of
phylogenetically independent contrasts. Systematic Biology 47: 654–672.

de Ricqlès, A., Padian, K., Horner, J.R., & Francillon-Vieillot, H. 2000. Palaeohistology of the
bones of pterosaurs (Reptilia: Archosauria): anatomy, ontogeny, and biomechanical
implications. *Zoological Journal of the Linnean Society* 129: 349-385.

1284

1285 de Sousa, A. A., Beaudet, A., Calvey, T., Bardo, A., Benoit, J., Charvet, C. J., ... & Wei, Y.
1286 (2023). From fossils to mind. *Communications Biology*, *6*(1), 636.

1287

1288 Druckenmiller, P. S., Erickson, G. M., Brinkman, D., Brown, C. M., & Eberle, J. J. (2021).
1289 Nesting at extreme polar latitudes by non-avian dinosaurs. *Current Biology*, *31*(16), 3469-
1290 3478.

1291

1292 Edinger T. (1929) *Die fossilen Gehirne.* J. Springer, Berlin.

1293

1294 Erickson, G.M., Rauhut, O.W.M., Zhou, Z., Turner, A.H., Inouye, B.D., Hu, D., and Norell,
1295 M.A. (2009). Was dinosaurian physiology inherited by birds? reconciling slow growth in
1296 *Archaeopteryx. PLoS ONE* 4(10): e7390.

1297

1298 Estienne, P., Simion, M., Hagio, H., Yamamoto, N., Jenett, A., & Yamamoto, K. (2024).
1299 Different ways of evolving tool-using brains in teleosts and amniotes. *Communications*
1300 *Biology*, *7*(1), 88.

1301

1302 Evans, D.C. 2005. New evidence on brain–endocranial cavity relationships in ornithischian
1303 dinosaurs. Acta Palaeontologica Polonica 50 (3): 617–622.

1304

1305 Evans, D. C., Ridgely, R., & Witmer, L. M. (2009). Endocranial anatomy of lambeosaurine
1306 hadrosaurids (Dinosauria: Ornithischia): a sensorineural perspective on cranial crest
1307 function. *The Anatomical Record*, 292(9), 1315-1337.

1308

1309 Eyal, G., Verhoog, M. B., Testa-Silva, G., Deitcher, Y., Lodder, J. C., Benavides-Piccione,
1310 R., ... & Segev, I. (2016). Unique membrane properties and enhanced signal processing in
1311 human neocortical neurons. *Elife*, *5*, e16553.

1312

1313 Fabbri, M., & Bhullar, B. A. S. (2022). The endocast of *Euparkeria* sheds light on the
1314 ancestral archosaur nervous system. *Palaeontology*, *65*(6), e12630.

1315

1316 Fragaszy, D. M., & Mangalam, M. (2018). Tooling. Advances in the Study of Behavior,
1317 2018(50), 177–241.

1318

1319 Franzosa, J., & Rowe, T. (2005). Cranial endocast of the Cretaceous theropod dinosaur
1320 *Acrocanthosaurus atokensis. Journal of Vertebrate Paleontology*, *25*(4), 859-864.

1321

1322   Fields, R. D., & Stevens-Graham, B. (2002). New insights into neuron-glia communication.
1323   *Science*, *298*(5593), 556-562.

1324

1325   Font, E., García-Roa, R., Pincheira-Donoso, D., & Carazo, P. (2019). Rethinking the effects
1326   of body size on the study of brain size evolution. *Brain Behavior and Evolution*, *93*(4), 182-
1327   195.

1328

1329   Font, E., Burghardt, G. M., & Leal, M. (2023). Brains, behaviour, and cognition: multiple
1330   misconceptions. In C. Warwick, P. C. Arena, & G. M. Burghardt (eds.), *Health and Welfare of*
1331   *Captive Reptiles* (pp. 211-238). Springer International Publishing, Cham.

1332

1333   Galton, P. M. (2001). Endocranial casts of the plated dinosaur *Stegosaurus* (Upper Jurassic,
1334   Western USA): a complete undistorted cast and the original specimens of Othniel Charles
1335   Marsh. In K. Carpenter (ed.), *The Armored Dinosaurs* (pp. 103–129). Indiana University
1336   Press, Bloomington.

1337

1338   Garland Jr, T., Harvey, P. H., & Ives, A. R. (1992). Procedures for the analysis of
1339   comparative data using phylogenetically independent contrasts. *Systematic Biology*, *41*(1),
1340   18-32.

1341

1342   Garland, Jr, T., & Ives, A. R. (2000). Using the past to predict the present: confidence
1343   intervals for regression equations in phylogenetic comparative methods. *The American*
1344   *Naturalist*, *155*(3), 346-364.

1345

1346   Gatesy, S. M. (1991). Hind limb scaling in birds and other theropods: implications for
1347   terrestrial locomotion. Journal of Morphology, 209(1), 83-96.

1348

1349   Giffin, E. B. (1989). Pachycephalosaur paleoneurology (Archosauria: Ornithischia). *Journal*
1350   *of Vertebrate Paleontology*, 9(1), 67-77.

1351

1352   Gignac, P.M. & Erickson, G.M. (2017). The biomechanics behind extreme osteophagy in
1353   *Tyrannosaurus rex*. *Scientific Reports* **7**(1), 2012.

1354

1355   Godfrey, R. K., Swartzlander, M., & Gronenberg, W. (2021). Allometric analysis of brain cell
1356   number in Hymenoptera suggests ant brains diverge from general trends. *Proceedings of*
1357   *the Royal Society B*, *288*(1947), 20210199.

1358

1359 Grigg, G., Nowack, J., Bicudo, J. E. P. W., Bal, N. C., Woodward, H. N., & Seymour, R. S.
1360 (2022). Whole-body endothermy: ancient, homologous and widespread among the
1361 ancestors of mammals, birds and crocodylians. *Biological Reviews*, *97*(2), 766-801.

1362

1363 Güntürkün, O. (2014). Is dolphin cognition special?. *Brain, Behavior and Evolution*, *83*(3),
1364 177-180.

1365

1366 Güntürkün, O., Ströckens, F., Scarf, D., & Colombo, M. (2017). Apes, feathered apes, and
1367 pigeons: differences and similarities. *Current Opinion in Behavioral Sciences*, *16*, 35-40.

1368

1369 Gutiérrez-Ibáñez, C., Iwaniuk, A. N., & Wylie, D. R. (2018). Parrots have evolved a primate-
1370 like telencephalic-midbrain-cerebellar circuit. *Scientific Reports*, *8*(1), 1-11

1371

1372 Gutiérrez-Ibáñez, C., Kettler, L., Pilon, M.C., Carr, C.E. & Wylie, D.R., (2023). Cerebellar
1373 inputs in the American alligator (*Alligator mississippiensis*). *Brain Behavior and Evolution*,
1374 98(1):.44-60.

1375

1376 Hackett, S. J., Kimball, R. T., Reddy, S., Bowie, R. C. K., Braun, E. L., Braun, M. J., … Yuri,
1377 T. (2008). A phylogenomic study of birds reveals their evolutionary history. *Science*,
1378 320(5884), 1763–1768.

1379

1380 Hansell, M., & Ruxton, G. (2008). Setting tool use within the context of animal construction
1381 behaviour. Trends in Ecology & Evolution, 23(2), 73–78.

1382

1383 Henderson, D. M. (2023). Growth constraints set an upper limit to theropod dinosaur body
1384 size. *The Science of Nature*, *110*(1), 4.

1385

1386 Herculano-Houzel, S. (2011). Brains matter, bodies maybe not: the case for examining
1387 neuron numbers irrespective of body size. *Annals of the New York Academy of Sciences*,
1388 *1225*(1), 191-199.

1389

1390 Herculano-Houzel, S. (2017). Numbers of neurons as biological correlates of cognitive
1391 capability. *Current Opinion in Behavioral Sciences*, 16(2017):1–7.

1392

1393 Herculano-Houzel, S. (2019). Longevity and sexual maturity vary across species with
1394 number of cortical neurons, and humans are no exception. *Journal of Comparative
1395 Neurology*, *527*(10), 1689-1705.

1396

1397 Herculano-Houzel, S. (2023). Theropod dinosaurs had primate-like numbers of telencephalic
1398 neurons. *Journal of Comparative Neurology*, *531*(9), 962-974.

1399

1400 Herculano-Houzel, S., & Kaas, J. H. (2011). Gorilla and orangutan brains conform to the
1401 primate cellular scaling rules: implications for human evolution. *Brain, Behavior and*
1402 *Evolution*, *77*(1), 33-44.

1403

Herculano-Houzel, S., Ribeiro, P., Campos, L., Valotta da Silva, A., Torres, L. B., Catania, K.
C., & Kaas, J. H. (2011). Updated neuronal scaling rules for the brains of glires
(Rodents/Lagomorphs). *Brain, Behavior and Evolution*, 78(4), 302-314.

Herculano-Houzel, S., Manger, P. R., & Kaas, J. H. (2014). Brain scaling in mammalian
evolution as a consequence of concerted and mosaic changes in numbers of neurons and
average neuronal cell size. *Frontiers in Neuroanatomy*, *8*, 77.

1410

1411 Herculano-Houzel, S., Catania, K., Manger, P. R., & Kaas, J. H. (2015). Mammalian brains
1412 are made of these: a dataset of the numbers and densities of neuronal and nonneuronal
1413 cells in the brain of glires, primates, scandentia, eulipotyphlans, afrotherians and
1414 artiodactyls, and their relationship with body mass. *Brain, behavior and evolution*, 86(3-4),
1415 145-163.

1416

1417 Holtz, T.R. (1996). Phylogenetic taxonomy of the Coelurosauria (Dinosauria: Theropoda).
1418 *Journal of Palaeontology, 70*(3), 536-538.

1419

1420 Hopson, J.A. (1979). Paleoneurology. In C. Gans, R. G. Northcutt, & P. Ulinski (eds.),
1421 *Biology of the Reptilia*, Volume 9 (pp. 39–146). Academic Press, New York.

1422

1423 Hothorn, T., Bretz, F., Westfall, P., Heiberger, R. M., Schuetzenmeister, A., & Scheibe, S.
1424 (2015). Package "multcomp". Available at:
1425 https://cran.r-project.org/web/packages/multcomp/multcomp.pdf

1426

1427 Hu, K., King, J. L., Romick, C. A., Dufeau, D. L., Witmer, L. M., Stubbs, T. L., ... & Benton,
1428 M. J. (2021). Ontogenetic endocranial shape change in alligators and ostriches and
1429 implications for the development of the non-avian dinosaur endocranium. *The Anatomical*
1430 *Record*, *304*(8), 1759-1775.

1431

1432 Hurlburt, G. R. (1996). *Relative brain size in recent and fossil amniotes: determination and*
1433 *interpretation.* PhD thesis, University of Toronto, Toronto, Canada (pp. 250).
1434
1435 Hurlburt, G.R., & Waldorf, L. (2002). Endocast volume and brain mass in a size series of
1436 alligators. *Journal of Vertebrate Paleontology*, *23*, (Supplement to 3), 69A.
1437
1438 Hurlburt, G. R., Ridgely, R. C., & Witmer, L. M. (2013) Relative size of brain and cerebrum in
1439 tyrannosaurid dinosaurs: an analysis using brain-endocast quantitative relationships in
1440 extant alligators. In J. M. Parrish, R. E. Molnar, P. J.Currie, & E. B. Koppelhus (eds.),
1441 *Tyrannosaurid Paleobiology* (pp. 134–154). Indiana University Press, Bloomington.
1442
1443 Iwaniuk, A. N., & Nelson, J. E. (2002). Can endocranial volume be used as an estimate of
1444 brain size in birds? *Canadian Journal of Zoology,* 80(1), 16-23.
1445
1446 Iwaniuk, A. N., & Wylie, D. R. (2020). Sensory systems in birds: What we have learned from
1447 studying sensory specialists. *Journal of Comparative Neurology*, *528*(17), 2902-2918.
1448
1449 Jenkins Jr, F. A., Shubin, N. H., Gatesy, S. M., & Padian, K. E. V. I. N. (2001). A diminutive
1450 pterosaur (Pterosauria: Eudimorphodontidae) from the Greenlandic Triassic. *Bulletin of the*
1451 *Museum of Comparative Zoology*, *156*(1), 151-170.
1452
1453 Jensen, T. R., Zeiträg, C., & Osvath, M. (2023). The selfish preen: absence of allopreening
1454 in Palaeognathae and its socio-cognitive implications. *Animal Cognition*, 26, 1467–1476.
1455
1456 Jerison, H. J. (1973). *Evolution of the Brain and Intelligence.* Academic Press, New York.
1457
1458 Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K., & Mooers, A. O. (2012). The global
1459 diversity of birds in space and time. *Nature*, *491*(7424), 444-448.
1460
1461 Jirak, D., & Janacek, J. (2017). Volume of the crocodilian brain and endocast during
1462 ontogeny. *PloS ONE*, *12*, 6, e0178491.
1463
1464 Kabadayi, C., Taylor, L. A., von Bayern, A. M., & Osvath, M. (2016). Ravens, New
1465 Caledonian crows and jackdaws parallel great apes in motor self-regulation despite smaller
1466 brains. *Royal Society Open Science*, *3*(4), 160104.
1467

1468 Kabadayi, C., Krasheninnikova, A., O'Neill, L., van de Weijer, J., Osvath, M., & von Bayern,
1469 A. M. (2017). Are parrots poor at motor self-regulation or is the cylinder task poor at
1470 measuring it?. *Animal Cognition*, *20*, 1137-1146.

1471

1472 Knoll, F., Buffetaut, E., & Buelow, M. (1999). A theropod braincase from the Jurassic of the
1473 Vaches Noires cliffs (Normandy, France); osteology and palaeoneurology. *Bulletin de la*
1474 *Société géologique de France*, *170*(1), 103-109.

1475

1476 Knoll, F., Lautenschlager, S., Kawabe, S., Martínez, G., Espílez, E., Mampel, L., & Alcalá, L.
1477 (2021). Palaeoneurology of the early cretaceous iguanodont *Proa valdearinnoensis* and its
1478 bearing on the parallel developments of cognitive abilities in theropod and ornithopod
1479 dinosaurs. *Journal of Comparative Neurology*, *529*(18), 3922-3945.

1480

1481 Knoll, F. & Schwarz-Wings, D. (2009) Palaeoneuroanatomy of *Brachiosaurus*. *Annales de*
1482 *Paléontologie*, *95*(3), 165–175.

1483

1484 Ksepka, D. T., Balanoff, A. M., Smith, N. A., Bever, G. S., Bhullar, B. A. S., Bourdon, E., ... &
1485 Smaers, J. B. (2020). Tempo and pattern of avian brain size evolution. *Current Biology*,
1486 *30*(11), 2026-2036.

1487

1488 Kumar, S., Stecher, G., Suleski, M., & Hedges, S. B. (2017). TimeTree: a resource for
1489 timelines, timetrees, and divergence times. *Molecular Biology and Evolution*, *34*(7), 1812-
1490 1819.

1491

1492 Kverková, K., Marhounová, L., Polonyiová, A., Kocourek, M., Zhang, Y., Olkowicz, S., ... &
1493 Němec, P. (2022). The evolution of brain neuron numbers in amniotes. *Proceedings of the*
1494 *National Academy of Sciences*, 119(11), e2121624119.

1495

1496 Larsson, H. C., Sereno, P. C., & Wilson, J. A. (2000). Forebrain enlargement among
1497 nonavian theropod dinosaurs. *Journal of Vertebrate Paleontology*, *20*(3), 615-618.

1498

1499 Legendre, L. J., & Davesne, D. (2020). The evolution of mechanisms involved in vertebrate
1500 endothermy. *Philosophical Transactions of the Royal Society B, 375*(1793), 20190136.

1501

1502 Legendre, L.J., Guénard, G., Botha-Brink, J., & Cubo, J. (2016). Palaeohistological evidence
1503 for ancestral high metabolic rate in archosaurs. *Systematic Biology* 65(6): 989-996.

1504

Lauters, P., Coudyzer, W., Vercauteren, M., & Godefroit, P. (2012). The brain of *Iguanodon*
and *Mantellisaurus*: perspectives on ornithopod evolution. In P. Godefroit (ed.), Bernissart
dinosaurs and early Cretaceous terrestrial ecosystems (pp 213–224). Indiana University
Press, Bloomington.

Lefebvre, L., Nicolakakis, N., & Boire, D. (2002). Tools and brains in birds. *Behaviour*,
139(7), 939–973.

Lockley, M.G., McCrea, R.T., Buckley, L.G., Lim, J.D., Matthews, N.A., Breithaupt, B.H.,
Houck, K.J., Gierliński, G.D., Surmik, D., Kim, K.S., Xing, L., Kong, D.Y., Cart, K., Martin, J.,
& Hadden, G. (2016). Theropod courtship: large scale physical evidence of display arenas
and avian-like scrape ceremony behaviour by Cretaceous dinosaurs. *Scientific Reports*
6:18952

Logan, C. J., Avin, S., Boogert, N., Buskell, A., Cross, F. R., Currie, A., ... & Montgomery, S.
H. (2018). Beyond brain size: Uncovering the neural correlates of behavioral and cognitive
specialization. *Comparative Cognition & Behavior Reviews*.

Maddison, W. P. & Maddison, D. R. (2023). Mesquite: a modular system for evolutionary
analysis. Available at: http://www.mesquiteproject.org

Manger, P. R. (2013). Questioning the interpretations of behavioral observations of
cetaceans: is there really support for a special intellectual status for this mammalian order?.
Neuroscience, *250*, 664-696.

Marsh, O. C. (1879). History and methods of palaeontological discovery. *American Journal*
of Science, *3*(107), 323-359.

Martins, E. P., & Housworth, E. A. (2002). Phylogeny shape and the phylogenetic
comparative method. *Systematic biology*, *51*(6), 873-880.

Medina, L. & Reiner, A. (2000). Do birds possess homologues of mammalian primary visual,
somatosensory and motor cortices?. *Trends in Neurosciences*, 23(1): 1-12.

Morhardt, A. C. (2016). *Gross anatomical brain region approximation (GABRA): assessing*
brain size, structure, and evolution in extinct archosaurs. PhD thesis, Ohio University, Ohio,
United States.

Müller, R. T., Ferreira, J. D., Pretto, F. A., Bronzati, M., & Kerber, L. (2021). The endocranial anatomy of *Buriolestes schultzi* (Dinosauria: Saurischia) and the early evolution of brain tissues in sauropodomorph dinosaurs. *Journal of Anatomy*, *238*(4), 809-827.

Nesbitt, S.J. (2011). The early evolution of archosaurs: relationships and the origin of major clades. *Bulletin of the American Museum of Natural History, 2011*(352), 1-292.

Ngwenya, A., Patzke, N., Spocter, M. A., Kruger, J. L., Dell, L. A., Chawana, R., ... & Manger, P. R. (2013). The continuously growing central nervous system of the Nile crocodile (*Crocodylus niloticus*). *The Anatomical Record*, *296*(10), 1489-1500.

Ngwenya, A., Patzke, N., Manger, P. R., & Herculano-Houzel, S. (2016). Continued growth of the central nervous system without mandatory addition of neurons in the Nile crocodile (*Crocodylus niloticus*). *Brain Behavior and Evolution*, *87*(1), 19-38.

O'Hara, M., Mioduszewska, B., Mundry, R., Yohanna, Haryoko, T., Rachmatika, R., Prawiradilaga, D. M., Huber, L., & Auersperg, A. M. I. (2021). Wild Goffin's cockatoos flexibly manufacture and use tool sets. *Current Biology*, *31*(20), 4512-4520.e6.

Olkowicz, S., Kocourek, M., Lučan, R. K., Porteš, M., Fitch, W. T., Herculano-Houzel, S., & Němec, P. (2016). Birds have primate-like numbers of neurons in the forebrain. *Proceedings of the National Academy of Sciences*, *113*(26), 7255-7260.

Osborn, H. F. (1912). Crania of *Tyrannosaurus* and *Allosaurus. Memoirs of the American Museum of Natural History, Volume. 1, New Series*, 1, 1-30.

Osmólska, H. (2004). Evidence on relation of brain to endocranial cavity in oviraptorid dinosaurs. *Acta Palaeontologica Polonica*, *49*(2), 321-324.

Padian, K., Horner, J.R., & De Ricqlès, A. 2004. Growth in small dinosaurs and pterosaurs: the evolution of the archosaurian growth strategies. *Journal of Vertebrate Paleontology* 24(3): 555-571.

Pagel, M. (1999). Inferring the historical patterns of biological evolution. *Nature*, *401*(6756), 877-884.

1578 Paradis, E., & Schliep, K. (2019). ape 5.0: an environment for modern phylogenetics and
1579 evolutionary analyses in R. *Bioinformatics*, *35*(3), 526-528.

1580

1581 Parker, S. T. (2015). Re-evaluating the extractive foraging hypothesis. *New Ideas in*
1582 *Psychology*, 37, 1–12.

1583

1584 Parker, S. T., & Gibson, K. R. (1977). Object manipulation, tool use and sensorimotor
1585 intelligence as feeding adaptations in cebus monkeys and great apes. *Journal of Human*
1586 *Evolution*, 6(7), 623–641.

1587

1588 Paulina-Carabajal, A., Bronzati, M., & Cruzado-Caballero, P. (2023) Paleoneurology of non-
1589 avian dinosaurs: an overview. In M. T. Dozo, A. Paulina-Carabajal, T. E. Macrini, & S. Walsh
1590 (eds.), *Paleoneurology of Amniotes : New Directions in the Study of Fossil Endocasts* (pp.
1591 267-332). Springer International Publishing, Cham.

1592

1593 Paulina-Carabajal, A., & Canale, J. I. (2010). Cranial endocast of the carcharodontosaurid
1594 theropod *Giganotosaurus carolinii* Coria & Salgado, 1995. *Neues Jahrbuch fur Geologie und*
1595 *Palaontologie*, *258*, 249-256.

1596

1597 Paulina-Carabajal, A., & Currie, P.J. (2012). New information on the braincase and endocast
1598 of *Sinraptor dongi* (Theropoda: Allosauroidea): Ethmoidal region, endocranial anatomy and
1599 pneumaticity. *Vertebrata PalAsiatica,* 50, 85–101

1600

1601 Paulina Carabajal, A., Carballido, J. L., & Currie, P. J. (2014). Braincase, neuroanatomy, and
1602 neck posture of *Amargasaurus cazaui* (Sauropoda, Dicraeosauridae) and its implications for
1603 understanding head posture in sauropods. *Journal of Vertebrate Paleontology*, *34*(4), 870-
1604 882.

1605

1606 Persons IV, W. S., Currie, P. J., & Erickson, G. M. (2020). An older and exceptionally large
1607 adult specimen of *Tyrannosaurus rex*. *The Anatomical Record*, 303(4), 656-672.

1608

1609 Picasso, M. B., Tambussi, C. P., & Degrange, F. J. (2011). Virtual reconstructions of the
1610 endocranial cavity of *Rhea americana* (Aves, Palaeognathae): postnatal anatomical
1611 changes. *Brain Behavior and Evolution*, *76*(3-4), 176-184.

1612

1613 Picasso, M. B. J. (2012). Postnatal ontogeny of the locomotor skeleton of a cursorial bird:
1614 greater rhea. *Journal of Zoology*, *286*(4), 303-311.

1615

1616   Pika, S., Sima, M. J., Blum, C. R., Herrmann, E., & Mundry, R. (2020). Ravens parallel great
1617   apes in physical and social cognitive skills. *Scientific Reports*, *10*(1), 1-19.

1618

1619   Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., Heisterkamp, S., and Van Willigen, B.
1620   (2017). Package "nlme". Available at: https://cran.r-project.org/web/packages/nlme/nlme.pdf.

1621

1622   Pontzer, H., Allen, V., & Hutchinson, J.R. (2009). Biomechanics of running indicates
1623   endothermy in bidepal dinosaurs. *PLoS ONE* 4(12): 10.1371.

1624

1625   Prondvai, E., Stein, K., Ősi, A., & Sander, P. M. (2012). Life history of *Rhamphorhynchus*
1626   inferred from bone histology and the diversity of pterosaurian growth strategies. *PLoS ONE*
1627   7(2): e31392.

1628

1629   R Core Team (2023). R: A language and environment for statistical computing. R
1630   Foundation for Statistical Computing, Vienna, Austria. Available at: https://www.R-
1631   project.org/.

1632

1633   Redelstroff, R. & Sander, P.M. (2009). Long and girdle bone histology of *Stegosaurus*:
1634   implications for growth and life history. *Journal of Vertebrate Paleontology* 29(4): 1087-1099.

1635

1636   Redelstroff, R., Hübner, T.R., Chinsamy, A., & Sander, P. M. (2013). Bone histology of the
1637   stegosaur *Kentrosaurus aethiopicus* (Ornithischia: Thyreophora) from the Upper Jurassic of
1638   Tanzania. *Evolutionary Biology* 296(6): 933-952.

1639

1640   Reiner, A. (2023). Could theropod dinosaurs have evolved to a human level of intelligence?.
1641   *Journal of Comparative Neurology*, 531(9), 975–1006.

1642

1643   Revell, L. J., Harmon, L. J., & Collar, D. C. (2008). Phylogenetic signal, evolutionary
1644   process, and rate. *Systematic Biology*, *57*(4), 591-601.

1645

1646   Roese-Miron, L., Jones, M. E. H., Ferreira, J. D., & Hsiou, A. S. (2023). Virtual endocasts of
1647   *Clevosaurus brasiliensis* and the tuatara: Rhynchocephalian neuroanatomy and the oldest
1648   endocranial record for Lepidosauria. *The Anatomical Record*, 1–24. https://doi.org/10.1002/
1649   ar.25212

1650

1651   Ridgway, S. H., Brownson, R. H., Van Alstyne, K. R., & Hauser, R. A. (2019). Higher neuron
1652   densities in the cerebral cortex and larger cerebellums may limit dive times of delphinids
1653   compared to deep-diving toothed whales. *PLoS One*, *14*(12), e0226206.

1654

1655   Rogers, S. W. (1998). Exploring dinosaur neuropaleobiology: computed tomography
1656   scanning and analysis of an *Allosaurus fragilis* endocast. *Neuron*, *21*(4), 673-679.

1657

1658   Rutz, C., Klump, B. C., Komarczyk, L., Leighton, R., Kramer, J., Wischnewski, S.,
1659   Sugasawa, S., Morrissey, M. B., James, R., St Clair, J. J. H., Switzer, R. A., & Masuda, B.
1660   M. (2016). Discovery of species-wide tool use in the Hawaiian crow. *Nature*, *537*(7620),
1661   403–407.

1662

1663   Saveliev, S. V., & Alifanov, V. R. (2007). A new study of the brain of the predatory dinosaur
1664   *Tarbosaurus bataar* (Theropoda, Tyrannosauridae). *Paleontological Journal*, *41*(3), 281-289.

1665

1666   Sakamoto, M. (2022). Estimating bite force in extinct dinosaurs using phylogenetically
1667   predicted physiological cross-sectional areas of jaw adductor muscles. *PeerJ,* 10: e13731.

1668

1669   Sampson, S. D. & Witmer, L. M. (2007) Craniofacial anatomy of *Majungasaurus*
1670   *crenatissimus* (Theropoda: Abelisauridae) From the Late Cretaceous Of Madagascar,
1671   *Journal of Vertebrate Paleontology*, 27:S2, 32-104.

1672

1673   Schliep, K., Paradis, E., de Oliveira Martins, L., Potts, A., White, T. W., Stachniss, C., &
1674   Kendall, M. (2019). Package 'phangorn'. Available at: *https://cran. r-project.*
1675   *org/web/packages/phangorn/phangorn. pdf*

1676

1677   Schmitt, V., Pankau, B., & Fischer, J. (2012). Old world monkeys compare to apes in the
1678   primate cognition test battery. *PloS one*, *7*(4), e32024.

1679

1680   Schweitzer, M. H., Wittmeyer, J. L., & Horner, J. R. (2005). Gender-specific reproductive
1681   tissue in ratites and *Tyrannosaurus rex. Science*, *308*(5727), 1456-1460.

1682

1683   Seymour, R. S., Bennett-Stamper, C. L., Johnston, S. D., Carrier, D. R., & Grigg, G. C.
1684   (2004). Evidence for endothermic ancestors of crocodiles at the stem of archosaur evolution.
1685   *Physiological and Biochemical Zoology*, *77*(6), 1051-1067.

1686

1687   Sereno, P.C., Tan, L., Brusatte, S.L., Kriegstein H.J., Zhao, X., & Cloward K. (2009).
1688   Tyrannosaurid skeletal design first evolved at small body size. *Science, 326*(5951), 418-422.

1689

1690   Shatkovska, O. V., & Ghazali, M. (2021). Relative skull size as one of the factors limiting
1691   skull shape variation in passerines. *Canadian Journal of Zoology*, *99*(12), 1054-1066.

1692

1693   Shumaker, R. W., Walkup, K. R., & Beck, B. B. (2011). *Animal Tool Behavior: The Use and*
1694   *Manufacture of Tools by Animals* (revised and updated edition edition). Johns Hopkins
1695   University Press, Baltimore.

1696

1697   Smaers, J. B., Rothman, R. S., Hudson, D. R., Balanoff, A. M., Beatty, B., Dechmann, D. K.,
1698   ... & Safi, K. (2021). The evolution of mammalian brain size. *Science Advances*, *7*(18),
1699   eabe2101.

1700

1701   Sol, D., Olkowicz, S., Sayol, F., Kocourek, M., Zhang, Y., Marhounová, L., ... & Němec, P.
1702   (2022). Neuron numbers link innovativeness with both absolute and relative brain size in
1703   birds. *Nature Ecology & Evolution*, *6*(9), 1381-1389.

1704

1705   Spence, R. D., Zhen, Y., White, S., Schlinger, B. A., & Day, L. B. (2009). Recovery of motor
1706   and cognitive function after cerebellar lesions in a songbird – role of estrogens. *European*
1707   *Journal of Neuroscience*, *29*(6), 1225-1234.

1708

1709   Stone, E. A. (2011). Why the phylogenetic regression appears robust to tree
1710   misspecification. *Systematic Biology*, *60*(3), 245-260.

1711

1712   Ströckens, F., Neves, K., Kirchem, S., Schwab, C., Herculano-Houzel, S., & Güntürkün, O.
1713   (2022). High associative neuron numbers could drive cognitive performance in corvid
1714   species. *Journal of Comparative Neurology*, *530*(10), 1588–1605.

1715

1716   Symonds, M. R. (2002). The effects of topological inaccuracy in evolutionary trees on the
1717   phylogenetic comparative method of independent contrasts. *Systematic Biology*, *51*(4), 541-
1718   553.

1719

1720   Tartarelli, G., & Bisconti, M. (2006). Trajectories and constraints in brain evolution in
1721   primates and cetaceans. *Human Evolution*, *21*, 275-287.

1722

1723 Tonini, J. F. R., Beard, K. H., Ferreira, R. B., Jetz, W., & Pyron, R. A. (2016). Fully-sampled
1724 phylogenies of squamates reveal evolutionary patterns in threat status. *Biological*
1725 *Conservation, 204,* 23-31.

1726

1727 Triki, Z., Aellen, M., van Schaik, C. P., & Bshary, R. (2021). Relative brain size and cognitive
1728 equivalence in fishes. *Brain, Behavior and Evolution*, *96*(3), 124-136.

1729

1730 Troscianko, J., von Bayern, A. M. P., Chappell, J., Rutz, C., & Martin, G. R. (2012). Extreme
1731 binocular vision and a straight bill facilitate tool use in New Caledonian crows. *Nature*
1732 *Communications*, *3*(1), 1110.

1733

1734 Tsuboi, M., van der Bijl, W., Kopperud, B. T., Erritzøe, J., Voje, K. L., Kotrschal, A., ... &
1735 Kolm, N. (2018). Breakdown of brain–body allometry and the encephalization of birds and
1736 mammals. *Nature Ecology & Evolution*, *2*(9), 1492-1500.

1737

1738 Ulinski, P. S. & Margoliash, D. (1990). Neurobiology of the reptile-bird transition. In E. G.
1739 Jones & A. Peters (eds.), *Comparative Structure and Evolution of Cerebral Cortex, Part I*
1740 (pp. 217-265). Springer, Boston.

1741

1742 Van Schaik, C. P., Triki, Z., Bshary, R., & Heldstab, S. A. (2021). A farewell to the
1743 encephalization quotient: a new brain size measure for comparative primate cognition. *Brain,*
1744 *Behavior and Evolution*, *96*(1), 1-12.

1745

1746 Varricchio, D.J., Martin, A.J., Katsura, Y. (2007). First trace and body fossil evidence of a
1747 burrowing, denning dinosaur. *Proceedings of the Royal Society B* 274:1361–1368.

1748

1749 Walsh, S., & Milner, A. (2011). *Halcyornis toliapicus* (Aves: lower Eocene, England)
1750 indicates advanced neuromorphology in Mesozoic Neornithes. *Journal of Systematic*
1751 *Palaeontology*, *9*(1), 173-181.

1752

1753 Wang, Y., Claessens, L.P.A.M., & Sullivan, C. (2023). Deep reptilian evolutionary roots of a
1754 major avian respiratory adaptation. *Communications Biology* 6:3.

1755

1756 Watanabe, A., Gignac, P.M., Balanoff, A.M., Green, T.L., Kley, N.J. & Norell, M.A. (2019).
1757 Are endocasts good proxies for brain size and shape in archosaurs throughout ontogeny?.
1758 *Journal of Anatomy*, *234*, 3, 291-305.

1759

1760 Wedel, M.J. (2006). Origin of postcranial skeletal pneumaticity in dinosaurs. *Integrative*
1761 *Zoology*, 1(2): 80-85.

1762

1763 Wiemann, J., Menéndez, I., Crawford, J. M., Fabbri, M., Gauthier, J. A., Hull, P. M., ... &
1764 Briggs, D. E. (2022). Fossil biomolecules reveal an avian metabolism in the ancestral
1765 dinosaur. *Nature, 606*(7914), 522-526.

1766

1767 Witmer, L. M., Chatterjee, S., Franzosa, J., & Rowe, T. (2003). Neuroanatomy of flying
1768 reptiles and implications for flight, posture and behaviour. *Nature, 425*(6961), 950-953.

1769

1770 Witmer, L. M., Ridgely, R. C., Dufeau, D. L., & Semones, M. C. (2008). Using CT to peer into
1771 the past: 3D visualization of the brain and ear regions of birds, crocodiles, and nonavian
1772 dinosaurs. In H. Endo & R.Frey (eds.), *Anatomical Imaging: Towards a New Morphology* (pp.
1773 67–87). Springer, Tokyo.

1774

1775 Witmer, L. M., & Ridgely, R. C. (2009). New insights into the brain, braincase, and ear region
1776 of tyrannosaurs (Dinosauria, Theropoda), with implications for sensory organization and
1777 behavior. *The Anatomical Record, 292*(9), 1266-1296.

1778

1779 Woodward, A., C. T. Moore, & M. F. Delaney. (1991). *Experimental Alligator Harvest: Final*
1780 *Report.* Study no. 7567. Bureau of Wildlife Research, Florida Game and Fresh Water Fish
1781 Commission, Tallahassee, 118pp.

1782

1783 Woodward, H. N., Tremaine, K., Williams, S. A., Zanno, L. E., Horner, J. R., & Myhrvold, N.
1784 (2020). Growing up *Tyrannosaurus rex*: Osteohistology refutes the pygmy "*Nanotyrannus*"
1785 and supports ontogenetic niche partitioning in juvenile *Tyrannosaurus. Science Advances*,
1786 *6*(1), eaax6250.

1787

1788 Yu, Y., Karbowski, J., Sachdev, R. N., & Feng, J. (2014). Effect of temperature and glia in
1789 brain size enlargement and origin of allometric body-brain size scaling in vertebrates. *BMC*
1790 *Evolutionary Biology, 14*(1), 1-14.

1791

1792 Zeiträg, C., Jensen, T. R., & Osvath, M. (2022). Gaze following: A socio-cognitive skill rooted
1793 in deep time. *Frontiers in Psychology*, 13:950935.

1794

Zeiträg, C., Reber, S. A., & Osvath, M. (2023). Gaze following in Archosauria — Alligators
and palaeognath birds suggest dinosaur origin of visual perspective taking. *Science*
Advances, *9*(20), eadf0405.

Zelenitsky, D. K., Therrien, F., Ridgely, R. C., McGee, A. R., & Witmer, L. M. (2011).
Evolution of olfaction in non-avian theropod dinosaurs and birds. *Proceedings of the Royal*
Society B: Biological Sciences, *278*(1725), 3625-3634.

Zhou, C. F., Gao, K. Q., Fox, R. C., & Du, X. K. (2007). Endocranial morphology of
psittacosaurs (Dinosauria: Ceratopsia) based on CT scans of new fossils from the Lower
Cretaceous, China. *Palaeoworld*, *16*(4), 285-293.

Supplementary Files

Suppl. File 1: Information on the selection and determination of brain endocast
measurements in this study, including comments on endocast data presented in Jerison
(1973), Hopson (1979) and Hurlburt (1996) and discussions on endocranial fill in non-
maniraptoriform dinosaurs.

Suppl. File 2: Brain and body mass data of extant sauropsid species, excluding the olfactory
tracts and bulbs. Brain mass data are derived from Hurlburt (1996), except for those of the
Siamese crocodile, which derive from Chentanez et al. (1983). All birds considered here
belong to non-telluravian taxa.

Suppl. File 3: Neuron count data, brain size measures and respective references for extant
sauropsid and mammalian species.

a

OcS   DSS   N.V   Pineal Loc.   Ch R

I C   Pit

b

OcS   OtC F   Art   Ch"R"   SaSR

c

SaSMe   Ch R   InHA   EthA

Ach   CCA

CbII

MO   N.II   OlTr

SaSR

SaSVe   Nn R   Pit   SaSVe

SaSVe   N.V   OlBu

5 cm

Lateral

Dorsal

XII

X

$V_{2,3}$ V_1

3.6   16.2   6.4

cm

"Cord"   "Brain"   "Bulb"