How smart was *T. rex*? Testing claims of exceptional cognition in dinosaurs and the
application of neuron count estimates in palaeontological research

Caspar, Kai R.[1,2*†], Gutiérrez-Ibáñez, Cristián[3*†], Bertrand, Ornella C.[4], Carr, Thomas[5],
Colbourne, Jennifer[6]; Erb, Arthur[7], George, Hady.[8], Holtz, Thomas R., Jr.,[9,10], Naish,
Darren[11]; Wylie, Douglas R.[3]; Hurlburt, Grant R.[12†]

*These authors contributed equally; † corresponding authors

[1]Institute of Cell Biology, Heinrich Heine University Düsseldorf, Düsseldorf, Germany;[2]Department of
Game Management and Wildlife Biology, Faculty of Forestry and Wood Sciences, Czech University of
Life Sciences, Prague, Czech Republic;[3]Department of Biological Sciences, University of Alberta,
Edmonton, Canada;[4]Institut Català de Paleontologia Miquel Crusafont, Universitat Autònoma de
Barcelona, Barcelona, Spain [5]Department of Biology, Carthage College, Kenosha, Wisconsin, USA
[6]Comparative Cognition Unit, Messerli Research Institute, University of Veterinary Medicine Vienna,
Vienna, Austria;[7] School of GeoSciences, Grant Institute, University of Edinburgh, Edinburgh,
Scotland, UK;[8]School of Earth Sciences, University of Bristol, Bristol, UK; [9]Department of Geology,
University of Maryland, College Park, Maryland, USA;[10]Department of Paleobiology, National Museum
of Natural History, Washington, DC, USA;[11] School of Biological Sciences, Faculty of Environment
and Life Sciences, University of Southampton, Southampton, United Kingdom;[12]Department of
Natural History, Royal Ontario Museum, Toronto, Ontario, Canada

Key words: endocast, palaeoneurology, brain evolution, comparative cognition, graphic
double integration

Abstract

Recent years have seen increasing scientific interest in whether neuron counts can
act as correlates of diverse biological phenomena. Lately, Herculano-Houzel (2022) argued
that fossil endocasts and comparative neurological data from extant sauropsids allow to
reconstruct telencephalic neuron counts in Mesozoic dinosaurs and pterosaurs, which might
act as proxies for behavior and life history traits in these animals. According to this analysis,
large theropods such as *Tyrannosaurus rex* were long-lived, exceptionally intelligent animals
with "macaque- or baboon-like cognition" whereas sauropods as well as many ornithischian
dinosaurs displayed an ectothermic physiology. Besides challenging established views on

Mesozoic dinosaur biology, these claims raise questions on whether neuron count estimates

could benefit research on fossil animals in general. Here, we address these findings by

revisiting Herculano-Houzel's (2022) neurological estimates and arguments. We present

revised estimates of encephalization and telencephalic neuron counts in dinosaurs, which

we derive from phylogenetically informed modeling and an amended dataset of endocranial

measurements. For large-bodied theropods in particular, we recover significantly lower

neuron counts than previously proposed. Furthermore, we review the suitability of

neurological variables such as neuron count estimates and relative brain size to predict

cognitive complexity, metabolic rate and life history traits in dinosaurs, coming to the

conclusion that they are flawed proxies of these biological phenomena. Instead of relying on

such neurological estimates when reconstructing Mesozoic dinosaur biology, we argue that

integrative approaches are needed to approach this complex subject.

Introduction

The Late Cretaceous North American theropod dinosaur *Tyrannosaurus rex* is a

superlative predator, being among the largest, heaviest, and most powerful (in terms of bite

force) terrestrial carnivores of all time (Gignac and Erickson 2017; Sakamoto 2022;

Henderson, 2023). Recently, Herculano-Houzel (2022) proposed that anthropoid primate-

level intelligence should be added to *T. rex*'s already impressive predatory resume based on

high estimated numbers of telencephalic neuron counts in large-bodied theropod taxa. This

conclusion emerged from a paradigm whereby neurological variables estimated from

endocasts can, so it is claimed, can be used to infer metabolic parameters, social behaviors,

and longevity in fossil species. Here, we test whether this approach and its remarkable

prospects withstand scrutiny.

The hypothesis of exceptional intelligence in dinosaurs such as *T. rex* challenges the

consensus of crocodile-like cognition in these animals, a position informed by comparative

anatomical data (Rogers, 1998; Witmer and Ridgely, 2009; Hurlburt et al., 2013). Moreover,

this claim bears ramifications that extend beyond specialized biological disciplines due to its

potential to create long-lasting impacts on the public's perspective on dinosaurs, evolution,

and the scientific process. Given the extreme contrast between Herculano-Houzel's (2022)

proposal and more traditional perspectives on dinosaur biology, we revisit the claim of

exceptional intelligence in these animals through an assessment of her methodology and a

reanalysis of the underlying data. By integrating perspectives from both paleontology and

neontology, we evaluate the potential benefits and limitations of neuron count estimation in

research on the behavior and physiology of fossil species. We begin with a brief review of

dinosaur paleoneurology and a discussion of how Herculano-Houzel's (2022) approach aims

to expand the field's methodological tool kit.

Dinosaur paleoneurology and the prospects of neuron count estimates for the field

Paleoneurology is a subfield of paleontology dedicated to research on the nervous

systems of extinct animals. Because soft tissues are not readily preserved in the fossil record,

paleobiologists rely on endocasts when studying the brains of extinct species (Paulina-

Carabajal et al. 2023). An endocast can be a natural (infilling), artificial (mold) or virtual

(digitally reconstructed) cast of the endocranial cavity that is formed by the bones of the

braincase.

The study of extinct species' endocasts, including those of dinosaurs, can be traced

back to the 1800s (e.g., Cuvier, 1812; Marsh, 1879). However, the field was truly defined by

Edinger (1929) who effectively introduced the concept of geological time to neurobiological

studies. Before her, anatomists made comparisons between endocasts and fresh brains, but

without considering the stratigraphic context (Buchholtz & Seyfarth, 2001). Jerison (1973)

followed up on Edinger's work by studying brain evolution in a quantitative manner and

developed the encephalization quotient (EQ) as an estimate of relative brain size, applicable

to both extant and extinct species. Later, the advent of X-ray computed tomography at the end

of the 1990s transformed the field and provided novel ways in which the neurosensory

systems of extinct species could be studied (e.g., Brochu, 2000; Witmer et al., 2008). Despite

these crucial innovations, however, paleoneurology has so far remained largely restricted to

the measurement and comparison of gross morphology, limiting our understanding of how the

brains of Mesozoic dinosaurs and other extinct animals worked.

Figure 1: Simplified phylogeny of the Sauropsida with a focus on the taxon Ornithodira and

representative color-coded brain morphologies, excluding the pituitary (not to scale). Blue: olfactory

bulb and tracts, Dark green: pallium (= cerebrum), Lime green: cerebellum, Yellow: diencephalon and

optic tectum; Violet: brain stem. Olfactory structures, pallium and subpallium comprise the

telencephalon. The gray overlay indicates extinct taxa, the brain morphologies of which are

approximated. Note that the brain morphology in tyrannosauroids is conspicuously plesiomorphic when

compared to the other ornithodirans pictured here (see e.g., Giffin, 1989). Silhouettes were taken from

PhyloPic (listed from top to bottom): *Morunasaurus* (in public domain), *Crocodylus* (in public domain),

Rhamphorhynchus (by Scott Hartman), *Olorotitan* (by ДиБгд, vectorized by T. Michael Keesey),

Tyrannosaurus (by Matt Dempsey), *Dromaeosaurus* (in public domain), *Dromaius* (by Darren Naish),

Corvus (in public domain).

Pterosaurs and dinosaurs (the latter including birds) form the clade Ornithodira (Fig.

1), the closest extant relatives of which are crocodylians (Fig. 1). Together, both lineages,

which separated about 250 million years ago, comprise the clade Archosauria (e.g., Legendre

et al., 2016). Next to birds, crocodylians therefore represent a critical reference point in

reconstructing the nervous systems of extinct ornithodirans.

Interestingly, highly disparate patterns of endocranial tissue organization are realized

in these two extant clades. One fundamental difference relates to the portion of the

endocranial cavity which is occupied by the brain rather than by the associated meningeal

tissues (including the dura mater and arachnoid mater) and cerebrospinal fluid (Fig. 2). In

crocodylians, nervous tissue only fills a fraction of the braincase (Hopson, 1979; Jirak &

Janacek, 2017; Watanabe et al. 2019). Longitudinal venous sinuses course along the dorsal

and ventral aspect of the brain, obscuring its true shape in casts of the braincase. Furthermore,

the size of the brain relative to both the endocranial volume and total body size, decreases

during crocodylian ontogeny, even over the course of adulthood (Hurlburt et al., 2013; but note

that the overall brain volume increases with body size, even in adults - Ngwenya et al., 2013).

Endocast morphology indicates that the endocranial cavity in most non-avian

dinosaurs was organized in crocodylian-like fashion and comparative studies suggest that this

configuration was indeed ancestral for the clade Archosauria (Witmer et al., 2008; Hurlburt et

al., 2013; Fabbri & Bhullar, 2022). For tyrannosaurids specifically, which are among the best-

studied dinosaurs when it comes to palaeoneurology, endocasts representing different

ontogenetic stages suggest that brain size (relative to endocranial volume) decreased with

age (Brusatte et al., 2009; Bever et al., 2013), as is the case in modern crocodylians.

Therefore, crocodylian and most dinosaurian endocasts do not faithfully capture the volume

and anatomy of the brain, particularly its posterior regions such as the cerebellum (Watanabe

et al., 2019). This contrasts with the situation in birds for which endocasts represent excellent

brain size proxies (e.g., Iwaniuk and Nelson, 2002), which is also the case for most mammals

(Bertrand et al., 2022).

The avian pattern probably evolved at the root of the theropod dinosaur clade

Maniraptoriformes, which includes ornithomimosaurs ('ostrich-mimic' dinosaurs) and

maniraptorans (the bird-like oviraptorosaurs, dromaeosaurids and kin, and birds themselves)

(Osmólska, 2004; Balanoff et al., 2013). Maniraptoriform brains have enlarged cerebral and

cerebellar regions that almost fully occupy the endocranial cavity. With the exception of at

least some pachycephalosaurs (Giffin, 1989), there is no evidence that the brains of other

dinosaurs similarly contacted the endocranial surface. Blood vessel impressions indicate that

the lateral surface of the cerebrum closely approached the endocranial surface in ornithopods

(Evans et al., 2009) but this does not necessarily indicate a bird-like endocranial fill, as the

same feature is present in crocodylians (G. R. Hurlburt, pers. obs.). Pterosaurs are similar to

maniraptoriforms in also possessing brains that fit tightly into the endocranial cavity (Witmer

et al., 2003).

Aside from general endocranial tissue organization, the neuroarchitecture of the

forebrain in birds and crocodylians differs notably from one another (Ulinski & Margoliash,

1990; Briscoe et al., 2018; Briscoe & Ragsdale, 2018). Comparisons with other sauropsids

demonstrate that again the crocodylian condition is the more plesiomorphic one (Briscoe &

Ragsdale, 2018). To which extent non-avian dinosaurs and pterosaurs resembled the two

extant archosaur groups in regards to neuroarchitecture cannot be reliably reconstructed,

since the latter lacks osteological correlates.

The inferred brain anatomy of various dinosaur groups has been discussed elsewhere

(Paulina-Carabajal et al. 2023) and reviewing it here is beyond the scope of this article. We

aim instead to focus on what endocast-based methods potentially reveal about the behavior

and cognition of extinct species. While considering the aforementioned limitations, endocasts

from fossil ornithodirans allow us to reasonably estimate basic neuroanatomical measures

such as EQ, as well as to deduce specific sensory specializations (e.g., Witmer et al., 2003;

Witmer & Ridgely, 2009; Zelenitsky et al., 2011). Nonetheless, it is generally assumed that the

predictive power of these data in elucidating the cognitive capacities of a fossil species is low

(Paulina-Carabajal et al., 2023). Researchers have long sought to identify robust

morphological correlates of cognition but have found traditional proxies such as EQ and

absolute brain size to be limited and problematic regarding their conceptual justifications (Van

Schaik et al., 2021). Current debates focus on whether refined neuroanatomical measures

such as "cognitive brain size" (Van Schaik et al., 2021) and brain region-specific neuron counts

(Herculano-Houzel, 2011; Kabadayi et al., 2016; Logan et al., 2018; Sol et al., 2022) might be

able to overcome these issues. The quantification of these, however, seemed out of reach for

vertebrate palaeontology.

With this in mind, the approach proposed by Herculano-Houzel (2022) is of great

potential significance: It entails that endocasts of extinct taxa can be used to model neuron

counts if neurological data from related extant species can be taken into account. If valid, this

technique would potentially allow researchers to elucidate aspects of brain physiology that

cannot be inferred from endocast morphology alone. Herculano-Houzel & Kaas (2011) and

Herculano-Houzel et al. (2011) pioneered this approach for fossil hominins and extinct giant

rodents, but Herculano-Houzel (2022) was first in applying this methodology to fossil sauropsid

groups separated from their extant relatives by hundreds of millions of years of evolution,

namely pterosaurs and Mesozoic dinosaurs. Indeed, Herculano-Houzel (2011, 2017, 2022)

has argued emphatically that neuron counts represent reliable estimates for cognitive abilities

in extant vertebrates, markedly outperforming other measures such as relative or absolute

brain size. If we accept this premise, accurate modeling of neuron counts in dinosaurs based

on endocast volumes and comparative neurological data might appear as a promising new

method to elucidate the behavior and cognitive capacities of these animals.

Figure 2: The endocast and endocranial tissue organization in the American alligator (*Alligator mississippiensis*), illustrating the plesiomorphic condition within the clade Archosauria. Scale bar = 2 cm in all cases. **a:** Endocast of a wild *A. mississippiensis* (Fla.F&G.HarvestTag 937095), Dorsal cranial length (DCL), 342.90 mm, right lateral view. Reduced in size to match proportions of brain in **b** & **c**. **b**: Dura mater around the brain of *A. mississippiensis*, specimen CITES FLM 12- 29409, DCL, 380 mm, left lateral view (reversed). **c**: Brain within arachnoid of FLM 12-29409. Brown-red material is dried blood filling the subarachnoid space (**SaS**), right lateral view. **Abbreviations: Ach**, arachnoid mater (covering the cerebellum); **Art**, artery on external wall of dura mater over the lateral pole of the

cerebrum; **CbII**, cerebellum; **CCA**, caudal cerebral artery; **Ch L** or **R**, left or right cerebral hemisphere;

DSS, dorsal sagittal sinus; **EthA**, common ethmoid artery; **I C**, internal carotid artery; **InHA**,

interhemispheric artery; **MO**, medulla oblongata; **N.II**, optic nerve; **N.V,** (cast of) trigeminal nerve; **Nn R**,

roots of nerves IX-XI; **OC**, occipital condyle; **OcS**, occipital sinus; **OlBu** & **Tr**, olfactory bulb & tract; **OtC**

F, fossa of otic capsule; **Pineal Loc**, pineal gland location; **Pit**, pituitary gland; **SaSMe**, mesencephalic

subarachnoid space; **SaSR**, rostral SaS; **SaSVe**, ventral SaS; **SN.I**, first spinal nerve. The rostral end

of the cerebrum is below the arrow for **SaSR** in **B**. Both specimens are housed in the private collection

of G. R. Hurlburt.

The methodology and rationale of Herculano-Houzel (2022)

Herculano-Houzel (2022) reconstructed relative brain size and neuron counts for 29

dinosaur and pterosaur species based on comparative data from extant non-avian

sauropsids ("reptiles" in the colloquial sense) and birds (Olkowicz et al., 2016; Kverková et

al., 2022). Although we want to avoid lengthy discussions about taxonomy, it is worth noting

that some of these are no longer considered valid taxonomic entities (see below; an updated

nomenclature for relevant dinosaurs is included in Tab. 1). For instance, *Rhamphorhynchus*

muensteri and *R. gemmingi* have long been synonymized (Bennett, 1995). Surprisingly,

Herculano-Houzel (2022) inferred an ectothermic metabolism for one, and endothermy for

the other based on assumptions about relative brain size.

Neuron count estimates for fossil taxa primarily concerned the telencephalon, a

major brain region which is critically involved in cognitive and motor functions as well as the

processing of sensory information. It encompasses the pallium (which equates to the

cerebral cortex in humans and other mammals) and subpallium as well as the olfactory bulbs

and tracts (Fig. 1). To understand the rationale behind the approach of reconstructing

neuron counts in fossil species, two important matters must be pointed out: firstly, among

jawed vertebrates, body size and brain size are highly correlated, exhibiting a constant

allometric relationship overall (Tsuboi et al., 2018). It should be noted however, that scaling

relationships can vary to some extent between major taxa as well as between early- and

late-diverging members of a clade (Ksepka et al., 2020; Bertrand et al., 2022). Secondly,

neuronal densities (the number of neurons in a given volume of nervous tissue) can differ

profoundly between different vertebrate taxa. For instance, most bird groups exhibit about

2.5 times more telencephalic neurons than a lizard or turtle with a telencephalon of equal

mass, and in members of the bird clade Telluraves (birds of prey, rollers, parrots, songbirds

and kin) the difference reaches 3.2 fold (Kverková et al., 2022). This means that brain size

alone is not a reliable predictor of neuron counts across distantly related clades (compare

Herculano-Houzel et al., 2014; Olkowicz et al., 2016), which makes their inference in fossil

groups inherently difficult.

To decide which neuronal density patterns apply to specific groups of dinosaurs and

pterosaurs, Herculano-Houzel (2022) relied on brain x body mass regressions. The brain

and body mass data used were taken from various literature sources and, as we attempt to

show here, both are problematic. In the resulting regression plot, she identified theropods

clustering distinctly from most other species. On average, they appeared to exhibit larger

brains for a given body size than the remaining dinosaur or pterosaur taxa. When comparing

the regression curves for extinct groups with those of living birds on the one hand and

squamates and turtles on the other, Herculano-Houzel (2022) noted that the theropod

regression fits with the avian one, while the remaining groups align more with the non-avian

sauropsid regression line.

Based on these analyses, Herculano-Houzel (2022) made two critical assumptions:

First, since only data from theropods align with those of endothermic extant sauropsids,

namely birds, the other groups (aside from specific pterosaurs and ornithischians that cluster

with theropods) should be considered ectothermic. Second, telencephalic neuron densities

in theropod brain tissue should have been comparable to those found in certain extant bird

taxa (precisely, to those found in a polyphyletic assemblage denoted as "pre-K-Pg birds" that

includes Palaeognathae, Galloanserae and Columbiformes and which is considered to form

a neurological grade: Kverková et al., 2022), whereas those of the other groups should

resemble densities encountered in squamates and turtles. No further justification for these

notions is provided.

Applying the avian scaling regime, Herculano-Houzel (2022) estimated remarkably

high telencephalic neuron counts in large-bodied theropods such as *Acrocanthosaurus*

atokensis (2.1 billion) and *T. rex* (3.3 billion) which would exceed those of any extant bird

and be comparable to large-bodied Old World monkeys such as baboons (Olkowicz et al.,

2016). Based on this apparent similarity to anthropoid primates, she further speculated that

these giant theropods would have crafted and used tools and exhibited cultural behaviors

(Herculano-Houzel, 2022).

We regard the methodology of Herculano-Houzel (2022) as problematic and disagree

with her physiological and behavioral interpretations. Before we attempt to replicate her

findings with a more refined analytical approach, we want to enumerate the most important

flaws of the article and how they affect the inferences made.

Issues with Herculano-Houzel's method and rationale

A key problem for paleoneurology lies in the fact that an endocast does not

necessarily reflect the morphology of an animal's brain. As discussed in previous sections,

the endocasts of most non-avian dinosaurs differ markedly in size and shape from the actual

brain, as is the case in crocodylians (Fig. 1). Unfortunately, not all studies that Herculano-

Houzel (2022) derived her raw data from considered this issue (see below). In addition to

that, the percentage of endocranial space filled by the brain, as well as its proportions may

be further influenced by ontogeny (Bever et al., 2013; Hurlburt et al., 2013; Ngwenya et al.,

2013; Jirak & Janacek, 2017; Hu et al., 2021). The latter point is relevant because

Herculano-Houzel (2022) includes several specimens which correspond to juveniles rather

than adults, and thus might have introduced biases to the dataset (see below). Interestingly,

at least in crocodylians, neuronal densities in the brain are also affected by ontogenetic

stage (Ngwenya et al., 2016).

To arrive at the estimated telencephalic neuron count of > 3 billion for *T. rex*,

Herculano-Houzel (2022) assumed a brain mass of 343 g. However, this presupposes that

the endocast accurately reflects the volume of the brain in this species. While it has indeed

been claimed that theropods such as *T. rex* had brains filling the entire endocranial cavity

(Balanoff et al., 2013), this hypothesis is, as previously discussed, contradicted by multiple

lines of evidence. More conservative inferences suggest a brain mass of approximately 200

g (Hurlburt, 1996; Hurlburt, et al, 2013; Morhardt, 2016) or possibly even lower (this study;

Tab. 1) for *T. rex*. Herculano-Houzel (2022) acknowledges these lower estimates but chose

to rely on the inflated values for large theropod brain masses in accompanying figures and

the article's Discussion section.

Besides that, the literature-derived brain mass estimates that were used for the

analyses did in some cases include the olfactory tracts and bulbs (Franzosa & Rowe, 2005,

Balanoff et al., 2013) while they omitted them in others (Hurlburt 1996; Hurlburt et al., 2013).

This incongruence creates critical biases, affecting both the inference of telencephalic

neuron counts and relative brain size estimates. The latter are additionally skewed by the

fact that body masses were not determined via a uniform methodology. Instead, they were

compiled from various sources that applied an array of approaches, varying greatly in

precision. When compared to body mass estimates derived from stylopodial circumference,

a well-established and robust method (Campione & Evans, 2020), some striking differences

become apparent (Tab. 1; Herculano-Houzel et al., 2022).

Another flaw of Herculano-Houzel's (2022) approach is the lack of consideration for

brain morphology to inform its analyses. To estimate telencephalic neuron numbers in fossil

species, the mass of the telencephalon needs to be approximated first. For theropods,

Herculano-Houzel (2022:6) extrapolates this variable from extant bird data while stating that

"within a clade, brain mass has strongly predictive power to arrive at estimates of numbers of

telencephalic neurons in a brain of known mass, once the neuronal scaling rules that

presumably apply are known." However, this statement can only hold true if the general

proportions of the telencephalon compared to the remaining brain are roughly constant in the

group of concern, which is a precondition that Herculano-Houzel (2022) did not test for in the

fossil sample. Indeed, avian brains only poorly reflect the brain morphologies found in the

majority of non-avian dinosaurs (reviewed by Paulina-Carabajal et al. 2023) and their

general proportions are only comparable to those found among maniraptoriform theropods

(Balanoff et al., 2013; Fig. 1). An important difference concerns the pallium, which crucially

contributes to higher cognitive functions, and greatly increased in size within the

maniraptoriform radiation (Balanoff et al., 2013). The same is true for the cerebellum, a part

of the brain which is not encompassed by the telencephalon but is also involved in various

aspects of cognition in amniotes (Spence et al., 2009). Thus, the telencephalic mass and

proportions of non-maniraptoriform theropods, such as *T. rex*, cannot be adequately

modeled based on extant birds. Similar limitations need to be considered when

reconstructing traits of the pterosaur or sauropodomorph telencephalon based on extant

non-avian sauropsids and they also apply to our own empirical approach.

Herculano-Houzel (2022) hypothesized that the inferred incongruence in relative

brain size between theropods and other dinosaurs reflects differences in thermobiology,

which would justify applying avian neuronal scaling schemes to the former.

Sauropodomorphs as well as selected ornithischians and pterosaurs are instead suggested

to be ectothermic due to their relatively smaller brains. Both of these assumptions are

problematic: First, multiple lines of evidence suggest that ornithodiran endothermy evolved

long before theropods emerged and was likely already present in the common ancestor of

dinosaurs and pterosaurs (e.g., Grigg et al., 2022). We will revisit this evidence and how it

challenges the brain size-derived hypothesis in more detail in the Discussion section of this

paper. Herculano-Houzel (2022) only referenced a single article on dinosaur thermobiology,

that of Wiemann et al. (2022), to defend her standpoint on the matter. The study in question

applies a promising but novel technique to infer endothermy based on lipoxidation end

products in fossil bone that still has to prove itself. Whereas it indeed suggests lowered

metabolic rates in some ornithischians, it also infers an endothermic metabolism for

pterosaurs and sauropodomorphs (Wiemann et al., 2022). Thus, its findings do not align with

Herculano-Houzel's (2022) assumptions.

Second, a comparison of birds and mammals demonstrates that there is no uniform

relationship between neuron density and relative brain size or elevated metabolic rates. The

neuronal density of non-primate mammals is only moderately increased compared to

ectothermic sauropsids (Kverková et al., 2022; note that the difference is still statistically

significant) and markedly lower than in birds, although mammals are large-brained. It is

therefore not straight-forward to assume avian neuron densities in theropods simply because

they exhibited endothermy or increased relative brain size. The other way around, the

extensive evidence for endothermy in other dinosaurs and pterosaurs does not negate that

these groups could have exhibited neuron densities similar to those found in extant

ectothermic sauropsids.

Other issues relate to the statistics employed by Herculano-Houzel (2022). Despite

dealing with a large multi-species dataset, the analyses do not take phylogeny into account,

which is argued to be a measure to prevent mathematical artifacts. However, phylogenetic

relationships between analyzed taxa need to be statistically addressed because shared

ancestry can result in non-independence of species-specific data points (Revell et al., 2008).

Such non-independence is known as the phylogenetic signal, and it has been prominently

recovered for relative brain size in extant sauropsids (Font et al., 2019). Hence,

phylogenetically-informed modeling is a necessity to adequately work with such datasets

(Font et al., 2019).

In light of these substantial shortcomings, we attempt to replicate the findings of

Herculano-Houzel (2022) with phylogenetically informed models of telencephalic neuron

counts in fossil dinosaurs that acknowledge the issues lined out above. We do not include

pterosaurs into this analysis due to difficulties with estimating their body mass (especially for

taxa with incomplete postcrania such as *Scaphognathus*) and because of the unclear

taxonomic and ontogenetic status of some of the few available endocasts.

Empirical part: modeling neurological variables in dinosaurs

Endocast sample composition, with notes on endocranial volumes provided by Hurlburt (1996)

Empirical part: modeling neurological variables in dinosaurs

Endocast sample composition, with notes on endocranial volumes provided by

Hurlburt (1996)

We estimated the mass of the brain (MBr, g; excluding the olfactory tracts and bulbs)

as well as its size relative to body mass (MBd, g) in 28 Mesozoic dinosaur taxa for which

data on endocranial volume (EV, ml) have been published (Tab. 1; Suppl. File 1). Note that

this study does not aim at providing a comprehensive dataset on dinosaur brain sizes. Given

the questions we want to address, its focus is on large-bodied theropods and an evaluation

of previously published EV datasets. We included one endocast per species, except for *T.*

rex, for which three adult endocasts (AMNH 5029, AMNH 5117, FMNH PR 2081) were

considered. We only considered species for which we could calculate body mass based on

stylopodial circumference (see below) to reliably infer encephalization. Due to this, our

analysis does not cover all dinosaur species for which complete endocasts are available, nor

all species that Herculano-Houzel (2022) included in her analyses (namely *Conchoraptor*

gracilis, *Tsaagan mangas*, *Zanabazar junior*, unnamed troodontid IGM 100/1126). Juvenile

specimens considered by that study (*Alioramus altai* IGM 100/1844, *Gorgosaurus libratus*

ROM 1247, *Tyrannosaurus rex* CMN 7541 = *"Nanotyrannus lancensis")* were omitted in this

analysis to eliminate the confounding variable of ontogeny.

The only juvenile we include is *Bambiraptor feinbergi* KUVP 129737, which is one of

the few maniraptoriform theropods that we can take into account. For this species, an adult

femur (FIP 002) is available, allowing us to estimate body mass in fully grown individuals.

Our method of body mass inference suggests that KUVP 129737 attained about 45% of

adult body mass when it died. Data from similar-sized extant rheas (*Rhea americana*),

palaeognath birds which are close neuroanatomical analogs to highly derived theropods

such as *Bambiraptor* (Balanoff et al., 2013), suggest that adult brain mass is already

approached at that point of somatic development (Picasso et al., 2011; Picasso, 2012;). We

therefore combine the juvenile endocranial measurement of *Bambiraptor* with adult body

mass estimates.

Just as Herculano-Houzel (2022) did, we derive a significant portion of our EV values

from Hurlburt (1996). However, many EV figures communicated in this reference must be

considered problematic and were carefully bypassed here. We give detailed reasons for

discarding or modifying data from Hurlburt (1996) and the references provided therein

(Jerison, 1973; Hopson, 1979) in Suppl. File 1 Part A. In cases where EV values appeared

doubtful but appropriate illustrations or photographs of specimens were available, we

recalculated EV using manual graphic double integration (GDI; see below for methodology).

This was done for 4 species (*Allosaurus fragilis, Euoplocephalus tutus, Kentrosaurus*

aethiopicus & Ornithomimus edmontonicus; see Suppl. File 1 for details on specimens).

Brain mass estimates

We estimate the brain mass (MBr, g) of fossil dinosaurs from endocast volume (EV,

mL). Because the specific gravity (density) of living amniote brain tissue approximates one

(1.036 g/mL - Iwaniuk & Nelson, 2002), we use brain volume and mass interchangeably

(Hurlburt et al., 2013; Herculano-Houzel, 2022). For maniraptoriform species, we assumed a

brain:endocast ratio of 100%, which is consistent with empirical data on the relationship

between MBr and EV in modern birds, which suggest contributions of meningeal tissue to

endocast volume to be negligible (Iwaniuk & Nelson, 2002). In contrast to that, we assumed

MBr:EV ratios of 31% and 42% to approximate brain mass in both non-maniraptoriform

theropods and non-theropod dinosaurs. For these groups, many previous studies have

assumed a 50% ratio (see Morhardt, 2016) while some even assumed 100% (Balanoff et al.,

2013) or advocated for intermediate values (e.g., Evans et al., 2008; Knoll & Schwarz-

Wings, 2009; Knoll et al., 2021). The widely adopted 50% ratio was originally proposed by

Jerison (1973) and based on measurements from a likely subadult green iguana (*Iguana*

iguana) and a mere visual estimate of endocranial filling in the tuatara (*Sphenodon*

punctatus; the endocranial fill in this species is now known to be 30% in adults - Roese-

Miron et al., 2023). We abandon the problematic 50% estimate and replace it here by the

two aforementioned ratios that are based on the morphology of extant crocodylians, the

closest extant analogs to most non-avian dinosaurs in regards to endocranial tissue

organization.

Excluding one anomalous value, the lowest MBr:EV ratio among the three longest

American alligators (*Alligator mississippiensis*) studied by Hurlburt et al (2013) was found to

be 31% (n_{total} = 12, note that this figure excludes the olfactory bulb and tract portions of the

endocranial cavity). This is consistent with observations on the largest Nile crocodile

(*Crocodylus niloticus;* a 16-year-old female) studied by Jirak & Janacek (2017) when

excluding the olfactory tracts and bulb portion of the endocast. The 42% ratio is derived

solely from American alligators. An endocranial fill of 42% was found in an adult female with

a total length of 2.870 m, which roughly approximates both (a) the maximum length for a

female American alligator and (b) the midpoint length within the size spectrum of sexually

mature alligators (Hurlburt et al, 2013; Hurlburt & Waldorf, 2002; Woodward et al, 1991).

The majority of EV data for Mesozoic dinosaurs were taken and modified from the

literature (detailed out in Tab. 1). In many cases, original sources communicated

measurements that correspond to total EV. This is the volume of the entire endocast, often

including the region of the olfactory tract and bulbs as well as portions of the cranial spinal

cord among other structures. For our analysis, we exclusively relied on the so-called "brain"

endocast volume instead (BrEV; Fig. 3), which was popularized by Jerison (1973) and has

been commonly used since then (e.g., Hurlburt, 1996; Larsson et al., 2000; Paulina-

Carabajal & Canale, 2010; Hurlburt et al., 2013). It excludes the spinal cord portion of the

endocast caudal to cranial nerve XII, the volume of nerve trunks from infillings of respective

foramina and blood vessel casts, the labyrinth of the inner ear, the infundibulum, the pituitary

gland, and especially the volume of the olfactory lobes and tracts (Fig. 4), which are often

only poorly preserved in fossil endocasts. In some dinosaurs, there is an obvious constriction

and/or a change in surface morphology at the junction of the cerebrum and olfactory tract

(e.g., *Euoplocephalus tutus* AMNH 5337 - Hopson, 1979; *Stegosaurus ungulatus* CM 106 -

Galton, 2001; *Diplodocus longus* CM 11161- Witmer et al., 2008). If present, this was used

as a landmark to delineate these brain regions from one another. In less obvious cases, the

junction between the cerebrum and olfactory tract portion was assumed to be where the

ventral curve of the rostral cerebrum flattens out to approach a horizontal orientation. When

selecting the boundary, we erred towards a more rostral location, to assign as much of the

endocast as part of the BrEV as seemed reasonable. In American alligators, the rostral

termination of the cerebrum within the rostral subarachnoid space is clearly visible (Fig. 2C;

SaSR) and consistent with the change in curvature referred to above.

We used manual GDI to extract BrEV from total EV (relevant details for each

specimen are provided in Supplementary File 1 Part B). The method involves drawing an

outline around two scaled orthogonal two-dimensional views of an endocast, and adding

equally spaced lines perpendicular to the endocast midline (Fig. 3). The mean length (cm) of

these lines in each view (i.e., dorsal, lateral) provides diameters D1 and D2. The volume

(mL) of the desired region is calculated using these two diameters and the length (L, cm) in

the formula for the volume (mL or cm^3) of a cylinder where (all lengths in cm):

$$Volume\ [mL]\ =\ (\pi)(0.25)(D1)(D2)(L)$$

GDI has been demonstrated to produce reasonable estimates of endocast volumes

(Fig. 3). For instance, Jerison (1973) used GDI to calculate a total endocast volume of 536

ml for a *T. rex* specimen (AMNH 5029), which was 101.13% of the 530 ml volume

determined for it by means of water displacement (Osborn, 1912; Fig. 3). For the same

specimen, Jerison (1973) calculated a 404 ml volume for the "brain region" of the endocast

(extending from cranial nerve XII to the rostral cerebral limit), which was 106.04 % of the 381

ml obtained by CT volumetry (Hurlburt et al., 2013).

Body mass estimates

We calculated the body mass (MBd, g) of the selected dinosaur taxa (and its mean

absolute percent prediction error - PPE or %PE - Campione & Evans, 2020) in a

standardized manner based on the minimum femoral circumference (as well as humeral

circumference in case of quadrupedal taxa) by aid of the *QE()* and *cQE()* functions from the

MASSTIMATE package (Campione, 2020) in R (R Core Team, 2022). Data on relevant long

bone dimensions were primarily obtained from Benson et al. (2017). Corresponding

specimens as well as additional sources and information on stylopodium circumference

measurements are listed in Tab. 1. To the best of our knowledge, all data correspond to

adult specimens.

Figure 3. Exemplary graphic double integration (GDI) of the endocast of *Tyrannosaurus rex* AMNH

5029. Equally spaced lines are drawn across the right lateral and dorsal views respectively. Mean

lengths of the lines drawn across the "brain" portion (BrEV) were 4.8 cm and 6.6 cm for dorsal and

lateral views respectively. BrEV = π x 0.25 x 4.8 x 6.6 x 16.2 = 404 mL (the volume of the entire

endocast was 536 mL). Abbreviations: "Bulb": Olfactory tract and bulbs; "Cord": spinal cord; V:

Trigeminal nerve with its three branches (V1, V2,& V3); X: vagus nerve; XII: hypoglossal nerve.

(Adapted from Fig. 2.7 in Jerison,1973, p. 51).

Table 1: Estimates of brain (MBr, g; derived from endocranial volume, EV) and body mass (MBd, kg) in Mesozoic dinosaurs. For maniraptoriform theropods,

we assumed that brain endocast volume equals brain volume. For non-maniraptoriform theropods and non-theropod dinosaurs, we assumed a

brain:endocast ratio of 31 - 42 %. Body mass was calculated based on stylopodial circumference (femoral circumference (FC) for bipedal and femoral as well

as humeral (HC) circumference for quadrupedal species). **Footnotes: [1]** *Bambiraptor feinbergi*. MBd from femur circumference, measured on a cast of the

right femur of an adult specimen now in the collection of the Vertebrate Paleontology Division, ROM (FIP 002). Original elements of this specimen are now in

the collection of the AMNH. **[2]** *Archaeopteryx lithographica*. MBd from FC (14.93 mm) estimated from femur length (60.5mm) of BMNH 37001 (Gatesy,

1991). FC was calculated using the equation from Benson et al. (2017): log10(Femur circumference estimate from femur length) = 1.132 * log10(Femur

length) - 0.8429. **[3]** *Carcharodontosaurus saharicus.* MBd from estimated femur circumference (FC = 417.52 mm) from Benson et al. (2017) for BSP 1922

X46. The specimen has been destroyed and is no longer accessible. **[4]** *Sinraptor dongi*. MBd based on FC (283 mm) from Campione & Evans (2020),

measurement taken from TMP 93.115.1 (cast of IVPP 10600). Benson et al. (2017) consider IVPP 10600 a subadult based on reported incomplete fusion of

cervical vertebrae; However, Paulina-Carabajal and Currie (2012) noted that the degree of cranial suture fusion indicates that the specimen is an adult or at

least a large subadult. We consider it an adult here. **[5]** *Tyrannosaurus rex* specimens. MBd based on FC values listed by Persons et al. (2020). For FMNH

PR 2081 ("Sue"), but not the other individuals considered here, both FC and EV are available. We associated the EV (313.636 mL) of AMNH 5117 with the

MBd (5515 kg) of BHI 3033 ("Stan"), as both specimens have been considered proxies for each other (G. M. Erickson, pers comm. To G.R. Hurlburt, 2005).

The EV (381.8 mL) of AMNH 5029 is here linked to the MBd (6430 kg) of CM 9380 (holotype specimen) because it fell between MBd's associated with EVs of

FMNH PR 3081 and AMNH 5117. Besides our brain:endocast ratios, we also apply the 57% ratio proposed by Morhardt (2016) to this species.

Species	Group	Body mass (MBd) (kg)	Lower 25% PPE (kg)	Upper 25% PPE (kg)	FC (mm)	HC (mm)	MBd Spec	MBd Source for FC & HC	Brain Mass (MBr) (g)	MBr/EV	Br EV (mL/cm3)	EV Specmen	EV Meth	Original
Bambiraptor feinbergi[1]	Non-avian maniraptoriform	8.0632	5.9966	10.1298	47	NA	FIP 002	Cast of right femur	14	1	14	KUVP 129737	Water displacement	Burnham
Citipati osmolskae	Non-avian maniraptoriform	123.993	92.214	155.772	127	NA	IGM 100/978	Benson et al. (2017), #321	22.05	1	22.05	IGM 100/978	CT	Balanoff
Khaan mckennai	Non-avian maniraptoriform	21.9198	16.3018	27.5378	67.62	NA	IGM 100/1127	Benson et al. (2017), #452	8.8	1	8.8	IGM 100/973	CT	Balanoff
Shuvuuia deserti	Non-avian maniraptoriform	3.0497	2.2681	3.8313	33	NA	IGM 100/1304	Benson et al. (2017), #574	0.78	1	0.78	IGM 100/977	CT	Balanoff
Ornithomimus edmontonicus	Non-avian maniraptoriform	83.5238	62.1167	104.9309	110	NA	ROM 851	Benson et al. (2017), #522	49.89	1	49.89	NMC 12228	GDI (this study)	Photos o
Stenonychosaurus inequalis	Non-avian maniraptoriform	47.3759	35.2335	59.52	89.5	NA	MOR 748 (MTC)	Benson et al. (2017), #621	45	1	45	RTMP 86.36.457 & 79.8.1	Water displacement	Currie &
Archaeopteryx lithographica[2]	Non-avian maniraptoriform	0.3446	0.2563	0.4329	14.93	NA	BMNH 37001	Gatesy, 1991	1.52	1	1.52	BMNH 37001	CT	Domingu
Acrocanthosaurus atokensis	Non-maniraptoriform theropod	3454.954	2569.449	4340.46	426	NA	NCSM 14345	Benson et al. (2017), #242	51.55	0.42	122.74	OMNH 10146	CT	Franzosa
Acrocanthosaurus atokensis	Non-maniraptoriform theropod	3454.954	2569.449	4340.46	426	NA	NCSM 14345	Benson et al. (2017), #242	38.05	0.31	122.74	OMNH 10146	CT	Franzosa
Allosaurus fragilis	Non-maniraptoriform theropod	2541.814	1890.347	3193.28	381	NA	AMNH 680	Benson et al. (2017), #253	41.37	0.42	98.5	UUVP 294	GDI (this study)	Photo of
Allosaurus fragilis	Non-maniraptoriform theropod	2541.814	1890.347	3193.28	381	NA	AMNH 680	Benson et al. (2017), #253	30.54	0.31	98.5	UUVP 294	GDI (this study)	Photo of
Carcharodontosaurus saharicus[3]	Non-maniraptoriform theropod	3269.147	2431.265	4107.03	417.52	NA	BSP 1922 X46	Benson et al. (2017), #307	94.08	0.42	224	SGM-Din 1	CT	Larsson
Carcharodontosaurus saharicus	Non-maniraptoriform theropod	3269.147	2431.265	4107.03	417.52	NA	BSP 1922 X46	Benson et al. (2017), #307	69.44	0.31	224	SGM-Din 1	CT	Larsson
Carnotaurus sastrei	Non-maniraptoriform theropod	1641.829	1221.028	2062.63	325	NA	MACN CH 894	Benson et al. (2017), #308	45.49	0.42	108.3	MACN CH-894	CT	Cerroni & (2019)
Carnotaurus sastrei	Non-maniraptoriform theropod	1641.829	1221.028	2062.63	325	NA	MACN CH 894	Benson et al. (2017), #308	33.57	0.31	108.3	MACN CH-894	CT	Cerroni & (2019)
Giganotosaurus carolinii	Non-maniraptoriform theropod	6136.771	4563.916	7709.625	525	NA	MUCPv-Ch1	Benson et al. (2017), #399	94.5	0.42	225	MUCPv-CH 1	CT	Paulina-C (2010)
Giganotosaurus carolinii	Non-maniraptoriform theropod	6136.771	4563.916	7709.625	525	NA	MUCPv-Ch1	Benson et al. (2017), #399	69.75	0.31	225	MUCPv-CH 1	CT	Paulina-C (2010)
Majungasaurus crenatissimus	Non-maniraptoriform theropod	1614.201	1200.481	2027.92	323	NA	FMNH PR 2278	Benson et al. (2017), #482	37.51	0.42	89.32	FMNH PR 2100	CT	Sampson
Majungasaurus crenatissimus	Non-maniraptoriform theropod	1614.201	1200.481	2027.92	323	NA	FMNH PR 2278	Benson et al. (2017), #482	27.69	0.31	89.32	FMNH PR 2100	CT	Sampson

Sinraptor dongi[4]	Non-maniraptoriform theropod	1122.287	834.645	1409.929	283	NA	TMP 93.115.1	Campione& Evans, 2020	39.9	0.42	91	IVPP 10600	CT	Paulina-((2012)
Sinraptor dongi	Non-maniraptoriform theropod	1122.287	834.645	1409.929	283	NA	TMP 93.115.1	Campione& Evans, 2020	29.45	0.31	91	IVPP 10600	CT	Paulina-((2012)
Tarbosaurus bataar	Non-maniraptoriform theropod	2345.113	1744.06	2946.165	370	NA	MPC-D 552/1	Benson et al. (2017), #610	66.86	0.42	159.2	PIN, no. 553-3/1	Estimated from latex half-cast	Saveliev
Tarbosaurus bataar	Non-maniraptoriform theropod	2345.113	1744.06	2946.165	370	NA	MPC-D 552/1	Benson et al. (2017), #610	49.35	0.31	159.2	PIN, no. 553-3/1	Estimated from latex half-cast	Saveliev
Tyrannosaurus rex [5]	Non-maniraptoriform theropod	8070.46	6002.001	10138.919	580	NA	FMNH PR 2081	Persons et al., 2020	173.96	0.42	414.19	FMNH PR 2081	CT	Hurlburt
Tyrannosaurus rex	Non-maniraptoriform theropod	8070.46	6002.001	10138.919	580	NA	FMNH PR 2081	Persons et al., 2020	128.40	0.31	414.19	FMNH PR 2081	CT	Hurlburt
Tyrannosaurus rex	Non-maniraptoriform theropod	6430.357	4782.256	8078.457	534	NA	CM 9380	Persons et al., 2020	160.34	0.42	381.76	AMNH 5029	CT	Hurlburt
Tyrannosaurus rex	Non-maniraptoriform theropod	6430.357	4782.256	8078.457	534	NA	CM 9380	Persons et al., 2020	118.35	0.31	381.76	AMNH 5029	CT	Hurlburt
Tyrannosaurus rex	Non-maniraptoriform theropod	6430.357	4782.256	8078.457	534	NA	CM 9380	Persons et al., 2020	217.11	0.57	381.76	AMNH 5029	CT	Morhardt
Tyrannosaurus rex	Non-maniraptoriform theropod	5515.247	4101.69	6928.805	505	NA	BHI 3033	Persons et al., 2020	131.73	0.42	313.64	AMNH 5117	CT	Hurlburt
Tyrannosaurus rex	Non-maniraptoriform theropod	5515.247	4101.69	6928.805	505	NA	BHI 3033	Persons et al., 2020	97.23	0.31	313.64	AMNH 5117	CT	Hurlburt
Amargasaurus cazaui	Non-theropod dinosaur	10194.61	7581.73	12807.49	505	388	MACN-N 15	Benson et al. (2017), #20	35.28	0.42	84	MACN-N 15	CT	Paulina (
Amargasaurus cazaui	Non-theropod dinosaur	10194.61	7581.73	12807.49	505	388	MACN-N 15	Benson et al. (2017), #20	26.04	0.31	84	MACN-N 15	CT	Paulina (
Apatosaurus sp.	Non-theropod dinosaur	41268.719	30691.546	51845.891	845	640	CM 3018	Benson et al. (2017), #33	43.04	0.42	102.48	BYU 17096	CT	Balanoff
Apatosaurus sp.	Non-theropod dinosaur	41268.719	30691.546	51845.891	845	640	CM 3018	Benson et al. (2017), #33	31.77	0.31	102.48	BYU 17096	CT	Balanoff
Diplodocus sp.	Non-theropod dinosaur	14813.081	11016.488	18609.673	563	460	USNM 10865	Benson et al. (2017), #86	42	0.42	100	CM 11161	CT	L. M. Wit (2023)
Diplodocus sp.	Non-theropod dinosaur	14813.081	11016.488	18609.673	563	460	USNM 10865	Benson et al. (2017), #86	31	0.31	100	CM 11161	CT	L. M. Wit (2023)
Giraffatitan brancai	Non-theropod dinosaur	34003.143	25288.137	42718.148	730	654	HMN SII	Benson et al. (2017), #107	130.2	0.42	310	MB.R.2223.1	Plasticine cast	Janensch
Giraffatitan brancai	Non-theropod dinosaur	34003.143	25288.137	42718.148	730	654	HMN SII	Benson et al. (2017), #107	96.1	0.31	310	MB.R.2223.1	Plasticine cast	Janensch
Euoplocephalus tutus	Non-theropod dinosaur	2329.632	1732.548	2926.717	278	244	AMNH 5404	Benson et al. (2017), #766	34.73	0.42	82.7	AMNH 5337	GDI (this study)	Hopson (
Euoplocephalus tutus	Non-theropod dinosaur	2329.632	1732.548	2926.717	278	244	AMNH 5404	Benson et al. (2017), #766	25.64	0.31	82.7	AMNH 5337	GDI (this study)	Hopson (

Kentrosaurus aethiopicus	Non-theropod dinosaur	1596.86	1187.585	2006.136	245	210	HMN composite specimen	Benson et al. (2017), #813	22.092	0.42	52.6	HMN Ki 124	GDI (this study)	Galton (1
Kentrosaurus aethiopicus	Non-theropod dinosaur	1596.86	1187.585	2006.136	245	210	HMN composite specimen	Benson et al. (2017), #813	16.306	0.31	52.6	HMN Ki 124	GDI (this study)	Galton (1
Stegosaurus ungulatus	Non-theropod dinosaur	6953.916	5171.627	8736.205	425	352	YPM 1853	Benson et al. (2017), #927	26.964	0.42	64.2	CM 106	GDI (this study)	Galton (2
Stegosaurus ungulatus	Non-theropod dinosaur	6953.916	5171.627	8736.205	425	352	YPM 1853	Benson et al. (2017), #927	19.902	0.31	64.2	CM 106	GDI (this study)	Galton (2
Protoceratops andrewsi	Non-theropod dinosaur	82.695	61.5	103.889	93	62	AMNH 6424	Benson et al. (2017), #892	12.6	0.42	30	AMNH 6466	GDI (Jerison, 1973)	Brown &
Protoceratops andrewsi	Non-theropod dinosaur	82.695	61.5	103.889	93	62	AMNH 6424	Benson et al. (2017), #892	9.3	0.31	30	AMNH 6466	GDI (Jerison, 1973)	Brown &
Triceratops sp.	Non-theropod dinosaur	13274.61	9872.328	16676.893	493	490	AMNH 5033	Benson et al. (2017), #950	58.8	0.42	140	USNM 2416	GDI (Jerison, 1973)	Hay (190
Triceratops sp.	Non-theropod dinosaur	13274.61	9872.328	16676.893	493	490	AMNH 5033	Benson et al. (2017), #950	43.4	0.31	140	USNM 2416	GDI (Jerison, 1973)	Hay (190
Edmontosaurus annectens	Non-theropod dinosaur	6610.079	4915.916	8304.243	512.3	250.5	AMNH 5730	Benson et al. (2017), #757	126	0.42	300	YPM 618	GDI (Jerison, 1973)	Lull & Wr
Edmontosaurus annectens	Non-theropod dinosaur	6610.079	4915.916	8304.243	512.3	250.5	AMNH 5730	Benson et al. (2017), #757	93	0.31	300	YPM 618	GDI (Jerison, 1973)	Lull & Wr
Hypacrosaurus altispinus	Non-theropod dinosaur	3689.151	2743.622	4634.681	395	222	CMN 8501	Benson et al. (2017), #800	85.53	0.31	275.9	ROM 702	CT	Evans et
Hypacrosaurus altispinus	Non-theropod dinosaur	3689.151	2743.622	4634.681	395	222	CMN 8501	Benson et al. (2017), #800	115.88	0.42	275.9	ROM 702	CT	Evans et
Iguanodon bernissartensis	Non-theropod dinosaur	8268.265	6149.108	10387.421	490	337.5	RBINS R51	Benson et al. (2017), #805	149.94	0.42	357	RBINS R51	CT	Lauters (
Iguanodon bernissartensis	Non-theropod dinosaur	8268.265	6149.108	10387.421	490	337.5	RBINS R51	Benson et al. (2017), #805	110.67	0.31	357	RBINS R51	CT	Lauters (

500 **Table 2**: Estimates of telencephalic neuron counts (N; excluding the olfactory system) in Mesozoic dinosaurs. Our inferences are compared with those

501 presented by Herculano-Houzel (2022; HH) if respected species were included in both studies (see text for the rationale of our sample composition).

502 Minimum and maximum estimates based on both avian and non-avian sauropsid regressions are provided.

Species	Group	MBr range (g)	N (non-avian_min)	N (non-avian_max)	N (avian_min)	N (avian_max)	MBr (g), HH	N (non-avian), HH	N (avian), HH
Acrocanthosaurus atokensis	Non-maniraptoriform theropod	38.05-51.55	133,214,329	162,531,415	568,823,512	707,197,873	191	311.3M	2,116M
Allosaurus fragilis	Non-maniraptoriform theropod	30.54-41.37	115,336,233	140,718,805	485,812,035	603,992,681	168	287.8M	1,921M
Amargasaurus cazaui	Non-theropod dinosaur	26.04-35.28	103,912,541	126,781,049	433,393,846	538,823,026			
Apatosaurus sp.	Non-theropod dinosaur	31.77-43.04	118,367,821	144,417,569	499,807,465	621,392,698			
Archaeopteryx lithographica	Non-avian maniraptoriform	1.52-1.52	16,163,324	16,163,324	56,526,229	56,526,229	1.47-1.76	15.8M-17.7M	54.2M-62.1M
Bambiraptor feinbergi	Non-avian maniraptoriform	14.00-14.00	69,204,922	69,204,922	277,742,256	277,742,256	14	62.9M	295.8M
Carcharodontosaurus saharicus	Non-maniraptoriform theropod	69.44-94.08	197,550,205	241,025,981	875,593,081	1,088,593,477			
Carnotaurus sastrei	Non-maniraptoriform theropod	33.57-45.49	122,728,852	149,738,351	519,999,703	646,496,982			
Citipati osmolskae	Non-avian maniraptoriform	22.05-22.05	93,187,090	93,187,090	384,673,367	384,673,367	22.62	84.4M	424.5M
Diplodocus sp.	Non-theropod dinosaur	31.00-42.00	116,483,666	142,118,759	491,105,154	610,573,426	57	148.5M	851.4M
Edmontosaurus annectens	Non-theropod dinosaur	93.00-126.00	239,210,068	291,854,121	1,079,618,782	1,342,251,315	150	268.5M	1,764M
Euoplocephalus tutus	Non-theropod dinosaur	25.64-34.73	102,856,358	125,492,426	428,574,120	532,830,834	41	121.4M	664.4M
Giganotosaurus carolinii	Non-maniraptoriform theropod	69.75-94.50	198,127,419	241,730,226	878,393,994	1,092,075,751			
Giraffatitan brancai	Non-theropod dinosaur	96.10-130.20	244,403,230	298,190,166	1,105,301,670	1,374,181,929	186	306.3M	2,075M

Species	Type	Range	Value 1	Value 2	Value 3	Value 4	N	Mass 1	Mass 2
Hypacrosaurus altispinus	Non-theropod dinosaur	85.53-115.88	226,443,950	276,279,513	1,016,710,016	1,264,044,148			
Iguanodon bernissartensis	Non-theropod dinosaur	110.67-149.94	268,078,950	327,076,310	1,223,031,388	1,520,551,064			
Kentrosaurus aethiopicus	Non-theropod dinosaur	16.31-22.09	76,473,452	93,303,314	309,828,433	385,198,580	24	87.5M	443.9M
Khaan mckennai	Non-avian maniraptoriform	8.80-8.80	51,057,468	51,057,468	199,095,862	199,095,862	8.83	47.4M	209.1M
Majungasaurus crenatissimus	Non-maniraptoriform theropod	27.69-37.51	108,177,375	131,984,465	452,902,502	563,077,439			
Ornithomimus edmontonicus	Non-avian maniraptoriform	49.89-49.89	159,082,339	159,082,339	690,786,497	690,786,497	87.85	193.6M	1,179M
Protoceratops andrewsi	Non-theropod dinosaur	9.30-12.60	52,939,454	64,590,081	207,143,081	257,533,564	28	96.1M	498.5M
Shuvuuia deserti	Non-avian maniraptoriform	0.78-0.78	10,441,072	10,441,072	35,034,828	35,034,828	0.83	11.2M	35.2M
Sinraptor dongi	Non-maniraptoriform theropod	29.45-39.90	112,635,163	137,423,299	473,371,729	588,526,094			
Stegosaurus ungulatus	Non-theropod dinosaur	19.90-26.96	87,136,573	106,313,117	357,418,565	444,365,684	22.5	84.1M	422.8M
Tarbosaurus bataar	Non-maniraptoriform theropod	49.35-66.86	157,956,586	192,718,814	685,437,196	852,179,485			
Triceratops sp.	Non-theropod dinosaur	43.40-58.80	145,204,105	177,159,837	625,098,265	777,162,256	72.2	171.7M	1,017M
Troodon formosus	Non-avian maniraptoriform	45.00-45.00	148,688,479	148,688,479	641,536,735	641,536,735	4.1	121.4M	664.4M
Tyrannosaurus rex AMNH 5029	Non-maniraptoriform theropod	118.35-160.34	280,116,860	341,763,457	1,283,275,361	1,595,450,236	343	445.5M	3,289M
Tyrannosaurus rex AMNH 5117 (Morhardt, 2016)	Non-maniraptoriform theropod	217.11-217.11	416,817,413	416,817,413	1,982,735,561	1,982,735,561			
Tyrannosaurus rex AMNH 5117	Non-maniraptoriform theropod	97.23-131.73	246,277,083	300,476,406	1,114,581,594	1,385,719,325			
Tyrannosaurus rex FMNH PR 2081	Non-maniraptoriform theropod	128.40-173.96	295,481,827	360,509,864	1,360,525,267	1,691,492,274	202	322.2M	2,207M

Phylogenetic modeling of neurological variables

We used data on extant sauropsids to place brain size variables for Mesozoic

dinosaurs into their phylogenetic context. To examine variations in the relative size of the

brain in fossil taxa and to calculate potential neuronal scaling regimes in extinct dinosaurs,

we relied on log-transformed published data on brain mass, telencephalon mass and

telencephalic neuronal numbers in extant groups (see below). Allometric equations were

calculated with least squares linear regressions using phylogenetic generalized least

squares (PGLS) to account for phylogenetic relatedness (Garland and Ives, 2000). PGLS

allows the covariance matrix to be modified to accommodate the degree to which trait

evolution deviates from Brownian motion, through a measure of phylogenetic correlation,

Pagel's λ (Pagel, 1999). PGLS and maximum likelihood estimates of λ were performed using

the *ape* (Paradis & Schliep, 2019) and *nlme* (Pinheiro et al., 2017) packages in R.

To compare differences in relative brain size across groups, phylogenetically

corrected ANCOVA with Tukey post hoc comparisons were performed using a modified

version of the *multcomp* package (Hothorn et al., 2015). Because of the uncertainty in

estimating both brain mass and body mass in Mesozoic dinosaurs, we opted to test for inter-

group differences in two datasets: One with the greatest possible relative brain size i.e. the

lowest body mass estimate (lower PPE) for each species and the brain mass estimated from

the highest assumed endocranial fill (42%), and one with the lowest relative brain size i.e.

the highest body mass estimates (upper PPE) and the lowest assumed endocranial fill

(31%). Since we assumed that the brain filled 100% of the endocranial cavity in

maniraptoriform theropods, inferred relative brain mass for these species was only affected

by differences in the applied body mass estimates.

As mentioned above, an important assumption of Herculano-Houzel (2022) is that

theropods in general had relative brain sizes similar to birds. However, there is notable

discontinuity in relative brain size and brain morphology between maniraptoriforms and more

basal non-maniraptoriform theropods (Fig. 1, Balanoff et al.,2013). Because of this, we

divided our sample of Mesozoic theropods into these two groups (Tabs. 1 & 2). For *T. rex*,

mean values for the three available adult brain mass and corresponding body mass

estimates were used. We grouped Sauropodomorpha and Ornithischia together as non-

theropod dinosaurs and compared relative brain size in this group with that in the two

theropod samples. PGLS models of brain mass vs. body mass with clade as a covariate

were used to test if relative brain size in these groups differs significantly between them and

from extant birds and/or non-avian sauropsids. Relative brain size data for 63 extant non-

avian sauropsids (including lepidosaurs, crocodylians, and turtles) and 84 bird species (not

including members of the large-brained clade Telluraves) were derived from Hurlburt (1996),

Chentanez et al. (1983) and Roese-Miron et al. (2023) and are listed in Supplementary File

2. Importantly, these sources provide brain mass estimates excluding the olfactory tracts and

bulbs and thus fit our brain size estimates for Mesozoic dinosaurs.

To test for differences in relative brain size we built a phylogenetic tree for all 175

fossil and extant species. To construct the phylogeny of bird species, we extracted 1,000

fully resolved trees from birdtree.org (Jetz et al., 2012) using the Hackett et al. (2008)

backbone, and built a maximum clade credibility (MCC) tree using *phangorn* (Schliep, 2011).

For lepidodaurs, we followed Kverková et al. (2022) and used a recently published species

level time-calibrated phylogeny (Tonini et al., 2016) and built a MCC tree the same way as

we did with the bird phylogeny. For turtles and crocodylians, we used the Timetree of Life

(Kumar et al., 2017). We then stitched the trees together manually, using the divergence

times from the Timetree of Life. For Mesozoic dinosaurs (28 species) we use an updated

version of the composite phylogeny of Benson et al. (2014, 2018). Phylogenies for fossil

dinosaurs, extant non-avian sauropsids, and birds were stitched together manually using

Mesquite (Maddison and Maddison, 2023). We opted to set all branch lengths to 1. This

provides adequately standardized branch lengths for other phylogenetic analyses, such as

independent contrasts (Garland et al., 1992).

Tree building procedures were the same for telencephalic neuron count regressions,

but trees used here included branch lengths. We calculate regression lines between brain

mass and telencephalic number of neurons for extant non-telluravian birds and non-avian

sauropsids. Analogous to Herculano-Houzel (2022), these regressions were then used to

estimate telencephalic neuron counts in dinosaurs, applying either an avian or a reptilian

scaling regime. Since our estimates are based on brain portion endocasts that exclude the

olfactory system, our telencephalic neuron counts correspond to the pallium and subpallium.

Data on whole brain and telencephalic neuron counts as well as on total telencephalic and

brain mass (including olfactory tract and bulbs, since neuron count data excluding these

structures are currently unavailable for the taxa considered here) for birds (n = 112) were

derived from Kverková et al. (2022) and Sol et al. (2022), for non-avian sauropsids (n = 108)

from Kverková et al. (2022) and for mammals (n = 39) from Herculano-Houzel et al. (2015).

The dataset is included in Supplementary File 3.

To get a more precise estimate of the possible number of telencephalic neurons in *T.*

rex, we also modeled scaling regimes for telencephalon mass vs. telencephalic neuron

numbers in extant sauropsids (non-avian sauropsids and non-telluravian birds), using the

same references listed above. We then calculated telencephalic neuron numbers in *T. rex*

using the obtained scaling regimes and applying the telencephalic volumes estimated with a

comparative 3D landmark approach by Morhardt (2016) (referred to as "cerebral

hemispheres" therein, excluding olfactory bulbs and tracts) for specimen AMNH FR 5117.

Based on the estimates of Morhardt (2016), we also comparatively assessed the mass of the

telencephalon and cerebellum in *T. rex*.

We want to note that our neuron count estimates might be biased by the fact that we

predict neuron numbers in the pallium and subpallium (telencephalon excluding the olfactory

system) based on total telencephalic neuron counts (including the olfactory system) here.

This is an issue that in parts also applies to Herculano-Houzel (2022) and which we cannot

circumvent due to limitations of the available raw data.

Results

Relative brain size.

Overall, we do not recover an unambiguous discontinuity in relative brain size between extant non-avian sauropsids and Mesozoic dinosaurs, including large theropods like *T. rex*. However, our analyses support the assumption that grade shifts in relative brain size occurred among non-avian dinosaurs. PGLS models showed a significant difference in relative brain size (intercept) between non-maniraptoriform theropods, such as *T. rex*, and the more bird-like Maniraptoriformes, which tend to have larger brains than other dinosaurs (PGLS, max: $F_{4,169}$= 18.11, $p \le 0.0001$, $\lambda = 0.601$); min: $F_{4,171}$= 23.50, $p \le 0.0001$, $\lambda = 0.601$). Post-hoc analysis shows that both the maximum and minimum relative brain size estimates for non-maniraptoriform theropods are not significantly different from what would be expected from extant non-avian sauropsids (Tab. 3). However, both minimum and maximum relative brain size estimates for these animals are significantly smaller than what would be expected for extant birds (Tab. 3; Fig. 4). On the other hand, we found that maniraptoriforms show no significant differences in relative brain size compared to birds (Tab. 3) and only exhibited significantly larger brains than non-avian sauropsids when the maximum relative brain size was assumed (Tab. 3; Fig 4; note that some maniraptoriforms such as *Shuvuuia deserti* cluster with non-avian sauropsids rather than with birds). Therefore, while most of the maniraptoriform species in our dataset exhibit relative brain sizes within the spectrum of living birds and larger than those of extant non-avian sauropsids, non-maniraptoriform theropods likely showed markedly smaller relative brain sizes within the range of the latter (Fig 4a-b). Both groups of theropods exhibit significantly larger brains than the non-theropod dinosaurs of the clades Sauropodomorpha and Ornithischia (Tab. 3), data for which we pool here. However, maximum relative brain sizes in these dinosaurs is also not significantly smaller than would be expected for non-avian sauropsids of their body mass. If minimum figures are assumed, relative brain size in these animals would have been significantly smaller than in extant non-avian sauropsids (Tab. 3).

Table 3: Tukey post hoc comparisons for a phylogenetic ANCOVA testing for differences in relative brain size between groups of fossil dinosaurs and extant sauropsids. Significant *p*-values are shown in bold.

Minimum relative brain size				
	Aves	Non-avian Maniraptoriformes	Non-maniraptoriform Theropoda	Non-theropod Dinosauria
Non-avian Maniraptoriformes	0.27			
Non-maniraptoriform Theropoda	**> 0.00001**	**0.00012**		
Non-theropod Dinosauria	**> 0.00001**	**> 0.00001**	**0.048**	
Non-avian Sauropsida	**> 0.00001**	0.075	0.93	**0.020**
Maximum relative brain size				
	Aves	Non-avian Maniraptoriformes	Non-maniraptoriform Theropoda	Non-theropod Dinosauria
Non-Avian Maniraptoriformes	0.73			
Non-Maniraptoriform Theropoda	**0.00061**	**0.0065**		
Non-theropod Dinosauria	**> 0.00001**	**> 0.00001**	**0.048**	
Non-avian Sauropsida	**> 0.00001**	**0.0081**	0.94	0.61

617 **Table 4**: Regression parameters for different models describing the scaling of neurological traits in

618 extant non-avian sauropsids (reptiles) and non-telluravian birds. Pagel's λ (ranging between 0 - 1)

619 was used to quantify the phylogenetic signal. See methods for details. SE = standard error.

	Model	Slope (SE)	Intercept (SE)	λ
Avian GLS	log(Tel Neurons N) ~ log(Brain Mass)	0.821 (0.043)	17.5 (0.063)	0
Avian PGLS	Log(Tel Neurons N) ~ log(Brain Mass)	0.717 (0.05)	17.55 (0.12)	0.50
Reptilian GLS	Log(Tel Neurons N) ~ log(Brain Mass)	0.615 (0.03)	16.347 (0.06)	0
Reptilian PGLS	Log(Tel Neurons N) ~ log(Brain Mass)	0.655 (0.03)	16.324 (0.17)	0.82
Avian PGLS	log(Brain Mass) ~ log(Body Mass)	0.584 (0.02)	-2.584 (0.25)	0.96
Reptilian PGLS	log(Brain Mass) ~ log(Body Mass)	0.56 (0.03)	-4.077 (0.26)	0.70

Numbers of neurons

We re-calculated estimates for the number of forebrain neurons in fossil dinosaurs

based on PGLS-derived regressions of brain size vs. number of telencephalic neurons in

extant non-avian sauropsids and birds. Our neuron count estimates are listed in Table 2 and

are compared to those of Herculano-Houzel (2022), whereas regression parameters are

provided in Table 4 . While many of the estimates do not differ notably from one another, the

differences for some taxa, especially large theropods, are striking. For *T. rex*, Herculano-

Houzel (2023) provided an estimate of 300-450M forebrain neurons if modeled based on

extant non-avian sauropsids, and 2-3B based on an avian regression. In contrast, we

estimated a range of 245-360M neurons with a reptilian regression and ~1-2B with an avian

one (Tab. 2, Fig. 5). Using the forebrain volumes estimated for *T. rex* by Morhardt (2016), we

predict 133-166M telencepahlic neurons in this species if applying a reptilian scaling and

0.989 to 1.25B based on an avian scaling (Fig. 5).

Figure 4: Relative brain size and forebrain neuronal numbers in Mesozoic dinosaurs and other

amniotes. a. The log-transformed mass of the brain is plotted as a function of the mass of the body for

extant and fossil sauropsids. In the case of fossil species, the mean body and/or brain size is shown

along with standard deviations. The orange dotted line represents the regression line for avian

species (excluding the large-brained clade Telluraves) obtained from PGLS while the pink one

represents the same for extant non-avian sauropsids ("reptiles" in the colloquial sense). **b.** A detail of

the plot shown in **a**. to illustrate the range of relative brain size in *Tyrannosaurus rex* and other

Mesozoic dinosaurs that we consider plausible. Besides our own brain size estimates, the plot

contains those from Morhardt (2016) (specimen AMNH FR 5117, endocranial fill = 57%) and Balanoff

et al. (2013) (specimen AMNH 5029, endocranial fill = 100%, assumed MBd = 5840 kg) **c.** Plot

showing log-transformed brain mass for different groups of extant amniotes plotted against body

mass. **d.** Plot showing log-transformed numbers of telencephalic neurons as a function of the mass of

the forebrain, illustrating neuronal density. Note that non-avian sauropsids and non-primate mammals

differ only moderately from one another here, although mammals have markedly larger brains relative

to body size, as shown in **c**. See methods for data sources. Silhouettes were taken from PhyloPic

(listed clockwise from top left): *Anas* (in public domain) *Morunasaurus* (in public domain),

Dromaeosaurus (in public domain), *Stegosaurus* (by Matt Dempsey), *Allosaurus* (by Tasman Dixon),

Tyrannosaurus (by Matt Dempsey), *Corvus* (in public domain), *Hylobates* (by Kai Caspar), *Antidorcas*

(by Sarah Wenig).

Figure 5: Predicted numbers of neurons in telencephalon (excluding olfactory tracts and bulbs) of *Tyrannosaurus rex*. Points represent the estimated number of neurons in three adult specimens of *T. rex* using different inference methods. Estimates from this study using a regression that takes phylogenetic relationships into account (PGLS, see methods, filled circle, triangle, square and cross) are plotted in cyan. The estimates from Herculano-Houzel (2022) based on a non-phylogenetic regression are shown in red (crossed square, asterisk). Different underlying ratios of brain volume:endocranial volume are annotated. On the left predicted numbers of forebrain neurons (based on either the avian or extant non-avian sauropsids scaling regime) based on the estimated volume of the brain portion of the endocast are shown. On the right, analogous to that, the predicted numbers of telencephalic neurons based on volumetric estimates by Morhart (2016) is plotted.

Figure 6: Relative size of telencephalon (excluding olfactory bulb and tracts) and cerebellum in

***T. rex.* a:** The log-transformed mass of the telencephalon in extant non-telluravian birds and non-

avian sauropsids is plotted as a function of the mass of the rest of the brain (total brain -

telencephalon - cerebellum volume). Green dots show the maximum, mid and minimum estimates for

the mass of the *T. rex* telencephalon as estimated from digital endocasts of AMNH 5117 by Morhardt

(2016). The orange dotted line represents the regression line for non-telluravian bird species obtained

from PGLS while the pink represents the same for non-avian sauropsids. **B**: Analogous plot to **a**, but

for the cerebellum. Note that telencephalic mass in extant species includes the olfactory bulb and

tracts.

Discussion of empirical results

We want to emphasize two aspects of our empirical findings that contrast with those

of Herculano-Houzel (2022). Firstly, we did not find relative brain size to separate large-

bodied non-maniraptoriform theropods such as *T. rex* from extant non-avian sauropsids;

rather, our data support a grade shift in this trait between maniraptoriform and non-

maniraptoriform theropods, which importantly relates to an increase in endocranial fill. As we

have argued beforehand, we see no support for the brains of non-maniraptoriform

theropods, sauropodomorphs, and most ornithischians to have filled the endocranial cavity in

a bird-like fashion. However, we are aware that such a condition, or one that is at least

intermediate between modern birds and crocodylians has been proposed for all of these

groups at one point (e.g., Knoll & Schwarz-Wings, 2009; Morhardt, 2016; Balanoff et al.,

2013; Knoll et al., 2021). Obviously, future research might still significantly change our

understanding of endocranial tissue organization in Mesozoic dinosaurs and thus challenge

the assumptions that we make here based on comparative evidence from adult crocodylians.

Our approach suggests that relative brain size in all dinosaurs, except for the majority of

maniraptoriform theropods, does not differ significantly from values present in extant non-

avian reptiles. These results agree with previous conclusions (Hurlburt et al., 2013;

Morhardt, 2016). Nonetheless, we want to stress that it remains unclear how meaningful the

transfer of brain size scaling rules established for the given extant bird (32g – 120kg) and

non-avian sauropsid (1g – 71kg) datasets to large-bodied dinosaurs actually is. Brain-body

size ratios in extant cetaceans drop dramatically in taxa that evolved multi-ton body masses

(Tartarelli & Bisconti, 2006), suggesting that allometric trajectories need to be accounted for.

However, the restricted body mass spectrum of extant birds and reptiles as well as the

limited availability of large-bodied crocodylians and turtles for neurological research hinders

the compilation of such datasets for sauropsids. Furthermore, the scarcity of complete and

adult dinosaur endocasts from taxa that also preserve stylopodial elements to derive body

mass estimates from, limits our understanding of differences in brain size scaling between

taxa. Different clades of mammals and birds have been shown to have distinct allometric

relationships for relative brain size (Ksepka et al., 2020; Smaers et al., 2021). The same

might have been the case in non-avian dinosaurs, biasing comparisons between the

groupings we selected here. In addition to that, there might also be temporal effects on

relative brain size. Such a phenomenon appears to be rampant in mammalian evolution

during the Cenozoic (Bertrand et al., 2022). To our knowledge, this pattern has not been

described yet in other vertebrate groups but should be considered in future studies on brain

evolution in long-lived clades such as dinosaurs.

Secondly, our empirical findings do not support Herculano-Houzel's (2022) claim of

exceptionally high telencephalic neuron counts in dinosaurs, particularly in *T. rex* and other

large theropods. Instead, *T. rex* likely did not exhibit more than approximately 1.5 bn (or at a

maximum 2 bn) telencephalic neurons, even when an avian neuronal density is assumed. If

we assume reptilian neuronal densities, it might even have exhibited neuron numbers an

order of magnitude lower than the 3.3 B suggested by Herculano-Houzel (2022). Apart from

the difficulty of estimating brain mass from a dinosaurian endocast, there is one additional

caveat to our neuron count estimates that needs to be acknowledged and that also applies

to Herculano-Houzel's (2022) study: Telencephalic neuron numbers can only be reliably

derived from total brain mass when the proportions of the studied brains are comparable.

Since brain morphology in many dinosaurian lineages differs significantly from both extant

birds and non-avian sauropsids (Fig. 1, Paulina-Carabajal et al., 2023), biases are thus

ingrained into such estimates. Volumetric modeling of brain regions from endocasts, on

which we relied here for *T. rex* exclusively, could potentially ameliorate this problem to some

degree (Morhardt, 2016) but it is challenging and not yet widely used. For *T. rex*, such

inferences yield lower telencephalic neuron numbers than would be hypothesized based on

our total brain volume estimates, if reptilian scaling rules are applied (Fig. 5). They overlap if

an avian neuron count scaling is assumed (Fig. 5).

We argue that there is little reason to assume that the brains of non-maniraptoriform

theropods such as *T. rex* had a telencephalic neuronal density similar to that of extant birds.

In living sauropsids, relative brain size is positively correlated with neural density (Kverková

et al., 2022). We showed that relative brain size in non-maniraptoriform theropods did not

differ significantly from that encountered in extant non-avian sauropsids. Consequently,

relative brain size cannot be used as an argument to defend elevated neuron densities in

these animals. The presence of endothermy in dinosaurs (see below) does also not entail

avian neuronal density (contra Herculano-Houzel, 2022), as mammals exemplify. Mammals

have not only evolved endothermy, but also exhibit large relative brain sizes (Fig. 4C; Tsuboi

et al., 2018) and a unique multilayered cerebral neuroarchitecture (Briscoe & Ragsdale,

2018), yet their forebrain neuronal density is only moderately elevated compared to extant

non-avian reptiles (Kverková et al., 2022). With respect to birds, it is remarkably low (at least

if not considering anthropoid primates, Fig. 4D). This suggests that metabolic rate and

neuronal density are not tightly coupled. Finally, the shape of the endocast and volumetric

estimates of its forebrain and cerebellar portions (compare Fig. 6) suggest that the brains of

T. rex and other large non-maniraptoriform theropods were not dissimilar to those of extant

crocodylians (Rogers, 1998; Hurlburt et al., 2013; Morhardt, 2016), which reflect the

plesiomorphic archosaurian condition (Fabbri & Bhullar, 2022). Given this morphological

conservatism, it appears appropriate to assume reptilian neuronal densities for these

animals. If we would strictly model the brains of Mesozoic dinosaurs after

Applying the same logic, it is tempting to speculate that the increased neuronal

density that sets birds apart from other sauropsids coevolved with the marked changes in

brain morphology and size that occurred in maniraptoriform theropods. If we indeed assume

that an avian-like brain organization and high neuronal density emerged early within this

clade's history, it is plausible that the dinosaurs with the highest neuron counts, perhaps

above the extant avian range, are represented by the largest-bodied taxa within this group

(for which no endocasts are currently available). These include bizarre animals such as the

ornithomimosaur *Deinocheirus mirificus* (~ 7 t), the immense, scythe-clawed *Therizinosaurus*

cheloniformis (~ 5 t) and the giant oviraptorosaur *Gigantoraptor erlianensis* (~ 2 t).

Alternatively, the emergence of volancy in small stem-birds similar to *Archaeopteryx* might

have driven the evolution of elevated neuron densities, since active flight likely imposes

constraints on skull and brain size (Olkowicz et al., 2016; Shatkovska & Ghazali, 2021).

However, the lack of reliable morphological markers to infer neuron density renders all these

notions speculative. Such a vagueness is inherent to predictions about the biology of extinct

taxa without close living relatives and obviously needs to be acknowledged. The main

argument for assuming avian neuronal densities in any group of Mesozoic dinosaurs is that

the emergence of this trait within the avian stem-group cannot be reliably dated and thus

might have significantly preceded the origins of crown birds. Hence, it needs to be

acknowledged that both a non-avian sauropsid and an avian neuron density (as well as

intermediate conditions) in these animals can principally be justified, although we strongly

advocate for the former, at least if taxa outside the Maniraptoriformes are concerned.

Importantly, however, even if we had robust evidence for high neuron counts in Mesozoic

dinosaurs, this would by no means automatically suggest exceptional cognitive capacities.

General discussion - implications for neuron count and brain size estimates for

vertebrate paleontology

1) Are neuron counts good predictors of cognitive performance?

To infer cognitive abilities in extinct animals from neuron count estimates for the

whole brain or pallium, we first need to be assured that this measure can give us meaningful

insight into behaviors of extant animals. However, while there is some evidence for effects of

pallial neuron counts on species-level cognitive performance in primates (Deaner 2007 - but

see below) and birds (impulse inhibition - Herculano-Houzel, 2017; but see Kabadayi et al.

2017 for conflicting evidence; foraging-related innovativeness – Sol et al., 2022; but consider

limitations on how innovativeness is measured – Logan et al., 2018), the available data do

not provide consistent support for the hypothesis (Barron & Mourmourakis, 2023). As an

example, Güntürkün et al. (2017) reviewed the performance of domestic pigeons (*Columba*

livia), corvids and anthropoid primates in a number of cognitive tasks with the aim to

determine if a "cognitive hierarchy" between the three groups exists. They note that pallial

neuron counts in corvids are about 2–6 times lower than in large-bodied monkeys and apes

but 6–17 times higher than in pigeons. Thus, one would predict major increases in cognitive

capacities from pigeons to corvids to anthropoids. Yet, corvids typically perform on par with

anthropoid primates (see also Kabadayi et al. 2016 and Pika et al. 2020), and pigeons do so

as well in some cognitive dimensions, such as numerical competence and short-term

memory (Güntürkün et al., 2017). In addition, standardized testing of various primate species

suggests that small-brained lemurs with comparatively low neuronal densities (Kverková et

al., 2022), monkeys and great apes rival each other in a number of cognitive dimensions

(Schmitt et al., 2012; Fichtel et al., 2020). In fact, findings that report the influence of

absolute brain size (and thus neuron number) on cognitive performance in primates (Deaner

et al. 2007) have failed replication so far (Fichtel et al. 2020). As a final example, we want to

point out large-bodied dolphin species, which have remarkably high neocortical neuron

counts (*Globicephala melas* - 37 bn, *Orcinus orca* - 43 bn; Ridgway et al., 2019). Although

neuron numbers in these animals vastly exceed those of humans (15-20 bn), there is no

evidence that cetacean cognition is on par or even superior to that of our species (e.g.,

Manger, 2013; Güntürkün, 2014). Hence, even immense differences in telencephalic neuron

numbers do not necessarily create cognitive divides and their value in predicting cognitive

performance is remarkably limited.

The case becomes even more untenable when we take more complex cognitive

phenomena into consideration, such as tool use. Remarkably, Herculano-Houzel (2022)

suggested that this might be within the realm of possibility for large theropods such as *T. rex*,

as it is for primates and telluravian birds. However, tool use even within these groups is rare,

especially if the more rigorous definition of "tooling" (requiring the deliberate management of

a mechanical interface, see Fragaszy & Mangalam, 2018) is employed: this occurs in only 9

avian and 20 primate genera (Colbourne et al., 2021). While it is true that telencephalon size

in birds has an association with tool use (Lefebvre et al., 2002), this correlation does not hold

any predictive power in the sense that all birds with a certain-sized telencephalon exhibit tool

use. Even within corvids, which telencephalic neuron counts and sophisticated cognitive

abilities overlap with those of anthropoid primates (Olkowicz et al., 2016; Ströckens et al.,

2022), New Caledonian crows (*Corvus moneduloides*), and Hawaiian 'alalā crows (*C.

hawaiiensis*) are the only species known to use and manufacture tools in the wild. Notably,

both species inhabit remote islands, and they share unusually straight beaks and greater

binocular overlap than other crows, which are thought to be specific morphological

adaptations to enable tool use (Rutz et al., 2016; Troscianko et al., 2012). A similar situation

can be observed in parrots. These birds display the highest avian telencephalic neuron

counts (Olkowicz et al., 2016; Kverková et al., 2022; Ströckens et al., 2022), and a greatly

enlarged medial spiriform nucleus, which acts as an interface between the pallium and the

cerebellum, enabling enhanced motor cognition (Gutiérrez-Ibáñez et al., 2018). However,

the Tanimbar corella (*Cacatua goffiniana*) is the only parrot known to be a sophisticated tool

user in the wild (O'Hara et al., 2021); tellingly, the Tanimbar corella also inhabits an isolated

Indonesian archipelago. Cases like these indicate that while there is a chance that a gross

neuronal threshold must be met for such sophisticated vertebrate tool use to evolve (a notion

we would reject since ants evolved remarkable tool use skills with brains that are small and

few in neurons even for the standard of hymenopteran insects - Godfrey et al., 2021), it is

highly unlikely to happen without sufficient ecological pressure, and the differences between

tool using and non-tool using species are likely too subtle to detect via measurement of

neuronal quantities.

Considering these findings, it is unsurprising that taxa converging in neuronal counts

often differ markedly in cognition and behavior. Herculano-Houzel (2022) ranked her

neuronal count estimates for large theropods against those of anthropoid primates, she

might as well have done so for giraffes (1.7 bn neurons), which exceed tool-proficient

capuchins (1.1 bn) and corvids (0.4-1.2 bn) in telencephalic neuron numbers, rivaling

macaques (0.8 - 1.7 bn) (Olkowicz et al., 2016). We know little about giraffes' cognitive

abilities (Caicoya et al., 2018), but it would be appropriate to be skeptical of any claim that

they might exhibit "macaque-like" cognition based simply on that measure. Too many other

biological traits divide these taxa, perhaps most strikingly body size. While we agree with

many contemporary authors that relative brain size per se is a flawed measure of cognitive

complexity (e.g., Van Schaik et al., 2021), it must not be totally ignored. This is especially

true if comparisons between primates and non-bird dinosaurs are drawn, since the species

concerned may differ in body mass by several orders of magnitude. Contrary to the

assumptions of Herculano-Houzel (2022), the size of the telencephalon and number of its

neurons must be related to the dimensions of the body, because it processes sensory,

visceral, and motoric information, which scale with body size (Chittka & Niven, 2009; Van

Schaik et al., 2021). Relative brain size and body size are thus not negligible variables in

comparative cognition and need to be considered in paleoneurology. This fact is clearly

reflected by the pronounced intra- as well as interspecific body size-dependent scaling of

brain size in vertebrates (Tsuboi et al., 2018; Ksepka et al., 2020; Van Schaik et al., 2021;

Bertrand et al., 2022), which can hardly be explained otherwise.

The confounding factor of body size on neurological measures might be mitigated by

calculating clade-specific portions of telencephalic mass dedicated to somatic functions

based on intraspecific variation (Triki et al., 2021; Van Schaik et al., 2021) or by focusing on

neuron counts in brain regions that are evidently not involved in somatic processing

(Herculano-Houzel, 2017; Logan et al., 2018). In fact, a number of studies, particularly in

birds, were able to associate intraspecific differences in certain cognitive dimensions to

localized neurological variation, making this approach a promising one (discussed by Logan

et al., 2018). At the same time however, the great intra- and interspecific heterogeneity in

brain tissue architecture and neurochemistry enormously complicates any interspecific

extrapolations (Logan et al., 2018; Barron & Mourmourakis, 2023). Thus, researchers cannot

translate these findings to extinct species with any tolerable degree of certainty. This issue is

of special relevance when comparing sauropsids with mammals. The mammalian forebrain

exhibits a layered cortex but the pallium of extant sauropsids (and thus likely Mesozoic

dinosaurs) is largely nuclear in organization. As the forebrain increases in size and neuron

counts, a cortical organization can reduce axon length (and therefore processing time and

energetic demands) by folding and bringing adjacent areas closer together, something that is

impossible in a nuclear organization (see Reiner, 2023 for an extensive review).

Neuron counts corresponding to major brain regions, whether empirically determined

or estimated, dramatically simplify neuronal tissue complexity, as do measures such as

absolute brain size or EQ. Based on current evidence, they also represent flawed cognitive

proxies that need to be viewed in the broader context of an animal's ecology, neuroanatomy,

connectomics, and neurochemistry (Fields & Stevens-Graham, 2002; Eyal et al., 2016;

Logan et al., 2018; Barron & Mourmourakis, 2023, Reiner, 2023). All in all, we want to

discourage attempts to predict cognitive performance in extinct species based on endocast-

derived neuron count estimates.

2) Inferring metabolic rate

Apart from inferences about cognition, Herculano-Houzel (2022) claims that relative

brain size should be established as a new thermobiological indicator in vertebrate

palaeontology: Relatively large brains, as inferred for theropods, should be viewed as

indicators of endothermy, while smaller ones, as are found in pterosaurs, sauropodomorphs

and many ornithischians, would indicate ectothermy. Whereas overall brain size in

vertebrates is indeed correlated with metabolic rate (e.g., Yu et al., 2014 - but also note the

extreme variability within ecto- as well as endothermic groups), Herculano-Houzel's (2022)

approach simplifies the matter and ignores a vast body of already available evidence on

dinosaur thermobiology. First, as we have extensively discussed here, relative brain size in

large theropods was markedly smaller than suggested by Herculano-Houzel (2022),

rendering all conclusions made for this group untenable. Second, it is important to point out

that there is a spectrum of metabolic rates in vertebrates (Legendre & Davesne, 2020) rather

than a dichotomy, as suggested by Herculano-Houzel (2022).

Where exactly certain ornithodiran taxa align within this spectrum continues to be

debated, but there is consensus that dinosaurs and pterosaurs, despite their in parts

iconically small brains, had metabolic rates well above the range of extant ectothermic

sauropsids (see references below). Rather than emerging with theropods, contemporary

evidence suggests that endothermy evolved in the ornithodiran stem-lineage or even earlier

(Legendre et al., 2016; Grigg et al., 2022) and hence was inherited by pterosaurs and

dinosaurs. The extensive data supporting the presence of endothermy across Ornithodira

has recently been reviewed by Grigg et al. (2022) and includes the presence of hair-like,

sometimes branched, integumentary structures (Benton et al., 2019; Campione et al., 2020),

the efficiency of the ornithodiran respiratory system (Wedel, 2006; Butler et al., 2009;

Aureliano et al., 2022; Wang et al., 2023), bone histology and high skeletal growth rates (de

Ricqlès et al., 2000; Padian et al. 2004; Prondvai et al., 2012; Redelstroff et al. 2013;

Legendre et al., 2016), paleoenvironmental data (Druckenmiller et al., 2021), models of

locomotor costs (Pontzer et al., 2009) and geochemically-derived thermometry (Barrick et

al., 1996; Dawson et al. 2020; Wiemann et al., 2022). Nevertheless, osteohistological

evidence suggests that both theropod and non-theropod ornithodiran taxa varied in their

growth and associated metabolic rates (Jenkins et al., 2001; Erickson et al., 2009;

Redelstroff et al. 2013; D'Emic et al., 2023) and a secondary reduction of metabolic rate in

some ornithischian groups is plausible (Padian et al., 2004; Redelstroff and Sander, 2009;

Wiemann et al., 2022), albeit still compatible with an endothermic physiology (Grigg et al.,

2022). Overall, we emphasize the need for a nuanced perspective on this trait; we cannot

simply rely on modern birds and squamates as analogs.

The assumption that relative brain size alone (even if inferred correctly) can

outperform all the aforementioned thermophysiological predictors to infer endothermy

appears at best improbable. Its utility to gauge metabolic rate across ornithodiran groups

therefore remains highly doubtful and must be viewed in the framework of other, more robust

lines of evidence.

3) Inferring life history traits

Finally, Herculano-Houzel (2022) suggests that neuron count estimates can be used

to model life history traits in Mesozoic ornithodiran taxa. This notion is based on previous

empirical work that showed an association between pallial neuron counts and selected

ontogenetic variables in extant mammals and birds (Herculano-Houzel, 2019). Applied to *T.*

rex, the respective equations predict that females reached sexual maturity at an age of 4–5

years and that the longevity of the species was 42-49 years (Herculano-Houzel, 2022).

These calculations rest on the assumption that *T. rex* had to have 2.2 – 3.3 billion

pallial neurons. As we have shown, this premise appears exceedingly unlikely. Furthermore,

the aforementioned life history predictions are contradicted by the fossil evidence: Sexual

maturity in extinct nonavian dinosaurs can be estimated histologically by the presence of

medullary bone, a tissue that forms as a calcium reservoir for egg shell production and which

is also seen in female birds (Schweitzer et al., 2005; Woodward et al., 2020). The earliest

estimate of sexual maturity in *T. rex*, as estimated by the presence of medullary bone, is 15

years (Woodward et al., 2020; Carr, 2020). If the life history of *T. rex* was similar to extant

American alligators where sexual maturity occurs in animals that attain half of adult size

(which would be in line with the available fossil data - Carr, 2020), then the earliest onset of

sexual maturity in *T. rex* happened in its 12th year of life. Based on these lines of evidence,

Herculano-Houzel's (2022) method greatly underestimates the onset of sexual maturity by 8

to 11 years. Based on the number of lines of arrested growth in its long bones, which are

thought to indicate annual cessations of growth, the chronologically oldest *T. rex* sampled so

far lived up to 33 years (Cullen et al., 2020). Although it is not unreasonable to assume that

T. rex lived longer than three decades, there is yet no histological evidence to support that

hypothesis. However, given that Herculano-Houzel's (2022) estimate is based on faulty

premises, it should not be considered a plausible alternative.

In fact, if applied to species other than *T. rex*, the limitations of the aforementioned

approach become even more visible. For instance, if the life history of the sauropod

Apatosaurus, a gigantic dinosaur with an adult weight exceeding 30 tonnes, is modeled

based on our own neuron count estimates derived from an avian regression and an

endocranial fill of 42%, the equations suggest a longevity of only 24.5 years and an onset of

sexual maturity at 2.2 years (note that assuming a non-avian sauropsid regression or smaller

brain size would result in an even more fast-paced life history prediction). These figures are

obviously unfeasible. We are aware that Herculano-Houzel (2022) assumes that sauropods

such as *Apatosaurus* were ectothermic animals so that the given equations could not be

applied to them. However, since this notion defies essentially all available evidence on the

biology of sauropods (see above), we choose to ignore it here. Perhaps unsurprisingly,

correlations between life history and neurology that were established from a limited number

of extant species of mammals and birds cannot be used to infer ontogenetic parameters

across non-avian dinosaurs. We strongly discourage relying on them in palaeontological

practice.

Beyond endocasts: What are the limits of inference on dinosaur cognition?

If neuron count estimates and other endocast-derived variables do not allow reliable

predictions about the cognitive abilities of non-avian dinosaurs to be made, what other

methods are available? First of all, trace fossils can provide direct evidence on how

dinosaurs exploited their environment and interacted with both con- and heterospecifics

(e.g., Carpenter et al. 2005; Varricchio et al. 2007; Lockley et al. 2016; Brown et al. 2021).

While trace fossils can provide precise and diverse insights into dinosaur behavior, obvious

limitations render perspectives gained from them extremely patchy, nonetheless.

One further way of inferring cognitive traits in dinosaurs is by comparatively studying

relevant behavioral phenomena in living crocodylians and birds, the groups that form their

extant phylogenetic bracket. While such approaches are starting to gain pace (Zeiträg et al.,

2023), we are not aware that ethological research could so far identify shared physical or

social cognitive skills in crocodylians and birds that have not also been found in turtles and

squamates (in case comparative data is indeed available - Zeiträg et al., 2022; Font et al.,

2023). Thus, the behavioral resolution of such approaches appears limited thus far.

Cognitive traits found exclusively in birds or crocodylians cannot simply be extrapolated to

Mesozoic dinosaurs with any degree of certainty since they might be crown group

apomorphies. Whereas it might be appealing to hypothesize that cognitive patterns found

among modern palaeognaths are representative for their maniraptoriform forerunners

(Jensen et al., 2023; Zeiträg et al., 2023), this idea is (in most cases) not testable and should

hence not be disseminated uncritically.

Inferences on dinosaur cognition are hindered by the fact that both extant

crocodylians and birds are highly derived in their own ways: Convergently to mammals, birds

have not only evolved an enlarged forebrain and cerebellum, but also extensive connections

between these two brain regions (Gutierrez-Ibanez et al., 2018) and descending projections

from the pallium to the brainstem and/or spinal cord (Ulinski and & Margoliash, 1990; Reiner

& Medina, 2000). These circuits are likely essential for enabling various avian behaviors,

including tooling, but are not present in extant non avian sauropsids (Ulinski and &

Margoliash 1990; Gutierrez-Ibanez et al., 2023). It remains obscure when they evolved.

Crocodylians on the other hand conserve a plesiomorphic brain morphology and cerebral

tissue organization (Briscoe et al., 2018; Briscoe & Ragsdale, 2018). Interestingly, the rather

sparse available data suggest that they exhibit a pronounced apomorphic decrease in

neuronal density (Kverková et al., 2022). They are also unusual in being secondary

ectotherms (e.g., Seymour et al., 2004; Legendre et al., 2016; Botha et al., 2023). It is

unclear how this might have affected their neurology and cognition. Thus, the extant

archosaurian groups leave us in a rather suboptimal position to infer cognitive traits in non-

avian dinosaurs.

Obviously, even the absence of a given cognitive trait in crocodylians and basal

extant birds like palaeognaths does not refute its existence in non-avian dinosaurs,

considering the diversity and long evolutionary history of this group. In fact, a species'

ecology is typically more indicative of certain behaviors and associated cognitive

phenomena than its phylogenetic affinities. Returning to the example of tool use, tool use is

an adaptation typically found in omnivorous extractive foragers (Parker & Gibson, 1977;

Parker, 2015) and only rarely reported in predators (Shumaker et al., 2011). This is reflected

by the fact that the most common types of tooling actions that have evolved comprise

reaching, probing or pounding, usually in order to access food (Colbourne et al., 2021). It

has long been observed that tool use appears when a species is found in an uncharacteristic

niche, for which it lacks the appropriate morphological adaptations, and thus compensates

by using tools to generate a functionally equivalent behavior (Alcock, 1972; Parker & Gibson,

1977). This is likely why a number of birds that use tools are found on islands, yet the ability

appears absent in their close mainland relatives (Rutz et al., 2016). Simply put, in order for

tool use to evolve, there needs to be a reason for it to evolve, and there are very few

ecological contexts where tool use is a superior adaptation to its morphological equivalent

(Hansell & Ruxton, 2008). Unfortunately, this type of extremely specific contextual

information is nearly absent in long extinct species. From its iconic tooth and jaw

morphology, one can confidently predict that a hypercarnivorous species like *T. rex* would

have no need for tools, but the problem remains that few assumptions about extinct animal

cognition are falsifiable.

In sum, reconstructing cognition in dinosaurs and other fossil taxa without close living

analogs is a challenging endeavor that requires integrative approaches if we are to provide

compelling inferences (de Sousa et al., 2023). Bare neuronal count estimates might be

considered a rather minor contribution to this effort and need to be aligned with data from

comparative anatomy and neurology, ecology, trace fossils, and comparative behavioral

studies on extant animals to offer a plausible picture of cognition. While communicating such

findings, researchers should acknowledge the limitations of the presented inferences to

allow their audience to delineate between reasoned conclusions and pure speculation. In a

field such as dinosaur research - avidly followed by popular media and the public eye - such

a nuanced view appears especially warranted.

1033

Conclusions

The dinosaurian neuronal count and relative brain size estimates presented by

Herculano-Houzel (2022) are inaccurate due to methodological shortcomings, in particular

for *T. rex*. Accordingly, the biological inferences drawn from them are implausible. As we

show here, there is no compelling evidence that relative brain size in large-bodied theropods

differed significantly from that of extant non-avian sauropsids, and their telencephalic neuron

counts were likely not exceptional, especially for animals of their size. Furthermore, we

highlight issues associated with neuron count estimates in vertebrate paleontology and

argue against their use in reconstructing behavioral and life history variables. The latter is

especially true for animals such as non-avian dinosaurs, for which disparate neuron

densities might be hypothesized based on different phylogenetic and morphological

arguments.

For obvious reasons, many inferences we might make about Mesozoic dinosaur

behavior will remain limited. Nevertheless, we can justify certain predictions - to a degree -

within integrative empirical frameworks to which neuron count estimates might well be added

in the future. Before such steps can be taken, however, a substantially improved

understanding of the relationship between neuron counts and other biological variables,

especially cognitive performance, in extant animals is required.

1052

Institutional abbreviations

AMNH = American Museum of Natural History, New York City, New York, United States;

BMNH / NHMUK = Natural History Museum, London, UK; BSP = Bayerische

Staatssammlung für Paläontologie und Historische Geologie, Munich, Germany; BYU =

Brigham Young University, Earth Science Museum, Provo, Utah, United States; CM =

Carnegie Museum of Natural History, Pittsburgh, Pennsylvania. CMN = Canadian Museum

of Nature, Ottawa, Ontario, Canada. DINO = Dinosaur National Monument, Jensen, Utah,

United States. FIP - Florida Institute of Paleontology, Palm Beach, Florida, United States;

FMNH = Field Museum of Natural History, Chicago, Illinois, United States; HMN / MB.R. =

Museum für Naturkunde, Berlin, Germany; IGM = Mongolian Institute of Geology, Ulaan

Bator, Mongolia; IRSNB / RBINS = Institut Royal des Sciences Naturelles de Belgique,

Brussels, Belgium; IVPP = Institute of Vertebrate Paleontology and Paleoanthropology,

Beijing, China; KUVP = Kansas University Natural History Museum, Lawrence, Kansas,

United States; MACN = Museo Argentino de Ciencias Naturales "Bernardino Rivadavia",

Buenos Aires, Argentina; MPC-D = Institute of Paleontology and Geology, Mongolian

Academy of Sciences, Ulaan Bator, Mongolia; MUCPv-CH = Museo de la Universidad

Nacional del Comahue, colección del Museo Ernesto Bachmann, Villa El Chocón,

Argentina; MOR = Museum of the Rockies, Bozeman, Montana, United States; NMC =

Canadian Museum of Nature, Ottawa, Canada; NCSM = North Carolina Museum of

Natural Sciences, Raleigh, North Carolina, United States; OMNH = Sam Noble Museum at

the University of Oklahoma, Norman, Oklahoma, United States; PIN = Paleontological

Institute, Russian Academy of Sciences, Moscow, Russia; ROM = Royal Ontario Museum,

Toronto, Ontario, Canada; RTMP/TMP = Royal Tyrrell Museum of Palaeontology,

Drumheller, Alberta, Canada; SGM = Ministere de l'Energie et des Mines, Rabat, Morocco;

USNM = Smithsonian National Museum of Natural History, Washington, D.C., United

States; UUVP = University of Utah, Salt Lake City, Utah, United States; YPM = Yale

Peabody Museum, New Haven Connecticut, United States.

1080

1081   **Acknowledgements**

1082   We want to thank David Burnham, Ariana Paulina-Carabajal, Gregory M. Erickson and

1083   Lawrence M. Witmer for sharing valuable information on fossil specimens. Andrew N.

1084   Iwaniuk is acknowledged for helpful discussions on the methodology and structure of the

1085   manuscript.

1086

1087   **References**

1088   Alcock, J. (1972). The evolution of the use of tools by feeding animals. *Evolution*, *26*(3), 464-
1089   473.

1090

1091   Alonso, P. D., Milner, A. C., Ketcham, R. A., Cookson, M. J., & Rowe, T. B. (2004). The
1092   avian nature of the brain and inner ear of *Archaeopteryx*. *Nature*, *430*(7000), 666-669.

1093

1094   Balanoff, A. M., Bever, G. S., & Ikejiri, T. (2010). The braincase of *Apatosaurus* (Dinosauria:
1095   Sauropoda) based on computed tomography of a new specimen with comments on variation
1096   and evolution in sauropod neuroanatomy. *American Museum Novitates*, *2010*(3677), 1-32.

1097

1098   Balanoff, A. M., Bever, G. S., Rowe, T. B., & Norell, M. A. (2013). Evolutionary origins of the
1099   avian brain. *Nature*, *501*(7465), 93-96.

1100

1101 Barrick, R. E., Showers, W. J., & Fischer, A. G. (1996). Comparison of thermoregulation of
1102 four ornithischian dinosaurs and a varanid lizard from the Cretaceous Two Medicine
1103 Formation: evidence from oxygen isotopes. *Palaios, 11*, 295-305.
1104
1105 Barron, A. B., & Mourmourakis, F. (2023). The relationship between cognition and brain size
1106 or neuron number. *Brain, Behavior and Evolution.* https://doi.org/10.1159/000532013
1107
1108 Bennett, S. C. (1995). A statistical study of *Rhamphorhynchus* from the Solnhofen Limestone
1109 of Germany: year-classes of a single large species. *Journal of Paleontology, 69*(3), 569-580.
1110
1111 Benton, M.J., Dhouailly, D., Jiang, B., & McNamara, M. (2019). The early origin of feathers.
1112 *Trends in Ecology and Evolution. 34*(9), 856-869.
1113
1114 Benson, R. B. J., Hunt, G., Carrano, M. T. & Campione, N. (2017). Data from: Cope's rule and
1115 the adaptive landscape of dinosaur body size evolution. *Dryad Digital Repository.*
1116 *https://doi.org/10.5061/dryad.1t3r4*
1117
1118 Benson, R.B., Campione, N. E., Carrano, M. T., Mannion, P. D., Sullivan, C., Upchurch, P. &
1119 Evans, D. C. (2014). Rates of dinosaur body mass evolution indicate 170 million years of
1120 sustained ecological innovation on the avian stem lineage. *PLoS Biology*, 12(5), p.e1001853.
1121
1122 Benson, R.B., Hunt, G., Carrano, M. T. & Campione, N. (2018). Cope's rule and the adaptive
1123 landscape of dinosaur body size evolution. *Palaeontology, 61*(1), 13-48.
1124
1125 Bertrand, O. C., Shelley, S. L., Williamson, T. E., Wible, J. R., Chester, S. G., Flynn, J. J., ...
1126 & Brusatte, S. L. (2022). Brawn before brains in placental mammals after the end-Cretaceous
1127 extinction. *Science, 376*(6588), 80-85.
1128
1129 Bever, G. S., Brusatte, S. L., Carr, T. D., Xu, X., Balanoff, A. M., & Norell, M. A. (2013). The
1130 braincase anatomy of the late Cretaceous dinosaur *Alioramus* (Theropoda:
1131 Tyrannosauroidea). *Bulletin of the American Museum of Natural History, 2013*(376)**,** 1-72.
1132
1133 Botha, J., Weiss, B. M., Dollman, K., Barrett, P. M., Benson, R. B., & Choiniere, J. N. (2023).
1134 Origins of slow growth on the crocodilian stem lineage. *Current Biology, 33*(19), 4261-4268.
1135

Briscoe, S. D., & Ragsdale, C. W. (2018). Homology, neocortex, and the evolution of
developmental mechanisms. *Science*, *362*(6411), 190-193.

Briscoe, S. D., Albertin, C. B., Rowell, J. J., & Ragsdale, C. W. (2018). Neocortical
association cell types in the forebrain of birds and alligators. *Current Biology*, *28*(5), 686-696.

Brochu, C. A. (2000). A digitally-rendered endocast for *Tyrannosaurus rex*. *Journal of*
Vertebrate Paleontology, 20(1), 1-6.

Brown, C.M., Currie, P.J., & Therrien, F. (2021) Intraspecific facial bite marks in tyrannosaurids
provide insight into sexual maturity and evolution of bird-like intersexual display. *Paleobiology*
48(1), 12-43.

Brusatte, S. L., Carr, T. D., Erickson, G. M., Bever, G. S., & Norell, M. A. (2009). A long-
snouted, multihorned tyrannosaurid from the Late Cretaceous of Mongolia. *Proceedings of the*
National Academy of Sciences, 106(41), 17261-17266.

Buchholtz, E. A., & Seyfarth, E.-A. (2001.) The study of "fossil brains": Tilly Edinger (1897–
1967) and the beginnings of paleoneurology. *BioScience*, 51(8), 674-682.

Burnham, D. A. (2004). New information on *Bambiraptor feinbergi* (Theropoda:
Dromaeosauridae) from the Late Cretaceous of Montana. In P. J. Currie, E. B. Koppelhus, M.
A. Shugar & J. L. Wright (eds.), *Feathered dragons: studies on the transition from dinosaurs*
to birds (pp. 67-111). Indiana University Press, Bloomington.
Butler, R. J., Barrett, P. M., & Gower, D. J. (2009). Postcranial skeletal pneumaticity and air-
sacs in the earliest pterosaurs. *Biology Letters*, *5*(4), 557-560.

Caicoya, Á. L., Amici, F., Ensenyat, C., & Colell, M. (2019). Object permanence in *Giraffa*
camelopardalis: First steps in giraffes' physical cognition. *Journal of Comparative*
Psychology, *133*(2), 207.

Campione, N. E. (2020) MASSTIMATE: Body Mass Estimation Equations for Vertebrates.
Available at: https://cran.r-project.org/web/packages/MASSTIMATE/MASSTIMATE.pdf

Campione, N. E., & Evans, D. C. (2020). The accuracy and precision of body mass estimation
in non-avian dinosaurs. *Biological Reviews*, *95*(6), 1759-1797.

Campione, N. E., Barrett, P. M., & Evans, D. C. (2020). On the ancestry of feathers in Mesozoic dinosaurs. In C. Foth & O. W. M. Rauhut (eds.), *The Evolution of Feathers: From Their Origin to the Present* (pp. 213-243). Springer International Publishing.

Carpenter, K., Sanders, F., McWhinney, L.A., & Wood, L. (2005). Evidence for predator-prey relationships: examples for *Allosaurus* and *Stegosaurus*. In K. Carpenter (ed.), *The Carnivorous Dinosaurs* (pp. 325-350). Indiana University Press, Bloomington.

Carr, T. D. (2020). A high-resolution growth series of *Tyrannosaurus rex* obtained from multiple lines of evidence. *PeerJ*, *8*, e9192.

Cerroni, M. A., & Paulina-Carabajal, A. (2019). Novel information on the endocranial morphology of the abelisaurid theropod *Carnotaurus sastrei*. *Comptes Rendus Palevol*, *18*(8), 985-995.

Chentanez, T., Huggins, S. E., & Chentanez, V. (1983). Allometric relationships for the Siamese crocodile, *Crocodylus siamensis*. *Journal of The Science Society of Thailand*, *9*, 5–26.

Chittka, L., & Niven, J. (2009). Are bigger brains better?. *Current Biology*, *19*(21), R995-R1008.

Colbourne, J. A. D., Auersperg, A. M. I., Lambert, M. L., Huber, L., & Völter, C. J. (2021). Extending the reach of tooling theory: a neurocognitive and phylogenetic perspective. *Topics in Cognitive Science*, *13*(4), 548–572.

Cullen, T. M., Canale, J. I., Apesteguía, S., Smith, N. D., Hu, D., & Makovicky, P. J. (2020). Osteohistological analyses reveal diverse strategies of theropod dinosaur body-size evolution. *Proceedings of the Royal Society B*, *287*(1939), 20202258.

Currie, P. J., & Zhao, X. J. (1993). A new troodontid (Dinosauria, Theropoda) braincase from the dinosaur park formation (Campanian) of Alberta. *Canadian Journal of Earth Sciences*, 30(10), 2231-2247.

Cuvier, G. (1812). Recherches sur les ossemens fossiles de quadrupèdes, Vol. 3. Deterville, Paris.

1210 Dawson, R. R., Field, D. J., Hull, P. M., Zelenitsky, D. K., Therrien, F., & Affek, H. P. (2020).
1211 Eggshell geochemistry reveals ancestral metabolic thermoregulation in Dinosauria. *Science*
1212 *Advances*, *6*(7), eaax9361.
1213
1214 Deaner, R. O., Isler, K., Burkart, J., & Van Schaik, C. (2007). Overall brain size, and not
1215 encephalization quotient, best predicts cognitive ability across non-human primates. *Brain,*
1216 *Behavior and Evolution*, *70*(2), 115-124.
1217
1218 D'Emic, M. D., O'Connor, P. M., Sombathy, R. S., Cerda, I., Pascucci, T. R., Varricchio, D.,
1219 ... & Curry Rogers, K. A. (2023). Developmental strategies underlying gigantism and
1220 miniaturization in non-avialan theropod dinosaurs. *Science*, *379*(6634), 811-814.
1221
1222 de Ricqlès, A., Padian, K., Horner, J.R., & Francillon-Vieillot, H. 2000. Palaeohistology of the
1223 bones of pterosaurs (Reptilia: Archosauria): anatomy, ontogeny, and biomechanical
1224 implications. *Zoological Journal of the Linnean Society* 129: 349-385.
1225
1226 de Sousa, A. A., Beaudet, A., Calvey, T., Bardo, A., Benoit, J., Charvet, C. J., ... & Wei, Y.
1227 (2023). From fossils to mind. *Communications Biology*, *6*(1), 636.
1228
1229 Druckenmiller, P. S., Erickson, G. M., Brinkman, D., Brown, C. M., & Eberle, J. J. (2021).
1230 Nesting at extreme polar latitudes by non-avian dinosaurs. *Current Biology*, *31*(16), 3469-
1231 3478.
1232
1233 Edinger T. (1929) *Die fossilen Gehirne.* J. Springer, Berlin.
1234
1235 Erickson, G.M., Rauhut, O.W.M., Zhou, Z., Turner, A.H., Inouye, B.D., Hu, D., and Norell,
1236 M.A. (2009). Was dinosaurian physiology inherited by birds? reconciling slow growth in
1237 *Archaeopteryx. PLoS ONE* 4(10): e7390.
1238
1239 Evans, D. C., Ridgely, R., & Witmer, L. M. (2009). Endocranial anatomy of lambeosaurine
1240 hadrosaurids (Dinosauria: Ornithischia): a sensorineural perspective on cranial crest
1241 function. *The Anatomical Record*, 292(9), 1315-1337.
1242
1243 Eyal, G., Verhoog, M. B., Testa-Silva, G., Deitcher, Y., Lodder, J. C., Benavides-Piccione,
1244 R., ... & Segev, I. (2016). Unique membrane properties and enhanced signal processing in
1245 human neocortical neurons. *Elife*, *5*, e16553.
1246

1247   Fabbri, M., & Bhullar, B. A. S. (2022). The endocast of *Euparkeria* sheds light on the
1248   ancestral archosaur nervous system. *Palaeontology*, *65*(6), e12630.
1249

1250   Fragaszy, D. M., & Mangalam, M. (2018). Tooling. Advances in the Study of Behavior,
1251   2018(50), 177–241.
1252

1253   Franzosa, J., & Rowe, T. (2005). Cranial endocast of the Cretaceous theropod dinosaur
1254   *Acrocanthosaurus atokensis*. *Journal of Vertebrate Paleontology*, *25*(4), 859-864.
1255

1256   Fields, R. D., & Stevens-Graham, B. (2002). New insights into neuron-glia communication.
1257   *Science*, *298*(5593), 556-562.
1258

1259   Font, E., García-Roa, R., Pincheira-Donoso, D., & Carazo, P. (2019). Rethinking the effects
1260   of body size on the study of brain size evolution. *Brain Behavior and Evolution*, *93*(4), 182-
1261   195.
1262

1263   Font, E., Burghardt, G. M., & Leal, M. (2023). Brains, behaviour, and cognition: multiple
1264   misconceptions. In C. Warwick, P. C. Arena, & G. M. Burghardt (eds.), *Health and Welfare of*
1265   *Captive Reptiles* (pp. 211-238). Springer International Publishing, Cham.
1266

1267   Galton, P. M. (2001). Endocranial casts of the plated dinosaur *Stegosaurus* (Upper Jurassic,
1268   Western USA): a complete undistorted cast and the original specimens of Othniel Charles
1269   Marsh. In K. Carpenter (ed.), *The Armored Dinosaurs* (pp. 103–129). Indiana University
1270   Press, Bloomington.
1271

1272   Garland Jr, T., Harvey, P. H., & Ives, A. R. (1992). Procedures for the analysis of
1273   comparative data using phylogenetically independent contrasts. *Systematic Biology*, *41*(1),
1274   18-32.
1275

1276   Garland, Jr, T., & Ives, A. R. (2000). Using the past to predict the present: confidence
1277   intervals for regression equations in phylogenetic comparative methods. *The American*
1278   *Naturalist*, *155*(3), 346-364.
1279

1280   Gatesy, S. M. (1991). Hind limb scaling in birds and other theropods: implications for
1281   terrestrial locomotion. Journal of Morphology, 209(1), 83-96.
1282

1283 Giffin, E. B. (1989). Pachycephalosaur paleoneurology (Archosauria: Ornithischia). *Journal*
1284 *of Vertebrate Paleontology*, 9(1), 67-77.

1285

1286 Gignac, P.M. & Erickson, G.M. (2017). The biomechanics behind extreme osteophagy in
1287 *Tyrannosaurus rex. Scientific Reports* **7**(1), 2012.

1288

1289 Godfrey, R. K., Swartzlander, M., & Gronenberg, W. (2021). Allometric analysis of brain cell
1290 number in Hymenoptera suggests ant brains diverge from general trends. *Proceedings of*
1291 *the Royal Society B*, *288*(1947), 20210199.

1292

1293 Grigg, G., Nowack, J., Bicudo, J. E. P. W., Bal, N. C., Woodward, H. N., & Seymour, R. S.
1294 (2022). Whole-body endothermy: ancient, homologous and widespread among the
1295 ancestors of mammals, birds and crocodylians. *Biological Reviews*, *97*(2), 766-801.

1296

1297 Güntürkün, O. (2014). Is dolphin cognition special?. *Brain, Behavior and Evolution*, *83*(3),
1298 177-180.

1299

1300 Güntürkün, O., Ströckens, F., Scarf, D., & Colombo, M. (2017). Apes, feathered apes, and
1301 pigeons: differences and similarities. *Current Opinion in Behavioral Sciences*, *16*, 35-40.

1302

1303 Gutiérrez-Ibáñez, C., Iwaniuk, A. N., & Wylie, D. R. (2018). Parrots have evolved a primate-
1304 like telencephalic-midbrain-cerebellar circuit. *Scientific Reports*, *8*(1), 1-11

1305

1306 Gutiérrez-Ibáñez, C., Kettler, L., Pilon, M.C., Carr, C.E. & Wylie, D.R., (2023). Cerebellar
1307 inputs in the American alligator (*Alligator mississippiensis*). *Brain Behavior and Evolution*,
1308 98(1):.44-60.

1309

1310 Hackett, S. J., Kimball, R. T., Reddy, S., Bowie, R. C. K., Braun, E. L., Braun, M. J., … Yuri,
1311 T. (2008). A phylogenomic study of birds reveals their evolutionary history. *Science*,
1312 320(5884), 1763–1768.

1313

1314 Hansell, M., & Ruxton, G. (2008). Setting tool use within the context of animal construction
1315 behaviour. Trends in Ecology & Evolution, 23(2), 73–78.

1316

1317 Henderson, D. M. (2023). Growth constraints set an upper limit to theropod dinosaur body
1318 size. *The Science of Nature*, *110*(1), 4.

1319

1320   Herculano-Houzel, S. (2011). Brains matter, bodies maybe not: the case for examining

1321   neuron numbers irrespective of body size. *Annals of the New York Academy of Sciences*,

1322   *1225*(1), 191-199.

1323

1324   Herculano-Houzel, S. (2017). Numbers of neurons as biological correlates of cognitive

1325   capability. *Current Opinion in Behavioral Sciences*, 16(2017):1–7.

1326

1327   Herculano-Houzel, S. (2019). Longevity and sexual maturity vary across species with

1328   number of cortical neurons, and humans are no exception. *Journal of Comparative*

1329   *Neurology*, *527*(10), 1689-1705.

1330

1331   Herculano-Houzel, S. (2022). Theropod dinosaurs had primate-like numbers of telencephalic

1332   neurons. *Journal of Comparative Neurology*, *531*(9), 962-974.

1333

1334   Herculano-Houzel, S., & Kaas, J. H. (2011). Gorilla and orangutan brains conform to the

1335   primate cellular scaling rules: implications for human evolution. *Brain, Behavior and*

1336   *Evolution*, *77*(1), 33-44.

1337

Herculano-Houzel, S., Ribeiro, P., Campos, L., Valotta da Silva, A., Torres, L. B., Catania, K.

C., & Kaas, J. H. (2011). Updated neuronal scaling rules for the brains of glires

(Rodents/Lagomorphs). *Brain, Behavior and Evolution*, 78(4), 302-314.

Herculano-Houzel, S., Manger, P. R., & Kaas, J. H. (2014). Brain scaling in mammalian

evolution as a consequence of concerted and mosaic changes in numbers of neurons and

average neuronal cell size. *Frontiers in Neuroanatomy*, *8*, 77.

1344

1345   Herculano-Houzel, S., Catania, K., Manger, P. R., & Kaas, J. H. (2015). Mammalian brains

1346   are made of these: a dataset of the numbers and densities of neuronal and nonneuronal

1347   cells in the brain of glires, primates, scandentia, eulipotyphlans, afrotherians and

1348   artiodactyls, and their relationship with body mass. *Brain, behavior and evolution*, 86(3-4),

1349   145-163.

1350

1351   Hopson, J.A. (1979). Paleoneurology. In C. Gans, R. G. Northcutt, & P. Ulinski (eds.),

1352   *Biology of the Reptilia*, Volume 9 (pp. 39–146). Academic Press, New York.

1353

1354 Hothorn, T., Bretz, F., Westfall, P., Heiberger, R. M., Schuetzenmeister, A., & Scheibe, S.
1355 (2015). Package "multcomp". Available at: https://cran.r-
1356 project.org/web/packages/multcomp/multcomp.pdf
1357
1358 Hu, K., King, J. L., Romick, C. A., Dufeau, D. L., Witmer, L. M., Stubbs, T. L., ... & Benton,
1359 M. J. (2021). Ontogenetic endocranial shape change in alligators and ostriches and
1360 implications for the development of the non-avian dinosaur endocranium. *The Anatomical*
1361 *Record*, *304*(8), 1759-1775.
1362
1363 Hurlburt, G. R. (1996). *Relative brain size in recent and fossil amniotes: determination and*
1364 *interpretation.* PhD thesis, University of Toronto, Toronto, Canada (pp. 250).
1365
1366 Hurlburt, G.R., & Waldorf, L. (2002). Endocast volume and brain mass in a size series of
1367 alligators. *Journal of Vertebrate Paleontology*, *23*, (Supplement to 3), 69A..
1368
1369 Hurlburt, G. R., Ridgely, R. C., & Witmer, L. M. (2013) Relative size of brain and cerebrum in
1370 tyrannosaurid dinosaurs: an analysis using brain-endocast quantitative relationships in
1371 extant alligators. In J. M. Parrish, R. E. Molnar, P. J.Currie, & E. B. Koppelhus (eds.),
1372 *Tyrannosaurid Paleobiology* (pp. 134–154). Indiana University Press, Bloomington
1373
1374 Iwaniuk, A. N., & Nelson, J. E. (2002). Can endocranial volume be used as an estimate of
1375 brain size in birds? *Canadian Journal of Zoology,* 80(1), 16-23.
1376
1377 Jenkins Jr, F. A., Shubin, N. H., Gatesy, S. M., & Padian, K. E. V. I. N. (2001). A diminutive
1378 pterosaur (Pterosauria: Eudimorphodontidae) from the Greenlandic Triassic. *Bulletin of the*
1379 *Museum of Comparative Zoology*, *156*(1), 151-170.
1380
1381 Jensen, T. R., Zeiträg, C., & Osvath, M. (2023). The selfish preen: absence of allopreening in
1382 Palaeognathae and its socio-cognitive implications. *Animal Cognition*, 26, 1467–1476.
1383
1384 Jerison, H. J. (1973). *Evolution of the Brain and Intelligence.* Academic Press, New York.
1385
1386 Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K., & Mooers, A. O. (2012). The global
1387 diversity of birds in space and time. *Nature*, *491*(7424), 444-448.
1388

Jirak, D., & Janacek, J. (2017). Volume of the crocodilian brain and endocast during ontogeny. *PloS ONE*, *12*, 6, e0178491.

Kabadayi, C., Taylor, L. A., von Bayern, A. M., & Osvath, M. (2016). Ravens, New Caledonian crows and jackdaws parallel great apes in motor self-regulation despite smaller brains. *Royal Society Open Science*, *3*(4), 160104.

Kabadayi, C., Krasheninnikova, A., O'Neill, L., van de Weijer, J., Osvath, M., & von Bayern, A. M. (2017). Are parrots poor at motor self-regulation or is the cylinder task poor at measuring it?. *Animal Cognition*, *20*, 1137-1146.

Knoll, F., Lautenschlager, S., Kawabe, S., Martínez, G., Espílez, E., Mampel, L., & Alcalá, L. (2021). Palaeoneurology of the early cretaceous iguanodont Proa valdearinnoensis and its bearing on the parallel developments of cognitive abilities in theropod and ornithopod dinosaurs. *Journal of Comparative Neurology*, *529*(18), 3922-3945.

Knoll, F. & Schwarz-Wings, D. (2009) Palaeoneuroanatomy of *Brachiosaurus*. *Annales de Paléontologie*, *95*(3), 165–175.

Ksepka, D. T., Balanoff, A. M., Smith, N. A., Bever, G. S., Bhullar, B. A. S., Bourdon, E., ... & Smaers, J. B. (2020). Tempo and pattern of avian brain size evolution. *Current Biology*, *30*(11), 2026-2036.

Kumar, S., Stecher, G., Suleski, M., & Hedges, S. B. (2017). TimeTree: a resource for timelines, timetrees, and divergence times. *Molecular Biology and Evolution*, *34*(7), 1812-1819.

Kverková, K., Marhounová, L., Polonyiová, A., Kocourek, M., Zhang, Y., Olkowicz, S., ... & Němec, P. (2022). The evolution of brain neuron numbers in amniotes. *Proceedings of the National Academy of Sciences*, 119(11), e2121624119.

Larsson, H. C., Sereno, P. C., & Wilson, J. A. (2000). Forebrain enlargement among nonavian theropod dinosaurs. *Journal of Vertebrate Paleontology*, *20*(3), 615-618.

Legendre, L. J., & Davesne, D. (2020). The evolution of mechanisms involved in vertebrate endothermy. *Philosophical Transactions of the Royal Society B*, *375*(1793), 20190136.

1425 Legendre, L.J., Guénard, G., Botha-Brink, J., & Cubo, J. (2016). Palaeohistological evidence
1426 for ancestral high metabolic rate in archosaurs. *Systematic Biology* 65(6): 989-996.
1427
1428 Lauters, P., Coudyzer, W., Vercauteren, M., & Godefroit, P. (2012). The brain of *Iguanodon*
1429 and *Mantellisaurus*: perspectives on ornithopod evolution. In P. Godefroit (ed.), Bernissart
1430 dinosaurs and early Cretaceous terrestrial ecosystems (pp 213–224). Indiana University
1431 Press, Bloomington.
1432
1433 Lefebvre, L., Nicolakakis, N., & Boire, D. (2002). Tools and brains in birds. *Behaviour*,
1434 *139*(7), 939–973.
1435
1436 Lockley, M.G., McCrea, R.T., Buckley, L.G., Lim, J.D., Matthews, N.A., Breithaupt, B.H.,
1437 Houck, K.J., Gierliński, G.D., Surmik, D., Kim, K.S., Xing, L., Kong, D.Y., Cart, K., Martin, J.,
1438 & Hadden, G. (2016). Theropod courtship: large scale physical evidence of display arenas
1439 and avian-like scrape ceremony behaviour by Cretaceous dinosaurs. *Scientific Reports*
1440 6:18952
1441
1442 Logan, C. J., Avin, S., Boogert, N., Buskell, A., Cross, F. R., Currie, A., ... & Montgomery, S.
1443 H. (2018). Beyond brain size: Uncovering the neural correlates of behavioral and cognitive
1444 specialization. *Comparative Cognition & Behavior Reviews*.
1445
1446 Maddison, W. P. & Maddison, D. R. (2023). Mesquite: a modular system for evolutionary
1447 analysis. Available at: http://www.mesquiteproject.org
1448
1449 Manger, P. R. (2013). Questioning the interpretations of behavioral observations of
1450 cetaceans: is there really support for a special intellectual status for this mammalian order?.
1451 *Neuroscience*, *250*, 664-696.
1452
1453 Marsh, O. C. (1879). History and methods of palaeontological discovery. *American Journal*
1454 *of Science*, *3*(107), 323-359.
1455
1456 Medina, L. & Reiner, A. (2000). Do birds possess homologues of mammalian primary visual,
1457 somatosensory and motor cortices?. *Trends in Neurosciences*, 23(1): 1-12.
1458
1459 Morhardt, A. C. (2016). *Gross anatomical brain region approximation (GABRA): assessing*
1460 *brain size, structure, and evolution in extinct archosaurs*. PhD thesis, Ohio University, Ohio,
1461 United States.

1462

1463 Ngwenya, A., Patzke, N., Spocter, M. A., Kruger, J. L., Dell, L. A., Chawana, R., ... &

1464 Manger, P. R. (2013). The continuously growing central nervous system of the Nile crocodile

1465 (*Crocodylus niloticus*). *The Anatomical Record*, *296*(10), 1489-1500.

1466

1467 Ngwenya, A., Patzke, N., Manger, P. R., & Herculano-Houzel, S. (2016). Continued growth

1468 of the central nervous system without mandatory addition of neurons in the Nile crocodile

1469 (*Crocodylus niloticus*). *Brain Behavior and Evolution*, *87*(1), 19-38.

1470

1471 O'Hara, M., Mioduszewska, B., Mundry, R., Yohanna, Haryoko, T., Rachmatika, R.,

1472 Prawiradilaga, D. M., Huber, L., & Auersperg, A. M. I. (2021). Wild Goffin's cockatoos flexibly

1473 manufacture and use tool sets. *Current Biology*, *31*(20), 4512-4520.e6.

1474

1475 Olkowicz, S., Kocourek, M., Lučan, R. K., Porteš, M., Fitch, W. T., Herculano-Houzel, S., &

1476 Němec, P. (2016). Birds have primate-like numbers of neurons in the forebrain. *Proceedings*

1477 *of the National Academy of Sciences*, *113*(26), 7255-7260.

1478

1479 Osborn, H. F. (1912). Crania of *Tyrannosaurus* and *Allosaurus*. *Memoirs of the American*

1480 *Museum of Natural History, Volume. 1, New Series*, 1, 1-30.

1481

1482 Osmólska, H. (2004). Evidence on relation of brain to endocranial cavity in oviraptorid

1483 dinosaurs. *Acta Palaeontologica Polonica*, *49*(2), 321-324.

1484

1485 Padian, K., Horner, J.R., & De Ricqlès, A. 2004. Growth in small dinosaurs and pterosaurs:

1486 the evolution of the archosaurian growth strategies. *Journal of Vertebrate Paleontology* 24(3):

1487 555-571.

1488

1489 Pagel, M. (1999). Inferring the historical patterns of biological evolution. *Nature*, *401*(6756),

1490 877-884.

1491 Paradis, E., & Schliep, K. (2019). ape 5.0: an environment for modern phylogenetics and

1492 evolutionary analyses in R. *Bioinformatics*, *35*(3), 526-528.

1493

1494 Parker, S. T. (2015). Re-evaluating the extractive foraging hypothesis. *New Ideas in*

1495 *Psychology*, 37, 1–12.

1496

1497   Parker, S. T., & Gibson, K. R. (1977). Object manipulation, tool use and sensorimotor
1498   intelligence as feeding adaptations in cebus monkeys and great apes. *Journal of Human*
1499   *Evolution*, 6(7), 623–641.

1500

1501   Paulina-Carabajal, A., Bronzati, M., & Cruzado-Caballero, P. (2023) Paleoneurology of non-
1502   avian dinosaurs: an overview. In M. T. Dozo, A. Paulina-Carabajal, T. E. Macrini, & S. Walsh
1503   (eds.), *Paleoneurology of Amniotes : New Directions in the Study of Fossil Endocasts* (pp.
1504   267-332). Springer International Publishing, Cham.

1505

1506   Paulina-Carabajal, A., & Canale, J. I. (2010). Cranial endocast of the carcharodontosaurid
1507   theropod *Giganotosaurus carolinii* Coria & Salgado, 1995. *Neues Jahrbuch fur Geologie und*
1508   *Palaontologie*, *258*, 249-256.

1509

1510   Paulina-Carabajal, A., & Currie, P.J. (2012). New information on the braincase and endocast
1511   of *Sinraptor dongi* (Theropoda: Allosauroidea): Ethmoidal region, endocranial anatomy and
1512   pneumaticity. *Vertebrata PalAsiatica,* 50, 85–101

1513

1514   Paulina Carabajal, A., Carballido, J. L., & Currie, P. J. (2014). Braincase, neuroanatomy, and
1515   neck posture of *Amargasaurus cazaui* (Sauropoda, Dicraeosauridae) and its implications for
1516   understanding head posture in sauropods. *Journal of Vertebrate Paleontology*, *34*(4), 870-
1517   882.

1518

1519   Persons IV, W. S., Currie, P. J., & Erickson, G. M. (2020). An older and exceptionally large
1520   adult specimen of *Tyrannosaurus rex*. *The Anatomical Record*, 303(4), 656-672.

1521

1522   Picasso, M. B., Tambussi, C. P., & Degrange, F. J. (2011). Virtual reconstructions of the
1523   endocranial cavity of *Rhea americana* (Aves, Palaeognathae): postnatal anatomical
1524   changes. *Brain Behavior and Evolution*, *76*(3-4), 176-184.

1525

1526   Picasso, M. B. J. (2012). Postnatal ontogeny of the locomotor skeleton of a cursorial bird:
1527   greater rhea. *Journal of Zoology, 286*(4), 303-311.

1528

1529   Pika, S., Sima, M. J., Blum, C. R., Herrmann, E., & Mundry, R. (2020). Ravens parallel great
1530   apes in physical and social cognitive skills. *Scientific Reports*, *10*(1), 1-19.

1531

1532   Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., Heisterkamp, S., and Van Willigen, B.
1533   (2017). Package "nlme". Available at: https://cran.r-project.org/web/packages/nlme/nlme.pdf.

1534

1535 Pontzer, H., Allen, V., & Hutchinson, J.R. (2009). Biomechanics of running indicates

1536 endothermy in bidepal dinosaurs. *PLoS ONE* 4(12): 10.1371.

1537

1538 Prondvai, E., Stein, K., Ősi, A., & Sander, P. M. (2012). Life history of *Rhamphorhynchus*

1539 inferred from bone histology and the diversity of pterosaurian growth strategies. *PLoS ONE*

1540 7(2): e31392.

1541

1542 R Core Team (2022). R: A language and environment for statistical computing. R

1543 Foundation for Statistical Computing, Vienna, Austria. Available at: https://www.R-

1544 project.org/.

1545

1546 Redelstroff, R. & Sander, P.M. (2009). Long and girdle bone histology of *Stegosaurus*:

1547 implications for growth and life history. *Journal of Vertebrate Paleontology* 29(4): 1087-1099.

1548

1549 Redelstroff, R., Hübner, T.R., Chinsamy, A., & Sander, P. M. (2013). Bone histology of the

1550 stegosaur *Kentrosaurus aethiopicus* (Ornithischia: Thyreophora) from the Upper Jurassic of

1551 Tanzania. *Evolutionary Biology* 296(6): 933-952.

1552

1553 Reiner, A. (2023). Could theropod dinosaurs have evolved to a human level of intelligence?.

1554 *Journal of Comparative Neurology*, 531(9), 975–1006.

1555

1556 Revell, L. J., Harmon, L. J., & Collar, D. C. (2008). Phylogenetic signal, evolutionary

1557 process, and rate. *Systematic Biology*, *57*(4), 591-601.

1558

1559 Roese-Miron, L., Jones, M. E. H., Ferreira, J. D., & Hsiou, A. S. (2023). Virtual endocasts of

1560 *Clevosaurus brasiliensis* and the tuatara: Rhynchocephalian neuroanatomy and the oldest

1561 endocranial record for Lepidosauria. *The Anatomical Record*, 1–24. https://doi.org/10.1002/

1562 ar.25212

1563

1564 Ridgway, S. H., Brownson, R. H., Van Alstyne, K. R., & Hauser, R. A. (2019). Higher neuron

1565 densities in the cerebral cortex and larger cerebellums may limit dive times of delphinids

1566 compared to deep-diving toothed whales. *PLoS One*, *14*(12), e0226206.

1567

1568 Rogers, S. W. (1998). Exploring dinosaur neuropaleobiology: computed tomography

1569 scanning and analysis of an *Allosaurus fragilis* endocast. *Neuron*, *21*(4), 673-679.

1570

1571 Rutz, C., Klump, B. C., Komarczyk, L., Leighton, R., Kramer, J., Wischnewski, S.,
1572 Sugasawa, S., Morrissey, M. B., James, R., St Clair, J. J. H., Switzer, R. A., & Masuda, B.
1573 M. (2016). Discovery of species-wide tool use in the Hawaiian crow. *Nature, 537*(7620),
1574 403–407.
1575
1576 Saveliev, S. V., & Alifanov, V. R. (2007). A new study of the brain of the predatory dinosaur
1577 *Tarbosaurus bataar* (Theropoda, Tyrannosauridae). *Paleontological Journal, 41*(3), 281-289.
1578
1579 Sakamoto, M. (2022). Estimating bite force in extinct dinosaurs using phylogenetically
1580 predicted physiological cross-sectional areas of jaw adductor muscles. *PeerJ,* 10: e13731.
1581
1582 Sampson, S. D. & Witmer, L. M. (2007) Craniofacial anatomy of *Majungasaurus*
1583 *crenatissimus* (Theropoda: Abelisauridae) From the Late Cretaceous Of Madagascar,
1584 *Journal of Vertebrate Paleontology*, 27:S2, 32-104.
1585
1586 Schliep, K., Paradis, E., de Oliveira Martins, L., Potts, A., White, T. W., Stachniss, C., &
1587 Kendall, M. (2019). Package 'phangorn'. Available at: *https://cran. r-project.*
1588 *org/web/packages/phangorn/phangorn. pdf*
1589
1590 Schmitt, V., Pankau, B., & Fischer, J. (2012). Old world monkeys compare to apes in the
1591 primate cognition test battery. *PloS one, 7*(4), e32024.
1592
1593 Schweitzer, M. H., Wittmeyer, J. L., & Horner, J. R. (2005). Gender-specific reproductive
1594 tissue in ratites and *Tyrannosaurus rex*. *Science, 308*(5727), 1456-1460.
1595 Seymour, R. S., Bennett-Stamper, C. L., Johnston, S. D., Carrier, D. R., & Grigg, G. C.
1596 (2004). Evidence for endothermic ancestors of crocodiles at the stem of archosaur evolution.
1597 *Physiological and Biochemical Zoology, 77*(6), 1051-1067.
1598
1599 Shatkovska, O. V., & Ghazali, M. (2021). Relative skull size as one of the factors limiting
1600 skull shape variation in passerines. *Canadian Journal of Zoology, 99*(12), 1054-1066.
1601
1602 Shumaker, R. W., Walkup, K. R., & Beck, B. B. (2011). *Animal Tool Behavior: The Use and*
1603 *Manufacture of Tools by Animals* (revised and updated edition edition). Johns Hopkins
1604 University Press, Baltimore.
1605

Smaers, J. B., Rothman, R. S., Hudson, D. R., Balanoff, A. M., Beatty, B., Dechmann, D. K.,
... & Safi, K. (2021). The evolution of mammalian brain size. *Science Advances*, *7*(18),
eabe2101.

Sol, D., Olkowicz, S., Sayol, F., Kocourek, M., Zhang, Y., Marhounová, L., ... & Němec, P.
(2022). Neuron numbers link innovativeness with both absolute and relative brain size in
birds. *Nature Ecology & Evolution*, *6*(9), 1381-1389.

Spence, R. D., Zhen, Y., White, S., Schlinger, B. A., & Day, L. B. (2009). Recovery of motor
and cognitive function after cerebellar lesions in a songbird – role of estrogens. *European*
Journal of Neuroscience, *29*(6), 1225-1234.

Ströckens, F., Neves, K., Kirchem, S., Schwab, C., Herculano-Houzel, S., & Güntürkün, O.
(2022). High associative neuron numbers could drive cognitive performance in corvid
species. *Journal of Comparative Neurology*, *530*(10), 1588–1605.

Tartarelli, G., & Bisconti, M. (2006). Trajectories and constraints in brain evolution in
primates and cetaceans. *Human Evolution*, *21*, 275-287.

Tonini, J. F. R., Beard, K. H., Ferreira, R. B., Jetz, W., & Pyron, R. A. (2016). Fully-sampled
phylogenies of squamates reveal evolutionary patterns in threat status. *Biological*
Conservation, 204, 23-31.

Triki, Z., Aellen, M., van Schaik, C. P., & Bshary, R. (2021). Relative brain size and cognitive
equivalence in fishes. *Brain, Behavior and Evolution*, *96*(3), 124-136.

Troscianko, J., von Bayern, A. M. P., Chappell, J., Rutz, C., & Martin, G. R. (2012). Extreme
binocular vision and a straight bill facilitate tool use in New Caledonian crows. *Nature*
Communications, *3*(1), 1110.

Tsuboi, M., van der Bijl, W., Kopperud, B. T., Erritzøe, J., Voje, K. L., Kotrschal, A., ... &
Kolm, N. (2018). Breakdown of brain–body allometry and the encephalization of birds and
mammals. *Nature Ecology & Evolution*, *2*(9), 1492-1500.

Ulinski, P. S. & Margoliash, D. (1990). Neurobiology of the reptile-bird transition. In E. G.
Jones & A. Peters (eds.), *Comparative Structure and Evolution of Cerebral Cortex, Part I*
(pp. 217-265). Springer, Boston.

1643

1644 Van Schaik, C. P., Triki, Z., Bshary, R., & Heldstab, S. A. (2021). A farewell to the

1645 encephalization quotient: a new brain size measure for comparative primate cognition. *Brain,*

1646 *Behavior and Evolution*, *96*(1), 1-12.

1647

1648 Varricchio, D.J., Martin, A.J., Katsura, Y. (2007). First trace and body fossil evidence of a

1649 burrowing, denning dinosaur. *Proceedings of the Royal Society B* 274:1361–1368.

1650

1651 Wang, Y., Claessens, L.P.A.M., & Sullivan, C. (2023). Deep reptilian evolutionary roots of a

1652 major avian respiratory adaptation. *Communications Biology* 6:3.

1653

1654 Watanabe, A., Gignac, P.M., Balanoff, A.M., Green, T.L., Kley, N.J. & Norell, M.A. (2019).

1655 Are endocasts good proxies for brain size and shape in archosaurs throughout ontogeny?.

1656 *Journal of Anatomy*, *234*, 3, 291-305.

1657

1658 Wedel, M.J. (2006). Origin of postcranial skeletal pneumaticity in dinosaurs. *Integrative*

1659 *Zoology*, 1(2): 80-85.

1660

1661 Wiemann, J., Menéndez, I., Crawford, J. M., Fabbri, M., Gauthier, J. A., Hull, P. M., ... &

1662 Briggs, D. E. (2022). Fossil biomolecules reveal an avian metabolism in the ancestral

1663 dinosaur. *Nature*, *606*(7914), 522-526.

1664

1665 Witmer, L. M., Chatterjee, S., Franzosa, J., & Rowe, T. (2003). Neuroanatomy of flying

1666 reptiles and implications for flight, posture and behaviour. *Nature*, *425*(6961), 950-953.

1667

1668 Witmer, L. M., Ridgely, R. C., Dufeau, D. L., & Semones, M. C. (2008). Using CT to peer into

1669 the past: 3D visualization of the brain and ear regions of birds, crocodiles, and nonavian

1670 dinosaurs. In H. Endo & R.Frey (eds.), *Anatomical Imaging: Towards a New Morphology*

1671 (pp. 67–87). Springer, Tokyo.

1672

1673 Witmer, L. M., & Ridgely, R. C. (2009). New insights into the brain, braincase, and ear region

1674 of tyrannosaurs (Dinosauria, Theropoda), with implications for sensory organization and

1675 behavior. *The Anatomical Record*, *292*(9), 1266-1296.

1676

1677 Woodward, A., C. T. Moore, & M. F. Delaney. (1991). *Experimental Alligator Harvest: Final*

1678 *Report.* Study no. 7567. Bureau of Wildlife Research, Florida Game and Fresh Water Fish

1679 Commission, Tallahassee, 118pp.

1680

1681 Woodward, H. N., Tremaine, K., Williams, S. A., Zanno, L. E., Horner, J. R., & Myhrvold, N.

1682 (2020). Growing up *Tyrannosaurus rex*: Osteohistology refutes the pygmy "*Nanotyrannus*"

1683 and supports ontogenetic niche partitioning in juvenile *Tyrannosaurus*. *Science Advances*,

1684 *6*(1), eaax6250.

1685

1686 Yu, Y., Karbowski, J., Sachdev, R. N., & Feng, J. (2014). Effect of temperature and glia in

1687 brain size enlargement and origin of allometric body-brain size scaling in vertebrates. *BMC*

1688 *Evolutionary Biology*, *14*(1), 1-14.

1689

1690 Zeiträg, C., Jensen, T. R., & Osvath, M. (2022). Gaze following: A socio-cognitive skill rooted

1691 in deep time. *Frontiers in Psychology*, 13:950935.

1692

1693 Zeiträg, C., Reber, S. A., & Osvath, M. (2023). Gaze following in Archosauria — Alligators

1694 and palaeognath birds suggest dinosaur origin of visual perspective taking. *Science*

1695 *Advances*, *9*(20), eadf0405.

1696

1697 Zelenitsky, D. K., Therrien, F., Ridgely, R. C., McGee, A. R., & Witmer, L. M. (2011).

1698 Evolution of olfaction in non-avian theropod dinosaurs and birds. *Proceedings of the Royal*

1699 *Society B: Biological Sciences*, *278*(1725), 3625-3634.

1700

Supplementary Files

Suppl. File 1: Information on the selection and determination of brain endocasts in this

study, including comments on endocast data presented in Jerison (1973), Hopson (1979)

and Hurlburt (1996)

Suppl. File 2: Brain and body mass data of extant sauropsid species, excluding the olfactory

tracts and bulbs. Brain mass data are derived from Hurlburt (1996), except for those of the

Siamese crocodile, which derive from Chentanez et al. (1983). All birds considered here

belong to non-telluravian taxa.

Suppl. File 3: Neuron count data, brain size measures and respective references for extant

sauropsid and mammalian species.

1711

Lepidosauria

Testudinata

Crocodylomorpha

Pterosauria

Ornithischia

Sauropodomorpha

Basal Theropoda

Tyrannosauroidea

Non-avian Maniraptora

Palaeognathae

Basal Neognathae

Telluraves

Archosauria

Ornithodira

Dinosauria

Aves