Title: Meta-metabolome ecology reveals that geochemistry and microbial functional potential

are linked to organic matter development across seven rivers

Authors: Robert E. Danczak[a,*], Amy E. Goldman[a], Mikayla A. Borton[b], Rosalie K. Chu[c], Jason

G. Toyoda[c], Vanessa A. Garayburu-Caruso[a], Emily B. Graham[c], Joseph W. Morad[d], Lupita

Renteria[a], Jacqueline R. Hager[a], Shai Arnon[e], Scott Brooks[f], Edo Bar-Zeev[e], Michael Jones[f], Nikki

Jones[f], Jorg Lewandowski[g,h], Christof Meile[i], Birgit M. Muller[g], John Schalles[j], Hanna Schulz[g,h],

Adam Ward[k], James C. Stegen[a]

[a]Pacific Northwest National Laboratory, Washington, USA

[b]Colorado State University, Colorado, USA

[c]Environmental Molecular Sciences Laboratory, Washington, USA

[d]New Jersey Department of Environmental Protection, New Jersey, USA

[e]Ben-Gurion University of the Negev, Israel

[f]Oak Ridge National Laboratory, Tennessee, USA

[g]Leibniz Institute of Freshwater Ecology and Inland Fisheries, Germany

[h]Humboldt University, Berlin, Germany

[i]University of Georgia, Georgia, USA

[j]Creighton University, Nebraska, USA

[k]Oregon State University, Oregon, USA

*Corresponding Author
Email: robert.danczak@pnnl.gov
Address: 902 Battelle Blvd.
24   P.O. Box 999
25   Richland, WA 99352
26
27

28
29

Abstract

Rivers receive substantial dissolved organic matter (DOM) input from the surrounding land which is transported to the ocean. As this DOM travels through watersheds, it undergoes biotic and abiotic transformations which impact biogeochemical cycles and release CO_2 into the atmosphere. While recent research has increased our mechanistic knowledge of DOM development within watersheds, DOM development across broad spatial distances and within divergent biomes is under investigated. Here, we combined DOM characterization, geochemical analyses, and shotgun metagenomics to analyze samples from seven rivers ranging from the U.S. Pacific Northwest to Berlin, Germany. Initial analyses revealed that many DOM properties separated based upon river type (e.g., wastewater, headwater), though paired geochemical analyses indicated that geochemistry often explained some variation. Analyses rooted in meta-metabolome ecology indicated that, at the global scale, DOM was structured overwhelmingly by deterministic selection. When controlling for scale, however, analyses indicated that ecological assembly dynamics were again structure, in part, by river type. Finally, microbial analyses revealed that many riverine microbes from our systems shared core metabolic functional potential while differing in peripheral capabilities in spatially resolved patterns. Further analyses in the carbon degradation potentials of the recovered metagenomically assembled genomes indicated that the sample rivers had strong taxonomically conserved niche differentiation regarding carbon degradation and that the diversity in carbon degradation potential was significantly related to organic matter diversity. Together, these results help us uncover interconnections between the development of DOM, riverine geochemistry, and microbial functional potential.

1. Introduction

Evaluating the principles underlying the composition of river corridor dissolved organic matter (DOM) has been an ongoing, cross-disciplinary endeavor (e.g., ecology, organic matter characterization, biogeochemistry, climatology) (Danczak et al., 2021; Gómez-Gener et al., 2021; Hu et al., 2022; Kaiser et al., 2017; Liu et al., 2019; Xenopoulos et al., 2021). This is because a significant quantity of organic matter is transported through rivers and undergoes a series of abiotic and biotic transformations that result in the release of CO2 into the atmosphere (Cole et al., 2007; Drake et al., 2018; Gómez-Gener et al., 2021; Graham et al., 2018; Regnier et al., 2013; Stegen et al., 2018). By identifying the principles responsible for promoting and constraining DOM composition and incorporating them into various modeling capabilities, we can improve our ability to evaluate watershed function under future conditions (Song et al., 2020; Wohl et al., 2017; Zarnetske et al., 2018).

Microbial communities are one of the more commonly studied factors responsible for influencing DOM dynamics because they can have major impacts on DOM composition. Most studies looking at the interconnections between microbial communities and riverine DOM examine a single watershed or reach (Garayburu-Caruso et al., 2020; Graham et al., 2018; Grunert et al., 2021; Osterholz et al., 2016; Saarela et al., 2022; Stegen et al., 2016a, 2018). Within the Columbia River Basin, researchers revealed that mixing dynamics and thermodynamic controls were the key intersections between DOM and microbiology (Graham et al., 2018; Stegen et al., 2016a, 2018). Salinity was an overarching control on both the DOM composition and microbial community structure within the Delaware Estuary, suggesting that external environmental pressures can outweigh the impacts of either assemblage (Osterholz et al., 2016). Studies investigating

respiration rates have revealed that DOM thermodynamics play a role in impacting microbial

activity, but only when carbon is a limiting factor (Garayburu-Caruso et al., 2020).

While each of these studies reveals different factors influencing DOM composition and microbial

community structure, these components are often analyzed separately. The recently proposed

analytical framework known as 'meta-metabolome ecology' represents a potential avenue to

develop connections between these data types (Danczak et al., 2020). In brief, this analysis treats

DOM as an assemblage of molecular formulas subject to processes akin to ecological community

assembly processes. By analyzing DOM data in this way, researchers can investigate factors

influencing DOM by identifying transferrable biogeochemical principles and connecting

biological and chemical data. For example, ecological null modeling helped reveal the potential

existence of thermodynamic redundancy, a parallel of functional redundancy observed in

microbial communities but for DOM, within a headwater stream (Danczak et al., 2021). The

application of these ecological tools has begun to reveal connections between DOM and microbial

communities as well. Similar analyses of both a riverine DOM assemblage and microbial

community revealed a lack of selective interconnections, contrary to expectation (Danczak et al.,

2020). A combination of DOM null modeling and functional trait analysis indicated that

deterministic assembly is predicted to increase under situations of increased global change (e.g.,

changing temperature/nutrient conditions) (Hu et al., 2022). Shifting away from null modeling,

functional diversity of DOM assemblages and microbial communities within a freshwater lake

revealed significant covariation indicating a potentially underappreciated role in the global carbon

cycle (Tanentzap et al., 2019). A study analyzing data at a global scale may help reveal transferable

principles concerning the behavior of organic matter is needed because many of these studies are

focused on specific freshwater ecosystems and revealed inconsistent dynamics.

Given the heterogeneity of river corridors across the globe, holistic studies that incorporate

information from many different sampling sites are necessary to uncover these principles. Looking

at 37 different aquatic environments, Kellerman et al. (2018) revealed that the age of organic

matter was consistently related to DOM composition indicating the importance of degradation

state. In support of these findings, D'Andrilli et al., (2015) demonstrated that organic matter

lability was an important differentiating factor across various aquatic environments, revealing that

glacial DOM was the most labile, followed by marine DOM, and finally freshwater. Hu et al.

(2022) studied aquatic microcosms across temperature and nutrient gradients and observed that

differential assembly processes affected the active and inactive components of DOM, with

significant implications for inherent functional traits of DOM molecules. By further expanding

both the range of samples and by integrating multiple data streams, we can better uncover

important factors impacting DOM composition.

Here, we investigated samples from seven rivers spanning a broad geographic extent from the

Middle East to the US Pacific Northwest, as part of the Worldwide Hydrobiogeochemical

Observation Network for Dynamic Rivers (WHONDRS) consortium. By combining DOM

characterization, geochemical analysis, and shotgun metagenomic sequencing partnered to

ecological analyses, we assessed environmental interactions which may explain global, river

corridor DOM variation and integrate prior observations. In doing so, we also sought to

characterize the microbes that interact with and alter river corridor DOM. The primary goal of

each of these analyses is to obtain a cross-ecosystem and transferrable understanding of river DOM

development and connections between DOM and the microbes that eat it. Ultimately, we observed

that the underlying geochemical and lithological conditions appear to be more strongly connected

to DOM composition than other factors and that globally distributed microbial species may have

conserved niches regarding carbon degradation.

2. Materials and Methods

2.1 Site Descriptions. This study obtained samples from the seven rivers spanning different biomes

and flow regimes (Figure 1). These rivers were the Altamaha River (Georgia, USA), Columbia

River (Washington, USA), East Fork Poplar Creek (Tennessee, USA), Erpe River (Germany), HJ

Andrews – Watershed 1 (Oregon, USA), Jordan River (Israel), and Nisqually River (Washington,

USA). The Altamaha River drains a large portion of the state of Georgia to the Atlantic Ocean and

has a watershed that extends over approximately 36,000 km2, encompassing the Northern

Piedmont and the Coastal Plain (Takagi et al., 2017). The site experiences semidiurnal tidal

variations in water levels, with a tidal amplitude on the order of 1-2 m, and occasionally

experiences periods of salinity during low flow periods of October. The Columbia River is a gravel

bedded river with some riparian vegetation and experiences variable flow associated with dam

activity (Arntzen et al., 2006; Stegen et al., 2016a; Villa et al., 2020). The East Fork Poplar Creek

has a streambed of gravel, cobble, and bedrock and is surrounded primarily by forest (Demers et

al., 2018). This site was historically impacted by mercury contamination because of nuclear

weapon development (Brooks and Southworth, 2011; Demers et al., 2018; Loar et al., 2011). The

Erpe River is an urban lowland stream that consists of up to 80% treated wastewater originating

from an upstream wastewater treatment plant (Mueller et al., 2021). The watershed at HJ Andrews

is a shallow, forested, headwater stream that experiences diel fluctuations driven by

evapotranspiration (Danczak et al., 2021). The Jordan River is a karstic fed stream found within

the northern section of the Syrian African Rift Valley and is subject to discharge fluctuations due

to groundwater pumping (Goldman et al., 2019). Finally, the Nisqually River is low-order stream

fed by that originates in Mount Rainier and experiences diel fluctuations in discharge due to glacial

freeze/thaw cycles (Curran et al., 2016). Additional information and site photos for each river can

be found in associated data packages published on ESS-DIVE and a summary of this information

can be found in Table 1.

2.2 Sample Collection and Coordination. Samples from the seven rivers were collected between

July and October 2018 as part of a WHONDRS initiative to study diel river dynamics (Stegen and

Goldman, 2018). As such, the sampling scheme was similar to the methodology described in

Danczak et al., 2021 and was originally devised to investigate temporal patterns. Three positions

separated by ~4m were selected at each river corridor to collect pore water samples.

Approximately 20 mL was collected from each of these locations every 3 h over a 48 h period.

Surface water was additionally collected in triplicate from the same position as the middle pore

water sampling site. In total, 690 total samples were collected over roughly 17 time points. Surface

water was collected using a 60 mL syringe through Teflon tubing while the pore water was

collected using a syringe attached via Teflon tubing to a 30 cm long stainless-steel sampling tube

(MHE Products, MI, USA) with a slotted screen across the bottom ~5 cm. One sampling tube was

installed to 30 cm depth at each pore water sampling location; these tubes sat in place for around

15 min before the first sample was collected and they remained in place during the 48-h time

course of sampling. Prior to sampling a given location, the syringe was flushed 3 times with the

source water to ensure only the desired water was collected. All samples were filtered through a

0.2 μm Sterivex filter (Millipore, MA, USA). At each time point, one filter was used for all three

pore water samples, and a different filter was used for all three surface water samples. Water

passing through the filters was collected for analysis using a needle attached to the filter and

injected through a septum. During sampling, water temperature, approximate water stage, and pH

were measured. Water samples for DOM analysis were injected into amber borosilicate glass vials.

Samples for cations and anions were injected into clear borosilicate glass vials. Once collected,

samples were stored in a cooler on blue ice until they could be frozen until they were processed in

the lab.

2.3 Geochemical Analyses. As with the sampling scheme, the geochemical analyses are similar to

the methods described in Danczak et al., 2021. In brief, anion concentrations were measured using

a Dionex ICS-2000 ion chromatograph with AS40 autosampler using an isocratic method (guard

column: IonPac AG18 guard, 4x50mm; analytical column: IonPac AS18, 4x250mm; suppressor:

RFIC ASRS, 300 rmm, self-regenerating; suppressor current: 57 mA). Standards were made from

Spex CertiPrep (Metuchen, NJ, 08840) 1000 mg/L anion standards and were diluted as follows:

NO_2 in the range of 0.04 to 20 ppm, F in the range of 0.2 to 10 ppm, Cl and SO_4 in the range of

0.16 to 80 ppm, and NO_3 in the range of 0.12 to 60 ppm. Cation samples were prepared with nitric

acid and were measured with a Perkin Elmer Optima 2100 DV ICP-OES with an AS93 auto

sampler. Calibration standards were made with Ultra Scientific ICP standards (Kingstown, RI): P,

Mg, Ca, K, and Na were diluted in the range of 5–4000 ppm, and Fe was diluted in the range of

0.5–400 ppm. Non-purgeable organic carbon (NPOC) was determined by a Shimadzu combustion

carbon analyzer TOC-L CSH/CSN E100V with ASI-L auto sampler calibrated to a range of 0.5 to

10 ppm NPOC as C via potassium hydrogen phthalate (Nacalia Tesque, lot M7M4380). An aliquot

of sample was acidified with 15% by volume of 2 N ultra-pure HCL and then sparged with carrier

gas for 5 min to remove inorganic carbon. The best 3 out of 4 injections replicates were averaged

to get the finalized result.

2.4 Dissolved organic matter sample analysis and data processing. All samples were analyzed

using a Fourier-Transform Ion Cyclotron Resonance mass spectrometer (FTICR-MS) following

methods established previously (Danczak et al., 2021, 2020; Garayburu-Caruso et al., 2020). In

brief, samples were standardized to a given carbon concentrations (NPOC 0.69 – 1.5 mg C/L),

acidified to pH 2 with 85% phosphoric acid, and extracted with PPL cartridges. High-resolution

mass spectra were collected from these samples using a 12 Tesla (12T) Bruker SolariX FTICR

mass spectrometer (Bruker, SolariX, Billerica, MA) equipped with an electrospray ionization

(ESI) source set to negative mode. BrukerDaltonik Data Analysis (version 5.0) was used to

generate a list of m/z values from the raw spectra using the FTMS peak picker module. Chemical

formulae were assigned using Formularity (Tolić et al., 2017) according to the Compound

Identification Algorithm (Kujawinski and Behn, 2006; Minor et al., 2012; Tfaily et al., 2017) with

the following settings: a signal-to-noise ratio >7, mass measurement error <0.5 ppm, and allowing

for CHONSP (Formularity formula settings: O > 0 AND (N + S + P) < 6 AND S < 3 AND P < 2).

The R package ftmsRanalysis (Bramer et al., 2020; Bramer and White, 2019) was used to remove

peaks that were outside the desired m/z range (200 m/z – 900 m/z) or had an isotopic signature,

calculate a number of derived metrics (double-bond equivalent (DBE), modified aromaticity index

(AI_{Mod}), nominal oxidation state of carbon (NOSC)), and organize the data (Hughey et al., 2001;

Koch and Dittmar, 2006; LaRowe and Van Cappellen, 2011; Tfaily et al., 2015). Given the

limitations associated with peak intensities, all intensities were converted to presence/absence

(Tfaily et al., 2017). Once converted to presence/absence, replicates for each sample were merged

such that a peak needed to be present in at least two of the three replicates for it to be considered

"present" in that sample. For each sample, we also calculated molecular formula richness as the

number of molecular formulas assigned in the respective sample.

2.5 Meta-Metabolome Ecology. Following the protocols established in Danczak et al., we

generated a transformation-weighted molecular characteristics dendrogram (TWCD) (Danczak et

al., 2020). The molecular characteristics dendrogram component of the dendrogram was generated

by first measuring the Euclidean distance between molecular formulas based on their derived

metrics (e.g., elemental composition, double-bond equivalents, modified aromaticity index, and

Kendrick's defect). These characteristic distances were then weighted by transformation distances

associated with the mass of each molecular formula. While described in detail elsewhere (Buser-

Young et al., 2023; Danczak et al., 2021, 2020; Fudyma et al., 2021; Graham et al., 2018), the

transformation analysis estimates the presence of putative biochemical transformations by

measuring the mass differences between identified peaks which are then related to a database of

known and commonly observed biochemical transformations. Relationships between peaks are

then established by measuring the number of transformations required for one peak to potentially

become another. These distances are normalized to a range between 0 and 1 and are used to down

weight the molecular characteristic distance. This transformation-weighted molecular

characteristic distance matrix is then used as input for an unweighted pair group method with

arithmetic mean (UPGMA) hierarchical cluster analysis. The output of this analysis represents

total potential relations that could exist across all DOM identified in this study, therefore, this

TWCD represents an approximation of global DOM relationships.

The TWCD was used primarily to perform a null modeling analysis known as β-nearest taxon

index (βNTI) to estimate the ecological assembly processes impacting the DOM (Danczak et al.,

2020; Stegen et al., 2015, 2013, 2012). To perform this analysis, a null distribution of 999 β-mean

nearest taxon distance (βMNTD) values were generated and compared to the observed βMNTD

value for a given pair of assemblages within each river (e.g., cross-river comparisons were not

analyzed). Pairwise comparisons with $|\beta NTI| > 2$ indicate that deterministic processes were

responsible for observed differences in metabolite composition. In contrast, pairwise comparisons

with $|\beta NTI| < 2$ indicate that stochastic processes were responsible for observed differences in

metabolite composition. Furthermore, the deterministic processes can be separated into two

classes. When $\beta NTI > 2$, differences in metabolite composition are greater than would be expected

by random chance (i.e., greater than the stochastic expectation). This is analogous to 'variable

selection,' which occurs when deterministic processes drive divergence in composition between a

pair of assemblages (Dini-Andreote et al., 2015; Stegen et al., 2015). When $\beta NTI < -2$, differences

in metabolite composition are less than the stochastic expectation. This is analogous to

'homogeneous selection,' which occurs when deterministic processes drive convergence in

composition between a pair of assemblages. We would like to highlight that while null modeling-

was performed on each river independently, the same dendrogram was used in each analysis and

was not pruned. This approach allowed us to take a more global perspective in assembly processes

and potentially overcome some of the scale limitations observed in other meta-metabolome

ecology studies (Danczak et al., 2023, 2021).

Using a similar approach to the microbial data above, we analyzed alpha diversity and beta

diversity using the diversity and vegdist/metaMDS functions, respectively, from the vegan R

package (v2.6.4). Additionally, we performed a distance-based redundancy analysis (dbRDA)

leveraging Jaccard dissimilarity to evaluate relationships between the meta-metabolome and

geochemical variables. First, we limited the effect of co-correlated variables by identifying them

using the vif.cca function and manually removing variables with high degrees of co-correlation

from our model; variables removed from the model were K, Na, and SO_4 Next, we performed

forward and reverse variable selection to identify the minimum set of geochemical variables that

could explain metabolome variation using the ordiR2step function. The final set of selected

variables were NPOC, Cl, Mg, TN, NO_3, F, P, Fe, and pH. Finally, we evaluated the significance

of the final dbRDA utilizing a permuted ANOVA and visualized the RDA with loadings.

2.6 DNA Extraction and Sequencing. DNA was extracted from filters using Zymobiomic's

Quick-DNA Fecal/Soil Microbe Microprep kit (cat# 6012) using the manufacturer's protocol and

the soil option. Metagenomic libraries were prepared using the Tecan Ovation Ultralow System

V2 and were sequenced on the NovaSEQ6000 platform on a S4 flow cell at Genomics Shared

Resource, Colorado Cancer Center, Denver, CO, USA.

2.7 Metagenome Assembly, Binning, and Annotation. All raw reads were trimmed for length and

quality using Sickle v1.33 (Joshi and Fass, 2011). To maximize bin recovery and quality, we used

multiple assembly methods which are denoted in each assembly and bin name as follows: (B)

megahit v1.1.1, (C) IDBA-UD 1.1.0, (D) metaspades v3.13.0, (E) bbcms followed by megahit

v1.1.1, and (F) subsample reads (25%) using bbtools followed by IDBA-UD 1.1.0 (Bushnell, 2020;

Li et al., 2015; Nurk et al., 2017; Peng et al., 2012). All assemblies were then individually binned

using Metabat2 v2.12.1 with default parameters to obtain microbial metagenome assembled

genomes (MAGs) (Kang et al., 2015).

MAG completion was estimated using CheckM v1.1.2 (Parks et al., 2015). To ensure only quality

MAGs were utilized for metabolic analyses, we discarded all MAGs that had completion <50%

and contamination >10%. This equates to "high quality" (HQ) bins and "medium quality" (MQ)

bin cutoffs outlined by the genome consortium standard. The resulting MAGs were manually

curated and then dereplicated using dRep with default parameters to result in a final set of 302

MAGs (>99% ANI which represents strain-level MAG distinctions) (Olm et al., 2017)

(Supplemental Table 1). To quantify MAG relative abundance in each temporal sample and

condition, trimmed metagenomic reads were mapped to the dereplicated MAG set using bbmap

(v38.70) at minid=95, and output as sam files which were converted to sorted bam files using

samtools (v1.9) (Bushnell, 2020; Danecek et al., 2021). We required reads to map to at least 75%

of a MAG in each sample, and second the MAG had to have at least 3x coverage in that sample.

To determine MAGs that had reads mapped to at least 75% of the MAG, we used CoverM (v0.3.2)

in genome mode to output MAGs that passed this threshold (–min-covered-fraction 75); MAGs

were considered present within a sample if they had mapping above this threshold (Woodcroft,

n.d.). To obtain reads per base for each MAG, we used CoverM (v0.3.2) in genome mode to output

reads_per_base (reads mapped/genome length), and from this calculated MAG coverage as

reads_per_base x 151 bp (Woodcroft, n.d.). MAGs were taxonomically classified and using the

Genome Taxonomy Database (GTDB) Toolkit v1.3.0 in August 2020 (Chaumeil et al., 2020).

MAG scaffolds were annotated using the DRAM v1.0 tool which uses PFAM (v33.1), KEGG

(v89.1), dbCAN (v9), MEROPS (v120), and VOGDB databases (https://vogdb.org/) for

annotations (Finn et al., 2016; Kanehisa et al., 2016; Rawlings et al., 2018; Yin et al., 2012). A

phylogenetic tree visualization was generated by 1) identifying ribosomal protein S3 (rps3) amino

acid sequences in each MAG by leveraging a Hidden Markov Model (HMM) from PFAM, 2)

aligning rps3 sequences using MUSCLE (Edgar, 2004), 3) trimming the MUSCLE alignment

using trimAl (Capella-Gutiérrez et al., 2009), 4) using the alignment to generate a phylogenetic

tree using FastTree (Price et al., 2010), and 5) finally visualizing the tree using the ggtree R

package (Yu et al., 2017). Additional analyses were performed on all MAGs using the dbCAN

standalone pipeline (Yin et al., 2012).

Using the MAG coverage data and the CAZyme count information obtained via the dbCAN

analyses, we performed alpha diversity analyses in each sample (diversity, vegan package v2.6.4;

Oksanen et al., 2019). Beta-diversity was measured using Bray-Curtis dissimilarity (vegdist, vegan

package v2.6.4) and was visualized using non-metric multidimensional scaling (NMDS;

metaMDS, vegan package v2.6.4). Microbial taxa at the Order-level and CAZyme identity were

subsequently fitted to their respective NMDS graphs using the envfit function in the vegan package

(v2.6.4) to variables influencing each sample or MAG.

2.8 Statistics and Plot Generation. Statistical analysis and plot generation were performed using

R v4.3.1 and RStudio (Posit team, 2023; R Core Team, 2023). Plotting relied on a mixture of

functions from the tidyverse collection of R packages (v2.0.0), ggthemes (v4.2.4), ggrepel (v0.9.3),

and ggpubr (v0.6.0) (Arnold, 2021; Kassambara, 2023; Slowikowski, 2023; Wickham et al., 2019).

Kruskal-Wallis tests were used to evaluate variation across multiple groups (e.g., whether

differences existed across all rivers) using the kruskal.test function or the stat_compare_means

function in ggpubr (v0.6.0), and Mann Whitney U tests were used to evaluate pairwise variation

using the wilcox.test function. Spearman correlations were performed using the rcorr function

from the Hmisc package (v5.1.0) (Harrell, 2023).

All code and relevant manuscript files can be found on ESS-DIVE at the following DOI:

[published when available]. All raw data used in this study can also be found on ESS-DIVE using

the following DOIs: https://data.ess-dive.lbl.gov/view/doi:10.15485/1509695, https://data.ess-

dive.lbl.gov/view/doi:10.15485/1577266,   https://data.ess-

dive.lbl.gov/view/doi:10.15485/1577260,   https://data.ess-

dive.lbl.gov/view/doi:10.15485/1577265,   https://data.ess-

dive.lbl.gov/view/doi:10.15485/1577278,   https://data.ess-

dive.lbl.gov/view/doi:10.15485/1577263,   https://data.ess-

dive.lbl.gov/view/doi:10.15485/1576995 (Chu et al., 2019; Danczak et al., 2019; Garayburu-

Caruso et al., 2019; Goldman et al., 2019; Renteria et al., 2019; Stegen et al., 2019; Wells et al.,

2019). Metagenomic data has been uploaded to NCBI and can be accessed via the Bioproject #

PRJNA946291.

3. Results and Discussion

3.1 Globally distributed rivers exhibit strong geographic environmental metabolome signatures

despite geochemical commonalities in similar surrounding environments. Geochemical analyses

revealed substantial environmental variation across the seven globally distributed rivers. Looking

first at the geochemical patterns using a principal component analysis (Supplemental Figure 2),

we see that each river has characteristic geochemical parameter values. Both surface water and

pore water from the Erpe River (ER) are characterized by higher NPOC, K, SO_4, Na, Cl, and NO_2

concentrations, while the surface water of East Fork Poplar Creek (EFPC) has higher Ca, Mg,

NO_3, Total N, and P concentrations. Given that both systems are considered either currently or

historically wastewater contaminated (Loar et al., 2011; Mueller et al., 2021), we would anticipate

elevated levels of many geochemical parameters like NPOC and NO_3/NO_2 while other

geochemical parameters like Ca and Mg are likely due to the underlying lithology. The Altamaha

River (AR) is characterized by elevated iron concentrations likely due to the dominant sediment

mineralogy, lower pH, and higher NPOC (Supplemental Figure 1) (Bhatti et al., 2009; Roebuck et

al., 2020; Shi et al., 2001). The 4 remaining rivers (and the pore water from EFPC) are defined by

lower concentrations of most parameters, particularly the HJ Andrews Watershed (HJA) and the

Nisqually River (NR) which are a headwater stream and a glacial melt run-off stream, respectively

(Curran et al., 2016; Danczak et al., 2021). Examining carbon data for combined pore and surface

water samples (Figure 1c), we observe significant differences in NPOC concentrations across the

seven rivers resulting in the formation of three broad groups: ER and AR constitute a high NPOC

group; the Jordan River (JR), HJA, and NR form a low NPOC group; the Columbia River (CR)

and EFPC have intermediate concentrations between these two groups.

Shifting focus to average environmental metabolome characteristics, each of the seven rivers has

a distinct organic matter mixture. The average nominal oxidation state of carbon (NOSC), a metric

which can track the thermodynamic bioavailability of organic matter, indicated no groups formed

among the rivers. Nevertheless, there were significant differences among some pairwise

comparisons (e.g., NR was significantly lower than AR), though few differences exist between

water types (e.g., EFPC surface water was significantly different from EFPC pore water)

(Supplemental Table 2, Figure 1d). Average double bond equivalents (DBE) and modified

aromaticity index (AI_{Mod}) were used to evaluate potential structural patterns of organic matter

across rivers and water types. Patterns based upon DBE are more apparent with AR, CR, and HJA

exhibiting higher values, ER, JR, and NR exhibiting lower values, each for both water types; DBE

for EFPC was different across the pore and surface water. No clear groupings across rivers are

apparent in AI_{Mod} in either surface water or pore water samples though EFPC, ER, and JR have

significantly different average aromaticity across surface and pore water. Overall, the less

urbanized, headwater rivers (e.g., NR, JR) have lower NOSC and DBE overall, while the larger

rivers with many tributaries (e.g., AR, CR) have higher NOSC and DBE. The two wastewater and

one of the headwater rivers (HJA) have NOSC and DBE values intermediate between the two end-

member groups. While similar river types (e.g., headwater, wastewater) group together on some

metrics, the specific location of each river plays a substantial role in shaping average

characteristics. This pattern is also in keeping with data indicating that the surrounding land (often

land coverage) plays a differentiating role in DOM composition (Danczak et al., 2023; Roebuck

et al., 2020). Moreover, these patterns are not exclusively a function of river NPOC concentration.

River type (e.g., wastewater vs. headwater) is a significant determinant for molecular formula

richness (e.g., the number of molecular formulas). Rivers fell into two groups based on molecular

formula richness – AR, CR, EFPC, and ER form a high richness group, while HJA, JR, and NR

constitute a low richness group and these groups held across both water types (Figure 1d). The

patterns in richness closely mirror the NPOC concentrations, with some disagreement; namely,

EFPC and the CR have higher richness relative to their NPOC concentrations. Likely, this is in

part driven by wastewater for EFPC and the large number of tributaries that feed into the CR

Specifically, wastewater is typically high in diverse types of organic matter while the many

tributaries of the CRB bring the potential for varied organic matter (e.g., pyrogenic DOM from

wildfires, farmland input, urban, etc.) despite comparatively low overall concentrations (Danczak

et al., 2023; Vaughn et al., 2023). The low formula richness group consists of samples that were

collected either in or close to the headwaters. This data may suggest that larger rivers under

agricultural/urban influence (CR and AR) and wastewater-impacted rivers (EFPC and ER) will

have higher richness than less urbanized headwater streams (HJA, JR, and NR). Such a pattern is

consistent with observations made within the Upper Mississippi River basin whereby forested

rivers had lower molecular formula richness than urban and agricultural rivers (Vaughn et al.,

2023). These results, in conjunction with previous studies in diverse basins, indicate that

incorporating the surrounding land into predictions regarding DOM development is critical.

However, the specific interactions between land type and environmental conditions and

subsequent impacts on DOM are not clear.

3.2 Ecological null modeling of environmental metabolomes reveals geographically distinct forces

drive the development of DOM in rivers. Multivariate analyses revealed that the organic matter in

each river and both water types was significantly different from the other (Figure 2a). The

exceptions to this pattern were HJA and NR which had statistically indistinguishable surface and

pore water samples. Both systems have low NPOC values and HJA has a high degree of

hydrological connectivity between surface and pore water (Ward et al., 2018; Wondzell, 2006;

Wondzell et al., 2007). Importantly, the HJA results differ from our previous study which indicated

that the pore water and surface water metabolomes were significantly divergent (Danczak et al.,

2021). The contrasting pattern for HJA between the two studies is likely driven by a shift in scale

– by providing more chemodiversity from additional rivers in this study, the surface and pore water

HJA metabolomes become much more similar in comparison. Similar results are noted in

ecological studies and emphasize that scale has important influences over perceived variation in

composition (Hernández, 2020; Ladau and Eloe-Fadrosh, 2019; Levin, 1992; Swenson et al.,

2006).

The DOM composition of each river was significantly, but weakly, related to the corresponding

geochemical profile (Mantel ρ: 0.1985, p-value: 0.001; PROTEST m_12^2: 0.7379, p-value:

0.001). A redundancy analysis (RDA) elucidated which geochemical parameters explained the

cross-river metabolome differences (Figure 2b). Initial forward/reverse variable selection prior to

the RDA suggested that 9 parameters (NPOC, Cl, Mg, TN, NO_3, F, P, Fe, and pH) sufficiently

captured variability across the samples. The RDA revealed that 6 parameters (Cl, Mg, NO_3, NPOC,

TN, and P) were associated with the samples from wastewater-impacted streams (ER and the

surface water of EFPC), while HJA, JR, and NR were associated with higher pH values and lower

concentrations of those 6 parameters. As in other analyses, the pore water of EFPC, AR, and CR

were intermediate to the other 4 rivers, though there was an iron signal associated with AR and a

weak fluoride signal for CR. These relationships indicate a cohesion between geochemical and

organic matter variability and suggest that geochemical condition might be a stronger predictor

than geographical location (e.g., NR and JR are more similar each other than other geographically

closer locations). While useful, these analyses do not tell us how the geochemistry is related to the

ecological processes which shape the environmental metabolome.

Following protocols established via meta-metabolome ecology, we observed that rivers
experienced significantly different ecological assembly processes (Danczak et al., 2020). Briefly,
this method leverages a null model called β-nearest taxon index (βNTI) to identify ecological
processes impacting the environmental metabolome. Based upon the βNTI value, we can assign
either a deterministic (e.g., selection; $|\beta NTI| > 2$) or stochastic (e.g., dispersal; $|\beta NTI| < 2$)
influence. Deterministic processes can be further separated based upon their sign. When βNTI < -
2, the communities were assembled due to homogenous selection and are more similar than would
be expected by random chance. When βNTI > 2, communities experienced variable selection
because they are more divergent than would be expected by random chance.

Comparing the environmental metabolomes among rivers, each river experiences variable
selection overall (βNTI > 2), with ER having the strongest variable selection and AR having the
weakest, bordering on stochastic assembly (Figure 3a). By relating βNTI values to geochemical
parameters, we can reveal which parameters are important in structuring environmental
metabolomes (Danczak et al., 2021, 2016; Stegen et al., 2016b). This analysis revealed that 8
parameters (Ca, SO_4, K, Mg, NPOC, Na, NO_2, and TN in order of correlation strength) were
significantly positively related to global βNTI values and iron was negatively related
(Supplemental Table 3). Looking to those parameters which were influential in the RDA (Figure
2b), we see overlap between these relationships suggesting that many of the influential parameters
may impact ecological processes likely through either metabolic or abiotic interaction. For
example, some parameters like calcium, potassium, or sodium might be indicative of underlying
lithology (e.g., calcium carbonate content) or anthropogenic influence and suggest that certain

geological characteristics might lead to deterministic, variable selection (e.g., larger differences in

sodium are related to higher βNTI). Alternatively, other metrics like sulfate, total nitrogen, or

nitrite may highlight specific nutrient, metabolic, or respiratory activity related to divergent

environmental metabolomes. Iron being the only negative relationship may point to a few

possibilities: 1) iron concentrations fluctuate inversely with one of the positively related

geochemical variables (e.g., pH), or 2) higher iron concentrations are associated with

homogeneous selection in environmental metabolomes via metabolic (e.g., iron reduction) or

abiotic (e.g., organometallic) interactions (Lv et al., 2016). Iron's interpretation is further

complicated in that many sites had iron concentrations that were below detection (Supplemental

Figure 1). Regardless of the specific interaction, geochemical conditions are significantly related

to the ongoing ecological processes shaping environmental metabolomes across seven

geographically distinct rivers.

By analyzing βNTI comparisons within a given group (here within rivers), we can evaluate local,

rather than global, processes (Figure 3b). EFPC and ER are structured by variable selection (βNTI

> 2), HJA and JR are structured by homogeneous selection (βNTI < -2), and AR, CR, and NR are

structured by stochastic processes (2 > βNTI > -2). Firstly, we see that the wastewater rivers in

this study are governed by distinct ecological assembly processes. Secondly, rivers with low

carbon content were more impacted by homogeneous processes. These low carbon rivers also have

lower geochemical variability as indicated by principal component and the redundancy analyses.

Combined, this suggests that low environmental variability may lead to homogenized organic

matter assemblages. Finally, we see stochastic (i.e., dispersal-based) processes impacting rivers

regardless of geochemical condition (e.g., low NPOC, low iron NR and high NPOC, high iron AR

were impacted by stochastic assembly). These results point to an observation whereby a priori

river type appears to be an important characteristic of DOM development in some cases (e.g., ER

as wastewater, HJA as headwaters) while dispersal may dominate in others. Similar patterns have

been discussed previously whereby land use has been related to DOM diversity (Danczak et al.,

2023; Roebuck et al., 2020; Vaughn et al., 2023). Results from these studies support our ecological

observations, and indicate that geochemical conditions and the specific location of a river

(including the historical influences up to the point in time when sampling occurred) are important

in understanding the development of DOM.

3.3 The global evolutionary history of river corridor microbes is more significant in dictating

carbon degradation potential than local pressures. Microbes are key drivers of biogeochemical

cycles and are linked to river corridor organic matter. By understanding the putative functions of

resident microbes, we can better understand the forces driving DOM development (Graham et al.,

2018, 2016). To evaluate the functional potential as it relates to organic matter diversity, a subset

of surface water and pore water samples from AR, CR, EFPC, ER, and JR (1 from each, 10 total)

were analyzed using shotgun metagenomic sequencing and a total of 302 metagenomic assembled

genomes (MAGs) of medium or higher quality were obtained. These MAGs belong to diverse

lineages which are detected across all river corridors (Figure 4a). At higher taxonomic levels,

members of the FCB Group and PVC Superphylum were broadly distributed. At lower taxonomic

levels, MAGs from the Order Burkholderiales represent at least 1% relative coverage in all rivers

aside from JR (Supplemental Figure 3). These Burkholderiales MAGs have consistent carbon,

nitrogen, and sulfur processing functional potential, though those found in ER often encoded both

an acetate kinase and pyruvate formate lyase and more diverse nitrogen metabolisms

(Supplemental Figure 4). This may indicate that these organisms encounter more diverse carbon

sources (e.g., short chain fatty acids) and nitrogen types due to their wastewater habitat (Cao et al.,

2013; Zhou et al., 2023). While no Burkholderiales were observed within JR, this river shared

members of the Order Flavobacteriales with AR, CR, and EFPC and the Order Rhizobiales with

AR. The core functional potential of these MAGs is also consistent across the sampled river

corridors (Supplemental Figure 4). However, sequences annotated as PEP carboxylase, an enzyme

involved in carbon fixation, are present in the Rhizobiales and Flavobacteriales in JR only.

A similar consistency in core metabolism is observed when examining less broadly distributed

taxa (Figure 4b; Supplemental Figure 5). Most Orders (aside from UBA9983) encode common

carbon, nitrogen, and sulfur metabolisms with limited variation. The 3 Campylobacterales MAGs

found in ER encode pyruvate:ferrodoxin oxidoreductase (PFOR), pyruvate water dikinase, formate

dehydrogenase. The Chitinophagales were observed both in CR (4 MAGs) and JR (2 MAGs), but

exhibited geographical variations: namely, the JR variants encoded PFOR, the RNF complex,

formate-tetrahydrofolate ligase, and a Na-transporting NADH:ubiquinone oxidoreductase. A

single Methanomicrobiales MAG was recovered from EFPC and encoded the potential for carbon

fixation via the Wood-Ljungdahl Pathway (CO-methylated acetyl-CoA synthase) and for

methanogenesis via CO_2. Many Nanopelagicales MAGs were primarily detected in AR (19

MAGs), though some were observed in CR (1 MAGs) and ER (3 MAGs) with low coverage. These

genomes were characterized by the presence of genes encoding an L-lactate dehydrogenase, a

partial malonate semialdehyde pathway, a PEP carboxylase, formate-tetrahydrofolate ligase, and

3-phenylpropionate dioxygenase. Finally, the UBA9983 completely lacks genes belonging to the

Citric Acid Cycle and has limited metabolic versatility in keeping with other members of the

Patescibacteria (also known as the Candidate Phyla Radiation) (Castelle et al., 2017; Vigneron et

al., 2023). UBA9983 also exhibits geographical patterns: the 4 MAGs detected in AR encode

pyruvate water dikinase and D-lactate dehydrogenases while the 15 ER MAGs encode fewer

SCFA enzymes and more diverse nitrogen cycling genes.

These patterns lead to two key observations for these microbial communities: 1) despite significant

geographical distances and phylogenetic variation, MAGs observed across the five river corridors

have largely similar core metabolic functions, and 2) differences that do arise appear to be related

to carbon-type utilization. This type of conservation in functional potential despite phylogenetic

variation has been observed in other ecosystems and likely indicates that microbial communities

across these five river corridors face many common pressures, though not necessarily dispersal

events (Isobe et al., 2020; Martiny et al., 2013; Walkup et al., 2023). In turn, this leads to the

development of a common set of functions encoded by varied taxa. However, while many

pressures may be similar, each river corridor has a set of unique conditions as evidenced by the

geochemical and organic C measurements above (Figure 1; Supplemental Figure 1). This gives

rise to a mixture of shared core metabolism and unique peripheral metabolisms, sometimes within

the same taxa. For example, while Rhizobiales and Flavobacterales are found in 3 and 4 river

corridors, respectively, only those in JR contain a gene characteristic of carbon fixation. The JR

also featured some of the lowest NPOC concentrations observed within the dataset. Taken

together, this suggests that conditions in the JR might select for carbon fixation due to low carbon

concentration. While this is only one example, the other variations in functional potential detected

across rivers suggest a more differentiating impact of carbon conditions than other variables (e.g.,

the differential presence of PFOR, formate-active sequences, etc.).

Diving more deeply into the carbon cycling potential of the Orders described above revealed potential taxonomically conserved niche differentiation across the seven rivers (Figure 5a). Specifically, we see that the Verrucomicrobiales and Rhizobiales (both present in JR) contain more diverse and significantly divergent CAZymes than the UBA9983 (characteristic of ER). An expanded CAZyme analysis for all MAGs revealed that potential niche differentiation exists across the taxonomic groups found in the five river corridors. This is captured by the high cross-taxonomic CAZyme beta-diversity (e.g., distance in the ordination) (Figure 5b). For example, while we see that the Burkholderiales and Nanopelagicales have comparable CAZyme alpha diversity, the distance between them in the ordination is high suggesting divergent roles in environmental carbon degradation. Meanwhile, phylogenetic conservation within taxonomic groups is also suggested by the relatively low within-group beta-diversity. For example, we see that MAGs from the Burkholderiales cluster closely together despite being obtained from 4 geographically distinct rivers (AR, CR, ER, and EFPC). These patterns of taxonomic niche differentiation and phylogenetic conservation of functional potential suggest that the global evolutionary history is more important to an organism's CAZyme pool than local evolutionary pressures. When combined with the above observation that many differences arise in the functional potential related to smaller carbon compounds, clades are selected for based upon their full carbon degradation potential which can be common across rivers (e.g., Burkholderiales) or specific to an individual river (e.g., UBA9983).

When various microbial diversity measurements (i.e., Shannon diversity for the overall community, for all CAZYmes contained in the genomes of the community, and for only glycosyl

hydrolases in those genomes) are related to DOM richness (i.e., the number of molecular formulas

within a sample), we observe significant correlations (Figure 5c). Specifically, the diversity of

genes putatively involved in carbon cycling (e.g., CAZymes, glycosyl hydrolases) have slightly

stronger relationships with DOM richness than overall diversity (Shannon's $\rho = 0.75$ vs. CAZyme

and GH $\rho = 0.82 – 0.83$). This highlights a connection between DOM variety within an ecosystem

and the taxonomic and functional potential of microorganisms. When considered in concert with

the potential niche partitioning described before, these results suggest that microorganisms may

assemble into a community based on the exposure to varied types of DOM, at least in part. Given

that carbon cycling potential evolved prior to the dissemination of specific microbial taxa as

suggested by the conservation of degradation capability within a given taxonomy (Figure 5ab),

DOM might act as a filter for selecting for present groups and may point to causality in the

composition of microbial communities. We suggest caution, however, in extrapolating these

results too broadly as they were obtained from a limited dataset, from limited data types.

Exploration of a greater number of samples across a wider range of biomes while leveraging varied

technologies (e.g., metatranscriptomics, metaproteomics, lipidomics, etc.) are necessary before

suggesting anything authoritative. Despite these limitations, these results offer a promising

glimpse into the ecological interconnections between microbes and the DOM they consume.

599   4. Conclusions

The combination of meta-metabolome ecology, in-depth geochemical analyses, and shotgun

metagenomics provided insights into the potential driving forces behind organic matter chemistry

and the interconnections between microbial communities and organic matter across seven rivers.

The geochemistry of an environment is significantly related to metabolome composition as

suggested by traditional multivariate analyses (e.g., Mantel and permuted PROCRUSTES tests)

and null modeling approaches. This pattern was expected due to the distinct geochemical

signatures of each location. However, our null modeling revealed that there are certain key

geochemical parameters that may play a role in structuring organic matter. Calcium, NPOC,

sulfate, potassium, sodium, magnesium, total nitrogen, and nitrate consistently were detected as

important predictors for variable selection (e.g., positively related to βNTI) (Supplemental Table

4). These geochemical characteristics largely reflect watershed characteristics (e.g. soil

characteristics, land cover) and geological processes (e.g., underlying lithology). For example,

calcium concentrations in circumneutral conditions are largely controlled by natural weathering

processes (Weyhenmeyer et al., 2019). Relationships with calcium (or other primarily weathering

derived characteristics) might provide insight into organic matter dynamics that are associated with

certain geological histories. They might also point to specific land-use or metabolic processes

driving organic matter compositions. Land-use and/or land type (i.e., forested, urban, wastewater,

etc.) significantly impacts carbon flux (Regnier et al., 2013) and potentially explains relationships

with NPOC. Many of these variables, however, are also involved in numerous metabolic pathways

and could point to organismal activity. For example, previous research in HJA and CR revealed

that nitrogen metabolism is an important component of DOM processing within those rivers

(Danczak et al., 2021; Graham et al., 2018; Stegen et al., 2018). Therefore, the relationships

between total nitrogen or nitrate and DOM may represent dynamics in associated nutrient or

energy pathways.

Relationships within the microbial communities were more challenging to identify due to the

limited sample number, but a few clear patterns emerged. First, we observed members of the

Burkholderiales in 4 of the 5 analysed rivers (AR, CR, EFPC, and ER) and members of the

Flavobacteriales in 3 of the 5 rivers (AR, CR, and JR). Focusing on carbon cycling potential, it

appears that the Burkholderiales play similar roles in each river despite significant geographical

distances (Supplementary Figure 3). Second, those lineages which are only found in fewer rivers

appear to play diversified roles, potentially related to environmental conditions of a given area.

For example, members of the Patescibacteria (UBA9983) are significantly functionally divergent

from the Verrucomicrobiales and Rhizobiales. Whereas the Verrucomicrobiales and Rhizobiales

appear to have similar carbon cycling potentials as many aerobic organisms, the UBA9983 MAGs

encode no respiratory capabilities, have reduced CAZyme diversity, and seem to be obligate

fermenters consistent with their taxonomic assignment (Castelle et al., 2017; Vigneron et al.,

2023). These differences in putative niche space represent the potential interactions between the

microbial community and DOM. While members with a broader potential niche space, like the

Rhizobiales or Flavobacteriales, may be able to access more DOM types due to a greater diversity

of encoded CAZymes, members with narrower potential niche space (e.g., UBA9983) would be

able to target less DOM. When DOM-based diversity measurements are related to these microbial

data, we observed significant interconnections highlighting putative role for DOM in selecting for

subsets of microbial taxa.

By examining organic matter chemistry in conjunction with geochemistry and microbial functional

potential, we were able to identify general trends across seven globally distributed rivers. Namely,

we observed that both geochemical conditions and the specific location of a river (i.e., the type of

the river, the land use history of the surrounding area, etc.) are important in understanding riverine

DOM composition. Additionally, we observed that these same factors significantly influenced the

underlying ecological assembly processes structuring DOM, often in a scale-dependent manner.

Finally, our metagenomic analyses indicated that members within the microbial communities are

differentiated by their carbon degradation potential with molecular diversity serving as a putative

selective filter. Together, these results give us a glimpse into the interconnections between DOM

development and microbial functional potential while also highlighting environmental parameters

that might be broadly important in understanding DOM in rivers.

References

Arnold, J.B., 2021. ggthemes: Extra Themes, Scales and Geoms for "ggplot2."

Arntzen, E.V., Geist, D.R., Dresel, P.E., 2006. Effects of fluctuating river flow on

groundwater/surface water mixing in the hyporheic zone of a regulated, large cobble bed

river. River Res. Appl. 22, 937–946. https://doi.org/10.1002/rra.947

Bhatti, A.M., Rundquist, D., Schalles, J., Ramirez, L., Nasu, S., 2009. A comparison between

above-water surface and subsurface spectral reflectances collected over inland waters.

Geocarto Int. 24, 133–141. https://doi.org/10.1080/10106040802460707

Bramer, L.M., White, A., 2019. ftmsRanalysis: Analysis and visualization tools for FT-MS data.

R package version 1.0.0. [WWW Document]. URL https://github.com/EMSL-

Computing/ftmsRanalysis

Bramer, L.M., White, A.M., Stratton, K.G., Thompson, A.M., Claborne, D., Hofmockel, K.,

McCue, L.A., 2020. FtmsRanalysis: An R package for exploratory data analysis and

interactive visualization of FT-MS data. PLoS Comput. Biol. 16, 1–12.

https://doi.org/10.1371/journal.pcbi.1007654

Brooks, S.C., Southworth, G.R., 2011. History of mercury use and environmental contamination

at the Oak Ridge Y-12 Plant. Environ. Pollut. 159, 219–228.

https://doi.org/10.1016/j.envpol.2010.09.009

Buser-Young, J.Z., Garcia, P.E., Schrenk, M.O., Regier, P.J., Ward, N.D., Biçe, K., Brooks,

S.C., Freeman, E.C., Lønborg, C., 2023. Determining the biogeochemical transformations

of organic matter composition in rivers using molecular signatures. Front. Water 5,

1005792. https://doi.org/10.3389/frwa.2023.1005792

Bushnell, B., 2020. BBTools software package.

Cao, S., Wang, S., Peng, Y., Wu, C., Du, R., Gong, L., Ma, B., 2013. Achieving partial

denitrification with sludge fermentation liquid as carbon source: The effect of seeding

sludge. Bioresour. Technol. 149, 570–574. https://doi.org/10.1016/j.biortech.2013.09.072

Capella-Gutiérrez, S., Silla-Martínez, J.M., Gabaldón, T., 2009. trimAl: a tool for automated

alignment trimming in large-scale phylogenetic analyses. Bioinformatics 25, 1972–1973.

https://doi.org/10.1093/bioinformatics/btp348

Castelle, C.J., Brown, C.T., Thomas, B.C., Williams, K.H., Banfield, J.F., 2017. Unusual

respiratory capacity and nitrogen metabolism in a Parcubacterium (OD1) of the Candidate

Phyla Radiation. Sci. Rep. 7, 40101. https://doi.org/10.1038/srep40101

Chaumeil, P.-A., Mussig, A.J., Hugenholtz, P., Parks, D.H., 2020. GTDB-Tk: a toolkit to

classify genomes with the Genome Taxonomy Database. Bioinformatics 36, 1925–1927.

https://doi.org/10.1093/bioinformatics/btz848

Chu, R.K., Goldman, A.E., Brooks, S.C., Danczak, R.E., Garayburu-Caruso, V.A., Graham,

E.B., Jones, M., Jones, N., Lin, X., Morad, J.W., Ren, H., Renteria, L., Resch, C.T., Tfaily,

M., Tolic, N., Toyoda, J.G., Wells, J.R., Stegen, J.C., 2019. WHONDRS 48 Hour Diel

Cycling Study at the East Fork Poplar Creek in Tennessee, USA.

https://doi.org/10.15485/1577278

Cole, J.J., Prairie, Y.T., Caraco, N.F., McDowell, W.H., Tranvik, L.J., Striegl, R.G., Duarte,

C.M., Kortelainen, P., Downing, J.A., Middelburg, J.J., Melack, J., 2007. Plumbing the

global carbon cycle: Integrating inland waters into the terrestrial carbon budget. Ecosystems

10, 171–184. https://doi.org/10.1007/s10021-006-9013-8

Curran, C.A., Grossman, E.E., Magirl, C.S., Foreman, J.R., 2016. Suspended Sediment Delivery

to Puget Sound from the Lower Nisqually River, Western Washington, July 2010–

November 2011. US Geol. Surv. Sci. Investig. Rep. 2016-5062 17.

http://dx.doi.org/10.3133/sir20165062

Danczak, R.E., Chu, R.K., Fansler, S.J., Goldman, A.E., Graham, E.B., Tfaily, M.M., Toyoda, J.,

Stegen, J.C., 2020. Using metacommunity ecology to understand environmental

metabolomes. Nat. Commun. 11, 6369. https://doi.org/10.1038/s41467-020-19989-y

Danczak, R.E., Garayburu-Caruso, V.A., Renteria, L., McKever, S.A., Otenburg, O.C., Grieger,

S.R., Son, K., Kaufman, M.H., Fulton, S.G., Roebuck, J.A., Myers-Pigg, A.N., Stegen, J.C.,

2023. Riverine organic matter functional diversity increases with catchment size. Front.

Water 5, 1087108. https://doi.org/10.3389/frwa.2023.1087108

Danczak, R.E., Goldman, A.E., Chu, R.K., Garayburu-Caruso, V.A., Graham, E.B., He, X., Lin,

X., Meile, C., Morad, J.W., Ren, H., Renteria, L., Resch, C.T., Rooze, J., Schalles, J.,

Tfaily, M., Thomle, J., Tolic, N., Toyoda, J.G., Wells, J.R., Stegen, J.C., 2019. WHONDRS

48 Hour Diel Cycling Study at the Altamaha River in Georgia, USA.

https://doi.org/10.15485/1577263

Danczak, R.E., Goldman, A.E., Chu, R.K., Toyoda, J.G., Garayburu-Caruso, V.A., Tolić, N.,

Graham, E.B., Morad, J.W., Renteria, L., Wells, J.R., Herzog, S.P., Ward, A.S., Stegen,

J.C., 2021. Ecological theory applied to environmental metabolomes reveals compositional

divergence despite conserved molecular properties. Sci. Total Environ. 788, 147409.

https://doi.org/10.1016/j.scitotenv.2021.147409

Danczak, R.E., Sawyer, A.H., Williams, K.H., Stegen, J.C., Hobson, C., Wilkins, M.J., 2016.

Seasonal hyporheic dynamics control coupled microbiology and geochemistry in Colorado

River sediments. J. Geophys. Res. Biogeosciences 121, 2976–2987.

https://doi.org/10.1002/2016JG003527

D'Andrilli, J., Cooper, W.T., Foreman, C.M., Marshall, A.G., 2015. An ultrahigh-resolution

mass spectrometry index to estimate natural organic matter lability. Rapid Commun. Mass

Spectrom. 29, 2385–2401. https://doi.org/10.1002/rcm.7400

Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O., Whitwham, A.,

Keane, T., McCarthy, S.A., Davies, R.M., Li, H., 2021. Twelve years of SAMtools and

BCFtools. GigaScience 10, giab008. https://doi.org/10.1093/gigascience/giab008

Demers, J.D., Blum, J.D., Brooks, S.C., Donovan, P.M., Riscassi, A.L., Miller, C.L., Zheng, W.,

Gu, B., 2018. Hg isotopes reveal in-stream processing and legacy inputs in East Fork Poplar

Creek, Oak Ridge, Tennessee, USA. Environ. Sci. Process. Impacts 20, 686–707.

https://doi.org/10.1039/C7EM00538E

Drake, T.W., Raymond, P.A., Spencer, R.G.M., 2018. Terrestrial carbon inputs to inland waters:

A current synthesis of estimates and uncertainty. Limnol. Oceanogr. Lett. 3, 132–142.

https://doi.org/10.1002/lol2.10055

Edgar, R.C., 2004. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. Nucleic Acids Res. 32, 1792–1797. https://doi.org/10.1093/nar/gkh340

Finn, R.D., Coggill, P., Eberhardt, R.Y., Eddy, S.R., Mistry, J., Mitchell, A.L., Potter, S.C.,

Punta, M., Qureshi, M., Sangrador-Vegas, A., Salazar, G.A., Tate, J., Bateman, A., 2016.

The Pfam protein families database: towards a more sustainable future. Nucleic Acids Res.

44, D279–D285. https://doi.org/10.1093/nar/gkv1344

Fudyma, J.D., Chu, R.K., Graf Grachet, N., Stegen, J.C., Tfaily, M.M., 2021. Coupled Biotic-

Abiotic Processes Control Biogeochemical Cycling of Dissolved Organic Matter in the

Columbia River Hyporheic Zone. Front. Water 2, 1–20.

https://doi.org/10.3389/frwa.2020.574692

Garayburu-Caruso, V.A., Goldman, A.E., Chu, R.K., Danczak, R.E., Graham, E.B., Lin, X.,

Morad, J.W., Ren, H., Renteria, L., Resch, C.T., Tfaily, M., Tolic, N., Toyoda, J.G., Wells,

J.R., Stegen, J.C., 2019. WHONDRS 48 Hour Diel Cycling Study at the Columbia River in

Washington, USA. https://doi.org/10.15485/1577265

Garayburu-Caruso, V.A., Stegen, J.C., Song, H.-S., Renteria, L., Wells, J., Garcia, W., Resch,

C.T., Goldman, A.E., Chu, R.K., Toyoda, J., Graham, E.B., 2020. Carbon Limitation Leads

to Thermodynamic Regulation of Aerobic Metabolism. Environ. Sci. Technol. Lett. 7, 517–

524. https://doi.org/10.1021/acs.estlett.0c00258

Goldman, A.E., Arnon, S., Bar-Zeev, E., Chu, R.K., Danczak, R.E., Garayburu-Caruso, V.A.,

Graham, E.B., Lin, X., Morad, J.W., Ren, H., Renteria, L., Resch, C.T., Tfaily, M., Tolic,

N., Toyoda, J.G., Wells, J.R., Stegen, J.C., 2019. WHONDRS 48 Hour Diel Cycling Study

at the Jordan River, Israel. https://doi.org/10.15485/1577266

Gómez-Gener, L., Rocher-Ros, G., Battin, T., Cohen, M.J., Dalmagro, H.J., Dinsmore, K.J.,

Drake, T.W., Duvert, C., Enrich-Prast, A., Horgby, Å., Johnson, M.S., Kirk, L., Machado-

Silva, F., Marzolf, N.S., McDowell, M.J., McDowell, W.H., Miettinen, H., Ojala, A.K.,

Peter, H., Pumpanen, J., Ran, L., Riveros-Iregui, D.A., Santos, I.R., Six, J., Stanley, E.H.,

Wallin, M.B., White, S.A., Sponseller, R.A., 2021. Global carbon dioxide efflux from rivers

enhanced by high nocturnal emissions. Nat. Geosci. 14, 289–294.

https://doi.org/10.1038/s41561-021-00722-3

Graham, E.B., Crump, A.R., Kennedy, D.W., Arntzen, E., Fansler, S., Purvine, S.O., Nicora,

C.D., Nelson, W., Tfaily, M.M., Stegen, J.C., 2018. Multi 'omics comparison reveals

metabolome biochemistry, not microbiome composition or gene expression, corresponds to

elevated biogeochemical function in the hyporheic zone. Sci. Total Environ. 642, 742–753.

https://doi.org/10.1016/j.scitotenv.2018.05.256

Graham, E.B., Knelman, J.E., Schindlbacher, A., Siciliano, S., Breulmann, M., Yannarell, A.,

Beman, J.M., Abell, G., Philippot, L., Prosser, J., Foulquier, A., Yuste, J.C., Glanville,

H.C., Jones, D.L., Angel, R., Salminen, J., Newton, R.J., Bürgmann, H., Ingram, L.J.,

Hamer, U., Siljanen, H.M.P., Peltoniemi, K., Potthast, K., Bañeras, L., Hartmann, M.,

Banerjee, S., Yu, R.Q., Nogaro, G., Richter, A., Koranda, M., Castle, S.C., Goberna, M.,

Song, B., Chatterjee, A., Nunes, O.C., Lopes, A.R., Cao, Y., Kaisermann, A., Hallin, S.,

Strickland, M.S., Garcia-Pausas, J., Barba, J., Kang, H., Isobe, K., Papaspyrou, S.,

Pastorelli, R., Lagomarsino, A., Lindström, E.S., Basiliko, N., Nemergut, D.R., 2016.

Microbes as engines of ecosystem function: When does community structure enhance

predictions of ecosystem processes? Front. Microbiol. 7, 1–10.

https://doi.org/10.3389/fmicb.2016.00214

Grunert, B.K., Tzortziou, M., Neale, P., Menendez, A., Hernes, P., 2021. DOM degradation by

light and microbes along the Yukon River-coastal ocean continuum. Sci. Rep. 11, 10236.

https://doi.org/10.1038/s41598-021-89327-9

Harrell, F.E., 2023. Hmisc: Harrell Miscellaneous.

Hernández, F., 2020. Ecological Discord and the Importance of Scale in Scientific Inquiry. J.

Wildl. Manag. 84, 1427–1434. https://doi.org/10.1002/jwmg.21942

Hu, A., Jang, K., Meng, F., Stegen, J., Tanentzap, A.J., Choi, M., Lennon, J.T., Soininen, J.,

Wang, J., 2022. Microbial and Environmental Processes Shape the Link between Organic

Matter Functional Traits and Composition. Environ. Sci. Technol. 56, 10504–10516.

https://doi.org/10.1021/acs.est.2c01432

Hughey, C.A., Hendrickson, C.L., Rodgers, R.P., Marshall, A.G., Qian, K., 2001. Kendrick Mass

Defect Spectrum: A Compact Visual Analysis for Ultrahigh-Resolution Broadband Mass

Spectra. Anal. Chem. 73, 4676–4681. https://doi.org/10.1021/ac010560w

Isobe, K., Bouskill, N.J., Brodie, E.L., Sudderth, E.A., Martiny, J.B.H., 2020. Phylogenetic

conservation of soil bacterial responses to simulated global changes. Philos. Trans. R. Soc.

B Biol. Sci. 375, 20190242. https://doi.org/10.1098/rstb.2019.0242

Joshi, N., Fass, J., 2011. Sickle: A sliding-window, adaptive, quality-based trimming tool for

FastQ files.

Kaiser, K., Canedo-Oropeza, M., McMahon, R., Amon, R.M.W., 2017. Origins and

transformations of dissolved organic matter in large Arctic rivers. Sci. Rep. 7, 13064.

https://doi.org/10.1038/s41598-017-12729-1

Kanehisa, M., Sato, Y., Kawashima, M., Furumichi, M., Tanabe, M., 2016. KEGG as a reference

resource for gene and protein annotation. Nucleic Acids Res. 44, D457–D462.

https://doi.org/10.1093/nar/gkv1070

Kang, D.D., Froula, J., Egan, R., Wang, Z., 2015. MetaBAT, an efficient tool for accurately

reconstructing single genomes from complex microbial communities. PeerJ 3, e1165.

https://doi.org/10.7717/peerj.1165

Kassambara, A., 2023. ggpubr: "ggplot2" Based Publication Ready Plots.

Koch, B.P., Dittmar, T., 2006. From mass to structure: an aromaticity index for high-resolution

mass data of natural organic matter. Rapid Commun. Mass Spectrom. 20, 926–932.

https://doi.org/10.1002/rcm.2386

Kujawinski, E.B., Behn, M.D., 2006. Automated analysis of electrospray ionization fourier

transform ion cyclotron resonance mass spectra of natural organic matter. Anal. Chem. 78,

4363–4373. https://doi.org/10.1021/ac0600306

Ladau, J., Eloe-Fadrosh, E.A., 2019. Spatial, Temporal, and Phylogenetic Scales of Microbial

Ecology. Trends Microbiol. 27, 662–669. https://doi.org/10.1016/j.tim.2019.03.003

LaRowe, D.E., Van Cappellen, P., 2011. Degradation of natural organic matter: A

thermodynamic analysis. Geochim. Cosmochim. Acta 75, 2030–2042.

https://doi.org/10.1016/j.gca.2011.01.020

Levin, S.A., 1992. The Problem of Pattern and Scale in Ecology: The Robert H. MacArthur

Award Lecture. Ecology 73, 1943–1967. https://doi.org/10.2307/1941447

Li, D., Liu, C.-M., Luo, R., Sadakane, K., Lam, T.-W., 2015. MEGAHIT: an ultra-fast single-

node solution for large and complex metagenomics assembly via succinct de Bruijn graph.

Bioinformatics 31, 1674–1676. https://doi.org/10.1093/bioinformatics/btv033

Liu, Q., Jiang, Y., Tian, Y., Hou, Z., He, K., Fu, L., Xu, H., 2019. Impact of land use on the

DOM composition in different seasons in a subtropical river flowing through a region

undergoing rapid urbanization. J. Clean. Prod. 212, 1224–1231.

https://doi.org/10.1016/j.jclepro.2018.12.030

Loar, J.M., Stewart, A.J., Smith, J.G., 2011. Twenty-five years of ecological recovery of east

fork poplar creek: Review of environmental problems and remedial actions. Environ.

Manage. 47, 1010–1020. https://doi.org/10.1007/s00267-011-9625-4

Lv, J., Zhang, S., Wang, S., Luo, L., Cao, D., Christie, P., 2016. Molecular-Scale Investigation

with ESI-FT-ICR-MS on Fractionation of Dissolved Organic Matter Induced by Adsorption

on Iron Oxyhydroxides. Environ. Sci. Technol. 50, 2328–2336.

https://doi.org/10.1021/acs.est.5b04996

Martiny, A.C., Treseder, K., Pusch, G., 2013. Phylogenetic conservatism of functional traits in

microorganisms. ISME J. 7, 830–838. https://doi.org/10.1038/ismej.2012.160

Minor, E.C., Steinbring, C.J., Longnecker, K., Kujawinski, E.B., 2012. Characterization of

dissolved organic matter in Lake Superior and its watershed using ultrahigh resolution mass

spectrometry. Org. Geochem. 43, 1–11. https://doi.org/10.1016/j.orggeochem.2011.11.007

Mueller, B.M., Schulz, H., Danczak, R.E., Putschew, A., Lewandowski, J., 2021. Simultaneous

attenuation of trace organics and change in organic matter composition in the hyporheic

zone of urban streams. Sci. Rep. 11, 4179. https://doi.org/10.1038/s41598-021-83750-8

Nurk, S., Meleshko, D., Korobeynikov, A., Pevzner, P.A., 2017. metaSPAdes: a new versatile

metagenomic assembler. Genome Res. 27, 824–834. https://doi.org/10.1101/gr.213959.116

Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.R.,

O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H., 2019.

vegan: Community Ecology Package [WWW Document]. URL https://cran.r-

project.org/package=vegan

Olm, M.R., Brown, C.T., Brooks, B., Banfield, J.F., 2017. dRep: a tool for fast and accurate

genomic comparisons that enables improved genome recovery from metagenomes through

de-replication. ISME J. 11, 2864–2868. https://doi.org/10.1038/ismej.2017.126

Osterholz, H., Singer, G., Wemheuer, B., Daniel, R., Simon, M., Niggemann, J., Dittmar, T.,

2016. Deciphering associations between dissolved organic molecules and bacterial

communities in a pelagic marine system. ISME J. 10, 1717–1730.

https://doi.org/10.1038/ismej.2015.231

Parks, D.H., Imelfort, M., Skennerton, C.T., Hugenholtz, P., Tyson, G.W., 2015. CheckM:

assessing the quality of microbial genomes recovered from isolates, single cells, and

metagenomes. Genome Res. 25, 1043–1055. https://doi.org/10.1101/gr.186072.114

Peng, Y.Y., Leung, H.C.M., Yiu, S.M., Chin, F.Y.L., 2012. IDBA-UD: a de novo assembler for

single-cell and metagenomic sequencing data with highly uneven depth. Bioinformatics 28,

1420–1428. https://doi.org/10.1093/bioinformatics/bts174

Posit team, 2023. RStudio: Integrated Development Environment for R. Posit Software, PBC,

Boston, MA.

Price, M.N., Dehal, P.S., Arkin, A.P., 2010. FastTree 2 – Approximately Maximum-Likelihood

Trees for Large Alignments. PLoS ONE 5, e9490.

https://doi.org/10.1371/journal.pone.0009490

R Core Team, 2023. R: A Language and Environment for Statistical Computing. R Foundation

for Statistical Computing, Vienna, Austria.

Rawlings, N.D., Barrett, A.J., Thomas, P.D., Huang, X., Bateman, A., Finn, R.D., 2018. The

MEROPS database of proteolytic enzymes, their substrates and inhibitors in 2017 and a

comparison with peptidases in the PANTHER database. Nucleic Acids Res. 46, D624–

D632. https://doi.org/10.1093/nar/gkx1134

Regnier, P., Friedlingstein, P., Ciais, P., Mackenzie, F.T., Gruber, N., Janssens, I.A., Laruelle,

G.G., Lauerwald, R., Luyssaert, S., Andersson, A.J., Arndt, S., Arnosti, C., Borges, A.V.,

Dale, A.W., Gallego-Sala, A., Goddéris, Y., Goossens, N., Hartmann, J., Heinze, C., Ilyina,

T., Joos, F., LaRowe, D.E., Leifeld, J., Meysman, F.J.R., Munhoven, G., Raymond, P.A.,

Spahni, R., Suntharalingam, P., Thullner, M., 2013. Anthropogenic perturbation of the

carbon fluxes from land to ocean. Nat. Geosci. 6, 597–607.

https://doi.org/10.1038/ngeo1830

Renteria, L., Goldman, A.E., Chu, R.K., Danczak, R.E., Garayburu-Caruso, V.A., Graham, E.B.,

Lin, X., Morad, J.W., Ren, H., Resch, C.T., Tfaily, M., Tolic, N., Toyoda, J.G., Wells, J.R.,

Znotinas, K.R., Stegen, J.C., 2019. WHONDRS 48 Hour Diel Cycling Study at the

Nisqually River, WA. https://doi.org/10.15485/1576995

Roebuck, J.A., Seidel, M., Dittmar, T., Jaffé, R., 2020. Controls of Land Use and the River

Continuum Concept on Dissolved Organic Matter Composition in an Anthropogenically

Disturbed Subtropical Watershed. Environ. Sci. Technol. 54, 195–206.

https://doi.org/10.1021/acs.est.9b04605

Saarela, T., Zhu, X., Jäntti, H., Ohashi, M., Ide, J., Siljanen, H., Pesonen, A., Aaltonen, H.,

Ojala, A., Nishimura, H., Kekäläinen, T., Jänis, J., Berninger, F., Pumpanen, J., 2022.

Dissolved organic matter composition regulates microbial degradation and carbon dioxide

production in pristine subarctic rivers (preprint). Biogeochemistry: Rivers & Streams.

https://doi.org/10.5194/bg-2022-225

Shi, W., Sun, M.-Y., Molina, M., Hodson, R.E., 2001. Variability in the distribution of lipid

biomarkers and their molecular isotopic composition in Altamaha estuarine sediments:

implications for the relative contribution of organic matter from various sources. Org.

Geochem. 32, 453–467. https://doi.org/10.1016/S0146-6380(00)00189-3

Slowikowski, K., 2023. ggrepel: Automatically Position Non-Overlapping Text Labels with

"ggplot2."

Song, H.-S., Stegen, J.C., Graham, E.B., Lee, J.-Y., Garayburu-Caruso, V.A., Nelson, W.C.,

Chen, X., Moulton, J.D., Scheibe, T.D., 2020. Representing Organic Matter

Thermodynamics in Biogeochemical Reactions via Substrate-Explicit Modeling. Front.

Microbiol. 11, 531756. https://doi.org/10.3389/fmicb.2020.531756

Stegen, J.C., Fredrickson, J.K., Wilkins, M.J., Konopka, A.E., Nelson, W.C., Arntzen, E.V.,

Chrisler, W.B., Chu, R.K., Danczak, R.E., Fansler, S.J., Kennedy, D.W., Resch, C.T.,

Tfaily, M., 2016a. Groundwater–surface water mixing shifts ecological assembly processes

and stimulates organic carbon turnover. Nat. Commun. 7, 11237.

https://doi.org/10.1038/ncomms11237

Stegen, J.C., Goldman, A.E., 2018. WHONDRS: a Community Resource for Studying Dynamic

River Corridors. mSystems 3. https://doi.org/10.1128/mSystems.00151-18

Stegen, J.C., Goldman, A.E., Chu, R.K., Danczak, R.E., Garayburu-Caruso, V.A., Graham, E.B.,

Lin, X., Morad, J.W., Ren, H., Renteria, L., Resch, C.T., Tfaily, M., Tolic, N., Toyoda, J.,

Wells, J., Herzog, S., Wondzell, S.M., 2019. WHONDRS 48 Hour Diel Cycling Study at HJ

Andrews Experimental Forest Watershed 1 (WS1). https://doi.org/10.15485/1509695

Stegen, J.C., Johnson, T., Fredrickson, J.K., Wilkins, M.J., Konopka, A.E., Nelson, W.C.,

Arntzen, E.V., Chrisler, W.B., Chu, R.K., Fansler, S.J., Graham, E.B., Kennedy, D.W.,

Resch, C.T., Tfaily, M., Zachara, J., 2018. Influences of organic carbon speciation on

hyporheic corridor biogeochemistry and microbial ecology. Nat. Commun. 9, 585.

https://doi.org/10.1038/s41467-018-02922-9

Stegen, J.C., Konopka, A., McKinley, J.P., Murray, C., Lin, X., Miller, M.D., Kennedy, D.W.,

Miller, E.A., Resch, C.T., Fredrickson, J.K., 2016b. Coupling among Microbial

Communities, Biogeochemistry, and Mineralogy across Biogeochemical Facies. Sci. Rep.

6, 1–14. https://doi.org/10.1038/srep30553

Stegen, J.C., Lin, X., Fredrickson, J.K., Chen, X., Kennedy, D.W., Murray, C.J., Rockhold,

M.L., Konopka, A., 2013. Quantifying community assembly processes and identifying

features that impose them. ISME J. 7, 2069–79. https://doi.org/10.1038/ismej.2013.93

Stegen, J.C., Lin, X., Fredrickson, J.K.J.K., Konopka, A.E., 2015. Estimating and mapping

ecological processes influencing microbial community assembly. Front. Microbiol. 6, 1–15.

https://doi.org/10.3389/fmicb.2015.00370

Stegen, J.C., Lin, X., Konopka, A.E., Fredrickson, J.K., 2012. Stochastic and deterministic

assembly processes in subsurface microbial communities. ISME J. 6, 1653–1664.

https://doi.org/10.1038/ismej.2012.22

Swenson, N.G., Enquist, B.J., Pither, J., Thompson, J., Zimmerman, J.K., 2006. The problem and

promise of scale dependency in community phylogenetics. Ecology 87, 2418–2424.

https://doi.org/10.1890/0012-9658(2006)87[2418:TPAPOS]2.0.CO;2

Takagi, K.K., Hunter, K.S., Cai, W.-J., Joye, S.B., 2017. Agents of change and temporal nutrient

dynamics in the Altamaha River Watershed. Ecosphere 8, e01519.

https://doi.org/10.1002/ecs2.1519

Tanentzap, A.J., Fitch, A., Orland, C., Emilson, E.J.S., Yakimovich, K.M., Osterholz, H.,

Dittmar, T., 2019. Chemical and microbial diversity covary in fresh water to influence

ecosystem functioning. Proc. Natl. Acad. Sci. U. S. A. 116, 24689–24695.

https://doi.org/10.1073/pnas.1904896116

Tfaily, M.M., Chu, R.K., Tolić, N., Roscioli, K.M., Anderton, C.R., Paša-Tolić, L., Robinson,

E.W., Hess, N.J., 2015. Advanced solvent based methods for molecular characterization of

soil organic matter by high-resolution mass spectrometry. Anal. Chem. 87, 5206–5215.

https://doi.org/10.1021/acs.analchem.5b00116

Tfaily, M.M., Chu, R.K., Toyoda, J., Tolić, N., Robinson, E.W., Paša-Tolić, L., Hess, N.J., 2017.

Sequential extraction protocol for organic matter from soils and sediments using high

resolution mass spectrometry. Anal. Chim. Acta 972, 54–61.

https://doi.org/10.1016/j.aca.2017.03.031

Tolić, N., Liu, Y., Liyu, A., Shen, Y., Tfaily, M.M., Kujawinski, E.B., Longnecker, K., Kuo,

L.J., Robinson, E.W., Paša-Tolić, L., Hess, N.J., 2017. Formularity: Software for

Automated Formula Assignment of Natural and Other Organic Matter from Ultrahigh-

Resolution Mass Spectra. Anal. Chem. 89, 12659–12665.

https://doi.org/10.1021/acs.analchem.7b03318

Vaughn, D.R., Kellerman, A.M., Wickland, K.P., Striegl, R.G., Podgorski, D.C., Hawkings, J.R.,

Nienhuis, J.H., Dornblaser, M.M., Stets, E.G., Spencer, R.G.M., 2023. Anthropogenic

landcover impacts fluvial dissolved organic matter composition in the Upper Mississippi

River Basin. Biogeochemistry 164, 117–141. https://doi.org/10.1007/s10533-021-00852-1

Vigneron, A., Cruaud, P., Guyoneaud, R., Goñi-Urriza, M., 2023. Into the darkness of the

microbial dark matter in situ activities through expression profiles of Patescibacteria

populations. Front. Microbiol. 13, 1073483. https://doi.org/10.3389/fmicb.2022.1073483

Villa, J.A., Smith, G.J., Ju, Y., Renteria, L., Angle, J.C., Arntzen, E., Harding, S.F., Ren, H.,

Chen, X., Sawyer, A.H., Graham, E.B., Stegen, J.C., Wrighton, K.C., Bohrer, G., 2020.

Methane and nitrous oxide porewater concentrations and surface fluxes of a regulated river.

Sci. Total Environ. 715. https://doi.org/10.1016/j.scitotenv.2020.136920

Walkup, J., Dang, C., Mau, R.L., Hayer, M., Schwartz, E., Stone, B.W., Hofmockel, K.S., Koch,

B.J., Purcell, A.M., Pett-Ridge, J., Wang, C., Hungate, B.A., Morrissey, E.M., 2023. The

predictive power of phylogeny on growth rates in soil bacterial communities. ISME

Commun. 3, 73. https://doi.org/10.1038/s43705-023-00281-1

Ward, A.S., Schmadel, N.M., Wondzell, S.M., 2018. Time-Variable Transit Time Distributions

in the Hyporheic Zone of a Headwater Mountain Stream. Water Resour. Res. 54, 2017–

2036. https://doi.org/10.1002/2017WR021502

Wells, J.R., Goldman, A.E., Chu, R.K., Danczak, R.E., Garayburu-Caruso, V.A., Graham, E.B.,

Lewandowski, J., Lin, X., Meinikmann, K., Morad, J.W., Müller, B.M., Ren, H., Renteria,

L., Resch, C.T., Schulz, H., Tfaily, M., Tolic, N., Toyoda, J.G., Stegen, J.C., 2019.

WHONDRS 48 Hour Diel Cycling Study at the Erpe River, Germany.

https://doi.org/10.15485/1577260

Weyhenmeyer, G.A., Hartmann, J., Hessen, D.O., Kopáček, J., Hejzlar, J., Jacquet, S., Hamilton,

S.K., Verburg, P., Leach, T.H., Schmid, M., Flaim, G., Nõges, T., Nõges, P., Wentzky,

V.C., Rogora, M., Rusak, J.A., Kosten, S., Paterson, A.M., Teubner, K., Higgins, S.N.,

Lawrence, G., Kangur, K., Kokorite, I., Cerasino, L., Funk, C., Harvey, R., Moatar, F., De

Wit, H.A., Zechmeister, T., 2019. Widespread diminishing anthropogenic effects on

calcium in freshwaters. Sci. Rep. 9, 10450. https://doi.org/10.1038/s41598-019-46838-w

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G.,

Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T., Miller, E., Bache, S., Müller, K.,

Ooms, J., Robinson, D., Seidel, D., Spinu, V., Takahashi, K., Vaughan, D., Wilke, C., Woo,

K., Yutani, H., 2019. Welcome to the Tidyverse. J. Open Source Softw. 4, 1686.

https://doi.org/10.21105/joss.01686

Wohl, E., Hall, R.O., Lininger, K.B., Sutfin, N.A., Walters, D.M., 2017. Carbon dynamics of

river corridors and the effects of human alterations. Ecol. Monogr. 87, 379–409.

https://doi.org/10.1002/ecm.1261

Wondzell, S.M., 2006. Effect of morphology and discharge on hyporheic exchange flows in two

small streams in the Cascade Mountains of Oregon, USA. Hydrol. Process. 20, 267–287.

https://doi.org/10.1002/hyp.5902

Wondzell, S.M., Gooseff, M.N., McGlynn, B.L., 2007. Flow velocity and the hydrologic

behavior of streams during baseflow. Geophys. Res. Lett. 34, L24404.

https://doi.org/10.1029/2007GL031256

Woodcroft, B.J., n.d. CoverM.

Xenopoulos, M.A., Barnes, R.T., Boodoo, K.S., Butman, D., Catalán, N., D'Amario, S.C.,

Fasching, C., Kothawala, D.N., Pisani, O., Solomon, C.T., Spencer, R.G.M., Williams, C.J.,

Wilson, H.F., 2021. How humans alter dissolved organic matter composition in freshwater:

relevance for the Earth's biogeochemistry. Biogeochemistry 154, 323–348.

https://doi.org/10.1007/s10533-021-00753-3

Yin, Y., Mao, X., Yang, J., Chen, X., Mao, F., Xu, Y., 2012. dbCAN: a web resource for

automated carbohydrate-active enzyme annotation. Nucleic Acids Res. 40, W445–W451.

https://doi.org/10.1093/nar/gks479

Yu, G., Smith, D.K., Zhu, H., Guan, Y., Lam, T.T.-Y.Y., 2017. ggtree: an R package for

visualization and annotation of phylogenetic trees with their covariates and other associated

data. Methods Ecol. Evol. 8, 28–36. https://doi.org/10.1111/2041-210X.12628

Zarnetske, J.P., Bouda, M., Abbott, B.W., Saiers, J., Raymond, P.A., 2018. Generality of

Hydrologic Transport Limitation of Watershed Organic Carbon Flux Across Ecoregions of

the United States. Geophys. Res. Lett. 45, 11,702-11,711.

https://doi.org/10.1029/2018GL080005

Zhou, Y., Zhu, Y., Zhu, J., Li, C., Chen, G., 2023. A Comprehensive Review on Wastewater

Nitrogen Removal and Its Recovery Processes. Int. J. Environ. Res. Public. Health 20,

3429. https://doi.org/10.3390/ijerph20043429

1021

Acknowledgments

Pacific Northwest National Laboratory is operated by Battelle Memorial Institute for the U.S.

Department of Energy under Contract No. DE-AC05-76RL01830. This research was supported by

the U.S. Department of Energy (DOE), Office of Biological and Environmental Research (BER),

as part of BER's Subsurface Biogeochemistry Research Program (SBR). This contribution

originates from the SBR Scientific Focus Area (SFA) at the Pacific Northwest National Laboratory

(PNNL). A portion of this work was performed at the Environmental Molecular Science

Laboratory User Facility.

1030

Author Contributions

R.E.D., A.E.G., and J.C.S. conceptualized the study. V.A.G., J.W.M., L.R., J.R.H, M.J., N.J.,

B.M.M., H.S., J.S. collected samples. V.A.G., J.W.M., L.R., J.R.H analyzed anions/cations. R.K.C

and J.G.T. collected FTICR-MS data and assisted with analyses. M.A.B. performed bioinformatic

analyses. R.E.D. aggregated data, performed ecological and statistical analyses, and analyzed

microbial functional potential. R.E.D. drafted the manuscript but all authors contributed to the

writing.

1038

1039 **Competing Interests**

1040 The authors declare no competing financial interests.

1041

Correspondence

Correspondence can be directed at robert.danczak@pnnl.gov

1044

Figure and Table Legends

Figure 1: Overview of sampling locations and dissolved organic matter (DOM) characteristics. a)
Map of sampling sites. b) Sampling efforts across the seven rivers. Solid circles indicate that all
samples were analyzed using the corresponding method, while empty circles indicate a subset of
samples was analyzed; no circle indicates no analysis. c) Boxplot of non-purgeable organic carbon
(NPOC) concentrations for each river. d) Boxplot of DOM molecular characteristics.

Figure 2: a) Non-metric multidimensional scaling (NMDS) ordination depicting the multivariate relationships of DOM across all seven rivers. b) A redundancy analysis (RDA) optimized to evaluate to what degree the geochemistry is related to DOM composition. Loading arrows indicate geochemical variables significantly related to DOM.

1098

1099
1100 **Figure 3**: Null modeling results. a) Density plots illustrating the distribution of raw pairwise βNTI
1101 values from each River. b) Average βNTI values calculated for pairwise comparisons within the
1102 surface water (top panel) or within pore water (bottom panel).

1129
1130 **Figure 4**: Microbial relationships. a) An rpsC phylogenetic tree generated using FastTree. Points
1131 on the tree are genomes obtained from this study and are colored based upon the river from which
1132 they were observed. b) Non-metric multidimensional scaling (NMDS) ordination indicating the
1133 beta-diversity of each sample. Loadings indicate microbial orders significantly related to the
1134 ordination structure.

1141
Figure 5: CAZyme analysis. a) Alpha-diversity analysis illustrating the CAZymes pools vary across microbial Orders. b) Non-metric multidimensional scaling (NMDS) ordination of the CAZyme pools illustrating multivariate dynamics across microbial Orders. c) Three different microbially derived diversity metrics plotted against molecular richness.

River	Abbreviation	Location	River Type	pH	NPOC (mg/L)
Altamaha River	AR	Georgia, United States	Large River	6 (0)	5.98 (1.34)
Columbia River	CR	Washington, United States	Large River	7.97 (0.3)	1.38 (0.32)
East Fork Poplar Creek	EFPC	Tennessee, United States	Wastewater	7.73 (0.37)	2.60 (0.40)
Erpe River	ER	Berlin, Germany	Wastewater	6.16 (0.37)	9.5 (1.39)
HJ Andrews Watershed 1	HJA	Oregon, United States	Headwater	7.79 (0.33)	0.57 (0.06)
Jordan River	JR	Israel	Headwater	7.72 (0.26)	0.57 (0.18)
Nisqually River	NR	Washington, United States	Headwater	7.50 (0.25)	0.06 (0.11)

1149 **Table 1**: Summary information for each river. Average pH and non-purgeable organic carbon are
1150 provided as contextual information (standard deviations are in parentheticals).
1151

Supplemental Information

1153
1154 **Supplemental Figure 1:** Boxplots displaying variation in geochemical parameters in each river,
1155 separated by river type. River abbreviations are AR = Altamaha River, CR = Columbia River,
1156 EFPC = East Fork Poplar Creek, ER = Erpe River, HJA = HJ Andrews Watershed 1, JR = Jordan
1157 River, and NR = Nisqually River.

1158
1159 **Supplemental Figure 2:** Principal component analysis (PCA) of river and pore water
1160 geochemistry.

Supplemental Figure 3: Bar plot depicting the number of bins with a given GTDB taxonomy detected in each river.

1200

1201 **Supplemental Figure 4:** Metabolic information and abundance of broadly distributed MAGs. A
1202 heatmap showing the presence (and relative completion) of the pathway listed on the y-axis in the
1203 metagenomic assembled genome (MAG) on the x-axis. Cells in the heatmap are colored based
1204 upon the taxonomy of the MAG listed on the x-axis. Below each MAG is a visual representation
1205 of the read mapping abundance in each river with rivers indicated by the colored bars.

1206

Supplemental Figure 5: Metabolic information and abundance of MAGs found in fewer rivers
(the inverse of Supplemental Figure 4). A heatmap showing the presence (and relative completion)
of the pathway listed on the y-axis in the metagenomic assembled genome (MAG) on the x-axis.
Cells in the heatmap are colored based upon the taxonomy of the MAG listed on the x-axis. Below
each MAG is a visual representation of the read mapping abundance in each river with rivers
indicated by the colored bars.

1213
1214   **Supplemental Table 1:** Table describing the summary statistics for each metagenomic assembled
1215   genome, including (but not limited to) measurements like completion/contamination percentage,
1216   MAG size in base pairs, and N50.
1217
1218   **Supplemental Table 2**: Pairwise Mann Whitney U statistics for DOM measurements across rivers
1219   and sample type.
1220
1221   **Supplemental Table 3:** Table of significant correlation values for βNTI and geochemical
1222   relationships.
1223
1224   **Supplemental Table 4:** Table of pairwise Mann Whitney U statistics for average βNTI values
1225   across rivers.
1226