Time dependent changes in feeding behavior and energy balance associated with weight gain in mice fed obesogenic diets

Payam A. Fathi[1], Michelle B. Bales[1,†], Julio E. Ayala[1,2,3*]

[1]Department of Molecular Physiology & Biophysics, Vanderbilt University School of Medicine, Nashville, TN, 37232

[2]Vanderbilt Mouse Metabolic Phenotyping Center, Nashville, TN, 37232

[3]Vanderbilt Center for Addiction Research, Nashville, TN, 37223

[†]Current address: Department of Psychology, Florida State University, Tallahassee, FL, 32306

*Corresponding author:

2201 Garland Avenue

702 Light Hall

Nashville, TN, 37232

Email: julio.e.ayala@vanderbilt.edu

Phone: (615) 323-7252

Disclosure: The authors declare no conflict of interest

Abstract

Obesity is characterized by dysregulated homeostatic mechanisms resulting in positive energy

balance, yet when this dysregulation occurs is unknown. We assessed the time course of

alterations to behaviors promoting weight gain in male and female mice switched to obesogenic

60% or 45% high fat diet (HFD). Switching mice to obesogenic diets promotes transient bouts of

hyperphagia during the first 2 weeks followed by persistent caloric hyperphagia. Energy

expenditure increases but not sufficiently to offset increased caloric intake, resulting in a

sustained net positive energy balance. Hyperphagia is associated with consumption of

calorically larger meals (impaired satiation) more frequently (impaired satiety) particularly

during the light-cycle. Running wheel exercise delays weight gain in 60% HFD-fed male mice

by enhancing satiation and increasing energy expenditure. However, exercise effects on

satiation are no longer apparent after 2 weeks, coinciding with weight gain. Thus, exposure to

obesogenic diets engages homeostatic regulatory mechanisms for ~2 weeks that ultimately fail,

and consequent weight gain is characterized by impaired satiation and satiety. Insights into the

etiology of obesity can be obtained by investigating changes to satiation and satiety mechanisms

during the initial ~2 weeks of HFD exposure.

Introduction

Obesity has reached epidemic proportions worldwide, and since 2013, the American Medical Association has recognized obesity as a disease state (Rosen, 2014). Significant effort has been given to uncovering factors that contribute to excessive weight gain and the mechanisms that become dysregulated in the setting of obesity. Put simply, obesity results from calories being consumed and absorbed in excess of calories expended. Several factors and mechanisms have been identified that play a role in the regulation of caloric consumption and energy expenditure. These discoveries have led to a general model for the regulation of body weight characterized by feedback mechanisms to maintain homeostasis (Myers et al., 2021). In this model, excess weight gain triggers changes in several hormonal and neural factors that primarily engage the central nervous system to subsequently reduce caloric intake and increase energy expenditure, thus reducing body weight. With this framework in mind, obesity has been characterized as a state associated with defects in many of these feedback mechanisms. However, much of the evidence that these mechanisms are dysregulated has been obtained by comparing subjects (mice, rats, humans) that are lean and those that already have obesity. Little is known about events leading to obesity, such as the timing for the changes in factors that contribute to excess weight gain. Characterizing the changes in parameters that regulate body weight when rodents are provided access to obesogenic diets can provide insights into potential behaviors and mechanisms associated with excess weight gain and obesity.

Studies in rodents have uncovered several factors associated with weight gain when exposed to obesogenic diets. Early studies examining the effects of high-fat diets on weight gain in rats

showed that differences in distinct feeding behaviors influence weight gain (Drewnowski et al.,

1984). Subsequent studies demonstrated that rats prone to greater weight gain consume more

calories on a per meal basis without altering circadian meal numbers (Farley et al., 2003). A

sustained increase in per meal caloric content is observed in rats exposed to a long time course

of high-fat diet, and this is associated with an in increase in meal consumption rates without

changing daytime meal numbers (Furnes et al., 2009). Sex is also a key contributor to propensity

for weight gain, with male rodents broadly increasing overall caloric intake and relative body

weight more so than female rodents (Yang et al., 2014). Timing of exposure to obesogenic diets

also influences weight gain. Mice fed obesogenic diets only during the light phase gained more

weight compared to mice fed the same diet only during the dark phase (Arble et al., 2009).

Similarly, access to high-fat diet rapidly disrupts the circadian variability in food intake, and

mice no longer consume the majority of meals during the dark cycle (Espinosa-Carrasco et al.,

2018). Taken together, these studies suggest that factors such as meal patterns, sex, and timing

of food intake can influence weight gain associated with obesogenic diets.

While these studies provide insight into factors contributing to weight gain, specifically when

these factors become relevant during the progression towards obesity is not known. This

requires comprehensive metabolic phenotyping of longitudinal changes in male and female

mice switched from standard to obesogenic diets. Multiplexed indirect calorimetry systems

generate high-resolution, real-time datasets of multiple components that contribute to the

regulation of body weight. Using this technology, the present studies investigated the time scale

and magnitude of changes to these components during the progression towards obesity. Our

findings demonstrate that upon exposure to high-fat diets, behaviors and metabolic phenotypes

that are engaged to prevent weight gain rapidly become dysregulated. Furthermore, the

magnitude of these changes is influenced by the type of obesogenic diet (i.e., 60% vs. 45% high-

fat diet) and sex and are more apparent during specific times of the 24h cycle. These findings

provide a blueprint for investigating specific molecular mechanisms whose dysregulation

contributes to the progression towards obesity.

Experimental Procedures

Animals

Male and female mice on a C57Bl6/J background at 10-12 weeks of age generated from in-house

breeding colonies were utilized in these experiments. Experimental procedures were approved

and conducted within the guidelines set by the Institutional Animal Care and Use Committee at

Vanderbilt University.

Diet Switch and Running Wheel Access Experiments

Diet switch experiments: male or female mice (n=8 per group) were removed from our animal

colony and single-housed in Promethion Metabolic chambers (Sable Systems International) with

ad-lib access to chow (2.86 kCal/gram; PicoLab Rodent Diet 5L0D). Following a 5-7 day data

acquisition period on chow diet, mice were switched to either 60% HFD (5.24 kCal/gram;

Research Diets Inc, D12492) for 4 weeks or 45% HFD 4.73 kCal/gram; Research Diets Inc.

D12451) for 3 weeks.

Wheel running experiments: male mice (n=16) were removed from our animal colony and single

housed in Promethion Metabolic chambers with access to chow as noted previously. All mice

had access to locked running wheels during the chow period. Mice where then switched to a

60% HFD and were randomly selected to either remain with running wheels locked (n=8) or

have the running wheels unlocked (n=8) for the duration of the 4-week study period. Mice

comprising both groups were evenly distributed throughout the Promethion cabinets to

minimize any potential impact of cage positioning within the cabinet on data acquisition.

Metabolic Measurements

Food intake, water intake, locomotor activity, respiratory exchange ratio, and energy

expenditure were continuously measured throughout the study period across five-minute

intervals. Energy balance was determined by taking the difference between energy expenditure

and intake across 12h light-dark cycles. Total energy expenditure over a given 12-hour

light:dark cycle was calculated by determining the average energy expenditure over the 12h

time interval and multiplying by 12. Energy intake was calculated by multiplying the mass of

food consumed over the corresponding 12h-intervals with the caloric density of diet consumed.

We would like to note that technical failure prevented us from capturing locomotor activity

beyond the first week in the female 60% HFD group.

Meal Parameters

Meals parameters were chosen based on previous studies in rodents.(Zorrilla et al., 2005) A

meal was defined as any removal of food from the food hopper spanning a duration of at least

30 seconds and weighing a minimum of 10 milligrams to account for inherent sensor limitations

and a maximum of 500 milligrams to account for instances of spillage. The threshold for the

intermeal interval was set at 5 minutes (i.e., a meal was defined as terminated when there was a

period of at least five minutes of no hopper activity).

Meal Type/Pattern Analysis

Meal types were defined by assembling all meal sizes consumed by male mice during the chow

feeding period in Promethion chambers across all experiments. Quartile distributions were used

to determine cut-offs for different meal types assigned from smallest (Type I) to largest (Type

IV) as follows: Type I: 10 mg – <60 mg; Type II: 60 mg – <160 mg; Type III: 160 mg – 280 mg; and

Type IV: >280 mg. Meal types were compared by determining the weekly proportion of a given

meal type within each animal across each week of diet access. Parts of whole analysis used

compiled averages of meal types across sex and diet groups. Histograms were generated using

bin sizes of 20 mg (mass analysis) or .2 kCal (calorie analysis) and curve-fit using non-linear

regression.

Statistical Analysis

Raw data were processed using the Sable Systems Macro Interpreter (v23.6), assembled and

analyzed using Microsoft Excel. Graphs were generated and data were statistically analyzed

using GraphPad Prism Software (v10). Figures were assembled using Adobe Illustrator (v28.0).

Data were statistically analyzed using Repeated Measures One-Way ANOVAs and multiple

comparisons were assessed using the Tukey Test or Two-Way ANOVAs with the Sidak multiple

comparisons tests as necessary. Significance thresholds were set to a p-value of 0.05. Data are

reported as means with the standard error of the mean unless stated otherwise.

Results

Food-intake and metabolic phenotyping of mice switched to a 60% high-fat diet

We first investigated the effect of switching mice from a chow diet to a 60% high-fat diet (60%

HFD) in male and female C57Bl/6J mice. Mice were switched from chow to 60% HFD while in

metabolic chambers and remained on the 60% HFD for 28 days. There is no appreciable increase

in the daily (**Figure 1A**) or weekly (**Figure 1B**) mass of food consumed throughout the 60% HFD

feeding period, although ~60% of male mice increased their mass intake during the first week

while the remaining ~40% actually reduced their mass intake (**Figure 1B**). Interestingly, when

analyzing the mass of food consumed by light:dark cycle, there was an increase in the daily and

average weekly mass of food consumed during the light cycle but not during the dark cycle in

male mice (**Supplemental Figure 1A-C**). Contrasting male mice, female mice decrease the mass

of food consumed throughout the 4-week exposure to 60% HFD and significantly so at weeks 2

and 3 (**Figure 1G, 1H**). This decrease in the mass of food consumed is primarily due to an effect

during the dark cycle (**Supplemental Figure 1D-F**).

When accounting for the differences in caloric density between the chow diet and the 60% HFD,

both sexes display a rapid and sustained increase in the total daily (**Figure 1C, 1I**) and weekly

(**Figure 1D, 1J**) calories consumed. This is accounted for by increased caloric consumption

during both the light and dark cycles (**Supplemental Figure 1G-L**). Analysis of daily caloric

intake reveals that after a rapid initial increase in caloric intake immediately following the

switch to the 60% HFD, there is a transient decrease during the first week of 60% HFD exposure

in both male and female mice (**Figure 1C, 1I**). This is followed by a second phase characterized

by a rapid increase in caloric intake and another attempt at lowering daily caloric intake over

the next 7 days. After these two cycles of elevated and then decreased caloric intake, there is a

sustained elevation in daily caloric intake for the remainder of the 60% HFD exposure period

(**Figure 1C, 1I**).

Switching to a 60% HFD increases daily and weekly average energy expenditure in both male

and female mice (**Figure 1E, 1F, 1K, 1L**) during both the light and dark cycles (**Supplemental**

Figure 1M-R). Interestingly, daily energy expenditure followed a similar pattern as daily caloric

intake, characterized by a rapid increase in daily energy expenditure upon switching to the 60%

HFD followed by a transient decrease, and then a sustained increase after the first two weeks of

the diet switch (**Figure 1E, 1K**). Utilizing measures of energy intake and energy expenditure we

calculated total energy balance across the diet exposure period. Male mice show a sustained

positive energy balance starting on the second week 60% HFD exposure (**Figure 1M, 1O,**

Supplemental Figure 2A), and this is associated with a significant increase in body weight

beginning on the second week of 60% HFD exposure (**Figure 1P**). Female mice also display a net

positive energy balance and weight gain upon switching to the 60% HFD, although the

magnitude of this effect is not as robust as in male mice (**Figure 1N, 1Q, 1R, Supplemental**

Figure 2B).

We assessed other behavioral and metabolic parameters in response to a switch to the 60% HFD

(**Supplemental Figure 2**). The respiratory exchange ratio significantly decreases upon exposure

to the 60% HFD in both males and females, reflecting the increased consumption of a diet

primarily composed of fat (**Supplemental Figure 2C-F**). This is observed during both the light

and dark cycles but is more robust during the dark cycle (**Supplemental Figure 1S-X**) Switching

to the 60% HFD also rapidly decreases water intake and locomotor activity in both males and

females (**Supplemental Figure 2G-N**) primarily due to changes during the dark cycle

(**Supplemental Figure 1Y-AJ**).

Meal pattern analysis of mice switched to a 60% high-fat diet

We next assessed effects of switching male and female mice to a 60% HFD on meal patterns.

Meal numbers rapidly increase in both sexes and are significantly sustained throughout the 60%

HFD feeding period (**Figure 2A-C**). This is due to increases in meal numbers during both the

light and dark cycles, with a more robust effect in male mice (**Supplemental Figure 3A-F**). The

increase in the number of meals results in decreased inter-meal interval and meal duration,

particularly in male mice (**Figure 2D-I**). The decrease in inter-meal interval is most robust

during the light cycle (**Supplemental Figure 3G-L**), whereas the decrease in meal duration is

observed during both the light and dark cycles (**Supplemental Figure 3M-R**). Upon switching

to a 60% HFD, average meal sizes by mass decrease in both male and female mice (**Figure 2J-L**),

and this is mostly accounted for by a decrease in meal sizes during the dark cycle

(**Supplemental Figure 3S-X**). However, when accounting for the caloric density of each diet, the

average meal size significantly increases in male mice after the second week of 60% HFD

exposure but does not significantly increase in female mice (**Figure 2M-O**). The increase in the

caloric content per meal observed in male mice is primarily due to an increase in meal size

during the light cycle (**Supplemental Figure 3Y-AA**). As with total caloric intake, there are

transient increases and attempts to restore meal size during the first two weeks of 60% HFD

exposure. The rate of meal consumption by either mass or caloric content increases upon

switching to a 60% HFD in both males and females, with males displaying a more robust effect

(**Figure 2P-U**). Following from other meal pattern results, these increases are primarily

accounted for by effects observed during the light cycle (**Supplemental Figure 3AE-AP**).

Food-intake and metabolic phenotyping of mice switched to a 45% high-fat diet

We next investigated the effect of switching mice to a 45% high fat diet (45% HFD) on

mechanisms and behaviors regulating body weight. When switched from chow to a 45% HFD

there is a consistent trend towards a decrease in total food intake by mass in both male and

female mice; however these differences are only significant at two weeks post diet switch

(**Figure 3B, 3H**). The decrease in mass consumed is primarily due to an effect during the dark

cycle (**Supplemental Figure 4A-F**). Accounting for the caloric value of each diet, there is a trend

towards an increase in caloric intake, although this is not significant in either sex (**Figure 3C,**

3D, 3I, 3J). This trend is evident during both light and dark cycles (**Supplemental Figure 4G-L**).

Similar to results obtained with the 60% HFD, albeit with a much lower magnitude and

duration, there is a transient hyperphagia followed by restoration to normal caloric

consumption that then gradually settles into a tendency for elevated caloric intake (**Figure 3C,**

3I).

When switched to a 45% HFD, both male and female mice rapidly increase energy expenditure

(**Figure 3E, 3F, 3K, 3L**), and this is evident during both the light and dark cycles (**Supplemental**

Figure 4M-R). Net energy balance in response to the 45% HFD switch shows a transient net

positive energy balance during the first week that is not significantly maintained in either sex

(**Figure 3M, 3N, 3O, 3Q**, **Supplemental Figure 5**). In contrast to observations made when mice

are switched to a 60% HFD, body weights significantly increase during each week of 45% HFD

diet exposure (**Figure 3P, 3R**) even though energy balance is not significantly different

compared to the chow period. The magnitude of weight gain in the 45% HFD fed mice is not as

robust as mice fed a 60% HFD suggesting that slight increases in overall energy balance are

sufficient to drive significant increases in body weight.

Similar to the observations in mice switched to a 60% HFD, switching mice to a 45% HFD

results in decreased respiratory exchange ratio and water intake in both males and females, but

there is only a significant decrease in activity in males (**Supplemental Figure 5C-N**). These

decreases are observed almost exclusively during the dark cycle (**Supplemental Figure 4S-AJ**).

Meal pattern analysis of mice switched to a 45% high-fat diet

Meal pattern analysis of mice switched to a 45% HFD shows a rapid, yet transient increase in

daily meal numbers during the first week of 45% HFD exposure followed by a slow, non-

significant increase during subsequent weeks (**Figure 4A-C**). This pattern is observed during

both light and dark cycles in male mice but primarily during the dark cycle in female mice

(**Supplemental Figure 6A-F**). There is no significant effect of switching to a 45% HFD on inter-

meal intervals in either sex (**Figure 4D-F**) and during either light or dark cycle (**Supplemental**

Figure 6G-L). Similar to observations in mice switched to a 60% HFD, average meal duration is

significantly decreased in mice of both sexes when switched to a 45% HFD (**Figure 4G-I**), and

this is observed during both the light and dark cycle (**Supplemental Figure 6M-R**). Also similar

to switching mice to a 60% HFD, meal sizes by mass decreases in both sexes during exposure to

45% HFD (**Figure 4J-L**) primarily due to an effect during the dark cycle (**Supplemental Figure**

6S-X). Accounting for the caloric density results in slight increases in the average caloric intake

per meal (**Figure 4M-O; Supplemental Figure 6Y-AD**). Meal rates by mass and caloric content

significantly increase in both male and female when switched to a 45% HFD, and these

increases are apparent during both the light and dark cycle (**Figure 4P-U, Supplemental Figure**

6AE-AP).

Meal-typing analysis of mice switched to obesogenic diets

Histograms capturing the percent frequency of meal sizes by mass and caloric content were

generated to more accurately describe changes that occur in feeding behaviors when mice are

switched to obesogenic diets. When fed a chow diet, meals of very low mass and caloric content

comprise the largest set of meals consumed (~15% of meals by mass and ~25% of meals by

calorie). The remaining meal sizes are almost equally distributed (**Figure 5A-D**). When switched

to a 60% HFD, male mice decrease the intake of meals of very low mass but also meals of larger

masses (**Figure 5A**). When accounting for the caloric density, there is a rightward shift towards a

larger proportion of meals with higher caloric content (**Figure 5B**). In female mice, switching to

60% HFD promotes a leftward shift indicating a decrease in the proportion of meals of large

mass and an increase the proportion of meals that are small in mass (**Figure 5C**). When

accounting for caloric density, there is a pronounced increase in histogram spread, indicating an

increase in the proportion of meals with larger caloric content (**Figure 5D**). When switched to a

45% HFD meal mass histograms show a consistent left-ward shift in both male and female mice,

and a corresponding flattening of histogram spread when examining the calories consumed per

meal (**Figure 5E-H**).

To further characterize the shifts in feeding behavior related to the mass of food consumed

when mice were switched to obesogenic diets, meal types were subcategorized based on the

quartile distribution of meal sizes by mass in chow-fed mice. These cut-offs yielded four meal

types: Type I (<60 mg), Type II (>60 mg – <160 mg), Type III (>160 mg < 280 mg) and Type IV

(>280 mg) (**Figure 5I**). Using these definitions of meal types we observe that when switched to

60% HFD, male mice increase the proportion of type II meals consumed while decreasing the

proportion of very small type I meals and large and very large type III and type IV meals,

respectively (**Figure 5J, 5N**). These changes are observed across both light and dark cycles

(**Supplemental Figure 7A-7D**). When switched to 60% HFD, female mice also increase the

proportion of type II meals consumed and decrease the proportion of very large type IV meals

consumed, but, unlike male mice, they also increase the proportion of very small type I meals

consumed (**Figure 5K, 5O**). These shifts are primarily observed during the dark cycle

(**Supplemental Figure 7E-7H**). Similar shifts than those observed in male and female mice fed

60% HFD are observed in mice switched to 45% HFD although the magnitude of the shift is not

significant (**Figure 5L, 5M, 5P, 5Q** and **Supplemental Figure 7I-7P**).

Metabolic phenotyping and meal patterns in mice switched to a 60% high-fat diet with

concurrent running wheel access

Switching male mice from a chow diet to a 60% HFD increases both caloric intake and energy

expenditure, yet the degree to which energy expenditure increases is not sufficient to offset the

increase in caloric intake (**Figure 1**). We investigated whether providing mice with access to a

running wheel concomitant with switching to a 60% HFD increases energy expenditure to

sufficiently offset the caloric content of the 60% HFD. Male mice were acclimated to cages with

locked running wheels during chow feeding. Mice were then switched to a 60% HFD, and

running wheel access was provided to a subset of mice at the time of the switch (Wheels

Unlocked) while the remaining mice had wheels locked through the study period (Wheels

Locked). Overall food intake was similar across both groups with similar patterns of transient

changes during the first ~2 weeks culminating in a sustained increase in daily caloric intake

(**Figure 6A-D**). Energy expenditure is increased to a greater degree in the Wheels Unlocked

group (**Figure 6E, 6F**) due exclusively to increased energy expenditure during the dark cycle

(**Supplemental Figure 8T**). This results in a lesser deviation towards positive energy balance in

mice with access to running wheels, although the net energy balance is not significantly

different between mice with and without access to running wheels (**Figure 6G, 6H,**

Supplemental Figure 8B). While the difference in energy balance is not statistically significant,

mice with running wheel access are protected against significant increases in body weight

during the first two weeks of 60% HFD exposure and overall gain significantly less weight

(**Figure 6I, Supplemental Figure 8C**). However, mice with access to running wheels do begin to

gain weight after the second week, and this is associated with increased caloric consumption

during the dark cycle (**Supplemental Figure 9A, 9B**).

Even though there were no differences in overall food intake, meal pattern analysis shows that

mice with running wheel access display a greater increase in meal number compared to mice

with wheels locked during the first week (**Figure 6J, 6K**). These increases are mainly driven by

effects during the dark cycle (**Supplemental Figure 9C**). Inter-meal interval (**Figure 6L, 6M**) and

meal duration (**Figure 6N, 6O**) decrease equally between both groups. Interestingly, access to

running wheel significantly lowers meal size by mass and calories during the first week of 60%

HFD exposure, and these differences are observed during both the light and dark cycles (**Figure**

6P-S; Supplemental Figure 9F, 9G). This difference in meal sizes is no longer evident from the

second week onwards, which is coincident with the greater deviations towards positive energy

balance and weight gain in mice with access to running wheels. Meal rates are not significantly

different between mice with and without access to running wheels (**Figure 6T-W, Supplemental**

Figure 9H, 9I). Histogram analysis of meal size by grams shows similar meal patterns across

both groups; however, the proportion of meals with a lower caloric content is higher in the mice

with running wheels blocked (**Figure 6X-AA**). Furthermore, both groups increase their

consumption of type II meals and decrease their consumption of very large type IV meals, but

mice with access to running wheels also increase the consumption of very small type I meals

and have a larger decrease in the proportion of very large type IV meals consumed during the

first week of 60% HFD exposure (**Supplemental Figure 9J-L**).

The respiratory exchange ratio is significantly lower in mice with running wheel access during

the first two weeks following diet switch (**Supplemental Figure 8D-G**). Locomotor activity (off

wheel) decreases in both groups, although mice with running wheel access have significantly

higher locomotor activity during the last two weeks of diet exposure, particularly during the

dark cycle (**Supplemental Figure 8L-O**). As expected, running wheel access results in

progressively increased wheel running activity over the first two weeks of the diet switch and

subsequently levels out by week four. Notably, daytime wheel running transiently increases

during the first two weeks (**Supplemental Figure 8P-R**).

Discussion

This study utilized metabolic phenotyping of male and female mice fed obesogenic diets to

observe time-dependent changes in parameters that influence body weight during the

progression towards obesity. Similar studies have investigated processes occurring during the

early phases of exposure to a high-fat diet. However, most of those studies are short-term (i.e.,

high-fat diet exposure for 3-7 days) and focus on effects on gross food intake as the primary

endpoint (Clyburn et al., 2023; Enriquez et al., 2022; Kless et al., 2017; Oraha et al., 2022). We

hypothesized that when placed on high-fat diets, mice would engage feedback mechanisms and

consequent behaviors to account for the increased caloric content of the diet in order to

maintain a normal body weight trajectory. Our results reveal the time course over which these

feedback processes are engaged and then ultimately fail, thus resulting in weight gain. These

important findings can be leveraged towards future studies focusing on mechanisms

characterizing the transition from attempts to maintain weight homeostasis to the failure of

these attempts. Furthermore, we provide detailed analyses of other parameters, such as meal

patterns, that we hypothesize are significant contributors to the propensity to gain weight.

When examining behavioral and metabolic differences in mice switched from chow to a 60%

HFD, a general pattern arises in which there is significant hyperphagia upon exposure to the

HFD that is rapidly followed by attempts to reduce food intake to pre-HFD exposure levels

during the first ~7 days, as previously shown.(Hwang et al., 2010) We extend these observations

to show that transient hyperphagia followed by attempts to restore normal food intake occurs

during the first ~14 days of HFD exposure. Following this ~2-week period, mice enter a period

of sustained hyperphagia. These results suggest that homeostatic mechanisms are engaged

during the first ~2 weeks of 60% HFD exposure, but they ultimately fail or are overtaken by

other contributors of feeding behavior such as hedonic mechanisms. Identifying not only the

mechanisms engaged during the first 2 weeks of exposure to a 60% HFD but also which of these

mechanisms fail or are overcome following this period will provide important information

about molecular targets that are essential for body weight control.

It is important to note that hyperphagia associated with exposure to a 60% HFD is in terms of

caloric intake but not mass consumed. Indeed, following a transient increase in the mass of food

consumed during the first 24h of 60% HFD exposure in male mice, as previously shown

(Clyburn et al., 2023; Enriquez et al., 2022), the average daily mass of food consumed gradually

decreases slightly. In female mice the average daily mass of food consumed decreases upon

switching to a 60% HFD, and this remains consistent for most of the study period. Similar

effects were observed in male and female mice switched to a 45% HFD, albeit to a lesser

magnitude. These observations suggest that mice can sense changes in diet caloric content and

adjust their intake accordingly. However, this adjustment is not sufficient to fully compensate

for the caloric density of the diet. Studies suggest that both humans and mice can perceive the

caloric density of ingested foods (Flynn et al., 2022; Hu et al., 2018; Su et al., 2017). Similar to

what we have shown, previous studies demonstrate that mice do not appreciably increase their

total food intake by mass when placed on obesogenic diets, and they actually decrease the total

mass of food consumed as the percentage of dietary fat exceeds 60% (Hu et al., 2018).

Identifying mass-sensing (e.g., mechanosensing) vs. calorie-sensing molecular mechanisms and

how these mechanisms are coupled or uncoupled could provide valuable insight into

approaches for restraining excess caloric intake in the presence of obesogenic diets.

Studies in rodents suggest that meal size by caloric content is a strong predictor of weight gain

(Furnes et al., 2009; Melhorn et al., 2010). Our findings show that meal size by calorie increases

immediately upon exposure to 60% HFD, with initial attempts to lower meal size during the

first ~2 weeks of exposure that eventually fail. This was most strongly evident in male mice

switched to a 60% HFD with a lesser effect in female mice and mice of both sexes switched to a

45% HFD. Interestingly, average meal size by mass actually tended to decrease, thus

contributing to the overall decrease in average daily mass consumed (see above). However,

analysis of meal type by mass shows that while chow-fed mice primarily eat very small meals

(what we refer to as Type I, <60 mg meals) with remaining meals being somewhat evenly

distributed by size, switching to a 60% HFD compresses this distributions by increasing the

relative consumption of what we refer to as Type II (<160 mg) meals at the expense of both very

small Type I meals and large/very large (Type III, <280 mg and IV, >280 mg) meals. When

accounting for the caloric content of the diet, this effect of exposure to a 60% HFD results in a

shift towards consumption of larger meals by calorie. This increase in meal size suggests that

switching to a HFD impairs mechanisms that regulate satiation. Switching mice to a HFD also

increased the number of meals consumed per day and, consequently, reduced the time between

meals. This suggests an impairment in mechanisms regulating satiety. Given the relationship

between meal patterns, particularly meal size, and propensity for weight gain (Furnes et al.,

2009; Melhorn et al., 2010), these findings further suggest that emphasis should be given to

investigating mechanisms regulating satiation and satiety and how these mechanisms become

dysregulated following ~2 weeks exposure to HFD.

Analysis of feeding behavior by light:dark cycle shows that, when switched to a 60% HFD, male

mice increase food intake primarily during the light cycle due to an increase in meal numbers

without compensatory changes in average meal size. The relative importance of meal timing has

been noted in multiple human and rodent studies. A study in rats showed that increases in

caloric intake were driven by inactive phase (light cycle) food consumption and removing food

access during this period blunts weight gain even though total caloric consumption remains

similar between groups (Olsen et al., 2017). Similarly mice fed a HFD during the inactive phase

(i.e., light cycle) gained more weight than those fed the same diet during the active phase (i.e.,

dark cycle) (Arble et al., 2009). Furthermore, human studies show that caloric consumption later

in the day is associated with greater weight gain (Bo et al., 2014; Jakubowicz et al., 2013;

Ruddick-Collins et al., 2022; Zhao et al., 2023). These results suggest that the increased caloric

consumption during the light phase in mice switched to HFD may contribute to weight gain in

male mice. Contrasting the findings in male mice, female mice increase meal numbers during

both cycles but reduce the mass of food consumed during the dark cycle, suggesting that female

mice are better able to recognize the caloric density of food and adjust nighttime feeding

accordingly (Goldstein et al., 2021; Su et al., 2017). Interestingly, meal duration during the light

cycle decreases in both male and female mice. This coupled with the fact that in male mice light

cycle meal numbers increased and average meal size did not change would indicate that male

mice eat larger meals (by calorie) in a shorter period. This is further reflected through a

significant increase in light cycle meal rates. Ultimately, the propensity of male mice to consume

more frequent shorter meals with higher caloric content per meal during the light cycle may be

a significant contributor to their greater weight gain. Cycle-dependent effects on food intake

and meal patterns were similar in mice switched to 45% HFD, although effects were less robust.

These smaller effects relative to those observed in mice fed 60% HFD likely contribute to the

delay in weight gain onset in mice fed a 45% HFD. In sum, changes to overall caloric intake

when fed HFD are driven by consumption of more frequent and shorter meals with higher

caloric content particularly during time spans during which significant food intake typically

does not occur in chow-fed mice (i.e., light cycle). Importantly, these changes follow a pattern

suggesting attempts to maintain homeostasis during the first two weeks of obesogenic diet

exposure that eventually fail.

Male mice, particularly of the C57Bl/6 background, are prone to weight gain on a HFD

compared to female mice. We observe that male mice display more robust caloric hyperphagia

and increases in meal size relative to female mice when switched to a HFD. Regarding meal

size, both sexes increase the relative intake of Type II meals, but this increase is comparatively

greater in male mice. This suggests that while exposure to 60% HFD impairs satiation in both

sexes, this effect is greater in male mice. A similar observation is made when examining the

impact of 45% HFD in that male mice show no significant change to the proportion of meal

types whereas females increase the proportion of very small Type I meals while decreasing the

proportion of larger Type III and IV meals. This suggests that in response to 45% HFD, satiation

is somewhat preserved in male mice but is actually enhanced in female mice. Overall, this

further suggests that the propensity towards weight gain in response to HFD feeding is

associated with impairments in both satiety and satiation that depend on the dietary fat content

and differ according to sex. These sex differences can be leveraged to investigate potential

mechanisms regulating meal patterns. For example, previous studies suggest that sex

differences in feeding behavior may due to steroid hormone signaling (Alonso-Caraballo and

Ferrario, 2019; Lenert et al., 2021; Petersen, 1976). Additionally, differences in energy

expenditure may also contribute to sex-dependent differences on weight gain (see below).

Our observed changes to satiety and satiation fall in line with the idea that obesogenic diet

exposure rapidly impairs: 1- the latency or ability to appropriately convey hunger signals

(satiety) and 2- the fidelity of- or ability to process signals that would appropriately account for

both mass and caloric density to regulate meal size (satiation). Obesogenic diets have been

shown to impact satiety by impairing vagal afferent responsiveness to distension as well as

change gastric emptying (Kentish et al., 2015; Loper et al., 2021). Previous studies also show

decreased responsiveness to cholecystokinin (CCK)-induced satiation in male mice fed a 45%

HFD (Huang et al., 2020). Importantly, HFD impairs vagal afferent circadian rhythms (Kentish

et al., 2018, 2016), and this may contribute in part to our observed increases in daytime food

intake. Interventions used to increase intestinal glucagon-like peptide-1 (GLP-1) levels have

been associated with decreased inactive phase food-intake in mice in a vagal GLP-1 receptor-

mediated manner, presumably through modulating meal termination (Borgmann et al., 2021;

Iwasaki et al., 2018). Thus, future studies should address whether defects in these mechanisms

occur in a time scale corresponding to the changes in feeding behavior we demonstrate in the

present studies.

Concomitant with an increase in caloric consumption, energy expenditure also increases in both

male and female mice when switched to a 60% HFD. This increase in energy expenditure

further adds to the notion that mice can sense the difference in the caloric content of HFD

relative to chow and attempt to compensate for this, much like the reduction in mass of food

consumed. Similarly to the latter, however, the increase in energy expenditure is not sufficient

to compensate for the increased caloric intake. The net result of this is that following the first ~2

weeks of exposure to 60% HFD, mice display a relative and consistent positive daily energy

balance of ~5 kCal/day for males and ~3 kCal/day for females. Interestingly, energy expenditure

also increases when mice are switched to a 45% HFD, and the magnitude of this increase is

similar to what is observed in mice switched to a 60% HFD. This, combined with a lesser degree

of caloric hyperphagia (see above) results in smaller deviations towards positive energy

balance, thus contributing to the reduced weight gain associated with 45% HFD relative to 60%

HFD exposure. Male and female mice experience quantitatively similar increases in energy

expenditure. Oraha et al. and Huang et al. (Huang et al., 2020; Oraha et al., 2022) suggest that

protection from weight gain in female relative to male mice is heavily driven by relative

increases in energy expenditure and the ability of female mice to better utilize fat. Our data

support this by showing lower relative respiratory exchange ratio in female compared to male

mice switched to a 60% HFD. When switched to a 45% HFD both males and females lower

respective respiratory exchange ratio to similar levels. This contrasts findings from Huang et al.

study (Huang et al., 2020) that used the same 45% HFD but suggest that female mice rely more

on fat oxidation. These discrepancies may stem from methodological differences since in our

studies, mice were switched to the 45% HFD while obtaining metabolic chamber measurements

whereas the studies of Huang et al. performed metabolic chamber measurements after mice had

been on the 45% HFD for 5 weeks. Nevertheless, we find that when exposed to a 45% HFD, the

propensity towards weight gain in 45% HFD male mice compared to female mice is associated

with changes in specific aspects of feeding behaviors in conjunction with alterations in energy

balance. This is in agreement with recent studies showing that female mice balance energy

expenditure and intake more efficiently than do male mice (Morris et al., 2020).

Energy expenditure increased in response to HFD exposure even as locomotor activity

decreased, as previously observed (Bjursell et al., 2008; Kless et al., 2017). Given the dramatic

sustained decrease in locomotor activity concurrent with switching to a HFD, we hypothesized

that this constrains energy expenditure and likely contributes to the observation that energy

expenditure does not increase sufficiently to account for increased caloric intake. A previous

study showed that decreased locomotor activity in mice fed a 40% HFD accounted for over 60%

of the bodyweight gain in these mice (Bjursell et al., 2008). We examined whether wheel

running access would increase energy expenditure sufficiently to modify weight gain

trajectories. Wheel running access provides a relative protection against weight gain in male

mice fed a 60% HFD (Cho et al., 2023), as we also show here. Our results show that the shortfall

between energy expenditure and caloric intake is partially mitigated in male mice given access

to a running wheel concomitant with exposure to a 60% HFD, and this is associated with a

greater increase in energy expenditure. However, this only leads to a delay in weight gain for

the first ~2 weeks of exposure to the HFD. This resistance to weight gain during early time

points of HFD exposure in the wheel running group was associated with specific changes in

feeding behaviors. While total energy intake was not different between the groups, wheel

running mice significantly increased meal numbers during the dark cycle with a corresponding

decrease in meal sizes across both the light and dark cycle. This suggests that wheel running

enhances satiation at the expense of satiety. In further support of this, mice with running wheel

access had an increased proportion of Type I very small meals. Following the first 2 weeks,

caloric intake was actually increased in mice with access to a running wheel during the dark

cycle relative to mice with no wheel access. This partially offset the significant increase in

energy expenditure observed in mice given access to running wheels and likely contributed to

the initiation of weight gain in this group. Meal patterns became indistinguishable between

mice with and without access to running wheels, and this corresponded to the period during

which mice with access to running wheels were no longer protected from weight gain. This

further emphasizes the potential important role of meal size as a contributor to weight gain in

the setting of consuming obesogenic diets. Untangling the mechanisms mediating the satiation

may prove useful in identifying molecular targets that can be leveraged to modulate weight

gain by controlling feeding behaviors.

Access to a running wheel is associated with subsequent decreases in off wheel locomotor

activity in mice fed a chow diet, suggesting the presence of a counterregulatory mechanism that

serves to stabilize the exercise driven increase in energy expenditure (Lark et al., 2018).

Similarly, the increase in energy expenditure resulting from wheel running plateaus after one

week even though on-wheel activity increases in mice fed a chow diet and decreases over time

in mice fed a HFD (O'Neal et al., 2017; Vellers et al., 2017). We expand on this by showing that

when switched to a 60% HFD, male mice rapidly increase energy expenditure resulting from an

increase in wheel running activity exclusively during the dark cycle that peaks by week 2. While

off-wheel locomotor activity decreased in mice with or without access to a running wheel, mice

with access to a running wheel displayed greater off-wheel locomotor activity, particularly

during the dark cycle, beginning around week 2 of the HFD exposure. Interestingly, this

coincided with a slight decrease in wheel running activity. The compensation of off-wheel

locomotor activity acts to preserve the increase in energy expenditure.

In sum, our study provides insights into how time-dependent changes in feeding behaviors

following obesogenic diet exposure can modify weight gain. Specifically, we found that

following exposure to HFD, mice engage mechanisms in an attempt to maintain homeostasis

and prevent weight gain during the first two weeks of diet exposure, but these mechanisms

eventually give way to the cumulative compounding effect of small increases in positive energy

balance and, thus, weight gain. Importantly, this positive energy balance is associated with

increases in the average calories consumed per meal. Wheel running activity increases energy

expenditure and prevents an increase in meal size, but this effect on meal size is temporary and

is only sufficient to delay but not prevent weight gain on obesogenic diets. These findings

provide valuable insight for future studies focusing on mechanisms that regulate satiation as

important regulators of weight gain and in particular highlight the value of investigating how

these mechanisms are affected during the transition from initial attempts to maintain neutral

energy balance to the eventual development of consistent positive energy balance.

Author Contributions

Conceptualization: JEA, MBB, PF; Methodology: JEA, MBB, PF; Analysis: JEA, PF; Writing: JEA,

PF; Funding: JEA. All authors reviewed and have approved the final version of this manuscript.

Acknowledgements and Funding

We would like to thank Dr. Louise Lantier and Merrygay James for their assistance. Promethion

metabolic chamber studies were conducted within the Vanderbilt Mouse Metabolic

Phenotyping Center (VMMPC) which is supported by NIH grant U2C0595637, shared

instrument grant S10OD028455 (JEA). This work was funded by NIH grants F31-DK127728

(PF), T32-DK07563 (PF, MBB), and R01-DK132852 (JEA).

References

Alonso-Caraballo Y, Ferrario CR. 2019. Effects of the estrous cycle and ovarian hormones on

cue-triggered motivation and intrinsic excitability of medium spiny neurons in the Nucleus

Accumbens core of female rats. *Horm Behav* **116**:104583. doi:10.1016/j.yhbeh.2019.104583

Arble DM, Bass J, Laposky AD, Vitaterna MH, Turek FW. 2009. Circadian timing of food intake

contributes to weight gain. *Obesity (Silver Spring)* **17**:2100–2. doi:10.1038/oby.2009.264

Bjursell M, Gerdin A-K, Lelliott CJ, Egecioglu E, Elmgren A, Törnell J, Oscarsson J, Bohlooly-Y

M. 2008. Acutely reduced locomotor activity is a major contributor to Western diet-induced

obesity in mice. *Am J Physiol Endocrinol Metab* **294**:E251-60. doi:10.1152/ajpendo.00401.2007

Bo S, Musso G, Beccuti G, Fadda M, Fedele D, Gambino R, Gentile L, Durazzo M, Ghigo E,

Cassader M. 2014. Consuming more of daily caloric intake at dinner predisposes to obesity.

A 6-year population-based prospective cohort study. *PLoS One* **9**:e108467.

doi:10.1371/journal.pone.0108467

Borgmann D, Ciglieri E, Biglari N, Brandt C, Cremer AL, Backes H, Tittgemeyer M, Wunderlich

FT, Brüning JC, Fenselau H. 2021. Gut-brain communication by distinct sensory neurons

differently controls feeding and glucose metabolism. *Cell Metab* **33**:1466-1482.e7.

doi:10.1016/j.cmet.2021.05.002

Cho C, Ji M, Cho E, Yi S, Kim JG, Lee S. 2023. Chronic voluntary wheel running exercise

ameliorates metabolic dysfunction via PGC-1α expression independently of FNDC5/irisin

pathway in high fat diet-induced obese mice. *J Physiol Sci* **73**:6. doi:10.1186/s12576-023-

00864-6

Clyburn C, Carson KE, Smith CR, Travagli RA, Browning KN. 2023. Brainstem astrocytes

control homeostatic regulation of caloric intake. *J Physiol* **601**:801–829. doi:10.1113/JP283566

Drewnowski A, Cohen AE, Faust IM, Grinker JA. 1984. Meal-taking behavior is related to

predisposition to dietary obesity in the rat. *Physiol Behav* **32**:61–67. doi:10.1016/0031-

9384(84)90071-4

Enriquez JR, McCauley HA, Zhang KX, Sanchez JG, Kalin GT, Lang RA, Wells JM. 2022. A

dietary change to a high-fat diet initiates a rapid adaptation of the intestine. *Cell Rep*

41:111641. doi:10.1016/j.celrep.2022.111641

Espinosa-Carrasco J, Burokas A, Fructuoso M, Erb I, Martín-García E, Gutiérrez-Martos M,

Notredame C, Maldonado R, Dierssen M. 2018. Time-course and dynamics of obesity-

related behavioral changes induced by energy-dense foods in mice. *Addiction Biology*

23:531–543. doi:10.1111/adb.12595

Farley C, Cook JA, Spar BD, Austin TM, Kowalski TJ. 2003. Meal pattern analysis of diet-

induced obesity in susceptible and resistant rats. *Obes Res* **11**:845–851.

doi:10.1038/OBY.2003.116

Flynn AN, Hall KD, Courville AB, Rogers PJ, Brunstrom JM. 2022. Time to revisit the passive

overconsumption hypothesis? Humans show sensitivity to calories in energy-rich meals.

Am J Clin Nutr **116**:581–588. doi:10.1093/ajcn/nqac112

Furnes MW, Zhao C-M, Chen D. 2009. Development of obesity is associated with increased

calories per meal rather than per day. A study of high-fat diet-induced obesity in young

rats. *Obes Surg* **19**:1430–8. doi:10.1007/s11695-009-9863-1

Goldstein N, McKnight AD, Carty JRE, Arnold M, Betley JN, Alhadeff AL. 2021. Hypothalamic

detection of macronutrients via multiple gut-brain pathways. *Cell Metab* **33**:676-687.e5.

doi:10.1016/J.CMET.2020.12.018

Hu S, Wang L, Yang D, Li L, Togo J, Wu Y, Liu Q, Li B, Li M, Wang G, Zhang X, Niu C, Li J, Xu

Y, Couper E, Whittington-Davies A, Mazidi M, Luo L, Wang S, Douglas A, Speakman JR.

2018. Dietary Fat, but Not Protein or Carbohydrate, Regulates Energy Intake and Causes

Adiposity in Mice. *Cell Metab* **28**:415-431.e4. doi:10.1016/j.cmet.2018.06.010

Huang K-P, Ronveaux CC, Knotts TA, Rutkowsky JR, Ramsey JJ, Raybould HE. 2020. Sex

differences in response to short-term high fat diet in mice. *Physiol Behav* **221**:112894.

doi:10.1016/j.physbeh.2020.112894

Hwang L-L, Wang C-H, Li T-L, Chang S-D, Lin L-C, Chen C-P, Chen C-T, Liang K-C, Ho I-K,

Yang W-S, Chiou L-C. 2010. Sex Differences in High-fat Diet-induced Obesity, Metabolic

Alterations and Learning, and Synaptic Plasticity Deficits in Mice. *Obesity* **18**:463–469.

doi:10.1038/oby.2009.273

Iwasaki Y, Sendo M, Dezaki K, Hira T, Sato T, Nakata M, Goswami C, Aoki R, Arai T, Kumari P,

Hayakawa M, Masuda C, Okada T, Hara H, Drucker DJ, Yamada Y, Tokuda M, Yada T.

2018. GLP-1 release and vagal afferent activation mediate the beneficial metabolic and

chronotherapeutic effects of D-allulose. *Nat Commun* **9**:113. doi:10.1038/s41467-017-02488-y

Jakubowicz D, Barnea M, Wainstein J, Froy O. 2013. High caloric intake at breakfast vs. dinner

differentially influences weight loss of overweight and obese women. *Obesity (Silver Spring)*

21:2504–12. doi:10.1002/oby.20460

Kentish SJ, Frisby CL, Kritas S, Li H, Hatzinikolas G, O'Donnell TA, Wittert GA, Page AJ. 2015.

TRPV1 Channels and Gastric Vagal Afferent Signalling in Lean and High Fat Diet Induced

Obese Mice. *PLoS One* **10**:e0135892. doi:10.1371/journal.pone.0135892

Kentish SJ, Hatzinikolas G, Li H, Frisby CL, Wittert GA, Page AJ. 2018. Time-Restricted Feeding

Prevents Ablation of Diurnal Rhythms in Gastric Vagal Afferent Mechanosensitivity

Observed in High-Fat Diet-Induced Obese Mice. *J Neurosci* **38**:5088–5095.

doi:10.1523/JNEUROSCI.0052-18.2018

Kentish SJ, Vincent AD, Kennaway DJ, Wittert GA, Page AJ. 2016. High-Fat Diet-Induced

Obesity Ablates Gastric Vagal Afferent Circadian Rhythms. *J Neurosci* **36**:3199–207.

doi:10.1523/JNEUROSCI.2710-15.2016

Kless C, Rink N, Rozman J, Klingenspor M. 2017. Proximate causes for diet-induced obesity in

laboratory mice: a case study. *Eur J Clin Nutr* **71**:306–317. doi:10.1038/ejcn.2016.243

Lark DS, Kwan JR, McClatchey PM, James MN, James FD, Lighton JRB, Lantier L, Wasserman

DH. 2018. Reduced Nonexercise Activity Attenuates Negative Energy Balance in Mice

Engaged in Voluntary Exercise. *Diabetes* **67**:831–840. doi:10.2337/db17-1293

Lenert ME, Chaparro MM, Burton MD. 2021. Homeostatic Regulation of Estrus Cycle of Young

Female Mice on Western Diet. *J Endocr Soc* **5**. doi:10.1210/jendso/bvab010

Loper H, Leinen M, Bassoff L, Sample J, Romero-Ortega M, Gustafson KJ, Taylor DM, Schiefer

MA. 2021. Both high fat and high carbohydrate diets impair vagus nerve signaling of

satiety. *Sci Rep* **11**:10394. doi:10.1038/s41598-021-89465-0

Melhorn SJ, Krause EG, Scott KA, Mooney MR, Johnson JD, Woods SC, Sakai RR. 2010. Acute

exposure to a high-fat diet alters meal patterns and body composition. *Physiol Behav* **99**:33–

9. doi:10.1016/j.physbeh.2009.10.004

Morris EM, Noland RD, Allen JA, McCoin CS, Xia Q, Koestler DC, Shook RP, Lighton JRB,

Christianson JA, Thyfault JP. 2020. Difference in Housing Temperature-Induced Energy

Expenditure Elicits Sex-Specific Diet-Induced Metabolic Adaptations in Mice. *Obesity*

(Silver Spring) **28**:1922–1931. doi:10.1002/oby.22925

Myers MG, Affinati AH, Richardson N, Schwartz MW. 2021. Central nervous system regulation

of organismal energy and glucose homeostasis. *Nat Metab* **3**:737–750. doi:10.1038/s42255-

021-00408-5

Olsen MK, Choi MH, Kulseng B, Zhao C-M, Chen D. 2017. Time-restricted feeding on weekdays

restricts weight gain: A study using rat models of high-fat diet-induced obesity. *Physiol*

Behav **173**:298–304. doi:10.1016/j.physbeh.2017.02.032

O'Neal TJ, Friend DM, Guo J, Hall KD, Kravitz A V. 2017. Increases in Physical Activity Result

in Diminishing Increments in Daily Energy Expenditure in Mice. *Curr Biol* **27**:423–430.

doi:10.1016/j.cub.2016.12.009

Oraha J, Enriquez RF, Herzog H, Lee NJ. 2022. Sex-specific changes in metabolism during the

transition from chow to high-fat diet feeding are abolished in response to dieting in

C57BL/6J mice. *Int J Obes (Lond)* **46**:1749–1758. doi:10.1038/s41366-022-01174-4

Petersen S. 1976. The temporal pattern of feeding over the oestrous cycle of the mouse. *Anim*

Behav **24**:939–955. doi:10.1016/S0003-3472(76)80023-1

Rosen H. 2014. Is Obesity A Disease or A Behavior Abnormality? Did the AMA Get It Right? *Mo*

Med **111**:104–108.

Ruddick-Collins LC, Morgan PJ, Fyfe CL, Filipe JAN, Horgan GW, Westerterp KR, Johnston JD,

Johnstone AM. 2022. Timing of daily calorie loading affects appetite and hunger responses

without changes in energy metabolism in healthy subjects with obesity. *Cell Metab* **34**:1472-

1485.e6. doi:10.1016/j.cmet.2022.08.001

Su Z, Alhadeff AL, Betley JN. 2017. Nutritive, Post-ingestive Signals Are the Primary Regulators

of AgRP Neuron Activity. *Cell Rep* **21**:2724–2736. doi:10.1016/j.celrep.2017.11.036

Vellers HL, Letsinger AC, Walker NR, Granados JZ, Lightfoot JT. 2017. High Fat High Sugar

Diet Reduces Voluntary Wheel Running in Mice Independent of Sex Hormone

Involvement. *Front Physiol* **8**:628. doi:10.3389/fphys.2017.00628

Yang Y, Smith DL, Keating KD, Allison DB, Nagy TR. 2014. Variations in body weight, food

intake and body composition after long-term high-fat diet feeding in C57BL/6J mice.

Obesity **22**:2147–2155. doi:10.1002/OBY.20811

Zhao D, Guallar E, Woolf TB, Martin L, Lehmann H, Coughlin J, Holzhauer K, Goheer AA,

McTigue KM, Lent MR, Hawkins M, Clark JM, Bennett WL. 2023. Association of Eating

and Sleeping Intervals With Weight Change Over Time: The Daily24 Cohort. *J Am Heart*

Assoc **12**:e026484. doi:10.1161/JAHA.122.026484

Zorrilla EP, Inoue K, Fekete EM, Tabarin A, Valdez GR, Koob GF. 2005. Measuring meals:

structure of prandial food and water intake of rats. *Am J Physiol Regul Integr Comp Physiol*

288:R1450-67. doi:10.1152/ajpregu.00175.2004

Figure Legends

Figure 1. Food intake and energy balance in mice fed a 60% high-fat diet

Mice were placed in metabolic chambers and were fed with chow for up to 6 days. Mice were

then switched to 60% HFD at day 0 (dashed line) and remained on 60% HFD for 28 days. (A-F,

M, O, P) represent data from male mice (black) and (G-L, N, Q, R) represent data from female

mice (pink). (A, G) Daily food intake by mass (grams). (B, H) Average weekly food intake by

mass. (C, I) Daily caloric intake. (D, J) Average weekly caloric intake. (E, K) Daily energy

expenditure. (F, L) Average weekly energy expenditure. (M, N) Heat map showing shifts in net

daily energy balance. (O, Q) Average weekly energy balance. (P, R) Average weekly body

weight. Data are shown as mean ± SEM for N=8 per sex and were statistically analyzed using a

repeated measures one-way ANOVA with Tukey multiple comparison tests. P-value definitions:

not-significant (ns) >0.05, * <0.05, **<0.01, ***<0.001, **** <0.00001.

Figure 2. Feeding behaviors in mice switched to a 60% high-fat diet

Feeding behavior analysis in male mice (black) and female mice (pink) switched from chow to

60% HFD in metabolic chambers at day 0 (dashed line). (A) Daily meal numbers. (B) Average

weekly meal number in male and (C) female mice. (D) Daily intermeal intervals. (E) Average

weekly intermeal intervals in male and (F) female mice. G) Daily meal duration. (H) Average

weekly meal duration in male and (I) female mice. (J) Daily meal size by mass. (K) Average

weekly meal size by mass in male and (L) female mice. (M) Daily meal size by calorie. (N)

Average weekly meal size by calorie in male and (O) female mice. (P) Daily meal rate in grams

consumed per minute. (Q) Average weekly meal rate in grams consumed per minute in male

and (R) female mice. (S) Daily meal rate in calories consumed per minute. (T) Average weekly

meal rate in calories consumed per minute in male and (U) female mice. Data are shown as

mean ± SEM for N=8 per sex and were statistically analyzed using a repeated measures one-way

ANOVA with Tukey multiple comparison tests. P-value definitions: not-significant (ns) >0.05, *

<0.05, **<0.01, ***<0.001, **** <0.00001.

Figure 3. Food intake and energy balance in male and female mice switched to a 45% high-fat

diet

Mice were placed in metabolic chambers and were fed with chow for 5 days. Mice were then

switched to 45% HFD at day 0 (dashed line) and remained on 45% HFD for 21 days. (A-F, M, O,

P) represent data from male mice (black) and (G-L, N, Q, R) represent data from female mice

(pink). (A, G) Daily food intake by mass (grams). (B, H) Average weekly food intake by mass.

(C, I) Daily caloric intake. (D, J) Average weekly caloric intake. (E, K) Daily energy expenditure.

(F, L) Average weekly energy expenditure. (M, N) Heat map showing shifts in net daily energy

balance. (O, Q) Average weekly energy balance. (P, R) Average weekly body weight. Data are

shown as mean ± SEM for N=8 per sex and were statistically analyzed using a repeated

measures one-way ANOVA with Tukey multiple comparison tests. P-value definitions: not-

significant (ns) >0.05, * <0.05, **<0.01, ***<0.001, **** <0.00001.

Figure 4. Feeding behaviors in mice switched to a 45% high-fat diet

Feeding behavior analysis in male mice (black) and female mice (pink) switched from chow to

45% HFD in metabolic chambers at day 0 (dashed line). (A) Daily meal numbers. (B) Average

weekly meal number in male and (C) female mice. (D) Daily intermeal intervals. (E) Average

weekly intermeal intervals in male and (F) female mice. G) Daily meal duration. (H) Average

weekly meal duration in male and (I) female mice. (J) Daily meal size by mass. (K) Average

weekly meal size by mass in male and (L) female mice. (M) Daily meal size by calorie. (N)

Average weekly meal size by calorie in male and (O) female mice. (P) Daily meal rate in grams

consumed per minute. (Q) Average weekly meal rate in grams consumed per minute in male

and (R) female mice. (S) Daily meal rate in calories consumed per minute. (T) Average weekly

meal rate in calories consumed per minute in male and (U) female mice. Data are shown as

mean ± SEM for N=8 per sex and were statistically analyzed using a repeated measures one-way

ANOVA with Tukey multiple comparison tests. P-value definitions: not-significant (ns) >0.05, *

<0.05, **<0.01, ***<0.001, **** <0.00001.

Figure 5. Meal patterns of male and female mice fed obesogenic diets

Meal size analysis in male and female mice switched from chow to 60% HFD (A-D, J-K, N-O)

for 4 weeks or 45% HFD (E-H, L-M, P-Q) for 3 weeks. Histograms of meal size by (A) grams and

(B) calories in male and (C) grams and (D) calories in female mice switched to a 60% HFD.

Histograms of meal size by (E) grams and (F) calories in male and (G) grams and (H) calories in

female mice switched to a 45% HFD. Histogram bin sizes of 0.02 (range: 0-0.5) for meals by

grams and bin size of 0.2 (range: 0-3) for meals by calories. Non-linear regression was used to fit

curves on each graph. (I) Range of meal sizes in grams consumed by mice (n=16) during the

chow feeding period. Quartiles were used to determine cut-offs for meal types. Type 1: <60 mg,

Type2: <160 mg, Type 3: <280 mg, and Type 4: >280 mg. Pie-charts depicting meal types across

diet switch week in (J) males and (K) females switched from chow to a 60% HFD and (L) males

and (M) females switched from chow to a 45% HFD. Percent frequency of meals by type across

each week of feeding in (N) males and (O) females switched from chow to a 60% HFD and (P)

males and (Q) females switched from chow to a 45% HFD. Data are shown as mean ± SEM for

N=8 per sex and were statistically analyzed using a repeated measures one-way ANOVA with

Tukey multiple comparison tests. P-value definitions: not-significant (ns) >0.05, * <0.05, **<0.01,

<0.001, * <0.00001.

Figure 6. Effects of running wheel access on feeding behavior and energy expenditure in

male mice switched to a 60% HFD

Male mice were placed in metabolic chambers with chow and locked running wheels for 5 days.

Mice were then switched to 60% HFD on day 0 (dashed line) and concomitantly remained with

locked wheels (wheels locked, black) or the running wheels were unlocked (wheels unlocked,

red) for 28 days. (A) Daily and (B) average weekly food intake in grams. (C) Daily and D)

average weekly caloric intake. (E) Daily and (F) average weekly energy expenditure. (G) Heat

map depicting net energy balance for each mouse across time after being switched from normal

chow to 60% HFD with locked or unlocked running wheels. (H) Average net daily energy

balance across each week of diet. (I) Average body weight across each week of diet. Feeding

behavior analysis depicting (J) daily and (K) average weekly average meal number, (L) daily

and (M) average weekly intermeal interval, N) daily and O) average weekly meal duration, (P)

daily and (Q) average weekly meal size in grams, (R) daily and (S) average weekly meal size in

calories, (T) daily and (U) average weekly meal rates in grams per minute, and (V) daily and

(W) average weekly meal rate in calories per minute. Meal size histograms by gram for (X) mice

with wheels locked and (Y) mice with wheels unlocked in bin sizes of 0.02 (Range 0-0.5). Meal

size histograms by calories for (Z) mice with wheels locked and (AA) mice with wheels

unlocked in bin sizes of 0.2 (Range 0-3). Non-linear regression was used to fit curves on each

graph. Data are shown as mean ± SEM for N=8 per group and were statistically analyzed using

a two-way ANOVA with Sidaks multiple comparison tests. P-value definitions: not-significant

(ns) >0.05, * <0.05, **<0.01, ***<0.001, **** <0.00001.

Supplemental Figure 1. Metabolic phenotyping of mice switched to a 60% high-fat diet

separated by light:dark cycle.

Metabolic phenotyping parameters in male and female mice switched from chow to 60% HFD

on day 0 (dashed lines) in metabolic chambers and divided by light (blue) and dark (beige)

cycles. (A) Daily food intake in grams in male and (D) female mice. Average weekly food intake

by mass in the light (B) and dark (C) cycle in male mice and in the light (E) and dark (F) cycle in

female mice. (G) Daily caloric intake in male and (J) female mice. Average weekly caloric intake

in the light (H) and dark (I) cycle in male mice and in the light (K) and dark (L) cycle in female

mice. (M) Daily energy expenditure in male and (P) female mice. Average weekly energy

expenditure in the light (N) and dark (O) cycles in male mice and in the light (Q) and dark (R)

cycle in female mice. (S) Daily respiratory exchange ratio in male and (V) female mice. Average

weekly respiratory exchange ratio in the light (T) and dark (U) cycle in male mice and in the

light (W) and dark (X) cycle in female mice. (Y) Daily water intake in male and (AB) female

mice. Average weekly water intake in the light (Z) and dark (AA) cycle in male mice and in the

light (AC) and dark (AD) cycle in female mice. (AE) Daily locomotor activity in male and (AH)

female mice. Average weekly locomotor activity in the light (AF) and dark (AG) cycle in male

mice and in the light (AI) and dark (AJ) cycle in female mice. Locomotor activity data are only

available for first seven days of diet switch in female mice due to instrument malfunction for

this feature. Data are shown as mean ± SEM for $N=8$ per sex and were statistically analyzed

using a repeated measures one-way ANOVA with Tukey multiple comparison tests. P-value

definitions: not-significant (ns) >0.05, * <0.05, **<0.01, ***<0.001, **** <0.00001.

Supplemental Figure 2. Metabolic phenotyping of mice switched to a 60% high-fat diet

Metabolic parameters in male (black) and female (pink) mice fed a chow diet and switched to

60% HFD at day 0 (dashed line) for 28 days. Average daily energy balance plots showing energy

intake (Energy In, circle), energy expenditure (Energy Out, square), and net energy balance

(triangle). (C) Daily and (D) average weekly respiratory exchange ratio in male and (E, F) female

mice. G) Daily and (H) average weekly water intake in male and (I, J) female mice. (K) Daily and

(L) average weekly locomotor activity in male and (M, N) female mice. Locomotor activity data

are only available for first seven days of diet switch in female mice due to instrument

malfunction for this feature. Data are shown as mean ± SEM for N=8 per sex and were

statistically analyzed using a repeated measures one-way ANOVA with Tukey multiple

comparison tests. P-value definitions: not-significant (ns) >0.05, * <0.05, **<0.01, ***<0.001, ****

<0.00001.

Supplemental Figure 3. Feeding behaviors in mice switched to a 60% high-fat diet separated

by light cycle.

Feeding behavior analysis in male and female mice switched from chow to 60% HFD on day 0

(dashed lines) in metabolic chambers and divided by light (blue) and dark (beige) cycles. (A)

Daily and (B, C) average weekly meal number in male mice in the light and dark cycle. (D)

Daily and (E, F) average weekly meal number in female mice in the light and dark cycle. (G)

Daily and (H, I) average weekly intermeal interval in male mice in the light and dark cycle. (J)

Daily and (K, L) average weekly intermeal interval in female mice in the light and dark cycle.

(M) Daily and (N, O) average weekly meal duration in male mice in the light and dark cycle. (P)

Daily and (Q, R) average weekly meal duration in female mice in the light and dark cycle. (S)

Daily and (T, U) average weekly meal size by mass in male mice in the light and dark cycle. (V)

Daily and (W, X) average weekly meal size by mass in female mice in the light and dark cycle.

(Y). Daily and (Z, AA) average weekly meal size by calorie in male mice in the light and dark

cycle. (AB) Daily and (AC, AD) average weekly meal size by calorie in female mice in the light

and dark cycle. (AE) Daily and (AF, AG) average weekly meal rate in grams per minute in male

mice in the light and dark cycle. (AH) Daily and (AI, AJ) average weekly meal rate in grams per

minute in female mice in the light and dark cycle. (AK) Daily and (AL, AM) average weekly

meal rate in calories per minute in male mice in the light and dark cycle. (AN) Daily and (AO,

AP) average weekly meal rate in calories per minute in female mice in the light and dark cycle.

Data are shown as mean ± SEM for N=8 per sex and were statistically analyzed using repeated

measures one-way ANOVA with Tukey multiple comparison tests. P-value definitions: not-

significant (ns) >0.05, * <0.05, **<0.01, ***<0.001, **** <0.00001.

Supplemental Figure 4. Metabolic phenotyping of mice switched to a 45% high-fat diet

separated by light:dark cycle.

Metabolic phenotyping parameters in male and female mice switched from chow to 45% HFD

on day 0 (dashed lines) in metabolic chambers and divided by light (blue) and dark (beige)

cycles. (A) Daily food intake in grams in male and (D) female mice. Average weekly food intake

by mass in the light (B) and dark (C) cycle in male mice and in the light (E) and dark (F) cycle in

female mice. (G) Daily caloric intake in male and (J) female mice. Average weekly caloric intake

in the light (H) and dark (I) cycle in male mice and in the light (K) and dark (L) cycle in female

mice. (M) Daily energy expenditure in male and (P) female mice. Average weekly energy

expenditure in the light (N) and dark (O) cycles in male mice and in the light (Q) and dark (R)

cycle in female mice. (S) Daily respiratory exchange ratio in male and (V) female mice. Average

weekly respiratory exchange ratio in the light (T) and dark (U) cycle in male mice and in the

light (W) and dark (X) cycle in female mice. (Y) Daily water intake in male and (AB) female

mice. Average weekly water intake in the light (Z) and dark (AA) cycle in male mice and in the

light (AC) and dark (AD) cycle in female mice. (AE) Daily locomotor activity in male and (AH)

female mice. Average weekly locomotor activity in the light (AF) and dark (AG) cycle in male

mice and in the light (AI) and dark (AJ) cycle in female mice. Data are shown as mean ± SEM for

N=8 per sex and were statistically analyzed using a repeated measures one-way ANOVA with

Tukey multiple comparison tests. P-value definitions: not-significant (ns) >0.05, * <0.05, **<0.01,

<0.001, * <0.00001.

Supplemental Figure 5. Metabolic phenotyping of mice switched to a 45% high-fat diet

Metabolic parameters in male (black) and female (pink) mice fed a chow diet and switched to

45% HFD at day 0 (dashed line) for 21 days. Average daily energy balance plots showing energy

intake (Energy In, circle), energy expenditure (Energy Out, square), and net energy balance

(triangle). (C) Daily and (D) average weekly respiratory exchange ratio in male and (E, F) female

mice. G) Daily and (H) average weekly water intake in male and (I, J) female mice. (K) Daily and

(L) average weekly locomotor activity in male and (M, N) female mice. Data are shown as mean

± SEM for N=8 per sex and were statistically analyzed using a repeated measures one-way

ANOVA with Tukey multiple comparison tests. P-value definitions: not-significant (ns) >0.05, *

<0.05, **<0.01, ***<0.001, **** <0.00001.

Supplemental Figure 6. Feeding behaviors in mice switched to a 45% high-fat diet separated

by light:dark cycle

Feeding behavior analysis in male and female mice switched from chow to 60% HFD on day 0

(dashed lines) in metabolic chambers and divided by light (blue) and dark (beige) cycles. (A)

Daily and (B, C) average weekly meal number in male mice in the light and dark cycle. (D)

Daily and (E, F) average weekly meal number in female mice in the light and dark cycle. (G)

Daily and (H, I) average weekly intermeal interval in male mice in the light and dark cycle. (J)

Daily and (K, L) average weekly intermeal interval in female mice in the light and dark cycle.

(M) Daily and (N, O) average weekly meal duration in male mice in the light and dark cycle. (P)

Daily and (Q, R) average weekly meal duration in female mice in the light and dark cycle. (S)

Daily and (T, U) average weekly meal size by mass in male mice in the light and dark cycle. (V)

Daily and (W, X) average weekly meal size by mass in female mice in the light and dark cycle.

(Y). Daily and (Z, AA) average weekly meal size by calorie in male mice in the light and dark

cycle. (AB) Daily and (AC, AD) average weekly meal size by calorie in female mice in the light

and dark cycle. (AE) Daily and (AF, AG) average weekly meal rate in grams per minute in male

mice in the light and dark cycle. (AH) Daily and (AI, AJ) average weekly meal rate in grams per

minute in female mice in the light and dark cycle. (AK) Daily and (AL, AM) average weekly

meal rate in calories per minute in male mice in the light and dark cycle. (AN) Daily and (AO,

AP) average weekly meal rate in calories per minute in female mice in the light and dark cycle.

Data are shown as mean ± SEM for N=8 per sex and were statistically analyzed using a repeated

measures one-way ANOVA with Tukey multiple comparison tests. P-value definitions: not-

significant (ns) >0.05, * <0.05, **<0.01, ***<0.001, **** <0.00001.

Supplemental Figure 7. Shifts in meal patterns across time and light-cycle in mice fed obesogenic diets

Frequency distribution of (A) Type I (<60 mg), (B) Type II (<160 mg), (C) Type III (<280 mg), (D) Type IV (>280 mg) meals in male and (E) Type I (<60 mg), (F) Type II (<160 mg), (G) Type III (<280 mg), and (H) Type IV (>280 mg) meals in female mice switched to a 60% HFD and separated by light:dark cycle. Frequency distribution of (I) Type I (<60 mg), (J) Type II (<160 mg), (K) Type III (<280 mg), (L) Type IV (>280 mg) meals in male and (M) Type I (<60 mg), (N) Type II (<160 mg), (O) Type III (<280 mg), and (P) Type IV (>280 mg) meals in female mice switched to a 45% HFD separated by light:dark cycle. Data are shown as mean ± SEM for N=8 per sex and were statistically analyzed using a repeated measures one-way ANOVA with Tukey multiple comparison tests. P-value definitions: not-significant (ns) >0.05, * <0.05, **<0.01, ***<0.001, **** <0.00001.

Supplemental Figure 8. Metabolic phenotyping of mice switched to a 60% HFD with running wheel access

Metabolic parameters in male mice switched from chow to a 60% HFD at day 0 (dashed line) in metabolic chambers with wheels locked or wheels unlocked. (A) Daily and (B) average weekly energy balance. (C) Average weekly body weight. (D) Daily and (E) average respiratory exchange ratio and average weekly respiratory exchange ratio during the (F) light and (G) dark

cycle. (H) Daily and (I) average water intake and average weekly water intake during the (J)

light and (K) dark cycle. (L) Daily and (M) average locomotor activity and average weekly

locomotor activity during the (N) light and (O) dark cycle. (P) Average daily and (Q) light cycle

and (R) dark cycle running wheel activity in mice with access to unlocked running wheels. (S)

Light and (T) dark cycle average weekly energy expenditure. Data are shown as mean ± SEM

for N=8 per group and were statistically analyzed using a two-way ANOVA with Sidaks

multiple comparison tests. P-value definitions: not-significant (ns) >0.05, * <0.05, **<0.01,

<0.001, * <0.00001.

Supplemental Figure 9. Effects of running wheel access on feeding behaviors and meal

patterns

Feeding behavior and meal type analysis by light:dark cycle in male mice switched from chow

to a 60% HFD with access to locked or unlocked running wheels. Average weekly (A) meal

numbers, (B) inter meal interval, (C) meal duration, (F) meal size by mass (G) meal size by

calorie, (H) meal rates by grams per minute, and (I) meal rates by calories per minute. Mice with

locked wheels are shown in light gray (light cycle) and dark gray (dark cycle) bars with data

points depicted by circles. Mice with unlocked wheels are shown in orange (light cycle) and

blue (dark cycle) with data points depicted by squares. Frequency of meal types [Type I (<60

mg), Type II (<160 mg), Type III (<280 mg), Type IV (>280] across time on diet in mice with (J)

wheels locked and (K) wheels unlocked. (L) Frequency of meal types across time on diet in mice

with wheels locked vs. wheels unlocked. Data are shown as mean ± SEM for N=8 per group and

were statistically analyzed using a two-way ANOVA with Sidaks multiple comparison tests. P-

value definitions: not-significant (ns) >0.05, * <0.05, **<0.01, ***<0.001, **** <0.00001.

Figure 1. Food intake and energy balance in mice fed a 60% high-fat diet

Figure 2. Feeding behaviors in mice switched to 60% high-fat diet

Figure 3. Food intake and energy balance in mice swithced to a 45% high-fat diet

Figure 4. Daily and weekly feeding behaviors in mice fed a 45% high-fat diet

Figure 5. Meal patterns of male and female mice fed obesogenic diets

Figure 6. Effects of running wheel access on feeding behavior and energy expenditure in male mice switched to a 60% high-fat diet

Supplemental Figure 1. Metabolic phenotyping of mice switched to a 60% high-fat diet [light:dark cycle]

Supplemental Figure 2. Metabolic phenotyping of mice switched to a 60% high-fat diet

Supplemental Figure 3. Feeding behavior of mice switched to a 60% high-fat diet [light:dark cycle]

Supplemental Figure 4. Metabolic Phenotyping of of mice switched to a 45% high-fat diet [light:dark cycle]

Supplemental Figure 5. Metabolic phenotyping of mice switched to a 45% high-fat diet

Supplemental Figure 6. Feeding behavior of mice switched to a 45% high-fat diet [light:dark cycle]

Supplemental Figure 7. Shifts in meal patterns across time and light-cycle in mice fed obesogenic diets

Supplemental Figure 8. Metabolic phenotyping of mice switched to a 60% HFD with running wheel access

Supplemental Figure 9. Effects of running wheel access on feeding behavior and meal patterns