A systematic analysis of the effects of splicing on the diversity of post-translational modifications in protein isoforms

Sam Crowl[1], Maeve Bella Coleman[1], Andrew Chaphiv[1], Ben T. Jordan[1], Kristen M. Naegle[1,*]

1 University of Virginia, Department of Biomedical Engineering and the Center for Public Health Genomics, Charlottesville, VA, 22903

* kmn4mj@virginia.edu

Abstract

Post-translational modifications (PTMs) and splicing are known to be important regulatory processes for controlling protein function and activity. Despite there being some examples of the interplay between alternative splicing and cell signaling in literature, there have been very few detailed analyses of the impacts of alternative splicing on PTMs, in part due to difficulties in extracting PTM information from splicing measurements. In this work, we bridged the protein- and genome-centric world views to map PTMs to genomic locations for subsequent projection of PTMs onto alternative isoforms, either from known, validated isoforms from Ensembl (ExonPTMapper) or from splice events quantified by RNA-sequencing (PTM-POSE). We then performed a systematic analysis of the diversification of PTMs by alternative splicing across the entire Ensembl transcriptome, including exploration of the modification-specific rates of inclusion across isoforms and how often the regulatory sequences directly flanking a PTM are impacted by splicing, which might indicate altered regulatory or binding interactions in the alternatively spliced isoform. We found that 6-51% of PTMs are excluded from at least one isoform, depending on the modification type. Further, approximately 2% of prospective PTM sites exhibited altered regulatory sequences surrounding the modification site, suggesting that regulatory or binding interactions might be diversified in these proteoforms. Further, to better understand how splicing diversification of PTMs may alter cell phenotype in specific biological contexts, we projected PTMs onto splice events identified within prostate patient tumors from The Cancer Genome Atlas (TCGA) as a result of ESRP1 expression. We identified protein interaction and regulatory networks that may be rewired as a result of both differential inclusion of PTM sites in ribosomal and cytoskeletal proteins and through altering flanking residues surrounding specific phosphorylation sites that may be targets of 14-3-3 proteins and SH2 domains. As a part of this work, we have provided the pipeline for annotating isoforms from either Ensembl (called ExonPTMapper) or novel splicing measurements (called PTM-POSE) with PTMs and their functional consequences as freely available python packages for use by the broader scientific community.

Introduction   1

Post-translational modifications (PTMs) are tightly regulated chemical modifications that play   2
important roles in protein activity, localization, interactions, and degradation. Across the human   3
proteome, there have been 395,822 experimentally observed post-translational modifications across   4
17,663 different proteins, encompassing 128 unique modification types, indicating that the vast majority   5
of proteins are modified post-translationally in some way [1]. Most post-translational modifications are   6
regulated by specific classes of enzymes that can either add or remove the modification on a specific   7
residue. For example, phosphorylation is added by kinases and removed by phosphatases. Additionally,   8
many post-translational modifications can trigger protein-interactions through PTM-specific recognition   9

domains: bromodomains bind to acetyl lysines [2], WW domains and 14-3-3 proteins bind ₁₀
phosphorylated serines and threonines [3], and ubiquitin-binding domains bind ubiquitinated lysines [4] ₁₁
(Supplementary Figure 1). The affinity of enzymes and PTM-specific binding domains for their targets is ₁₂
largely driven by the flanking residues directly surrounding the modification site, a phenomenon often ₁₃
referred to as domain-motif interactions [5] [6]. For example, not all 90 tyrosine kinases within the ₁₄
human proteome can phosphorylate all known phosphotyrosines. Instead, kinases are 'wired' to ₁₅
phosphorylate only certain sequences (called the enzyme or domain 'motif'), which allows different ₁₆
kinases to elicit different physiological effects in a cell. In this way, post-translational modifications allow ₁₇
for both specific and controlled regulation of protein function and interactions, and loss of a PTM site or ₁₈
change to the flanking sequence can have large implications on protein function and regulation. ₁₉

Nearly all multi-exonic genes are expected to undergo alternative splicing to produce multiple unique ₂₀
protein isoforms from the same gene, suggesting the potential for PTMs to be modulated by alternative ₂₁
splicing [7] [8]. In fact, there are several examples of isoform-specific PTMs throughout literature that ₂₂
have resulted in distinct functional differences between isoforms, such as a glycosylation site found in an ₂₃
isoform of neuroligin-1 important for regulating its interaction with neurexin, altering synapse ₂₄
formation [9]. Globally, PTM sites, particularly those found within disordered regions of a protein, have ₂₅
been suggested to be enriched within tissue-specific exons (differentially expressed across ₂₆
tissues) [10] [11], although there is some work that disputes this idea [15]. Further, alternatively spliced ₂₇
regions of proteins are enriched for linear motifs corresponding to several PTM-specific binding domains ₂₈
important for faciliatating protein interaction and signaling networks, including SH2 and WW ₂₉
domains [12]. However, most proteomic experimental pipelines cannot capture different protein isoforms ₃₀
from the information in small peptide fragments. Hence, the majority of published proteomic ₃₁
experiments, which then get wrapped into PTM-databases, assign PTMs to a single reference protein, ₃₂
typically the "canonical" isoform [13]. When considered together, it is typically based on a small subset ₃₃
of data (tissue-specific exons) [10] [11] [15], reliant on sparse UniProt annotations [12] [16], and/or looks ₃₄
at PTMs as one entity, limiting the broad utility of these results. While there are some recently ₃₅
developed tools and databases that annotate splice events with impacted PTMs like Exon Ontology [53] ₃₆
and ExonSkipDB [80], they are not easily implementable across the entire transcriptome, work with a ₃₇
smaller repository of modifications, and tend to provide limited contextual information on specific PTM ₃₈
sites (function, regulation, interactions, etc.). Further, they do not take into account the potential of the ₃₉
flanking residues surrounding PTMs to be impacted by splice events and disrupt linear motifs required ₄₀
for certain protein interactions. As a result, pre-mRNA splicing and post-translational modifications are ₄₁
often thought of as distinct mechanisms of altering protein function and behavior both within and ₄₂
between tissues [14].As such, there is a need for more comprehensive analyses of the relationship between ₄₃
splicing and PTMs that considers the diversity of PTMs, their potential functions, and the different ways ₄₄
in which splicing can impact PTMs in specific biological contexts. ₄₅

To overcome the limitations of the proteomic annotations of PTMs and assess the relationship with ₄₆
splicing, we developed a reproducible computational pipeline to merge proteome-level annotations of ₄₇
post-translational modifications with genome-level annotations of alternative transcripts, identifying ₄₈
"prospective" PTM sites in alternative isoforms annoted in Ensembl or identified from transcriptomic ₄₉
experiments. To the best of our knowledge, this is the most extensive global analysis of the relationship ₅₀
between splicing and post-translational modifications to date and the first to consider the importantance ₅₁
of changes to the flanking residues surrounding PTMs. We also assess different modification types ₅₂
separately, whereas most prior literature focused on PTMs as a whole or only phosphorylation. We ₅₃
predominantly focused on two mechanisms by which alternative splicing can modulate PTMs – exclusion ₅₄
of the modification site from an isoform or alteration of the short linear motif surrounding the ₅₅
modification site. We illustrated that splicing plays a significant role in regulating the presence of PTM ₅₆
sites in alternative isoforms ($\approx 30\%$ of PTMs are excluded from at least one isoform), but that certain ₅₇
modification types are more likely to be maintained across different isoforms, such as glycosylation. ₅₈
Further, we found that a subset of prospective PTM sites ($\approx 2\%$, amounting to 14,850 prospective PTM ₅₉
sites) exhibit altered flanking sequences that may serve to change the regulatory enzymes responsible for ₆₀
the modification, as well as the binding domains that interact with that site. ₆₁

Ultimately, we wished to understand the role of splicing control of PTM sites on defining protein interactions, signaling pathways, and cell phenotype in specific biological contexts. As a case study, we applied a PTM-focused analysis of splicing patterns in prostate cancer patients with increased ESRP1-expression using data from The Cancer Genome Atlas (TCGA), as this splicing factor has been shown to be detrimental to prostate patient outcomes. We identified protein-interaction and signaling networks that may be altered due to differential inclusion of PTMs, potentially explaining increased cell proliferation through increased SGK1-mTOR signaling. Further, we identified several cases in which potential interactions with SH2 domains or 14-3-3 proteins were disrupted by altered flanking sequences as a result of ESRP1 expression. From this work, we have highlighted the importance of including post-translational modifications in the analysis of splicing and provided easily implementable tools for annotating isoforms and splice events with PTMs and their functional consequences, called ExonPTMapper (for Ensembl-defined transcripts) and PTM-POSE (for transcriptomic datasets).

Results

In order to perform a systematic analysis of when and how alternative splicing diversifies PTMs across protein isoforms, we first needed to expand our knowledge of the post translational modifications present in alternative isoforms. The majority of post-translational modifications in modern-day PTM compendia were discovered by mass spectrometry-based approaches, which typically rely on trypsin-based proteolysis to yield MS-amenable peptide fragments. PTM annotations coming from tryptic-fragment identification are usually assigned to the canonical isoform, unless the tryptic fragment can be uniquely assigned to an alternative isoform (Supplementary Figure 2). In fact, only 15.3% of alternative isoforms have any recorded PTMs, compared to only 93.1% of canonical isoforms, despite the fact that a peptide containing a modified residue could often be mapped to more than one possible isoform. In order to overcome this limitation, we integrated three key resources using a Python-based framework. We used genomic information from Ensembl to provide the set of all human protein-coding transcripts supported by experimental evidence and the genomic location of their exons [17]. We used the protein database UniProt to establish the list of unique human protein identifiers, the canonical isoform of each human gene, and any reviewed alternative protein isoforms [18]. We used ProteomeScout and PhosphoSitePlus as the resources for post-translational modifications, with the former aggregating PTMs from five major compendia, creating a comprehensive and integrated resource of PTMs [1] [25]. Using these resources, we ultimately sought to translate between the fields of genomics and proteomics for a comprehensive analysis of the control of PTMs by splicing, avoiding aggregating PTMs into a single class of information. Importantly, we developed this framework for reusability and expansion as our knowledge of both splicing and post-translational modifications continues to improve, providing the tools for annotation of the entire Ensembl transcriptome (ExonPTMapper) or transcripts measured in specific biological contexts and quantified by tools like SpliceSeq or MATS (PTM-POSE) [43] [78].

Projecting Prospective Post-translational Modification Sites onto Alternative Isoforms

To expand the data available for analysis, we mapped post-translational modification sites in canonical protein isoforms to their location in their corresponding exon, transcript and gene recorded in Ensembl. With the genomic location of each PTM, we can project PTMs to the exons or spliced regions that contain its genomic location, allowing us to account for whether final transcripts include some or all of the PTM-carrying exon (Fig. 1, Supplementary Figure 3). We refer to the PTM sites projected onto isoforms as "prospective", indicating that the genomic location is known to code for a modifiable residue, though there may not be direct annotated evidence of the modification existing in the alternative isoform. Using this pipeline for Ensembl-defined protein-coding transcripts, we successfully expanded the annotations of PTMs on alternative isoforms across the proteome and transcriptome – 98.8% of prospective PTM sites identified in alternative isoforms from UniProt and/or Ensembl were not previously annotated in protein databases (Supplementary Figure 2C).

Figure 1. Mapping and Projecting Post-Translational Modification (PTMs) onto Alternative Transcripts. In order to expand the knowledge of PTMs in alternative isoforms, we mapped PTMs from comprehensive databases of post translational modifications (ProteomeScout and PhosphoSitePlus) onto the genomic coordinates of Ensembl **(A)**, then project those genomic locations onto Ensembl-defined protein coding transcripts or splice events identified from transcriptomic data **(B)**. We refer to these projected PTMs as "prospective" PTMs in alternative isoforms. We defined three key outcomes of splicing on PTMs based on the predominant events observed in our analysis (PTM denoted as the gray circle on the third exon in the canonical transcript): **(1)** the PTM was successfully mapped onto the alternative isoform, with no changes to its surrounding region, **(2)** the PTM was excluded from the alternative isoform, or **(3)** the PTM was successfully projected onto the alternative isoform, but residues surrounding the PTM site were altered, which may influence the specificity of enzymes and binding partners.

Given the successful projection of PTMs onto alternative isoforms, we next assessed the effect of splicing on PTMs based on a set of key, possible effects (also described in Fig. 1). First, a PTM may be included in an isoform, maintaining any potential interactions that have been shown for the canonical isoform. Second, a PTM could be excluded from an isoform altogether, removing any potential functional changes or interactions induced by the modification. Lastly, if a prospective PTM is included in the alternative isoform, it may contain a unique flanking sequence directly surrounding the PTM distinct from the canonical isoform, potentially leading to different interactions with regulatory enzymes or PTM-specific binding domains. The effect of these changes was also annotated with the splicing event that resulted in the change relative to the canonical isoform, such as a skipped exon, alternative splice site, or mutually exclusive exons. The transcriptome-wide datasets, prospective PTMs, and regulation through splicing are freely available to the research community (see data availability secton for details).

PTMs are differentially included through alternative splicing

We sought to evaluate how splicing might impact the presence of PTMs in alternative isoforms across the entire transcriptome and whether the relationship between splicing and PTMs is dependent on the specific modification type. To address this question, we first considered the number of PTMs within Ensembl-defined constitutive exons – exons found in every transcript associated with a gene. We found constitutive exons exhibited a lower density of PTM sites than non-constitutive exons, with the most densely populated exons tending to be non-constitutive (Supplementary Figure 4). This matches previous reports comparing tissue-specific exons to constitutive exons measured from RNA sequencing data and provides further evidence that splicing plays a role in the inclusion of PTM sites in protein isoforms [10] [11]. To identify which PTMs in particular are or are not being regulated by splicing, we identified "constitutive PTMs", or PTMs that are included in all protein-coding transcripts of a gene and are therefore not controlled by splicing. Overall, 68.23% of PTM sites could be defined as

constitutive across Ensembl transcripts. However, not all types of PTMs are controlled by splicing at the same rate, with constitutive rates ranging from 49.1% (Palmitoylation, 167 unique observations) to 93.8% (Crotonylation, 192 unique observations) (Fig. 2A,B, Supplementary Figure 5). PTMs found within structural domains were slightly more likely to be included in isoforms, but this alone failed to explain the variation across modification types (Supplementary Figure 6). When compared against a null model where splicing is entirely random and non-specific to modifiable residues, we found that most modifications, except palmitoylation, exhibited slightly higher constitutive rates than would be expected by ranomd chance, but the observed variability across modification types is unlikely to be observed by randomness alone (Fig. 2B). Phosphorylation, sumoylation, hydroxylation and palmitoylation were among the modification types with the lowest constititive rates. On the other hand, ubiquitination, glycosylation, crotonylation, carboxylation and sulfation had much higher constitutive rates than in the null models (Fig. 2B).

Given the broader classes of PTMs demonstrated variability, we also assessed the subtypes within each modification type, and found further variability in splicing regulation (Supplementary Figure 7). For example, glycosylation as whole exhibits a higher constitutive rate than most other modifications (74.6%). This is largely due to N-linked glycosylation (75.0%), while O-linked glycosylation has a constitutive rate closer to the average across all modifications (68.9%). Similarly, phosphoserines exhibited slightly lower constitutive rates than either phosphothreonine or phosphotyrosine sites. This indicates that there is variability in splicing regulation even amongst similar modification types.

As many Ensembl transcripts have not yet been shown to produce functional proteins, it is possible that these results are influenced by transcripts that do not ultimately produce functional proteins. To account for this possibility, we repeated the same PTM-specific analysis using different definitions of functional transcripts, including functionality predictions from TRIFID [19]. As we implemented more restrictive criteria, we observed a decrease in the number of isoforms per protein and an increase in the constitutive PTM rate, suggesting that the true constitutive rate is likely somewhere between 70-90% (Supplementary Figure 8). Regardless of functional criteria used, the rates continued to retain similar variability across most modification types, suggesting that control of PTMs by splicing is PTM-specific and PTMs should always be evaluated in a detailed, PTM-specific manner, even within the same modification class – such as seen in O- vs. N-linked glycosylation and serine vs tyrosine phosphorylation.

To determine the potential functional relevance of splicing-regulated PTMs, we obtained molecular functions and biological processes associated with PTMs from PhosphoSitePlus [25] and tested for enriched annotations within constitutive and non-constitutive PTMs. PTMs controlled by splicing (non-constitutive) were found to be significantly associated with many different functions that are important for regulating protein interactions and activity, including PTM sites important for intracellular localization and regulating molecular associations, indicating that alternative splicing may rewire protein-interaction networks through exclusion of the PTM sites driving these protein interactions (Supplementary Figure 9, Supplementary Table 1). Constitutive PTMs, on the other hand, were only found to be enriched for sites regulating receptor desensitization, and a larger than expected percentage of constitutive PTM sites did not have a known function. Similarly, constitutive PTM sites were not found to be enriched for any biological processes, while non-constitutive PTMs were enriched for a variety of processes including cell growth and gene transcription (Supplementary Figure 9). In addition to protein function, differences in included PTM sites may alter the regulation of PTMs. For example, when looking at known kinase-substrate interactions from PhosphoSitePlus [25] and RegPhos [74], we found that some kinases' substrates were more impacted by splicing (RAF1, ERBB2, SRPK1) while others had substrates that tended to be constitutive within that protein (BUB1, GRK2, CAMK2A) (Supplementary Figure 9). Combined, this points to the idea that splicing control of PTMs across isoforms will impact protein function and the regulatory networks associated with the spliced protein.

Next, we wished to understand if the way in which PTMs were excluded had a relationship with the types of splicing that can occur – such as by skipped exons, alternative splice sites, or mutually exclusive exons. While the rate of splicing control of PTMs was modification-specific, most modifications were impacted by different splice events in similar proportions, with most excluded PTMs being a result of a skipped exon event. There were a few small exceptions, as hydroxylation and sulfation sites were lost

Figure 2. Splicing controls the presence of post-translational modifications in alternative isoforms. Modification-specific statistics of PTM inclusion across human alternative transcripts. Each panel shares the same x-axis, sorted from the largest number of identified modifications (phosphorylation) to the least (sulfation). Here, we only show modifications with at least 140 instances across the human proteome for visualization purposes. For data of all modification types, see Supplementary Figure 5. **A)** The number of modifications documented in canonical UniProt proteins for each class of post-translational modification. Each bar is colored based on what type of molecule is being added: small molecules, proteins, sugars, or fatty acids. **B)** The fraction of modification sites that were found across all isoforms of a given gene, defined as a constitutive PTM rate. We compared the results for the real PTM data to a null model generated by randomly selecting residues across the proteome in matching proportions and treating them as real PTM sites. **C)** Breakdown of the types of alternative splicing events that result in the loss of a PTM site relative to the canonical isoform, considering skipped exon events, alternative splice sites (ASS), and mutually exclusive exons (MXE). Alternative splice sites are defined as described in Wang et al. and elsewhere [7]. Mutually exclusive exons were identified based on criteria from Pillmann et al. [20]. **D)** Density of each type of PTM in tissue-specific exons identified across three different publications [10] [22] [21]. PTM density is defined as the number of PTMs per 1000 amino acids and has been normalized by the density of that same PTM across the entire proteome (regardless of tissue-specific expression). A normalized density greater than 1 indicates more PTMs than expected within tissue-specific exons, a normalized density less than 1 indicates fewer PTMs than expected.

due to an alternative splice site more often than other modifications (Fig. 2C). In rare cases, mutually exclusive exon events may also impact the presence of PTMs. We identified 430 mutually exclusive exon events associated with PTM sites, using criteria defined previously by Pillman et al. [20]. These events encompass 153 different PTM sites and 13 modification types (Supplementary Figure 10). Based on global alignment of mutually exclusive exon pairs, we found that PTM sites were likely to be conserved in only 51.6% of cases, perhaps indicating that mutually exclusive exon events may be a mechanism by which PTM sites can be removed from alternative isoforms without significantly disrupting the overall structure of a protein.

While these results suggest a potential role of splicing in controlling PTM inclusion, we did not consider expression of transcripts within and across tissues, instead looking at all protein-coding alternative transcripts that have been recorded in Ensembl. To assess differential inclusion of PTM sites across tissues, we compiled a set of tissue-specific exons identified across three different publications [10] [21] [22]. Overall, PTMs tended to be found at a higher density in tissue specific exons than across the entire proteome, in line with prior results [10] [11]. As when looking at the entire transcriptome, PTM density in tissue-specific exons was largely dependent on the modification type. Glycosylation and nitrosylation exhibit lower density within tissue-specific exons than found across the entire proteome, matching their higher constitutive rate (Fig. 2D). Interestingly, sumoylation and palmitoylation, which were observed to have low constitutive rates, were not found to be abundant within these tissue-specific exons. Although there are some discrepancies between PTM density within tissue-specific exons and constitutive rates across the transcriptome for certain modification types, these results confirm that alternative splicing is an important mechanism for modulating the presence of certain PTMs, in particular those involving phosphorylation and methylation sites.

Alternative Splicing affects PTM-regulatory sequences

PTMs are important components of signaling pathways, cellular communication, and protein interactions, all of which are facilitated by direct interactions with a PTM by its corresponding modifying enzyme(s) and recognition domain(s). The specificity of these interactions are driven by the residues immediately upstream and downstream of the modification site, with the linear motif directly surrounding the PTM being a prerequisite for recognition of the PTM site [5] [6] [23]. Given this, we hypothesized that splicing may alter PTM function by changing the flanking sequence surrounding the PTM site. There are several mechanisms by which an alternative splice event could lead to changes in the flanking sequence of a PTM, including a skipped/retained exon adjacent to the PTM, an alternative splice site in a nearby exon or in the PTM-containing exon, or a mutually exclusive exon event (Fig. 3A). With this in mind, we sought to characterize how often alternative splicing leads to changes in the flanking sequence surrounding PTM sites, defining the flanking sequence as the five amino acids on either side of the PTM, unless otherwise stated. Although the specific residues important for PTM recognition are variable, depending on the modification and interacting protein, a regulatory window size of five residues likely captures the most common residues important for facilitating interactions across most modifications/proteins.

When comparing flanking sequences of a PTM site in the canonical and corresponding alternative isoforms, we found that 2.09% of prospective PTM sites observed changes to the flanking residues (Supplementary Table 2). Importantly, 25.8% of these sites have a matching tryptic fragment compared to the canonical tryptic fragment, indicating that 0.54% of prospective PTM sites have altered regulatory regions that would not be detected by trypsin-mediated MS (Fig. 3B). Across the possible altered regulatory regions, we observed that acetylation, trimethylation, carboxylation, and palmitoylation exhibited the highest rate of alteration (Fig. 3C). When compared against the null model, we found that the high alteration rate could not be explained by randomness alone (Supplementary Figure 11). In the majority of cases, we observed that altered regulatory windows are caused by changes to the adjacent exon, rather than the PTM-containing exon itself, usually via a skipped/retained exon (Fig. 3D). Both glycosylation and nitrosylation, which also had higher constitutive rates, have lower rates of altered flanking sequences, which were at rates that could be attributed to the distribution of amino acids across the transcriptome (Supplementary Figure 11).

We further explored the residue positions in the flanking sequences that were most affected by

Figure 3. Alternative splicing rewires PTM-driven interaction networks by altering flanking sequences. A) Illustrative examples of the modes of alternative splicing that result in changes to the flanking sequence around a PTM, comparing two transcripts with differing flanking sequences (highlighted in red) surrounding the same PTM. **B)** Number of prospective PTMs found in alternative isoforms with an unchanged flanking sequence (+/- 5 amino acids), an altered flanking sequence but unchanged tryptic fragment, or both an altered flanking sequence and tryptic fragment. **C)** Fraction of prospective PTMs with altered flanking sequences, broken down by modification type, with the total number of events indicated above each bar. The proximity of the change to the PTM is denoted by the different bar colors, ranging from immediately next to (1) or up to 5 amino acids away from the PTM on either side. **D)** Breakdown of which events described in panel A most commonly lead to altered regulatory regions. **E)** Sequence identity between differing flanking sequences in the canonical and alternative isoforms, using different window sizes of amino acids on either side of the PTM. **F)** Change in the likelihood of kinase interaction from the canonical to the alternative flanking sequence, based on predicted site percentiles derived from Kinase Library software [23]. We only considered PTM sites with an altered flanking sequence and conserved tryptic fragment (orange bar in panel B) that also had at least one known kinase interaction annotated in PhosphoSitePlus [25] **G)** An example of an altered flanking sequence surrounding the phosphorylation site at S159 in SIRT3. The change in Kinase Library site percentile between the canonical and alternative isoform of SIRT3 for various cyclin-dependent kinases are shown, with positive changes (red arrows) indicating a preference for the alternative isoform, while negative changes (blue arrows) indicating a preference for the canonical isoform.

splicing and the potential consequence of these changes. Due to the nature of splicing-induced changes, most regulatory regions are altered predominantly only on one side of the modifications flanking sequence, and the farther you move from the PTM the more likely it is to see changes to the regulatory region (Supplementary Figure 11). Among the $\approx 2\%$ prospective PTMs with altered linear motifs, 7.5% of these PTMs (2,381 prospective PTMs) have a change in the +/-1 residue, immediately adjacent to the PTM site, which has the potential to significantly impact enzyme or binding domain interaction (Fig 3C, Supplementary Figure 11). For example, proline-directed kinases are a large subset of serine/threonine kinases with a strong preference for a proline in the +1 position, which includes cyclin dependent kinases (CDKs) and ERK [23]. Similarly, the CDY family of proteins contain a chromodomain that binds to methylated lysines with a strong preference for lysines that immediately follow an arginine [24]. In both cases, the remaining residues in the regulatory window serve to further drive specificity and the strength of the interaction, with more minor effects. We further confirmed that splicing-induced regulatory sequence changes typically preserve information in the motif, but create smaller changes. Regardless of the regulatory window size chosen (3, 5, 7, or 10 amino acids on either side of the PTM), regulatory sequences generally still maintained > 60% sequence identity (Fig. 3E). Altogether, these results indicate that splicing can alter regulatory sequences in ways that could impact PTM-specificity and recognition, but suggests that it is unlikely to fully disrupt the modifiability of a PTM.

We next sought to understand how changes in the regulatory sequences might affect PTM regulation and protein interactions. Given that there is minimal data regarding isoform-specific binding or enzymatic activity, we compared the predicted affinity of various serine/threonine kinases for both canonical sequences and alternative regulatory sequences using a kinase motif enrichment tool, Kinase Library [23]. We focused on serine and threonine phosphorylation sites with an altered flanking sequence and a conserved tryptic fragment, as these are the modifications that are most likely to be missed by trypsin-mediated MS and be misannotated on the canonical isoform of the protein. In total, we looked at 328 phosphotyrosine events and 1484 events (82 of which had a known annotated kinase interaction). We used the Kinase Library tool to predict the likelihood of a given kinase interacting with either the canonical or alternative regulatory sequence and then extracted the change in the Kinase Library percentile. Interestingly, in many cases, flanking sequence changes had a wide range of possible impacts on both annotated kinase interactions (kinases that had been previously observed to phosphorylate a given site) and on kinases not annotated as a regulatory kinase: decreased likelihood in some cases, increased it in others, and in an equal number of cases it had no effect on the predicted likelihood (Fig. 3F). Kinases that are not known to interact with a particular phosphorylation site exhibited more extreme cases than the annotated, known kinase interactions. In some cases, the predicted ranking for unannotated kinases differed by more than 100 ranks between the canonical and alternative isoform, while predicted changes in the annotated interaction tended to be more subtle. However, there are also cases where the annotated kinase's predicted interaction differs when looking at the alternative isoform. For example, SIRT3, a deacetylase that regulates energy metabolism, contains an activating phosphorylation site at S159 that has been shown to be phosphorylated by CDK1. In an alternative isoform of SIRT3, a skipped exon event results in a unique regulatory sequence preceding S159 (Fig. 3G). While the proline in the +1 position remains present, a key feature of proline-directed kinases like CDKs, the surrounding changes result in a decreased likelihood of phosphorylation by CDK1 while increasing affinity for other CDK family members like CDK2, suggesting a potential difference in the regulation of phosphorylation at S159, moving away from CDK1 in the alternative isoform (Fig. 3G). It appears that splicing-induced changes to regulatory sequences causes subtle changes in PTM-specificity and, in some cases, may lead to altered regulatory networks. Although limited by constraints on known enzyme interactions for extending confidently to more phosphorylation sites or interaction changes with other differentially spliced PTMs, this analysis indicates that altered flanking sequences of PTMs introduced by splicing has the capacity to diversify regulatory networks of a protein.

Overexpression of splicing factor, ESRP1, leads to tumor-specific post-translational modifications that drive cell growth and chromosomal instability in prostate cancer

Having assessed the global relationship between PTMs across all possible isoforms recorded in Ensembl, we wished to understand how alternative splicing within specific biological contexts regulates the inclusion of PTMs to alter protein function and drive changes to cell phenotype. Aberrant splicing is involved in several diseases, including cystic fibrosis, muscular dystrophy, and cancer [26] [27] [28] [29] [30]. In our analysis of the entire transcriptome, we found that non-constitutive PTM sites were enriched for sites important to carcinogenesis, suggesting that changes to splicing patterns may influence the presence of functionally important PTM sites for the development and progression of cancer (Supplementary Figure 9). Further, many splicing factors important for regulating splicing of pre-mRNA have been implicated in altering cancer progression, such as epithelial splicing regulatory proteins (ESRPs), splicing factors responsible for the splicing of epithelial-specific isoforms [31]. In prostate cancer, it has been reported that elevated levels of ESRP1 leads to worsened prognosis [32] [33] [30]. For this reason, we selected prostate cancer as a system to study the impact of differential inclusion of PTMs and their flanking sequences as a mechanism for disease progression.

To identify potential mechanisms by which ESRP1 may worsen prognosis in prostate cancer, we first used PTM-POSE to project PTMs onto all splice events identified by SpliceSeq within prostate cancer patient samples from The Cancer Genome Atlas (TCGA) [43]. By applying PTM-POSE to TCGA SpliceSeq data, we could compare both exon and PTM utilization across samples within the prostate cancer cohort, defined by the percent spliced in (PSI) rates of each exon and PTM. By comparing PSI for patients with low and high ESRP1 expression, we identified a total of 1,125 PTM sites (0.5% of PTMs found in dataset) that were differentially included based on ESRP1 expression, with 476 PTM sites excluded and 649 PTM sites included when ESRP1 is highly expressed (Fig. 4A, Supplementary Table 3). Across the 6.8% genes in the dataset associated with significant splice events as a result of ESRP1 expression, 54% of these genes contained PTMs impacted by ESRP1-associated splicing, with around of half of impacted exons containing PTMs (Fig 4B). To ensure that we captured PTM sites that are directly impacted by ESRP1, we also applied PTM-POSE to splice events previously identified upon knockdown of ESRP1 in H358 cells by Yang et al., finding that a subset of the PTMs were found in both ESRP1 knockdown and TCGA datasets, including phosphorylation sites found on TSC2, CTNND1, CD46, and MAP3K7, indicating that we successfully captured at least some ESRP1-mediated splicing in these patient samples (Supplementary Figure 12) [34]. Phosphorylation makes up the the majority of differentially included modifications, but we also identified several acetylation, ubiquitination, and methylation sites, among others (Supplementary Figure 12). Another 334 PTMs, or 0.2% of PTMs assessed, were also found to have altered flanking sequences due to changes to adjacent exons (Fig. 4B). In sum, 1489 different PTM sites across 562 different exons were impacted by differences in ESRP1 expression, indicating extensive regulation of PTMs as a function of ESRP1.

In order to better understand the role of ESRP1-mediated splicing of PTMs on protein function and disease progression, we performed a multi-level analysis which harnessed information from gene-level (Gene Ontology [48], Reactome [73]), exon-level (Exon Ontology [53]), and PTM-level databases (PhosphoSitePlus [25], RegPhos [74], PTMcode [77]). We found that genes associated with spliced PTMs were enriched for cytoskeletal and actin-binding proteins, which matches expected results given ESRP1's role in regulating cell motility and the epithelial-to-mesenchymal transition (Fig. 4C, Supplementary Figure 13). Over 75% of spliced cytoskeletal genes contained PTMs that were impacted by ESRP1-related splicing. Intriguingly, within the spliced cytoskeletal genes, there was a subset of genes that were also involved in the apoptotic execution phase pathway (CASP8, ADD1, MAPT, PTK2, SPTAN1) which was only enriched when considering genes with impacted PTMs rather than all spliced genes, indicating this to be a PTM-driven regulation of apoptosis although the specific function of the impacted sites has yet to be elucidated (Fig. 4C, Supplementary Figure 13). We did also observe several additional individual PTM sites that are associated with cell growth, cell cycle regulation, and apoptosis based on PhosphoSitePlus annotations. Several of these PTMs, but not all, suggest increased

Figure 4. ESRP1 drives changes to protein function and interactions in prostate cancer by altering the presence and flanking sequences of specific PTM sites. Using data from The Cancer Genome Atlas and SpliceSeq, we explored the relationship between ESRP1 expression and splicing-controlled PTM sites in prostate cancer [43] [44]. **A)** Patient-specific percent spliced in (PSI) of individual PTM sites that were found to be differentially included between ESRP1-low patients ($z \leq -1$) and ESRP1-high patients ($z \geq 1$). **B)** Breakdown of the fraction of genes, exons, and PTMs impacted by splicing, including whether the impact is due to changes in inclusion or alterations to the surrounding flanking sequence of a PTM. **C)** Selected gene sets from Gene Ontology (GO) [48] and Reactome (R) [73] that are enriched among spliced genes with PTMs that are impacted (orange and green groups from Panel B). Each point indicates the fraction of spliced genes associated with the gene set belonging to each group in Panel B. **D)** Exon Ontology terms [53] that are associated with the highest number of impacted PTMs, broken down by the type of impact (excluded, included, or altered by ESRP1) **E)** Subset of the PTM-associated interactions impacted by differential inclusion of PTMs due ESRP1 expression, and the proteins with the highest number of impacted interactions in the network. Interactions are based on data from PhosphoSitePlus [25], RegPhos [74], and PTMcode [77]. Orange nodes indicate proteins with spliced PTMs outside domains, and black nodes indicate proteins with spliced PTMs within domains. **F)** Phosphorylated tyrosine sites with different flanking sequences based on ESRP1 expression, which only match the preferred linear motif of select SH2 domains depending on whether an adjacent exon is included in the protein isoform. Each row indicates a different phosphorlated tyrosine site and the specific residue change is denoted as "Inclusion AA -$\leq$ Exclusion AA". Changes that specifically impact the motif are bolded and in black.

proliferation and decreased apoptosis including CRK Y221 (induces apoptosis, decreased inclusion with ESRP1 expression) and PAK4 S291 (induces cell growth, increased inclusion with ESRP1 expression) (Supplementary Figure 14). However, there were also other identified PTMs, such as EEF1A1 S21 (induces apoptosis, increased inclusion in ESRP1 expressing cells), highlighting the importance of context on PTM function and the need to consider all altered proteins and PTMs holistically.

Across the PTMs impacted by ESRP1 expression, the majority of PTMs were found in exonic regions associated with intrinsically unstructured regions of the protein, rather than distinct domains (59.1%) (Fig. 4D). Further, spliced PTM-containing exons were in regions of protein binding and nuclear localization signals, suggesting that ESRP1 control of PTMs may lead to altered protein interaction and regulatory networks (Fig. 4D). In support of the role of these exons and PTMs in facilitating protein interactions, there were 24 and 7 of the ESRP1-related PTMs that have previously been annotated as being important for regulating molecular associations and intracellular localization, respectively (Supplementary Figure 14). Given the importance of spliced PTMs in facilitating protein interactions, we pulled PTM-associated interactions from PhosphoSitePlus [25], RegPhos [74], and PTMcode [77]. We found a total of 476 PTM-associated interactions that are impacted by ESRP1-mediated splicing out of 25,510 potential interactions, with most PTMs facilitating these interactions falling outside of specific structural domains (Fig. 4E). Upon ESRP1 expression, we noted a high number of decreased interactions of ABL/SRC interacting proteins (CRK, ABI1), increased interactions of several ribosomal proteins (RPL6, RPS14, RPS3A), and variable changes to interactions of cytoskeletal proteins (MAPT, CTNND1) (Fig. 4E). Increased proliferation is often observed among ESRP1-expressing cancer cells, which may align with the increase in PTM-associated interactions of ribosomal proteins [31]. Further, previous literature has reported a strong link between increased ESRP1 expression, high copy numbers, and chromosomal deletions, which has been hypothesized to be the result of chromosomal instability and dedifferentiation. Several ribosomal proteins, including RPL6 and RPS14, have also been linked to chromosomal instability and DNA damage response, although it is not clear how the affected phosphorylation, methylation and ubiquitination sites may or may not regulate these processes [75] [81]. In further support of splicing-control of PTMs leading to increased translation and chromosomal instability, we also identified altered interaction networks for several transcription, splicing, and translation regulators including heterogenous nuclear ribonucleoprotein A1 (HNRNPA1), Poly(Rc)-binding protein 2 (PCBP2), and Elongation factor 1-alpha 1 (EEF1A1) (Supplementary Figure 15). An included methylation site in EEF1A1 at K165, methylated by METTL21B upon activation of stress-related pathways commonly observed in different cancers, has been show to alter which mRNA becomes translated without disrupting translation altogether [82]. Based on our results, this may be due to changes in interactions with RPS5, XPO5, and other elongation factors (EEF1G,EEF1D) important for regulation mRNA translation. In addition, phosphorylation sites at S199 in HNRNPA1 and S213/S272 in PCBP2 have been shown to be important for regulating differentiation in various cell types. Combined with the aforementioned apoptosis and cell growth-related PTM sites, this suggests that ESRP1-mediated splicing of PTMs may drive worsened prognosis in prostate cancer by rewiring protein interactions that drive protein translation, cell growth, and chromosomal instability, rather than through the more commonly studied EMT pathways.

In addition to differential inclusion of PTM sites, we also assessed cases where changes to flanking sequences may alter protein interactions by disrupting preferred linear motifs of specific binding domains. We found similar patterns of disruption as observed when looking across the entire transcriptome, where most changes to the flanking sequence retained 60-80% sequence identity after exclusion of the adjacent exon (Supplementary Figure 16A-C). Among the altered flanking sequence events, we found many events that disrupted potential targets of various binding domains with motifs annotated in the Eukaryotic Linear Motif (ELM) database, including the motifs for 14-3-3 proteins (bind phosphorylated serines and threonines) and three different SH2 domains (bind to phosphorylated tyrosines) (Supplementary Figure 16D-E) [76]. For example, the SH2 domain of CRK preferentially binds to proline, leucine, isoleucine, or valine in the +3 position, which are changed to an alanine for CLTCL1 Y1477 and KIAA1217 Y1776 when ESRP1 is expressed (Fig. 4F). In the case of CLTCLC1, the flanking sequence change also causes a shift towards the preferred binding motif of SRC SH2 domain. In this way, CRK interactions appear to

be disrupted in ESRP1-high patients both directly through loss of PTMs within and nearby its SH3 domain and indirectly by changes to CRK SH2 domain binding motifs. Alternatively, the SH2 domain of STAP1 was most impacted by changes at the +4 position for TPM1 Y60, PHPT1 Y93, and TECR Y4 (Fig. 4F). PTMs with disrupted motifs for 14-3-3 proteins were more varied in how they were disrupted, but most impacted genes were related to cytoskeletal genes like MARK2 and SORBS2, which have been shown to interact with 14-3-3 protein epsilon based on interaction data from STRING (Supplementary Figure 16) [46]. Combined with the disrupted interactions by altered inclusion of PTMs, these results highlight how splicing control of PTMs by ESRP1 can rewire whether and how proteins interact with each other, centered around specific biological processes (cell growth, cytoskeletal changes, regulation of translation).

ESRP1 expression is correlated with PTEN and SGK1 signaling in prostate cancer

In addition to altering the function and interactions of proteins, changes to the presence and flanking sequence of specific PTMs will change how proteins are regulated. As phosphorylation is the most commonly impacted type of modification, we asked if the targets of specific kinase were more impacted by ESRP1-mediated splicing than others. Among the known kinase-substrate interactions from PhosphoSitePlus [25] and RegPhos [74], we observed that multiple substrates of AKT1, SGK1, and PRKACA were included with ESRP1 expression, substrates of ABL1 were excluded, and several substrates of MAPK1/3 were altered (Fig. 5A, Supplementary Figure 17). However, given that kinase-substrate information tends to be sparse (less than 5% of phosphorylation sites have an annotated kinase), we employed a kinase-substrate enrichment analysis adapted from a kinase activity inference tool we previously developed, called KSTAR, to determine if there were global changes to kinase-substrate availability due to splicing. We found that substrates of SGK1 and other SGKs were enriched in ESRP1-high patients, while AKT1 substrates were actually more commonly found in ESRP1-low patients. PRKACA, MAPK1/3, and ABL1 were not significantly enriched in either group (Fig. 5B). We also assessed if altered flanking sequences around phosphorylation sites may be further altering these kinase networks, particularly for MAPK1/3 which had two known substrates with altered flanking sequences. Based on the change in PSI and predicted scores from Kinase Library, SGK1 and PRKACA exhibited the highest variability in the likelihood it would interact with phosphorylation sites containing altered flanking sequences as a result of ESRP1 expression, whereas as MAPK3 actually seemed to be least impacted by flanking sequence changes (Fig 5C). However, it should be noted that many of the same phosphorylation sites observed increased preference for multiple of the assessed kinases, with only DOCK7 S958 and SVIL S1177 being uniquely increased in preference for SGK1. Altogether, though, the targets of SGK1 appear to be most impacted by ESRP1 expression in prostate cancer patients.

Given the potential relationship between ESRP1 and SGK1, we sought to understand how changes to SGK1 signaling may lead to different phenotypes depending on ESRP1 expression levels. Potential targets of SGK1 impacted by ESRP1 expression in prostate cancer patient samples many genes relevant to the cytoskeleton, cell growth, and/or mTOR signaling, including sites with increased inclusion in ESRP1-high patients (TSC2 S981, FNIP1 S232, and FLNA S1981) or altered flanking sequences (LIMCH1 S325, SCRIB T728, DOCK7 S958) (Supplementary Figure 17, Supplementary Table 3). Tuberin (TSC2) is a particularly interesting case, as TSC2 is a negative regulator of mTOR signaling and loss of TSC2 has been shown to lead to aberrant S-phase progression upon DNA damage and increased sensitivity to stress, in large part due to sustained mTOR signaling [40] [41] [42]. In addition to loss of expression, TSC2 inhibition of mTORC1 signaling can be abrogated by phosphorylation of S981 by SGK1, which leads to interaction with 14-3-3 proteins (whose targets we also previously identified as being impacted by ESRP1 expression) and localization away from mTORC1 [79]. Exon 26 of TSC2, which contains S981, is found in a higher percentage of TSC2 transcripts from ESRP1-high patients, a trend that was observed across both prior ESRP1 knockdown experiments and other TCGA cancer cohorts (Fig 5D). Given that both SGK1 and TSC2 are linked to the PI3K/PTEN/mTOR pathway, we were curious if PI3K/PTEN pathway activity was related to the oncogenic effects of ESRP1.

Figure 5. ESRP1 rewires kinase signaling networks through changes to their downstream targets. Given the phosphorylation sites identified to be impacted by ESRP1 splicing in prostate cancer in Figure 4, we assessed whether these sites tended to be targets of specific kinases. **A)** Kinases with the highest number of differentially included substrates, based on either known kinase-substrate interactions from PhosphoSitePlus [25] and RegPhos [77] **B)** Enrichment of kinase-substrate interactions among phosphorylation sites that are either included or excluded when ESRP1 is highly expressed. Kinase-substrate enrichment was performed using an adapted version of kinase activity inference algorithm, called KSTAR [70]. The size of the dots represent the strength of enrichment, with color indicating significance. **C)** Change in the likelihood of a kinase-substrate interaction due to changed flanking sequences. Interaction change is defined as the change in kinase library percentile between the inclusion and exclusion isoforms multiplied by the change in percent spliced in (PSI), such that a positive change indicates the kinase is more likely to interact in ESRP1-high patients and a negative change indicates that is less likely to interact. We have highlighted interations that fall in the 90th percentile in green (most likely to increase upon ESRP1 expression). **D)** Percent Spliced In (PSI) of tuberin S981 across different cancers from TCGA and ESRP1 knockdown experiments from Yang et al. [34]. **E)** Progression-free survival of prostate cancer patients with high ESRP1 expression ($z \geq 1$) and/or low PTEN expression ($z \leq 1$).

In prostate cancer, PTEN deletions are a common alteration, found in 17.3% of the TCGA cohort. When comparing progression-free survival in patients that express high levels of ESRP1 but low levels of PTEN, we found that these patients exhibited significantly worse outcomes than other patients, even for those who had high ESRP1 or low PTEN alone (Fig. 5E). This suggests a synergistic relationship between ESRP1 and the PI3K/PTEN pathway, likely in part due to SGK1 activity, phosphorylation of TSC2, and subsequent activation of mTORC1. Thus, patients with low expression of ESRP1 may contain a constitutively activated isoform of TSC2, leading to suppressed mTORC1 activity and less aggressive prostate cancers, while high expression of ESRP1 may lead to increased mTORC1 activity and proliferation in response to PI3K/PTEN pathway activity.

Discussion

This comprehensive analysis of the effect of splicing on PTMs and their linear motifs demonstrated a few key considerations for future studies. First, inclusion/exclusion of PTMs by splicing is extensive. Second, regulatory or protein interaction connections might be diversified through differential inclusion or alterations in the linear motif of PTMs nearby splice junctions or by mutually exclusive exon events. Lastly, it is important to consider different types of PTMs separately when assessing splicing, given their distinct biochemical properties, regulation, and different relationships to splicing regarding constitutive rates, tissue specificity, flanking sequence regulation, and even the type of splicing events that alter the PTMs. In addition to the systematic analysis of the global interplay between splicing and PTMs, exploring protein functional diversity within specific biological contexts driven by isoform-specific PTMs yields fruitful and new hypotheses. Here, we explored ESRP1-related prostate cancer differences, generating new hypotheses regarding the complexity that governs the behaviors of ESRP1 within prostate cancer.

Of course, both these global and cancer-specific studies highlighted a major limitation stemming from the lack of PTM annotation, including context-specific functional roles and regulatory connections of individual PTM sites, with less than 5% of PTMs being annotated with additional functional context in PhosphoSitePlus [25]. This inherently limits our current understanding of the full scope of the impact of isoform-specific PTMs on protein functional diversity. Even still, larger patterns between splicing and PTM-driven regulation still emerge. It has previously been suggested that transcription and alternative splicing are functionally coupled, and the activity of many splicing factors are regulated by post-translational modifications to induce splicing changes [55] [56]. We found distinct examples of this where many of the ESRP1-related splicing events were related to ribosomal proteins and other translation regulators, each of which was altered through differential inclusion of PTMs that may alter their interaction partners. More examples of potential signaling-transcription-splicing cross-talk exist throughout literature, such as HRAS-mutant dependent splicing via SRSF proteins in lung cancer [72] and combined control of NF-κB signaling by splicing and PTMs [71]. In this work, we found a predicted coordination amongst possible SGK1/2/3 substrates that were differentially spliced in ESRP1-expressing prostate cancers, suggesting that ESRP1-induced splicing may facilitate SGK signaling in tumors where SGK is present and that there might be larger patterns in how signaling and protein interaction networks are controlled by splicing. Although we did not find a strong correlation between ESRP1 and SGK expression in the TCGA cohort, SGK1 has previously been shown to be upregulated in response to androgen receptor (AR) activity, a transcription factor that is one of the most common targets for prostate cancer treatments and shares many of the same transcriptional targets as ESRP1/2's splicing targets [57] [58]. This suggests a potential dual mechanism of regulation where AR and ESRP1 activity stimulates the expression of both SGKs and SGK's substrates (via splicing), altering SGK-related pathway activity. This relationship between SGKs and ESRP1 may not be limited to prostate cancer, as in breast cancer where ESRP1 expression also worsens patient prognosis, SGKs are amplified in some TCGA breast cancer patients and have been linked to estrogen receptor activity [59] [60]. Both ESRP1 and SGKs have been linked to resistance to therapies targeting estrogen or the estrogen receptor in breast cancer, which may be a result of both functioning in the same pathway [38] [61]. This suggests there may be a consistent mechanism of regulation that explains the tissue-dependent role of ESRP1.

Further study is needed to confirm the link between ESRP1 and SGK signaling in vitro, but these initial findings provide an excellent picture as to how splicing, post-translational modifications, and their regulatory networks may work in tandem to produce phenotypic changes in cells.

Finally, although we have lowered the barrier for integrating PTMs into splicing anaysis by developing tools for annotating experimentally observed splice events with PTMs and their functions, experimental connection at the protein level will remain a challenge. Unfortunately, the vast majority of differential spliced PTMs occur in such a way that they are effectively indistinguishable amongst isoforms by traditional mass spectrometry workflows. For the same reason that databases of PTMs predominantly annotate canonical isoforms, the current experimental approaches will be unable to attribute the peptide-wise modification to the full protein details of an isoform. However, this work also highlights the interesting flip side of this conundrum – that the relative quantification of PTM measurements by mass spectrometry could potentially be a reflection of changes in overall modifiability of that target peptide across isoforms. For example, increases in a phosphorylation site, often attributed to an increase in a phosphorylation site on a single protein isoform, might also be a reflection of increased exon inclusion of that modifiable residue, rather than an overall increase in the kinase responsible for that site. Together, this study suggests key avenues for advancement include improving annotations of modifications (their function and interaction partners) and developing the experimental and computational approaches to connect modifications to specific isoforms. Additionally, we know that the field is rapidly improving and expanding in identification of new modification sites and transcript isoform identification. Given that growth, we have developed and provided the computational pipeline developed in this work as an easily implementable tool for annotating RNA-sequencing datasets with PTM sites, as well as allow for new global analyses as our knowledge base expands.

Methods

Data acquisition and processing

Exon and coding sequences for protein-coding genes belonging to version GRCh38.p14 of Ensembl were downloaded using the Biomart web interface. Additional exon, transcript, and gene meta information was downloaded from Biomart using the pybiomart package. Transcript sequences were derived from exon sequences and exon rank information, and the coding sequence was matched to its location in the transcript. Transcripts producing a protein isoform with fewer than twenty amino acids were removed from analysis. Post-translational modifications and other protein-level information associated with UniProt proteins were downloaded from ProteomeScout [1] and PhosphoSitePlus [25]. Protein sequences from Ensembl and ProteomeScout were compared and any discrepancies were removed from analysis. Transcripts that coded for the same protein sequence were collapsed into a single isoform entry. TRIFID functional scores associated with each transcript were downloaded from the APPRIS database [64].

Mapping PTMs to the Genome and Projecting them onto Alternative Isoforms

In order to map post-translational modifications to their location in the genome, coding sequences were first translated into the corresponding amino acid sequences using Biopython. Coding sequences that did not start with a start codon or had incomplete codons were removed from analysis. Based on the position of the PTM site in the protein and the location of the coding sequence in the transcript, the transcript location of the codon responsible for the modifiable residue was obtained and the corresponding exon containing that codon identified. Using the genomic coordinates of the exon of interest, the PTM could then be mapped to its location in the genome. Next, to project PTM sites onto alternative transcripts and protein isoforms in Ensemble, we identified all exons in the transcriptome whose genomic coordinates contained the mapped PTM location. For each of these exons, the position of the PTM in the alternative protein was determined using location of the exon and start of the coding sequence in the alternative transcript. With PTMs projected, we validated that no frame shifts were

introduced that would lead to a change in the residue associated with that position. In the rare case where a PTM site exists at the splice junction and is coded for by two exons, we checked for conservation of at least one of the two contributing exons in the alternative transcript, then validated that the residue remained unchanged. We call the PTMs that were successfully projected on an alternative transcript a "prospective" PTM site, as the majority of these sites have not been experimentally validated. See Supplementary Figure 3 for a toy example of this process. For the whole transcriptome, this pipeline has been provided as a python package called ExonPTMapper. For projecting PTMs onto splice events, such as those in SpliceSeq, we use a streamlined version of this approach (PTM-POSE) which projects PTMs onto the genomic region of an identified splice event, rather than a distinct exon, and do not consider cases where a PTM might exist directly on a splice junction.

Identifying Splice Events

When thinking about the different types of splice events that may impact PTMs, we focused predominantly on skipped exon events, alternative splice sites, and mutually exclusive exon events, as these are the events that can lead to loss of PTM sites and are a direct result of splicing. Given that most PTM-level information used in this work is documented on canonical UniProt isoforms, splice events were defined relative to the transcripts associated with the canonical UniProt isoform. Genes for which there was not a clear canonical Uniprot isoform were excluded from this analysis. For each exon associated with the canonical isoform, we first checked to see if the Ensembl exon ID matched any exons in the alternative transcripts. In cases where there were no matching IDs, we compared the genomic location of the exon in the canonical isoform to the genomic locations of exons in the alternative transcript. From this, we identified cases where 1) an alternative exon had the same genomic coordinates ("Conserved"), 2) no alternative exons have any overlapping genomic coordinates ("Skipped"), 3) or an alternative exon has partially overlapping genomic coordinates ("Alternative Splice Site"). For "Conserved" cases, we checked to see if the protein sequences coded for by the exon within both the canonical and alternative isoform were the same, or if they had been altered due to a frameshift or alternative promoter. These cases were removed from analysis, so that focus was specifically on splicing mechanisms. For exons that were identified as "Skipped" in an alternative transcript, we assessed whether there was potential for a mutually exclusive exon event, using criteria defined by Pillman et al. [20]. In brief, we identified exons in the alternative transcript as candidate mutually exclusive exons if they were adjacent to the skipped exon location and not found in the canonical isoform. We then defined true mutually exclusive exons as those that had a length difference of no more than 20 amino acids (60 nucleotides) and a sequence similarity of 15% or more. Sequence similarity was determined through global pairwise alignment between the canonical exon and MXE candidate, using gap penalties of -10 when introducing gaps and -2 for extending gaps. Alignment was performed using Biopython [65].

Calculating PTM and flanking sequence conservation across isoforms

For each known PTM, we calculated the fraction of alternative transcripts associated with the canonical isoform for which the prospective PTM site was found. PTM sites that were found in all isoforms of a given gene were defined as "constitutive". Given that multiple transcripts can code for the same protein sequences, we collapsed transcripts with matching protein sequences into a single isoform entry prior to calculating conservation. We could then calculate a "constitutive rate" for different modification types, which is the fraction of PTMs of that modification type that could be defined as constitutive. For cases in which transcripts were filtered out based on different functional criteria, potentially dysfunctional transcripts were removed from analysis prior to calculating constitutive rates (Supplementary Figure 8).

In order to compare flanking sequences between canonical and alternative isoforms in the Ensembl genome, we obtained the flanking sequences based on the projected location of the PTM in the alternative isoform, usually considering the five amino acids on either side of the PTM unless mentioned otherwise. To obtain tryptic fragments, we identified the nearest lysine and/or arginine residues to the PTM that are not proceeded by a proline, as this is the preferential cut sites for the enzyme trypsin [66]. We then compared the flanking sequences between the canonical and alternative isoform and identified

cases in which the flanking sequence did not match (at least one amino acid was different). When calculating the overall rate of alteration for these flanking sequences, we only considered PTM sites that were successfully projected onto the alternative isoform of interest (i.e. it does not incorporate PTMs that are excluded from the isoform entirely).

In order to identify cases of changed flanking sequences in prostate cancer, we projected PTMs onto the flanking exons (exons immediately upstream or downstream of the spliced regions) of each significant event in the TCGASpliceSeq data. For PTMs that were within fifteen base pairs of the splice junction (five amino acids), we used the Ensembl REST API to extract the DNA sequence associated with both the flanking regions and the spliced region. We then stitched together two DNA sequences, one which includes the spliced region (inclusion region) and one which excludes the spliced region (exclusion region). We translated these DNA sequences into amino acid sequences, using the PTM location to define the reading frame for translation (assume no frame shifts introduced). If extracted flanking sequences were too short (less than five amino acids on either side), the PTM was removed. We then compared the flanking sequences of the inclusion and exclusion regions and identified cases in which the flanking sequence was different.

Generating the null splicing model for comparison with patterns of modifications

In order to determine if the observed splicing control of PTMs was unique to modifiable residues and not a general property of all residues, we developed a null model in which "modifiable" residues are randomly distributed across the transcriptome. For each modification type, we randomly sampled an identical number of residues as there are the modification of interest, with the same distribution of the specific amino acids that can be modified. For example, for the phosphorylation null model, we randomly sampled a total of 223,659 residues (131,236 serines, 54,358 threonines, 38,025 tyrosines), the same number of phosphorylation sites that have been observed in the proteome. With the random sample of residues, we then repeated the same analysis as done for the true modifiable residues (constitutive rate, flanking sequence identity, etc.). We repeated this process 100 times for each modification type to create a null model distribution of constitutive rates and altered flanking sequence rates. Finally, we considered the rate statistically significant if the observed value was found to have a higher or lower value than all but five or fewer of the random null models ($p \leq 0.05$).

Identifying tissue-specific PTM sites

Tissue-specificity data was downloaded from three different publications which used different approaches to identifying for tissue-specific exons and/or transcripts [21] [10] [22]. In cases in which only tissue-specific transcripts were reported, we compared the two transcript isoforms to identify tissue-specific exons, excluding alternative splice site events. Across all tissue-specific exons, density of PTMs, defined as the number of PTM sites per 1000 residues, were calculated by counting the number of PTM sites whose genomic coordinates fall within the tissue-specific exons and dividing that number by the amino acid length of each exon. Amino acid length excluded residues which were found at splice junctions and were coded for by two exons. PTM density was then normalized to the density of PTMs found across the entire proteome, defined as the $\frac{total\ number\ of\ PTM\ sites\ in\ canonical\ isoforms}{total\ number\ of\ residues\ in\ canonical\ isoforms}$. Thus, a normalized density greater than one indicates a higher density of PTM sites then expected relative to the entire proteome.

Predicting changes in kinase interactions using Kinase Library

To assess how changes to a PTM's flanking sequence may impact could impact which proteins interact with that site, we turned to a kinase motif enrichment tool, Kinase Library [23]. To get scores from Kinase Library, we first restricted our analysis to prospective PTMs in alternative isoforms with altered flanking sequences (based on five amino acids) and a matching tryptic fragment. In total, we scored a total of 1812 different events. For each, we used the Kinase Library web interface to score the flanking

sequence in both the canonical and alternative isoforms, and then extracted the 'site percentile', which indicates where the phosphorylation site ranks relative to all other phosphorylation sites in the phosphoproteome scored for that kinase. We then calculated the change in site percentile from the canonical to alternative sequence for the entire transcriptome, or between the inclusion and exclusion flanking sequences for specific splice events in prostate cancer. For the latter, we further normalized the change in site percentile by the change in percent spliced in for the given exon to capture the effect of the splicing event on the kinase interaction.

Identifying PTM sites with variable inclusion in ESRP1-high and -low prostate cancer patients

In order to identify PTM sites specific to individual prostate cancer patients from The Cancer Genome Atlas (TCGA), "percent spliced in" values (PSI) specific to each patient were downloaded from the TCGASpliceSeq database, which indicate the fraction of transcripts within a given patient for which the exon of interest is included [43]. PSI values were calculated using SpliceSeq, an algorithm designed to predict alternative splicing patterns from RNA sequencing data, like the ones generated by TCGA [43]. While the majority of this work relies on Ensembl hg38 coordinates, TCGASpliceSeq reports exons in terms of hg19 coordinates, so genomic coordinates of PTM sites were converted into the hg19 coordinates system using the pyliftover python package. From these genomic locations, we projected PTMs onto SpliceSeq exons using the streamlined version of the procedure used for all Ensembl transcripts, PTM-POSE, which only requires the genomic coordinates of the splice event quantified by SpliceSeq. Next, we downloaded patient-specific expression of ESRP1 mRNA across the from cBioPortal [44]. We also downloaded survival data and alteration data for ESRP1 and PTEN from cBioPortal. ESRP1-high and low patients were defined as patients with ESRP1 mRNA expression greater or less than one standard deviation from the mean expression across the TCGA cohort, respectively. Finally, we compared inclusion of SpliceSeq exons across the ESRP1-low and -high groups and identified statistically significant differences using a Mann Whitney U test and Benjamini-Hochberg FDR correction ($p \leq 0.01$, $r \geq 0.25$). PTM sites projected onto these differentially included exons were then also defined as differentially included. To identify PTMs that have altered flanking sequences as a result of each splicing event, we instead identified the PTMs associated with flanking exon regions (regions immediately adjacent to spliced region). For each PTM found within 5 amino acids (or 15 nucleotides) from the splice junction, we translated the flanking sequence based on whether the spliced region was included or excluded, assuming no frame shifts had been introduced into the protein isoform, which we found to be a rare event across the entire transcriptome.

PTM- and gene-level enrichment analysis

PTM-site specific enrichment analysis was performed utilizing annotations downloaded from PhosphoSitePlus [25] and Exon Ontology [53]. Enrichment for a function or process was assessed using a Fisher's Exact test, where the background population either included all PTM sites in canonical isoforms (as in Fig. 2) or all PTMs identified in the TCGA prostate cancer SpliceSeq dataset (as in Fig4). P-values were corrected using Benjamini-Hochberg false positive correction [45]. Gene-specific enrichment analysis was performed using the Enrichr wrapper from the gseapy python package [67] [68] [69]. Gene sets from Gene Ontology and Reactome were used [48] [73]. The background population was defined as all genes identified in the TCGA prostate cancer SpliceSeq dataset (as in Fig. 4).

To identify kinases with enriched substrates regulated by ESRP1 (Supplementary Figure 17), we utilized kinase-substrate networks of known connections from PhosphoSitePlus [25] and RegPhos [74] or predicted connections from KSTAR [70]. In both cases, a one-tailed hypergeometric test was used to assess statistically significant enrichment of a kinase's substrates across ESRP1-regulated PTMs, with all phosphorylation sites identified in the TCGA SpliceSeq data used as the background population. For the KSTAR networks, the median p-value enrichment was extracted across the ensemble of 50 predicted kinase-substrate networks to obtain a final enrichment value.

Constructing a PTM-associated interaction network

In order to identify protein-interactions potentially impacted by splicing-controlled PTMs, we pulled interaction information from PhosphoSitePlus [25], RegPhos [74], and PTMcode [77]. For each resource, we converted the information into common UniProt accession numbers and then combined the three resources into a single interaction network, with each node being a specific PTM site facilitating the interaction. We then identified the differentially spliced PTMs from the TCGA prostate cancer cohort that could be identified in the interaction network. From this network, we condensed the PTM sites into single protein nodes and defined as either a gained interaction (PTM sites are included in ESRP1-high patients) or lost interaction (PTM sites are excluded in ESRP1-high patients). For the few cases where the PTM was indicated to disrupt the interaction, we defined the interaction as anti-correlated with inclusion. The proteins with the highest number of edges (interactions gained or lost) were then reported.

To determine the impacts of altered flanking sequences on protein interactions, we utilized linear motif classes annotated in the Eukaryotic Linear Motif database (ELM), including the preferred linear motifs for common ligand binding domains [76]. ELM reports each classes motif as a regular expression, which we then used to search for the presence of each motif within the five amino acid flanking sequence of a PTM site, repeating this for the flanking sequence that occurs during the inclusion and exclusion events. We identified cases in which a specific linear motif was found in only one event (i.e. either when the adjacent exon is excluded or when it is included, but not both). Further, for linear motifs that specifically bind to a type of PTM, such as SH2 domains or 14-3-3 proteins, we checked to make sure the motif aligned with the position of the PTM site in the flanking sequence, as ELM motifs do not account for modified residues.

Data and Code Availability

All data generated for this work has been deposited in Figshare at `https://doi.org/10.6084/m9.figshare.24969993`. The computational pipeline used to map PTMs onto transcripts and their genomic location and project them onto alternative transcripts is freely available at the following github repository: `https://github.com/NaegleLab/ExonPTMapper`, v0.2.0. Further, we have developed a streamlined version this code for use with most splicing quantification tools, called PTM-POSE `https://github.com/NaegleLab/PTM-POSE`, v0.1.0. All additional code and analysis done for this work can be found at a separate github repository here: `https://github.com/NaegleLab/PTM_Splicing_Analysis`, v2.0.0.

Acknowledgements

We would like to thank Noah Perry and Mete Civelek for their guidance and input throughout the development of this work. Additionally, we would like to thank Ben Jordan for early mentorship of undergraduates working on this project. We would also like to thank the Systems Biology Community at the University of Virginia for their invaluable feedback on the data analysis and the manuscript figures. The results shown here are in part based upon data generated by the TCGA Research Network: https://www.cancer.gov/tcga. Research reported in this publication was supported by the National Institute Of General Medical Sciences of the National Institutes of Health under Award Number R35GM138127 and by the National Science Foundation Graduate Research Fellowship under Grant No. 1842490. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or the National Science Foundation.

Competing Interests

The authors declare that they have no competing interests.

References

1. Matlock, M. K., Holehouse, A. S. & Naegle, K. M. Proteomescout: A repository and analysis resource for post-translational modifications and proteins. *Nucleic Acids Research* (2015).

2. Mujtaba, S., Zeng, L. & Zhou, M. M. Structure and acetyl-lysine recognition of the bromodomain. *Oncogene 2007 26:37* **26**, 5521–5527 (2007). URL https://www.nature.com/articles/1210618.

3. Reinhardt, H. C. & Yaffe, M. B. Phospho-ser/thr-binding domains: navigating the cell cycle and dna damage response. *Nature Reviews Molecular Cell Biology 2013 14:9* **14**, 563–580 (2013). URL https://www.nature.com/articles/nrm3640.

4. Dikic, I., Wakatsuki, S. & Walters, K. J. Ubiquitin-binding domains — from structures to functions. *Nature Reviews Molecular Cell Biology 2009 10:10* **10**, 659–671 (2009). URL https://www.nature.com/articles/nrm2767.

5. Akiva, E., Friedlander, G., Itzhaki, Z. & Margalit, H. A dynamic view of domain-motif interactions. *PLOS Computational Biology* **8**, e1002341 (2012). URL https://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.1002341.

6. Tompa, P., Davey, N. E., Gibson, T. J. & Babu, M. M. A million peptide motifs for the molecular biologist (2014). URL http://dx.doi.org/10.1016/j.molcel.2014.05.032.

7. Wang, E. T. *et al.* Alternative isoform regulation in human tissue transcriptomes. *Nature* **456**, 470–476 (2008). URL http://www.nature.com/articles/nature07509.

8. Pan, Q., Shai, O., Lee, L. J., Frey, B. J. & Blencowe, B. J. Deep surveying of alternative splicing complexity in the human transcriptome by high-throughput sequencing. *Nature Genetics 2008 40:12* **40**, 1413–1415 (2008). URL https://www.nature.com/articles/ng.259.

9. Comoletti, D. *et al.* Characterization of the interaction of a recombinant soluble neuroligin-1 with neurexin-1β. *Journal of Biological Chemistry* **278**, 50497–50505 (2003). URL http://www.jbc.org/article/S0021925820755705/fulltexthttp://www.jbc.org/article/S0021925820755705/abstracthttps://www.jbc.org/article/S0021-9258(20)75570-5/abstract.

10. Buljan, M. *et al.* Tissue-specific splicing of disordered segments that embed binding motifs rewires protein interaction networks. *Molecular Cell* **46**, 871–883 (2012).

11. Merkin, J., Russell, C., Chen, P. & Burge, C. B. Evolutionary dynamics of gene and isoform regulation in mammalian tissues. *Science* **338**, 1593–1599 (2012). URL https://www.science.org/doi/10.1126/science.1228186.

12. Weatheritt, R. J., Davey, N. E. & Gibson, T. J. Linear motifs confer functional diversity onto splice variants. *Nucleic Acids Research* **40**, 7123–7131 (2012). URL https://dx.doi.org/10.1093/nar/gks442.

13. Wang, X. *et al.* Detection of proteome diversity resulted from alternative splicing is limited by trypsin cleavage specificity. *Molecular & Cellular Proteomics : MCP* **17**, 422 (2018). URL https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5836368/https://onedrive.live.com/view.aspx?resid=D7F68FF2A2AD0CF4%2125440&id=documents&wd=target%28Papers.one%7C2571AE6F-B743-44CC-925E-F28A5791D905%2FDiversityLimitedbyTrypsinCleavage%7CD4FF7E86-F077-4AF.

14. Goldtzvik, Y., Sen, N., Lam, S. D. & Orengo, C. Protein diversification through post-translational modifications, alternative splicing, and gene duplication. *Current Opinion in Structural Biology* **81**, 102640 (2023). URL https://linkinghub.elsevier.com/retrieve/pii/S0959440X23001148.

15. Colak, R. *et al.* Distinct types of disorder in the human proteome: Functional implications for alternative splicing. *PLOS Computational Biology* **9**, e1003030 (2013). URL https://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.1003030.

16. Zhou, J., Zhao, S. & Dunker, A. K. Intrinsically disordered proteins link alternative splicing and post-translational modifications to complex cell signaling and regulation. *Journal of Molecular Biology* **430**, 2342–2359 (2018).

17. Cunningham, F. *et al.* Ensembl 2022. *Nucleic Acids Research* **50**, D988–D995 (2022). URL https://dx.doi.org/10.1093/nar/gkab1049.

18. Bateman, A. *et al.* Uniprot: the universal protein knowledgebase in 2021. *Nucleic Acids Research* **49**, D480–D489 (2021). URL https://academic.oup.com/nar/article/49/D1/D480/6006196.

19. Pozo, F. *et al.* Assessing the functional relevance of splice isoforms. *NAR Genomics and Bioinformatics* **3**, 1–16 (2021). URL https://academic.oup.com/nargab/article/3/2/lqab044/6281449.

20. Pillmann, H., Hatje, K., Odronitz, F., Hammesfahr, B. & Kollmar, M. Predicting mutually exclusive spliced exons based on exon length, splice site and reading frame conservation, and exon sequence homology. *BMC Bioinformatics* **12**, 1–16 (2011). URL https://bmcbioinformatics.biomedcentral.com/articles/10.1186/1471-2105-12-270. Constraints on MXEs¡br/¿- MXE's must have the same length (difference less than 20aa)¡br/¿- Must have conserved splice site patterns (GT-AG, GC-AG, GG-AG, AT-AC) OR ¡br/¿- Reading frame must be conserved¡br/¿- They must show sequence similarity (¿15% similarity)¡br/¿¡br/¿.

21. Gonzàlez-Porta, M., Frankish, A., Rung, J., Harrow, J. & Brazma, A. Transcriptome analysis of human tissues and cell lines reveals one dominant transcript per gene. *Genome Biology* **14**, 1–11 (2013). URL https://genomebiology.biomedcentral.com/articles/10.1186/gb-2013-14-7-r70.

22. Rodriguez, J. M., Pozo, F., Domenico, T. D., Vazquez, J. & Tress, M. L. An analysis of tissue-specific alternative splicing at the protein level. *PLOS Computational Biology* **16**, e1008287 (2020). URL https://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.1008287.

23. Johnson, J. L. *et al.* An atlas of substrate specificities for the human serine/threonine kinome. *Nature 2023 613:7945* **613**, 759–766 (2023). URL https://www.nature.com/articles/s41586-022-05575-3.

24. Fischle, W., Franz, H., Jacobs, S. A., Allis, C. D. & Khorasanizadeh, S. Specificity of the chromodomain y chromosome family of chromodomains for lysine-methylated ark(s/t) motifs. *Journal of Biological Chemistry* **283**, 19626–19635 (2008). URL http://www.jbc.org/article/S0021925820814161/fulltexthttp://www.jbc.org/article/S0021925820814161/abstracthttps://www.jbc.org/article/S0021-9258(20)81416-1/abstract.

25. Hornbeck, P. V. *et al.* Phosphositeplus: a comprehensive resource for investigating the structure and function of experimentally determined post-translational modifications in man and mouse. *Nucleic acids research* **40**, D261–70 (2012). URL http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3245126&tool=pmcentrez&rendertype=abstract.

26. Scotti, M. M. & Swanson, M. S. Rna mis-splicing in disease (2016).

27. Wang, G. S. & Cooper, T. A. Splicing in disease: disruption of the splicing code and the decoding machinery. *Nature Reviews Genetics 2007 8:10* **8**, 749–761 (2007). URL https://www.nature.com/articles/nrg2164.

28. Fletcher, S. *et al.* Antisense suppression of donor splice site mutations in the dystrophin gene transcript. *Molecular Genetics and Genomic Medicine* **1**, 162–173 (2013).

29. Voena, C. *et al.* Oncogenic alk regulates emt in non-small cell lung carcinoma through repression of the epithelial splicing regulatory protein 1. *Oncotarget* **7**, 33316 (2016). URL /pmc/articles/PMC5078097//pmc/articles/PMC5078097/?report=abstracthttps: //www.ncbi.nlm.nih.gov/pmc/articles/PMC5078097/.

30. Lee, H. H. *et al.* Epithelial splicing regulatory protein (espr1) expression in an unfavorable prognostic factor in prostate cancer patients. *Frontiers in Oncology* **10**, 2274 (2020).

31. Liu, Y. *et al.* Underlying mechanisms of epithelial splicing regulatory proteins in cancer progression. *Journal of Molecular Medicine 2022 100:11* **100**, 1539–1556 (2022). URL https://link.springer.com/article/10.1007/s00109-022-02257-5.

32. Freytag, M. *et al.* Epithelial splicing regulatory protein 1 and 2 (esrp1 and esrp2) upregulation predicts poor prognosis in prostate cancer. *BMC Cancer* **20**, 1–12 (2020). URL https://bmccancer.biomedcentral.com/articles/10.1186/s12885-020-07682-8.

33. Gerhauser, C. *et al.* Molecular evolution of early-onset prostate cancer identifies molecular risk markers and clinical trajectories. *Cancer Cell* **34**, 996–1011.e8 (2018). - ESRP1 duplication correlated with ESRP1 expression (¿1.5 fold change)¡br/¿- Some patients exhibited both ESRP1 and Myc amplifications, but only ESRP1 showed changes at mRNA level¡br/¿- ESRP1 duplications exhibited higher Gleason scores (indicative of worse tumor stages)¡br/¿- Increased ESRP1 protein correlated with higher proliferation right by Ki67 labeling¡br/¿- Showed that it is an independent biomarker¡br/¿- Showed strong relationship between PEPCI (Purity-Adjusted Epigenetic Prostate CAncer Index) and ESRP1 amplification.

34. Yang, Y. *et al.* Determination of a comprehensive alternative splicing regulatory network and combinatorial regulation by key factors during the epithelial-to-mesenchymal transition. *Molecular and Cellular Biology* **36**, 1704–1719 (2016). URL https://www.tandfonline.com/doi/abs/10.1128/MCB.00019-16.

35. Warzecha, C. C., Sato, T. K., Nabet, B., Hogenesch, J. B. & Carstens, R. P. Esrp1 and esrp2 are epithelial cell-type-specific regulators of fgfr2 splicing. *Molecular Cell* **33**, 591–601 (2009).

36. Brown, R. L. *et al.* Cd44 splice isoform switching in human and mouse epithelium is essential for epithelial-mesenchymal transition and breast cancer progression. *The Journal of Clinical Investigation* **121**, 1064–1074 (2011). URL http://www.jci.org.

37. Ishii, H. *et al.* Epithelial splicing regulatory proteins 1 (esrp1) and 2 (esrp2) suppress cancer cell motility via different mechanisms. *Journal of Biological Chemistry* **289**, 27386–27399 (2014). URL http://www.jbc.org/article/S0021925820483784/fulltexthttp: //www.jbc.org/article/S0021925820483784/abstracthttps: //www.jbc.org/article/S0021-9258(20)48378-4/abstract.

38. Gökmen-Polar, Y. *et al.* Splicing factor esrp1 controls er-positive breast cancer by altering metabolic pathways. *EMBO reports* **20**, e46078 (2019). URL https://onlinelibrary.wiley.com/doi/full/10.15252/embr.201846078.

39. Nanni, M., Ranieri, D., Persechino, F., Torrisi, M. R. & Belleudi, F. The aberrant expression of the mesenchymal variant of fgfr2 in the epithelial context inhibits autophagy. *Cells 2019, Vol. 8, Page 653* **8**, 653 (2019). URL https://www.mdpi.com/2073-4409/8/7/653/htm.

40. Soucek, T., Pusch, O., Wienecke, R., DeClue, J. E. & Hengstschläger, M. Role of the tuberous sclerosis gene-2 product in cell cycle control. *Journal of Biological Chemistry* **272**, 29301–29308 (1997). URL http://www.jbc.org/article/S0021925818508863/fulltext.

41. Zhang, H. *et al.* Loss of tsc1/tsc2 activates mtor and disrupts pi3k-akt signaling through downregulation of pdgfr. *The Journal of Clinical Investigation* **112**, 1223–1233 (2003).

42. Pai, G. M. *et al.* Tsc loss distorts dna replication programme and sensitises cells to genotoxic stress. *Oncotarget* **7**, 85365 (2016). URL /pmc/articles/PMC5356742//pmc/articles/PMC5356742/?report=abstracthttps://www.ncbi.nlm.nih.gov/pmc/articles/PMC5356742/.

43. Ryan, M. *et al.* Tcgaspliceseq a compendium of alternative mrna splicing in cancer. *Nucleic Acids Research* **44**, D1018–D1022 (2016). URL https://academic.oup.com/nar/article/44/D1/D1018/2503095.

44. Cerami, E. *et al.* The cbio cancer genomics portal: An open platform for exploring multidimensional cancer genomics data. *Cancer Discovery* **2**, 401–404 (2012).

45. Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B (Methodological)* **57**, 289–300 (1995). URL https://onlinelibrary.wiley.com/doi/full/10.1111/j.2517-6161.1995.tb02031.xhttps://onlinelibrary.wiley.com/doi/abs/10.1111/j.2517-6161.1995.tb02031.xhttps://rss.onlinelibrary.wiley.com/doi/10.1111/j.2517-6161.1995.tb02031.x.

46. Szklarczyk, D. *et al.* String v11: protein–protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Research* **47**, D607 (2019). URL /pmc/articles/PMC6323986//pmc/articles/PMC6323986/?report=abstracthttps://www.ncbi.nlm.nih.gov/pmc/articles/PMC6323986/.

47. Kanehisa, M. & Goto, S. Kegg: Kyoto encyclopedia of genes and genomes. *Nucleic Acids Research* **28**, 27–30 (2000). URL https://dx.doi.org/10.1093/nar/28.1.27.

48. Ashburner, M. *et al.* Gene ontology: tool for the unification of biology. *Nature Genetics 2000 25:1* **25**, 25–29 (2000). URL https://www.nature.com/articles/ng0500_25http://fruitfly.bdgp.berkeley.eduhttp://genome-www.stanford.eduhttp://www.informatics.jax.org.

49. Kjer-Hansen, P. & Weatheritt, R. J. The function of alternative splicing in the proteome: rewiring protein interactomes to put old functions into new contexts. *Nature Structural & Molecular Biology 2023 30:12* **30**, 1844–1856 (2023). URL https://www.nature.com/articles/s41594-023-01155-9.

50. Louadi, Z. *et al.* Digger: exploring the functional role of alternative splicing in protein interactions. *Nucleic Acids Research* **49**, D309 (2021). URL /pmc/articles/PMC7778957//pmc/articles/PMC7778957/?report=abstracthttps://www.ncbi.nlm.nih.gov/pmc/articles/PMC7778957/.

51. Louadi, Z. *et al.* Functional enrichment of alternative splicing events with nease reveals insights into tissue identity and diseases. *Genome Biology 2021 22:1* **22**, 1–22 (2021). URL https://genomebiology.biomedcentral.com/articles/10.1186/s13059-021-02538-1.

52. Fuente, L. D. L. *et al.* Tappas: A comprehensive computational framework for the analysis of the functional impact of differential splicing. *Genome Biology* **21**, 1–32 (2020). URL https://genomebiology.biomedcentral.com/articles/10.1186/s13059-020-02028-w.

53. Tranchevent, L. C. *et al.* Identification of protein features encoded by alternative exons using exon ontology. *Genome Research* **27**, 1087–1097 (2017). URL https://genome.cshlp.org/content/27/6/1087.fullhttps://genome.cshlp.org/content/27/6/1087https://genome.cshlp.org/content/27/6/1087.abstract.

54. ŠoŠtarić, N. & van Noort, V. Molecular dynamics shows complex interplay and long-range effects of post-translational modifications in yeast protein interactions. *PLOS Computational Biology* **17**, e1008988 (2021). URL https://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.1008988.

55. Kornblihtt, A. R., Mata, M. D. L., Fededa, J. P., Muñoz, M. J. & Nogués, G. Multiple links between transcription and splicing. *RNA* **10**, 1489–1498 (2004). URL http://rnajournal.cshlp.org/content/10/10/1489.fullhttp://rnajournal.cshlp.org/content/10/10/1489http://rnajournal.cshlp.org/content/10/10/1489.abstract.

56. Kretova, M., Selicky, T., Cipakova, I. & Cipak, L. Regulation of pre-mrna splicing: Indispensable role of post-translational modifications of splicing factors. *Life 2023, Vol. 13, Page 604* **13**, 604 (2023). URL https://www.mdpi.com/2075-1729/13/3/604/htmhttps://www.mdpi.com/2075-1729/13/3/604.

57. Shanmugam, I. *et al.* Serum/glucocorticoid-induced protein kinase-1 facilitates androgen receptor-dependent cell survival. *Cell Death & Differentiation 2007 14:12* **14**, 2085–2094 (2007). URL https://www.nature.com/articles/4402227.

58. Shah, K. *et al.* Androgen receptor signaling regulates the transcriptome of prostate cancer cells by modulating global alternative splicing. *Oncogene 2020 39:39* **39**, 6172–6189 (2020). URL https://www.nature.com/articles/s41388-020-01429-2.

59. Toska, E., Castel, P., Leslie, C. S., Scaltriti, M. & Baselga, J. Pi3k inhibition activates sgk1 via a feedback loop to promote chromatin-based regulation of er-dependent gene expression. *CellReports* **27**, 294–306.e5 (2019). URL https://doi.org/10.1016/j.celrep.2019.02.111.

60. Wang, Y. *et al.* Sgk3 is an estrogen-inducible kinase promoting estrogen-mediated survival of breast cancer cells. *Molecular Endocrinology* **25**, 72–82 (2011). URL https://dx.doi.org/10.1210/me.2010-0294.

61. Wang, Y. *et al.* Sgk3 sustains erα signaling and drives acquired aromatase inhibitor resistance through maintaining endoplasmic reticulum homeostasis. *Proceedings of the National Academy of Sciences of the United States of America* **114**, E1500–E1508 (2017). URL https://www.pnas.org/doi/abs/10.1073/pnas.1612991114.

62. Oltean, S. & Bates, D. O. Hallmarks of alternative splicing in cancer. *Oncogene 2014 33:46* **33**, 5311–5318 (2013). URL https://www.nature.com/articles/onc2013533.

63. Wu, X.-J. *et al.* The splicing factor sr2 is an important virulence factor of toxoplasma gondii. *Frontiers in Microbiology* **14**, 1302512 (2023). URL https://www.frontiersin.org/articles/10.3389/fmicb.2023.1302512/full.

64. Rodriguez, J. M. *et al.* Appris: annotation of principal and alternative splice isoforms. *Nucleic acids research* **41** (2013). URL https://pubmed.ncbi.nlm.nih.gov/23161672/.

65. Cock, P. J. *et al.* Biopython: Freely available python tools for computational molecular biology and bioinformatics. *Bioinformatics* **25**, 1422–1423 (2009).

66. Olsen, J. V., Ong, S. E. & Mann, M. Trypsin cleaves exclusively c-terminal to arginine and lysine residues. *Molecular and Cellular Proteomics* **3**, 608–614 (2004). URL http://www.mcponline.org/article/S1535947620338779/fulltexthttp://www.mcponline.org/article/S1535947620338779/abstracthttps://www.mcponline.org/article/S1535-9476(20)33877-9/abstract.

67. Fang, Z., Liu, X. & Peltz, G. Gseapy: a comprehensive package for performing gene set enrichment analysis in python. *Bioinformatics* **39** (2023). URL https://dx.doi.org/10.1093/bioinformatics/btac757.

68. Chen, E. Y. *et al.* Enrichr: interactive and collaborative html5 gene list enrichment analysis tool. *BMC bioinformatics* **14** (2013). URL https://pubmed.ncbi.nlm.nih.gov/23586463/.

69. Kuleshov, M. V. *et al.* Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. *Nucleic acids research* **44**, W90–W97 (2016). URL https://pubmed.ncbi.nlm.nih.gov/27141961/.

70. Crowl, S., Jordan, B. T., Ahmed, H., Ma, C. X. & Naegle, K. M. Kstar: An algorithm to predict patient-specific kinase activities from phosphoproteomic data. *Nature Communications 2022 13:1* **13**, 1–16 (2022). URL https://www.nature.com/articles/s41467-022-32017-5.

71. Leeman, J. R. & Gilmore, T. D. Alternative splicing in the nf-κb signaling pathway. *Gene* **423**, 97–107 (2008). URL https://dx.doi.org/10.1016/J.GENE.2008.07.015.

72. Lo, A., McSharry, M. & Berger, A. H. Oncogenic kras alters splicing factor phosphorylation and alternative splicing in lung cancer. *BMC Cancer* **22**, 1–13 (2022). URL https://bmccancer.biomedcentral.com/articles/10.1186/s12885-022-10311-1.

73. Jassal, B. *et al.* The reactome pathway knowledgebase. *Nucleic Acids Research* **48**, D498 (2020). URL https://academic.oup.com/nar/article/48/D1/D498/5613674.

74. Huang, K. Y. *et al.* Regphos 2.0: an updated resource to explore protein kinase–substrate phosphorylation networks in mammals. *Database* **2014**, 1–12 (2014). URL https://dx.doi.org/10.1093/database/bau034.

75. Kim, T. H., Leslie, P. & Zhang, Y. Ribosomal proteins as unrevealed caretakers for cellular stress and genomic instability. *Oncotarget* **5**, 860 (2014). URL https://www.oncotarget.com/article/1784/text/.

76. Kumar, M. *et al.* Elm—the eukaryotic linear motif resource in 2020. *Nucleic Acids Research* **48**, D296–D306 (2020). URL https://dx.doi.org/10.1093/nar/gkz1030.

77. Minguez, P., Letunic, I., Parca, L. & Bork, P. Ptmcode: a database of known and predicted functional associations between post-translational modifications in proteins. *Nucleic Acids Research* **41**, D306–D311 (2013). URL https://dx.doi.org/10.1093/nar/gks1230.

78. Shen, S. *et al.* rMATS: robust and flexible detection of differential alternative splicing from replicate RNA-Seq data. *Proceedings of the National Academy of Sciences of the United States of America* **111**, E5593–E5601 (2014). URL https://www.pnas.org/doi/abs/10.1073/pnas.1419161111.

79. Cai, S. L. *et al.* Activity of TSC2 is inhibited by AKT-mediated phosphorylation and membrane partitioning. *The Journal of Cell Biology* **173**, 279 (2006). URL https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2063818/

80. Kim, P., Yang, M., Yiya, K., Zhao, W. & Zhou, X. ExonSkipDB: functional annotation of exon skipping event in human. *Nucleic Acids Research* **48**, D896–D907 (2020). URL https://dx.doi.org/10.1093/nar/gkz917.

81. Yang, C., Zang, W., Ji, Y., Li, T., Yang, Y. & Zheng, X. Ribosomal protein L6 (RPL6) is recruited to DNA damage sites in a poly(ADP-ribose) polymerase– dependent manner and regulates the DNA damage response. *Journal of Biological Chemistry* **294**, 2827–2838 (2019). URL http://www.jbc.org/article/S0021925820393789/fulltext

82. Malecki, J., Aileni, V. K., Ho, A. Y. Y., Schwarz, J., Moen, A., Sørensen, V., Nilges, B. S., Jakobsson, M. E., Leidel, S. A. & Falnes, P. O. The novel lysine specific methyltransferase METTL21B affects mRNA translation through inducible and dynamic methylation of Lys-165 in human eukaryotic elongation factor 1 alpha (eEF1A). *Nucleic Acids Research* **45**, 4370–4389 (2017). URL https://dx.doi.org/10.1093/nar/gkx002.

Supporting Information

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3

Supplementary Figure 4

Supplementary Figure 5

Supplementary Figure 6

Supplementary Figure 7

Supplementary Figure 8

Supplementary Figure 9

Supplementary Figure 10

Supplementary Figure 11

Supplementary Figure 12

Supplementary Figure 13

Supplementary Figure 14

Supplementary Figure 15

Supplementary Figure 16

Supplementary Figure 17

Supplementary Table 1

Supplementary Table 2

Supplementary Table 3