Systematic analysis of nonsense variants uncovers peptide release rate as a novel
modifier of nonsense-mediated mRNA decay efficiency

Divya Kolakada[1,2], Rui Fu[3#], Nikita Biziaev[4], Alexey Shuvalov[4], Mlana Lore[2], Amy E. Campbell[1],
Michael A. Cortázar[1], Marcin P. Sajek[1,5], Jay R. Hesselberth[1,3], Neelanjan Mukherjee[1,3], Elena
Alkalaeva[4], and Sujatha Jagannathan[1,3,6]*

[1]Department of Biochemistry and Molecular Genetics, University of Colorado Anschutz Medical
Campus, Aurora, CO 80045, USA.

[2]Molecular Biology Graduate Program, University of Colorado Anschutz Medical Campus, Aurora,
CO 80045, USA.

[3]RNA Bioscience Initiative, University of Colorado Anschutz Medical Campus, Aurora, CO 80045,
USA.

[4]Engelhardt Institute of Molecular Biology, The Russian Academy of Sciences, 119991 Moscow,
Russia.

[5]Institute of Human Genetics, Polish Academy of Sciences, Poznan, Poland

[6]Lead contact

*Corresponding Author: sujatha.jagannathan@cuanschutz.edu

[#]Current Address: New York Genome Center, New York, NY

Keywords: NMD, PTC, translation, termination, splicing

ABSTRACT

Nonsense variants underlie many genetic diseases. The phenotypic impact of nonsense variants is determined by Nonsense-mediated mRNA decay (NMD), which degrades transcripts with premature termination codons (PTCs). NMD activity varies across transcripts and cellular contexts via poorly understood mechanisms. Here, by leveraging human genetic datasets, we uncover that the amino acid preceding the PTC dramatically affects NMD activity in human cells. We find that glycine codons in particular support high levels of NMD and are enriched before PTCs but depleted before normal termination codons (NTCs). Gly-PTC enrichment is most pronounced in human genes that tolerate loss-of-function variants. This suggests a strong biological impact for Gly-PTC in ensuring robust elimination of potentially toxic truncated proteins from non-essential genes. Biochemical assays revealed that the peptide release rate during translation termination is highly dependent on the identity of the amino acid preceding the stop codon. This release rate is the most critical feature determining NMD activity across our massively parallel reporter assays. Together, we conclude that NMD activity is significantly modulated by the "window of opportunity" offered by translation termination kinetics. Integrating the window of opportunity model with the existing framework of NMD would enable more accurate nonsense variant interpretation in the clinic.

INTRODUCTION

Mutations that introduce a premature stop codon account for over 30% of inherited human diseases due to the action of nonsense-mediated RNA decay (NMD) (Miller & Pearce, 2014). NMD eliminates transcripts that contain premature termination codons (PTCs) that would otherwise generate truncated protein products. Much of our understanding of NMD comes from studies of naturally occurring nonsense mutations and genotype-phenotype correlations in genes involved in Mendelian disorders such as β-globin (Neu-Yilik et al., 2011) (*HBB*), Dystrophin (Kerr et al., 2001) (*DMD*), and Lamin A/C (van Engelen et al., 2005) (*LMNA*). For example, individuals with a nonsense mutation in early exons of *HBB* develop a recessive form of β-thalassemia due to degradation of the mutant mRNA via NMD (Chang & Kan, 1979), whereas nonsense mutations in the last exon of *HBB* cause a dominant form of the disease due to the mutant mRNA escaping NMD and leading to the production of a C-terminally truncated beta-globin protein (Kazazian et al., 1992). Such studies have helped refine the rules of NMD, including the "55-nt rule", which describes the NMD insensitivity of nonsense mutations in the last and penultimate exons up to 55 nucleotides upstream of the final exon junction complex (Popp & Maquat, 2016).

Several exceptions to these rules have also been observed: nonsense mutations found close to the start codon often escape NMD due to downstream reinitiation of translation (Inacio et al., 2004; Pereira et al., 2015; Zhang & Maquat, 1997); stop codon readthrough has been reported to contribute to the escape of some nonsense variants from NMD (Hogg & Goff, 2010; Keeling et al., 2004); and NMD efficiency has been found to vary across cell types (Linde et al., 2007), tissues (Bateman et al., 2003; Zetoune et al., 2008), and individuals (Kerr et al., 2001; Nguyen et al., 2014), offering potential explanations for tissue tropism and varying penetrance across the population of disease-causing nonsense variants. A systematic study of PTCs in the Cancer Genome Atlas database found that the 55-nt rule only predicted the fate of ~50% of PTCs; while additional factors such as exon length could account for another ~20% of NMD variability, ~30% of NMD variability is still unexplained (Lindeboom et al., 2016; Lindeboom et al., 2019). Another study investigating nonsense variants that are highly frequent in the healthy human population found that ~50% of PTC-containing transcripts escape NMD via mechanisms including alternative splicing, stop codon readthrough, and alternative translation initiation (Jagannathan & Bradley, 2016). If every premature termination event that satisfies the 50-55 nt rule does not lead to NMD, what other factors influence the probability and activity of NMD?

As premature translation termination is the key signal for NMD, termination efficiency may impact NMD. Translation termination efficiency can be conceptualized in two ways: (1) the fidelity

of termination, reflecting the likelihood of stop codon readthrough, and (2) the kinetics of
termination, reflecting the dwell time of the terminating ribosome at the stop codon. Both these
aspects are influenced by cis-acting elements (Baker & Hogg, 2017; Loughran et al., 2014;
McCaughan et al., 1995; Pierson et al., 2016; Tork et al., 2004). For example, specific elements
within the stop codon sequence context and certain viral RNA structures can reduce termination
fidelity (Baker & Hogg, 2017; Bonetti et al., 1995; Cridge et al., 2018; Loughran et al., 2014;
Mangkalaphiban et al., 2021; Tork et al., 2004). Additionally, the identity of the nucleotide
immediately after the stop codon and the peptide-tRNA hydrolysis rates of the amino acid
preceding the stop codon impact the rate of peptide release (McCaughan et al., 1995; Pierson et
al., 2016). However, whether and how local sequence context and translation termination fidelity
or kinetics influence the probability that a given transcript will undergo NMD remains to be
determined.

In this study, we hypothesized that the sequence contexts surrounding PTCs may hold the
key to understanding NMD variability among transcripts. Indeed, we found that glycine codons
are highly enriched immediately upstream of PTCs among healthy individuals and that this
sequence context potently enhances NMD activity. To systematically assess the impact of PTC
sequence context on NMD activity, we performed a massively parallel reporter assay (MPRA)
and found a broad range of NMD activities associated with different sequences. Statistical
modeling of the MPRA data identified peptide release rate during translation termination as the
major predictor of NMD activity, which we verified using biochemical assays. We propose that
variable peptide release rates alter NMD activity by modulating the kinetic window available for
NMD factors to bind and enact NMD during translation termination. Our data suggest that the Gly-
PTC context coevolved with NMD to efficiently eliminate transcripts containing premature stop
codons.

RESULTS

Glycine codons are enriched preceding PTCs in non-essential genes. Healthy individuals
harbor ~100 putative loss of function genetic variants, most of which are protein-truncating
(MacArthur et al., 2012). In previous work, we showed that such PTCs often escape NMD via
mechanisms including alternative splicing or stop codon readthrough, allowing them to rise to high
frequencies in the healthy population (minor allele frequency, MAF > 5%) (Jagannathan &
Bradley, 2016). In this study, we focused on rare variants with MAF $\leq$ 1% in healthy individuals,

reasoning that such variants are more likely to undergo NMD and yet may have varying

Figure 1. Gly is enriched preceding a PTC in non-essential genes. (**A**) Schematic for analyzing LoF
scores for rare PTCs in healthy individuals and PTCs in disease contexts. (**B-C**) Amino acid enrichment -1
to a PTC (B) or an NTC (C). (**D**) Each amino acid's frequency delta (enrichment before a PTC-NTC) at the
-1 position. (**E**) The -1 amino acid enrichment of PTCs binned into 3 categories of probability of loss-of-

function intolerance (pLI). The lower the pLI, the more tolerant the PTCs are to loss of
function. (**F**) The -1 amino acid enrichment of PTCs in disease contexts.

efficiencies. Using the gnomAD database (n=151,332, v2.1.1), which contains genetic variant
data for >700,000 healthy humans (Karczewski et al., 2020), we analyzed amino acid enrichment
within the 10 amino acids preceding PTCs (**Fig. 1A, Supp. Fig. 1A**). We compared the enrichment
of these amino acids to that of normal termination codon (NTC) contexts in the reference genome
(n=23,066, **Fig. 1A, Supp. Fig. 1B**) using the method similar to Koren *et al* (Koren et al., 2018).
Gly codons at the -1 position stood out among all examined positions as the most enriched before
PTCs and least enriched before the NTC (**Fig. 1B-C,** and **Supp. Fig. 1**). Threonine (Thr) and
proline (Pro) were the next most enriched in PTC compared to NTC at the -1 position (**Fig. 1D**).
Other amino acids show the opposite trend – enriched before an NTC and depleted before a PTC
– including asparagine (Asn), valine (Val), and aspartic acid (Asp). These data suggest that the
enrichment of Gly immediately upstream of a PTC (Gly-PTC) has a functional impact that allowed
it to rise to high frequency in healthy individuals.

To further explore the functional impact of Gly-PTC enrichment in healthy individuals, we
analyzed the relative frequency of PTC-inducing variants binned in terms of the probability of LoF
intolerance (pLI) score of their host genes (**Fig. 1A** and **1E**). The lower the pLI, the more tolerant
to loss of function a gene is and the less essential. We found that Gly-PTC contexts were enriched
in genes with lower pLI, suggesting that these PTC-containing variants were more tolerant to loss
of function (**Fig. 1E**). We also examined the amino acids enriched immediately upstream of PTCs
from the ClinVar database, which contains human genetic variants associated with disease
(n=48,847, **Fig. 1A**) (Landrum et al., 2018). Intriguingly, we found no enrichment of Gly at the -1
position of disease-associated PTCs (**Fig. 1F**). In summary, the Gly-PTC enrichment in non-
essential genes in comparison to essential or disease-relevant genes suggests a non-pathogenic
and homeostatic role for this sequence context.

Massively parallel reporter assay (MPRA) captures widespread sequence-dependent
variation in NMD activity. To systematically examine the relationship between NMD activity and
the PTC sequence context, we conducted an MPRA using a set of NMD reporters we previously
developed (Kolakada et al., 2024). These reporters contain two types of NMD targets: one
triggered by the presence of exon-junction complexes downstream of a PTC ("EJC-enhanced")
and the other by the presence of a long 3' UTR ("EJC-independent"). Because these targets rely

on related but distinct cues to trigger NMD, they might respond differently to PTC sequence
variation.

We varied the 6 nucleotides before the PTC, the PTC itself, and 1 nucleotide after the PTC in
an unbiased fashion (N_6TRRN; corresponding to 65,536 unique sequences; **Fig. 2A**) and asked
how these sequences impacted NMD activity in both EJC-independent and EJC-enhanced
reporters. Because we employed stop codons with "TRR" for array-based oligonucleotide
synthesis ("R" being a purine), 25% of our sequence library consists of a tryptophan (Trp) codon
(TGG) instead of a PTC – thereby serving as control sequences that do not undergo NMD. All
sequences were incorporated into both EJC-independent and EJC-enhanced fluorescent reporter
backbones (**Supp. Fig. 2A-B**) and integrated into a single genomic location in HEK293T cells via
Cre-Lox mediated recombination (Khandelia et al., 2011). These cells were then treated with
either DMSO or SMG1i, harvested for RNA, and subjected to targeted sequencing to quantify the
levels of each sequence context in the pool of cells. The NMD activity of an individual reporter
sequence was calculated as the $\log_2$ transcript fold change between DMSO-treated cells (NMD
active) and SMG1i-treated cells (NMD-inhibited) as measured by DESeq2 (Love et al., 2014),
thus a lower (negative) value indicates higher NMD activity, and vice versa. The ~1/4th of control
sequences containing the Trp codon rather than a stop codon were used for normalization.

The NMD activity of both EJC-independent and EJC-enhanced reporters had a bimodal
distribution with clear separation between constructs containing stop codons and the Trp-
encoding TGG codon (**Fig. 2B**): sequences containing a stop codon were left-shifted compared
to the sequences containing the Trp control, indicating that the former are targets of NMD. In the
population of sequences undergoing NMD (i.e. those containing a stop codon), there was a wide
range of NMD activity for both the EJC-independent and the EJC-enhanced reporters. EJC-
enhanced reporter expression was reduced by an average of 4.8-fold compared to the cells
treated with the NMD inhibitor (log2FC: -2.27), while the EJC-independent reporters experienced
an average 3.3-fold decrease (log2FC: -1.71) (**Fig. 2B**). These results indicate that EJC-
enhanced targets undergo more efficient NMD compared to EJC-independent targets, consistent
with prior research (Buhler et al., 2006; Chu et al., 2021; Singh et al., 2008).

Next, we investigated if the PTC sequence context could explain the wide range of observed
NMD activity. Specifically, we were interested in the contribution of the codons and amino acids
at the -1 and -2 positions, the stop codon, and the nucleotide at the +4 position of the PTC. We
constructed a linear model for each of these biological variables to determine how much it

explained the variation in NMD activity (i.e. transcript fold change) of the associated sequences.

Figure 2. MPRA shows a wide range of PTC context-dependent NMD efficiencies. (A) Design of
the MPRA. **(B)** Histogram of transcript fold change EJC-independent and EJC-enhanced NMD+ reporters
comparing DMSO treated cells to SMG1i treated cells. **(C-D)** Dot plots of the beta coefficient representing
the effect of the stop codon **(C)** and the +4 nt **(D)** on EJC-independent and EJC-enhanced NMD activity.

(E-H) Plots depicting a variable's effect on EJC-independent NMD: (**E**) Heat map for the -2 codon, (**F**) Dot
plot for the -2 amino acid, (**G**) Heat map for the -1 codon, and (**H**) Dot plot for the -1 amino acid.

The model allowed us to evaluate the effect size (beta coefficient) and statistical significance (p-
value) of individual features within a variable on NMD activity relative to a control feature. For
example, in the case of stop codons, we evaluated the effect of TAA, TAG, or TGA on NMD
activity relative to TGG (Trp). Negative beta coefficients indicated stronger NMD activity, while
positive ones indicated weaker NMD activity. It is important to note that each model (represented
by individual plots) is unique, and the beta coefficients cannot be quantitatively compared across
the plots.

As expected, our modeling showed that all stop codons caused significantly stronger NMD
activity compared to the Trp control for both EJC-independent and EJC-enhanced reporters (**Fig.
2C**). Notably, the TGA stop codon had a beta coefficient 0.2 lower than TAG, indicating a 20%
increase in NMD activity compared to the TAG stop codon for the EJC-independent reporters
(**Supp. Fig. 3A**). The NMD activity of TAA was also higher than TAG (~5%) but not to the extent
of TGA for the EJC-Independent reporters. For the EJC-enhanced reporters, both TGA and TAA
had higher NMD activity than TGA (~5%) (**Supp. Fig. 3A**). For subsequent analyses, only the
transcripts expected to undergo NMD (i.e. transcripts with stop codons) were analyzed. When
determining the effect of the +4 nt identity on NMD, we found that T, G, and C promoted ~6%
weaker NMD activity compared to A for the EJC-independent reporters. A similar effect was
observed for G compared to A for the EJC-enhanced reporters (**Fig. 2D**). Overall, the nucleotide
after the PTC only subtly impacted NMD activity.

The identity of the codon at the –2 position had a wide range of effects on NMD activity for
the EJC-independent reporters. As the –2 codon position is coded by randomized nucleotides
(NNN), it can also include all three stop codons. We found that the effects of the codons were
largely grouped by the identity of the encoded amino acid, with only a few that were significantly
different (**Fig. 2E**; all codons compared to Ala-GCA). Therefore, we next looked at the codons
based on their encoded amino acid identity in a new linear model. We found that a stop codon at
the –2 position caused the strongest increase (~40%) in NMD activity compared to the Ala control
(**Fig. 2E-F**). More subtle effects were observed across various amino acids, with an increase up
to ~10% in NMD activity with Val at the -2 position, or a reduction of up to 20% when Arg was
encoded at the -2 position.

When examining the effect of the codon at the -1 position on NMD activity, all codons except
one Leu codon, were again grouped by the encoded amino acid (**Fig. 2G**; all codons compared
to Ala-GCA). Remarkably, we found that Gly-PTC (the same context enriched in the gnomAD
database; **Fig. 1**) has the strongest effect on NMD activity, reducing it by 41% compared to Ala
(**Fig. 2G-H**). Thr exhibited almost as potent an effect as Gly (~38% enhanced compared to Ala):
both Gly-PTC and Thr-PTC resulted in much stronger NMD than a stop codon in the -1 position
(~14% enhanced compared to Ala). Indeed, Thr was the second-most enriched amino acid
encoded immediately upstream to PTCs in our gnomAD analysis (**Fig. 1D**). In contrast, Tyr-PTC
had the least efficient NMD, weakening NMD by ~25% compared to Ala-PTC (**Fig. 2G-H**).

For the EJC-enhanced reporters, there were fewer significant changes at both -1 and -2
positions. Note that the control feature for the -2 position is Arg or an Arg codon and for the -1
position it is Ala or an Ala codon. Control features were picked based on amino acids or codons
with beta coefficients in the middle range of the model. One Ala and one Lys codon at the -2
position have significant effects compared to Arg-AGA, while one Arg codon shows a significant
difference in the -1 position compared to Ala-GCA. That said, the NMD effect size range was
much smaller overall compared to the EJC-independent models. Upon aggregation of codons to
the encoded amino acids, most amino acids did not affect the efficiency of NMD at the -2 or -1
positions, except Lys and Pro at the -2 position, which decreased NMD activity by ~17% or
increased it by ~6%, respectively (**Supp Fig. 3A-D**). Interestingly, a stop codon in the -2 position
weakened NMD by ~8% compared to the control variable Arg – the opposite effect that we
observed with the EJC-independent reporters. Altogether, some codons at the -2 position can
marginally influence NMD activity in a manner that appears to depend on whether NMD proceeds
via an EJC-independent or EJC-enhanced pathway.

At first glance, the discrepancy in results between the EJC-independent and EJC-enhanced
reporters is puzzling. However, it is possible that the highly efficient EJC-enhanced NMD is
insensitive to other modifying factors or that a narrow range of biological effects may not emerge
as statistically significant in a large pool of ~65k sequences. To further explore the basis for this
discrepancy, we switched to an independent, luciferase-based reporter system to test a limited
number of sequence contexts and their relative impact on EJC-independent and EJC-enhanced
NMD activity (Kolakada et al., 2024). In this reporter system, the transfection control is Renilla
luciferase and the NMD readout is provided by Firefly luciferase (**Supp. Fig. 3F-G**).

As the -1 position of the PTC had the most significant impact on EJC-independent NMD
activity in our MPRA, we checked its effect on EJC-independent and EJC-enhanced NMD activity

in the luciferase reporter system. We varied the amino acid in the -1 position of either the PTC or

the NTC, choosing Gly, Val, Ala, and Tyr, which display a range of NMD efficiencies from highest

to lowest (**Supp. Fig. 3F**). As expected, luciferase activity was ~10-fold lower for the EJC-

enhanced reporters compared to the EJC-independent reporters (**Supp. Fig. 3G**). Notably, the

relative trend in -1 amino acid-dependent NMD variability was the same for both the EJC-

independent and EJC-enhanced reporters. The NMD activity varied depending on the -1 amino

acid, with Gly having ~22% or ~29% more efficient NMD compared to Tyr for the EJC-independent

and EJC-enhanced reporters, respectively (**Supp. Fig. 3G**). Compared to the EJC-independent

and EJC-enhanced reporters, the NMD– reporters did not vary with the change in the -1 amino

acid (**Supp. Fig. 3G**). Taken together, these data demonstrate that the -1 amino acid effect can

extend to both forms of NMD.

Key NMD and translation factors do not influence the -1 amino acid-dependent variation in

NMD activity. To determine the mechanistic basis of the -1 amino acid-dependent variation in

NMD, we next asked whether candidate trans factors played a role in this effect. To this end, we

utilized shRNAs to knock down various NMD- and translation-related factors and test the effect

of the knockdown on reporter levels using the EJC-independent luminescent reporters.

We investigated the UPF proteins (UPF1, UPF2, and UPF3B), which are key effectors of NMD

(Kashima et al., 2006; Yamashita et al., 2009); SMG1, which phosphorylates UPF1 to trigger

NMD; and SMG6, which is responsible for cleaving the transcript when NMD is initiated (Arias-

Palomo et al., 2011; Yamashita et al., 2009) (**Fig. 3A-B**). Reporter protein levels increased upon

knockdown of all NMD factors except for UPF3B, likely because UPF3B is known to be redundant

to UPF3A (Wallmeroth et al., 2022). However, the relative difference in expression of the reporters

based on -1 amino acid identity largely remained the same (**Fig. 3B**; **Supp. Fig. 4A**).

Next, using the same approach, we investigated key translation and protein quality control-

related factors, CUL2, ZNF598, ABCE1, EIF3J, and EIF5A (**Fig. 3C-D**). CUL2 forms a key part of

the protein complex that recognizes C-terminal Gly degrons and could contribute to the decrease

observed in Gly-PTC luciferase reporters (Koren et al., 2018; Lin et al., 2018). ZNF598 recognizes

the interface of collided disomes, thus playing a role in ribosome-associated quality control

substrate recognition (Juszkiewicz et al., 2018). It is possible that some sequences would promote

ribosome collision at the end of a transcript, which would then influence NMD. ABCE1 is a

ribosome recycling factor that acts during normal and premature termination (**Fig. 3C**) (Hellen,

2018; Jackson et al., 2012; Zhu et al., 2020). EIF3J facilitates the binding of the release factors

to the ribosome, and thus stimulates peptide release (Egorova et al., 2021). Finally, EIF5A is an
elongation factor that aids the ribosome in translating Gly/Pro sequences. EIF5A accumulates at
the ribosome during translation termination and is hypothesized to bind at the E-site when the
deacylated tRNA leaves (Pelechano & Alepuz, 2017; Schrader et al., 2006; Schuller et al., 2017).
EIF5A binding to the termination complex may be influenced by the transcript sequence within
the ribosome. Similar to the NMD factors, none of these conditions eliminated the differences in
reporter level expression based on -1 amino acid identity (**Fig. 3D**; knockdown validation **Supp.**
Fig. 4B).

Figure 3. Key NMD and translation factors do not influence the -1 amino acid-dependent variation
in NMD activity. (A) Model of NMD factors tested. **(B)** shRNA knockdown of key NMD factors and their
effect on the luminescence-based EJC-independent reporters. **(C)** Models of the mechanisms of translation
termination, ribosome recycling, and protein degradation factors tested. **(D)** shRNA knockdown of key
translation-related factors and their effect on the luminescence-based EJC-independent reporters.

These data show that key NMD, translation, and protein quality control-related trans-factors
do not have a role in the -1 amino acid-dependent variation of NMD. Instead, the mechanistic
basis of this effect may lie within the intrinsic properties of the sequence.

Peptide release rate is the most important feature that classifies EJC-independent NMD substrate activity. To discover sequence-intrinsic factors that influence NMD, we constructed a random forest classifier trained to discriminate between constructs with high versus low NMD activity (**Fig. 4A**). We selected the 3000 sequences with the highest and lowest NMD activity for the EJC-independent and EJC-enhanced reporters. We then added sequence-intrinsic variables that could influence the classification of high and low NMD activity, including the identities of the codons as well as their properties such as usage frequency (GenScript), tRNA binding strength, and optimality (Bazzini et al., 2016; Grosjean & Westhof, 2016); the identities of the amino acids as well as their physical properties such as charge, solubility, dissociation constants, and hydropathy (ThermoFisher Scientific); and the GC and pyrimidine content of each of the sequences. As termination and NMD are intrinsically linked, we also included the rate of peptide release, which is a measure of the time taken for a release factor to trigger peptidyl-tRNA hydrolysis within the P-site (i.e. the -1 position of a stop codon) and the release of the polypeptide itself. These release rates vary based on the identity of the amino acid present within the P-site during termination. Since release rates are not available for eukaryotic systems, we used the prokaryotic peptide release rates (Pierson et al., 2016). Prokaryotes have two release factors RF1 and RF2 that each recognize a subset of the three stop codons, and therefore, we included the rates calculated for both RF1 and RF2 in our classifier.

We trained two separate classifiers via train (80%)/ test (20%) cross-validation for the EJC-independent and EJC-enhanced reporters using the same input variables. Classification performance for EJC-independent NMD activity was much better than for EJC-enhanced NMD activity (**Fig. 4B**), as indicated by the substantially higher area under the receiver operating characteristic curve and F1 scores for the EJC-independent (0.75 and 0.73, respectively) compared to the EJC-enhanced (0.57 and 0.37, respectively) classifiers. We concluded that only the classifier for EJC-independent NMD activity was robust and suitable for further interpretation (refer to **Supplementary Tables 1-2** for the confusion matrices and other model metrics). The poor performance of the EJC-enhanced classifier suggests that hitherto unknown variables are missing in the classifier that may better predict these reporters' NMD activity.

We focused on the features that were most important for the EJC-independent model. Prokaryotic release factor 1 (RF1) and release factor 2 (RF2) peptide release rates emerged as the most important predictor of NMD activity (**Fig 4C**). Although these rates are measured from a prokaryotic context, the similarity in peptide release mechanisms between prokaryotes and eukaryotes suggests that the relative magnitudes of release rates may also apply to eukaryotic

systems (Hellen, 2018). To visualize the effect of peptide release rates on NMD activity, we

compared the subpopulation of sequences with the fastest and slowest dipeptide release rates

versus all NMD targets (**Fig. 4D**). The two populations were well-separated for RF1 peptide

release rates and less robustly for the RF2 peptide release rates (**Fig. 4D**). To examine this

relationship more quantitatively, we compared the beta coefficients of the effect of the peptide

release rates for the -1 amino acid on NMD activity (**Fig 4E**). We observed a strong correlation

for the RF1 peptide release rates ($r = 0.8$; $p = 1.95 \times 10^{-5}$) and a weaker but significant correlation

for the RF2 peptide release rates ($r = 0.55$; $p = 0.01$) (**Fig 4E**). Taken together, these data suggest

that peptide release rate may be a major factor contributing to the -1 amino acid-dependent impact

on NMD activity.

Figure 4. Peptide release rates predict NMD activity (A) Data preparation and variables input into the

Random Forest model. **(B)** ROC curves, AUC values, and F1 scores for the EJC-independent and EJC-

enhanced classifiers. **(C)** Ranked feature importance for the EJC-independent model. **(D)** Density curves
for sequences with slow and fast peptide release rates overlaid on the NMD activity distribution for the EJC-
independent NMD+ sequences. Prokaryotic release rates used (Pierson et al., 2016), left: RF1 release
rates, right: RF2 release rates. For RF1, slow release rates are defined as log (RF1 release rate) £ –0.5,
and fast release rates are defined as log (RF1 release rate) > 1. For RF2, slow release rates are defined
as log (RF2 release rate) £ 0, and fast release rates are defined as log (RF2 release rate) > 1.25. p,
Wilcoxon test p-value **(E)** Scatterplot of the beta coefficients for each -1 amino acid obtained for the EJC-
independent reporters against the RF1 (left) and RF2 (right) peptide release rates. "r" refers to the
Pearson's correlation coefficient and "p" is the corresponding P-value.

Eukaryotic translation termination efficiency varies with the identity of the amino acid
upstream of the stop codon *in vitro*. The peptide release rate values used in the Random
Forest classifier (**Fig. 4**) came from a prokaryotic system and we wanted to determine whether
they reflect the relative peptide release rate in a eukaryotic system (Pierson et al., 2016). To
determine if the rate of peptide release varied based on the identity of the -1 amino acid in a
eukaryotic system, we used the *in vitro* Termi-Luc assay (**Fig. 5A**). In this assay, we provide an
excess of mutated eukaryotic release factor 1 (eRF1(AGQ)), which can recognize stop codons
but unable to induce peptidyl-tRNA hydrolysis (Frolova et al., 1999). eRF1(AGQ) binding thereby
traps translation at the termination stage, allowing us to purify pre-termination ribosomal
complexes (preTCs) from cells. PreTCs are then exposed to purified, functional release factors
(working concentrations determined in **Supp. Fig. 5**), and the real-time release kinetics of the
synthesized NLuc are then measured to determine the translation termination rate of the
transcript. We used this system to compare termination rates when Gly, Val, Ala, or Tyr codons
(representing the full range of observed NMD efficiencies from high to low) were present at the -
1 position to the stop codon.

To check if the substitution of an amino acid residue affected NLuc activity, translation
efficiency, or preTC stability of NLuc, control experiments were first performed. To determine
whether NLuc activity was impacted by altering the last amino acid, we purified all tested peptide
variants as recombinant proteins and determined their maximum luminescence. All NLuc peptide
variants had the same activity (**Fig. 5B**). To determine whether translation efficiency was
impacted by changing the last amino acid, we calculated the maximal derivative (slope) of the
linear portion of the luminescence production curve. The identity of the last amino acid affected
translation efficiency with Tyr having the lowest and Ala having the highest efficiency (**Fig. 5C**).
To determine preTC stability, the rate of peptide release of each peptide variant was calculated
without adding any release factors. These data showed that Gly had the most unstable preTC,
while Tyr had the most stable preTC (**Fig. 5D**). Finally, we performed the Termi-Luc assay on

each peptide variant starting with equal amounts of preTC complexes to account for the variability

in translation efficiency and preTC stability. We found that Gly had the slowest release rate, and

Tyr had the fastest release rate. This difference was statistically significant and correlated with

high and low NMD activity, respectively (**Fig. 5E**). Finally, the release rates observed via the

TermiLuc assay were highly correlated with the values obtained in the prokaryotic system by

Pierson *et al.* (**Fig. 5F**; r_{RF1} = 0.98; r_{RF2} = 0.94) (Pierson et al., 2016). These data show that peptide

release rates indeed vary based on the identity of the last amino acid in a eukaryotic system, with

glycine showing the slowest termination rate and tyrosine the fastest, among the amino acids

tested.

390
Figure 5. Peptide release rate in an *in vitro* eukaryotic system varies with the identity of the amino
acid upstream of the stop codon. (A) Schematic of the Termi-Luc peptide release assay. This assay
was performed for nanoluciferase reporters with different amino acids -1 to the stop codon. **(B)** Tested *in*

vitro luminescence activity of recombinant NLuc variants. **(C)** Translation efficiency of each reporter in the
in vitro system. **(D)** Relative rate of peptide release when no release factors are added to the termination
reaction i.e., preTC instability. **(E)** Relative peptide release rates for each reporter **(F)** A comparison of
Termi-Luc determined eukaryotic release rates with Pierson et al prokaryotic release rates (Pierson et al.,
2016). "r" refers to the Pearson's correlation coefficient and "p" is the corresponding P-value.

Taken together, our data supports a model where a slower peptide release rate offers a longer
kinetic window for NMD factors to assemble and degrade a transcript, thus causing highly efficient
NMD (**Fig. 6**).

403
404

DISCUSSION

NMD is often considered a binary outcome that relies on the presence of triggering factors
downstream of the PTC. While recent results increasingly suggest that NMD is tunable
(Jagannathan & Bradley, 2016; Lindeboom et al., 2019), the factors influencing this variability are
incompletely understood. Our data support a window of opportunity model where the time of
residence of the terminating ribosome, which serves as the platform for the assembly of NMD
machinery, determines the effectiveness of NMD on a given transcript (**Fig. 6**). Consistent with
this model, a slower peptide release rate enhances NMD activity, while a faster rate leads to less
efficient NMD. Together with the RBP cues that trigger NMD, termination kinetics are a key
modifier of the probability with which a prematurely terminating ribosome leads to NMD.

The key finding that led us to the window of opportunity model was the strong enrichment
of glycine in the position immediately before the PTC in prematurely terminating transcripts. While
Gly codons are the most enriched before a PTC, they are the least enriched before an NTC. Even
further, PTCs with Gly-PTC contexts are enriched among variants that are tolerant to loss of
function in healthy individuals, while they are not enriched among disease-causing variants.
These enrichment patterns became logically coherent once we showed that Gly-PTC induces the
most efficient NMD among all possible amino acid contexts and does so through its slow peptide
release rate from the terminating ribosome. Interestingly, Gly is also a C-terminal degron,
meaning that Gly-end proteins, truncated or otherwise, are more efficiently cleared (Koren et al.,
2018; Lin et al., 2018). Thus, the preponderance of glycine codons at the ends of truncated
proteins may be the result of the co-evolution of PTCs alongside processes that efficiently
eliminate Gly-end proteins at both the RNA and protein levels.

Figure 6. A kinetic model for the variability in NMD activity dependent upon the codon preceding the PTC. A window of opportunity model for how peptide release rates influence NMD activity. A slower peptide release rate gives NMD factors a larger window to bind an act on a target, and vice versa.

Other amino acids having patterns similar to Gly are also consistent with our model. For example, Thr and Pro are also enriched before a PTC in healthy individuals and depleted before an NTC. They both support efficient NMD and exhibit slow peptide release rates. On the other end of the spectrum, Tyr had the lowest NMD activity in our experiments and was enriched before NTCs but not PTCs; Tyr also has the fastest peptide release rate. PTCs with Tyr at the -1 of a PTC are depleted among variants tolerant to loss of function and enriched in disease contexts.

Apart from the -1 amino acid position, our massively parallel assessment of NMD activity allowed us to uncover other local sequence preferences for efficient NMD. Surprisingly, we found that the stop codon TGA caused 20% more efficient EJC-independent NMD than TAG. TGA also emerged as an important predictor of EJC-independent NMD in the random forest classifier. This observation is counterintuitive since TGA is known to be the stop codon with the highest probability of readthrough, which usually stabilizes an NMD target (Tate & Mannering, 1996). Our results suggest that there are kinetic differences in termination at different stop codons. Additional factors, such as RBPs or other sequence elements, may influence whether a stop codon is read through or whether it causes efficient NMD.

Our study highlights the power of human genetics to discover novel molecular mechanisms that can explain the real-world variability observed in the expression of genes containing nonsense variants. Our results point to a key role for the kinetics of translation termination in determining NMD activity, ultimately determining the efficiency with which prematurely terminating transcripts are cleared from the cell. Future work will be required to determine whether the influence of translation termination kinetics may also extend to other translation-associated quality

453 control processes. The impact of amino acid identity upstream of the PTC on NMD activity and

454 patterns of amino acid-PTC enrichment in healthy and diseased populations suggest that NMD

455 has played a key role in shaping the evolution of nonsense variants. These findings can be used

456 to better model the phenotypic outcomes of PTCs, and thus develop more effective strategies to

457 counteract PTC-associated diseases.

ACKNOWLEDGEMENTS

We thank Matthew Taliaferro for the gift of HEK293T cells with the loxP cassette. We thank Timothy Stasevich, Tatsuya Morisaki, and Laura White for exploratory studies on glycine's impact on NMD. We thank Monkol Lek and Sander Pajusalu for helpful discussions on gnomAD analysis. We thank Srinivas Ramachandran, Olivia Rissland, and all members of the Jagannathan laboratory for insightful manuscript feedback. This work was supported by the RNA Bioscience Initiative (R. F.), AHA Award #831183 (D.K.), 5T32GM136444-03 (M.L.), Polish National Agency for Academic Exchange Bekker Program PPN/BEK/2019/1/00173 (M.P.S.), R35 GM119550 (J.H.), R35GM147025 (N.M.), University of Colorado School of Medicine Translational Research Scholars Program (S.J.), and the National Institutes of Health grant R35GM133433 (S.J.), The study of the effect of the stop codon 5' context on the efficiency of translation termination was supported by the Russian Science Foundation grant 22-14-00279 (E.A.).

AUTHOR CONTRIBUTIONS

Conceptualization (D.K., R.F., S.J.); Data curation (D.K., R.F., A.E.C., S.J.); Formal analysis (D.K., R.F., S.J.); Funding acquisition (D.K., E.A., S.J.); Investigation (D.K., R.F., N.B., A.S., M.L., A.E.C., S.J.); Project administration (S.J.); Software (D.K., R.F., M.P.S., M.A.C., N.M., S.J.); Supervision (J.H., N.M., E.A., S.J.); Validation (D.K., A.E.C.); Visualization (D.K., R.F., A.E.C., S.J.); Writing – original draft (D.K., S.J.); Writing – review and editing (D.K., R.F., A.E.C., M.A.C, M.P.S., J.H., N.M., E.A., J S.J.).

DECLARATION OF INTERESTS

The authors declare no competing interests.

METHODS

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Sujatha Jagannathan (sujatha.jagannathan@cuanschutz.edu).

Materials availability

All unique reagents generated in this study are available from the lead contact with a completed Material Transfer Agreement.

Data and code availability

- All sequencing data have been deposited at GEO (GSE261151) and will be publicly available as of publication.
- All original code is available at the GitHub repository: https://github.com/jagannathan-lab/2024-kolakada_et_al.
- Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

Cell lines and culture conditions

HEK293T cells (female) were obtained from ATCC (CRL-3216; RRID:CVCL_0063). HEK293T Lox2272/LoxP cells were obtained from Taliaferro laboratory. All cell lines were determined to be free of mycoplasma by PCR screening. HEK293T and HEK293T cells with Lox2272/LoxP cells were maintained in Dulbecco's Modified Eagle Medium (DMEM) (Thermo Fisher Scientific) supplemented with 10% EqualFETAL (Atlas Biologicals).

METHOD DETAILS

gnomAD and ClinVar analysis

Rare (frequency < 0.01) SNP variants from gnomAD v2.1.1 exome database that pass random forest classification quality filtering were analyzed, filtering for "Stop-gained" consequence via bcftools. Using "cDNA_position" field pre-annotated by VEP, the sequence and codons surrounding variants from Gencode v19 transcript FASTA file was then extracted using bedtools getfasta. For other contexts such as normal stop codons, custom code (found in: https://github.com/jagannathan-lab/2024-kolakada_et_al) was used to extract sequence and codon information using Biostrings and BSgenome in R. Codon enrichment is calculated as observed frequency of a particular codon/amino acid normalized against normal occurrence frequency of the codon/amino acid in the last 10 positions of all normal coding genes, similar to Koren et al (Koren et al., 2018). Gene-level loss of function scores were retrieved from gnomAD LoF constraint scores. Clinvar data from 2022/07/30 was processed similarly to gnomAD, but with sequence extraction done in R.

Cloning

The NMD reporter plasmids have been previously described (Kolakada et al., 2024). For individual fluorescent and luminescent reporters containing Gly, Val, Ala, and Tyr amino acids before the PTC, oligos containing these sequences (Integrated DNA Technologies) and matching the EcoRI (NEB, R3101S) and XhoI (NEB, R0146S) restriction sites were synthesized. These oligos were annealed and ligated using the Quick Ligation Kit (NEB, M2200L) into EJC-independent and EJC-enhanced fluorescent and luminescent backbones digested with EcoRI and XhoI. To make the NMD– reporters for each sequence, each EJC-independent reporter was digested with EcoRI and MfeI (NEB, R3589S), the sticky ends were filled in using Klenow Fragment (NEB, M0212S) and the blunt ends were ligated together using the Quick Ligation Kit (NEB), removing the GFP 3′ UTR. The oligos synthesized for each sequence tested are as follows.

Oligo	Sequence (5' – 3')
G1_Top	AATTCAGAACCACCAGAACCACCTTAGCCAGAACCACCAGAACCACCC
G2_Top	AATTCAGAACCACCAGAACCACCTTAACCAGAACCACCAGAACCACCC
A_Top	AATTCAGAACCACCAGAACCACCTTAGGCAGAACCACCAGAACCACCC
V_Top	AATTCAGAACCACCAGAACCACCTTACACAGAACCACCAGAACCACCC
Y_Top	AATTCAGAACCACCAGAACCACCTTAGTAAGAACCACCAGAACCACCC
G1_Bottom	TCGAGGGTGGTTCTGGTGGTTCTGGCTAAGGTGGTTCTGGTGGTTCTG
G2_Bottom	TCGAGGGTGGTTCTGGTGGTTCTGGTTAAGGTGGTTCTGGTGGTTCTG

A_Bottom	TCGAGGGTGGTTCTGGTGGTTCTGCCTAAGGTGGTTCTGGTGGTTCTG
V_Bottom	TCGAGGGTGGTTCTGGTGGTTCTGTGTAAGGTGGTTCTGGTGGTTCTG
Y_Bottom	TCGAGGGTGGTTCTGGTGGTTCTTACTAAGGTGGTTCTGGTGGTTCTG

MPRA plasmid library

An oligo library (Eurofins) of sequence CTA GCA AAC TGG GGC ACA GCC TCG AGG GTG GTT CTG GTG GTN NNN NNT RRN GTG GTT CTG GTG GTT CTG AAT TCG ACT ACA AGG ACC ACG ACG G was synthesized, where N refers to equal proportions of the A, C, G, and T nucleotides, and R refers to equal portions of the A and G nucleotides. This library was resuspended in water to 100 μM. The oligo libraries were then filled in (1 amplification cycle) using the Q5 High-Fidelity DNA Polymerase (NEB, M0492S) according to manufacturer's instructions, using 100 μmols of oligos, and 100 μmols of reverse primer (sequence: CACCGTCGTGGTCCTTGTAGTC), in 2x PCR reactions of 25 μL each. After amplification, the PCR reaction was digested with Exonuclease I, at 37°C for 3 h to digest any remaining single stranded DNA. The DNA was purified using AMPure XP beads (Beckman, A63882), as per manufacturer's instructions.

The EJC-independent and EJC-enhanced fluorescent reporter backbones were linearized using EcoRI (NEB) and XhoI (NEB) at room temperature overnight. Digested plasmid DNA was gel purified using the NucleoSpin Gel and PCR Clean-up kit (Takara Bio, 740609.250). The oligo library was then cloned into the digested backbones using the Gibson Assembly Master Mix (NEB, E2611L) using an insert:vector molar ratio of 7:1 for the EJC-independent backbone and 3:1 for the EJC-enhanced backbone. The reactions were incubated at 50°C for 1 h. The reactions were then purified using AMPure XP beads to remove excess salts and a total of 200 ng of plasmid DNA was transformed into MegaX DH10B T1R Electrocompetent Cells (ThermoFisher, C640003), using a Biorad GenePulser electroporator. The transformed cells were recovered in the recovery medium provided with the electrocompetent cells at 37°C for 1 h, then plated on 33 pre-warmed 15 cm Luria broth (LB) agar-Carbenicillin (RPI, C46000-5.0) plates and incubated at 37°C overnight. The next day colonies were collected by scraping plates, spun-down, and then midi-prepped (NucleoBond Xtra Midi EF kit, 740420.50) to extract the plasmid DNA.

Plasmid transfections

Transfections for fluorescent and luminescent reporters were performed differently since
fluorescent proteins have longer maturation rates than luciferase proteins (Mazo-Vargas et al.,
2014). For the fluorescent reporters, cells were split 24 h after transfection equally into two new
6-wells. Forty-eight hours after transfection, one set of cells was harvested for RNA and the other
set for flow cytometry. For the luminescent reporters, transfections were performed in 24-well
plates at a 40% confluency with 500 ng of plasmid DNA using Lipofectamine 2000 (ThermoFisher
Scientific, 11668027), as per manufacturer's instructions. All cells were harvested for luciferase
assays.

Generating NMD+ reporter cell lines for the MPRA

To create the EJC-independent and EJC-enhanced reporter cell lines HEK293T cells
containing a Lox2272 and LoxP enclosing a blasticidin cassette (courtesy Taliaferro Lab, CU
Anschutz), were co-transfected with 97.5% NMD reporter plasmid library and 2.5% Cre-plasmid
(Addgene plasmid #27493). These transfections were performed in 60-70% confluent 10 cm
plates using 11,800 ng of plasmid library and 295 ng of Cre plasmid per plate for 20 plates.
Twenty-four hours after transfection, each plate of cells was split equally into two new plates and
puromycin (2 μg/mL) was added to the media to begin selection. Puromycin selection lasted
between 1-2 weeks, during which the media was replaced with fresh media supplemented with
puromycin every 1-2 days and cells were consolidated into fewer plates when their confluency
got too low. Once selection was complete, cells were frozen down in medium containing 10%
DMSO. These cells were thawed at a later time for the MPRA.

Targeted sequencing of MPRA libraries

To obtain MPRA sequencing libraries, frozen EJC-independent and EJC-enhanced
reporter cell lines were thawed. These cells were split into 6, 15 cm plates, 3 of which were treated
with DMSO and the other 3 treated with 0.5 μM SMG1i (technical replicates). These plates were
harvested for RNA in TRIzol Reagent (ThermoFisher Scientific, 15596018) after 24 hrs. TRIzol
extractions were performed for RNA as per manufacturer's instructions. A DNase digest was
conducted on 10 μg of RNA from each sample using the Turbo DNA-free kit (ThermoFisher
Scientific, AM1907), including reaction inactivation. Following this, 2 μg of digested RNA was
used to make cDNA using Superscript II Reverse Transcriptase in duplicate 20 μL reactions
(ThermoFisher Scientific, 18064014).

For library preparation, each cDNA sample was split into 20 PCR reactions (2 μL
cDNA/PCR) and amplified using reporter-specific forward and reverse primers with UMIs (forward

primer: ACACTCTTTCCCTACACGACGCTCTTCCGATCTGCAGACTTCCTCTGCCCTC;
reverse primer: AGACGTGTGCTCTTCCGATCTNNNNNNNNNtggggcacagcctcga, where the Ns
refer to the nucleotides for the UMIs). PCR reactions were performed using Kapa HotStart PCR
Kit (Roche, KAPA KK2502) using an annealing temperature of 69°C and 32x cycles. The 20 PCR
reactions were pooled and 100 µL of all reactions were purified using AMPure XP beads. A
second PCR reaction was performed on 25 ng of the purified product with primers containing
Illumina adaptors. Once again, the Kapa HotStart PCR Kit was used, this time with an annealing
temperature of 65°C and 8x cycles. Reactions were purified using AMPure XP beads and samples
were run on an agarose gel to verify the library size. Libraries were sequenced using the Illumina
sequencing platform NovaSeq 6000 (paired-end 2x150 cycles). A total of 50 million and 30 million
reads were requested for the EJC-independent and EJC-enhanced libraries, respectively.

MPRA data analysis

To process the MPRA sequencing libraries, the UMI-tools repository was used to add the
UMIs to the name of each read in the FASTA files (Smith et al., 2017). Following this, Cutadapt
was used to trim the 5′ and 3′ends of the reads in a sequence dependent manner, up to the 10 nt
context of interest (Martin, 2011); ~25% of reads were lost in this process. SeqPrep was used to
merge sequencing reads 1 and 2, only accepting reads that perfectly align
(https://github.com/jstjohn/SeqPrep); ~5% of reads were lost in this process. Bowtie was used to
map the reads to a reference file of all sequence contexts (Langmead & Salzberg, 2012); 1% of
reads were lost in this process. Samtools was used to create a BAM file of mapped reads, followed
by deduplicating the BAM files based on UMIs and converting the BAM files to BED files (Li et al.,
2009). Around 10% of reads were lost upon deduplication of UMIs. A custom python script was
used to remove reads on the negative strand and to get a bed file of reads in the correct
orientation.

The NMD activity of an individual reporter transcript was the log2-fold change in between
DMSO-treated cells to SMG1i-treated cells calculated with DESeq2 (Love et al., 2014). The
reporter transcripts that do not undergo NMD, i.e. Trp codon rather than a stop codon at the PTC
position, were used similar to housekeeping normalization controls within DESeq2. Both linear
models and random forest classifiers were performed in R. Random Forest models were built
utilizing the caret R package (doi:10.18637/jss.v028.i05). First, features with variance close to
zero were eliminated, after which the linear correlation for all numeric features was calculated.
For the features with correlation ≥ 0.85 one feature was randomly selected, and the others
discarded from the model. Data were divided into training and validation set using 0.8:0.2 ratio.

Hyperparameters tuning was performed for train set using 5x train-test cross validation for the
following model parameters: mtry, maxnodes and ntree. Performance of final model was
measured using F1 scores and on validation set. The code and packages used for these analyses
can be found in this repository: https://github.com/jagannathan-lab/2024-kolakada_et_al.

Luciferase assays

For cells harvested from 24-well transfections, the media was aspirated, 400 µL of 1x PLB
from the Dual Luciferase Reporter Assay System (Promega, E1980) was added to the cellular
monolayer per well, and the plate was frozen for subsequent luciferase assays. The frozen cells
were brought to room temperature, transferred to tubes, and lysed thoroughly via vortexing for 30
s per sample. 20µL of this dilution was pipetted into a 96-well plate, in triplicate, per sample. The
LAR II and Stop & Glo reagents were prepared as per the manufacturer's instructions, using 50
µL per sample, each. Luciferase assay was performed using the Glomax Navigator (Promega,
GM2010) as per the Dual Luciferase Assay Protocol.

Lentiviral transductions

Knockdowns were conducted using shRNAs. Plasmids containing shRNAs were first
packaged into lentivirus. To do this, HEK293T cells were transfected with 4 µg of the relevant
shRNA vector (Functional Genomics Core, CU Anschutz), 3 µg of psPAX2 (Addgene plasmid
#12260), and 1 µg of pMD2.G (Addgene plasmid #12259), in 80% confluent 6 cm plates. The
shRNAs used were for scrambled (SCH016), UPF1 (TRCN0000022254), UPF2
(TRCN0000151381), UPF3B (TRCN0000152769), SMG1 (TRCN0000194827), SMG6
(TRCN0000040014), ZNF598 (TRCN0000073159), EIF5A (TRCN0000062552), EIF3J
(TRCN0000062013), CUL2 (TRCN0000006523), and ABCE1 (TRCN0000158417). Twenty-four
hours after transfection the media was replaced with 3 mL of fresh media. Forty-eight hours after
transfection media containing lentivirus was harvested and spun down at 1200 rpm for 5 minutes.
Aliquots of the supernatant were frozen at -80°C for later use.

For the knockdown experiments with the luciferase reporters, cells were seeded at a 40%
confluency in 6-wells and infected with 50 µL of virus. Twenty-four hours post transduction, the
cells were split into 24-wells at a 20% confluency. Forty-eight hours post transduction, the
luciferase reporters were transfected into cells as described above. Seventy-two hours post-
transduction the cells were harvested for luciferase assays. For knockdown validation
experiments, the same procedure was followed without the transfection of luciferase reporters.

RNA extraction and RT-qPCR

For RT-qPCR used to validate knockdowns of SMG1, ZNF598, and EIF3J, RNA was

extracted from cells using TRIzol Reagent (ThermoFisher Scientific). DNase digestion was

performed with DNaseI (ThermoFisher Scientific, 18068-015) on 1 µg of extracted RNA. This was

followed by cDNA synthesis using the SuperScript III First-Strand Synthesis kit, using random

hexamers. A no-RT sample was included as a control to make sure there was no genomic DNA

contamination. The cDNA was then diluted 1:4 and 2 µL were used per 10 µL qPCR reaction.

qPCR was performed using iTaq Universal SYBR Green Supermix and primers specific for

SMG1, ZNF598, EIF3J, and the housekeeping gene RPL27, were used at a final concentration

of 0.25 µM. The following primers were used for each gene:

Oligo	Sequence (5' – 3')
RPL27_F	GCAAGAAGAAGATCGCCAAG
RPL27_R	TCCAAGGGGATATCCACAGA
SMG1_F	TGGGAAAGACCACCACTGCACA
SMG1_R	TGCATGTGTTGACTGGCCTGCT
ZNF598_F	TCGTTGGTGGCGAAGACTAC
ZNF598_R	TCGGTCCTCTTCTTCCCTTT
EIF3J_F	GCAGATAAACTGCGGCTAAAGA
EIF3J_R	TCTCTTGAAGATGGGTTCATAGCA

Reactions were set up in triplicate per sample and plated in 384-well plates. The plates

were then run on the OPUS Bio-Rad qPCR machine (Bio-Rad, 12011319) using the 2-step

Amplification and melting curve protocol. For the knockdown validations, the Livak method (Livak

& Schmittgen, 2001) was used to quantify differences in RNA levels: SMG1, ZNF598, EIF3J, were

normalized to RPL27. Mean RNA levels of 3 different transfections or 3 technical replicates of

transductions were plotted, respectively. Error was calculated using standard error of the mean

(fluorescent/luminescent reporters) or standard deviation (knockdowns).

Protein isolation and western blotting

Cells were harvested in RIPA buffer supplemented with complete protease inhibitor

(Roche, 11836170001). Protein was run on NuPAGE Bis-Tris precast polyacrylamide gels

(Thermo Fisher Scientific, NP0323BOX) alongside PageRuler Plus Prestained Protein Ladder

(Thermo Fisher Scientific, 26619) and transferred to Odyssey nitrocellulose membrane (LI-COR

Biosciences, 926-31092). Membranes were blocked in Intercept (PBS) Blocking Buffer (LI-COR

Biosciences, 927-70001) before overnight incubation at 4°C with primary antibodies diluted in

Blocking Buffer containing 0.2% Tween 20. Membranes were incubated with IRDye-conjugated

secondary antibodies (LI-COR Biosciences) for 1 h and fluorescent signal visualized using a Sapphire Biomolecular Imager (Azure Biosystems) and Sapphire Capture software (Azure Biosystems). When appropriate, membranes were stripped with Restore Western Blot Stripping Buffer (ThermoFisher Scientific) before being re-probed. Primary antibodies include anti-UPF1 (Abcam, ab109363), anti-UPF2 (Cell Signaling, 11875S), anti-UPF3B (Abcam, ab134566), anti-SMG6 (ABclonal, A10141), anti-CUL2 (Bethyl Laboratories, A302-476A), anti-ABCE1 (Abcam, ab185548), anti-EIF5A (Abcam, ab32407), and anti-GAPDH (Abcam, ab9484). Secondary antibodies used include IRDye 650 Goat anti-Mouse IgG Secondary Antibody (LI-COR Biosciences, 926-65010) and IRDye 800CW Goat anti-Rabbit IgG Secondary Antibody (LI-COR Biosciences, 926-32211).

Luminescent reporters for *in vitro* assays

A pNL-globine vector derived from pNL1.1 vector (Promega), with a β-globin 5′UTR addition before nanoluciferase (NLuc) coding sequence was used (Shuvalov et al., 2021). Additional constructs were created based on pNL-globin, encoding NLuc with various substitutions of the codon before the stop codon using the QuikChange Site-Directed Mutagenesis Kit (Agilent Technologies, cat. 200518-5) was used. Primers were selected in the manufacturer's recommended web-based QuikChange Primer Design Program (www.agilent.com/genomics/qcpd). For obtaining mRNA, the fragment of the plasmid were amplified using RV3L (CTAGCAAAATAGGCTGTCCCCAG) and FLA50 (TTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTAACTTGTTTATTGCAGC TTATAATGG) primers, as described in Shuvaolv et al., 2021 (Shuvalov et al., 2021). Templates were run-off transcribed with the T7 RiboMAX ™ Large Scale RNA Production System (Promega, P1320) kit according to the manufacturer's protocol. The mRNA was then purified sequentially by isolation in acidic phenol, precipitation with 3 M LiCl followed by 80% ethanol wash.

Expression and purification of eRF1 and eRF3

Recombinant eRF1 was expressed from the plasmid pET-SUMO-eRF1. To obtain pET-SUMO-eRF1, the eRF1 coding sequence was amplified from the plasmid pET23b-eRF1 (Frolova et al., 2002) using primers petSUMO_eRF1_F (GAGAACAGATTGGTGGTATGGCGGACGACCCCAG) and petSUMO_eRF1_R (CCGAATAAATACCTAAGCTCTAGTAGTCATCAAGGTCAAAAAATTCATCGTCTCCTCC). eRF1 was expressed in E. coli BL21(DE3) cells. The lysate was prepared by ultrasonification of the pelleted cells in a buffer composed of 20 mM Tris-HCl pH 7.5, 500 mM KCl, 10% glycerol, 0.1% Triton X-100, 0.5 mM PMSF, and 1 mM DTT. Following this, His-SUMO tag was cleaved by

His-tagged Ulp1 protease. Untagged eRF1 was further purified by anion-exchange chromatography (HiTrap Q HP, Cytiva). Fractions enriched with eRF1 were collected, dialyzed in storage buffer, frizzed in liquid nitrogen, and stored at −70 ºC. Recombinant eRF3A cloned into baculovirus vector EMBacY from a MultiBac expression system was expressed in the insect cell line Sf21. Following this, recombinant proteins were purified using Ni-NTA agarose and ion-exchange chromatography, as described previously (Ivanov et al., 2016).

Expression, purification, and determination of luminescence or recombinant nanoluciferase

To obtain recombinant NLuc with varying amino acids in the position -1 to the stop codon, corresponding constructs of pNL-globine were created using the petSUMO vector (Invitrogen, cat. K30001). Each vector was assembled via Gibson Assembly (NEB, cat. E2611L) using PCR products of the petSUMO backbone (forward primer: AGCTTAGGTATTTATTCGGCGCAAAGTG; reverse primer: ACCACCAATCTGTTCTCTGTGAGC) and pNL-globine inserts (forward primer: GAGAACAGATTGGTGGTATGGTCTTCACACTCGAAGATTTCGTTGG; reverse primer: CCGAATAAATACCTAAGCTTACGCCAGAATGCGTTCGCA) amplified using Q5 High-Fidelity DNA Polymerase (NEB, cat. M0491L). The resulting plasmids were expressed in *E. coli* BL21(DE3) and His-SUMO-Nluc was purified using Ni-affinity chromatography. Recombinant 6xHis-ULP1 protease was added to purify 6xHisSUMO-Nluc and incubated for 2 hours. The 6xHisSUMO fragment and 6xHis-ULP1 were then removed with Ni-NTA agarose to obtain purified Nluc without tags. 0.1 femtomole of recombinant Nluc was incubated in storage buffer, with the addition of BSA up to 500 μg/mL and 0.5 % NanoGlo (Promega). Luminescence was measured at 30°C using a Tecan Infinite 200 Pro (Tecan, Männedorf, Switzerland) in a 40 min period. The luminescence of NLuc was calculated as a maximum of relative luminescence units (RLU_{max}).

Purification of the preTC-NLuc and Termi-Luc Assay

For the Termi-Luc assay, preTCs translating NLuc (preTC-NLuc) were purified using previously published methods (Shuvalov et al., 2021; Susorov et al., 2020). 100% RRL lysate was preincubated in a mixture containing 1 mM $CaCl_2$ and 3 U/μL Micrococcal nuclease (Fermentas) at 30°C for 10 min, followed by the addition of EGTA to a final concentration of 4 mM. The lysate was then diluted to 70% (v/v) and supplemented with 20 mM HEPES-KOH (pH 7.5), 80 mM KOAc, 0.5 mM $Mg(OAc)_2$, 0.3 mM ATP, 0.2 GTP, 0.04 mM of each of 20 amino acids (Promega), 0.5 mM spermidine, 0.45 μM aminoacylated total rabbit tRNA, 10 mM creatine

phosphate, 0.003 U/μL creatine kinase (Sigma), 2 mM DTT, and 0.2 U/μL Ribolock
(ThermoFisher) (70% RRL mix).

For preTC-NLuc assembly, 220 μL of 70% RRL mix was preincubated in the presence of
1.7 μM ERF1 G183A mutant (Alkalaeva et al., 2006) at 30° C for 10 min, followed by the addition
of 10.5 pmol of NLuc mRNA. The mixture was incubated at 30°C for 40 min. The KOAc
concentration was then adjusted to 300 mM and the mixture was layered on 5 ml of a 10–35%
linear sucrose gradient in a buffer containing 50 mM HEPES-KOH, pH 7.5, 7.5 mM Mg(OAc)$_2$,
300 mM KOAc, 2 mM DTT. The gradient was centrifuged in a SW55-Ti (Beckman Coulter) rotor
at 55 000 rpm (367 598 g$_{max}$) for 1 h. The gradient was fractionated in 15 fractions of 150 μL from
bottom to top and the remaining sucrose was collected separately. Fractions of the first peak,
enriched with the preTC-Nluc were analyzed by the Termi-Luc assay, merged, flash-frozen in
liquid nitrogen, and stored at −70°C.

A peptide release assay with the preTC-NLuc (Termi-Luc) was performed as previously
described with some modifications (Susorov et al., 2020). Peptide release was conducted in a
solution containing 1.5 pM preTC-Nluc, 45 mM HEPES-KOH pH 7.5, 1.4 mM Mg$_2$OAc, 56 mM
KOAc pH 7.0, 1 mM DTT, 177 μM spermidine, 1.5 % (w/w) sucrose, 0.8 mM MgCl$_2$, 0.2 mM GTP
supplemented with equimolar MgCl$_2$, and 0.5 % NanoGlo (Promega), in the absence or the
presence of release factors eRF1 and eRF3a at various concentrations. Luminescence was
measured at 30°C using a Tecan Infinite 200 Pro (Tecan, Männedorf, Switzerland) for 12 min.
The translation efficiency was calculated as the maximal derivative of the growing linear section
of the luminescence curve (v$_0$, RLU/min). For comparison of different preTCs, a single working
concentration of 3 nM of eRF1-eRF3A was chosen in the linear section. The instability of the
preTC-NLuc was evaluated in Termi-Luc assay in the absence of release factors.

The amount of preTC-Nluc was calculated as a maximum of relative luminescence
(RLU$_{max}$). To determine concentration of preTC-Nluc, 1.25 μL of preTC sample were incubated in
the presence of excess of eRF1 and eRF3a (100 nM). Luminescence was measured at 30°C
using a Tecan Infinite 200 Pro (Tecan, Männedorf, Switzerland) for 40 min. PreTC concentration
was considered in the translation termination rate experiments.

Translation of NLuc in RRL lysates

For translation of NLuc, 19 μL of 70% RRL lysate prepared as described in Termi-Luc
section, was mixed with 0.19 μL of NanoGlo and 0.25 pmol of NLuc mRNA. Luminescence was
measured at 30°C using a Tecan Infinite 200 Pro (Tecan, Männedorf, Switzerland) for 60 min.

The translation efficiency was calculated as a maximal derivative of the growing linear section of

the luminescence curve (v_0, RLU/min).

QUANTIFICATION AND STATISTICAL ANALYSIS

Data analysis, statistical tests, and visualization

All experiments were performed with a sample size of at least n=3. For transient

transfections, each replicate represents an independent transfection. For stably integrated cells,

the same cell line was plated in 3 different wells, from which RNA and protein were harvested.

Statistical significance between various populations was calculated using a student's t-test in R

and p-values were two-sided (*$p<0.05$, **$p<0.005$, ***$p<0.0005$). Statistical details of specific

experiments can be found in the Results, Methods, and/or Figure Legends. Plots were generated

using R plotting functions and/or the ggplot2 package. The exact code and packages used for

these analyses can be found in this repository: https://github.com/jagannathan-lab/2024-

kolakada_et_al.

SUPPLEMENTAL INFORMATION

SUPPLEMENTARY FIGURES

Figure S1

A

B

805
806
807
808 **Figure S1. Amino acids enriched from the -10 to the -1 position before a termination codon.**
809 (**A**) Left: Schematic to indicate PTCs analyzed in the gnomAD database, right: amino acid
810 enrichment before the PTC. (**B**) Left: Schematic to indicate NTCs analyzed in the reference
811 genome, right: amino acid enrichment before the NTC. Gly is colored in red.
812

813 **Figure S2**

815 **Figure S2.** MPRA library representation. (**A**) Plasmid library representation for the EJC-
816 independent NMD**+** reporter library. (**B**) The same as (A) but for the EJC-enhanced NMD**+**
817 reporter library.

Figure S3

Figure S3. The codon identity at the -1 position of the PTC alters NMD activity of EJC-enhanced luminescence-based reporters. (**A**) Effect of stop codon identity on EJC-

independent and EJC-enhanced NMD without TGG. (**B-E**) Plots depicting a variable's effect on
EJC-enhanced NMD: (**B**) heat map for the -2 codon, (**C**) dot plot for the -2 amino acid, (**D**) heat
map for them -1 codon, and (**E**) dot plot for the -1 amino acid. (**F**) Schematic of luminescence
based NMD reporters used (right) as well as the amino acids tested in the -1 position to the
PTC/NTC (left). (**G**) Luciferase activity for each of the reporters in EJC-independent, EJC-
enhanced, and NMD– contexts.

Figure S4

Figure S4. Knockdown validations for all shRNAs used. (**A**) Top: western blot validating
UPF1, UPF2, and UPF3B knockdowns; middle: western blot validating SMG6 knockdown;
bottom: qPCR validation of SMG1 knockdown. (**B**) Top left: western blot for CUL2 knockdown
validation; top right: western blot for ABCE1 knockdown validation; middle left: western blot for
EIF5A knockdown validation; bottom left: qPCR validation of ZNF598 knockdown; bottom right:
qPCR validation for EIF3J knockdown. GAPDH was the loading control for all western blots.
RPL27 was the housekeeping gene for all qPCRs. qPCRs were performed in triplicate.

Figure S5.

Figure S5. Determining the optimum concentration of release factors to use for the Termi-
Luc assay.

SUPPLEMENTARY TABLES

Table 1: Confusion matrices for the EJC-independent and EJC-enhanced classifier

Prediction	Reference	n
EJC-Independent		
NMD Escape	NMD Escape	504
Strong NMD	NMD Escape	96
NMD Escape	Strong NMD	271
Strong NMD	Strong NMD	329
EJC-Enhanced		
NMD Escape	NMD Escape	166
Strong NMD	NMD Escape	434
NMD Escape	Strong NMD	131
Strong NMD	Strong NMD	469

866 **Table 2:** Model metrics for the EJC-independent and EJC-enhanced classifier

EJC-Independent	
Sensitivity	0.84
Specificity	0.55

Pos Pred Value	0.65
Neg Pred Value	0.77
Precision	0.65
Recall	0.84
F1	0.73
Prevalence	0.50
Detection Rate	0.42
Detection Prevalence	0.65
Balanced Accuracy	0.69
EJC-Enhanced	
Sensitivity	0.28
Specificity	0.78
Pos Pred Value	0.56
Neg Pred Value	0.52
Precision	0.56
Recall	0.28
F1	0.37
Prevalence	0.50
Detection Rate	0.14
Detection Prevalence	0.25
Balanced Accuracy	0.53

867 **SUPPLEMENTARY DATA**

869 **Supplementary Data Table 1.** DEseq2 results for the EJC-independent library of sequences
870 representing the fold change of DMSO over SMG1i treatment of cells.
871
872 **Supplementary Data Table 2.** DEseq2 results for the EJC-enhanced library of sequences
873 representing the fold change of DMSO over SMG1i treatment of cells.

REFERENCES

Alkalaeva, E. Z., Pisarev, A. V., Frolova, L. Y., Kisselev, L. L., & Pestova, T. V. (2006, Jun 16). In vitro reconstitution of eukaryotic translation reveals cooperativity between release factors eRF1 and eRF3. *Cell, 125*(6), 1125-1136. https://doi.org/10.1016/j.cell.2006.04.035

Arias-Palomo, E., Yamashita, A., Fernandez, I. S., Nunez-Ramirez, R., Bamba, Y., Izumi, N., Ohno, S., & Llorca, O. (2011, Jan 15). The nonsense-mediated mRNA decay SMG-1 kinase is regulated by large-scale conformational changes controlled by SMG-8. *Genes Dev, 25*(2), 153-164. https://doi.org/10.1101/gad.606911

Baker, S. L., & Hogg, J. R. (2017). A system for coordinated analysis of translational readthrough and nonsense-mediated mRNA decay | PLOS ONE. *PLoS ONE, 12(3): e0173980*. https://doi.org/10.1371/journal.pone.0173980

Bateman, J. F., Freddi, S., Nattrass, G., & Savarirayan, R. (2003, Feb 1). Tissue-specific RNA surveillance? Nonsense-mediated mRNA decay causes collagen X haploinsufficiency in Schmid metaphyseal chondrodysplasia cartilage. *Hum Mol Genet, 12*(3), 217-225. https://www.ncbi.nlm.nih.gov/pubmed/12554676

Bazzini, A. A., Del Viso, F., Moreno-Mateos, M. A., Johnstone, T. G., Vejnar, C. E., Qin, Y., Yao, J., Khokha, M. K., & Giraldez, A. J. (2016, Oct 4). Codon identity regulates mRNA stability and translation efficiency during the maternal-to-zygotic transition. *EMBO J, 35*(19), 2087-2103. https://doi.org/10.15252/embj.201694699

Bonetti, B., Fu, L., Moon, J., & Bedwell, D. M. (1995, Aug 18). The efficiency of translation termination is determined by a synergistic interplay between upstream and downstream sequences in Saccharomyces cerevisiae. *J Mol Biol, 251*(3), 334-345. https://doi.org/10.1006/jmbi.1995.0438

905 Buhler, M., Steiner, S., Mohn, F., Paillusson, A., & Muhlemann, O. (2006, May). EJC-
906   independent degradation of nonsense immunoglobulin-mu mRNA depends on 3' UTR
907   length. *Nat Struct Mol Biol, 13*(5), 462-464. https://doi.org/10.1038/nsmb1081
908

909 Chang, J. C., & Kan, Y. W. (1979, Jun). beta 0 thalassemia, a nonsense mutation in man. *Proc*
910   *Natl Acad Sci U S A, 76*(6), 2886-2889. https://www.ncbi.nlm.nih.gov/pubmed/88735
911 https://www.ncbi.nlm.nih.gov/pmc/articles/PMC383714/pdf/pnas00006-0406.pdf
912

913 Chu, V., Feng, Q., Lim, Y., & Shao, S. (2021, Dec 1). Selective destabilization of polypeptides
914   synthesized from NMD-targeted transcripts. *Mol Biol Cell, 32*(22), ar38.
915   https://doi.org/10.1091/mbc.E21-08-0382
916

917 Cridge, A. G., Crowe-McAuliffe, C., Mathew, S. F., & Tate, W. P. (2018, Feb 28). Eukaryotic
918   translational termination efficiency is influenced by the 3' nucleotides within the
919   ribosomal mRNA channel. *Nucleic Acids Res, 46*(4), 1927-1944.
920   https://doi.org/10.1093/nar/gkx1315
921

922 Egorova, T., Biziaev, N., Shuvalov, A., Sokolova, E., Mukba, S., Evmenov, K., Zotova, M.,
923   Kushchenko, A., Shuvalova, E., & Alkalaeva, E. (2021, Nov 8). eIF3j facilitates loading of
924   release factors into the ribosome. *Nucleic Acids Res, 49*(19), 11181-11196.
925   https://doi.org/10.1093/nar/gkab854
926

927 Frolova, L., Seit-Nebi, A., & Kisselev, L. (2002, Feb). Highly conserved NIKS tetrapeptide is
928   functionally essential in eukaryotic translation termination factor eRF1. *RNA, 8*(2), 129-
929   136. https://doi.org/10.1017/s1355838202013262
930

931   Frolova, L. Y., Tsivkovskii, R. Y., Sivolobova, G. F., Oparina, N. Y., Serpinsky, O. I., Blinov, V.
932   M., Tatkov, S. I., & Kisselev, L. L. (1999, Aug). Mutations in the highly conserved GGQ
933   motif of class 1 polypeptide release factors abolish ability of human eRF1 to trigger
934   peptidyl-tRNA hydrolysis. *RNA, 5*(8), 1014-1020.
935   https://doi.org/10.1017/s135583829999043x
936

937   Grosjean, H., & Westhof, E. (2016, Sep 30). An integrated, structure- and energy-based view of
938   the genetic code. *Nucleic Acids Res, 44*(17), 8020-8040.
939   https://doi.org/10.1093/nar/gkw608
940

941   Hellen, C. U. T. (2018, Oct 1). Translation Termination and Ribosome Recycling in Eukaryotes.
942   *Cold Spring Harb Perspect Biol, 10*(10). https://doi.org/10.1101/cshperspect.a032656
943

944   Hogg, J. R., & Goff, S. P. (2010, Oct 29). Upf1 senses 3'UTR length to potentiate mRNA decay.
945   *Cell, 143*(3), 379-389. https://doi.org/10.1016/j.cell.2010.10.005
946

947   Inacio, A., Silva, A. L., Pinto, J., Ji, X., Morgado, A., Almeida, F., Faustino, P., Lavinha, J.,
948   Liebhaber, S. A., & Romao, L. (2004, Jul 30). Nonsense mutations in close proximity to
949   the initiation codon fail to trigger full nonsense-mediated mRNA decay. *J Biol Chem,*
950   *279*(31), 32170-32180. https://doi.org/10.1074/jbc.M405024200
951

952   Ivanov, A., Mikhailova, T., Eliseev, B., Yeramala, L., Sokolova, E., Susorov, D., Shuvalov, A.,
953   Schaffitzel, C., & Alkalaeva, E. (2016, Sep 19). PABP enhances release factor
954   recruitment and stop codon recognition during translation termination. *Nucleic Acids*
955   *Res, 44*(16), 7766-7776. https://doi.org/10.1093/nar/gkw635
956

Jackson, R. J., Hellen, C. U., & Pestova, T. V. (2012). Termination and post-termination events

in eukaryotic translation. *Adv Protein Chem Struct Biol, 86*, 45-93.

https://doi.org/10.1016/B978-0-12-386497-0.00002-5

Jagannathan, S., & Bradley, R. K. (2016, Dec). Translational plasticity facilitates the

accumulation of nonsense genetic variants in the human population. *Genome Res,*

26(12), 1639-1650. https://doi.org/10.1101/gr.205070.116

Juszkiewicz, S., Chandrasekaran, V., Lin, Z., Kraatz, S., Ramakrishnan, V., & Hegde, R. S.

(2018, Nov 1). ZNF598 Is a Quality Control Sensor of Collided Ribosomes. *Mol Cell,*

72(3), 469-481 e467. https://doi.org/10.1016/j.molcel.2018.08.037

Karczewski, K. J., Francioli, L. C., Tiao, G., Cummings, B. B., Alfoldi, J., Wang, Q., Collins, R.

L., Laricchia, K. M., Ganna, A., Birnbaum, D. P., Gauthier, L. D., Brand, H., Solomonson,

M., Watts, N. A., Rhodes, D., Singer-Berk, M., England, E. M., Seaby, E. G., Kosmicki,

J. A., Walters, R. K., Tashman, K., Farjoun, Y., Banks, E., Poterba, T., Wang, A., Seed,

C., Whiffin, N., Chong, J. X., Samocha, K. E., Pierce-Hoffman, E., Zappala, Z.,

O'Donnell-Luria, A. H., Minikel, E. V., Weisburd, B., Lek, M., Ware, J. S., Vittal, C.,

Armean, I. M., Bergelson, L., Cibulskis, K., Connolly, K. M., Covarrubias, M., Donnelly,

S., Ferriera, S., Gabriel, S., Gentry, J., Gupta, N., Jeandet, T., Kaplan, D., Llanwarne,

C., Munshi, R., Novod, S., Petrillo, N., Roazen, D., Ruano-Rubio, V., Saltzman, A.,

Schleicher, M., Soto, J., Tibbetts, K., Tolonen, C., Wade, G., Talkowski, M. E., Genome

Aggregation Database, C., Neale, B. M., Daly, M. J., & MacArthur, D. G. (2020, May).

The mutational constraint spectrum quantified from variation in 141,456 humans. *Nature,*

581(7809), 434-443. https://doi.org/10.1038/s41586-020-2308-7

Kashima, I., Yamashita, A., Izumi, N., Kataoka, N., Morishita, R., Hoshino, S., Ohno, M.,

Dreyfuss, G., & Ohno, S. (2006, Feb 1). Binding of a novel SMG-1-Upf1-eRF1-eRF3

complex (SURF) to the exon junction complex triggers Upf1 phosphorylation and

nonsense-mediated mRNA decay. *Genes Dev, 20*(3), 355-367.

https://doi.org/10.1101/gad.1389006

Kazazian, H. H., Jr., Dowling, C. E., Hurwitz, R. L., Coleman, M., Stopeck, A., & Adams, J. G.,

3rd. (1992, Jun 1). Dominant thalassemia-like phenotypes associated with mutations in

exon 3 of the beta-globin gene. *Blood, 79*(11), 3014-3018.

https://www.ncbi.nlm.nih.gov/pubmed/1586746

http://www.bloodjournal.org/content/bloodjournal/79/11/3014.full.pdf

Keeling, K. M., Lanier, J., Du, M., Salas-Marco, J., Gao, L., Kaenjak-Angeletti, A., & Bedwell, D.

M. (2004, Apr). Leaky termination at premature stop codons antagonizes nonsense-

mediated mRNA decay in S. cerevisiae. *RNA, 10*(4), 691-703.

https://www.ncbi.nlm.nih.gov/pubmed/15037778

Kerr, T. P., Sewry, C. A., Robb, S. A., & Roberts, R. G. (2001, Oct). Long mutant dystrophins

and variable phenotypes: evasion of nonsense-mediated decay? *Hum Genet, 109*(4),

402-407. https://doi.org/10.1007/s004390100598

1003

1004   Khandelia, P., Yap, K., & Makeyev, E. V. (2011, Aug 2). Streamlined platform for short hairpin

1005   RNA interference and transgenesis in cultured mammalian cells. *Proc Natl Acad Sci U S*

1006   *A, 108*(31), 12799-12804. https://doi.org/10.1073/pnas.1103532108

1007

1008   Kolakada, D., Campbell, A. E., Galvis, L. B., Li, Z., Lore, M., & Jagannathan, S. (2024, Feb 20).

1009   A system of reporters for comparative investigation of EJC-independent and EJC-

1010   enhanced nonsense-mediated mRNA decay. *Nucleic Acids Res*.

1011   https://doi.org/10.1093/nar/gkae121

1012

1013 Koren, I., Timms, R. T., Kula, T., Xu, Q., Li, M. Z., & Elledge, S. J. (2018, Jun 14). The
1014   Eukaryotic Proteome Is Shaped by E3 Ubiquitin Ligases Targeting C-Terminal Degrons.
1015   *Cell, 173*(7), 1622-1635 e1614. https://doi.org/10.1016/j.cell.2018.04.028
1016

Landrum, M. J., Lee, J. M., Benson, M., Brown, G. R., Chao, C., Chitipiralla, S., Gu, B., Hart, J.,
Hoffman, D., Jang, W., Karapetyan, K., Katz, K., Liu, C., Maddipatla, Z., Malheiro, A.,
McDaniel, K., Ovetsky, M., Riley, G., Zhou, G., Holmes, J. B., Kattman, B. L., & Maglott,
D. R. (2018, Jan 4). ClinVar: improving access to variant interpretations and supporting
evidence. *Nucleic Acids Res, 46*(D1), D1062-D1067.
https://doi.org/10.1093/nar/gkx1153
1023

1024 Langmead, B., & Salzberg, S. L. (2012, Mar 4). Fast gapped-read alignment with Bowtie 2. *Nat*
1025   *Methods, 9*(4), 357-359. https://doi.org/10.1038/nmeth.1923
1026

1027 Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,
1028   Durbin, R., & Genome Project Data Processing, S. (2009, Aug 15). The Sequence
1029   Alignment/Map format and SAMtools. *Bioinformatics, 25*(16), 2078-2079.
1030   https://doi.org/10.1093/bioinformatics/btp352
1031

1032 Lin, H. C., Yeh, C. W., Chen, Y. F., Lee, T. T., Hsieh, P. Y., Rusnac, D. V., Lin, S. Y., Elledge,
1033   S. J., Zheng, N., & Yen, H. S. (2018, May 17). C-Terminal End-Directed Protein
1034   Elimination by CRL2 Ubiquitin Ligases. *Mol Cell, 70*(4), 602-613 e603.
1035   https://doi.org/10.1016/j.molcel.2018.04.006
1036

1037 Linde, L., Boelz, S., Neu-Yilik, G., Kulozik, A. E., & Kerem, B. (2007, Nov). The efficiency of
1038   nonsense-mediated mRNA decay is an inherent character and varies among different
1039   cells. *Eur J Hum Genet, 15*(11), 1156-1162. https://doi.org/10.1038/sj.ejhg.5201889
1040

Lindeboom, R. G., Supek, F., & Lehner, B. (2016, Oct). The rules and impact of nonsense-mediated mRNA decay in human cancers. *Nat Genet, 48*(10), 1112-1118. https://doi.org/10.1038/ng.3664

Lindeboom, R. G. H., Vermeulen, M., Lehner, B., & Supek, F. (2019, Nov). The impact of nonsense-mediated mRNA decay on genetic disease, gene editing and cancer immunotherapy. *Nat Genet, 51*(11), 1645-1651. https://doi.org/10.1038/s41588-019-0517-5

Livak, K. J., & Schmittgen, T. D. (2001, Dec). Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods, 25*(4), 402-408. https://doi.org/10.1006/meth.2001.1262

Loughran, G., Chou, M. Y., Ivanov, I. P., Jungreis, I., Kellis, M., Kiran, A. M., Baranov, P. V., & Atkins, J. F. (2014, Aug). Evidence of efficient stop codon readthrough in four mammalian genes. *Nucleic Acids Res, 42*(14), 8928-8938. https://doi.org/10.1093/nar/gku608

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol, 15*(12), 550. https://doi.org/10.1186/s13059-014-0550-8

MacArthur, D. G., Balasubramanian, S., Frankish, A., Huang, N., Morris, J., Walter, K., Jostins, L., Habegger, L., Pickrell, J. K., Montgomery, S. B., Albers, C. A., Zhang, Z. D., Conrad, D. F., Lunter, G., Zheng, H., Ayub, Q., DePristo, M. A., Banks, E., Hu, M., Handsaker, R. E., Rosenfeld, J. A., Fromer, M., Jin, M., Mu, X. J., Khurana, E., Ye, K., Kay, M., Saunders, G. I., Suner, M. M., Hunt, T., Barnes, I. H., Amid, C., Carvalho-Silva, D. R., Bignell, A. H., Snow, C., Yngvadottir, B., Bumpstead, S., Cooper, D. N., Xue, Y., Romero, I. G., Genomes Project, C., Wang, J., Li, Y., Gibbs, R. A., McCarroll, S. A.,

1070   Dermitzakis, E. T., Pritchard, J. K., Barrett, J. C., Harrow, J., Hurles, M. E., Gerstein, M.
1071   B., & Tyler-Smith, C. (2012, Feb 17). A systematic survey of loss-of-function variants in
1072   human protein-coding genes. *Science, 335*(6070), 823-828.
1073   https://doi.org/10.1126/science.1215040
1074

1075   Mangkalaphiban, K., He, F., Ganesan, R., Wu, C., Baker, R., & Jacobson, A. (2021, Apr).
1076   Transcriptome-wide investigation of stop codon readthrough in Saccharomyces
1077   cerevisiae. *PLoS Genet, 17*(4), e1009538. https://doi.org/10.1371/journal.pgen.1009538
1078

1079   Martin, M. (2011, 2011-05-02). Cutadapt removes adapter sequences from high-throughput
1080   sequencing reads [next generation sequencing; small RNA; microRNA; adapter
1081   removal]. *2011, 17*(1), 3. https://doi.org/10.14806/ej.17.1.200
1082

1083   Mazo-Vargas, A., Park, H., Aydin, M., & Buchler, N. E. (2014, Nov 5). Measuring fast gene
1084   dynamics in single cells with time-lapse luminescence microscopy. *Mol Biol Cell, 25*(22),
1085   3699-3708. https://doi.org/10.1091/mbc.E14-07-1187
1086

1087   McCaughan, K. K., Brown, C. M., Dalphin, M. E., Berry, M. J., & Tate, W. P. (1995, Jun 6).
1088   Translational termination efficiency in mammals is influenced by the base following the
1089   stop codon. *Proc Natl Acad Sci U S A, 92*(12), 5431-5435.
1090   https://doi.org/10.1073/pnas.92.12.5431
1091

1092   Miller, J. N., & Pearce, D. A. (2014, Oct-Dec). Nonsense-mediated decay in genetic disease:
1093   friend or foe? *Mutat Res Rev Mutat Res, 762*, 52-64.
1094   https://doi.org/10.1016/j.mrrev.2014.05.001
1095

1096   Neu-Yilik, G., Amthor, B., Gehring, N. H., Bahri, S., Paidassi, H., Hentze, M. W., & Kulozik, A. E.
1097   (2011, May). Mechanism of escape from nonsense-mediated mRNA decay of human

1098   beta-globin transcripts with nonsense mutations in the first exon. *RNA, 17*(5), 843-854.

1099   https://doi.org/10.1261/rna.2401811

1100

1101 Nguyen, L. S., Wilkinson, M. F., & Gecz, J. (2014, Oct). Nonsense-mediated mRNA decay:

1102   inter-individual variability and human disease. *Neurosci Biobehav Rev, 46 Pt 2*, 175-186.

1103   https://doi.org/10.1016/j.neubiorev.2013.10.016

1104

1105 Pelechano, V., & Alepuz, P. (2017, Jul 7). eIF5A facilitates translation termination globally and

1106   promotes the elongation of many non polyproline-specific tripeptide sequences. *Nucleic*

1107   *Acids Res, 45*(12), 7326-7338. https://doi.org/10.1093/nar/gkx479

1108

1109 Pereira, F. J., Teixeira, A., Kong, J., Barbosa, C., Silva, A. L., Marques-Ramos, A., Liebhaber,

1110   S. A., & Romao, L. (2015, Jul 27). Resistance of mRNAs with AUG-proximal nonsense

1111   mutations to nonsense-mediated decay reflects variables of mRNA structure and

1112   translational activity. *Nucleic Acids Res, 43*(13), 6528-6544.

1113   https://doi.org/10.1093/nar/gkv588

1114

1115 Pierson, W. E., Hoffer, E. D., Keedy, H. E., Simms, C. L., Dunham, C. M., & Zaher, H. S. (2016,

1116   Sep 27). Uniformity of Peptide Release Is Maintained by Methylation of Release Factors.

1117   *Cell Rep, 17*(1), 11-18. https://doi.org/10.1016/j.celrep.2016.08.085

1118

1119 Popp, M. W., & Maquat, L. E. (2016, Jun 2). Leveraging Rules of Nonsense-Mediated mRNA

1120   Decay for Genome Engineering and Personalized Medicine. *Cell, 165*(6), 1319-1322.

1121   https://doi.org/10.1016/j.cell.2016.05.053

1122

1123 Schrader, R., Young, C., Kozian, D., Hoffmann, R., & Lottspeich, F. (2006, Nov 17).

1124   Temperature-sensitive eIF5A mutant accumulates transcripts targeted to the nonsense-

1125   mediated decay pathway. *J Biol Chem, 281*(46), 35336-35346.

1126   https://doi.org/10.1074/jbc.M601460200

1127

1128 Schuller, A. P., Wu, C. C., Dever, T. E., Buskirk, A. R., & Green, R. (2017, Apr 20). eIF5A

1129   Functions Globally in Translation Elongation and Termination. *Mol Cell, 66*(2), 194-205

1130   e195. https://doi.org/10.1016/j.molcel.2017.03.003

1131

1132 Shuvalov, A., Shuvalova, E., Biziaev, N., Sokolova, E., Evmenov, K., Pustogarov, N.,

1133   Arnautova, A., Matrosova, V., Egorova, T., & Alkalaeva, E. (2021, Nov 12). Nsp1 of

1134   SARS-CoV-2 stimulates host translation termination. *RNA Biol, 18*(sup2), 804-817.

1135   https://doi.org/10.1080/15476286.2021.1999103

1136

1137 Singh, G., Rebbapragada, I., & Lykke-Andersen, J. (2008, Apr 29). A competition between

1138   stimulators and antagonists of Upf complex recruitment governs human nonsense-

1139   mediated mRNA decay. *PLoS Biol, 6*(4), e111.

1140   https://doi.org/10.1371/journal.pbio.0060111

1141

1142 Smith, T., Heger, A., & Sudbery, I. (2017, Mar). UMI-tools: modeling sequencing errors in

1143   Unique Molecular Identifiers to improve quantification accuracy. *Genome Res, 27*(3),

1144   491-499. https://doi.org/10.1101/gr.209601.116

1145

1146 Susorov, D., Egri, S., & Korostelev, A. A. (2020, Dec). Termi-Luc: a versatile assay to monitor

1147   full-protein release from ribosomes. *RNA, 26*(12), 2044-2050.

1148   https://doi.org/10.1261/rna.076588.120

1149

1150 Tate, W. P., & Mannering, S. A. (1996, Jul). Three, four or more: the translational stop signal at

1151   length. *Mol Microbiol, 21*(2), 213-219. https://doi.org/10.1046/j.1365-

1152   2958.1996.6391352.x

1153

1154 Tork, S., Hatin, I., Rousset, J. P., & Fabret, C. (2004). The major 5' determinant in stop codon
1155   read-through involves two adjacent adenines. *Nucleic Acids Res, 32*(2), 415-421.
1156   https://doi.org/10.1093/nar/gkh201
1157

1158 van Engelen, B. G., Muchir, A., Hutchison, C. J., van der Kooi, A. J., Bonne, G., & Lammens, M.
1159   (2005, Jan 25). The lethal phenotype of a homozygous nonsense mutation in the lamin
1160   A/C gene. *Neurology, 64*(2), 374-376.
1161   https://doi.org/10.1212/01.WNL.0000149763.15180.00
1162

1163 Wallmeroth, D., Lackmann, J. W., Kueckelmann, S., Altmuller, J., Dieterich, C., Boehm, V., &
1164   Gehring, N. H. (2022, May 16). Human UPF3A and UPF3B enable fault-tolerant
1165   activation of nonsense-mediated mRNA decay. *EMBO J, 41*(10), e109191.
1166   https://doi.org/10.15252/embj.2021109191
1167

1168 Yamashita, A., Izumi, N., Kashima, I., Ohnishi, T., Saari, B., Katsuhata, Y., Muramatsu, R.,
1169   Morita, T., Iwamatsu, A., Hachiya, T., Kurata, R., Hirano, H., Anderson, P., & Ohno, S.
1170   (2009, May 1). SMG-8 and SMG-9, two novel subunits of the SMG-1 complex, regulate
1171   remodeling of the mRNA surveillance complex during nonsense-mediated mRNA decay.
1172   *Genes Dev, 23*(9), 1091-1105. https://doi.org/10.1101/gad.1767209
1173

1174 Zetoune, A. B., Fontaniere, S., Magnin, D., Anczukow, O., Buisson, M., Zhang, C. X., &
1175   Mazoyer, S. (2008, Dec 5). Comparison of nonsense-mediated mRNA decay efficiency
1176   in various murine tissues. *BMC Genet, 9*, 83. https://doi.org/10.1186/1471-2156-9-83
1177

1178 Zhang, J., & Maquat, L. E. (1997, Feb 17). Evidence that translation reinitiation abrogates
1179   nonsense-mediated mRNA decay in mammalian cells. *EMBO J, 16*(4), 826-833.
1180   https://doi.org/10.1093/emboj/16.4.826

1181

1182 Zhu, X., Zhang, H., & Mendell, J. T. (2020, Jul 14). Ribosome Recycling by ABCE1 Links

1183   Lysosomal Function and Iron Homeostasis to 3' UTR-Directed Regulation and

1184   Nonsense-Mediated Decay. *Cell Rep, 32*(2), 107895.

1185   https://doi.org/10.1016/j.celrep.2020.107895

1186

1187