1 **Impacts of increasing isolation and environmental variation on Florida Scrub-Jay**
2 **demography**
3

4 Jeremy Summers[1] (jsummer7@ur.rochester.edu), Elissa J. Cosgrove[2] (ecj87@cornell.edu),
5 Reed Bowman[3] (rbowman@archbold-station.org), John W. Fitzpatrick[4] (jwf7@cornell.edu),
6 Nancy Chen[1] (nchen11@ur.rochester.edu)
7
8 [1]Department of Biology, University of Rochester, Rochester, NY
9 [2]Department of Molecular Biology & Genetics, Cornell University, Ithaca, NY
10 [3]Avian Ecology Program, Archbold Biological Station, Venus, FL
11 [4]Cornell Lab of Ornithology, Cornell University, Ithaca, NY

12
13 **Abstract**
14

Isolation caused by anthropogenic habitat fragmentation can destabilize populations.
Populations relying on the inflow of immigrants can face reduced fitness due to inbreeding
depression as fewer new individuals arrive. Empirical studies of the demographic consequences
of isolation are critical to understand how populations persist through changing conditions. We
used a 34-year demographic and environmental dataset from a population of cooperatively-
breeding Florida Scrub-Jays (*Aphelocoma coerulescens*) to create mechanistic models linking
environmental and demographic factors to population growth rates. We found that the
population has not declined despite both declining immigration and increasing inbreeding, owing
to a coinciding response in breeder survival. We find evidence of density-dependent
immigration, breeder survival, and fecundity, indicating that interactions between vital rates and
local density play a role in buffering the population against change. Our study elucidates the
impacts of isolation on demography and how long-term stability is maintained via demographic
responses.

28
29 **Introduction**
30
Identifying the potential demographic and environmental drivers of population dynamics
is a fundamental goal of ecology (Cappuccino & Price 1995; Iles *et al.* 2019). Population growth
rates are influenced by density-dependent (*e.g.*, resource competition) and independent (*e.g.*,
climate) factors that interact in ways that remain poorly understood (Goswami *et al.* 2011; Oro
2013). Furthermore, the dynamics of small populations can be affected by intrinsic genetic
factors such as inbreeding depression, or decreased fitness of inbred individuals (Fagan &
Holmes 2006; Kardos *et al.* 2023). As more populations worldwide are threatened by habitat
fragmentation and climate change (Marzluff *et al.* 2001; Simberloff 1999; Young *et al.* 2016),
elucidating the effects of temporal changes in immigration, inbreeding, and environmental
factors on the demography of wild populations is increasingly urgent (Kardos *et al.* 2023;
Schaub & Ullrich 2021).
As populations become more isolated, theory predicts that immigration will decrease and
levels of inbreeding will increase, but downstream effects on population dynamics are more
complicated to predict. Empirical studies have found that immigration may play an important role
in governing population dynamics (Lieury *et al.* 2015; Santoro *et al.* 2016; Tenan *et al.* 2023),
but the demographic impact of immigration varies depending on recent changes in
metapopulation structure (Reichert *et al.* 2021), resource availability (Horswill *et al.* 2022),
temporal variation in the environment (Millon *et al.* 2019), and population density (Harman *et al.*
2020). Decreasing immigration may not always lead to population declines, as immigration may
not be a large driver of variability in population growth rates or may be buffered by other
demographic and environmental factors. For instance, in classic source-sink systems, source

populations that are net producers of immigrants should be insensitive to changes in incoming
immigration (Pulliam 1988). Immigration may be negatively density-dependent, which inherently
stabilizes populations against fluctuations (Sæther *et al.* 1999; Wilson *et al.* 2017). Finally,
temporal variation in immigration could be compensated by changes in other vital rates through
density-dependent processes or differential responses to environmental drivers, leading to
buffering via demographic compensation (Villellas *et al.* 2015). High local population regulation
relative to immigration will make a population more asynchronous from its metapopulation
(Lande *et al.* 1999), which will result in greater local stability when the surrounding
metapopulation is in decline (Fox *et al.* 2017; Rowland *et al.* 2022). Characterizing the relative
contributions of immigration and local survival or fecundity has important implications on the
spatial scale required for conservation (Paquet *et al.* 2021), yet few studies have quantified
temporal changes in immigration and subsequent effects on population dynamics (Horswill *et al.*
2022).
Gene flow into populations via immigration may counter the negative demographic
effects of inbreeding depression (Hedrick & Garcia-Dorado 2016; Niskanen *et al.* 2020; Whitlock
et al. 2000). The extent of population-level effects of inbreeding depression remains
controversial (Bell *et al.* 2019; Keller *et al.* 2007; Kyriazis *et al.* 2020; Mathur & DeWoody 2021;
Szulkin *et al.* 2013), and relatively few studies have documented negative impacts of inbreeding
on population growth rates (Bozzuto *et al.* 2019; Johnson *et al.* 2011; Kardos *et al.*
2023).Theory predicts that the demographic consequences of inbreeding depression depend on
the ecology and life history of a species. Effects of inbreeding on vital rates that do not
contribute much to population growth will not result in large changes in population size (Johnson
et al. 2011; Mills & Smouse 1994). Populations that are primarily regulated by density- or
frequency-dependent (soft) selection may not experience reduced population growth in
response to inbreeding depression (Saccheri & Hanski 2006). Finally, as the strength of
inbreeding depression can vary with environmental conditions, the population-level
consequences of inbreeding may also depend on other ecological conditions. More empirical
studies that integrate the environmental and demographic contexts of both immigration and
inbreeding are needed to clarify the role of increasing isolation on population dynamics.
Quantifying the demographic effects of immigration and inbreeding requires large
amounts of data that are rarely available in natural populations. Direct measures of immigration
require tracking all new individuals in a population to identify immigrants and residents
(Greenwood & Harvey 1982; Hixon *et al.* 2002; Millon *et al.* 2019). While recent advances in
population modeling have allowed for the inference of immigration rates in situations where data
are limited (Abadi *et al.* 2010), these inferences may overestimate the impact of immigration by
failing to account for temporal variation in immigration rates (Nater 2024; Paquet *et al.* 2021).
Similarly, investigating the consequences of inbreeding requires measures of relatedness
between individuals and individual fitness measures (Johnson *et al.* 2011; Sin *et al.* 2021).
Long-term, detailed, population and ecological data obviate the need to infer immigration rates
and provide relatedness and fitness data for multiple individuals, and, while rare, provide the
best opportunity to empirically test expectations of the consequences of changing immigration
rates and inbreeding.
The large amounts of data available through long-term ecological studies can be
analyzed with recently developed transient life table response experiments (tLTRE) (Koons *et*
al. 2016, 2017). tLTREs retrospectively quantify the contribution of different vital rates to
observed variation in population growth rate while accounting for temporal variation in vital rates
and demographic structure (Koons *et al.* 2016, 2017). As opposed to prospective sensitivity
analyses (Caswell 2000), tLTREs consider both the *potential* impact of vital rates on population
growth and the *realized* variation in these rates, making them ideal for attributing observed
change in growth rates to underlying change in vital rates (Coulson *et al.* 2005), including
immigration (Christian *et al.* 2024). These methods can be built upon to link environmental

factors to vital rates through a path analysis (Van Tienderen 2000), ultimately linking variation in
the environment to variation in population growth rate (Coulson & Tuljapurkar 2008; Dahlgren &
Ehrlén 2009; Knape *et al.* 2023; Nater *et al.* 2021).
Here we empirically test how a natural population of Federally Threatened Florida Scrub-
Jays (*Aphelocoma coerulescens*) has remained robust despite increasing isolation. Florida
Scrub-Jays are facultative cooperative breeders: most individuals delay dispersal and assist
with rearing young on their natal territories, though some disperse and help on other territories
before establishing as a breeder (Woolfenden & Fitzpatrick 1984). Females are more likely to
disperse earlier and farther than males, but few other vital rates differ between sexes (Aguillon
et al. 2017; Suh *et al.* 2020; Woolfenden & Fitzpatrick 1984). Florida Scrub-Jays occupy
territories year-round and are restricted to fire-mediated xeric oak scrub habitat (Fitzpatrick &
Bowman 2016). Habitat loss and degradation has led to declines across their range (Boughton
& Bowman 2011; Coulon *et al.* 2008). One color-banded population of Florida Scrub-Jays at
Archbold Biological Station has been monitored for >50 years, accumulating complete life
histories for thousands of individuals, territory and burn area maps, climate and acorn
abundance data, and a 14-generation population pedigree (Chen *et al.* 2019; Fitzpatrick &
Bowman 2016). This population has experienced a significant decline in immigration over the
past 30 years, which has contributed to increased inbreeding and inbreeding depression (Chen
et al. 2016), yet census population sizes have not declined (Figure 1).
We characterized the factors underlying variation in population growth rates over a
period of 34 years (1989-2021) to (1) determine the impact of sex-biased immigration on
population growth rate, (2) quantify the impact of inbreeding, population density, and other
relevant environmental factors on vital rates and population growth rate, and (3) test whether
the demographic impacts of decreased immigration and increased inbreeding have been
counteracted by other vital rates or environmental factors. We constructed both female-specific
and male-specific stage-structured matrix population models and used tLTREs to assess the
contribution of different vital rates to population growth. We used generalized linear models
(GLMs) to link vital rates to population-wide levels of inbreeding, population density, and
environmental factors, and performed environmental tLTREs to determine how these factors
have influenced population growth. These analyses allow us to detect density dependence
directly and quantify the realized impact of changes in immigration and inbreeding, as well as
any other coinciding temporal trend. We predicted that density-dependent responses in the vital
rates of non-breeders would compensate for declining immigration, given the abundance of
resident non-breeders available to fill breeding vacancies (Woolfenden & Fitzpatrick 1984), and
that inbreeding would not strongly influence population growth because of density dependence.
Our study is among the few that use direct measures of immigration and inbreeding to
determine the impact of isolation on the demography of a natural population, which is crucial for
understanding how to preserve threatened populations.
141
142 **Methods**
143
1. Data Collection
1.1 Study Population
We used data from a long-term study of Florida Scrub-Jays at Archbold Biological
Station (Woolfenden & Fitzpatrick 1984). The population is censused monthly, and all nests are
tracked during the breeding season (March-June), providing detailed location, survival, and
breeding records for all individuals within the study area and allowing identification of
immigrants over time. We limited our study area to 14.8 km^2 of land that has been consistently
monitored between 1988-2021. Our dataset includes 4,267 individuals (1,948 female, 1,820
male, and 499 unsexed) from 1,894 territory-years. Please see Supplementary Methods for
details on demographic data collection.

154
1.2 Covariates

We considered several covariates that likely impact Florida Scrub-Jay demography: pedigree-based inbreeding and relatedness coefficients, population density, annual acorn abundance, burned area, and the equatorial southern oscillation index during the dry (breeding) season (EQSOI) (Supplementary Methods). We used mean inbreeding coefficients or pairwise relatedness coefficients estimated from the population pedigree to capture the effect of known inbreeding depression on fecundity and survival in our study population (Chen *et al.* 2016). We calculated population density as the total number of individuals per occupied hectare. As a proxy for habitat quality, we calculated the area burned within the past 2-9 years (the frequency of prescribed fires that generate optimal Florida Scrub-Jay habitat; (Fitzpatrick & Bowman 2016). We used acorn abundance as a metric of resource availability, as acorns are the most important over-winter food source for the Florida Scrub-Jay and vary greatly in abundance over time (DeGange *et al.* 1989; Pesendorfer *et al.* 2021). EQSOI is a commonly used climate index that captures atmospheric conditions that influence El Niño events (Shi & Su 2020), which heavily impact precipitation and temperature in Florida.

2. Contribution of Variation in Vital Rates to Population Growth
2.1 Stage-Structured Matrix Population Model

We constructed matrix population models with four stages (juveniles, helpers, new pairs, and established pairs; Figure 1) and a post-breeding census (Supplementary Methods). We defined juveniles as any individuals between 11 days and 1 year of age, which includes all sexually immature individuals. Helpers are individuals of 1 year of age or older who are not established as a socially dominant breeder. Previous work showed that pairs that had bred together previously had higher fecundity than newly-formed pairs (Woolfenden & Fitzpatrick 1984); therefore we divided breeding pairs into new and established pairs based on whether the two individuals had previously paired during a breeding season. Immigrants (all individuals who were not born within the study area) can join the population as either helpers or breeders, forming new pairs. Extensive population monitoring allowed us to fit both female-only and male-only models using direct measures of immigration and sex-specific vital rates for adults. To account for uncertainty caused by unsexed individuals, we randomly assigned these individuals a sex based on a 1:1 sex ratio (Woolfenden & Fitzpatrick 1984) 100 times for each model and calculated annual vital rates for each iteration to parameterize annual projection matrices (Supplementary Methods).

2.2 Transient Life Table Response Experiment

To quantify how vital rates result in observed variation in population growth rates over time, we performed a tLTRE (Koons *et al.* 2016). This method does not assume that the population is near its stable stage distribution and allows variation in stage structure to contribute to variation in population growth rate (Koons *et al.* 2016). We calculated the realized growth rate as the product of the projection matrix ($A(t)$) that is parameterized with the observed vital rates (θ_i, for all vital rates i) and the stage distribution vector for each year (n). We calculated the sensitivities of the realized growth rate to the vital rates using the *Deriv* package (Clausen *et al.* 2021). We weighted the sensitivity of the population growth rate to each vital rate by the observed variation in vital rates to calculate the variation in growth caused by variation and covariation in vital rates. Covariation can produce a negative contribution when opposite trends in two vital rates result in more similar growth rates over time (*e.g.,* negative covariation between survival and fecundity means high survival years have low fecundity, reducing fluctuations in population size). To include time-lagged effects, we considered the contribution of variation in vital rates from the same year (t) and from the previous year ($t - 1$) to variation in the growth rate. Please see Supplementary Methods for details.

205
3. Impact of Inbreeding, Density, and Environmental Factors on Vital Rates

To determine how environmental variation influences population growth rate, we modeled the underlying vital rates in terms of the covariates described above. We used GLMs with a binomial distribution to model survival for each social class (juveniles, helpers, and breeders) and transition rates from non-breeding to breeding status. We ran separate models for male and female helpers and breeders but not for juveniles due to the large number of unsexed juveniles (480). Note that because Florida Scrub-Jays typically disperse as helpers (Suh *et al.* 2020; Woolfenden & Fitzpatrick 1984), helper mortality may be inflated by emigration, but annual surveys of surrounding habitat patches suggest that emigration rates from our study population are low (Supplementary Methods). We used a gaussian distribution to model log-transformed fecundity and a Poisson distribution to model immigration. We modeled immigration separately for males and females to capture the impact of female-biased dispersal (Woolfenden & Fitzpatrick 1984). Please see Supplementary Methods for full details.

4. Contribution of Inbreeding, Density, and Environmental Variation to Population Growth

To quantify the contributions of our covariates to variation in the population growth rate, we performed a path analysis (Van Tienderen 2000) as an extension of our tLTRE (Knape *et al.* 2023). We decomposed variation in growth caused by each vital rate (quantified in our tLTRE) by the covariate associated with these rates. We analytically solved for the sensitivities of the vital rates to each environmental factor using the link functions and estimated β values from our GLMs. We then used these sensitivities along with the chain rule to convert the sensitivities of the population growth rate to the observed vital rates into the sensitivities of the population growth rate to each environmental factor and performed another tLTRE using the covariates in place of the vital rates. We summed the contributions to variation in the population growth rate based on each covariate. Please see Supplementary Methods for details.

5. Simulations

To confirm our hypothesis that changes in other vital rates compensated for changes in immigration and inbreeding, we ran post-hoc simulations to project population sizes under different scenarios. We projected our initial population vectors from 1988 using our projection matrices, assigning values for vital rates using predictions from our GLMs. This approach allows us to simulate vital rates under different conditions. We detrended vital rates by using the same value for the time fixed effect in all years. To incorporate density dependence, we recalculated population density each year based on the projected population size and used that value to predict vital rates for the next year. We simulated three scenarios: (1) decreasing immigration and increasing inbreeding with increasing breeder survival, (2) no temporal trends in breeder survival, and (3) no temporal trends in breeder survival, immigration, or inbreeding. Please see Supplemental Methods for details.

Results

1. Demographic and Environmental Trends

Despite previously-documented decreases in immigration (Mann Kendall tau = -0.337, *p* = 0.007) and increases in population-wide inbreeding (Mann Kendall tau = 0.405, *p* = 0.001), the census population size remained relatively stable (coefficient of variance = 0.167; Figure 1; Reed & Hobbs 2004). In fact, we found an increase in population size over time (Mann Kendall tau = 0.273, *p* = 0.024), largely driven by an increase in the number of established pairs (Mann Kendall tau = 0.414, *p* = 0.007; Figure 1A; Table S2). Annual growth rates fluctuated rapidly between growth ($\lambda > 1$) and decline ($\lambda < 1$; Figure S1). We found no significant temporal trend in

the annual growth rate (Mann Kendall tau = -0.004, *p* = 0.987), and the geometric mean of λ is
0.998 (Figure S1, Table S1). When we considered sex-specific vital rates, we only found a
significant decline in male immigration after correcting for multiple tests (Mann Kendall tau = -
0.405, *q* = 0.010, Figure S2, Table S2). Inbreeding significantly increased over time (Figure S5,
Table S3), consistent with previous work (Chen *et al.* 2016).

2. Contribution of Variation in Vital Rates to Population Growth
We used tLTREs to quantify how different vital rates contribute to differences in
population growth over time (Figure S7). Results for our female-only and male-only models
were similar, thus we report the results for the female-only model except when noted. Temporal
variation in population growth rates was primarily driven by fecundity (55.1%) and survival
(32.8%, Figure 2), specifically established pair fecundity (28.9%) and breeder survival (18.9%,
Figure S7). Immigration accounted for 4.8% of variation in population growth rate in the female-
only model and -4.3% in the male-only model due to negative covariation between male
immigration and both survival and fecundity (Figure 2). Probability to pair accounted for 4.5% of
the variation in the population growth rate (Figure 2).

3. Impact of Inbreeding, Density, and Environmental Factors on Vital Rates
To determine how inbreeding, population density, and environmental conditions
influence annual variation in vital rates, we ran GLMs for each vital rate. Consistent with
previous work, we found that higher inbreeding in the population was significantly associated
with lower juvenile survival (β = -0.133, p = 0.01), male breeder survival (β = -0.259, p = 0.008,
Table S3), and lower fecundity (β = -0.084, p = 0.03, Table S5). We found negative density
dependence in male breeder survival (β = -0.170, *p* = 0.007, Table S3), female immigration (β =
-0.166, *p* = 0.009, Table S4), both new (β = -0.267, *p* < 0.001) and established pair fecundity (β
= -0.123, *p* = 0.003, Table S5), and the probability of female helpers to form new pairs (β = -
0.198, *p* = 0.046). Of our environmental factors, only acorn abundance had a significant
association with any vital rates. Greater acorn abundance was associated with increased
survival of juveniles (β = 0.196, *p* = 0.001), female breeders (β = 0.240, *p* = 0.002), and male
breeders (β = 0.154, *p* = 0.029, Figure 3, Table S3). To account for any temporal trends that
couldn't be explained by our other fixed effects, we included a fixed effect of time. We found a
significant negative effect of time on female (β = -0.185, *p* = 0.007) and male immigration (β = -
0.378, *p* < 0.001, Table S4). Though we did not find any temporal trends in breeder survival
(Table S2), we found significant positive impacts of time on female (β = 0.338, *p* = 0.001) and
male breeder (β = 0.417, *p* < 0.001) survival (Table S3).

4. Contribution of Inbreeding, Density, and Environmental Variation to Population
Growth
To quantify the effect of inbreeding, density, and environmental factors on population
growth, we calculated the sensitivities of population growth to these factors and summed the
contribution of each factor through their impact on vital rates. Population growth rate had similar
sensitivities to inbreeding, population density, acorn abundance, and temporal trends, with
negative effects of inbreeding and density and positive effects of acorn abundance and time
(Figure 4A). Sensitivity analysis indicates that vital rates with positive temporal trends (breeder
survival) more than compensated for those with negative temporal trends (immigration) (Figure
3). When we weighted sensitivities by the observed variation in each factor, we found that
variation in population density contributed the most to variation in population growth rate
(12.1%), followed by variation in acorn abundance (7.1%) and EQSOI (0.6%, Figure 4B).
Variation in inbreeding and temporal trends (variation in time) both contributed very little (>0.1%
and 0.5%, respectively; Figure 4), largely because negative covariation between them canceled
out any potential contributions to population growth (Figure 4B). Negative covariation between

inbreeding and density as well as a slight decline in acorn abundance over time (Figure S5,
Table S2) also reduced fluctuations in population size (Figure 4).
309
5. Simulations
We simulated density-dependent population growth to test whether the observed
increase in breeder survival was sufficient to compensate for the observed decline in
immigration and increase in inbreeding. We first performed simulations that included all
observed temporal trends as a control. While these simulations consistently underestimated
population size in later years, they accurately predicted the observed population stability overall
(Figure 5). Divergence between observed and simulated population dynamics is driven by a few
unusually high fecundity years. When we simulated dynamics without a temporal increase in
breeder survival, the population declined in both sex-specific models, with a more dramatic
decrease in the male-only model (Figure 5). This decline in population size is rescued in both
models when we also removed temporal trends in immigration and inbreeding (Figure 5).
321
322 **Discussion**
323
We investigated how a wild population has maintained population size despite
increasing isolation by quantifying the effect of variation in local demographic rates, immigration,
inbreeding, population density, and environmental factors on variation in population growth. The
stability of our Florida Scrub-Jay population results from density dependence and compensation
by breeder survival. Though immigration does contribute to variability in population growth rates,
we found a positive trend in breeder survival over time (after accounting for annual fluctuations
in inbreeding, population density, and acorn abundance) that outweighs the demographic
impact of declines in immigration and increasing inbreeding. This result contradicts our initial
prediction that flexibility in the demography of non-breeders would buffer the population at
saturation during changing conditions (Porter & Coulson 1987). We found direct evidence of
density dependence in female immigration and covariation between male immigration and both
survival and fecundity, suggesting that density dependent immigration also contributed to
stability. Finally, despite evidence of inbreeding depression in vital rates with high contributions
to population growth (survival and fecundity), inbreeding had a negligible effect on variation in
population growth rate because of compensation by the temporal trend in breeder survival and
negative covariation between inbreeding and density.
Our results demonstrate the power of long-term ecological studies to provide highly
detailed insight into population regulation during changing conditions. The compensation of
both declining immigration and increasing inbreeding by breeder survival, resulting in a
significantly greater number of established breeders over time (Table S2), demonstrates that
significant shifts in local demography were required to maintain population stability through
increasing isolation. Multiple studies have documented how increasing immigration drove
increases in other vital rates that allowed newly-established populations to become self-
sufficient source populations (Herman & Lynch 2022; Santoro *et al.* 2016; Tenan *et al.* 2017).
Our study presents a rare example of the opposite occurrence: an existing stable population
adjusting to decreased immigration and becoming more self-sufficient via compensatory
increases in other vital rates. Characterizing how populations can remain robust to increasing
isolation is important for identifying the appropriate spatial scale for effective management of
populations threatened by habitat fragmentation (Paquet *et al.* 2021)..
The observed increase in breeder survival could be a result of decreased immigration or
of other trends. Because Florida Scrub-Jays engage in potentially risky territorial defense
behaviors (Woolfenden & Fitzpatrick 1984), decreased immigration could have increased
breeder survival through relaxed competition for breeding territories. Alternatively, immigrants
frequently have reduced fitness compared to residents in birds (Barbraud & Delord 2021; Dickel

et al. 2023), thus declining immigration could lead to increased average breeder survival.
However, residents and immigrants do not have significantly different breeder lifespans in our
population (Wilcoxon signed-rank test, W = 62241, *p* = 0.22). It is also possible that increased
breeder survival has contributed to decreased immigration instead, via more limited breeding
vacancies. The observed temporal change in breeder survival could also be caused by a
reduction in negative density dependence, decreased reliance on acorns, or decreased effects
of inbreeding, or as a result of successional change in the habitat due to the prescribed fire
regime (Pesendorfer *et al.* 2021). While we detected no impact of burned area on breeder
survival, prescribed fires can increase the availability of arthropod prey (Goud 2017) and create
more optimal habitat structure that can increase breeder survival (Breininger *et al.* 2009;
Fitzpatrick & Bowman 2016).
We found that inbreeding contributed relatively little to variation in the population growth
rate, consistent with previous findings that inbreeding depression may not directly translate into
population decline (Johnson *et al.* 2011). The overall impact of inbreeding was reduced by
covariation between inbreeding and density (Figure 4B), which supports the soft selection
hypothesis (Bozzuto *et al.* 2019; Keller *et al.* 2007; Wallace 1975). Populations experiencing
soft selection primarily include those where intraspecific competition is important (Bell *et al.*
2019), including highly territorial species such as the Florida Scrub-Jay. Our work provides an
empirical example of density dependent regulation maintaining population growth despite
observable inbreeding depression. However, we note that the level of inbreeding observed in
our populations remains relatively low when compared to populations experiencing extinction
vortices, a negative feedback loop of population decline and inbreeding depression (Fagan &
Holmes 2006; Frankham 2015; Kardos *et al.* 2023; Pimm *et al.* 2006). It remains to be seen
whether demographic effects of inbreeding will change as population-wide inbreeding
coefficients continue to increase. Future work should test the level of inbreeding required for a
significant demographic response, with particular attention paid to effects on population density
and competition for limited resources, such as breeding opportunities.
Our study emphasizes the importance of considering both males and females when
modeling population dynamics when – as is often the case – sex differences exist among
demographic processes. For example, we found evidence of density dependence in immigration
of females but not males, which could be a result of female-biased dispersal. Similar sex-
specific population dynamics have been found in sexually monomorphic species that exhibit
some sex differences in vital rates (Kraus *et al.* 2008), although density dependence has been
found in immigration of both sexes for other bird species that also exhibit female-biased
dispersal (Wilson *et al.* 2017). Our simulations predicted greater decline without compensation
by breeder survival in males compared to females, likely because of the greater decline in male
immigration over time and lack of density dependent buffering in male immigration. Had we
used the standard approach of only fitting female-only models, we would have underestimated
the role of demographic compensation in maintaining stability and overestimated the importance
of density dependent immigration.
Combining tLTREs and a path analysis is a powerful approach for evaluating the effects
of inbreeding, population density, and environmental factors on population growth; however,
successful application of this approach requires careful evaluation of model assumptions. Our
GLMs cannot specify causation, but the tLTRE provides direct information on how observed
variation in vital rates results in observed variation in population growth rate. Our retrospective
analysis incorporates the contributions of covariances between vital rates or covariates to
observed population dynamics, making it difficult to quantify the effects of single variables in
isolation and make predictions about dynamics in unobserved conditions (an advantage of
prospective analyses). We added simulations to better elucidate how changes in individual
factors lead to different potential outcomes, though our simulations assume that modeled
relationships between vital rates and cofactors remain consistent. Our GLMs and thus the

environmental tLTRE assume a linear relationship between individual covariates and vital rates.
Non-linear relationships could alter the contributions of the covariates as their values change
over time, leading to over- or underestimates of the effect of that covariate on population growth
depending on the functional form of the relationship. For example, Allee effects could result in a
reversal of density dependence at low densities, causing the sensitivity of the population to
change sign (Kramer *et al.* 2009). None of our models violated the linearity assumption
(Supplementary Information). We suggest that future work consider the functional relationships
between covariates and vital rates. Our matrix population models assume uniformity in vital
rates between resident and immigrant individuals. This assumption does not impact predictions
of population growth, but can affect estimates of the impact of immigration. We found no
significant differences in vital rates between residents and immigrants in our data
(Supplementary Information). If that were not the case, immigration rates could impact future
vital rates and thus population growth over time. Finally, we note that uncertainty in our
analyses is not randomly distributed over our study period. The proportion of unsexed
individuals and pedigree incompleteness is greater during the first 10 years of our dataset
(Figure 1). We do not include sex-specific vital rate estimates for juveniles (the majority of our
unsexed individuals), and we found no significant difference between our pedigree-derived
inbreeding coefficients for all individuals vs. a subset of individuals with more complete pedigree
information (Figure S8). Future work would benefit from more precise estimates of emigration
from our focal population, as density dependent emigration is another important aspect of
dispersal-mediated population regulation (Doncaster *et al.* 1997; Harman *et al.* 2020).
Additionally, identifying the source populations of immigrants would allow us to perform
metapopulation analyses and study the impact of dispersal at different geographic scales.
We provide much needed empirical evidence on the demographic consequences of
temporal variation in immigration and inbreeding. Our results show that compensation by other
vital rates can maintain population growth rates as immigration declines or inbreeding
increases. Our focal population thus represents a case where local density dependence is
sufficient to regulate the population as it becomes more independent of the surrounding
metapopulation. This result emphasizes the importance of maintaining high quality habitat that
is capable of supporting such robust populations in the face of habitat fragmentation.Our study
demonstrates the power of tLTREs and path analyses for characterizing potential demographic
and environmental drivers of population dynamics in changing environments.
441
442 **Data Accessibility**
443   All data and scripts used in this study are publicly available at
444 github.com/jtsummers53/LTRE_project.
445
446
447 **Acknowledgments**
448

449   We thank the staff and interns at Archbold Biological Station who have contributed
450 greatly over many years to collecting the ecological and demographic data that makes this
451 research possible. We also thank the fellow members of the Chen Lab (Kristin Hardy, Bailey
452 Jones, Faye Romero, Shailee Shah, and Daniel Seidman) for guidance that shaped the
453 structure of this manuscript. J.S. was supported by NIH grant 1R35GM133412 to N.C.

References

Abadi, F., Gimenez, O., Ullrich, B., Arlettaz, R. & Schaub, M. (2010). Estimation of immigration rate using integrated population models. *J. Appl. Ecol.*, 47, 393–400.

Aguillon, S.M., Fitzpatrick, J.W., Bowman, R., Schoech, S.J., Clark, A.G., Coop, G., *et al.* (2017). Deconstructing isolation-by-distance: The genomic consequences of limited dispersal. *PLoS Genet.*, 13, 1–27.

Barbraud, C. & Delord, K. (2021). Selection against immigrants in wild seabird populations. *Ecol. Lett.*, 24, 84–93.

Bell, D.A., Robinson, Z.L., Funk, W.C., Fitzpatrick, S.W., Allendorf, F.W., Tallmon, D.A., *et al.* (2019). The Exciting Potential and Remaining Uncertainties of Genetic Rescue. *Trends Ecol. Evol.*, 34, 1070–1079.

Boughton, R.K. & Bowman, R. (2011). *State wide assessment of Florida Scrub-Jays on managed areas : A comparison of current populations to the results of the 1992-93 survey*.

Bozzuto, C., Biebach, I., Muff, S., Ives, A.R. & Keller, L.F. (2019). Inbreeding reduces long-term growth of Alpine ibex populations. *Nat. Ecol. Evol.*, 3, 1359–1364.

Breininger, D.R., Nichols, J.D., Carter, G.M. & Oddy, D.M. (2009). Habitat-specific breeder survival of Florida Scrub-Jays: inferences from multistate models. *Ecology*, 90, 3180–3189.

Cappuccino, N. & Price, P.W. (1995). *Population Dynamics: New Approaches and Synthesis*. Elsevier.

Caswell, H. (2000). Prospective and Retrospective Perturbation Analyses : Their Roles in Conservation Biology. *Ecology*, 81, 619–627.

Chen, N., Cosgrove, E.J., Bowman, R., Fitzpatrick, J.W. & Clark, A.G. (2016). Genomic Consequences of Population Decline in the Endangered Florida Scrub-Jay. *Curr. Biol.*, 26, 2974–2979.

Chen, N., Juric, I., Cosgrove, E.J., Bowman, R., Fitzpatrick, J.W., Schoech, S.J., *et al.* (2019). Allele frequency dynamics in a pedigreed natural population. *Proc. Natl. Acad. Sci. U. S. A.*, 116, 2158–2164.

Christian, M., Oosthuizen, W.C., Bester, M.N. & de Bruyn, P.J.N. (2024). Robustly estimating the demographic contribution of immigration: Simulation, sensitivity analysis and seals. *J. Anim. Ecol.*, 93, 632–645.

Clausen, A., Sokol, S. & Rappold, A. (2021). Deriv: Symbolic Differentiation.

Coulon, A., Morellet, N., Goulard, M., Cargnelutti, B., Angibault, J.M. & Hewison, A.J.M. (2008). Inferring the effects of landscape structure on roe deer (Capreolus capreolus) movements using a step selection function. *Landsc. Ecol.*, 23, 603–614.

Coulson, T., Gaillard, J.M. & Festa-Bianchet, M. (2005). Decomposing the variation in population growth into contributions from multiple demographic rates. *J. Anim. Ecol.*, 74, 789–801.

Coulson, T. & Tuljapurkar, S. (2008). The dynamics of a quantitative trait in an age-structured population living in a variable environment. *Am. Nat.*, 172, 599–612.

Dahlgren, J.P. & Ehrlén, J. (2009). Linking environmental variation to population dynamics of a forest herb. *J. Ecol.*, 97, 666–674.

DeGange, A.R., Fitzpatrick, J.W., Layne, J.N. & Woolfenden, G.E. (1989). Acorn harvesting by Florida scrub jays. *Ecology*, 70, 348–356.

Dickel, L., Arcese, P., Keller, L.F., Nietlisbach, P., Goedert, D., Jensen, H., *et al.* (2023). Multigenerational Fitness Effects of Natural Immigration Indicate Strong Heterosis and Epistatic Breakdown in a Wild Bird Population. *Am. Nat.*, 000–000.

Doncaster, C.P., Clobert, J., Doligez, B., Gustafsson, L. & Danchin, E. (1997). Balanced Dispersal between Spatially Varying Local Populations: An Alternative to the Source-

Sink Model. *Am. Nat.*, 150, 425–445.

Fagan, W.F. & Holmes, E.E. (2006). Quantifying the extinction vortex. *Ecol. Lett.*, 9, 51–60.

Fitzpatrick, J.W. & Bowman, R. (2016). Florida scrub-jays: oversized territories and group
defense in a fire-maintained habitat. In: *Cooperative breeding in vetebrates: studies of*
ecoogy, evolution, and behavior. Cambridge University Press, Cambridge, UK, pp. 77–
96.

Fox, J.W., Vasseur, D., Cotroneo, M., Guan, L. & Simon, F. (2017). Population extinctions can
increase metapopulation persistence. *Nat. Ecol. Evol.*, 1, 1271–1278.

Frankham, R. (2015). Genetic rescue of small inbred populations: meta-analysis reveals large
and consistent benefits of gene flow. *Mol. Ecol.*, 24, 2610–2618.

Goswami, V.R., Getz, L.L., Hostetler, J.A., Ozgul, A. & Oli, M.K. (2011). Synergistic influences
of phase, density, and climatic variation on the dynamics of fluctuating populations.
Ecology, 92, 1680–1690.

Goud, E.M. (2017). Diversity and abundance of litter-dwelling arthropods increase with time-
since-burn in a Florida scrub ecosystem. *Biodiversity*, 18, 151–155.

Greenwood, P.J. & Harvey, P.H. (1982). The natal and breeding dispersal of birds. *Annu. Rev.*
Ecol. Syst. Vol. 13, 13, 1–21.

Harman, R.R., Goddard, J., Shivaji, R. & Cronin, J.T. (2020). Frequency of Occurrence and
Population-Dynamic Consequences of Different Forms of Density-Dependent
Emigration. *Am. Nat.*, 195, 851–867.

Hedrick, P.W. & Garcia-Dorado, A. (2016). Understanding Inbreeding Depression, Purging, and
Genetic Rescue. *Trends Ecol. Evol.*, 31, 940–952.

Herman, R.W. & Lynch, H.J. (2022). Age-structured model reveals prolonged immigration is key
for colony establishment in Gentoo Penguins. *Ornithol. Appl.*, 124, duac014.

Hixon, M.A., Pacala, S.W. & Sandin, S.A. (2002). Population regulation: Historical context and
contemporary challenges of open vs. closed systems. *Ecology*, 83, 1490–1508.

Horswill, C., Wood, M.J. & Manica, A. (2022). Temporal change in the contribution of
immigration to population growth in a wild seabird experiencing rapid population decline.
Ecography, 2022, e05846.

Iles, D.T., Rockwell, R.F. & Koons, D.N. (2019). Shifting Vital Rate Correlations Alter Predicted
Population Responses to Increasingly Variable Environments. *Am. Nat.*, 193, E57–E64.

Johnson, H.E., Mills, L.S., Wehausen, J.D., Stephenson, T.R. & Luikart, G. (2011). Translating
Effects of Inbreeding Depression on Component Vital Rates to Overall Population
Growth in Endangered Bighorn Sheep. *Conserv. Biol.*, 25, 1240–1249.

Kardos, M., Zhang, Y., Parsons, K.M., A, Y., Kang, H., Xu, X., *et al.* (2023). Inbreeding
depression explains killer whale population dynamics. *Nat. Ecol. Evol.*, 1–12.

Keller, L.F., Biebach, I. & Hoeck, P.E.A. (2007). The need for a better understanding of
inbreeding effects on population growth. *Anim. Conserv.*, 10, 286–287.

Knape, J., Paquet, M., Arlt, D., Kačergytė, I. & Pärt, T. (2023). Partitioning variance in
population growth for models with environmental and demographic stochasticity. *J.*
Anim. Ecol., 92, 1979–1991.

Koons, D.N., Arnold, T.W. & Schaub, M. (2017). Understanding the demographic drivers of
realized population growth rates: *Ecol. Appl.*, 27, 2102–2115.

Koons, D.N., Iles, D.T., Schaub, M. & Caswell, H. (2016). A life-history perspective on the
demographic drivers of structured population dynamics in changing environments. *Ecol.*
Lett., 19, 1023–1031.

Kramer, A.M., Dennis, B., Liebhold, A.M. & Drake, J.M. (2009). The evidence for Allee effects.
Popul. Ecol., 51, 341–354.

Kraus, C., Eberle, M. & Kappeler, P.M. (2008). The costs of risky male behaviour: sex
differences in seasonal survival in a small sexually monomorphic primate. *Proc. R. Soc.*
B Biol. Sci., 275, 1635–1644.

Kyriazis, C., Wayne, R. & Lohmueller, K. (2020). Strongly deleterious mutations are a primary
determinant of extinction risk due to inbreeding depression. *Evol. Lett.*, 678524.
Lande, R., Engen, S. & Sæther, B. (1999). Spatial Scale of Population Synchrony:
Environmental Correlation versus Dispersal and Density Regulation. *Am. Nat.*, 154,
271–281.
Lieury, N., Gallardo, M., Ponchon, C., Besnard, A. & Millon, A. (2015). Relative contribution of
local demography and immigration in the recovery of a geographically-isolated
population of the endangered Egyptian vulture. *Biol. Conserv.*, 191, 349–356.
Marzluff, J.M., Bowman, R. & Donnelly, R. (2001). *Avian ecology and conservation in an*
urbanizing world.
Mathur, S. & DeWoody, J.A. (2021). Genetic load has potential in large populations but is
realized in small inbred populations. *Evol. Appl.*, 14, 1540–1557.
Millon, A., Lambin, X., Devillard, S. & Schaub, M. (2019). Quantifying the contribution of
immigration to population dynamics: a review of methods, evidence and perspectives in
birds and mammals. *Biol. Rev.*, 94, 2049–2067.
Mills, L.S. & Smouse, P.E. (1994). Demographic Consequences of Inbreeding in Remnant
Populations. *Am. Nat.*, 144, 412–431.
Nater, C.R. (2024). Redefining 'state-of-the-art' for integrated population models with
immigration. *J. Anim. Ecol.*, 93, 520–524.
Nater, C.R., Eide, N.E., Pedersen, Å.Ø., Yoccoz, N.G. & Fuglei, E. (2021). Contributions from
terrestrial and marine resources stabilize predator populations in a rapidly changing
climate. *Ecosphere*, 12, e03546.
Niskanen, A.K., Billing, A.M., Holand, H., Hagen, I.J., Araya-Ajoy, Y.G., Husby, A., *et al.* (2020).
Consistent scaling of inbreeding depression in space and time in a house sparrow
metapopulation. *Proc. Natl. Acad. Sci. U. S. A.*, 117, 14584–14592.
Oro, D. (2013). Grand challenges in population dynamics. *Front. Ecol. Evol.*, 1, 1–2.
Paquet, M., Knape, J., Arlt, D., Forslund, P., Pärt, T., Flagstad, Ø., *et al.* (2021). Integrated
population models poorly estimate the demographic contribution of immigration.
Methods Ecol. Evol., 12, 1899–1910.
Pesendorfer, M.B., Bowman, R., Gratzer, G., Pruett, S., Tringali, A. & Fitzpatrick, J.W. (2021).
Fire history and weather interact to determine extent and synchrony of mast-seeding in
rhizomatous scrub oaks of Florida. *Philos. Trans. R. Soc. B Biol. Sci.*, 376, 20200381.
Pimm, S.L., Dollar, L. & Bass, O.L. (2006). The genetic rescue of the Florida panther. *Anim.*
Conserv., 9, 115–122.
Porter, J.M. & Coulson, J.C. (1987). Long-term Changes in Recruitment to the Breeding Group,
and the Quality of Recruits at a Kittiwake Rissa tridactyla Colony. *J. Anim. Ecol.*, 56,
675–689.
Pulliam, H.R. (1988). Sources , Sinks , and Population Regulation. *Am. Nat.*, 132, 652–661.
Reed, D.H. & Hobbs, G.R. (2004). The relationship between population size and temporal
variability in population size. *Anim. Conserv. Forum*, 7, 1–8.
Reichert, B.E., Fletcher, R.J. & Kitchens, W.M. (2021). The demographic contributions of
connectivity versus local dynamics to population growth of an endangered bird. *J. Anim.*
Ecol., 90, 574–584.
Rowland, F.E., Schyling, E.S., Freidenburg, L.K., Urban, M.C., Richardson, J.L., Arietta, A.Z.A.,
et al. (2022). Asynchrony, density dependence, and persistence in an amphibian.
Ecology, 103, e3696.
Saccheri, I. & Hanski, I. (2006). Hard and soft selection. *Trends Ecol. Evol.*, 6, 341–347.
Sæther, B.-E., Engen, S. & Lande, R. (1999). Finite metapopulation models with density–
dependent migration and stochastic local dynamics. *Proc. R. Soc. Lond. B Biol. Sci.*,
266, 113–118.
Santoro, S., Green, A.J. & Figuerola, J. (2016). Immigration enhances fast growth of a newly

established source population. *Ecology*, 97, 1048–1057.
Schaub, M. & Ullrich, B. (2021). A drop in immigration results in the extinction of a local
woodchat shrike population. *Anim. Conserv.*
Shi, Y. & Su, J. (2020). A New Equatorial Oscillation Index for Better Describing ENSO and
Westerly Wind Bursts. *J. Meteorol. Res.*, 34, 1025–1037.
Simberloff, D. (1999). Habitat fragmentation and population extinction of birds. *NCASI Tech.*
Bull., 2, 491–492.
Sin, S.Y.W., Hoover, B.A., Nevitt, G.A. & Edwards, S.V. (2021). Demographic History, Not
Mating System, Explains Signatures of Inbreeding and Inbreeding Depression in a Large
Outbred Population. *Am. Nat.*, 197, 658–676.
Suh, Y.H., Pesendorfer, M.B., Tringali, A., Bowman, R. & Fitzpatrick, J.W. (2020). Investigating
social and environmental predictors of natal dispersal in a cooperative breeding bird.
Behav. Ecol., 1–10.
Szulkin, M., Stopher, K.V., Pemberton, J.M. & Reid, J.M. (2013). Inbreeding avoidance,
tolerance, or preference in animals? *Trends Ecol. Evol.*, 28, 205–211.
Tenan, S., Fasola, M., Volponi, S. & Tavecchia, G. (2017). Conspecific and not performance-
based attraction on immigrants drives colony growth in a waterbird. *J. Anim. Ecol.*, 86,
1074–1081.
Tenan, S., Moulins, A., Tepsich, P., Bocconcelli, A., Verga, A., Ballardini, M., *et al.* (2023).
Immigration as the main driver of population dynamics in a cryptic cetacean. *Ecol. Evol.*,
13, e9806.
Van Tienderen, P.H. (2000). Elasticities and the Link Between Demographic and Evolutionary
Dynamics. *Ecology*, 81, 666–679.
Villellas, J., Doak, D.F., García, M.B. & Morris, W.F. (2015). Demographic compensation among
populations: what is it, how does it arise and what are its implications? *Ecol. Lett.*, 18,
1139–1152.
Wallace, B. (1975). Hard and Soft Selection Revisited. *Evolution*, 29, 465–473.
Whitlock, M.C., Ingvarsson, P.K. & Hatfield, T. (2000). Local drift load and the heterosis of
interconnected populations. *Heredity*, 84, 452–457.
Wilson, S., McKellar, A.E., Reudink, M.W., Marra, P.P. & Ratcliffe, L.M. (2017). Density-
dependent immigration promotes population stability in a long-distance migratory bird.
Popul. Ecol., 59, 169–178.
Woolfenden, G.E. & Fitzpatrick, J.W. (1984). *The Florida Scrub Jay: Demography of a*
Cooperative-breeding Bird. Princeton University Press.
Young, H.S., McCauley, D.J., Galetti, M. & Dirzo, R. (2016). Patterns, Causes, and
Consequences of Anthropocene Defaunation. *Annu. Rev. Ecol. Evol. Syst.*, 47, 333–
358.

646
Figure 1. Demography of the Florida Scrub Jay. (A) Total population size over the study period
divided by sex and social class. (B) Mean inbreeding coefficient of all individuals over time. (C)
Total number of new male and female immigrants arriving to the population over time. (D) Life
cycle diagram for the Florida Scrub Jay that describes the movement of individuals across
social classes. Individuals who survive to the next breeding season may remain in their existing
pair status (pink arrows), form new pairs (dark blue arrows), or lose a mate through divorce or
death (gray arrows). New individuals enter the population either through reproduction (orange
arrows) or immigration (purple arrows).

Figure 2. Contributions of variation in vital rates to variation in population growth rate for the
female-only model (left) and male-only model (right). Bars are colored by the covariance
involved (*e.g.*, pink bars in the fecundity column are contributions made by covariance between
fecundity and survival). Negative contributions indicate that the covariation between two vital
rates diminished the variation in the population growth rate (*i.e.*, one rate promoted growth while
the other promoted decline).

665 **Figure 3.** Impact of inbreeding, density, environmental variables, and year on vital rates. Error
666 bars represent the 95% confidence interval. Colors indicate the type of vital rate modeled (pink:
667 survival, purple: immigration, yellow: fecundity, dark blue: probability to pair). Juvenile vital rates
668 are not divided between males and females due to the relatively large number of unsexed
669 juveniles. Asterisks indicate significance (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).

Figure 4. Contributions of variation in inbreeding, density, and environmental factors to variation in population growth rate. (A) Sensitivity of the population growth rate to inbreeding, density, acorn abundance, burned area, EQSOI, and year. The distributions are the result of resampling the effect sizes from the GLMs. The mean sensitivity is indicated with a white bar. (B) Inbreeding, density, and environmental factors contributed to the population growth rate through their association with vital rates. Bars are colored by covariances (*e.g.*, the yellow bar in the acorn abundance column shows the contribution made by covariance between acorn abundance and year). Negative contributions indicate that the covariation between two factors diminished the variation in the population growth rate (*i.e.*, one factor promoted growth while the other promoted decline).

Figure 5. Observed and simulated population dynamics for females (left) and males (right). Density-dependent simulations that used vital rates predicted with observed temporal trends in immigration, inbreeding, and breeder survival (pink; Modeled with Temporal Trends) matched observed dynamics (black). Simulations that removed the temporal trend in breeder survival (purple; Modeled with Breeder Survival Detrended) resulted in population decline, and we recovered observed population dynamics when we removed temporal trends in immigration, inbreeding, and breeder survival (blue; Modeled with No Temporal Trends). Ribbons indicate the 95% confidence interval for simulated population size.