Binding domain mutations provide insight into CTCF's relationship with chromatin and its contribution to gene regulation

Catherine Do[1,2,*], Guimei Jiang[1,2,*], Giulia Cova[1,2*], Christos C. Katsifis[3,4,5,φ], Domenic N. Narducci[3,4,5.φ,] Theodore Sakellaropoulos[1,2], Raphael Vidal[1,2], Priscillia Lhoumaud[1,2], Aristotelis Tsirigos [1,2,6], Faye Fara D. Regis[1,2], Nata Kakabadze[1,2], Elphege P Nora[7], Marcus Noyes[8], Anders S. Hansen[3,4,5], Jane A Skok[1,2,†]

1. Department of Pathology, NYU Grossman School of Medicine, New York, NY 10016, USA
2. Perlmutter Cancer Center, NYU Langone Health, New York, NY 10016, USA
3. MIT Department of Biological Engineering, Cambridge, MA 02139, USA
4. Gene Regulation Observatory, Broad Institute of MIT and Harvard, Cambridge, MA 02139, USA
5. Koch Institute for Integrative Cancer Research, Cambridge, MA, 02139, USA
6. Applied Bioinformatics Laboratories, Office of Science & Research, NYU Grossman School of Medicine, New York, NY 10016, USA
7. Cardiovascular Research Institute, and Department of Biochemistry and Biophysics, University of California San Francisco, CA 94143, USA
8. Institute for Systems Genetics, Department of Biochemistry and Molecular Pharmacology, NYU Langone Health, New York, NY 10016, USA

[*,φ] These authors contributed equally.
[†] Lead contact
Corresponding author:
Jane.Skok@nyulangone.org

Summary 150 words.

Here we used a series of CTCF mutations to explore CTCF's relationship with chromatin and its
contribution to gene regulation. CTCF's impact depends on the genomic context of bound sites
and the unique binding properties of WT and mutant CTCF proteins. Specifically, CTCF's signal
strength is linked to changes in accessibility, and the ability to block cohesin is linked to its binding
stability. Multivariate modelling reveals that both CTCF and accessibility contribute independently
to cohesin binding and insulation, however CTCF signal strength has a stronger effect. CTCF and
chromatin have a bidirectional relationship such that at CTCF sites, accessibility is reduced in a
cohesin-dependent, mutant specific fashion. In addition, each mutant alters TF binding and
accessibility in an indirect manner, changes which impart the most influence on rewiring
transcriptional networks and the cell's ability to differentiate. Collectively, the mutant perturbations
provide a rich resource for determining CTCF's site-specific effects.

Introduction

The CCCTC-binding factor (CTCF) is an eleven-zinc finger DNA-binding protein that together with its binding partner, cohesin plays a key role in organizing chromatin into TAD structures by promoting the formation of loops and boundaries that are important for gene regulation. This involves a loop-extrusion mechanism in which cohesin complexes create chromatin loops by actively extruding DNA until movement of the complex is blocked by two CTCF binding sites in convergent orientation[1,2]. CTCF, in conjunction with cohesin, is enriched at TAD boundaries that function as insulators, contributing to gene regulation by restricting the interaction of regulatory elements to promoters of target genes located within the same TAD[3,4].

CTCF has been degraded by auxin in numerous cell types, and although these studies provide important insight into its role in gene regulation, degron models are insufficient for (i) analyzing the contribution of genomic context to site-specific CTCF binding and function, or (ii) distinguishing direct from indirect effects that can be a confounding issue for interpreting site specific impact. Instead, information related to CTCF's direct effects on individual loci has come from the genetic ablation of CTCF binding sites at specific regions in the genome in a given cell type. However, interrogating the general mechanisms underlying the crosstalk between CTCF function and the cell- and locus-specific context of its binding by genetic manipulation of individual CTCF binding sites is laborious and has the limitation of providing insight into the control of only a limited number of loci in the neighboring region.

The *CTCF* locus, located on chromosome 16q band 22, corresponds to one of the smallest regions of overlap for common deletions in breast and prostate cancers[5,6]. Moreover, point mutations and deletions have been identified in many other tumors. Together these findings indicate that CTCF acts as a tumor suppressor[7]. Additionally, genetic alterations in *CTCF* and changes in dosage are associated to varying degrees with intellectual disability and microcephaly, and thus CTCF is thought to play an important role in brain development and neurological disorders[8-10].

CTCF binds to chromatin through a subset of its eleven zinc fingers (ZFs)[11,12]. Specifically, ZFs3-7 make sequence specific contacts with a 12-15 base-pair (bp) core consensus binding site. While the crystal structure of CTCF in complex with a known DNA binding site[13] has revealed new insights into the contribution of each ZF to sequence-specific binding in an *in vitro* setting, there has been no complementary, comprehensive analysis of cancer and neurological development

disorders (NDDs) associated CTCF mutants and it is thus not known how mutations within each
ZF impact binding stability and DNA sequence specificity, and the consequences this has on
binding profile, cohesin overlap, accessibility, chromosome architecture and gene regulation.
Given this lack of information, we have no context in which to determine the functional relevance
of mutations.

Using an inducible mouse ESC complementation system, we combined genetic, imaging,
molecular, AlphaFold3 and bioinformatic analyses to examine the impact of nine, high frequency
cancer associated CTCF mutations, a subset of which are also associated with NDDs. Selected
mutations occur in eight amino acid residues of ZFs that contact the core consensus binding motif.
Compared to a complete knockout of CTCF or deletion of a CTCF binding site, the mutants offer
a more subtle and in-depth approach for teasing out (i) the relationship between WT and mutant
CTCF and chromatin, and (ii) direct or indirect site-specific effects on accessibility, TF binding,
cohesin overlap, chromatin interactivity and gene expression, since each mutation produces its
own unique effect.

Here we demonstrate that the functional impact of WT and mutant CTCF depends both on the
genomic context of bound sites and the unique binding properties of each WT and mutant CTCF
protein. Analyses of the relationship between CTCF, cohesin and accessibility reveals that
although CTCF can bind both accessible and inaccessible sites, inaccessible bound sites
predominantly have weaker signals and are less competent at blocking cohesin. Multivariate
modelling demonstrates that both CTCF and accessibility contribute independently to cohesin
binding, however CTCF signal strength has a dominant effect, linked to a stronger insulation
score. Indeed, CTCFs ability to block cohesin is linked to the residence time or 'OFF' rate of each
mutant on chromatin as determined by FRAP experiments, while occupancy or 'ON' rate is
correlated with their binding at common accessible sites, indicating that accessibility dominates
the search space. Together, the mutants highlight the fact that occupancy and residence time can
be uncoupled such that although some mutants have fewer binding sites than WT CTCF, they
have a stronger residence time at these locations which increases their ability to block cohesin
and contribute to chromatin folding. The reverse situation also occurs. Our analyses also highlight
that in contrast to CTCF-bound sites, cohesin binding at CTCF-free sites is weak and does not
track with accessibility even at enhancers and promoters, perhaps because TFs that block
cohesin have considerably shorter residence times compared to CTCF.

The unique properties of WT and mutant CTCF perturb TF binding and cancer, brain, immune
and metabolic gene expression pathways. Mutants have a more pronounced effect on
transcriptional networks than auxin inducible degradation, because of indirect effects that can be
explained by altered TF binding profiles. Indeed, indirect effects that do not overlap changes in
CTCF binding impart the most influence on rewiring chromatin looping and transcriptional
networks. Changes in each mutant's transcriptional program in turn have an impact on the cell's
ability to exit from pluripotency and differentiate. Collectively, the mutants offer a rich resource to
investigate site specific CTCF-mediated effects on chromatin folding and gene regulation, while
distinguishing cause from consequence and direct from indirect outcomes.
123
RESULTS
CTCF complementation system
To explore CTCF's relationship with chromatin and its contribution to gene regulation, we made
use of a small-molecule auxin inducible degron (AID) mouse ESC (mESC) system[14,15]. In these
cells, both endogenous CTCF alleles are tagged with AID as well as eGFP (CTCF-AID-eGFP)
and the auxin-activated ubiquitin ligase TIR1 (from *Oryza sativa*) is constitutively expressed from
the *Rosa26* locus (**Figure 1A**). Addition of indole acetic acid (IAA), an analogue of auxin leads to
rapid poly-ubiquitination and proteasomal degradation of the AID tagged protein[16].
132
To study the impact of CTCF mutations, we established a rescue system, modifying the mESC
degron cell line to express either a stable doxycycline-inducible control wild-type *Ctcf* or a mutant
Ctcf (*mCtcf*) transgene in the absence of endogenous CTCF[17]. Transgenes, knocked into the
Tigre locus, have a 3 x FLAG tag, which allows us to distinguish mutant or wild-type transgenic
CTCF from endogenous CTCF using a FLAG antibody in Western blot and ChIP-seq analysis.
An mRuby fluorescent tag enables us to accurately determine expression levels of mutant or wild-
type transgenic CTCF by flow cytometry (**Figure 1A**). The three conditions used for our analysis
are shown in **Figure 1B**, and an example of the FACS profile for cells expressing a WT CTCF
transgene in each condition is shown (**Figure 1C**).
142
We selected a subset of missense mutations (identified from cBioPortal [18] and COSMIC[19]) that
occur at high frequency in cancer patients (**Figure S1**), focusing on those found in ZFs that
contact the 12-15bp consensus CTCF binding motif (**Figure 1D** and highlighted with an asterisk
in **Figure S2**). We analyzed mutations in **(i)** amino acids that make base-specifying contacts with
the alpha helix of DNA (at locations -2, 2, 3 and 6 on the ZF), **(ii)** residues that coordinate the zinc

ion which are essential for providing stability to ensure the proper folding of the ZF domain, **(iii)**
two other classes of amino acid residues in the ZFs: boundary residues (that are important for
interactions between ZFs), and residues that contact the sugar phosphate backbone of DNA, and
(iv) other highly mutated residues in the region. Each class of mutation is color-coded as shown
in **Figure 1D** (left) and **Figures S1-2** and the color code is maintained throughout the figures.
153
In total, we analyzed mutations in eight amino acid residues and for R377, the most highly mutated
residue in cancer, we included two distinct mutations of R377, both of which (R377H and R377C)
are equally represented in patients (**Figure S2**). Selected mutations (R339Q, R342C, R448Q,
R377H and R377C) are also implicated in NDDs (**Figure S2**)[10,20]. Individual clones with
comparable levels of transgene expression (mutant and WT) were selected based on flow
cytometry and Western blot analysis (**Figure 1E, F and Figures S3-4**). To assess rescue of
endogenous CTCF by WT transgenic CTCF, we performed ChIP-seq using an anti-CTCF
antibody and RNA-seq in WT untreated and WT transgene expressing (ID) cells. This analysis
showed that about half of the overall CTCF binding sites (but 97% of the CTCF motif-containing
sites) and gene expression changes are rescued by transgenic CTCF (**Figure S5**), consistent
with the lower expression of the transgene reported previously[17]. This data provides a rationale
for comparing the effect of the mutant and WT transgenic CTCF.
166
Retained binding sites are enriched in accessible regions with strong CTCF signals
We next analyzed the profiles of mutant versus WT CTCF using cells expressing transgenic WT
and mutant CTCF in the absence of endogenous CTCF (ID). Briefly, we degraded endogenous
CTCF-mAID-GFP by IAA addition, induced WT or mutant transgene expression with Dox for 48
hours, and then performed FLAG ChIP-seq. For each mutation we identified WT only sites (bound
by WT CTCF that mutants no longer bind), common sites (bound by both WT and mutant proteins)
and mutant only sites (*de novo* sites that only mutant proteins bind)**.**
174
The location of each mutation and the complementary DNA triplet that the relevant ZF binds is
marked by a box in the consensus motif shown in **Figure 2A** and **Figure S6A**. Below that we
highlight the most frequent motif identified by MEME[21] for the set of *de novo* sites**.** These motifs
differ from that of the consensus sequence by either (i) a diminished requirement for DNA bases
that contact the mutated zinc finger (**Figure 2A**), or (ii) an alteration to the sequence of the
consensus motif, which could be explained by the impact of mutations on DNA-base interacting
zinc finger residues. For example, K365 binds to the second guanine in the triplet GGC of the

consensus motif and the *de novo* motif found in the K365T mutant is consistent with loss of

hydrogen bonds with the central guanine due to the threonine substitution, as previously

demonstrated[13]. Motifs found at WT only and common sites were similar to the consensus motif

(**Figure S6B**), with the exception of R377C for which few WT only sites were observed. On

average, the percentages of CTCF containing motifs at common sites were higher than at WT

only sites, suggesting that the latter might be lower affinity sites with degenerate motifs, consistent

with the weaker CTCF signal observed. Each mutant displays their own unique binding profiles

including the two distinct mutations of residue R377. R377C binds to less common and more *de*

novo sites compared to R377H and each mutant binds a motif that is similar to the consensus

motif but lacking bases associated with ZF3 (**Figure 2A** and **Figure S6**).

The bar graphs show the percentage of WT only, common and mutant only binding sites, while

the heatmaps reveal the number of binding sites for each subset and highlight differences in

CTCF ChIP-seq signal between WT and mutants at the three locations. Differential binding

analysis using DiffBind (**Figure S7)** showed that only a fraction of the mutant and WT only peaks

reached statistical significance, which can partly be explained by the fact that these peaks are

weaker and have a smaller effect size compared to the stronger, common peaks. However, the

ratios of gained versus lost CTCF binding sites are consistent with the proportion of 'WT only' and

'mutant only' peaks (**Figure 2A** and **S6A**). CTCF signal strength is strongest at common sites,

however, CTCF this is altered in a mutant specific manner, suggesting that a 'binding dosage

effect' contributes to their functional impact. As expected, the H455R mutation in the zinc

coordinating residue of ZF7 loses the most binding sites. OmniATAC[22] profiles for the different

binding groups indicate that CTCF ChIP-seq signal strength is a function of chromatin

accessibility, with common sites being the most accessible (**Figure 2A and Figure S6A**).

In sum, ChIP-seq analysis reveals that CTCF mutants lose, retain and gain a subset of binding

sites throughout the genome with mutant specific profiles. Loss and gain of CTCF binding occurs

predominantly at weak inaccessible CTCF sites, while binding is retained at the stronger common

binding sites which have high accessibility, although CTCF ChIP-seq signal and accessibility can

vary depending on the mutant.

CTCF reduces ATAC-seq signal in a cohesin-dependent manner

To further investigate the link between accessibility and CTCF binding, we compared the ATAC-

seq signal at CTCF-bound versus unbound sites in the presence or absence of SMC3, a subunit

of the cohesin complex (**Figure 2B** and **Figure S8**). Accessibility was reduced in an SMC3-
dependent manner at both WT and mutant CTCF-bound sites. Each CTCF mutant exhibited a
differential ability to decrease the height and breadth of ATAC-seq peaks, suggesting that CTCF
is driving the change in accessibility and that the relationship between CTCF and accessibility is
bidirectional. We do not observe the same phenomenon at sites without SMC3, which lends
support to the idea that loop formation decreases accessibility, indicating a unique chromatin
structure at loop bases. The same phenomenon is observed at enhancer and promoter sites,
although the effect is weaker, particularly at promoters (**Figure S9**). The weakened effect may be
due to interference of TF binding which can affect CTCF-dependent nucleosome phasing and
insulation at CTCF bound sites or competitive binding with CTCF[23]. These results highlight the
usefulness of the mutants in distinguishing cause from effect.
227
Mutations reduce the chromatin residence time and bound fraction of CTCF
Fluorescence recovery after photobleaching (FRAP) was performed on the 9 mutants and WT
CTCF[24,25], using the same auxin and Dox conditions as described for the ChIP-seq and ATAC-
seq. For the FRAP a circular 1 µm diameter circle was bleached, and recovery monitored over 10
minutes, recording 30 movies per condition in three replicates. While the recovery of WT CTCF
largely matched prior CTCF FRAP results[24], all mutants exhibited faster recovery consistent with
reduced and/or less stable DNA-binding (**Figure 3A and S10**). We fit the FRAP curves to a
reaction-dominant kinetic model [24,25] and estimated the residence time (duration of CTCF binding
to DNA) and bound fraction (proportion of total CTCF bound to DNA) of each mutant with
subsequent comparison to the WT (**Figure 3B, C**). We found a general reduction in the residence
time and/or bound fraction across each mutant, with WT CTCF exhibiting the largest specific
bound fraction and residence time (**Figure 3B, C**). Each mutant displayed unique parameters,
regardless of the mutant category. The R377H and R377C mutants – mutations in the phosphate
contacts – have similar bound fractions, but distinct residence times. The mutations in the amino
acids making direct contact with DNA (R339Q, K365T, Q418R, and R448Q) have varied
residence times and bound fractions, despite belonging to the same group. R448Q showed the
second lowest bound fraction, despite exhibiting a residence time comparable to WT. Q418R had
the lowest residence time of all mutants examined, with an intermediate bound fraction.
Conversely, K365T exhibited a bound fraction and residence time most closely comparable to
WT. H455R, which assists in coordinating the zinc, had the lowest bound fraction and the second
lowest residence time of all examined mutants, accurately reflecting the most deleterious mutation
to canonical CTCF function. R342C, which occurs in a boundary residue between the two zinc-

ligand histidine residues of ZF3, showed a residence time similar to that of wild type, but had a lower bound fraction, possibly suggesting a role for boundary residues in binding efficiency. Lastly, S354F, located between the two zinc-ligand cysteine residues of ZF4, which substitutes a bulky phenylalanine in place of a serine in ZF4, showed both a moderate decrease in bound fraction as well as residence time.

In summary, our FRAP results demonstrate that all mutants show changes in DNA binding either through an altered residence time or bound fraction. Surprisingly, some mutations strongly affect either the residence time or the bound fraction, with minimal or less of an effect on the other. This suggests that the CTCF target search (ON-rate) can be decoupled from binding stability (OFF rate) [26], such that some mutants affect the efficiency of the search for binding sites, without affecting the residence time of CTCF.

The impact of mutations on the DNA-CTCF complex

To assess the impact of each mutation on the DNA-CTCF complex, we generated predictions using AlphaFold3. First, we compared the WT and K365T mutant DNA-CTCF complexes between the predicted and experimental structure and found a high concordance (**Figure S11**). We then predicted the DNA-CTCF (ZF1-11) structure for each mutant and compared the alignment against WT (**Figure S12**). Like K365T, the mutations have little effect on the overall conformation of CTCF. However, Q418R the mutant with the lowest residence time has the highest deviation from WT (RMSD=1.6 Å), although the RMSD is still <2Å, which is considered a good alignment with the WT structure. It is of note that AlphaFold3 does not take account of the effect of ions, therefore, the prediction for H455R might not be accurate since the mutation disrupts the coordination of H455 with the zinc ion. Some of the mutations found at residues making direct contact with the DNA, have lost or weakened hydrogen bonds (K365T, R339Q and R448Q). Q418R gains hydrogen bonds (**Figure S12**), which could explain the paradox of this mutation: that it has the lowest residence time but strong binding at common sites.

CTCF mutations have a graded impact on binding and function

To understand the relationship between FRAP and ChIP-seq data, we asked whether the residence time is linked to CTCF's binding stability and ability to block cohesin. In **Figure S13A** we show the stratified overlap of cohesin/FLAG at WT only, common and mutant only CTCF sites. Overlap with cohesin is lower at WT only and mutant only sites, consistent with their weaker CTCF and ATAC-seq signals. These changes do not reflect changes in cohesin protein levels as

Western blots demonstrate that CTCF mutants do not impact the levels of RAD21 or SMC1, two
subcomponents of the cohesin complex (**Figure S13B**). The proportion of global CTCF-cohesin
versus CTCF only binding sites shown in the heatmaps (**Figure 3D**), demonstrate that CTCF-
cohesin overlap is graded across mutants, to some extent mirroring a gene dosage effect that
could occur as a consequence of altered residence time (**Figure 3C**). Indeed, analysis of SMC3
ChIP-seq revealed that the residence time was strongly associated (R^2=0.8, p=0.002) with the
percentage of CTCF-cohesin overlap for each mutant (**Figure 3E**) which reflects the impact of
binding stability on CTCF's function in loop extrusion. Conversely, the proportion of SMC3
overlapping CTCF might reflect changes in the number of CTCF peaks across the mutants since
the expression level of components of the cohesin complex, SMC1 and RAD 21 do not change
in CTCF mutant expressing cells. Indeed, **Figure S14A** shows a strong correlation between the
percentage of SMC3 peaks overlapping CTCF and the number of CTCF peaks (**Figure S14B**).
296
We next compared the specific bound fraction estimated by FRAP, to the bound fraction using
FLAG ChIP-seq and ATAC-seq peaks to define all potential binding sites. While we did not find a
good correlation when assessing the overall CTCF ChIP-seq bound fraction (**Figure S14C**), there
was a significant correlation (R^2=0.6, p=0.008) between the chromatin bound fraction of the
mutants and the fraction of mutant binding at common CTCF sites (**Figure 3F**). These data
suggest that the FRAP bound fraction mostly captures the effect of the mutations on the strong,
accessible binding sites. Furthermore, we found a strong correlation with the fraction bound and
the overall ability of CTCF-cohesin overlapping sites to reduce accessibility, (R^2=0.7, p=0.003),
in line with CTCF functioning in this context in a dose dependent manner (**Figure 3G**).
306
The relationship between cohesin and accessibility at CTCF-bound and CTCF-free sites
Consistent with previous findings[27], our data links accessibility with CTCF binding signal strength.
Furthermore, the FRAP data support an association between CTCF residence time and cohesin
overlap, which is also supported by simulation analysis[28]. To separate out the contribution of
CTCF and ATAC strength on SMC3 signals, we performed multivariate modelling to determine (i)
the effect of CTCF strength on SMC3 overlap, adjusted for the effect of ATAC signal, and (ii) the
effect of ATAC strength on SMC3 overlap, adjusted for the effect of CTCF signal. **Figure 4A** (top)
shows that SMC3 enrichment independent of ATAC-seq signal, is strongly associated with CTCF
signal strength in a dose-dependent manner. Only strong accessibility significantly influences
SMC3 binding independent of CTCF, however, accessibility has much less contribution compared
to CTCF binding stability **(Figure 4A**, bottom). Thus, CTCF signal strength plays a dominant role

in determining the probability of CTCF-cohesin overlap, consistent with the FRAP and ChIP-seq

data (**Figure 3E**). The heatmap in **Figure 4B** and **Table S1**, further shows that, in contrast to

CTCF-bound sites, the percentage of SMC3 overlap at CTCF-free sites is increased with

increasing accessibility, however signal strength at highly accessible sites is much lower than

sites overlapping even weak CTCF-bound sites. The same effect is observed for the mutants, but

in some cases SMC3 overlap at CTCF-bound sites is weaker than WT CTCF, in line with their

reduced residence times.

Overall SMC3 profiles at CTCF-bound sites confirmed that CTCF is most effective at blocking

cohesin at accessible sites enriched with strong CTCF signals, and further highlight that each

mutant CTCF performs this function with variable ability depending on their residence time

(**Figure 3E, 4C and Figure S15**). A comparison of SMC3 profiles at CTCF bound versus unbound

sites, demonstrates that other factors such as TFs, contribute very little compared to CTCF likely

because residence times of TFs on chromatin are much shorter than CTCF's[29]. Indeed, at CTCF-

free sites, the SMC3 profiles are similar at CTCF-free accessible and inaccessible sites,

suggesting that accessibility has little contribution to cohesin binding in the absence of CTCF. A

similar trend was observed at enhancers and promoters where accessibility is higher and TFs are

known to be enriched (**Figure S16**). These data suggest that for a given cell type, the subset of

strong and accessible CTCF binding sites are predominantly involved in loop extrusion. However,

weaker, inaccessible CTCF binding sites might also have a role in more transient, dynamic

looping events with a reduced loop lifetime[30,31]. Moreover, a subset of inaccessible sites (16% in

the WT) have strong CTCF signals.

Each CTCF mutation alters gene expression and TF binding in a unique manner.

To determine how CTCF mutations impact gene expression we performed RNA-seq and did an

unsupervised clustering analysis, comparing gene expression in cells in which endogenous CTCF

was degraded in the absence (IAA) or presence of the WT or mutant CTCF transgenes (ID).

Changes in transcriptional output were determined by comparison with cells expressing WT

transgenic CTCF (ID) (**Figure 5A**). Overall, 6474 genes had altered gene expression across all

mutants including the IAA condition (**Table S2**). **Figure S17A** shows that the majority of

differentially expressed genes (DEGs) are found only in one mutant.

Gene set enrichment analysis using the KEGG database[32] showed a strong enrichment in

functional pathways related to cancer, brain, immune and metabolic processes among DEGs

across mutants (color coded in **Figure 5B**, with genes outside these pathways represented in
black**)**. This finding is compatible with the human diseases associated with CTCF mutations. The
top DEGs of these pathways are shown in the bar graphs below. Examples in **Figure 5B** show
changes in pathway and gene expression in the absence of CTCF (IAA condition) and in cells
expressing the mutant CTCF transgenes, H455R, S354F, R448Q and R339Q (ID condition). All
other mutants are shown in **Figure S17B**.

We also observed an enrichment of dysregulated genes observed in patients with CTCF
associated neurodevelopmental disorders (NDDs[9]) (**Figure S17C and Table S2**). CTCF plays an
important role in imprinting, and alterations in imprinted genes could contribute to the CTCF-
dependent neurodevelopmental phenotype as suggested by congenital imprinting disorders[33].
Indeed, we found a strong enrichment of imprinted genes across all mutants (**Figure S17D and**
Table S2).

Together, our analyses demonstrate that although each mutant exhibits a unique impact on
binding and gene expression, all mutants affect common pathways, consistent with phenotypes
linked to human CTCF-associated disorders. Differences in the impact of each mutant provide
insight into CTCF-associated NDD inter-patient variability. Interestingly, cells in which
endogenous CTCF is degraded (IAA condition) show fewer gene expression changes compared
to mutant expressing cells, except for the R377C mutation (**Figure S17B**). Some mutants have
alterations in many pathways (R399Q), while others highlight changes in only a few (Q418R and
R377C) as shown in **Figure 5B and Figure S17B**, suggesting that in the latter, the alteration in
gene expression may not accumulate in specific pathways.

Mutant specific alterations in transcriptional networks impact exit from pluripotency
To investigate, whether mutant specific transcriptional networks could potentially affect exit from
pluripotency and the ability to differentiate we compared expression profiles of WT and a selection
of mutant CTCF expressing cells before and after 72 hours of LIF withdrawal[34]. The mutants
analyzed were selected because they cover a range of CTCF residence times as measured by
FRAP. R342C exhibits a similar residence time to WT, K365T an intermediate residence time and
Q418R has one of the lowest residence times. RNA-seq revealed that each mutant demonstrated
a distinct differentiation trajectory compared to WT cells, which are most skewed towards a
neuroectodermal fate (**Figure 5C** and **Table S3**). A subset of the DEGs is shown in **Figure S18**.

In the mutants, we identified several differentially expressed imprinted genes , such as H19 and
PEG10, as well as ZFP57, which plays an important role in regulating imprinting. To further assess
the biological relevance of the differentiation behavior, we annotated DEGs using the DECIPHER
developmental disorder database (https://www.deciphergenomics.org/ddd) (**Table S3**). We found
44 genes associated with developmental disorders, including NDDs such as *Camk2a*, *Cnksr2*,
Sobp that are involved in intellectual disability, and *Dchs1* which is responsible for periventricular
neuronal heterotopia. In addition, we detected genes responsible for congenital connective tissue
syndromes (*Fras1* in Fraser syndrome and *Fbn1* in Marfan syndrome) as well as genes involved
in skeletal dysplasia (*Sox9*) and congenital cardiomyopathy (*Speg*). While CTCF-related
congenital disorders are most commonly characterized by neurodevelopmental symptoms,
patients can also exhibit other congenital anomalies such as heart defects, craniofacial
malformations or cleft palate[10], suggesting the biological relevance of the changes in
differentiation behavior observed in our system.

Transcriptional changes correspond to predicted alterations in TF binding at promoters

To investigate whether mutant specific transcriptional changes could be explained by altered TF
binding, we performed a footprinting analysis of ATAC-seq data using the TOBIAS pipeline [35] in
cells which have endogenous CTCF degraded in the absence (IAA) or presence of WT or mutant
transgenes (ID). For this analysis replicates were merged and comparisons made with WT ID
cells. Overall, 195 TFs had predicted differential binding across all mutants including the IAA
condition. As a proof of principle, we detected predicted differential binding for CTCF, which has
altered binding in the mutants. Numerous other predicted differentially bound TFs (n = 194) were
identified as shown in **Figure 5D, S19A** and **Table S4.** About one third were found in only one
mutant (**Figure S19B and Table S4**). Annotation of differentially bound TFs revealed a significant
enrichment for TFs associated with NDDs (Odd Ratio =2.2, p=0.02) (**Table S4**).

We overlapped predicted differentially bound TFs with a region of 2kb around the promoters of
DEGs in cells expressing different CTCF mutants, to analyze the impact of predicted differential
binding of TFs on gene expression and observed a strong enrichment (**Figure 5E, Figure S20**
and **Table S4**). Differentially expressed target genes were also significantly enriched in genes
associated with developmental disorders (OR=3.6, $p=7 \times 10^{-36}$) and in differentially expressed
genes found in CTCF associated NDDs (OR=4, $p=10^{-63}$). It is of note, that the majority of the
predicted changes in TF binding at the promoters of DEGs do not overlap CTCF ChIP-seq peaks
or predicted differential CTCF binding. This data suggests that many of the mutant-mediated

transcriptional changes could be explained by indirect effects resulting from the disruption of
specific TF pathways.

Figure 5E and Figure S20 highlight the impact of two of the top predicted differentially bound
TFs (CTCF and MBD2) at promoters of DEGs in different mutants. These are shown in volcano
plots for cells in which endogenous CTCF is degraded (IAA) and in cells expressing different
CTCF mutations (ID). MBD2 is a methyl-sensitive factor that belongs to the NurD chromatin
remodeling complex, that has been implicated in cancer and NDDS. NurD is also required for exit
from pluripotency and lineage commitment[36-39]. For each CTCF mutant, predicted differentially
bound TFs target the promoters of a distinct set of up (red) and down (blue) regulated genes.
Both CTCF and MBD2 exhibit an overall activator profile with a positive association between
binding and expression (**Figure 5E and Figure S20**). However, for some mutants, a weaker but
significant negative association was also observed suggesting that both factors can act as
activators or repressors depending on the cell-specific genomic context, consistent with previous
reports[38,40].

Examples of altered CTCF binding at the promoter of differentially expressed *Rerg* and predicted
differentially bound MYC at the promoter of differentially expressed *Brdt*, are shown in **Figure 5F.**
Loss of CTCF binding at the promoter of the *Rerg* gene leads to a decrease in its expression,
while expression of *Brdt* increases in line with increased predicted binding of MYC at its promoter.
The latter is an example of an indirect effect because CTCF is not bound at the promoter of this
gene. In sum, combined analysis of RNA-seq and ATAC-seq highlight the variable effects of each
CTCF mutation, connecting gene expression changes with 195 predicted differentially bound TFs,
to provide a deeper understanding of the factors that underlie the unique transcriptomic profile of
each CTCF mutation and their impact on developmental processes particularly relevant to brain
disorders.

Chromatin interactivity and insulation are linked to mutant binding properties
Hi-C demonstrated that each mutant has a distinct interaction profile and exhibits (i) variable loss
of intra TAD interactions and boundary strength as well as (ii) aggregated loop strength. The IAA
condition, in which CTCF is degraded shows a loss of intra TAD interactions and a gain of inter
TAD interactions as well as the most reduction in aggregated loop strength (**Figure 6A**), while
mutants display their own unique effects. No major compartment changes were observed as
shown by the interaction matrices at the chromosome level (**Figure S21A**).

Consistent with our finding that cohesin overlap is correlated with CTCF mutant residence time
as determined by FRAP analysis (**Figure 3E**), we found that residence time is also associated
with chromatin interactivity and insulation score (**Figure 6B, C**).The correlation with the latter was
modest and did not reach significance, probably because of the lower Hi-C resolution (10 kb)
compared to ChIP-seq. The loop extrusion length (P'max[41]) for each mutant is associated with
the fraction that is bound to chromatin (**Figure 6D**), with the longest loops associated with mutants
that have the lowest bound fraction, indicating a lower density of CTCF-cohesin anchors in these
mutants (**Figure 3B**). The variability in the interaction profile of each mutant can be clearly seen
across a 10Mb region of chromosome (**Figure 6E and Figure S21B**).

To further examine the impact of each mutant on chromatin organization, we first compared the
aggregated insulation score at lost sites upon CTCF degradation (comparison of IAA versus WT-
ID condition) (**Figure S21C**) using a 2Mb window centered on CTCF peak summits. The insulation
scores were generated from Hi-C data at a 10kb resolution. This analysis confirmed that the
mutants partially retain insulation at lost sites depending on their binding stability. We then
analyzed the insulation score at WT only, common and mutant only binding sites (**Figure 6F and**
Figure S22A). This analysis revealed that the strong, common accessible binding sites with the
highest cohesin overlap, have the highest insulation compared to WT only and mutant only sites.
At the WT only sites (comparison of WT-ID versus mutant-ID), some mutants exhibit weak
insulation which is not detected when CTCF is degraded (**Figure 6F, Figure S22A** and **Figure**
S22B), suggesting there might be weak mutant CTCF binding (non-significantly detected by ChIP-
seq) at some of these sites. Whether insulation scores at WT only and mutant only sites are lower
or higher than each other depends on the residence time and cohesin overlap of each mutant, in
line with what we observed in **Figure 3C, E and 6C**. These studies further demonstrate that
imaging and molecular analyses can be functionally integrated to provide new insight into mutant
specific effects.

Since WT and mutant only sites are associated with weaker CTCF peaks and lower accessibility,
we examined the insulation score at CTCF binding sites, stratified by CTCF and ATAC-seq
strength. The profiles in **Figure 6G, 6H, S22C and S22D** indicate that stronger CTCF signals are
associated with stronger insulation, while accessibility has a more modest impact. To quantify
their independent effects, we performed a mixed multivariate model, the results of which are
consistent with the analysis of SMC3 overlap shown in **Figure 4A**.

Changes in gene expression are linked to changes in chromatin interactivity

To determine if changes in gene expression are linked to altered chromatin folding, we analyzed interactions involving the promoters of genes that were differentially expressed and found that the promoters of overexpressed genes were enriched in gained chromatin loops (**Figure 7A**). Enrichment was not observed among the under-expressed genes, suggesting that under-expression observed in mESC expressing mutant CTCF proteins might be an indirect effect, while overexpression might reflect a direct CTCF-mediated effect on loop extrusion. This is supported by the sub-analysis of overexpressed and underexpressed genes shown in **Figure S23,** focusing on loops with both anchors overlapping a CTCF binding site, which show a similar result. These loops are more likely to capture the direct effect of CTCF on loop extrusion. This is shown in examples in **Figure 7B** and **Figure S24A** which depict gain of interactions from CTCF binding sites toward the promoters of overexpressed genes, including *Msh6*, *Epcam*, *Foxn2* and *Cox7a2l* (**Figure 7B** and **Figure S24B)**. Interestingly, all these genes are implicated in tumorigenesis[42-48]

Most of the change in expression are the result of indirect effects.

Our data show that most CTCF mutations have more impact on gene expression than the IAA condition. Indeed, CTCF degradation has been shown to induce limited gene expression changes [14,15]. To ask whether this could be explained by indirect effects which depend on cell-specific transcriptomes, we estimated the direct versus indirect effects of CTCF mutations on gene expression. Since CTCF can act as a traditional transcription and chromatin regulator [40], we defined indirect CTCF effects as (i) differentially expressed genes (DEGs) with a predicted differentially bound TF at the promoters that do not overlap a CTCF ChIP-seq peak or a predicted differentially bound CTCF site, and (ii) DEGs with differential looping of the promoter without CTCF binding at either loop anchor. Conversely, for direct effects, we required (i) the presence of a CTCF peak and/or a predicted differentially bound CTCF site at the DEG promoter or (ii) loops anchoring the DEG promoter with at least one loop anchor overlapping a CTCF binding site.

Figure 7C shows that indirect effects account for approximately twice as many DEGs compared to the direct effects, except for Q418R. This mutant has a short residence time but a high number of peaks, supporting the hypothesis that unstable peaks might be involved in dynamic and functional looping. Furthermore, a higher percentage of DEGs can be explained by changes in TF or CTCF binding at gene promoters, rather than by changes in chromatin interactivity in all the mutants. For the IAA condition, the contribution of differential loops to expression changes is greater. A similar result was observed when segregating up- and down-regulated genes but with

a higher proportion of direct versus indirect effects in the IAA condition for down-regulated genes

consistent with loss of loops upon CTCF degradation (**Figure S24B**). These findings suggest that

although CTCF mutations induce changes in chromatin looping, the primary mechanism driving

alterations in gene expression is loop-independent and mediated by changes in transcription

factor binding.

DISCUSSION

Using a combination of genetic, imaging, molecular, AlphaFold3 and bioinformatic approaches

we have examined the impact of nine, high frequency cancer associated CTCF mutations in ZFs

that make contact with the core consensus binding motif. Our studies reveal that the specificity of

interaction between the ZFs in CTCFs binding domain and its target DNA sequence is a key

determinant of attachment. However, if the same sequence is found in both accessible and

inaccessible chromatin, CTCF will preferentially bind accessible sites as highlighted by the fact

that mutants bind common accessible sites in preference to less accessible WT only sites. This

is underscored by the strong correlation between the fraction of chromatin bound mutants

detected by FRAP, and mutant binding at common, accessible CTCF sites. These findings are

consistent with studies from Dirk Schubeler's group showing that CTCF can bind both accessible

and inaccessible sites, but that the latter are less likely to be involved in insulation [27,49]. We also

observed a good correlation between each mutant's residence time and CTCF-cohesin overlap,

indicating that binding stability is important for blocking cohesin. A similar, but more modest

correlation was found between residence time and chromatin interactivity as well as insulation

score, linking binding stability with cohesin overlap as well as loop and boundary formation.

By dissecting out the contribution of CTCF signal strength and accessibility to SMC3 overlap, we

were able to reveal the dominant effect of CTCF. SMC3 overlap at accessible and inaccessible

CTCF-bound sites is stronger than overlap at accessible, CTCF-free sites, including enhancers

and promoters, likely because TFs have a short residence time compared to CTCF [29]. Our data

also demonstrate that strong accessible CTCF bound sites are associated with higher insulation,

consistent with higher SMC3 stalling and more stable loops.

In principle, these features could have been uncovered by analyzing WT CTCF in the context of

cohesin overlap and accessibility. However, collectively the graded effects of the mutants act as

perturbations that lend support to our models, underscoring how different aspects of CTCFs

binding properties contribute. Moreover, they reveal that each mutant's search for binding sites

can be decoupled from its chromatin bound residence time, as previously demonstrated for a
mutant form of CTCF with ZF8 deleted [50]. For example, the Q418R mutant binds an intermediate
number of sites but at these sites it is has a very low residence time and is less competent at
blocking cohesin. In contrast, R448Q binds fewer sites with a similar residence time compared to
WT CTCF, and it is slightly better at blocking cohesin than WT protein. The low chromatin bound
fractions of H455R and R448Q, which have very different chromatin residence times, are
correlated to an increased loop length of both mutants (P'-max), which can be explained by the
unobstructed passage of cohesin past what would normally be a bound CTCF site.

The crosstalk between accessibility and CTCF binding goes in both directions, such that
accessibility affects binding and binding affects accessibility. Binding at accessible sites, narrows
and reduces the ATAC-seq peak. Evidence for this being a CTCF driven effect comes from
analysis of the CTCF mutants. Those with low residence times (H455R and Q418R) are unable
to function in this capacity, while the R342C mutant has a stronger effect compared to WT CTCF.
We speculate that these differences reflect decreased accessibility at the bases of CTCF-SMC3-
mediated loops, because in the absence of SMC3, where loops cannot form there is no reduction
in accessibility.

While loss or gain of CTCF binding at promoters can account for direct mutant specific effects on
accessibility and gene expression, we could only demonstrate a direct effect of gained CTCF
binding associated with gained loops involving the promoters of over-expressed genes. Under-
expression could mostly reflect indirect effects of CTCF binding alterations. Indeed, all mutants
have their own unique indirect impact in globally changing accessibility at sites where CTCF is
not bound. This effect links changes in TF footprinting with altered transcriptional output,
explaining most of the mutant specific effects that we observed. We also identified a higher
proportion of DEGs mediated by changes in TF or CTCF binding at the gene promoters than by
changes in chromatin looping, suggesting that the primary mechanism driving alterations in gene
expression might be loop independent. Indeed, the impact of each mutant on TF expression and
altered TF binding at the promoters of differentially expressed genes is less likely to give rise to
detectable changes in interactivity by Hi-C, consistent with our finding that SMC3 signals at CTCF-
free sites are far weaker than at CTCF-bound sites, and thus likely to be involved in weaker, more
dynamic loops.

Aside from the R377C mutant, we observed fewer changes in gene expression in IAA treated
cells compared to CTCF mutant expressing cells, which can be explained by the predominant
contribution of indirect changes mediated by TF binding on the mutant-specific transcriptome.
Although the mutants all uniquely affect gene expression pathways, we detected enrichment in
pathways affecting the brain, immune system, cancer and metabolism, which is consistent with
the clinical setting in which the mutations are found, namely cancer and brain disorders. The
added effect on immune and metabolic pathways provides insight into other changes that could
occur in CTCF mutant-mediated diseases.
597
CTCF mutations also affect the trajectory of exit from pluripotency. Indeed, while both WT and
CTCF mutant cells tend to acquire a neuroectodermal expression profile upon LIF withdrawal, we
observed mutant-specific transcriptional signatures of other germ layers, suggesting a more
heterogenous differentiation trajectory, which could be confirmed in future work by scRNA-seq.
This finding provides a potential mechanism for CTCF mutation-associated neurodevelopmental
disorders. Indeed, while CTCF has been involved in neuro-progenitor maintenance and survival[51],
our analyses suggest that CTCF mutations might also alter brain development by affecting germ
layer commitment.
606
607 Taken together the binding domain mutants we have analyzed here, provide a new appreciation
608 of CTCF's bidirectional relationship with chromatin, its ability to bind and function in different
609 contexts as well as the potential impact of each mutation in clinical settings. Furthermore, our
610 analyses provide a better understanding of how any genetic or epigenetic disorder that alters the
611 landscape of chromatin can in turn impact CTCF binding and function.
612
613 **LIMITATIONS OF THE STUDY**
614
In our complementation system, the expression of the transgene is lower than the endogenous
CTCF. Since CTCF binding and function is dose-dependent, our analyses of CTCF binding sites
might be biased toward higher affinity sites. To distinguish the binding of mutant transgene from
residual endogenous WT CTCF in the ID condition, we performed ChIP-seq using a FLAG
antibody which resulted in multiple non-specific binding sites. To address this issue, we used a
stringent peak calling resulting in a lower number of CTCF binding sites compared to previous
reports[15].
622

In this study, we found decreased accessibility at CTCF-SMC3 sites and hypothesize a link with
the loop extrusion process. However, despite the evidence from the mutant perturbation, the
reduction in overall accessibility at these sites could reflect a preference for CTCF binding
involved in insulation at less open chromatin regions, which we cannot entirely rule out.

We found that most of the differences in transcriptomic profiles observed in CTCF mutant cells
were the result of indirect and loop-independent effects. However, our Hi-C data has a resolution
of 10 kb, which could lead to an underestimation of CTCF loop-dependent expression changes.

RESOURCE AVAILABILITY
Lead contact
Further information and requests for resources should be directed to and will be fulfilled by the
lead contact, Jane Skok (Jane.Skok@nyulangone.org).

Materials availability
The WT and mutant CTCF mESC lines are available upon reasonable request.

Data and code availability
All raw and processed sequencing data files will be deposited at NCBI's Gene Expression
Omnibus (GEO) upon publication. FRAP analysis was performed using custom Python scripts
available   at   https://github.com/ahansenlab/ZNF143_analysis_code/tree/main   and
https://zenodo.org/doi/10.5281/zenodo.14056922 [52]. No other custom code/software was used in
the study. The publicly available software used is indicated in the Method details.

Acknowledgements
This work was supported by 1R35GM122515 (J.S) and NIH P01CA229086 (J.S). A.T. was
supported by the American Cancer Society (RSG-15-189-01-RMC) and St. Baldrick's foundation
(581357). ASH additionally acknowledges support from US National Institutes of Health grants
R00GM130896, DP2GM140938, R33CA257878 and UM1HG011536, National Science
Foundation grant 2036037, the and a Pew-Stewart Cancer Research Scholar grant. GC and GJ
were supported by fellowships from the NCC.

The authors thank Skok lab members for helpful scientific discussions, New York University
School of Medicine High Performance Computing Facility (HPCF) for computing technical

support, Adriana Heguy and the Genome Technology Center (GTC) core for sequencing efforts,
NYU Flow Cytometry and Cell Sorting Center for FACS analysis and sorting. GTC is a shared
resource partially supported by the Cancer Center Support Grant P30CA016087 at the Laura and
Isaac Perlmutter Cancer Center.

Author contributions

These studies were designed by J.A.S., C.D. and M.N. The analysis was performed by C.D and
T.S. G.J, P.L and F.F.D.R. established the CTCF mutant cell lines with the guidance of E.P.N.
Sequencing libraries were performed by G.J., G.C., R.V., N.K. A.T. reviewed the manuscript.
FRAP analysis was performed by C.C.K, D.N.N and A.S.H. The paper was written by J.A.S and
C.D.

Declaration of interests

The authors declare no competing interests.

MAIN FIGURE TITLES AND LEGENDS

Figure 1: **CTCF complementation system**. (**A)** Scheme of the CTCF doxycycline-inducible
degron system. (**B**) Experimental strategy for the expression of WT and mutant transgenic CTCF.
(**C**) Flow cytometry showing the level of GFP (endogenous CTCF) and mRuby (transgenic WT
CTCF). (**D**) Left, scheme showing the locations of the different types of CTCF mutations within a
ZF. Amino acids making contacts with the DNA are shown in shades of pink, residues that
coordinate the zinc ion in red, boundary residues in purple, and residues that contact the sugar
phosphate backbone of DNA in blue. Right, representation of CTCF showing the locations of each
mutation under investigation. (**E**) Bar graph showing the expression levels (mean fluorescence
intensity) of endogenous CTCF (GFP) and transgenic WT or mutant CTCF (mRuby2) after ID
treatment. The error bars represent the standard deviation between the two replicates. (**F**)
Western blot of endogenous CTCF (CTCF antibody) and transgenic CTCF (FLAG antibody) in
untreated (WT-U), IAA, ID conditions. See also **Figures S1-5**.

Figure 2: **CTCF mutations have unique chromatin binding profiles**. (**A)** Scheme showing the
locations of the CTCF mutations within the ZF. The consensus CTCF motif highlights the triplet
to which each mutant ZF binds (top). The most common motif for the *de novo* binding sites is
shown below.. Each bar graph shows the percentage of WT only, common and mutant only CTCF

binding sites. The heatmaps show the profile of CTCF and ATAC-seq signal at those sites. The

UN condition corresponds to the FLAG control. (**B**) Profiles of ATAC-seq in WT and mutants.

Wilcoxon p-values were coded as follow: NS, not significant, * $5x10^{-2}$-$5x10^{-3}$, ** $5x10^{-3}$-$5x10^{-4}$, ***

$5x10^{-4}$-$5x10^{-5}$, **** $<5x10^{-5}$. Data were generated on 2 replicates. See also **Figures S6-9.**

Figure 3: **Each mutation uniquely impacts CTCF's chromatin bound fraction, residence**

time and interaction with DNA. (**A**) Plots of FRAP dynamics for WT and mutant CTCF. The bold

lines show the fitted model of the average recovery, and the outlines give the 95% confidence

intervals (95%CIs). (**B**) Violin plots of specific bound fractions. (**C**) Violin plots of specific residence

times (min). p-values were determined by bootstrapping (n=2500). (**D**) Heatmaps show the

proportion of CTCF-cohesin versus CTCF only binding sites. UN corresponds to the FLAG control

in untreated cells. (**E**) Correlation between residence time and the percentage of CTCF-cohesin

overlap. (**F**) Correlation between the FRAP specific bound fraction relative to WT and the fraction

of common CTCF sites relative to all potential binding sites. (**G**) Correlation between the FRAP

specific bound fraction relative to WT and the effect of CTCF binding on ATAC-seq signal at

CTCF-SMC3 sites (**Figure 2B**). For **E**, **F** and **G,** data were generated in 2 replicates, the p-values

were calculated using linear regression, the shaded area corresponds to the 95% CI. See also

Figures S10-14.

Figure 4. Effect of CTCF binding and accessibility on SMC3 overlap. (A) Bar graph showing

the independent effect of CTCF (top) and ATAC-seq (bottom) signal on SMC3 enrichment in

WT. The error bars correspond to the 95%CIs. The p-values were calculated using a

multivariate logistic model. (**B**) Percentage of SMC3 overlap in WT and CTCF mutants, stratified

by ATAC-seq and CTCF signals. **(C)** SMC3 profiles in WT and mutant CTCF. p-values were

calculated using Wilcoxon tests. See also **Figures S15-16** and **Table S1**.

Figure 5: **CTCF mutations alter gene expression, cellular reprogramming and TF binding.**

(**A**) Heatmap showing supervised clustering of the cell lines based on the expression levels of

DEGs identified across the comparisons to WT. (**B**) Gene set enrichment analysis of DEGs in IAA

condition and in H455R. The volcano plots below highlight the DEGs belonging to these enriched

pathways. (**C**) Radar plots showing the averaged expression of developmental germ layer genes

in WT and mutant mESCs cultured in LIF and no LIF conditions. (**D**) Heatmap showing predicted

differentially bound TFs in WT, mutants and IAA. CTCF is highlighted with a *. (**E**) Volcano plots

highlighting the differentially expressed target genes of CTCF and MBD2 in IAA and CTCF

mutants. The metrics for enrichment of the target genes among the DEGs are reported on top on

the volcanos (ORs and logistic p-values). (**F**) Examples of altered CTCF binding and footprinting

at *Rerg* promoter (left) and altered MYC footprinting at *Brdt* promoter (right). All data in this figure

were generated on 2 replicates. See also **Figures S17-20** and **Table S2-4**.

Figure 6: CTCF mutations alter chromatin interactivity. (A) The top panels show the

aggregated differential TAD analysis between mutants and WT**. (a)** Reflects the intra-TAD

interaction, **(b)** and **(c)** highlight the inter-TAD interactions**.** The lower panels show the aggregated

differential peak analysis. The data was generated by Hi-C on 2 replicates. (**B**) Correlation

between the interaction counts and the FRAP residence times. (**C**) Correlation between the

insulation score at CTCF peaks and the FRAP residence times. (**D**) Correlation between the loop

extrusion length and the FRAP bound fractions. (**E**) Example of differential interactions between

mutants and WT (right). The left matrix shows the interaction in WT within a 10 Mb region (40 kb

resolution) with the insulation score on the side. (**F**) Profiles show the averaged insulation score

at WT only, common and mutant only binding sites. (**G**) Profiles show the insulation score at

CTCF binding sites stratified by CTCF signals. The bar graph shows the independent effect of

CTCF signal on insulation score. (**H**) Profiles show the insulation score at CTCF binding sites

stratified by ATAC signals. The bar graph shows the independent effect of ATAC signal on

insulation score. For **F, G** and **H,** p-values reported in the profiles were calculated using Kruskal-

Wallis tests. The fitted estimates for the bar graphs in (**G**) and (**H**) were obtained using a mixed

multivariate model. The error bars correspond to the 95%CI. See also **Figures S21-24**.

Figure 7: Changes in gene expression are linked to changes in chromatin interactivity but

to a lesser extent) than changes in TF binding at gene promoters: (**A**) Bar graphs showing

the enrichment of over-expressed (top) or under-expressed (bottom) genes in gained (top) or lost

(bottom) loops in IAA and mutants compared to WT. (**B**) Example of 2 loci (blue and red

rectangles) with a direct effect of gain in CTCF binding and chromatin interactivity. The interaction

matrices (left) show both gain of intra- and inter-TAD interactions in some mutants compared to

WT. The left panels show the zoom-in tracks of these loci. Only significant differential chromatin

loops are shown. Overexpressed genes are highlighted in red. **C**) Bar graph showing the

percentage of DEGs resulting from direct or indirect effect of CTCF binding distinguishing loop

dependent and independent effect.

STAR Methods

Key resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
mouse anti FLAG	Sigma	Cat#F1804
mouse anti GAPDH	BioLegend	Cat#W17079A
mouse anti CTCF (WB)	Active Motif	Cat#61311
mouse anti CTCF (Chip-seq)	Cell signaling	Cat#3418
FLAG M2 Magnetic Beads	Sigma	Cat#M8823
mouse anti SMC3	abcam	Cat#ab9263
Mouse anti RAD21	abcam	Cat#ab992
Mouse anti SMC1	ThermoFisher	Cat#MA5-15583
Rabbit IgG	abcam	Cat#ab37415
Critical commercial assays		
Tagment DNA Enzyme and Buffer Kit	Illumina	Cat#20034198
Illumina Stranded Total RNA Prep, Ligation with Ribo-Zero Plus		Cat#20040529
Arima Hi-C kit	Arima	Cat#A510008
Oligonucleotide		
Nextera barcodes	Schmidl et al [54]	NA
Deposited data		
RNA-seq	This paper	GEO accession number available upon publication
HiC	This paper	GEO accession number available upon publication
OmniATAC-seq	This paper	GEO accession number available upon publication
SMC3 Chipmentation	This paper	GEO accession number available upon publication
FLAG Chipmentation	This paper	GEO accession number available upon publication
H3K27ac peaks from 120/Ola mESCs	ENCODE	ENCFF519QMV
HOCOMOCO v11	Kulakovskiy et al. [74]	https://hocomoco11.autosome.org/
Developmental germ layer gene sets	Hutchins et al. [68]	NA
CTCF associated NDD genes	Konrad et al. [9]	NA
Developmental genes	DECIPHER	https://www.deciphergenomics.org/ddd/overview
Imprinted genes	geneimprint	https://www.geneimprint.com/
ENCODE blacklisted regions	Amemiya et al. [61]	https://github.com/Boyle-Lab/Blacklist
KEGG genesets	Kanehisa et al. [32]	https://www.genome.jp/kegg/genes.html
CTCF-DNA complex (WT ZF1-7)	PDB	8SSS
CTCF-DNA complex (K365T ZF1-7)	PDB	8SST

CTCF-DNA complex (WT ZF3-11)	PDB	8SSQ
CTCF-DNA complex (K365T ZF3-11)	PDB	8SSR
Experimental models: Cell lines		
Mouse embryonic stem cells E14Tg2a (129/Ola isogenic background) expressing transgene WT or mutant CTCF	This paper	NA
pEN366 - pTRE3G-CTCF-mRuby2-BGHpA-CAGGS-rtta3G-rbgpA-Frt-PGK-EM7-PuroR-bpA-Frt TIGRE donor	Nora EP et al. [15]	Addgene #156432
pX330-EN1201	Nora EP et al. [15]	Addgene #92144
pEN114 - pCAGGS-Tir1-V5-BpA-Frt-PGK-EM7-PuroR-bpA-Frt-Rosa26	Nora EP et al. [15]	Addgene #92143
pASH40-H2B-mRuby-neo (H2B-mRuby)	Hansen et al. [24]	NA
Recombinant DNA		
Super piggyBac Transposase expression vector (replaces PB200A-1)	SBI (System Biosciences)	PB210PA-1
Software and algorithms		
FlowJo	BD	https://www.bdbiosciences.com/en-us/products/software/flowjo-v10-software
PyMol 3.1.0	Schrödinger	https://www.pymol.org/
Alphafold3	Abramson et al [53]	https://alphafoldserver.com/
Bowtie2	Tarasov et al [59]	https://github.com/BenLangmead/bowtie2
MACS2	Amemiya et al [61]	https://pypi.org/project/MACS2/
trimGalore	Krueger [57]	https://zenodo.org/record/7598955
Sambamba	Tarasov et al. [59]	https://lomereiter.github.io/sambamba/
MEME	Bailey et al. [21]	https://meme-suite.org/meme/
Deeptools	Ramirez et al. [63]	https://deeptools.readthedocs.io/en/develop/
pheatmap	Kolde et al. [66]	https://www.rdocumentation.org/packages/pheatmap/versions/1.0.12/topics/pheatmap
fgsea	Sergushichev et al. [67]	https://github.com/ctlab/fgsea
STAR aligner	Dobin et al. [64]	https://github.com/alexdobin/STAR
Seq-N-Slide pipeline	Dolgalev [56]	https://doi.org/10.5281/zenodo.5550459
HiC-Pro	Servant et al. [69]	https://github.com/nservant/HiC-Pro

HiCExplorer	Wolff et al. [70]	https://hicexplorer.readthedocs.io/en/latest/
GENOVA	van der Weide et al. [44]	https://github.com/robinweide/GENOVA
FitHic	Ay et al. [77]	https://github.com/ay-lab/fithic
pairToBed	Quinlan et al. [78]	https://bedtools.readthedocs.io/en/latest/content/tools/pairtopair.html
samtools	Danecek et al. [62]	https://github.com/samtools/samtools
DESeq2	Love et al. [65]	https://bioconductor.org/packages/release/bioc/html/DESeq2.html
cooler	Abdennur et al. [71]	https://cooler.readthedocs.io/en/latest/
STATA v17	stataCorp	https://www.stata.com/stata17/
R	CRAN	https://www.r-project.org/
TOBIAS	Bentsen et al. [35]	https://github.com/loosolab/TOBIAS
UROPA	Kondili et al. [75]	https://uropa-manual.readthedocs.io/custom.html
MultiHiCCompare	Stansfield et al. [79]	https://github.com/dozmorovlab/multiHiCcompare

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Cell lines

Mouse embryonic stem cells E14Tg2a (karyotype 19, XY; 129/Ola isogenic background) and all clones derived from these were cultured under feeder-free conditions in 0.1% gelatin (Sigma ES-006-B) coated dishes (Falcon, 353003) at 37°C and 5% CO_2 in a humidified incubator. The cells were grown in DMEM (Thermo Fisher, 11965-118) supplemented with 15% fetal bovine serum (Thermo Fisher, SH30071.03), 100 U/ml penicillin - 100 µg/ml streptomycin (Sigma, P4458), 1 X GlutaMax supplement (Thermo Fisher, 35050-061), 1 mM sodium pyruvate (Thermo Fisher, 11360-070), 1 X MEM non-essential amino-acids (Thermo Fisher, 11140-50), 50 µM b-mercaptoethanol (Sigma, 38171), 10^4 U/ml leukemia inhibitory factor (Millipore, ESG1107), 3 µM CHIR99021 (Sigma, SML1046) and 1 µM MEK inhibitor PD0325901 (Sigma, PZ0162). The cells were passaged every alternate day by dissociation with TrypLE (Thermo Fisher, 12563011). For the exit from pluripotency experiments, mESCs were cultured for 3 days in the same medium but

without the addition of LIF (leukemia inhibitory factor, CHIR99021 (Sigma, SML1046)) and MEK

inhibitor (PD0325901).

METHOD DETAILS

DNA constructs

Construction of vector for cloning transgenic, doxycycline-inducible expression of WT and mutant

mouse *Ctcf* cDNA were obtained from GenScript in pUC19 vectors. The cDNA was amplified such

that it harbors an AflII sequence at the 3' end of the gene and fused to a FLAG tag (that harbors

NotI sequence) at the 5' end with the help of a fusion PCR. The resultant fragment was digested

with NotI and AflII. The *Ctcf* gene was removed from pEN366[15] by digesting with the same

enzymes. This backbone was used for insertion of each *Ctcf* mutant.

In brief, the cDNA region corresponding to each of the C and N terminals and zinc fingers were

PCR amplified in such a way that it included a short stretch of the 5′ and/or 3′ region of the

neighboring fragment to be connected. The desired PCR products were then annealed, amplified

by PCR and cloned into the NotI and AflII sites of the pEN366 backbone (Addgene #156432). All

the constructs were verified by DNA sequence analysis. For all transgenes, the final vector

harbors an N terminal 3 X FLAG tag and a C terminal *mRuby* as in-frame fusion to WT and mutant

Ctcf. The vector also harbors a *TetO-3G* element and *rtTA3G* for doxycycline induced expression

of the transgene, and homology arms surrounding the sgRNA target site of the *Tigre* locus for

locus-specific insertion. The selection of stable integrants was achieved by virtue of *FRT-PGK-*

puro-FRT cassette. Further details of the vector are described elsewhere [15]. The vector pX330-

EN1201 [15] harboring spCas9 nuclease and *Tigre*-targeting sgRNA was used for targeting of *Tigre*

locus (Addgene #92144).

Gene targeting

Mouse embryonic stem cell E14Tg2a harboring *Ctcf-AID-eGFP* on both alleles and a knock-in of

pEN114 - *pCAGGS-Tir1-V5-BpA-Frt-PGK-EM7-PuroR-bpA-Frt-Rosa26* at *Rosa26* locus was

used as the parental cell line for making all the transgenes [15]. pEN366 derived vectors harboring

the rescue transgenes (WT and mutant *Ctcf*) were used for targeting transgenes to the *Tigre* locus

(clone ID# EN156.3.5) [14]. For nucleofections, 15 µg each of plasmids harboring the transgenes

and 2.5 µg of those with sgRNA targeting the *Tigre* locus was used. Nucleofection were performed

using Amaxa P3 Primary Cell kit (Lonza, V4XP-3024) and 4D-transfector. 2 million cells were

transfected with program CG-104 in each case. The cells were recovered for 48 h with no

antibiotic followed by selection in puromycin (1 µg/mL) (Thermo Fisher, A1113803). Single

colonies were manually picked and expanded in 96 well plates. Clones were genotyped by PCR

and flow cytometry was performed to confirm that the level of expression of transgenes were

comparable. All the clones that were used for the analyses were homozygous for the integration

of the transgenes and their levels of expression were comparable.

Induction of auxin inducible degradation of CTCF and doxycycline induced expression

For degradation of endogenous CTCF, the auxin-inducible degron was induced by adding 500

µM indole-3-acetic acid (IAA, chemical analog of auxin) (Sigma, I5148) to the media. Expression

of transgenes was achieved by the addition of doxycycline (Dox, 1 µg/ml) (Sigma, D9891) to the

media. The cells were treated with IAA and/or Dox for 2 days.

Western Blotting

mESCs were dissociated using TrypLE, washed in PBS, pelleted and used for western blotting.

Approximately 2 million cells were used to prepare cell extract. Cell pellets were resuspended in

RIPA lysis buffer (Thermo Fisher, 89900) with 1X HALT protease inhibitors (Thermo Fisher,

78430), incubated on ice for 30 min, spun at 4^OC at 13,000 rpm for 10 min and supernatant was

collected. For the western blot of CTCF, low salt lysis buffer (0.1 M NaCl, 25 mM HEPES, 1 mM

$MgCl_2$, 0.2 mM EDTA and 0.5% NP40) was used supplemented with 125 U/ml of benzonase

(Sigma E1014). Protein concentration was measured using the Pierce BCA assay kit (Thermo

Fisher, 23225). 20 µg of protein were mixed with Laemmli buffer (Biorad, 1610737) and β-

mercaptoethanol, heated at 95^OC for 10 min and run on a Mini-protean TGX 4%-20%

polyacrylamide gel (Biorad, 456-1095). The proteins were transferred onto PVDF membranes

using the Mini Trans-Blot Electrophoretic Transfer Cell (Bio-Rad, 170-3930) at 80 V, 120 mA for

90 min. PVDF membranes were blocked with 5% BSA in 1 X TBST prior to the addition of

antibody. The membranes were probed with appropriate antibodies overnight at 4^OC, mouse anti-

FLAG antibody (Sigma, F1804; 1: 1,000 dilution), GAPDH (BioLegend, W17079A; 1:5,000

dilution), CTCF (active motif, 61311; 1: 2,000 dilution), SMC1 (ThermoFischer, MA5-15583.

1:1000 dilution), RAD21 (abcam, ab992,1:2000 dilution). Membranes were washed five times in

PBST (1 × PBS and 0.1% Tween 20) for 5 min each and incubated with respective secondary

antibodies in 5% BSA at room temperature for 1 h. The blots were rinsed in PBST and developed

using enhanced chemiluminescence (ECL) and imaged by Odyssey LiCor Imager (Kindle

Biosciences).

Flow cytometric analysis

Cells were dissociated with TrypLE, washed and resuspended in MACS buffer for flow cytometric analysis on LSRII UV (BD Biosciences). Analysis was performed using the FlowJo software.

Cell culture – FRAP

Control cells expressing H2B-mRuby were engineered using the same background as the CTCF transgene expressing cells, E14Tg2a mESCs. Briefly, pEN114 - *pCAGGS-Tir1-V5-BpA-Frt-PGK-EM7-PuroR-bpA-Frt-Rosa26* was used as the parental cell line to be nucleofected by the pASH40-H2B-mRuby-neo with the PiggyBac transposase vector (SBI, PB210PA-1). Halo-H2B plasmid was previously published in Hansen *et al,* [24] and we replaced the Halotag with mRuby. For the FRAP experiment, CTCF and H2B expressing cells were cultured for two days on 35 mm no 1.5H glass-bottom imaging dishes (MatTek, Ashland, MA, P35G-1.5-14C) coated with Geltrex (Gibco, A1413201) according to manufacturer's instructions. 48 hours prior to the experiment, the culture medium was changed to medium supplemented with 500 µM 3-Indoleacetic acid (IAA; Sigma-Aldrich, I2886-5G) and 1 µg/mL doxycycline (Sigma-Aldrich, D3072-1ML). Just prior to imaging, the medium was changed to imaging medium: phenol red free DMEM with all other aspects of the medium the same.

Fluorescence recovery after photobleaching (FRAP)

FRAP experiments were performed on an LSM900 confocal microscope (Zeiss, Germany) equipped with a full, humidified incubation chamber that was maintained at 37°C with 5.5% CO_2. The time series were acquired in confocal mode using two-stage acquisition with the following excitation parameters: 561 nm excitation laser at 0.8% power, 53 µm pinhole size corresponding to 1 AU, scan speed 7, and 3x crop factor corresponding to a pixel dwell time of 3.06 µs. During the first phase, we imaged 25 frames with 0.5 seconds between frames, and during the second phase we imaged 160 frames with 4 seconds between frames, resulting in a total movie length of 657 seconds. For bleaching, a 1µm diameter ROI was selected. Bleaching was performed after frame 8 using both the 488nm and 561nm lasers at 100% power, 53µm pinhole size, and scan speed 7 corresponding to a pixel dwell time of 3.06µs. For each condition, 30 movies were acquired and between 3 and 11 were excluded due to drift.

ChIPmentation

mESCs were dissociated using TrypLE (Thermofisher #12605010) and washed once in 1X PBS. After counting, cells were divided in 10 million aliquots and resuspended in fresh 1X PBS (1million cells/1ml). For double cross linking 25mM EGS (ethylene glycol bis(succinimidyl succinate);

Thermofisher #21565) were added and cells were put in rotation for 30 min at room temperature,

followed by addition of 1% formaldehyde (Tousimis #1008A) for 10 min also in rotation at room

temperature. Quenching was performed by adding glycine to a final concentration of 0.125 M

followed by incubations of 5 min at room temperature in rotation. Fixed cells were washed twice

with 5 ml of 1 X PBS containing 0.5% BSA and centrifuged at 3000 rpm for 5 min at 4°C. Pellets

were finally resuspended in 500 µl 1X PBS containing 0.5% BSA, transferred to 1.5 ml Eppendorf

and centrifuged at 3000rpm for 3 min at 4°C. Supernatant was completely removed, pellets were

snap-frozen in liquid nitrogen and stored at -80°C. Fixed cells (10 million) were thawed on ice,

resuspended in 350 µl ice cold lysis buffer (10 mM Tris-HCl (pH 8.0), 100 mM NaCl, 1 mM EDTA

(pH 8.0), 0.5 mM EGTA (pH 8.0), 0.1% sodium deoxycholate, 0.5% N-lauroysarcosine and 1X

protease inhibitors) and lysed for 10 min by rotating at 4° C. Chromatin was sheared using a

bioruptor (Diagenode) for 15 minutes (30 sec on, 30 sec off, high output level). 100 µl of cold

lysis buffer and 50 µl of 10% Triton X-100 (final concentration of 1) were then added and the

samples were centrifuged for 5 min at full speed at 4°C. Supernatant was collected, transferred

to a new tube (Protein Low Binding tube) and shearing was continued for another 10 min, then

the chromatin was quantified. FLAG M2 Magnetic Beads (Sigma, M8823) were used for FLAG

immunoprecipitation. In other cases, SMC3 (abcam, ab9263), CTCF (cell Signaling Technology,

3418) and Rabbit IgG (abcam, ab37415) antibodies were bound to protein A magnetic beads by

incubation on a rotator for one hour at room temperature. 10 µl each of antibody was bound to 50

µl of protein-A magnetic beads (Dynabeads) and added to the sonicated chromatin for

immunoprecipitation at 4°C overnight. Next day, samples were washed and tagmentation were

performed as per the original ChIPmentation protocol [54]. In short, the beads were washed

successively twice in 500 µl cold low-salt wash buffer (20 mM Tris-HCl (pH 7.5), 150 mM NaCl, 2

mM EDTA (pH 8.0), 0.1% SDS, 1% tritonX-100), twice in 500 µl cold LiCl-containing wash buffer

(10 mM Tris-HCl (pH 8.0), 250 mM LiCl, 1 mM EDTA (pH 8.0), 1% triton X-100, 0.7% sodium

deoxycholate) and twice in 500 µl cold 10 mM cold Tris-Cl (pH 8.0) to remove detergent, salts

and EDTA. Subsequently, the beads were resuspended in 25 µl of the freshly prepared

tagmentation reaction buffer (10 mM Tris-HCl (pH 8.0), 5 mM MgCl2, 10% dimethylformamide)

and 1 µl Tagment DNA Enzyme from the Tagment DNA Enzyme and Buffer Kit (Illumina

#20034198) and incubated at 37°C for 10 min in a thermomixer. Following tagmentation, the

beads were washed successively twice in 500 µl cold low-salt wash buffer (20 mM Tris-HCl (pH

7.5), 150 mM NaCl, 2 mM EDTA (pH8.0), 0.1% SDS, 1% triton X-100) and twice in 500 µl cold

Tris-EDTA-Tween buffer (0.2% tween, 10 mM Tris-HCl (pH 8.0), 1 mM EDTA (pH 8.0)). Chromatin

was eluted and de-crosslinked by adding 70 µl of freshly prepared elution buffer (0.5% SDS, 300

mM NaCl, 5 mM EDTA (pH 8.0), 10 mM Tris-HCl (pH 8.0) and 10 ug/ml proteinase K for 1 hour

at 55°C 850rpm and overnight at 65°C 850rpm. Next day, the supernatant was collected,

transferred to new DNA Low Binding tubes and supplemented with an additional 30 µl of elution

buffer. DNA was purified using MinElute Reaction Cleanup Kit (Qiagen #28204) and eluted in 20

ul. Purified DNA (20 µl) was amplified as per the ChIPmentation protocol (Schmidl, Rendeiro et

al. 2015) using indexed and non-indexed primers and NEBNext High-Fidelity 2X PCR Master Mix

(NEB M0541) in a thermomixer with the following program: 72°C for 5 m; 98°C for 30 s; 14 cycles

of 98°C for 10 s, 63°C for 30 s, 72°C for 30 s and a final elongation at 72°C for 1 m. DNA was

purified using Agencourt AMPure XP beads (Beckman, A63881) to remove fragments larger than

700 bp as well as the primer dimers. Library quality and quantity were estimatred using

Tapestation (Agilent High Sensitivity D1000 ScreenTape #5067-5584 and High Sensitivity D1000

reagents #5067-5585) and quantified by Qubit (Life Technologies Qubit™ 1X dsDNA High

Sensitivity (HS) #Q33230). Libraries were then sequenced with the Novaseq6000 Illumina

technology according to the standard protocols and with around 200 million 150bp paired-end

total per sample.

Total RNA-seq

mESCs were dissociated using TrypLE, washed in 1X PBS, pelletted and 2.5 million cells were

used for extracting RNA with RNeasy plus kit (Qiagen #74134) in each case. RNA quality was

checked in Tapestation (Agilent High Sensitivity RNA ScreenTape #5067-5579, High Sensitivity

RNA Sample Buffer #5067-5580 and High Sensitivity RNA Ladder #5067-5581) and quantified

by Nanodrop. 500 ng were used for Total RNA libraries preparation using Illumina kit (Illumina

Stranded Total RNA Prep, Ligation with Ribo-Zero Plus #20040529). Library concentrations were

estimated using Tapestation (Agilent High Sensitivity D1000 ScreenTape #5067-5584 and High

Sensitivity D1000 reagents #5067-5585) and quantified by Qubit (Life Technologies Qubit™ 1X

dsDNA High Sensitivity (HS) #Q33230). Libraries were then sequenced with the Novaseq6000

Illumina technology according to the standard protocols and with around 100 million 150bp paired-

end total per sample.

Hi-C

Hi-C was performed in duplicates using around 1 million cells each. mESCs were dissociated

using TrypLE (Thermofishe #12605010), washed once in 1X PBS and resuspended in fresh 1X

PBS (1 million cells/ 1 ml). For double cross linking 25 mM EGS (ethylene glycol bis(succinimidyl

succinate); Thermofisher #21565) were added and cells were put in rotation for 30 min at room

temperature, followed by addition of 1% formaldehyde (Tousimis #1008A) for 10 min also in

rotation at room temperature. Quenching was performed by adding glycine to a final concentration

of 0.125M followed by incubations of 5 min at room temperature in rotation. Fixed cells were

washed twice with 5 ml of 1X PBS containing 0.5% BSA and centrifuged at 3000 rpm for 5 min at

4°C. Pellets were finally resuspended in 500 µl 1X PBS containing 0.5% BSA, transferred to 1.5ml

Eppendorf and centrifuged at 3000 rpm for 3 min at 4°C. Supernatant was completely removed,

pellets were snap-frozen in liquid nitrogen and stored at -80°C. Samples were subsequently

processed using the Arima Hi-C kit (A510008) as per the manufacturer's protocol and sequenced

with the Novaseq6000 Illumina technology according to the standard protocols and with around

600 million 150bp paired-end reads per sample.

ATAC-seq

We used an improved ATAC-seq protocol (Omni-ATAC) adopted from Corces et al., [22] with small

adaptations. Briefly, cells in culture were treated with 200 U/ml DNase (Worthington # LS002007)

for 30 min at 37°C to remove free-floating DNA and any DNA from dead cells. Cells were then

harvested via trypsinization and resuspended in regular medium. After counting, 500,000 cells

were collected, resuspended in 1X cold PBS and spin down at 500 g for 5 min at 4°C in a fixed-

angle centrifuge. After centrifugation, the supernatant was removed and the pellet resuspended

in 500 µl of cold ATAC-seq resuspension buffer 1 (10 mM Tris-HCl pH 7.4, 10 mM NaCl, 3 mM

$MgCl_2$, 0.1% NP-40 and 0.1% Tween-20 in water). 50 µl (50.000 cells) were transferred to a new

1.5 ml Eppendorf tube and 0.5 µl l 1% Digitonin (Promega #G9441) was added by pipetting well

few times. The cell lysis reaction was incubated on ice for 3 min. After lysis, 1 ml of cold ATAC-

seq resuspension buffer 2 (10 mM Tris-HCl pH 7.4, 10 mM NaCl, 3 mM $MgCl_2$ and 0.1% Tween-

20 in water) was added. Tube was inverted three times to mix and nuclei were pelleted by

centrifugation for 10 min at 500 g at 4°C. Supernatant was very carefully removed (pellet could

be almost invisible at this step) and nuclei resuspended in 45 µl of transposition mix (25 µl 2X TD

buffer (Illumina #20034198), 16.5 µl 1X PBS, 0.5 µl 1% digitonin, 0.5 µl 10% Tween-20 and 2.5

µl water) by pipetting up and down few times. 5 µl of transposase enzyme (Illumina #20034198)

was then added and the transposition reaction was incubated at 37°C for 30 min in a thermomixer

with shaking at 1000 rpm. Reaction was cleaned up with Zymo DNA Clean and Concentrator-5

kit (Zymo#11-302C). Transposed DNA fragments were then amplified as described previously in

Buenrostro et al., [55]. Final libraries were purified with with Zymo DNA Clean and Concentrator-5

kit (Zymo#11-302C; Note: better to use two separate kits for pre- and post-amplification clean up),

checked by Tapestation (Agilent High Sensitivity D1000 ScreenTape #5067-5584 and High

Sensitivity D1000 reagents #5067-5585) and quantified by Qubit (Life Technologies Qubit™ 1X

dsDNA High Sensitivity (HS) #Q33230). Libraries were then sequenced with the Novaseq6000

Illumina technology according to the standard protocols and with around 50 million 150bp paired-

end reads per sample.

QUANTIFICATION AND STATISTICAL ANALYSIS

Fluorescence recovery after photobleaching (FRAP) analysis

FRAP analysis was performed using custom Python scripts. Briefly, movies were loaded into a

custom Python GUI (https://github.com/ahansenlab/FRAP-drift-correction-GUI). The bleached

nucleus was segmented using manually selected segmentation parameters and was used to

estimate the average nuclear intensity and the average background intensity. To correct for cell

drift, the ROI positions were manually updated at a number of frames and the ROI position was

linearly interpolated between manual updates. Cells with excessive drift were manually excluded

and the mean fluorescence intensity of the bleach was taken as the average intensity within the

ROI.

To correct for photobleaching, $I_{nonbleach}(t)$ was smoothed using an averaging filter, and

a series of correction factors ($C(t)$) were computed as the ratio of these values to their average

prebleach value as follows:

$$C(t) = \frac{I_{nonbleach,smooth}(t) - I_{background}(t)}{\langle I_{nonbleach}(t) - I_{background}(t)\rangle_{prebleach}}$$

1000
1001 Subsequently, the FRAP curves were normalized according to the following equation:
1002
1003

1004 $$\underline{I_{bleach}(t)} = \frac{C(t)(I_{bleach}(t) - I_{background}(t))}{\langle C(t)(I_{bleach}(t) - I_{background}(t))\rangle_{prebleach}}$$

1005
1006 where $\underline{I_{bleach}(t)}$ is the normalized, background-subtracted, and photobleaching corrected FRAP

1007 signal.

1008 Finally, since previous measurements of CTCF's free diffusion coefficient have indicated

1009 a reaction-dominant model is most appropriate, we fit the following, previously demonstrated

1010 model (Sprague, Pego et al. 2004, Hansen, Pustova et al. 2017).

1011
1012 $$FRAP(t) = 1 - C_{eq,fast}e^{-k_{off,fast}t} - C_{eq,slow}e^{-k_{off,slow}t}$$

1013

where $k_{off,slow} < k_{off,fast}$. $C_{eq,slow}$ was taken as the specific bound fraction and the specific

residence time was taken as $\frac{1}{k_{off,slow}}$. To estimate parameter distributions, we implemented a

bootstrapping routine in which a set of 16 movies were randomly resampled from the full dataset

2500 times with replacement. For each sample, the above equation was fit and the parameter

values were taken as one data point. The 2500 resampled points were then used as distributions

for plotting the specific bound fraction and residence time. For ease of comparison, the specific

bound fractions were normalized to the average WT value when plotting.

1021

ChIP-seq data processing and quality control

Data were processed using Seq-N-Slide pipeline [56]. Briefly, after trimming for NEXTERA adaptor

sequence using trimGalore [57]. The reads were aligned to mm10 genome with Bowtie2 [58].

Ambiguous reads were filtered to use uniquely mapped reads in the downstream analysis. PCR

duplicates were removed using Sambamba [59]). Narrow peaks were called using MACS2 [60] in pair-

end mode and with IgG as control for SMC3 and CTCF ChIP-seq or WT untreated condition (in

which FLAG CTCF is not expressed) for CTCF FLAG to control for unspecific binding, Peaks

overlapping ENCODE blacklisted regions were filtered out [61]. These procedures allowed us to

minimize the false positive calls, as reflected by the high percentage of called peaks containing

CTCF consensus motifs as detected by MEME [21] with 63% for WT. To ensure comparability

between conditions, FLAG and SMC3 chip-seq samples were down-sampled to the same number

of aligned (100 millions reads per replicates), except for 2 samples R377H-rep1 and R339Q-rep

which were under-sequenced (with 68 and 90 millions reads, respectively) and good quality reads

before peak calling using samtools [62]. Bigwigs were obtained for visualization on individual as well

as merged bam files using Deeptools [63]. For CTCF FLAG visualization, the differential signals

were generated after subtracting non-specific signal present in the WT untreated condition.

Heatmaps and average profiles were performed on merged bigwig files using Deeptools.

1039

RNA-seq data processing and quality control

Data were processed using Seq-N-Slide pipeline [56]. Briefly, after trimming for the Illumina adaptor

and the T overhang as per Illumina recommendation using TrimGalore [57], reads were aligned

against the mouse reference genome (mm10) using the STAR [64] aligner and differentially

expressed genes were called using DESeq2 [65] with an adjusted p-value of 0.05 and a fold change

cutoff of 1.5. Heatmap was performed using pheatmap R package [66] and GSEA using fgsea R

package [67] and KEGG genesets [32]. Developmental germ layer gene sets were downloaded from

Hutchins *et a*l.,[68]. Radarplots were generated using the R package fsmb after averaging the
expression level of the genes of each germ layers and of the 2 replicates. CTCF-associated
neurodevelopmental disorder (NDD) genes were retrieved from Konrad *et al.* [9], while genes
associated with developmental disorders were downloaded from DECIPHER
(https://www.deciphergenomics.org/ddd) and imprinted genes from geneimprint
(https://www.geneimprint.com/). Enrichment analyses for those sets were performed using
logistic regressions. To compare the change in gene expression during the exit from pluripotency
between WT and mutant (longitudinal analysis), linear regressions on the difference of log2
between LIF versus no LIF condition were performed.

1056

ATAC-seq data processing and quality control
Data were processed using Seq-N-Slide pipeline [56]. Briefly, Reads were aligned to mm10 genome
with Bowtie2 [58]. Ambiguous reads were filtered to use uniquely mapped reads in the downstream
analysis. PCR duplicates were removed using Sambamba [59]. Mitochondrial contamination was
assessed using samtools [62]. All the samples showed less than 0.6% of mitochondrial DNA, Peaks
were called using MACS2 [60] and peaks overlapping ENCODE blacklisted regions were filtered out
[61]. Bigwigs were obtained for visualization on individual as well as merged bam files using
Deeptools [63]. Heatmaps and average profiles were performed on merged bigwig files using
Deeptools.

1066

Hi-C Processing and Quality Control.
HiC-Pro [69] was used to align and filter the Hi-C data. To generate Hi-C filtered contact matrices,
the Hi-C reads were aligned against the mouse reference genome (mm10) by bowtie2 (version
2.3.1) using local mode for the second step of HiC-Pro alignment (--very-sensitive -L 20 --score-
min G,20,8 --local) to ignore the molecular barcode UMI sequences and dark bases introduced
by Arima library prep kit at the 5' end of the read. Singleton, ambiguous and duplicated reads
were filtered out. Ligation sites for Arima HiC kit were set up using
GATCGATC,GANTGATC,GANTANTC,GATCANTC. Low mapping quality (MAPQ<30), self
circle, dangling ends and re-ligation reads were filtered out through the HiC-pro pipeline. Valid
pairs represented more than 80% for all the samples. Samples were downscaled to the same
number of good quality uniquely aligned reads using HiCExplorer [70]. Of note, we did not use the
number of valid pairs to downscale the samples to not overcorrect some conditions such as IAA
and mutant H455R which were expected to show a dramatic biological decrease in valid
interactions.

1081

Downscaled interaction matrices were generated separately for each replicate and merged for

visualization and downprocessing as mcool files using cooler [71]. Samples were balanced using

the ICE correction method [72] with HiCExplorer [70]. HiC maps were generated from the merged

downscaled and corrected matrices using GENOVA and HiCExplorer [44,70]. RCP (relative contact

probabilities) and insulation score were generated using GENOVA [44]. After log-log transformation

and Loess smoothing using R, the first derivative (P'max) was estimated as described in Gassler

et al., [41].

1089

1090   **CTCF peak motif analysis**

1091   Motifs were called using MEME [21] with a 1st order background model on CTCF peaks called using

1092   MACS2 after separating the peaks into 3 groups for each pairwise comparison between WT and

1093   mutant CTCF: WT only, common and mutant only peaks. Peak intersection sets were generated

1094   using the intervene package [73] and motifs were searched within 100 bp of each peak summit.

1095

AlphaFold3 prediction of the DNA-CTCF complexes in the mutants

As a quality control, to compare the prediction to experimental data, we downloaded the structure

deposited by Yang *et al*. [13] from PDB for WT CTCF ZF1-ZF7 (accession number PDB 8SSS, with

oligo CCAGCAGGGGGCGCTAGTGAG and CTCACTAGCGCCCCCTGCTGG) , K365T CTCF

ZF1-ZF7 (accession number PDB 8SST with the same oligos as WT), WT CTCF ZF3-ZF11

(accession number PDB 8SSQ with oligo

GTGCAGTACCACATTTAACCAGCAGGGGGCGCTAA and

TTAGCGCCCCCTGCTGGTTAAATGTGGTACTGCAC), K365T CTCF ZF3-ZF11 (accession

number PDB 8SSR with oligo with the same oligos as WT). For the Alphafold3 predictions we

used the same DNA sequences and the CTCF ZF1-11 positions obtained from Yang *et al* [13].

AlphaFold3 predictions [53] were also carried out for WT and K365T CTCF ZF3-ZF11 and CTCF

ZF1-7 using the same DNA and CTCF sequence to compare the prediction to the experimental

data. We, then, predicted the CTCF (ZF1-11)-DNA complex for WT and each mutant CTCF using

the concatenated DNA sequences from Yang *et al* [13]

(GTGCAGTACCACATTTAACCAGCAGGGGGCGCTAGTGAG and

CTCACTAGCGCCCCCTGCTGGTTAAATGTGGTACTGCAC). The predicted protein-DNA

structures were visualized in PyMOL 3.1.0. The hydrogen bonds between DNA and CTCF were

identified using PyMOL dist function with a cutoff of 5 Å. The root mean square deviation (RMSD)

between the structures was calculated using PyMOL 3.1.0.

1115

Stratified analysis at enhancers and promoters

1116

1117 H3K27ac peaks from 120/Ola mESCs were downloaded from ENCODE (ENCFF519QMV).

1118 CTCF, cohesin and ATAC-seq peaks were classified as active promoter sites when they overlap

1119 with an H3K27ac peak and are located in a gene promoter region, defined as 2kb upstream and

1120 1 kb downstream gene TSSs. Active enhancer sites when they overlap with an H3K27ac peak

1121 and are located outside gene promoter regions.

1122

Multivariate CTCF/SMC3 overlap and insulation score analysis

1123

To test the independent of CTCF and Accessibility on SMC3 overlap, CTCF ChIP-seq and ATAC-

seq signal strengths were categorized into weak, intermediate and strong signal using the inter-

quartile values of the signal across WT and mutant samples. Weak binding or accessibility was

defined as sites with signals inferior to the 1^{st} quartile, intermediate binding or accessibility as

sites with signals between the 2^{nd} and 3^{rd} quartiles and strong binding or accessibility as sites with

signals superior to the 3^{rd} quartile. For SMC3 overlap, a multivariate logistic model testing the

effect of CTCF strength, ATAC strength and their interaction was performed in WT mESCs after

averaging the replicates. The log2 odd ratios were computed to estimate the enrichment in SMC3.

For insulation score a mixed multivariate model testing the effect of CTCF strength, ATAC

strength and their interaction was performed across sample. For both models the marginal effects

and 95th confidence intervals were computed. These analyses were performed using STATA v17.

1135

ATAC-seq footprinting analysis

1136

Footprinting analysis was performed on ATAC-seq signal using TOBIAS [35]. Briefly, ATAC-seq

signals were first corrected for Tn5 cutting sequence bias using the ATACorrect function and

footprinting score calculated using FootprintScores at peaks detected by MACS. Finally

differential footprinting estimated pariwise between WT and mutant CTCF using BINDetect and

HOCOMOCO v11 core motif probability weight matrices [74]. Peaks were annotated for gene

promoter using UROPA [75] and GENCODE version M25 [76]. Gene promoters were defined as

regions within 2kb upstream and 1 kb downstream the gene transcription starting site.

1144

Loop calling and annotation

1145

1146 Significant loops (q-value<0.05) were called on the downscaled and corrected interaction

1147 matrices using FitHIC at 10kb resolution [77] and annotated for CTCF peaks and gene TSS using

1148 pairToBed [78].

1149

Differential aggregated peak and TAD analyses

Aggregated peak and TAD analysis on downscaled and corrected interaction matrices was performed using GENOVA at 10kb resolution [44]. For aggregated peaks analyses, the union of loops called by FitHIC [77] in WT and mutant CTCF for each pairwise comparison was used to generate and compare aggregated interaction at loop anchors. For aggregated TAD analyses, the insulation score was first calculated to identify TAD and TAD boundaries. The union of TADs called by GENOVA in WT and mutant CTCF was used to generate and compare aggregated intra and inter-TAD interactions.

Differential loop analysis

Differential loops (adjusted p-value<0.05) were called for each pairwise comparison between WT and mutant CTCF using MultiHiCCompare which performed a cross-sample normalization before testing for differential loops using an exact test [79]. Enrichment for differentially expressed genes among differential loops was tested using logistic regression after annotating loop anchors with gene TSS.

Direct vs indirect effect analysis

Direct effect was defined as differential TF binding with a predicted differential CTCF binding identified using TOBIAS [35] or ChIP-seq peak at the gene promoter and/or differential loops identified using MultiHiCCompare [79] with a CTCF peak at either anchor. Gene promoters were defined as regions spanning 2 kb upstream and 1 kb downstream gene TSSs. Overlap between gene promoters and differential loop were performed using pairToBed [78].

SUPPLEMENTAL INFORMATION:

Excel table title and legends

Table S1: Number and percentage of SMC3 peaks overlapping a CTCF or ATAC-seq peak, stratified by CTCF and ATAC signal strength. This table is related to **Figure 4B.**

Table S2: Differentially expressed genes (DEGs) identified using DE-seq. DEGs were defined as adjusted genes with p-value<0.05 and absolute log2 fold change $\geq$ log2(1.5) compared to WT. This table is related to **Figure 5A**.

1183

Table S3: CTCF mutations altered cell differentiation. Tab A, Expression of germ layer genes

(TPM) in LIF versus no LIF condition in WT and the mutant. **Tab B,** Differentially expressed genes

during in LIF condition in mutant compared to WT. **Tab C,** Differentially expressed genes during

in no LIF condition in mutant compared to WT. **Tab D,** Differentially expressed genes during the

exit from pluripotency (LIF to no LIF) in mutant compared to WT. This table is related to **Figure**

5C.

1190

1191 **Table S4: CTCF mutations altered TF binding. Tab A,** Differentially bound transcription factors

1192 predicted by footprinting analyses. This tab is related to **Figure 5D**. **Tab B,** Enrichment of the

1193 target genes of differentially bound TFs among the DEGs. This tab is related to **Figure 5E**. **Tab**

1194 **C,** List of the target genes of differentially bound TFs with differential expression. This tab is

1195 related to **Figure 5E**.

References:

1. Davidson, I.F., Bauer, B., Goetz, D., Tang, W., Wutz, G., and Peters, J.M. (2019). DNA loop extrusion by human cohesin. Science *366*, 1338-1345. 10.1126/science.aaz3418.

2. Fudenberg, G., Imakaev, M., Lu, C., Goloborodko, A., Abdennur, N., and Mirny, L.A. (2016). Formation of Chromosomal Domains by Loop Extrusion. Cell Rep *15*, 2038-2049. 10.1016/j.celrep.2016.04.085.

3. Hansen, A.S. (2020). CTCF as a boundary factor for cohesin-mediated loop extrusion: evidence for a multi-step mechanism. Nucleus *11*, 132-148. 10.1080/19491034.2020.1782024.

4. Dekker, J., and Mirny, L. (2016). The 3D Genome as Moderator of Chromosomal Communication. Cell *164*, 1110-1121. 10.1016/j.cell.2016.02.007.

5. Filippova, G.N., Lindblom, A., Meincke, L.J., Klenova, E.M., Neiman, P.E., Collins, S.J., Doggett, N.A., and Lobanenkov, V.V. (1998). A widely expressed transcription factor with multiple DNA sequence specificity, CTCF, is localized at chromosome segment 16q22.1 within one of the smallest regions of overlap for common deletions in breast and prostate cancers. Genes Chromosomes Cancer *22*, 26-36.

6. Xiang, Y., Zhou, X., Hewitt, S.L., Skok, J.A., and Garrard, W.T. (2011). A multifunctional element in the mouse Igkappa locus that specifies repertoire and Ig loci subnuclear location. J Immunol *186*, 5356-5366. 10.4049/jimmunol.1003794.

7. Debaugny, R.E., and Skok, J.A. (2020). CTCF and CTCFL in cancer. Curr Opin Genet Dev *61*, 44-52. 10.1016/j.gde.2020.02.021.

8. Gregor, A., Oti, M., Kouwenhoven, E.N., Hoyer, J., Sticht, H., Ekici, A.B., Kjaergaard, S., Rauch, A., Stunnenberg, H.G., Uebe, S., et al. (2013). De novo mutations in the genome organizer CTCF cause intellectual disability. Am J Hum Genet *93*, 124-131. 10.1016/j.ajhg.2013.05.007.

9. Konrad, E.D.H., Nardini, N., Caliebe, A., Nagel, I., Young, D., Horvath, G., Santoro, S.L., Shuss, C., Ziegler, A., Bonneau, D., et al. (2019). CTCF variants in 39 individuals with a

variable neurodevelopmental disorder broaden the mutational and clinical spectrum.
Genet Med *21*, 2723-2733. 10.1038/s41436-019-0585-z.

10. Valverde de Morales, H.G., Wang, H.V., Garber, K., Cheng, X., Corces, V.G., and Li, H.
(2023). Expansion of the genotypic and phenotypic spectrum of CTCF-related disorder
guides clinical management: 43 new subjects and a comprehensive literature review. Am
J Med Genet A *191*, 718-729. 10.1002/ajmg.a.63065.

11. Chen, H., Tian, Y., Shu, W., Bo, X., and Wang, S. (2012). Comprehensive identification
and annotation of cell type-specific and ubiquitous CTCF-binding sites in the human
genome. PLoS One *7*, e41374. 10.1371/journal.pone.0041374.

12. Maurano, M.T., Wang, H., John, S., Shafer, A., Canfield, T., Lee, K., and
Stamatoyannopoulos, J.A. (2015). Role of DNA Methylation in Modulating Transcription
Factor Occupancy. Cell Rep *12*, 1184-1195. 10.1016/j.celrep.2015.07.024.

13. Yang, J., Horton, J.R., Liu, B., Corces, V.G., Blumenthal, R.M., Zhang, X., and Cheng, X.
(2023). Structures of CTCF-DNA complexes including all 11 zinc fingers. Nucleic Acids
Res *51*, 8447-8462. 10.1093/nar/gkad594.

14. Nishana, M., Ha, C., Rodriguez-Hernaez, J., Ranjbaran, A., Chio, E., Nora, E.P., Badri,
S.B., Kloetgen, A., Bruneau, B.G., Tsirigos, A., and Skok, J.A. (2020). Defining the relative
and combined contribution of CTCF and CTCFL to genomic regulation. Genome Biol *21*,
108. 10.1186/s13059-020-02024-0.

15. Nora, E.P., Goloborodko, A., Valton, A.L., Gibcus, J.H., Uebersohn, A., Abdennur, N.,
Dekker, J., Mirny, L.A., and Bruneau, B.G. (2017). Targeted Degradation of CTCF
Decouples Local Insulation of Chromosome Domains from Genomic
Compartmentalization. Cell *169*, 930-944 e922. 10.1016/j.cell.2017.05.004.

16. Holland, A.J., Fachinetti, D., Han, J.S., and Cleveland, D.W. (2012). Inducible, reversible
system for the rapid and complete degradation of proteins in mammalian cells. Proc Natl
Acad Sci U S A *109*, E3350-3357. 10.1073/pnas.1216880109.

17. Nora, E.P., Caccianini, L., Fudenberg, G., So, K., Kameswaran, V., Nagle, A., Uebersohn,
A., Hajj, B., Saux, A.L., Coulon, A., et al. (2020). Molecular basis of CTCF binding polarity
in genome folding. Nat Commun *11*, 5612. 10.1038/s41467-020-19283-x.

18.   Cerami, E., Gao, J., Dogrusoz, U., Gross, B.E., Sumer, S.O., Aksoy, B.A., Jacobsen, A.,
1254   Byrne, C.J., Heuer, M.L., Larsson, E., et al. (2012). The cBio cancer genomics portal: an
1255   open platform for exploring multidimensional cancer genomics data. Cancer Discov *2*,
1256   401-404. 10.1158/2159-8290.CD-12-0095.

1257   19.   Tate, J.G., Bamford, S., Jubb, H.C., Sondka, Z., Beare, D.M., Bindal, N., Boutselakis, H.,
1258   Cole, C.G., Creatore, C., Dawson, E., et al. (2019). COSMIC: the Catalogue Of Somatic
1259   Mutations In Cancer. Nucleic Acids Res *47*, D941-D947. 10.1093/nar/gky1015.

1260   20.   Price, E., Fedida, L.M., Pugacheva, E.M., Ji, Y.J., Loukinov, D., and Lobanenkov, V.V.
1261   (2023). An updated catalog of CTCF variants associated with neurodevelopmental
1262   disorder phenotypes. Front Mol Neurosci *16*, 1185796. 10.3389/fnmol.2023.1185796.

1263   21.   Bailey, T.L., Johnson, J., Grant, C.E., and Noble, W.S. (2015). The MEME Suite. Nucleic
1264   Acids Res *43*, W39-49. 10.1093/nar/gkv416.

1265   22.   Corces, M.R., Trevino, A.E., Hamilton, E.G., Greenside, P.G., Sinnott-Armstrong, N.A.,
1266   Vesuna, S., Satpathy, A.T., Rubin, A.J., Montine, K.S., Wu, B., et al. (2017). An improved
1267   ATAC-seq protocol reduces background and enables interrogation of frozen tissues. Nat
1268   Methods *14*, 959-962. 10.1038/nmeth.4396.

1269   23.   Do, C., Jiang, G., Zappile, P., Heguy, A., and Skok, J.A. (2024). A genome wide code to
1270   define cell-type specific CTCF binding and chromatin organization. bioRxiv.
1271   10.1101/2024.11.02.620823.

1272   24.   Hansen, A.S., Pustova, I., Cattoglio, C., Tjian, R., and Darzacq, X. (2017). CTCF and
1273   cohesin regulate chromatin loop stability with distinct dynamics. Elife *6*.
1274   10.7554/eLife.25776.

1275   25.   Sprague, B.L., Pego, R.L., Stavreva, D.A., and McNally, J.G. (2004). Analysis of binding
1276   reactions by fluorescence recovery after photobleaching. Biophys J *86*, 3473-3495.
1277   10.1529/biophysj.103.026765.

1278   26.   Hansen, A.S., Amitai, A., Cattoglio, C., Tjian, R., and Darzacq, X. (2020). Guided nuclear
1279   exploration increases CTCF target search efficiency. Nat Chem Biol *16*, 257-266.
1280   10.1038/s41589-019-0422-3.

27. Iurlaro, M., Masoni, F., Flyamer, I.M., Wirbelauer, C., Iskar, M., Burger, L., Giorgetti, L.,
and Schubeler, D. (2024). Systematic assessment of ISWI subunits shows that NURF
creates local accessibility for CTCF. Nat Genet *56*, 1203-1212. 10.1038/s41588-024-
01767-x.

28. Rahmaninejad, H., Xiao, Y., Tortora, M.M.C., and Fudenberg, G. (2024). Dynamic barriers
modulate cohesin positioning and genome folding at fixed occupancy. bioRxiv.
10.1101/2024.10.08.617113.

29. de Jonge, W.J., Patel, H.P., Meeussen, J.V.W., and Lenstra, T.L. (2022). Following the
tracks: How transcription factor binding dynamics control transcription. Biophys J *121*,
1583-1592. 10.1016/j.bpj.2022.03.026.

30. Mach, P., Kos, P.I., Zhan, Y., Cramard, J., Gaudin, S., Tunnermann, J., Marchi, E.,
Eglinger, J., Zuin, J., Kryzhanovska, M., et al. (2022). Cohesin and CTCF control the
dynamics of chromosome folding. Nat Genet *54*, 1907-1918. 10.1038/s41588-022-01232-
7.

31. Gabriele, M., Brandao, H.B., Grosse-Holz, S., Jha, A., Dailey, G.M., Cattoglio, C., Hsieh,
T.S., Mirny, L., Zechner, C., and Hansen, A.S. (2022). Dynamics of CTCF- and cohesin-
mediated chromatin looping revealed by live-cell imaging. Science *376*, 496-501.
10.1126/science.abn6583.

32. Kanehisa, M., and Goto, S. (2000). KEGG: kyoto encyclopedia of genes and genomes.
Nucleic Acids Res *28*, 27-30. 10.1093/nar/28.1.27.

33. Isles, A.R. (2022). The contribution of imprinted genes to neurodevelopmental and
neuropsychiatric disorders. Transl Psychiatry *12*, 210. 10.1038/s41398-022-01972-4.

34. Hirai, H., Karian, P., and Kikyo, N. (2011). Regulation of embryonic stem cell self-renewal
and   pluripotency   by   leukaemia   inhibitory   factor.   Biochem   J   *438*,   11-23.
10.1042/BJ20102152.

35. Bentsen, M., Goymann, P., Schultheis, H., Klee, K., Petrova, A., Wiegandt, R., Fust, A.,
Preussner, J., Kuenne, C., Braun, T., et al. (2020). ATAC-seq footprinting unravels kinetics
of transcription factor binding during zygotic genome activation. Nat Commun *11*, 4267.
10.1038/s41467-020-18035-1.

36.   Lax, E., Do Carmo, S., Enuka, Y., Sapozhnikov, D.M., Welikovitch, L.A., Mahmood, N.,
1311   Rabbani, S.A., Wang, L., Britt, J.P., Hancock, W.W., et al. (2023). Methyl-CpG binding
1312   domain 2 (Mbd2) is an epigenetic regulator of autism-risk genes and cognition. Transl
1313   Psychiatry *13*, 259. 10.1038/s41398-023-02561-9.

1314   37.   Mahmood, N., Arakelian, A., Szyf, M., and Rabbani, S.A. (2024). Methyl-CpG binding
1315   domain protein 2 (Mbd2) drives breast cancer progression through the modulation of
1316   epithelial-to-mesenchymal transition. Exp Mol Med *56*, 959-974. 10.1038/s12276-024-
1317   01205-2.

1318   38.   Menafra, R., and Stunnenberg, H.G. (2014). MBD2 and MBD3: elusive functions and
1319   mechanisms. Front Genet *5*, 428. 10.3389/fgene.2014.00428.

1320   39.   Reynolds, N., Latos, P., Hynes-Allen, A., Loos, R., Leaford, D., O'Shaughnessy, A.,
1321   Mosaku, O., Signolet, J., Brennecke, P., Kalkan, T., et al. (2012). NuRD suppresses
1322   pluripotency gene expression to promote transcriptional heterogeneity and lineage
1323   commitment. Cell Stem Cell *10*, 583-594. 10.1016/j.stem.2012.02.020.

1324   40.   Holwerda, S.J., and de Laat, W. (2013). CTCF: the protein, the binding partners, the
1325   binding sites and their chromatin loops. Philos Trans R Soc Lond B Biol Sci *368*,
1326   20120369. 10.1098/rstb.2012.0369.

1327   41.   Gassler, J., Brandao, H.B., Imakaev, M., Flyamer, I.M., Ladstatter, S., Bickmore, W.A.,
1328   Peters, J.M., Mirny, L.A., and Tachibana, K. (2017). A mechanism of cohesin-dependent
1329   loop extrusion organizes zygotic genome architecture. EMBO J *36*, 3600-3618.
1330   10.15252/embj.201798083.

1331   42.   Edelmann, W., Yang, K., Umar, A., Heyer, J., Lau, K., Fan, K., Liedtke, W., Cohen, P.E.,
1332   Kane, M.F., Lipford, J.R., et al. (1997). Mutation in the mismatch repair gene Msh6 causes
1333   cancer susceptibility. Cell *91*, 467-477. 10.1016/s0092-8674(00)80433-x.

1334   43.   Chen, Y., Liu, P., Sun, P., Jiang, J., Zhu, Y., Dong, T., Cui, Y., Tian, Y., An, T., Zhang, J.,
1335   et al. (2019). Oncogenic MSH6-CXCR4-TGFB1 Feedback Loop: A Novel Therapeutic
1336   Target of Photothermal Therapy in Glioblastoma Multiforme. Theranostics *9*, 1453-1473.
1337   10.7150/thno.29987.

44. van der Weide, R.H., van den Brand, T., Haarhuis, J.H.I., Teunissen, H., Rowland, B.D.,
and de Wit, E. (2021). Hi-C analyses with GENOVA: a case study with cohesin variants.
NAR Genom Bioinform *3*, lqab040. 10.1093/nargab/lqab040.

45. Liu, X.H., Liu, L.P., Xu, X.M., Hua, M., Kang, Q., Li, A., and Huang, L. (2021). FOXN2
suppresses the proliferation and invasion of human hepatocellular carcinoma cells. Eur
Rev Med Pharmacol Sci *25*, 731-737. 10.26355/eurrev_202101_24634.

46. Ma, J., Lu, Y., Zhang, S., Li, Y., Huang, J., Yin, Z., Ren, J., Huang, K., Liu, L., Yang, K.,
et al. (2018). beta-Trcp ubiquitin ligase and RSK2 kinase-mediated degradation of FOXN2
promotes tumorigenesis and radioresistance in lung cancer. Cell Death Differ *25*, 1473-
1485. 10.1038/s41418-017-0055-6.

47. Ye, H., and Duan, M. (2019). FOXN2 is downregulated in breast cancer and regulates
migration, invasion, and epithelial- mesenchymal transition through regulation of SLUG.
Cancer Manag Res *11*, 525-535. 10.2147/CMAR.S176938.

48. Ikeda, K., Horie-Inoue, K., Suzuki, T., Hobo, R., Nakasato, N., Takeda, S., and Inoue, S.
(2019). Mitochondrial supercomplex assembly promotes breast and endometrial
tumorigenesis by metabolic alterations and enhanced hypoxia tolerance. Nat Commun *10*,
4108. 10.1038/s41467-019-12124-6.

49. Do, C., and Skok, J.A. (2024). Factors that determine cell type-specific CTCF binding in
health and disease. Curr Opin Genet Dev *88*, 102244. 10.1016/j.gde.2024.102244.

50. Soochit, W., Sleutels, F., Stik, G., Bartkuhn, M., Basu, S., Hernandez, S.C., Merzouk, S.,
Vidal, E., Boers, R., Boers, J., et al. (2021). CTCF chromatin residence time controls three-
dimensional genome organization, gene expression and DNA methylation in pluripotent
cells. Nat Cell Biol *23*, 881-893. 10.1038/s41556-021-00722-w.

51. Watson, L.A., Wang, X., Elbert, A., Kernohan, K.D., Galjart, N., and Berube, N.G. (2014).
Dual effect of CTCF loss on neuroprogenitor differentiation and survival. J Neurosci *34*,
2860-2870. 10.1523/JNEUROSCI.3769-13.2014.

52. Narducci, D.N., and Hansen, A.S. (2025). Putative looping factor ZNF143/ZFP143 is an
essential transcriptional regulator with no looping function. Mol Cell *85*, 9-23 e29.
10.1016/j.molcel.2024.11.032.

53.   Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger, O.,
Willmore, L., Ballard, A.J., Bambrick, J., et al. (2024). Accurate structure prediction of
biomolecular interactions with AlphaFold 3. Nature *630*, 493-500. 10.1038/s41586-024-
07487-w.

54.   Schmidl, C., Rendeiro, A.F., Sheffield, N.C., and Bock, C. (2015). ChIPmentation: fast,
robust, low-input ChIP-seq for histones and transcription factors. Nat Methods *12*, 963-
965. 10.1038/nmeth.3542.

55.   Buenrostro, J.D., Wu, B., Chang, H.Y., and Greenleaf, W.J. (2015). ATAC-seq: A Method
for Assaying Chromatin Accessibility Genome-Wide. Curr Protoc Mol Biol *109*, 21 29 21-
21 29 29. 10.1002/0471142727.mb2129s109.

56.   Dolgalev,   I.   (2022).   Seq-N-Slide   [Computer   software].
https://doi.org/10.5281/zenodo.5550459.

57.   Krueger, F. (2021). TrimGalore. https://zenodo.org/record/7598955.

58.   Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat
Methods *9*, 357-359. 10.1038/nmeth.1923.

59.   Tarasov, A., Vilella, A.J., Cuppen, E., Nijman, I.J., and Prins, P. (2015). Sambamba: fast
processing   of   NGS   alignment   formats.   Bioinformatics   *31*,   2032-2034.
10.1093/bioinformatics/btv098.

60.   Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E., Nusbaum,
C., Myers, R.M., Brown, M., Li, W., and Liu, X.S. (2008). Model-based analysis of ChIP-
Seq (MACS). Genome Biol *9*, R137. 10.1186/gb-2008-9-9-r137.

61.   Amemiya, H.M., Kundaje, A., and Boyle, A.P. (2019). The ENCODE Blacklist:
Identification of Problematic Regions of the Genome. Sci Rep *9*, 9354. 10.1038/s41598-
019-45839-z.

62.   Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O., Whitwham,
A., Keane, T., McCarthy, S.A., Davies, R.M., and Li, H. (2021). Twelve years of SAMtools
and BCFtools. Gigascience *10*. 10.1093/gigascience/giab008.

63.   Ramirez, F., Ryan, D.P., Gruning, B., Bhardwaj, V., Kilpert, F., Richter, A.S., Heyne, S.,
Dundar, F., and Manke, T. (2016). deepTools2: a next generation web server for deep-
sequencing data analysis. Nucleic Acids Res *44*, W160-165. 10.1093/nar/gkw257.

64.   Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner.
Bioinformatics *29*, 15-21. 10.1093/bioinformatics/bts635.

65.   Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550. 10.1186/s13059-014-
0550-8.

66.   Kolde, R. (2019). pheatmap: Pretty Heatmaps. R package version 1.0.12.

67.   Sergushichev, A. (2016). An algorithm for fast preranked gene set enrichment analysis
using cumulative statistic calculation. bioRxiv. doi: 10.1101/060012.
http://biorxiv.org/content/early/2016/06/20/060012.

68.   Hutchins, A.P., Yang, Z., Li, Y., He, F., Fu, X., Wang, X., Li, D., Liu, K., He, J., Wang, Y.,
et al. (2017). Models of global gene expression define major domains of cell type and
tissue identity. Nucleic Acids Res *45*, 2354-2367. 10.1093/nar/gkx054.

69.   Servant, N., Varoquaux, N., Lajoie, B.R., Viara, E., Chen, C.J., Vert, J.P., Heard, E.,
Dekker, J., and Barillot, E. (2015). HiC-Pro: an optimized and flexible pipeline for Hi-C
data processing. Genome Biol *16*, 259. 10.1186/s13059-015-0831-x.

70.   Wolff, J., Bhardwaj, V., Nothjunge, S., Richard, G., Renschler, G., Gilsbach, R., Manke,
T., Backofen, R., Ramirez, F., and Gruning, B.A. (2018). Galaxy HiCExplorer: a web
server for reproducible Hi-C data analysis, quality control and visualization. Nucleic Acids
Res *46*, W11-W16. 10.1093/nar/gky504.

71.   Abdennur, N., and Mirny, L.A. (2020). Cooler: scalable storage for Hi-C data and other
genomically labeled arrays. Bioinformatics *36*, 311-316. 10.1093/bioinformatics/btz540.

72.   Imakaev, M., Fudenberg, G., McCord, R.P., Naumova, N., Goloborodko, A., Lajoie, B.R.,
Dekker, J., and Mirny, L.A. (2012). Iterative correction of Hi-C data reveals hallmarks of
chromosome organization. Nat Methods *9*, 999-1003. 10.1038/nmeth.2148.

73.   Khan, A.M., A (2017). Intervene: a tool for intersection and visualization of multiple
genomic region sets.

74.   Kulakovskiy, I.V., Vorontsov, I.E., Yevshin, I.S., Sharipov, R.N., Fedorova, A.D.,
Rumynskiy, E.I., Medvedeva, Y.A., Magana-Mora, A., Bajic, V.B., Papatsenko, D.A., et al.
(2018). HOCOMOCO: towards a complete collection of transcription factor binding models
for human and mouse via large-scale ChIP-Seq analysis. Nucleic Acids Res *46*, D252-
D259. 10.1093/nar/gkx1106.

75.   Kondili, M., Fust, A., Preussner, J., Kuenne, C., Braun, T., and Looso, M. (2017). UROPA:
a tool for Universal RObust Peak Annotation. Sci Rep *7*, 2593. 10.1038/s41598-017-
02464-y.

76.   Frankish, A., Diekhans, M., Jungreis, I., Lagarde, J., Loveland, J.E., Mudge, J.M., Sisu,
C., Wright, J.C., Armstrong, J., Barnes, I., et al. (2021). Gencode 2021. Nucleic Acids Res
49, D916-D923. 10.1093/nar/gkaa1087.

77.   Ay, F., Bailey, T.L., and Noble, W.S. (2014). Statistical confidence estimation for Hi-C data
reveals regulatory chromatin contacts. Genome Res *24*, 999-1011.
10.1101/gr.160374.113.

78.   Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for comparing
genomic features. Bioinformatics *26*, 841-842. 10.1093/bioinformatics/btq033.

79.   Stansfield, J.C., Cresswell, K.G., and Dozmorov, M.G. (2019). multiHiCcompare: joint
normalization and comparative analysis of complex Hi-C experiments. Bioinformatics *35*,
2916-2923. 10.1093/bioinformatics/btz048.

1443

Figure 1

Figure 2

A

B

ATAC profiles at sites <u>overlapping</u> with Smc3 peaks, with or without CTCF binding

ATAC profiles at sites <u>not overlapping</u> with Smc3 peaks, with or without CTCF binding

Figure 3

Figure 4

A

Effect of CTCF adjusted for the effect of ATAC

p=1e-11 p=1.2e-19 p=2.5e-59

Enrichment in Smc3 (Log2 Odd ratios)

Weak   Interm.   Strong
Strength

Effect of ATAC adjusted for the effect of CTCF

NS   NS   p=8e-06

Enrichment in Smc3 (Log2 Odd ratios)

Weak   Interm.   Strong
Strength

B

Percentage of SMC3 overlap stratified by CTCF and ATAC signal strengths

Strong ATAC
Interm. ATAC
Weak ATAC
No ATAC

WT   H455R   K365T   Q418R   R339Q   R342C   R377C   R377H   R448Q   S354F

C

Smc3 profiles at accessible sites with or without CTCF binding

Smc3 ChIP-seq signal (RPKM)

WT — CTCF, Smc3, ATAC 8162 / Smc3, ATAC 36152
R342C — CTCF, Smc3, ATAC 6430 / Smc3, ATAC 53654
R377H — CTCF, Smc3, ATAC 8284 / Smc3, ATAC 59195
R377C — CTCF, Smc3, ATAC 15430 / Smc3, ATAC 44420
S354F — CTCF, Smc3, ATAC 10043 / Smc3, ATAC 29739

Smc3 profiles at inaccessible sites with or without CTCF binding

Smc3 ChIP-seq signal (RPKM)

WT — CTCF, Smc3 6956 / Smc3 23447
R342C — CTCF, Smc3 9092 / Smc3 46758
R377H — CTCF, Smc3 8820 / Smc3 49731
R377C — CTCF, Smc3 14451 / Smc3 40499
S354F — CTCF, Smc3 11884 / Smc3 36232

Figure 5

Figure 6

Figure 7

SUPPLEMENTAL FIGURES

Figure S1. Prevalence of missense CTCF mutations on ZF3-ZF7 in human cancers.
Schematic representation of CTCF showing the location and frequency of each mutation under investigation. This figure relates to **Figure 1**.

Figure S2

Mutations reported in > than 2 samples	Zinc Finger	No. of samples	Associated cancers	Associated neurological disorder
Mutations in phosphate contacts (positions -5, 4th position of the linker between adjacent ZFs)				
R377C*	5	27	Breast, Colorectal, Endometrial, Haematopoietic And Lymphoid, Skin Cancer	ASD, CRD
R377H*	5	26	Breast, Colorectal, Endometrial, Esophagogastric, Haematopoietic And Lymphoid, Urinary Tract	ASD, CRD
R320C	3	6	Colorectal, Endometrial, Prostate Cancer	
R320H	3	3	Colorectal, Esophagogastric, Head And Neck Cancer	
K349E	4	2	Endometrial Cancer	
R377L	5	2	Breast, Esophagogastric Cancer	
Y358C	4	2	Colorectal, Endometrial Cancer	
Mutations in the amino acid that make direct contact with DNA (positions -1, +2, +3 and +6)				
R339W	3	20	Colorectal, Endometrial, Haematopoietic And Lymphoid, Renal Cancer	
R339Q*	3	12	Endometrial, Esophagogastric, Haematopoietic And Lymphoid Cancer	CRD,ASD,DD
Q418R*	6	5	Embryonal Tumor, Glioma, Hepatobiliary Cancer	
R448Q*	7	5	Endometrial, Haematopoietic And Lymphoid, Renal Cancer	CRD,ASD,DD
K365T*	4	4	Endometrial Cancer	
R368C	4	4	Colorectal, Esophagogastric, Haematopoietic And Lymphoid Cancer	DD,NDD,CRD,ASD
D451H	7	3	Skin, Urinary Tract Cancer	
D451V	7	3	Haematopoietic And Lymphoid Cancer	
R368H	4	3	Breast, Ovary, Prostate Cancer	CRD,ASD,DD
E362K	4	2	Haematopoietic And Lymphoid Cancer	
D390N	5	2	Haematopoietic And Lymphoid Cancer	CRD,ASD
Mutations in the amino acids in the other regions of the zinc finger				
S354F*	4	21	Breast, Endometrial, Esophagogastric, Head And Neck, Hepatobiliary, Skin, Urinary Tract Cancer	
P378L	5	10	Endometrial, Haematopoietic And Lymphoid Cancer	NDD, CRD, DD
S388N	5	6	Haematopoietic And Lymphoid, Hepatobiliary, Urinary Tract Cancer	
S402L	5	6	Breast Cancer	
P378A	5	5	Breast, Urinary Tract Cancer	
F351L	4	5	Breast, Urinary Tract Cancer	
G375V	4	5	Endometrial, Haematopoietic And Lymphoid Cancer	
A359T	4	4	Colorectal, Ovary, Prostate Cancer	
A387V	5	4	Breast, Endometrial Cancer	
R415W	6	4	Colorectal, Endometrial, Penile Cancer	
F416L	6	4	Endometrial, Glioma Cancer	
E376K	4	4	Breast, Haematopoietic And Lymphoid, Urinary Tract Cancer	
K352N	4	3	Endometrial Cancer	
D328N	3	3	Colorectal, Endometrial Cancer	
S461F	7	3	Skin Cancer	
E348K	3	3	Colorectal, Haematopoietic And Lymphoid Cancer	
F379I	5	2	Haematopoietic And Lymphoid Cancer	
P440S	7	2	Skin Cancer	
I411V	6	2	Urinary Tract Cancer	
H413Y	6	2	Colorectal Cancer	
D443G	7	2	Hepatobiliary Cancer	
F331Y	3	2	Haematopoietic And Lymphoid Cancer	
R341C	3	2	Colorectal, Skin Cancer	
R341H	3	2	Head And Neck Cancer	
Q428E	6	2	Haematopoietic And Lymphoid Cancer	
K344E	3	2	Breast, Haematopoietic And Lymphoid Cancer	
G375A	4	2	Haematopoietic And Lymphoid, Urinary Tract Cancer	
G375R	4	2	Head And Neck, Lung Cancer	
Mutations in boundary residues (position -2 and +9)				
R342C*	3	13	Breast, Endometrial, Esophagogastric, Haematopoietic And Lymphoid, Skin, Urinary Tract Cancer	NDD,CRD,DD
R371H	4	8	Colorectal, Endometrial, Haematopoietic And Lymphoid, Lung Cancer	
R342H	3	7	Endometrial, Haematopoietic And Lymphoid Cancer	CRD
R371C	4	4	Endometrial, Skin, Urinary Tract Cancer	
R457Q	7	4	Colorectal, Esophagogastric Cancer	
R389S	5	2	Breast, Skin Cancer	
Mutations in the amino acid that co-ordinate the zinc (position -10, -7, +7, +11 or +12 depending on ZF)				
C356R	4	3	Esophagogastric Cancer	
H373L	4	3	Endometrial Cancer	NDD,ASD
H455R*	7	2	Haematopoietic And Lymphoid Cancer	
C324Y	3	2	Haematopoietic And Lymphoid Cancer	
C356F	4	2	Haematopoietic And Lymphoid, Head And Neck Cancer	
H345P	3	2	Breast Cancer	
H345R	3	2	Colorectal, Prostate Cancer	

Figure S2: Table of CTCF mutations. Table showing CTCF mutations identified from cBioPortal [S1] and COSMIC [S2]. We selected a subset of mutations that occur at high frequency in cancer patients, focusing on those found in ZFs that make contact with the 12-15bp consensus CTCF binding motif. These are highlighted with an asterisk and color coded as in **Figure 1D.** Mutations analyzed included those found in **(i)** amino acids that make base-specifying contacts with the alpha helix of DNA (at locations -2, 2, 3 and 6 on the ZF), **(ii)** residues that coordinate the zinc ion which are essential for providing stability to ensure the proper folding of the ZF domain, **(iii)** two other classes of amino acid residues in the ZFs: boundary residues (that are important for interactions between ZFs), and residues that contact the sugar phosphate backbone of DNA, and **(iv)** other highly mutated residues in the region. The table identifies which zinc finger each mutation is found in, the number of patients it was identified in and the cancers each is associated with. Mutations were annotated for neurodevelopmental or neurological disorders using the CTCF variant catalog published in Price et al [S3]. ASD, autism spectrum disorder; CRD, CTCF related neurodevelopmental disorder; DD developmental disorder; NDD, neurodevelopmental disorder. This figure relates to **Figure 1**.

Figure S3

Figure S3: **Flow cytometry analysis of mutant expression.** (**A**) Flow cytometry showing the level of GFP (endogenous CTCF) and mRuby (transgenic CTCF) in the IAA and Dox treated cells (ID). (**B**) Bar graph showing the percentage of GFP (endogenous CTCF) and mRuby2 (transgene WT and mutant CTCF) positive cells after treatment with IAA and Dox (ID). The presence of the IAA leads to degradation of GFP labelled endogenous CTCF. The error bars represent the standard deviation between the 2 replicates. This Figure relates to **Figure 1.**

Figure S4

Figure S4: **Western Blot of endogenous and transgenic CTCF.** Western blot showing degradation of endogenous CTCF as detected with an antibody to CTCF, and the induction of transgenic CTCF as detected using an antibody to FLAG. The first replicate is shown in **Figure 1**. The uncropped pictures of the WB gels are shown below. This figure relates to **Figure 1**.

Figure S5. Rescue of CTCF binding and gene expression by the transgene WT CTCF. (**A**) Heatmap and profiles showing the rescued CTCF binding sites in WT-ID compared to WT-UN. (**B**) Alluvial plot showing the number of DEGs in IAA versus WT untreated cells and WT-ID versus WT untreated fells. The plot shows the percentage of genes that return to baseline in the WT-ID condition (no significant expression level changes). DEGs were defined as fold change>1.5 and adjusted p-value<0.05. This figure relates to **Figure 1.**

Figure S6

Figure S6: **CTCF mutations have unique chromatin binding profiles. (A)** Scheme showing the locations of the different CTCF mutations within the ZF. The consensus CTCF motif highlights the triplet to which each mutant zinc finger binds (top) as well as the motif identified by MEME for the *de novo* mutant only binding sites (bottom). Each graph shows the percentage of WT and mutant CTCF binding at WT only, common and mutant only sites and heatmaps show the profile of CTCF binding and ATAC-seq signal for WT only, common and mutant only sites. The remaining mutants are shown in **Figure 2**. (**B**) Most prevalent motifs found at WT only, common and mutant only sites. The motifs were identified using MEME de novo motif search. The percentages correspond to the percentage of peaks containing the motif. This figure relates to **Figure 2**.

Figure S7

Figure S7. Differential CTCF binding in mutant versus WT-ID. Volcano plots showing significant differences in WT and mutant transgenic CTCF binding (p<0.05 and fold change > 1.5 (red) or fold change < 1.5 (blue). The subthreshold of differentially bound sites are shown in light blue and pink. The p-values were computed using DiffBind. This figure relates to **Figure 2.**

Figure S8

Figure S8: At CTCF WT and mutant sites, accessibility is more or less reduced in a cohesin-dependent manner. Profiles of ATAC-seq in WT and mutant CTCF expressing cells are shown for sites, with (color-coded for each mutant) or without (green) CTCF binding in the presence or absence of SMC3. The remaining mutants are shown in **Figure 2B**. Wilcoxon p-values were coded as follow: * 5×10^{-2}-5×10^{-3}, ** 5×10^{-3}-5×10^{-4}, *** 5×10^{-4}-5×10^{-5}, **** $<5 \times 10^{-5}$. This figure relates to **Figure 2.**

Figure S9

ATAC-seq signal at enhancers

ATAC profiles at sites <u>with</u> Smc3, with or without CTCF binding

ATAC profiles at sites <u>without</u> Smc3, with or without CTCF binding

ATAC-seq signal at promoters

ATAC profiles at sites <u>with</u> Smc3, with or without CTCF binding

ATAC profiles at sites <u>without</u> Smc3, with or without CTCF binding

Figure S9: At CTCF WT and mutant sites overlapping gene promoters and enhancers, accessibility is more or less reduced in a cohesin-dependent manner. Profiles of ATAC-seq in WT and mutant CTCF expressing cells are shown for sites, with (color-coded for each mutant) or without (green) CTCF binding in the presence or absence of SMC3 at gene enhancers (top) and gene promoters (bottom). In the absence of SMC3, for WT and most of the mutants, very few accessible CTCF peaks were observed at enhancers or promoters. Those data should be interpreted with caution. Wilcoxon p-values were coded as follow: * 5×10^{-2}-5×10^{-3}, ** 5×10^{-3}-5×10^{-4}, *** 5×10^{-4}-5×10^{-5}, **** $<5\times10^{-5}$. This figure relates to **Figure 2.**

Figure S10

Figure S10. **Plots of FRAP dynamics for WT and mutant mRuby-CTCF for each pairwise comparison between mutant and WT**. The scatterplots show the average recovery across all movies analyzed, and the outlines give the 95% CI. The bold line shows the fitted model. This figure is related to **Figure 3**.

Figure S11

A AlphaFold prediction CTCF, full protein

B Experimental (8SSQ) versus alpha WT, ZF3-11
RSMD=1.2 Å

C Experimental (8SSS) versus alpha WT, ZF1-7
RSMD=1.5 Å

D Experimental (8SST) versus alpha K365T, ZF1-7
RSMD=1.4 Å

E Experimental WT (8SSS) versus K365T (8SST), ZF1-7
RSMD=0.3 Å

F AlphaFold WT versus K365T, ZF1-7
RSMD=0.3 Å

G Experimental WT (8SSS) versus K365T (8SST), K365T
RSMD=0.3 Å

H AlphaFold WT versus K365T, K365T
RSMD=0.2 Å

Figure S11: Comparison between the prediction the CTCF-DNA complex and the experimental structure. (**A-H**) Comparison of Alphafold3 prediction versus experimental crystal structure of WT and K365T-DNA complex. (**A**) Prediction of the full CTCF structure. On the right the pLDDT matrix shows a high confidence prediction of CTCF ZF1-11 and its positions relative to the DNA double helix. (**B**) Comparison between the experimental and predicted DNA-WT CTCF ZF3-11 complex. (**C**) Comparison between the experimental and predicted DNA-WT CTCF ZF1-7 structure. (**D**) Comparison between the experimental and predicted DNA-K365T mutant CTCF ZF1-7 structure. For **B-D**, the experimental structures are shown in grey and the prediction in color. Each ZF is color-coded. (**E**) Comparison of the DNA-CTCF complex between K365T mutant and WT from the experimental data and (**F**) between K365T mutant and WT from the AlphaFold3 prediction. For **E** and **F**, the WT structures are shown in grey and the mutant in color. (**G**) Comparison of the DNA-CTCF structure between the K365T mutant and WT showing the hydrogen bond between the mutated or WT residue and interacting DNA nucleotide from the experimental data and (**H**) from the prediction. The experimental crystal structures of the DNA-WT and DNA-K365T complexes were generated in Yang et al [29]. This figure is related to **Figure 3**.

Figure S12: **Comparison of the predicted DNA-CTCF complexes between WT and CTCF mutants.** Comparison of the DNA-CTCF (ZF1-11) complexes between WT and each mutant CTCF predicted by AlphaFold3. For mutated residues making direct contact with the DNA, the Zoom-in shows the hydrogen bond between the mutated residue and interacting DNA nucleotide compared to WT. The WT structures are shown in grey and the mutant in color. Each ZF is color-coded. The ZFs bearing the mutations are labelled in red. This figure is related to **Figure 3**.

Figure S13

A

Percentage of FLAG-CTCF and Smc3 overlap

Figure S13: The ability of the CTCF mutants to block cohesin is mostly retained at common sites. Each mutation uniquely. (A) Bar graphs showing the percentage of CTCF-cohesin overlap for each mutant at WT only, common and mutant only sites. For common sites, the CTCF peaks were overlapped with either the WT or mutant SMC3 peaks. (**B**) Western blot showing the protein levels of 2 components of the cohesin complex (SMC1 and RAD21) in WT untreated (UN), IAA (CTCF degraded) and WT and CTCF mutant ID treated cells (cells expressing WT and mutant transgenes in the absence of endogenous CTCF). The uncropped pictures of the WB gels are shown below. This figure is related to **Figure 3**.

Figure S14: The percentage of SMC3 overlapping CTCF and the overall fraction of bound CTCF sites are less informative . (A) Correlation between the percentage of SMC3 overlapping CTCF peaks and the number of CTCF peaks. (**B**) Heatmaps of CTCF (blue) and SMC3 (green) in CTCF WT and mutant expressing cells at CTCF only, CTCF and SMC3, and SMC3 only sites. (**C**) Non-significant correlation between the chromatin bound mutant versus WT fraction detected by FRAP and the fraction of all CTCF binding sites relative to all FLAG and SMC3 ChIP-seq and ATAC-seq peaks. This figure is related to **Figure 3**.

Figure S15

Figure S15: **The relationship between CTCF binding, cohesin overlap and accessibility. (A)** Profiles of SMC3 binding in WT and mutant CTCF expressing cells is shown for accessible (top) and inaccessible (bottom) sites with (color-coded for each mutant) or without (green) CTCF binding. The remaining mutants are shown in **Figure 4C**. Wilcoxon p-values were coded as follow: * 5×10^{-2}-5×10^{-3}, ** 5×10^{-3}-5×10^{-4}, *** 5×10^{-4}-5×10^{-5}, **** $<5 \times 10^{-5}$. **(B)** Profiles of SMC3 binding in WT and mutant CTCF expressing cells compared to IgG at accessible (top) and inaccessible (bottom) sites with or without CTCF (right and left profiles, respectively). Wilcoxon p-values were coded as follow: * 5×10^{-2}-5×10^{-3}, ** 5×10^{-3}-5×10^{-4}, *** 5×10^{-4}-5×10^{-5}, **** $<5 \times 10^{-5}$. This figure relates to **Figure 4**.

Figure S16

Figure S16: **The relationship between CTCF binding, cohesin overlap and accessibility at enhancers and promoters.** Profiles of SMC3 binding in WT and mutant CTCF expressing cells is shown for accessible and inaccessible CTCF binding at enhancers (top) and gene promoters (bottom) with (color-coded for each mutant) or without (green). Wilcoxon p-values were coded as follow: * $5x10^{-2}$-$5x10^{-3}$, ** $5x10^{-3}$-$5x10^{-4}$, *** $5x10^{-4}$-$5x10^{-5}$, **** $<5x10^{-5}$. This figure relates to **Figure 4**.

Figure S17

Figure S17: **CTCF mutations alter gene expression.** (**A**) Bar graph showing the percentage of DEGs found in 1 to 10 mutants, including IAA. The list of genes is reported in **Table S2**. (**B**) Gene set enrichment analysis of DEGs. All other mutants are shown in **Figure 5B**. For each condition, bar graphs show significantly enriched KEGG pathways. The volcano plots below highlight the DEGs belonging to these enriched pathways with genes outside of these pathways shown in black. Brain related pathways are shown in purple, cancer in pink, immune in green and metabolism in mustard. (**C**) Enrichment of the DEGs in over- and under-expressed genes reported in CTCF associated human neuro-developmental disorders [S4]. (**D**) Enrichment of the DEGs in imprinted genes reported in Geneimprint (https://www.geneimprint.com). This figure relates to **Figure 5**.

Figure S18

Figure S18: **Examples of genes implicated in altered differentiation trajectories in WT versus CTCF mutant expressing cells.** Bar graphs showing the comparison of WT versus mutant CTCF expression levels (TPM) in LIF and no LIF medium. The p-values assessing the significance of the longitudinal differences between mutant and WT were computed using linear regressions at the gene level. p-values are coded as followed *: 5×10^{-2}-5×10^{-3}, **5×10^{-3}-5×10^{-4}, *** 5×10^{-4}-5×10^{-5}, ****$<5\times10^{-5}$. This figure relates to **Figure 5**.

Figure S19

A

Log2 Fold chang bindinding score (Mutant vs WT)

TF	Motif	IAA	H455R	Q418R	K365T	R342C	S354F	R448Q	R339Q	R377C	R377H
MBD2		0.1	0.2	0.2	0.2	0.3	0.5	0.5	0.4	0.4	0.3
CTCFL		-0.2	-0.1	0.0	0.2	0.1	0.0	0.5	0.0	0.2	0.1
NRF1		0.2	0.2	0.1	0.3	0.2	0.3	0.5	0.3	0.3	0.3
NFYB		0.2	0.0	0.2	0.3	0.1	0.1	0.4	0.1	0.1	0.5
CTCF		-0.2	-0.2	0.0	0.2	0.1	0.0	0.4	0.0	0.1	0.0
NFYC		0.2	0.0	0.2	0.3	0.1	0.1	0.4	0.1	0.1	0.4
KAISO		0.1	0.2	0.2	0.3	0.3	0.3	0.4	0.3	0.3	0.3
NFYA		0.2	0.0	0.2	0.3	0.1	0.1	0.4	0.1	0.1	0.4
E2F2		0.1	0.2	0.1	0.2	0.3	0.3	0.4	0.3	0.3	0.2
MECP2		0.1	0.1	0.1	0.2	0.2	0.3	0.4	0.2	0.3	0.2
SP2		0.1	0.2	0.1	0.2	0.3	0.3	0.4	0.3	0.3	0.2
PBX3		0.2	0.0	0.2	0.2	0.1	0.0	0.3	0.1	0.1	0.3
E2F1		0.1	0.2	0.1	0.2	0.2	0.3	0.3	0.3	0.3	0.2
AP2B		0.1	0.2	0.1	0.1	0.2	0.3	0.3	0.2	0.2	0.2
E2F4		0.2	0.2	0.2	0.3	0.2	0.3	0.3	0.2	0.3	0.2

B

Figure S19: **CTCF mutations alter TF binding. (A)** Top 15 predicted differentially bound TFs. The TFs were sorted by the maximum absolute log2 fold change of binding score between mutant versus WT observed across mutants. The total number of sites and p-value for these TFs, as well as the full list of differentially bound TFs for each mutant are reported in **Table S4A**. (**B**) Bar graph showing the percentage of differentially bound TFs found in 1 to 9 mutants including IAA (none were found in all the conditions). The list of significant differentially bound TFs is reported in **Table S4A**. This figure relates to **Figure 5**.

Figure S20

Figure S20: **The DEGs are enriched in the target genes of differentially bound TF.** Volcano plots show examples of two differentially bound TFS (CTCF, MBD2) overlapping the promoters of DEGs in CTCF mutant expressing cells (ID). The remaining mutant comparisons are shown in **Figure 5E**. The enrichment of the TF target genes among the up- and down-regulated genes are reported on top on the volcanos (Odds Ratios (ORs), and p-values). Enrichment for all the differentially bound TFs and their differentially expressed target genes are reported in **Table S4B** and **S4C**. This figure relates to **Figure 5**.

Figure S21

Figure S21: **CTCF mutations alter insulation and chromatin interactivity.** (**A**) Absolute Hi-C matrices of chromosome 1 using a 100kb resolution for IAA and 3 mutants (bottom left) versus WT (top right) showing the minimal changes in chromatin compartmentalization. The compartment scores are shown on the left for the mutant and top for WT. (**B**) Absolute Hi-C matrix for IAA and the mutant shown in Figure **6E** compared to WT. The insulation scores are shown on the left for the mutant and top for WT. (**C**) Profiles comparing the insulation score in WT-IAA versus mutant-ID at lost CTCF sites upon IAA degradation. This figure relates to **Figure 6**.

Figure S22

Figure S22: **Insulation is a function of CTCF and ATAC-seq signal strength** . (**A**) Profiles show the aggregated insulation score at WT only, common and mutant only binding sites for four mutants. The profiles for the remaining mutants are shown in **Figure 6F.** (**B**) Profiles comparing the insulation score in mutant-ID versus IAA conditions at WT only sites. (**C**) Profiles show the aggregated insulation score at CTCF binding sites with weak, intermediate and strong CTCF signals for different mutants. The profiles for the remaining mutants are shown in **Figure 6G.** (**D**) Profiles show the aggregated insulation score at CTCF binding sites with no, weak, intermediate and strong ATAC signals for different mutants. The profiles for the remaining mutants are shown in **Figure 6H.** For (**C**) and (**D**)**,** Kruskal-Wallis p-values were coded as follow: * $5x10^{-2}$-$5x10^{-3}$, ** $5x10^{-3}$-$5x10^{-4}$, *** $5x10^{-4}$-$5x10^{-5}$, **** $<5x10^{-5}$. This figure relates to **Figure 6.**

Figure S23: Differentially expressed genes are enriched in gain of loops with a CTCF binding site at both anchors. Bar graphs showing the enrichment (calculated over WT ID) of over-expressed (top) or under-expressed (bottom) genes in gained (top) or lost (bottom) loops with both anchors overlapping a CTCF binding sites and at least one anchor overlapping the promoter of DEGs in IAA and CTCF mutants. This figure relates to **Figure 7.**

Figure S24

Figure S24: Indirect effects resulting from changes in TF binding at gene promoters account for the majority of CTCF mutation-mediated changes in expression. (**A**) Example of a locus on chromosome 7 (red rectangles) with a direct effect of gain in CTCF binding and chromatin interactivity. The left panel shows the Hi-C interaction matrices in WT, Q418R and K365T with both gain of intra- and inter-TAD interactions in the mutant compared to WT. The left panels show the zoom-in tracks of CTCF (blue) SMC3 (green), ATAC (red) and significant differential chromatin loops (purple) with one anchor overlapping the differential CTCF binding sites. Overexpressed genes are highlighted in red. Of note, only gained interactions were detected at these loci. This analysis was performed at 10 kb resolution. (**B**) Bar graphs showing the percentage of UP and DOWN DEGs associated with differential TF/CTCF binding, differential loops or both, stratified by direct and indirect effect. A direct effect was defined as predicted differential CTCF binding at the gene promoter and/or differential loops anchoring the DEG promoters with a CTCF peak at either anchor. This figure relates to **Figure 7.**

REFERENCES

S1.   Cerami, E., Gao, J., Dogrusoz, U., Gross, B.E., Sumer, S.O., Aksoy, B.A., Jacobsen, A., Byrne, C.J., Heuer, M.L., Larsson, E., et al. (2012). The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. Cancer Discov *2*, 401-404. 10.1158/2159-8290.CD-12-0095.

S2.   Tate, J.G., Bamford, S., Jubb, H.C., Sondka, Z., Beare, D.M., Bindal, N., Boutselakis, H., Cole, C.G., Creatore, C., Dawson, E., et al. (2019). COSMIC: the Catalogue Of Somatic Mutations In Cancer. Nucleic Acids Res *47*, D941-D947. 10.1093/nar/gky1015.

S3.   Price, E., Fedida, L.M., Pugacheva, E.M., Ji, Y.J., Loukinov, D., and Lobanenkov, V.V. (2023). An updated catalog of CTCF variants associated with neurodevelopmental disorder phenotypes. Front Mol Neurosci *16*, 1185796. 10.3389/fnmol.2023.1185796.

S4.   Konrad, E.D.H., Nardini, N., Caliebe, A., Nagel, I., Young, D., Horvath, G., Santoro, S.L., Shuss, C., Ziegler, A., Bonneau, D., et al. (2019). CTCF variants in 39 individuals with a variable neurodevelopmental disorder broaden the mutational and clinical spectrum. Genet Med *21*, 2723-2733. 10.1038/s41436-019-0585-z.