Integrative analysis in head and neck cancer reveals distinct role of miRNome and methylome as tumour epigenetic drivers

Katarina Mandić[1]*, Nina Milutin Gašperov[2]*, Ksenija Božinović[2], Emil Dediol[3], Jure Krasić[4], Nino Sinčić[4,5,6], Magdalena Grce[2], Ivan Sabol[2]**, Anja Barešić[1]**

*/**Equally contributing authors

Division of Electronics, Ruđer Bošković Institute, Zagreb, Croatia

Division of Molecular Medicine, Ruđer Bošković Institute, Zagreb, Croatia

Clinical Hospital Dubrava, Department of Maxillofacial Surgery, Zagreb, Croatia

Department of Medical Biology, University of Zagreb School of Medicine, Zagreb, Croatia

Centre of Excellence in Reproductive and Regenerative Medicine, University of Zagreb School of Medicine, Zagreb, Croatia

Biomedical Research Center Šalata, University of Zagreb School of Medicine, Zagreb, Croatia

Corresponding author: Ivan Sabol, PhD; Division of Molecular Medicine, Ruđer Bošković Institute, Bijenička 54, 10000 Zagreb, Croatia; E-mail address: isabol@irb.hr; Phone:

+38514561110; Fax: +38514561010;

Abstract

Background: Head and neck cancer is the sixth most common malignancy worldwide, with

the relatively low 5-year survival rate, mainly because it is diagnosed at a late stage.

Infection with HPV is well known etiology, which affects nature of these cancers and patients'

survival. Besides, it is considered that the main driving force for this type of cancer could be

epigenetics. In this study we aimed to find potential epigenetic biomarkers, by integrating

miRNome, methylome, and transcriptome analyses.

Methods: From the fresh head and neck cancer tissue samples, we chose a group for

miRNome, methylome and transcriptome profiling, in comparison to adequate control

samples. Bioinformatics analyses are performed in R v4.2.2. Count normalization and group

differential expression for mRNA and the previously obtained miRNA count data was

performed with DESeq2 v1.36. Gene set enrichment analysis was performed and visualized

using gProfiler2 v0.2.1 Identification of miRNA targets was performed by querying in

miRTarBase using multiMiR v1.18.0. Annotation of CpG sites merging into islands was

obtained from RnBeads.hg19 v1.28.0. package. For the integrative analysis we performed

kmeans clustering using stats v4.2.2 package, using 8-12 clusters and nstart 100.

Results: We found that transcriptome analysis divides samples into cancers and controls

clusters, with no relation to HPV status or cancer anatomical location. Differently expressed

genes (n=2781) were predominantly associated with signaling pathways of tumour

progression. We identified a cluster of genes under control of the transcription factor E2F,

that are significantly underexpressed in cancer tissue, as well as T cell immunity genes and

genes related to regulation of transcription. Among overexpressed genes in tumours we

found those that belong to cell cycle regulation and vasculature. A small number of genes

were found significantly differently expressed in HPV positive *versus* HPV negative tumours

(for example NEFH, ZFR2, TAF7L, ZNF541, and TYMS).

Conclusions: In this comprehensive study on an overlapping set of samples where the

integration of miRNome, methylome and transcriptome analysis were performed for head

and neck cancer, we demonstrated that the majority of genes were associated exclusively

with miRNome or methylome and, to a lesser extent, under control of both epigenetic

mechanisms.

Keywords: head and neck cancer, HPV, miRNome, methylome, transcriptome, integrative

analysis

Background

Head and neck cancer is the sixth most common malignancy worldwide, with head and neck

squamous cell carcinoma (HNSCC) accounting for more than 90% of cases. Unfortunately,

HNSCC often gets diagnosed in a late phase when it is challenging to treat the tumour, which

contributes to poor 5-year survival rate, currently at 66% (Johnson et al. 2020). Even though

this type of cancer arises from a single cell type, i.e. the squamous cell, HNSCC is surprisingly

of very heterogeneous nature (Canning et al. 2019). Primary site of tumour origin

contributes to tumour heterogeneity, with most common sites being oral cavity, oropharynx,

larynx, and sinonasal tract (Yan et al. 2011). Besides tumour location, the infection with

high-risk human papilloma virus (HPV) types, especially in oropharyngeal area predominately

affects HNSCC nature and patients' survival (Kobayashi et al. 2018). HPV positive and HPV

negative HNSCC can substantially differ in terms of etiology, genetics and epigenetics

(Joseph and D'Souza 2012; Johnson et al. 2020). HPV-positive HNSCC mostly affect younger

population, with no or low levels of smoking and/or alcohol use (Gillison et al. 2015).

Moreover, unlike the HPV-negative group, preferable site of tumour origin for HPV-positive

HNSCC is oropharynx and it has been reported in many Western countries that this group of

patients is associated with better overall survival and better response to therapy (Marur et

al. 2010). There has been significant shift in the incidence of HPV-positive oropharyngeal

squamous cancer in Western countries, where HPV is associated to HNSCC in almost 70% of

cases (Marur and Forastiere 2016), and in fact, increase in 5-year survival of overall HNSCC

cases could be contributed to higher prevalence of HPV-associated HNSCC (Johnson et al.

2020).

Up to date, many studies reported better outcomes for HPV-positive HNSCC (Perri et al.

2020; Kofler et al. 2014; Sun et al. 2021), however the consensus on specific biomarkers,

which could improve diagnostic, prognostic and/or therapeutic approaches is still missing.

Lack of consensus biomarkers could partially be owed to differences in studied populations

sometimes inadequately registered tumour site of origin and possibly to the lack of

stratification by the HPV status. Therefore, we consider the cohort presented below to be a

specific case of tumours in terms of lifestyle, i.e. higher smoking and alcohol intake rates.

In this study, we aimed to identify epigenetic patterns in HNSCC since the main driving force

for the HNSCC development is considered to be epigenetics, rather than genetics. We have

previously analyzed the content and transcriptional levels of whole genome microRNA

(miRNA) profile (miRNome) in HPV-positive and HPV-negative oral and oropharyngeal

HNSCC samples using next generation sequencing and validated with qRT-PCR. We have also

analyzed the whole genome methylation profile (methylome) in the same set of samples

using the whole genome methylation array and validated by pyrosequencing. In this study,

we assessed the whole genome transcriptome analysis on the same sample set and

combined the mRNA, miRNA and methylation information to elucidate relevant epigenetic

mechanisms and their interactions. Motivated by the limited literature regarding integrative

studies on epigenetic changes, we set out to elucidate whether miRNAs or DNA methylation

have more prominent impact on HNSCC development. Moreover, we aimed to identify key

mechanisms and/or cellular signaling pathways of genes that have been differentially

expressed in HNSCC compared to healthy controls. Consequently, we provide a better

understanding of the epigenetic influence on the development and progression of HPV-

positive and HPV-negative HNSCC, shedding light on mechanisms underlying prognostic,

diagnostic and therapeutic strategies. In addition, integrative analysis points out key target

genes and signaling pathways deregulated in HNSCC in Croatian and similar populations.

Material and Methods

Patient material

A total of 61 fresh oropharyngeal (OP) and oral (O) HNSCC primary tumour samples have

been obtained from Dubrava Clinical Hospital in the 2015-2018 time period. Twenty-two of

them and three non-malignant tonsil control samples were included in the miRNA profiling

(Božinović et al. 2019), column 2 in **Table 1**. Of 22 sequenced samples, two were excluded,

due to one sample being a recurrent tumour and one outlier. miRNome analysis and

detailed patients' socioepidemiologic and clinical data have been described in our previous

study . From the previously published analysis of HNSCC methylome (Milutin Gašperov et al.

2020), of 22 samples included in miRNA study, 12 HNSCC samples were included in the

whole genome methylation profiling (column 3, **Table 1**), as well as one additional sample

that had methylome but no transcriptome and miRNome profile. Eight healthy swabs of oral

mucosa were included in methylome profiling, which served as control samples .

For the transcriptome profiling, we selected 10 fresh HNSCC samples (6 oral and 4

oropharyngeal) with previous miRNome and methylome data (column 1, **Table 1**). Three

fresh tonsil tissues from the miRNome dataset were included in transcriptome profiling as

control tissue. HPV status was assessed by PCR in all samples as described previously .

Table 1. Samples used in integrative analysis. Sequenced samples are shown in white. O=oral,
OP=oropharyngeal, P= HPV-positive, N=HPV-negative.

Transcriptome profiling

As stated above, 13 samples were selected for transcriptome analysis in the current study to

complement previously obtained omics results. Sample annotation for samples used in

transcriptome analysis and subsequent integrative epigenetic study is presented in **Table 1**.

Transcriptome was assessed using TrueSeq Stranded mRNA (Illumina) kit for library

preparation, followed by sequencing on Nextseq500 instrument using NextSeq HIGH 150

cycles (Illumina) reagents on high output flow cell in paired end mode. Debarcoding was

performed using Basespace (Illumina) platform. Quality control of sequencing data was

performed with FastQC v0.11.9. Sequences were aligned to the human reference genome

GRCh37 downloaded from GENCODE (released in 2013) and quantified using salmon v1.5.0,

(Patro et al. 2017) reporting estimated fragment counts per transcript.

Bioinformatics preprocessing

All bioinformatics analyses listed hereafter were performed in R v4.2.2, and refer to R

language packages, unless otherwise stated. The preprocessing consisted of transcriptome

gene count determination, reprocessing of miRNA counts (for a subset of samples from

Božinović 2019) and reprocessing of differentially methylated CpGs (for a subset of samples

from Milutin Gašperov 2020), here expanded to CpG islands.

Count normalization and group differential expression for mRNA and the previously

obtained miRNA count data was performed with DESeq2 v1.36. Pre-filtering of the data set

was performed to reduce genes that have nearly no information about gene expression by

filtering out genes and miRNAs with total counts across all samples less than 10. Count

variance stabilization transformation was performed with regularized logarithm

transformation (DESeq2 rlog function) and used as input to calculate principal components

with plotPCA function from DESeq2 and Euclidean sample distances with dist function from

the stats v4.2.2 package. Counts normalization was performed with DESeq function with log

fold change shrinkage. We calculated differential expression in tumour *versus* control and

HPV positive *versus* negative samples (**Table 1**). Differentially expressed genes (DEGs) and

miRNA (DEmiR) were selected with BH-adjusted $p < 0.05$ set as significant.

Gene set enrichment analysis (GSEA) was performed and visualized using gProfiler2 v0.2.1

on DEGs using annotated human genes as the background set for the hypergeometric test.

We applied the tailor-made multiple test correction method Set Counts and Sizes (SCS)

(Raudvere et al. 2019) and a p-value threshold of 0.05. GSEA was performed across multiple

databases: Gene Ontology: Biological Processes (GO:BP) (Ashburner et al. 2000), Kyoto

Encyclopedia of Genes and Genomes (KEGG) (Kanehisa et al. 2019), Reactome (Fabregat et

al. 2018), miRTarBase (Chou et al. 2018) and TRANSFAC (Matys et al. 2006). We used these

parameters for all subsequent GSEA.

Identification of miRNA targets was performed by querying significant DEGs against DEmiRs

using multiMiR v1.18.0 and the miRTarBase database (Huang et al. 2022), selecting only

experimentally validated miRNA-gene interactions.

Previously published DNA methylation (DNAm) data (Milutin Gašperov et al. 2020) was

loaded with RnBeads v2.14.0. Differential methylation (DM) was calculated comparing

tumour *versus* control groups on CpG sites and islands (CGIs). Annotation of CpG sites

merging into islands was obtained from RnBeads.hg19 v1.28.0 package. All results are

presented on CGIs unless otherwise stated. CGI DM was defined as the quotient of

methylation between groups across all CpG sites in a CGI, with significance level set to FDR-

adjusted p<0.05. Promoter region annotations were obtained from the RnBeads package,

defined as 1500 bases upstream and 500 bases downstream of the transcription start site.

CGI targets were assigned where CGI overlapped a promoter region.

Integrative analysis

Workflow for comparing all omics data and identifying specific mechanisms in HNSCC is

summarized in **Figure 1**.

Integrative epigenetic analysis of HNSCC

transcriptome

Fresh 6 OSCC (3 HPV+ and 3 HPV-)
Fresh 4 OPSCC (2 HPV+ and 2 HPV-)
Fresh 3 healthy OP controls (HPV-)

Illumina TrueSeq stranded mRNA kit
BaseSpace platform
Salmon tool, DESeq2 (R)

Exploratory analysis, PCA,
differential expression,
gene set enrichment

miRnome

Fresh 6 OSCC (3 HPV+ and 3 HPV-)
Fresh 4 OPSCC (2 HPV+ and 2 HPV-)
Fresh 3 healthy OP controls (HPV-)

Illumina TrueSeq Small RNA
Library prep kit
NextSeq 500 Mid output kit

BaseSpace platform, DESeq2 (R)

Exploratory analysis, PCA, differential expression,
gene set enrichment, miRTarBase

methylome

Fresh 6 OSCC (3 HPV+ and 3 HPV-)
Fresh 4 OPSCC (2 HPV+ and 2 HPV-)
Fresh 8 healthy O controls (HPV-)

Illumina Infinium
MethylationEPIC BeadChip array

RnBeads (R), CpG islands

Exploratory analysis, PCA,
differential methylation, gene set
enrichment

Figure 1. Schematic representation of integrative epigenetic HNSCC study.

Three -omics datasets were compared, centered on identified DEGs (**Supplemental table S1**).

Multiple miRNA-gene pairs were reduced by selecting only the miRNA with highest absolute

value of Spearman correlation of counts (mirRNA_Rho) for the given DEG. Multiple gene–CGI

pairs were not as common, and the CGI with the lowest p-value was selected for every DEG.

Resulting table contained DEGs and their associated miRNA and/or CGI. We assessed the

interplay of DEmiRs, DM CGIs and DEGs in two ways: by obtaining simple overlap of DEGs

targeted by DEmiRS or DM CGIs and kmeans clustering. The simple overlap was based on

whether DEG's expression levels were associated with DEmiR's activity and/or CGI

methylation levels. Kmeans clustering was performed on log-transformed, centered and

scaled values of fold change for DEGs and DEmiRs, and DNAm quotient. We performed

kmeans clustering using stats v4.2.2 package, using 8-12 clusters and nstart 100, with the

best performing result for 11 clusters.

Results

Transcriptome profiling

The whole transcriptome profiling was performed on tissue samples from 13 individuals with

different anatomical locations and HPV infection status (**Table 1**). The top two principal

components of the transcriptome (**Figure 2a**), which explain 55% of the variance in the

samples, splits samples into two clusters of tumour and control samples, with no obvious

contribution of HPV status or tumour anatomical location. We observe the same pattern in

miRNome and methylome data (**Figure 2b** and **c**).

Figure 2. Exploratory analysis of three omics data sets: a) PCA plot for transcriptome data; b)
sample distance heatmap for miRNome data; c) sample distance heatmap for methylome. N
- HPV-negative, P - HPV-positive O - oral tumours, OP - oropharyngeal tumours.

Differential expression and GSEA

Exactly 18104 genes had no less than 10 detected reads across samples. Differential

expression analysis revealed that 1248 (6.9%) of genes were significantly overexpressed and

1533 (8.5%) were significantly underexpressed in tumour (BH-adjusted p<0.05), when

compared to transcriptomes of control samples. GSEA on DEGs showed enrichment of

signaling pathways associated with tumour progression, such as cell cycle regulation and

oncogenic signaling pathways such as NF-kB (**Figure 3**), for a full list of terms see

Supplemental table S2.

id	source	term id	term name	term size	p value
1	GO:BP	GO:0032502	developmental process	4582	2.0e-09
2	GO:BP	GO:0070661	leukocyte proliferation	235	1.8e-04
3	KEGG	KEGG:04064	NF-kappa B signaling pathway	102	1.6e-02
4	REAC	REAC:R-HSA-453276	Regulation of mitotic cell cycle	88	2.4e-02
5	REAC	REAC:R-HSA-8852276	The role of GTSE1 in G2/M progression after G2 checkpoint	79	7.0e-04

Figure 3. Gene enrichment analysis of tumour *vs.* control samples. Databases shown are GO:BP (orange), KEGG (pink), and REACTOME (REAC, blue). The highlighted terms correspond to key concepts associated with cancer progression and regulation.

Differential miRNA and DNA methylation assessment

Integration of 514 significant DEmiRs based on MiRNome data (Božinović et al. 2019) with the database containing experimentally supported miRNA-target gene interactions yielded 3042 interactions among 90 different miRNAs and 1227 target genes, hereafter termed DEmiR-DEGs. In our analysis Most of these genes were regulated by a single miRNA, however some were associated with multiple miRNAs. Notably, two genes with the highest number of associated DEmiRs, *BCL2* and *MYC*, which play a role in regulating cell proliferation and apoptosis, and were associated with 18 and 16 miRNAs, respectively. In addition, some miRNAs were associated with many target genes, harboring potential therapeutic value by potentially affecting many cancer-related signaling pathways. An extreme example is a frequently reported tumour suppressor hsa-miR-26b-5p putatively

forming interactions with 255 unique DEGs. Within our dataset hsa-miR-26b-5p showed a

high negative expression correlation with known oncogenes such as *MMP10* (Rho= -0.81,

p<0.005), *IGSF3* (Rho= -0.79, p<0.005) and *ARNTL2* (Rho= -0.79, p<0.005) (**Supplemental**

table S1).

Overall methylation levels in our samples showed that more than 300,000 CpG sites were

hypomethylated in tumour samples compared to controls (FDR-adjusted p<0.05), while

72,000 were hypermethylated (**Supplemental table S3**). We also noted a larger number of

hypomethylated CGIs (6557) than hypermethylated (5457), although the difference is not as

pronounced as in CpG sites (**Supplemental table S4**). The DM CpGs predominantly fell into

gene bodies, but also some fell in intergenic regions corresponding to repeated elements.

These regions consistently showed a larger number of hypomethylated CpGs compared to

hypermethylated ones (**Supplemental figure S1**).

To identify deregulated tumour suppressors in our samples, we used the TSGene database

2.0 (Zhao et al. 2016) containing tumour suppressors curated from several thousand

publications combined with expression and mutational profiles. We pinpointed 10 genes

that were underexpressed and contained hypermethylated CGIs in the promoter region in

our samples (**Supplemental table S5**).

To assess the effects of transcription factors on DEGs that had differentially methylated CGIs

(DM-DEGS) within our dataset we performed GSEA using transcription factor database

TRANSFAC. We found associations with several groups of transcription factors including E2F,

SP/KLF and AP-2 families (**Supplemental table S6**).

Of note we found the enrichment of genes regulated by the E2F family of TFs (SCS-adjusted

$p<10^{-43}$), which play a critical role in cell cycle control. Furthermore, GSEA on the same

dataset using the Reactome database showed enrichment of genes involved in cell cycle

regulation such as G1/S Transition and G2/M Checkpoints (SCS-adjusted $p<10^{-4}$), supporting

the above findings.

Integrative analysis: Influence of miRNome and methylome on transcription on

transcriptional activity in HNSCC

Each DEG was grouped based on the association with DEmiR or DM CGI, shown in **Figure 4**.

We obtained four groups: DEGs associated only with DEmiR (DEmiR-DEGs), DEGs associated

only with DM CGIs (DM-DEGs), DEGs associated with both DEmiR and DM (DEmiR+DM-

DEGs), DEGs not associated with either DEmiR or DM CGIs (noReg-DEGs). The groups had

different proportion of differentially represented GSEA terms, i.e. GO:BP, KEGG, miRTarBase

(MIRNA), Reactome and TRANSFAC (TF) (**Figure 4**, **Supplemental table S7**). The DM-DEGs

group were enriched by genes regulated by TFs but no significant results of GO biological

pathways was detected. The DEmiR-DEGs group were enriched with pathways related to

signaling and immunological systems. The noReg-DEGs group was enriched with immunological

processes and muscle system processes. DEmiR+DM-DEGs were enriched with general regulatory

processes: biological, metabolic and cellular processes. These terms are high level terms in GO

biological processes, suggesting that genes in the intersection have broad regulatory roles.

Figure 4. DEGs between tumour and control samples, in association with DEmiR or DM
promoters. Four panels show number of terms from GSEA for each group.

Clustering of DEGs with accompanying DEmiR and DM levels revealed a critical distinction in

the epigenetic factors driving HNSCC. The identified clusters predominantly demonstrated

either significant DEmiRs or DM. This suggests that in HNSCC, gene expression is

predominantly influenced by either miRNA regulation or methylation, but not both

concurrently. This key finding illuminates the mutually exclusive roles of miRNA and

methylation in shaping the gene expression landscape in HNSCC.

It is generally believed that miRNA and DNAm have an inhibitory effect on gene expression,

and this pattern was evident in clusters 3, 4, 7 and 10. In addition, we observed miRNA

expression and methylation patterns in clusters 1, 5, 6, and 9 that were indicative of

noncanonical regulatory activity. It is important to mention that due to multiple target genes

of miRNAs there was a significant overlap of underexpressed miRNAs in clusters 3 and 9

(73%) and overexpressed miRNAs in clusters 5 and 7 (77%).

GSEA analysis of each cluster linked expression and epigenetic patterns to biological function.

Representative terms that best described each cluster are shown in **Figure 5** and the full

GSEA results for each cluster are shown in **Supplemental table S7**. DM-DEGs in clusters 1, 4,

6 and 10 were highly enriched with genes regulated by the E2F family of TFs. Most of the

genes from the E2F family were not differentially expressed in our samples (**Supplemental**

figure S2), although it has been previously shown that the methylation status of CpG sites of

E2F binding motif influences the binding of the transcription factor (Campanero, Armstrong,

and Flemington 2000; Yin et al. 2017). In addition, cluster 6 contained underexpressed genes

with hypomethylated CGIs and was enriched with genes regulated by the ZF5 transcription

factor, which can function as a repressor (Numoto, Yokoro, and Koshi 1999). Clusters 4 and 5

were associated with similar terms related to various cell cycle processes but have different

patterns of miRNA expression and DNAm.

Figure 5. Kmeans clustering on integrated transcriptome, miRNome and methylome data on

DEGs. Fold change values for transcriptome and miRNome and DNAm quotient were scaled

and normalized prior to clustering. Eleven clusters are functionally described by applying

GSEA over several databases (GO:BP, KEGG, REACTOME, MIRNA and TRANSFAC).

Influence of HPV

HPV influence on gene expression was found to be inconclusive (**Figure 2**). We observed a

small number of genes and miRNAs, and no DM CGIs or sites that were significantly changed

while comparing HPV positive and negative tumour sample groups (FDR-adjusted p<0.1).

Number of DEGs associated with HPV status was 44, of which 25 were overexpressed

(**Supplemental table S9** and **S10).** Among these, we focused on the selection of DEGs found

to be functionally relevant and/or previously found to be differentially expressed in HPV-

associated cancers (**Figure 6**).

Figure 6. Top 20 DEGs in HPV positive compared to HPV negative tumour patients.

Discussion

In this study, we aimed to integrate miRNome and methylome analysis with transcriptome

analysis on the same samples to elucidate whether miRNAs or DNAm have more prominent

impact on HNSCC development. The main goal was to identify key mechanisms and/or

cellular signaling pathways of DEGs in HNSCC compared to healthy control samples.

Herein, we observed that transcriptome analysis divided samples into two major clusters:

tumour and control samples, with no obvious contribution of HPV status or tumour

anatomical location. Similar results were seen in the miRNA expression profile study and

DNAm study in an overlapping set of samples (**Figure 2 b** and **c**) as previously published

(Božinović et al. 2019; Milutin Gašperov et al. 2020). This suggests that these tumours may

share similar cells of origin or common pathways during tumourigenesis.

The emergence of tumour progression pathways among DEGs (n=2781, **Figure 3**) resembles

transcriptomes of radioresistant HNSCCs, where hypoxia response, p53 pathway, NF-kB

pathway and inflammatory response were found abnormally activated. Interestingly, some

authors found that NF-kB and other transcription factors signatures differentiate HPV-

positive and HPV-negative HNSCCs.

Using unsupervised clustering, we found a cluster of genes under control of the transcription

factor E2F, that were significantly underexpressed in tumour samples in comparison to

controls (**Figure 5**, cluster 6). This is in line with overall hypomethylation in tumours and

points to exclusive control of overexpressed miRNAs. The genome-wide hypomethylation of

CpG sites is a known phenomenon that occurs in most cancers (Gama-Sosa et al. 1983;

Debernardi et al. 2021). E2F family members expression was also found to be negatively

regulated by miRNAs in other cancers, like lung cancers, pediatric brain tumours, and

chronic myeloid leukemia. In addition, genes that were significantly associated with the E2F

were found in clusters 1, 5 and 10 (**Supplemental table S6**).

In cluster 8, we found genes related to T cell immunity underexpressed in tumour samples in

comparison to control samples (**Figure 5**), mostly found to be targets of overexpressed

miRNAs, and in less extent associated with hypermethylation. This finding is in line with

trends detected in several previous methylation studies of head and neck cancers (Moody et

al. 2020; Anić et al. 2023), and in cervical cancers (Farkas et al. 2013; Milutin Gašperov et al.

2014).

Among underexpressed genes in tumours, we observed a cluster of DM-DEGs related to

regulation of transcription (**Figure 5**, cluster 10). Our result showed that the cluster mostly

contains genes with hypermethylation of CGIs in promoter regions and some genes that

were targeted by overexpressed miRNAs. This implies that the hypermethylation of CGIs in

promoter region of genes in this cluster may have disrupted binding of transcription factors

required for their expression, that is a well-studied phenomena (Eden and Cedar 1994;

Pikaart, Recillas-Targa, and Felsenfeld 1998; Bird 2002).

Among overexpressed genes in tumours were those that belong to cell cycle regulation

(**Figure 5**, clusters 4 and 5) and vasculature (**Figure 5**, cluster 3). Herein, we found cell cycle

genes have mostly hypomethylated promoter regions. On the other hand, our result showed

that genes involved in vascularization in tumours are DEmiR+DM-DEGs, likely influenced by

both epigenetic drivers. Moreover, other authors found a link with DNAm and cardiovascular

diseases directly rather than changes driven by miRNA expression (Goossens et al. 2019), or

that both epigenetic changes are involved in vascular diseases, claiming that, the most

prominent gene cluster is activated via hypomethylation (Aavik et al. 2015). Interestingly,

the same authors found keratin mRNAs upregulated in atherosclerosis and concluded that

DNA hypomethylation may play a role in keratin gene activation and ectopic keratin

expression (Aavik et al. 2015). Keratins are characteristic to epithelial cells and are not

expressed in normal arteries and induction of keratin expression in atherosclerosis has been

detected even earlier (Slomp et al. 1997). Our study detected similar findings in HNSCC;

keratin mRNAs were found upregulated, that was partially under control of DNAm and

partially under control of miRNome (**Figure 5**, cluster 2).

Herein, we found that cluster 5 (**Figure 5**) contained upregulated miRNA and their

corresponding upregulated target genes. Conversely, cluster 9 displayed downregulated

miRNA and corresponding downregulated target genes. This positive correlation of miRNAs

and target genes has been previously observed in multiple studies (Tan et al. 2019; Xu et al.

2020). While our results could imply either a direct or indirect effect of miRNA on DEmiR-

DEGs potentially through negating the action of repressive miRNA, protein complexes (Tan

et al. 2019), it is important to note that our findings represent correlation, not causation.

Further studies are required to establish the causal mechanisms involved. In addition, it is

worth considering that miRNAs often have multiple targets, and the interactions in existing

databases may be derived from high-throughput methods that may not reflect the full

complexity of in vivo biological interactions.

A small number of genes were found significantly differentially expressed in HPV positive

versus HPV negative tumours (**Figure 6**) in our study, a subset of which was previously found

differentially expressed in head and neck cancer and related to HPV status, i.e. *NEFH*, *ZRF2*,

TAF7L, ZNF541 and *TYMS*. Notably, some authors have already found hypermethylation of

NEFH (protein kinase binding and microtubule binding), in pharyngeal squamous cell

carcinoma, which significantly correlated with HPV positivity (Nakagawa et al. 2017). *ZFR2*

gene codes for RNA-binding protein characterized by its domain associated with zinc fingers

and zinc ions. It was found overexpressed in HPV-positive HNSCC patients when compared

to HPV-negative patients and has been flagged as prognostic for the cervical cancer cases

(Tripathi et al. 2020). The expression of the transcription factor *TAF7L* (TAF7-like RNA

polymerase II) has been already found increased, even 220-fold, in HPV positive HNSCCs

(Slebos et al. 2006). *ZNF541* encodes a zinc finger protein supposed to be a component of

chromatin remodeling complexes. The hypomethylation and higher expression of this gene

was also observed in HPV-positive oropharyngeal squamous cell carcinoma (OPSCC) and

significantly associated with a better overall survival, independent of HPV status (Camuzi et

al. 2021). *TYMS* gene codes for thymidylate synthetase that catalyzes the methylation of

deoxyuridylate   to   deoxythymidylate.   In   HPV-positive   cancers   they   have   found   it

overexpressed, indicating that HPV-positive oropharyngeal cancers may be more resistant to

5-fluorouracil chemotherapy than HPV-negative (Lohavanichbutr et al. 2009).

To our knowledge, we presented the most comprehensive study on an overlapping set of

samples where the integration of miRNome, methylome and transcriptome analyses in

HNSCC was conducted. We acknowledge certain limitations of our study, including the

relatively small sample size, which may affect the validity of our findings. In addition,

obtaining control samples that are genuinely representative of healthy tissue poses its own

set of challenges. Associating miRNome and methylome with transcriptome profiles, we

found many DEGs in HNSCC in comparison to control samples, that were mainly associated

with cell cycle regulation, transcription, immunity and tumour development. We

demonstrated the majority of genes belong to clusters associated exclusively with miRNome

or methylome and, to a lesser extent, under control of both epigenetic mechanisms.

408 References

Aavik, Einari, Henri Lumivuori, Olli Leppänen, Thomas Wirth, Sanna-Kaisa Häkkinen, Jan-
Hinrich Bräsen, Ulrich Beschorner, et al. 2015. 'Global DNA Methylation Analysis of
Human Atherosclerotic Plaques Reveals Extensive Genomic Hypomethylation and
Reactivation at Imprinted Locus 14q32 Involving Induction of a miRNA Cluster'.
European Heart Journal 36 (16): 993–1000.
https://doi.org/10.1093/eurheartj/ehu437.
Anić, Petra, Jasminka Golubić Talić, Ksenija Božinović, Emil Dediol, Marinka Mravak-Stipetić,
Magdalena Grce, and Nina Milutin Gašperov. 2023. 'Methylation of Immune Gene
Promoters in Oral and Oropharyngeal Cancer'. *International Journal of Molecular
Sciences* 24 (9): 7698. https://doi.org/10.3390/ijms24097698.
Ashburner, M., C. A. Ball, J. A. Blake, D. Botstein, H. Butler, J. M. Cherry, A. P. Davis, et al.
2000. 'Gene Ontology: Tool for the Unification of Biology. The Gene Ontology
Consortium'. *Nature Genetics* 25 (1): 25–29. https://doi.org/10.1038/75556.
Bird, Adrian. 2002. 'DNA Methylation Patterns and Epigenetic Memory'. *Genes &
Development* 16 (1): 6–21. https://doi.org/10.1101/gad.947102.
Božinović, Ksenija, Ivan Sabol, Emil Dediol, Nina Milutin Gašperov, Spomenka Manojlović,
Zuzana Vojtechova, Ruth Tachezy, and Magdalena Grce. 2019. 'Genome-Wide
miRNA Profiling Reinforces the Importance of miR-9 in Human Papillomavirus
Associated Oral and Oropharyngeal Head and Neck Cancer'. *Scientific Reports* 9 (1):
2306. https://doi.org/10.1038/s41598-019-38797-z.

Campanero, Miguel R., Monica I. Armstrong, and Erik K. Flemington. 2000. 'CpG Methylation
as a Mechanism for the Regulation of E2F Activity'. *Proceedings of the National*
Academy of Sciences 97 (12): 6481–86. https://doi.org/10.1073/pnas.100340697.
Camuzi, Diego, Luisa Aguirre Buexm, Simone de Queiroz Chaves Lourenço, Davide Degli
Esposti, Cyrille Cuenin, Monique de Souza Almeida Lopes, Francesca Manara, et al.
2021. 'HPV Infection Leaves a DNA Methylation Signature in Oropharyngeal Cancer
Affecting Both Coding Genes and Transposable Elements'. *Cancers* 13 (14): 3621.
https://doi.org/10.3390/cancers13143621.
Canning, Madison, Gang Guo, Miao Yu, Calvin Myint, Michael W. Groves, James Kenneth
Byrd, and Yan Cui. 2019. 'Heterogeneity of the Head and Neck Squamous Cell
Carcinoma Immune Landscape and Its Impact on Immunotherapy'. *Frontiers in Cell*
and Developmental Biology 7 (April): 52. https://doi.org/10.3389/fcell.2019.00052.
Chou, Chih-Hung, Sirjana Shrestha, Chi-Dung Yang, Nai-Wen Chang, Yu-Ling Lin, Kuang-Wen
Liao, Wei-Chi Huang, et al. 2018. 'miRTarBase Update 2018: A Resource for
Experimentally Validated microRNA-Target Interactions'. *Nucleic Acids Research* 46
(D1): D296–302. https://doi.org/10.1093/nar/gkx1067.
Debernardi, Carla, Laura Libera, Enrico Berrino, Nora Sahnane, Anna Maria Chiaravalli,
Cristiana Laudi, Mattia Berselli, et al. 2021. 'Evaluation of Global and Intragenic
Hypomethylation in Colorectal Adenomas Improves Patient Stratification and
Colorectal Cancer Risk Prediction'. *Clinical Epigenetics* 13 (1): 154.
https://doi.org/10.1186/s13148-021-01135-0.
Eden, Sharon, and Howard Cedar. 1994. 'Role of DNA Methylation in the Regulation of
Transcription'. *Current Opinion in Genetics & Development* 4 (2): 255–59.
https://doi.org/10.1016/S0959-437X(05)80052-8.
Fabregat, Antonio, Steven Jupe, Lisa Matthews, Konstantinos Sidiropoulos, Marc Gillespie,
Phani Garapati, Robin Haw, et al. 2018. 'The Reactome Pathway Knowledgebase'.
Nucleic Acids Research 46 (D1): D649–55. https://doi.org/10.1093/nar/gkx1132.
Farkas, Sanja A, Nina Milutin-Gašperov, Magdalena Grce, and Torbjörn K Nilsson. 2013.
'Genome-Wide DNA Methylation Assay Reveals Novel Candidate Biomarker Genes in
Cervical Cancer'. *Epigenetics* 8 (11): 1213–25. https://doi.org/10.4161/epi.26346.
Gama-Sosa, M. A., V. A. Slagel, R. W. Trewyn, R. Oxenhandler, K. C. Kuo, C. W. Gehrke, and M.
Ehrlich. 1983. 'The 5-Methylcytosine Content of DNA from Human Tumors'. *Nucleic*
Acids Research 11 (19): 6883–94. https://doi.org/10.1093/nar/11.19.6883.
Gillison, Maura L., Anil K. Chaturvedi, William F. Anderson, and Carole Fakhry. 2015.
'Epidemiology of Human Papillomavirus-Positive Head and Neck Squamous Cell
Carcinoma'. *Journal of Clinical Oncology: Official Journal of the American Society of*
Clinical Oncology 33 (29): 3235–42. https://doi.org/10.1200/JCO.2015.61.6995.
Goossens, Eveline A. C., Margreet R. de Vries, Karin H. Simons, Hein Putter, Paul H. A. Quax,
and A. Yaël Nossent. 2019. 'miRMap: Profiling 14q32 microRNA Expression and DNA
Methylation Throughout the Human Vasculature'. *Frontiers in Cardiovascular*
Medicine 6 (August): 113. https://doi.org/10.3389/fcvm.2019.00113.
Huang, Hsi-Yuan, Yang-Chi-Dung Lin, Shidong Cui, Yixian Huang, Yun Tang, Jiatong Xu,
Jiayang Bao, et al. 2022. 'miRTarBase Update 2022: An Informative Resource for
Experimentally Validated miRNA-Target Interactions'. *Nucleic Acids Research* 50 (D1):
D222–30. https://doi.org/10.1093/nar/gkab1079.
Johnson, Daniel E., Barbara Burtness, C. René Leemans, Vivian Wai Yan Lui, Julie E. Bauman,
and Jennifer R. Grandis. 2020. 'Head and Neck Squamous Cell Carcinoma'. *Nature*
Reviews Disease Primers 6 (1): 1–22. https://doi.org/10.1038/s41572-020-00224-3.
Joseph, Andrew W., and Gypsyamber D'Souza. 2012. 'Epidemiology of Human

Papillomavirus-Related Head and Neck Cancer'. *Otolaryngologic Clinics of North*
America 45 (4): 739–64. https://doi.org/10.1016/j.otc.2012.04.003.

Kanehisa, Minoru, Yoko Sato, Miho Furumichi, Kanae Morishima, and Mao Tanabe. 2019.
'New Approach for Understanding Genome Variations in KEGG'. *Nucleic Acids*
Research 47 (D1): D590–95. https://doi.org/10.1093/nar/gky962.

Kobayashi, Kazuhiro, Kenji Hisamatsu, Natsuko Suzui, Akira Hara, Hiroyuki Tomita, and
Tatsuhiko Miyazaki. 2018. 'A Review of HPV-Related Head and Neck Cancer'. *Journal*
of Clinical Medicine 7 (9): 241. https://doi.org/10.3390/jcm7090241.

Kofler, B., S. Laban, C. J. Busch, B. Lörincz, and R. Knecht. 2014. 'New Treatment Strategies
for HPV-Positive Head and Neck Cancer'. *European Archives of Oto-Rhino-*
Laryngology: Official Journal of the European Federation of Oto-Rhino-Laryngological
Societies (EUFOS): Affiliated with the German Society for Oto-Rhino-Laryngology -
Head and Neck Surgery 271 (7): 1861–67. https://doi.org/10.1007/s00405-013-
2603-0.

Lohavanichbutr, Pawadee, John Houck, Wenhong Fan, Bevan Yueh, Eduardo Mendez, Neal
Futran, David R. Doody, et al. 2009. 'Genomewide Gene Expression Profiles of HPV-
Positive and HPV-Negative Oropharyngeal Cancer: Potential Implications for
Treatment Choices'. *Archives of Otolaryngology--Head & Neck Surgery* 135 (2): 180–
88. https://doi.org/10.1001/archoto.2008.540.

Marur, Shanthi, Gypsyamber D'Souza, William H. Westra, and Arlene A. Forastiere. 2010.
'HPV-Associated Head and Neck Cancer: A Virus-Related Cancer Epidemic – A Review
of Epidemiology, Biology, Virus Detection and Issues in Management'. *The Lancet.*
Oncology 11 (8): 781–89. https://doi.org/10.1016/S1470-2045(10)70017-6.

Marur, Shanthi, and Arlene A. Forastiere. 2016. 'Head and Neck Squamous Cell Carcinoma:
Update on Epidemiology, Diagnosis, and Treatment'. *Mayo Clinic Proceedings* 91 (3):
386–96. https://doi.org/10.1016/j.mayocp.2015.12.017.

Matys, V., O. V. Kel-Margoulis, E. Fricke, I. Liebich, S. Land, A. Barre-Dirrie, I. Reuter, et al.
2006. 'TRANSFAC and Its Module TRANSCompel: Transcriptional Gene Regulation in
Eukaryotes'. *Nucleic Acids Research* 34 (Database issue): D108-110.
https://doi.org/10.1093/nar/gkj143.

Milutin Gašperov, Nina, Sanja A. Farkas, Torbjörn K. Nilsson, and Magdalena Grce. 2014.
'Epigenetic Activation of Immune Genes in Cervical Cancer'. *Immunology Letters* 162
(2 Pt B): 256–57. https://doi.org/10.1016/j.imlet.2014.09.019.

Milutin Gašperov, Nina, Ivan Sabol, Ksenija Božinović, Emil Dediol, Marinka Mravak-Stipetić,
Danilo Licastro, Simeone Dal Monego, and Magdalena Grce. 2020. 'DNA Methylome
Distinguishes Head and Neck Cancer from Potentially Malignant Oral Lesions and
Healthy Oral Mucosa'. *International Journal of Molecular Sciences* 21 (18).
https://doi.org/10.3390/ijms21186853.

Moody, Laura, Sylvia L. Crowder, Andrew D. Fruge, Julie L. Locher, Wendy Demark-
Wahnefried, Laura Q. Rogers, Ashley Delk-Licata, et al. 2020. 'Epigenetic
Stratification of Head and Neck Cancer Survivors Reveals Differences in Lycopene
Levels, Alcohol Consumption, and Methylation of Immune Regulatory Genes'.
Clinical Epigenetics 12 (September): 138. https://doi.org/10.1186/s13148-020-
00930-5.

Nakagawa, Takuya, Keisuke Matsusaka, Kiyoshi Misawa, Satoshi Ota, Kiyoko Takane, Masaki
Fukuyo, Bahityar Rahmutulla, et al. 2017. 'Frequent Promoter Hypermethylation
Associated with Human Papillomavirus Infection in Pharyngeal Cancer'. *Cancer*
Letters 407 (October): 21–31. https://doi.org/10.1016/j.canlet.2017.08.008.

Numoto, M., K. Yokoro, and J. Koshi. 1999. 'ZF5, Which Is a Kruppel-Type Transcriptional

Repressor, Requires the Zinc Finger Domain for Self-Association'. *Biochemical and*
Biophysical Research Communications 256 (3): 573–78.
https://doi.org/10.1006/bbrc.1999.0375.
Patro, Rob, Geet Duggal, Michael I. Love, Rafael A. Irizarry, and Carl Kingsford. 2017. 'Salmon
Provides Fast and Bias-Aware Quantification of Transcript Expression'. *Nature*
Methods 14 (4): 417–19. https://doi.org/10.1038/nmeth.4197.
Perri, Francesco, Francesco Longo, Francesco Caponigro, Fabio Sandomenico, Agostino
Guida, Giuseppina Della Vittoria Scarpati, Alessandro Ottaiano, Paolo Muto, and
Franco Ionna. 2020. 'Management of HPV-Related Squamous Cell Carcinoma of the
Head and Neck: Pitfalls and Caveat'. *Cancers* 12 (4): 975.
https://doi.org/10.3390/cancers12040975.
Pikaart, Michael J., Félix Recillas-Targa, and Gary Felsenfeld. 1998. 'Loss of Transcriptional
Activity of a Transgene Is Accompanied by DNA Methylation and Histone
Deacetylation and Is Prevented by Insulators'. *Genes & Development* 12 (18): 2852–
62. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC317165/.
Raudvere, Uku, Liis Kolberg, Ivan Kuzmin, Tambet Arak, Priit Adler, Hedi Peterson, and Jaak
Vilo. 2019. 'G:Profiler: A Web Server for Functional Enrichment Analysis and
Conversions of Gene Lists (2019 Update)'. *Nucleic Acids Research* 47 (W1): W191–98.
https://doi.org/10.1093/nar/gkz369.
Slebos, Robbert J. C., Yajun Yi, Kim Ely, Jesse Carter, Amy Evjen, Xueqiong Zhang, Yu Shyr, et
al. 2006. 'Gene Expression Differences Associated with Human Papillomavirus Status
in Head and Neck Squamous Cell Carcinoma'. *Clinical Cancer Research: An Official*
Journal of the American Association for Cancer Research 12 (3 Pt 1): 701–9.
https://doi.org/10.1158/1078-0432.CCR-05-2017.
Slomp, J., A. C. Gittenberger-de Groot, V. E. Koteliansky, M. A. Glukhova, A. J. Bogers, and R.
E. Poelmann. 1997. 'Cytokeratin Expression in Human Arteries Pertinent to Intimal
Thickening Formation in the Ductus Arteriosus'. *Differentiation; Research in*
Biological Diversity 61 (5): 305–11. https://doi.org/10.1046/j.1432-
0436.1997.6150305.x.
Sun, Yingming, Zhe Wang, Sufang Qiu, and Ruoyu Wang. 2021. 'Therapeutic Strategies of
Different HPV Status in Head and Neck Squamous Cell Carcinoma'. *International*
Journal of Biological Sciences 17 (4): 1104–18. https://doi.org/10.7150/ijbs.58077.
Tan, Hua, Shan Huang, Zhigang Zhang, Xiaohua Qian, Peiqing Sun, and Xiaobo Zhou. 2019.
'Pan-Cancer Analysis on microRNA-Associated Gene Activation'. *EBioMedicine* 43
(May): 82–97. https://doi.org/10.1016/j.ebiom.2019.03.082.
Tripathi, Nimisha, Sneha Keshari, Pallavi Shahi, Poonam Maurya, Atanu Bhattacharjee,
Kushal Gupta, Sanjay Talole, and Mukesh Kumar. 2020. 'Human Papillomavirus
Elevated Genetic Biomarker Signature by Statistical Algorithm'. *Journal of Cellular*
Physiology 235 (12): 9922–32. https://doi.org/10.1002/jcp.29807.
Xu, Peng, Qian Wu, Jian Yu, Yongsheng Rao, Zheng Kou, Gang Fang, Xiaolong Shi, Wenbin Liu,
and Henry Han. 2020. 'A Systematic Way to Infer the Regulation Relations of miRNAs
on Target Genes and Critical miRNAs in Cancers'. *Frontiers in Genetics* 11.
https://www.frontiersin.org/articles/10.3389/fgene.2020.00278.
Yan, Wusheng, Ignacio I. Wistuba, Michael R. Emmert-Buck, and Heidi S. Erickson. 2011.
'Squamous Cell Carcinoma - Similarities and Differences among Anatomical Sites'.
American Journal of Cancer Research 1 (3): 275–300. https://doi.org/10.1158/1538-
7445.am2011-275.
Yin, Yimeng, Ekaterina Morgunova, Arttu Jolma, Eevi Kaasinen, Biswajyoti Sahu, Syed Khund-
Sayeed, Pratyush K. Das, et al. 2017. 'Impact of Cytosine Methylation on DNA

Binding Specificities of Human Transcription Factors'. *Science (New York, N.Y.)* 356
(6337): eaaj2239. https://doi.org/10.1126/science.aaj2239.
Zhao, Min, Pora Kim, Ramkrishna Mitra, Junfei Zhao, and Zhongming Zhao. 2016. 'TSGene
2.0: An Updated Literature-Based Knowledgebase for Tumor Suppressor Genes'.
Nucleic Acids Research 44 (D1): D1023-1031. https://doi.org/10.1093/nar/gkv1268.
581

582 Declarations

583 Ethics approval and consent to participate

584 All patients provided informed consent to participate at the time of primary cancer

585 treatment. The collection of samples was approved by Bioethical Bord of the Ruđer Bošković

586 Institute (BEP-3748/2-2014) and the Ethical Board of the Clinical Hospital Dubrava (EP- KBD-

587 10.06.2014).

588 Consent for publication

589 Not applicable - no personally identifiable information is presented in the manuscript.

590 Availability of data and materials

Code used is available at https://github.com/PKatarina/HNSCC_Grce

Raw datasets analyzed during the current study are available in the Figshare repository

under the following DOI identfiers:

Transcriptome https://doi.org/10.6084/m9.figshare.24721746

miRNome https://doi.org/10.6084/m9.figshare.24721731

Methylome https://doi.org/10.6084/m9.figshare.24721728

All datasets and raw data will be deposited in ArrayExpress repository as well. Upon

publication links will be replaced by accession numbers

Due to the file size restrictions Supplementary table S3 is deposited at Figshare repository

https://doi.org/10.6084/m9.figshare.24722268

Competing interests

603 The authors declare that they have no competing interests.

Funding

605 This study was funded by the Croatian Science Foundation grants HRZZ-IP-2013-11-4758 and

606 UIP-2020-02-1623. KM is supported by Croatian Science Foundation doctoral student

607 scholarship DOK-2021-02-7831

Authors' contributions

609 NMG, IS, MG and AB designed the study. KB, ED, JK, NS, IS and KM collected and processed

610 the data. KM performed the analyses. KM, KB and NMG wrote the first draft of the

611 manuscript. MG, IS and AB supervised and acquired funding for the study. All authors

612 contributed to the writing of the final manuscript. All authors read and approved the final

613 manuscript.

Acknowledgement

615 The authors are thankful to Prof. Marinka Mravak-Stipetić for oral normal tissue samples

616 collection, and to Dr. Ruth Tachezy for providing reference non-malignant tonsil control

617 tissues. The authors are also grateful to Jasminka Golubić Talić for her technical assistance.

Supplemental figures

Integrative analysis in head and neck cancer reveals distinct role of miRNome and methylome as tumour epigenetic drivers

Katarina Mandić[1]*, Nina Milutin Gašperov[2]*, Ksenija Božinović[2], Emil Dediol[3], Jure Krasić[4], Nino Sinčić[4,5], Magdalena Grce[2], Anja Barešić[1]**, Ivan Sabol[2]**

Suplemental figure 1. Significant number of hypermethylated and hypomethylated CpGs in cancer vs. control groups across all genomic elements (Chi squared p < 2x10^-16). Labels: Gene body – across exons and introns; Promoter region – from 500 to 2000 distance from canonical transcription start site; LINE – Long interspersed nuclear elements; SINE – Short interspersed nuclear elements; LTR – Long terminal repeat elements which include retroposons; DNA – DNA repeat elements; Satellite – Satellite repeats. Annotation for genomic elements were used from Jin et al., 2015.

Supplemental figure 2. Expression of E2F family genes in cancer vs control samples.