The relationship between gut and nasopharyngeal microbiome composition can predict the severity of COVID-19.

Benita Martin-Castaño[1,2,#], Patricia Diez-Echave[2,3,#], Jorge García-García[3,*] , Laura Hidalgo-García[2,3], Antonio Jesús Ruiz-Malagon[2,3], José Alberto Molina-Tijeras[2,3], María Jesús Rodríguez-Sojo[2,3], Anaïs Redruello-Romero[2], Margarita Martínez-Zaldívar[2,4], Emilio Mota[4], Fernando Cobo[5], Xando Díaz-Villamarin[2], Emilio Fernández-Varón[2,3], Marta Alvarez-Estevez[2,6,7], Federico García[2,6,7], Concepción Morales-García[8], Silvia Merlos[8], Paula García-Flores[8], Manuel Colmenero-Ruiz[2,9], Andrés Ruiz-Sancho[2,10], María Nuñez[2,11,12], María Elena Rodríguez-Cabezas[2,3], Ángel Carazo[2,6], Javier Martín[13], Rocío Morón[2,11,*], Alba Rodríguez-Nogales[2,3,‡], Julio Galvez[2,3,14, ‡].

[1] Centro de Salud Las Gabias, Distrito Granada-Metropolitano, 18110-Granada, Spain.

[2] Instituto de Investigación Biosanitaria de Granada (ibs.GRANADA), 18012-Granada, Spain.

[3] Department of Pharmacology, Center for Biomedical Research (CIBM), University of Granada, 18071-Granada, Spain.

[4] Centro de Salud "Salvador Caballero", Distrito Granada-Metropolitano, 18012-Granada, Spain.

[5] Servicio Microbiología, Hospital Universitario Virgen de las Nieves, 18014-Granada, Spain.

[6] Servicio Microbiología, Hospital Universitario Clínico San Cecilio, 18016-Granada, Spain.

[7] CIBER de Enfermedades Infecciosas (CIBER-Infecc), Instituto de Salud Carlos III, 28029- Madrid, Spain.

[8] Respiratory Medicine Department, Hospital Universitario Virgen de las Nieves, 18014- Granada, Spain.

[9] Servicio de Medicina Intensiva, Hospital Universitario Clínico San Cecilio, 18016- Granada, Spain.

[10] Servicio de Enfermedades Infecciosas, Hospital Universitario Clínico San Cecilio,
18016- Granada, Spain.

[11] Servicio Farmacia Hospitalaria, Hospital Universitario Clínico San Cecilio, 18016-
Granada, Spain.

[12] CIBER de Epidemiología y Salud Pública (CIBER-ESP), Instituto de Salud Carlos
III, 28029- Madrid, Spain.

[13] Department of Cell Biology and Immunology, Institute of Parasitology and
Biomedicine López-Neyra, CSIC, 18016-Granada, Spain.

[14] CIBER de Enfermedades Hepáticas y Digestivas (CIBER-EHD), Instituto de Salud
Carlos III, 28029- Madrid, Spain.

[#] Both authors contributed equally to this manuscript.

[*] Correspondence: jgarcia.51@ugr.es (J. G.-G.); rmoronr@gmail.com (R.R.-M).

[‡] Both authors contributed equally to this work.

ABSTRACT

Background: Coronavirus disease 2019 (COVID-19) is a respiratory illness caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) that displays great variability in clinical phenotype. Many factors have been described to be correlated with its severity but no specific determinants of infection outcome have been identified yet, maybe due the complex pathogenic mechanisms. The microbiota could play a key role in the infection and in the progression and outcome of the disease. Hence, SARS-CoV-2 infection has been associated with nasopharyngeal and gut dysbiosis and higher abundance of opportunistic pathogens. **Methods:** To identify new prognostic markers for the disease, a multicenter prospective observational cohort study was carried out in COVID-19 patients that were divided in three cohorts according to their symptomatology: mild (n=24), moderate (n=51) and severe/critical (n=31). Faecal and nasopharyngeal samples were taken and the microbiota was analysed. **Results:** Microbiota composition could be associated with the severity of the symptoms and the linear discriminant analysis identified the genera *Mycoplasma* and *Prevotella* as severity biomarkers in nasopharyngeal samples, and *Allistipes*, *Enterococcus* and *Escherichia* in faecal samples. Moreover, *M. salivarium* was defined as a unique microorganism in COVID-19 patients' nasopharyngeal microbiota while *P. bivia* and *P. timonensis* were defined in faecal microbiota. A connection between faecal and nasopharyngeal microbiota in COVID-19 patients was also identified as a strong positive correlation between *P. timonensis* (faeces) towards *P. dentalis* and *M. salivarium* (nasopharyngeal) was found in critically ill patients. **Conclusions:** This ratio could be used as a novel prognostic biomarker for severe COVID-19 patients.

Keywords: COVID-19; Gut microbiota; Nasopharyngeal microbiota; SARS-CoV-2; Severity.

INTRODUCTION

Coronavirus disease 2019 (COVID-19) is a respiratory illness caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). The data reported in November 2023 revealed that almost 700 million people have been infected with the virus [1]. Even though the majority of COVID-19 cases are mild, disease has been also shown to cause long-term effects on human health. Therefore, a remarkable feature of SARS-CoV-2 infection is the great variability in clinical phenotype among infected people. Many factors can correlate with COVID-19 disease severity, including age, gender, body mass index, previous comorbidities, immune responses, and genetics [4-6], but, unfortunately, the determinants of infection outcome and the pathogenic mechanisms are not completely understood yet [3].

SARS-CoV-2 primarily infects the respiratory tract by binding to angiotensin-converting enzyme 2 (ACE2) receptor [7], and a growing body of evidence suggests that it can also infect other organs since viral particles and nucleic acids have been found in various biological samples, like sputum, bronchoalveolar lavage fluid, faeces, blood, and urine [8-10]. Thus, ACE2 has been detected by single-cell RNA sequencing in various organs and tissues, like the gastrointestinal tract, where they are highly expressed [11], suggesting a substantial involvement of the gastrointestinal tract in the pathogenesis of the disease, including the ability of SARS-CoV-2 to infect and replicate in intestinal enterocytes [12], increased expression of the viral entry receptor (ACE2 receptor) and several membrane-bound serine proteases (such as transmembrane protease serine 2 (TMPRSS2) and TMPRSS4) in intestinal epithelial cells [13].

Moreover, SARS-CoV-2 infection has been extensively reported to induce dysbiosis the in the respiratory tract and the colon [17-20], characterized by increased presence of opportunistic pathogens, including *Staphylococcus*, *Corynebacterium* and *Acinetobacter* bacteria [14, 15], which can raise the risk of secondary infections, morbidity and mortality [16]. Thus, it is evident that there is a relevant connection between the microbiome from the respiratory and gastrointestinal tracts and the development and progression of this disease, and also the recovery processes [14, 20]. However, there is limited understanding of its precise association with the establishment of different

symptomatic profiles in this condition, and to date, few studies have focused on the
relationships between the severity of COVID-19 and the microbiome composition of the
nasopharyngeal and intestinal tracts contemplated simultaneously.

Considering that the emergence of mutations and variants has caused several additional
waves of infection and threatens to compromise the efficacy of existing vaccines and
anti-viral drugs [2], new therapeutic approaches and prognostic tools are necessary.
Therefore, the characterization of the nasopharyngeal and intestinal microbiome will
allow identifying predictive biomarkers for the diagnosis and prognosis of the disease,
as well as possible therapeutic targets in the management of SARS-CoV-2.

MATERIALS AND METHODS

Ethics approval.

The study was conducted in accordance with the declaration of Helsinki and the
protocol approved by the Clinical Research Ethics Committee of Granada (CEIC) (ID
of the approval omicovid-19 1133-N-20). All patients provided written informed
consent before being included in the study. The samples were managed by the
ibs.GRANADA Biobank following the protocols approved by the Andalusian
Biomedical Research Ethics Coordinating Committee.

Subject recruitment and sample collection
A multicentre prospective observational cohort study was carried out between
September 2020 and July 2021. Patients with SARS-CoV-2 infection were recruited
from the University Hospital San Cecilio, the University Hospital Virgen de las Nieves,
and the Primary Care centres, Salvador Caballero and Las Gabias in Granada (Spain).
These patients were laboratory-confirmed SARS-CoV-2 positive by quantitative reverse
transcription polymerase chain reaction (RT-qPCR) performed on nasopharyngeal
swabs collected by healthcare practitioners. Patients were classified in three groups
based on severity profile following the described guidelines [21] mild cohort (n=24),
subjects with moderate symptomatology (n=51) and severe/critically ill patients (n=31).
Mild illness included individuals who have any of the various signs and symptoms of
COVID-19 (e.g., fever, cough, sore throat, malaise, headache, muscle pain, nausea,

vomiting, diarrhoea, loss of taste and smell) but do not have shortness of breath,

dyspnoea, or abnormal chest imaging. Moderate cases were those showing fever and

respiratory symptoms with radiological findings of pneumonia. Severe group was

composed of patients with any of the following criteria: respiratory distress ($\geq$30

breaths/min), oxygen saturation $\leq$93% at rest, arterial partial pressure of oxygen

(PaO$_2$)/fraction of inspired oxygen (FiO$_2$) $\leq$300 mmHg, respiratory failure and requiring

mechanical ventilation, shock, and with other organ failures that required intensive care.

Healthcare staff collected Nasopharyngeal swabs and stools samples from patients while

asymptomatic patients provided stools self-sampled at home. Stools and nasopharyngeal

swabs were collected in collection tubes containing preservative media

(OMNIgene®•GUT, DNAGENOTEK®, Ottawa, Ontario, Canada) and stored at −80°C

until processing.

Microbial DNA extraction, library preparation and next generation sequencing

For all faecal and nasopharyngeal samples, DNA was isolated according to the protocol

reported by *Rodríguez-Nogales et al.* [22] and using Qiagen Allprep PowerFecal DNA

kit (Qiagen, Hilden, Germany). DNA was quantified using Qubit dsDNA HS assay kit

(Yeason Biotechnology, Shanghai, China) and total DNA was amplified by targeting

variable regions V4-V5 of the bacterial 16 S rRNA gene. Quality control of amplified

products was achieved by running a high-throughput Invitrogen 96-well-E-gel (Thermo

Fisher Scientific, Waltham, MA, USA). PCR products from the same samples were

pooled in one plate and normalised with the high-throughput Invitrogen SequalPrep 96-

well Plate kit. Then, the samples were pooled into a library prior to sequencing. Lastly,

Next-Generation Sequencing (NGS) techniques were performed using an Illumina

MiSeq machine.

Bioinformatic tools and statistical analysis

Bioinformatic analysis of demultiplexed raw data from nasopharyngeal and stool

microbiota samples was performed with QIIME2 software (open access, Northern

Arizona University, Flagstaff, AZ, USA). Trimming and filtering taking into account

their quality scores before specific taxa identification achieved quality control of the

samples. DADA2 software was employed to carry out denoising steps and to obtain

amplicon sequence variants (ASVs). SILVA reference database was used for taxonomic

assignment [23]. The remaining analyses were performed R software [24].

For numerical clinical variables analysis, data was displayed as mean ± SD when it

followed a normal distribution and median and interquartile range were represented for

non-normal distributions. Categorical variables were set out as percentages. In these

cases, statistical differences were calculated by ANOVA and Kruskal Wallis test for

numerical variables and Fisher's exact test for categorical.

Alpha and beta diversity and relative abundance were appraised with the Phyloseq

package. Normality and homogeneity of variance were examined by the Nortest and

LeveneTest packages, respectively. When these assumptions were reached, an ANOVA

test was carried out. Otherwise, the Kruskal Wallis test was employed.

Beta diversity differences were analysed with a Permutational Multivariate Analysis of

Variance (PERMANOVA) included in the Vegan package. Euler and microbial

packages were utilised for constructing Venn diagrams and to perform linear

discriminant analysis (LDA) effect size (LEfSe) with an LDA score of 3. The Corrplot

package was applied for correlation analysis using the Spearman's correlation

coefficient.

RESULTS

Study patients characteristics

A total of 106 patients (52 women and 54 men) who had laboratory confirmation of

SARS-CoV-2 infection were included in the present study. The patients had a median

age of 54 years (range, 40 to 68). Based on the clinical spectrum criteria reported in the

COVID-19 treatment guidelines, patients were categorised into 3 cohorts: mild

symptomatology (24 patients), moderate illness and hospitalised in Respiratory Unit (51

patients) and severe symptomatology and admitted in the intensive care units (ICU) (31

patients) (**Table 1**). As expected, the age of the patients significantly increased with the

severity of the symptoms, and therefore, the patients included in the severe symptoms

group were significantly older than those with mild or moderate symptoms (**Table 1**).

Patient inclusion was carried out evenly in terms of gender; nevertheless, a gender-

related impact on the clinical course of these patients can be observed since the group of

patients with severe symptoms was predominantly composed of males when compared

with the mild illness group (**Table 1**). Correspondingly, the clinical course of patients classified according to severity was different. Most mild patients showed symptoms of a mild respiratory infection, but a third of them also displayed dyspnoea and low oxygen saturation, and a quarter reported the existence of gastrointestinal complaints (like stomach ache, digestive discomfort or diarrhoea); and a low percentage of patients (4%) reported high respiratory and heart rates. Moderate and severe patients showed higher frequencies of the evaluated symptoms: dyspnoea, low oxygen saturation and increased respiratory or heart rates (p < 0.05). However, no significant differences were observed in the percentage of the gastrointestinal complaints among the three groups of patients (**Table 1**). When the different comorbidities were considered, only those patients with severe symptoms showed a higher percentage of cardiomyopathy compared to those from mild or moderate symptomatology (p < 0.05). Additionally, no significant differences were found in the prevalence of the other pathologies among groups. Regarding the counts of lymphocytes and neutrophils did not show meaningful differences between the three groups of patients. However, the plasmatic determinations of platelets, D-dimer, ferritin and C reactive protein correlated with the severity of the symptoms, being the severe group significantly different (p<0.05) (**Table 1**).

	Mild (n=24)	Moderate (n=51)	Severe (n=31)
Clinical variables			
Age (Years)	43 ± 12 [a]	54 ± 14 [b]	62 ± 11 [c]
Gender (Male)	33% [a]	47% [b]	71% [c]
Symptoms			
Dyspnoea (Yes)	33% [a]	75% [b]	84% [b]
Gastrointestinal alteration (Yes)	26%	33%	33%
Respiratory rate (≥ 20 bpm)	4% [a]	22% [a]	63% [b]
SpO_2 (Low)	33% [a]	43% [a]	71% [b]
Heart rate (≥ 100 bpm)	4% [a]	27% [b]	55% [c]
Comorbidities			
Obesity (Yes)	26%	27%	32%
Diabetes (Yes)	20%	18%	26%
Asthma (Yes)	3%	6%	7%
Cardiomyopathy (Yes)	5% [a]	6% [a]	42% [b]
Plasma determinations			
Lymphocytes (10^3/µL)	1.1 ± 0.6	1.4 ± 0.6	1.2 ± 2.7
Neutrophils (10^3/µL)	6 [5.5;6.6]	6.4 [4.2;8.6]	7.9 [5.4;10.9]
Platelets (10^3/µL)	329.6 ± 8.5 [a]	257.4 ± 115 [b]	276.4 ± 93 [b]
D dimer (mg/L)	0.39 [0.2;0.8] [a]	0.6 [0.3;1] [a]	1.6 [0.9;4.3] [b]
Ferritin (ng/L)	157 [126;179] [a]	487 [274;1027] [b]	829 [488;1376] [c]
C reactive protein (mg/L)	3.4 [2.6;4] [a]	18.2 [7.8;41.9] [b]	162 [65;210] [c]

222 **Table 1. Clinical data description of enrolled patients.** Normal distributions are represented as mean
223 $\pm$ SD while non normal distributions are represented by median and interquartile range. Categorical
224 variables are represented with percentage. Groups with different letters statistically differ ($p < 0.05$).
225
226

Bacterial composition differs between sample type and severity index in SARS-CoV-2

infected patients

Nasopharyngeal swabs and faeces were obtained from all the patients included in the

study in the first seven days after symptom onset, and used for characterization of the

microbiota composition. Microbiome diversity showed alterations that could be

associated with the disease severity (**Figure 1**). Specifically, the α-diversity in the

nasopharyngeal microbiota was reduced in the moderate and severe groups, in

comparison with the mild group although it was only significant in patients with

moderate symptoms (**Figure 1A**). Conversely, when α-diversity was examined in stool

samples, no significant modifications were observed between groups (**Figure 1B**). On

the other hand, ß-diversity analysis revealed statistical differences between groups for

both samples, nasopharyngeal swabs and stools ($p < 0.001$) (**Figure 1C** and **1D**).

Nasopharyngeal microbial populations could be grouped based on the severity of the

symptoms and appear like three distinct and separate clusters corresponding to the

patients with mild, moderate and severe symptoms (**Figure 1C**). Remarkably, faecal

microbial communities of patients with severe symptoms differed significantly from

those of mild and moderate ill patients using the unweighted Bray-Curtis metric, which

compares samples based on bacterial presence-absence information (**Figure 1D**).

248
249 **Figure 1. Nasopharyngeal and gut microbiota composition is modified depending on the severity of**
250 **COVID-19 symptoms.** (**A**) Alpha diversity analysis of nasopharyngeal swab samples microbiota. (**B**)
251 Alpha diversity analysis of stool samples microbiota. (**C**) PCoA for Bray-Curtis index of nasopharyngeal
252 swab microbiota. (**D**) PCoA for Bray-Curtis index of stool samples microbiota. Values are represented as
253 mean ± SD. Significant differences are represented as * = p < 0.05.
254

Similarly, the characterization of microbiota composition revealed heterogeneity in the
microbiota profile associated with severity and disease progression in these patients
(**Figure Sx**). At phylum level, in nasopharyngeal microbiota, the abundance of
Bacillota was increased while the abundance of *Bacteroidota* and *Actinobacteroidota*
was reduced in patients with severe symptomatology (**Figure 2A**). Conversely, the three
groups presented a more homogeneous distribution of faecal microbiota than the
nasopharyngeal one, being the most abundant phyla *Bacillota* and *Bacteroidota* (**Figure
2B**). Only the patients that had a worse prognosis showed a decrease of abundance in
Bacteroidota (**Figure 2B**).

At genus level, symptom severity was associated with a higher number of detected
genera (**Figure 2C-D**). Of note, the nasopharyngeal microbiome composition revealed
significant differences between groups in genus abundance. The mild group presented
significantly higher abundance of *Alistipes*, *Muribaculaceae* and *Lachnospiraceae* (p <
0.001),   the moderate group showed a significant increase in *Alcaligenes* and
Pseudorobacter (p < 0.001) while the severe group had significantly higher relative

abundance of *Acinetobacter*, *Actinomyces*, *Anaerococcus*, *Atopobium*, *Campylobacter*, *Dolosigranulum*, *Enterobacter*, *Enterococcus*, *Finegoldia*, *Fusobacterium*, *Gemella*, *Haemophilus*, *Lawsonella*, *Leptotrichia*, *Megasphaera*, *Neisseria*, *Serratia*, *Rotia* and *Veillonella* (p < 0.001) (**Figure 2C**). However, a reduction in the number of detected genera was observed in stool samples as symptom severity increased (**Figure 2D**). Concretely, mild patients showed more presence of *Barnesiella*, *Muribaculaceae* and different members of the *Clostridia* class (*Clostridia*, *Coprococcus*, *Dorea*, *Lachnospiraceae*, *Roseburia* and *Ruminococcus*) (p < 0.001). Although moderate ill patients presented different genera, only *Streptococcus* was significantly increased in this group (p < 0.001). Remarkably, *Anaerococcus*, *Dialister*, *Lachnocostridium* or *Peptoniphilus* were more abundant in patients with severe symptoms (p < 0.001) (**Figure 2D**).

Figure 2. Microbiota composition of nasopharyngeal and stool samples at phylum level is slightly modified by COVID-19 symptoms severity. In contrast, at genus level, severity increases the total amount of detected bacteria in nasopharyngeal swabs while in stool samples it is reduced. (**A**) Representation of the most abundant phyla in nasopharyngeal swab samples. (**B**) Representation of the most abundant phyla in stool samples. (**C**) Taxa identification of the most abundant genera in nasopharyngeal swab samples. (**D**) Taxa identification of the most abundant genera in stool samples.

292
293
294 *Differences in bacteria abundance could be used as biomarkers to predict disease*
295 *severity and outcome in SARS-CoV-2 infection*
296

We investigated if some specific taxa could contribute to the severity of the symptoms.
ASVs were evaluated to determine core taxa along with the specific bacteria of each
group of patients and samples (**Figure 3A,B**). In nasopharyngeal swabs, Venn diagram
analysis revealed that the three groups of study shared 51 core taxa. 60 specific bacteria
were identified in patients with mild symptoms while 32 were seen in patients with
moderate symptoms and 8 in patients with severe symptoms. In stool, 159 core taxa
were shared by the three groups of patients, being 27 specific for mild patients
symptoms, 33 for moderate patients and 27 for severe patients (For more details see
Table S1).
306

Figure 3. Differential analysis expression of microbiota composition from nasopharyngeal and stool samples revealed the presence of specific bacteria related to COVID-19 severity index. **(A)** Venn diagram showing ASVs distribution in nasopharyngeal swab samples. **(B)** Venn diagram showing ASVs distribution in stool samples. **(C)** LEfSe plot of taxonomic biomarkers present in nasopharyngeal swab samples (p value = 0.01 and LDA value = 4). **(D)** LEfSe plot of taxonomic biomarkers present in stool samples (p value = 0.01 and LDA value =4).

Besides, the linear discriminant analysis (LEfSe) was performed to identify differential microorganisms for each group of patients (**Figure 3C,D**). In nasopharyngeal samples, *Burkholderia sp.*, *Paraburkholderia sp.* and *Massilia sp.* were identified in mild patients; *Pseudomonas veronii*, *Stenotrophomonas rhizophila* and *Azotobacter chroococcum* in moderate patients; and *Mycoplasma salivarium*, *Prevotella dentalis*, *Leptotrichia* and *Haemophilus parainfluenzae* in severe patients. In stool samples, *Bacteroides coprocola*, *Veillonella sp.*, *Ruminococcus bicirculans* and *Sutterella stercoricanis* were identified as predictors of mild condition; *Prevotella stercorea*, *Bacteroides cellulosilyficus*, *Streptococcus salivarus*, *Bacteroides stercoris* and

Prevotella copri as predictors of moderate symptoms; and *Escherichia*, *Enterococcus*

durans, *Alistipes onderdonkii*, *Prevotella timonensis* and *Prevotella bivia* as markers of

severe condition.

To further assess the role of these potential biomarkers in the prediction of COVID-19

severity, a correlation analysis was performed (**Figure 4**). In summary, biomarkers for

mild symptomatology (*B. coprocola*, *R. bicirculans*, *S. stercoricanis* and *Veillonella*

sp.) presented a negative correlation profile with the different clinical features or

biochemical parameters evaluated. In contrast, biomarkers associated with severe

symptoms in nasopharyngeal swabs (*M. salivarium* and *Leptotrichia*) showed a positive

correlation with D dimer and cardiomyopathy, respectively. In addition, the other two

biomarkers linked to the highest severity (*H. parainfluenzae* and *P. dentalis*) also

showed a tendency related to CRP, D dimer and cardiomyopathy (**Figure 4A**).

Interestingly, similar results were found in stool samples. Severe biomarkers revealed a

positive correlation towards D dimer and CRP levels, especially *P. bivia* and *P.*

timonensis. These two bacteria also presented a positive association with ferritin levels,

age and respiratory rate, and a negative correlation with lymphocyte count (**Figure 4B**).

Figure 4. Whereas mild biomarkers showed negative correlations towards clinical variables, severe biomarkers presented positive correlations. (A) Correlation plot of nasopharyngeal swab biomarkers and clinical variables. **(B)** Correlation plot of stool samples biomarkers and clinical variables. RR: respiratory rate; HR: heart rate; GI: gastrointestinal alterations.

Identification of a novel microbiome-based COVID-19 prognosis approach.

Considering the LEfSe results, we propose a new approach to predict disease severity in patients suffering SARS-CoV-2 infection based on establishing a pattern of nasopharyngeal-gut microbiota. The Spearman's correlation analysis revealed no important associations between nasopharyngeal and faecal microbiota in mild and moderate groups (**Figure 5A,B**). However, in patients with severe symptoms the Spearman's rho coefficient showed a significant positive correlation between *P. timonensis* towards *P. dentalis* and *M. salivarium* (**Figure 5C**). Consequently, the ratio between the abundance of these bacteria could serve as reliable predictors of severity of COVID-19. The results revealed a significant increase in the ratios *P. timonensis* / *M. salivarium* and *P. timonensis*/*P. dentalis* in patients with severe symptoms compared to those with mild or moderate symptoms (**Figure 5D,E**).

Figure 5. The existence of a relationship between the abundance of nasopharyngeal severe biomarkers and stool severe biomarkers allow the employment of an abundance ratio between them as a new tool for predicting COVID-19 severity. (A) Correlation plot among biomarkers found in nasopharyngeal swab and stool samples in each condition (mild, moderate and severe from left to right) **(B)** Ratio of the abundance between *P. timonensis* (stool) and *M. salivarium* and *P.dentalis* (nasopharyngeal swab) biomarkers. Groups with different letters statistically differ (p < 0.05).

370
DISCUSSION

Recent findings have evidenced the prominent role of the microbiome in viral
infections, and it can either promote or supress viral them [25, 26]. In fact, different
studies have explored the interplay between the host microbiota and SARS-CoV-2
infection [27, 28]. However, few studies have investigated the nasopharyngeal-faecal
axis as a potential biomarker of severity in patients infected with SARS-CoV-2. Hence,
the present study provides several nasopharyngeal and gut microbiota-based biomarkers
that could help to predict COVID-19 severity.

According to the National Institutes of Health (NIH), COVID-19 severity is classified
depending on the associated symptoms [21] that include age, gender, D-dimer levels,
dyspnoea and higher SpO_2 score, which are predictors of worse disease progression
[29]. The current study confirmed that older age as well as a higher percentage of
dyspnoea; increased heart and respiratory rates together with lower oxygen saturation
were associated with severe symptoms. In fact, ageing is related to immune response
decline as well as higher incidence of systemic, chronic and low-grade inflammation
called inflammaging. Gender is also considered a risk factor, as a recent meta-analysis
has shown that men tended to have higher risk of developing severe symptoms, being
hospitalised, admitted to the intensive care units and die [34] for more severe disease.
The results in the present study support these previous studies as it is found that men
and women are disproportionately affected since males suffered from more severe
disease than females, including higher ICU admission rates, dyspnoea, increased heart
rate. Sex disparities in symptoms severity has been attributed to higher rates of
hazardous behaviours and existence of comorbidities, such as cardiomyopathy, in males
than in females. In fact, the incidence of cardiovascular complications in COVID-19
pathology appears to be associated with sex and gender differences, thus contributing to
the greater severity and poorer outcomes of the SARS-CoV2-mediated disease in male
patients compared to women [35]. This relationship has also been demonstrated in this
study since a higher rate of cardiovascular condition is evidenced in male patients as the
severity of symptoms increases. In this context, there is few and controversial sex-
stratified data investigating the role of cardiovascular complications in the prognosis
and outcome of COVID-19 disease in men and women. However, it has been reported

that women exhibit higher expression and activation of angiotensin type 2 (AT2)

receptors, which have been associated with a more robust anti-inflammatory immune

response against SARS-CoV-2 infection and are also involved in the control of blood

pressure and renal function, thereby providing protection for cardiovascular

complications in female patients [36]. Remarkably, the extent of cardiac cell mortality is

more noticeable in males than in females under various conditions [35], and this could

be linked to an augmented protection against cardiovascular complications in female

patients. Additionally, women produce high levels of regenerative white blood cells and

epoxy-eicosatrienoic acids, which display antihypertensive and anti-inflammatory

properties on blood vessels [37]. Consequently, this leads to restricted cardiac

remodelling and a more effective restoration of functionality [38]. Regarding some

biochemical parameters typically described as biomarkers for COVID-19 severity (D-

dimer, ferritin and CRP) [39, 40], the findings obtained in this study found that those

groups of patients with moderate or severe symptoms showed significantly increased

levels of these plasma parameters, which are consistent with previous reports [41-43].

The association of then microbiota composition with these clinical variables has been

widely studied [15] and it is well described that the microbiota can modulate host

immunity and physiological functions [47], Consequently, the microbiota could be key

in the clinical phenotype of these patients although the specific contribution of the

microbiota to the progression of the infection and a poor prognosis is not yet fully

understood. This study addresses for the first time the implication of nasopharyngeal

and faecal microbiota in the prognosis of COVID-19. Firstly, when the alpha and beta

diversity were evaluated, the results revealed only substantial changes in richness and

Shannon diversity index in nasopharyngeal microbiome associated with severe

symptoms. In this sense, controversial results have been previously reported, and

although most of the studies have proposed that SARS-CoV-2 infection is associated

with lower microbial diversity in nasopharyngeal samples [14, 48, 49], others did not

find differences in alpha diversity composition among groups with different

symptomatology [50, 51].

Furthermore, in terms of beta diversity, previous studies have reported modifications in

the microbiome composition from the respiratory or gastrointestinal tract in COVID-19

patients when compared to healthy subjects [19, 52]. In the present study, in both

nasopharyngeal swabs and stools, every group of patients presented distinctly
differentiated clusters. As previously reported, COVID-19 disease severity is more
dependent on the presence or absence of certain bacteria rather than alterations in
bacterial diversity and richness [14, 53]. Supporting this, characterization of bacterial
microbiota composition at phyla and genera levels for nasopharyngeal and stool
samples indicated a more evident association between changes on them and the severity
of the disease. Specifically, in nasopharyngeal swabs, the presence of *Bacteroidota* and
Actinobacteriota has been previously linked to a better prognosis of SARS-CoV-2
infection, since these bacteria have been proposed to exert beneficial effects by
preventing respiratory diseases, including COVID-19 [54-58]. Moreover, in
nasopharyngeal samples, the abundance of *Bacillota* and *Pesudomonadota* was
increased in patients with severe symptomatology, thus supporting previous studies in
which higher counts of *Bacillota* (*Staphylococcus* sp. and *Streptococcus* sp.) and
Pesudomonadota (*Pseudomonas* sp.) were associated with moderate and severe
symptoms of COVID-19 [59].

The evaluation of genera abundance composition showed that *Alistipes* and
Muribaculaceae were highly abundant in mild patients. While these bacteria have been
well characterised in gut microbiota, little information regarding their presence in
nasopharyngeal microbiota has been provided up to date. Different experimental studies
in mice have suggested their role in viral infections. Thus, *Muribaculaceae* was found
in the lung microbiota in SARS-CoV-2 infected mice that were treated with a selective
inhibitor of the main protease (M^{pro}) [60]. In the case of *Alistipes,* in a study conducted
in children infected with respiratory syncytial virus (RVS), these bacteria were more
abundant in the nasopharyngeal microbiota of non RVS-infected subjects [61]. Overall,
these genera could be associated with a protective role against viral infection, and their
higher presence in nasopharyngeal samples from mild COVID-19 patients could
prevent the progression to severe disease.

Interestingly, mild patients have also shown a higher content of *Lactobacillus*, similarly
to that reported previously in asymptomatic COVID-19 patients [62]. Of note, it is well
described that the microorganisms forming the protective microbiota are fundamentally
represented by *Lactobacillus* species. Correspondingly, the use of *Lactobacillus* strains
as probiotics for preventing viral infections has been previously explored [63], and
hence, its administration in SARS-CoV-2 infection could be considered to avoid
complications. The increased presence of other genera, such as *Corynebacterium*,

Acinetobacter, Staphylococcus and *Veillonella*, positively correlated with the severity of SARS-CoV-2 infection. This correlation is supported by previous studies in which these genera were associated with both disease severity and systemic inflammation [14, 64]. In addition, higher abundance of *Enterococcus* was observed in severe patients, thus confirming other studies in critically ill patients [65].

In contrast to the findings in the nasopharyngeal microbiota, the analysis at phylum level in stool samples did not reveal notable modifications among patient groups. However, in mild ill patients, different genera from *Clostridia* class (*Clostridia*, *Coprococcus*, *Dorea*, *Lachnospiraceae*, *Roseburia* and *Ruminococcus*), *Barnesiella* and *Muribaculaceae* were identified as highly abundant. While the class *Clostridia* was associated with a reduced production of proinflammatory cytokines in COVID-19 patients and in those who recovered from the infection [66, 67], *Barnesiella* prevents colonisation by antibiotic-resistant bacteria such as *Enterococcus*, which is involved in bloodstream infection in critically ill COVID-19 patients [68, 69]. Furthermore, studies performed in mice have shown that *Muribaculaceae* abundance is reduced in mice coinfected with different respiratory viruses, suggesting that it may play a protective role under viral infection [70]. Therefore, it seems that under SARS-CoV-2 infection, the reduction of *Barnesiella, Clostridia* and *Muribaculaceae* members were associated with more severe symptoms. Nonetheless, the detected genera for the severe illness group, *Lachnocostridum*, *Anaerococcus* and *Peptoniphilus*, have been recognised as opportunistic pathogens and could contribute to a poor prognosis through inducing gut inflammation [71].

Regarding these differences, both nasopharyngeal and gut microbiota composition could be used to identify specific bacteria to predict COVID-19 severity. In the present study, unique ASVs for each condition were identified. For nasopharyngeal samples, species belonging to the genus *Lactobacillus* (*L. fermentum* or *L. reuteri*) or *Prevotella* (*P. pallens, P. ori* and *P. shahii*) have been identified. The role of *Prevotella sp.* in COVID-19 infection has not been clearly elucidated. Published microbiome analysis have revealed that its abundance was higher in mild patients [72], although others have suggested that it could be a biomarker of critical phenotype in COVID-19 patients [73, 74]. In spite of the controversial results, the results obtained in this study would confirm the potential use of this species as a biomarker for mild symptomatology. Interestingly, *Anaerococcus prevotii* was one of the exclusive species found in stools in mild patients.

This species has been linked to lower inflammation in COVID-19 patients [75].
Conversely, *Coprobacillus cateniformis* was solely found in severe patients, which
could be involved in the development of a worse condition in these patients through
ACE2 upregulation [18].

Even though these bacteria are unique for each group, LEfSe was performed to obtain
specific biomarkers [76]. For mild patients, *Burkholderia* and *Paraburkholderia* were
identified in nasopharyngeal swabs and *B. coprocola* and *R. bicirculans* in stool
samples. Although the information regarding the first two species in humans is limited,
a few studies have reported their presence in the commensal human microbiota [77, 78].
Contrarily, *B. coprocola* and *R. bicirculans* have been found in both healthy and
COVID-19 patients [79] although the abundance of *R. bicirculans* was reduced in
infected subjects [80].

In patients with moderate symptoms, *P. veronii* was detected in nasopharyngeal samples
whereas *P. stercorea*, *B. cellulosilyficus*, *B. stercoris* and *P. copri* were identified in
stool samples. In general, these findings agree with previous studies in COVID-19
patients [81]. Thus, *Xu et al.* found that infected patients showed higher abundance of *B.*
cellulosilyficus [82], whereas *B. stercoris* and *P. copri* were associated with ACE2
upregulation and increased proinflammatory cytokine production, respectively, in
COVID-19 patients [79, 83].

In critically ill patients, the biomarkers found for nasopharyngeal microbiota were *M.*
salivarium, *P. dentalis*, *Leptotrichia* and *H. parainfluenzae*. In stool samples,
Escherichia sp., *E. durans*, *P. timonensis* and *P. bivia* were the species recognised as
biomarkers. In general, all of them have been observed in the microbiota of SARS-
CoV-2 infected patients. Moreover, both *P. bivia* and *P. timonensis* have been defined
as unique microorganisms in COVID-19 patients' microbiota [79, 84], whilst *M.*
salivarium, *H. parainfluenzae* and *E. durans* were related to a higher abundance and
poor outcome in these patients [85-87]. Of note, the use of these bacteria as biomarkers
of severity in SARS-CoV-2 infection is further supported by the fact that these species
exhibited positive correlations with various clinical variables. Specifically, *M.*
salivarium, *H. parainfluenzae, P. dentalis*, *P. bivia* and *P. timonensis* showed positive
correlation with ferritin, CRP and D-dimer levels, as well as cardiomyopathy and
respiratory rates. Several studies have revealed both the relationship between CRP
levels, gut microbiota and COVID-19 severity [88], as well as the positive correlation of
specific bacteria with D-dimer, CRP and the levels of pro-inflammatory mediators in

plasma [89]. When considering *A. onderdonkii*, it has been reported that this bacteria do not aggravate the symptomatology of the COVID-19 patients due to its anti-inflammatory properties [18]; however, there are conflicting evidences regarding its pathogenicity that indicate that *A. onderdonkii* may have protective effects against some diseases, including liver fibrosis, colitis and cardiovascular disease, as well as in cancer immunotherapy, while it may be *involved* in colorectal cancer development and affective disorders like depression [90]. Moreover, *Alistipes* is a relatively recent subdivision genus of the *Bacteriodota*, which is commonly associated with chronic intestinal inflammation [90]. Therefore, taking into account that *Zuo T et al.* employed a different methodology to analyse microbiota composition from stool samples [18], *A. onderdonkii* could be considered as a biomarker of severe condition in SARS-CoV-2 infected subjects.

Finally, the implication of a connection between faecal and nasopharyngeal microbiota in COVID-19 patients has been previously proposed [91]. In the present study, and to maximise the potential use of these biomarkers, the relationship of specific bacteria from nasopharyngeal and stool samples was analysed. Concretely, a strong positive correlation between *P. timonensis* (stool) towards *P. dentalis* and *M. salivarium* (nasopharyngeal) was found in severe condition. Accordingly, the ratio of the abundance of these species was also significantly increased within the highest severity of this condition. As a result, the ratio proposed in this study could be used as a novel predictor to identify critically ill COVID-19 patients as the ratio *Bacillota* and *Bacteroidetes* has been used as a marker of dysbiosis [92]. In this case, this ratio *P. timonensis*/*P. dentalis* and *M. Salivarium* could be a prognostic tool for severe SARS-CoV-2, and an increase in it could be associated with a higher risk to develop a severe condition.

CONCLUSION

This inter-individual variability between the COVID-19 patients could contribute to the different symptomatology observed. This study has identified a correlation between changes in the nasopharyngeal and stool microbiota with COVID-19 severity. A novel biomarker linked to severity of COVID-19 infection has been described based on changes in the abundance of bacterial species in nasopharyngeal and faecal samples. This knowledge can support the design of novel therapeutic strategies to mitigate adverse outcomes. Further investigations are imperative to explore how the association

573 between nasopharyngeal and faecal microbiota can be modulated to uncover its role in
574 enhancing immune health, preventing or treating SARS-CoV-2 infections, and fostering
575 immunity.
576

AUTHOR CONTRIBUTIONS
Benita Martín-Castaño, Margarita Martínez-Zaldívar, Emilio Mota, Fernando Cobo,
Concepcion Morales-García, Marta Alvarez-Estevez, Federico García, Silvia Merlos,
Paula García-Flores, Manuel Colmenero-Ruiz, José Hernández-Quero, María Nuñez
were involved in the sample collection. Patricia Diez-Echave, Jorge García-García,
Alba Rodríguez-Nogales, Maria Elena Rodríguez-Cabezas, Laura Hidalgo-García,
Antonio Jesús Ruiz-Malagon, José Alberto Molina-Tijeras, María Jesús Rodríguez-Sojo
and Anaïs Redruello were involved in the processing of simples and obtaining results.
Rocio Morón and Emilio Fernández-Varón had access to the data and were involved in
the conception and data analysis and interpretation. Alba Rodríguez-Nogales, Javier
Martin, Maria Elena Rodriguez-Cabezas, Benita Martín-Castaño, Rocio Morón, Jorge
García-García, Ángel Carazo and Julio Gálvez had access to the data and were involved
in the conception and design of the work, data analysis and interpretation, critical
revision of the article and final approval before submission. All authors reviewed the
final manuscript and agreed to be account able for all aspects of the work.
592

593 **ACKNOWLEDGEMENTS**
594 We acknowledge the collaboration of all the participants who voluntarily and selflessly
595 participated in the study.
596

597 **CONFLICT OF INTEREST STATEMENT**
598 All authors declare no interest.
599

600 **FUNDING INFORMATION**
601 The research project was sup-ported by Government of Andalucia (Spain) (CV20-
602 99908).
603

604 **DATA AVAILABILITY STATEMENT**
605 Participant data cannot be made publicly available due to the sensitive nature of the
606 personal health data and privacy and confidentiality reasons. However, under certain

607 conditions, these data could be accessible for statistical and scientific research. For
608 further information, please contact the corresponding authors.
609
610
611
612 **REFERENCES**
613

1.　Geneva: World Health Organization, A.o.h.c.w.i., *WHO COVID-19 Dashboard.*
2020.
2.　Harvey, W.T., et al., *SARS-CoV-2 variants, spike mutations and immune escape.*
Nat Rev Microbiol, 2021. **19**(7): p. 409-424.
3.　Crook, H., et al., *Long covid-mechanisms, risk factors, and management.* BMJ,
2021. **374**: p. n1648.
4.　Yang, J., et al., *Prevalence of comorbidities and its effects in patients infected*
with SARS-CoV-2: a systematic review and meta-analysis. Int J Infect Dis, 2020.
94: p. 91-95.
5.　Severe Covid, G.G., et al., *Genomewide Association Study of Severe Covid-19*
with Respiratory Failure. N Engl J Med, 2020. **383**(16): p. 1522-1534.
6.　Bastard, P., et al., *Autoantibodies against type I IFNs in patients with life-*
threatening COVID-19. Science, 2020. **370**(6515).
7.　Zhou, P., et al., *A pneumonia outbreak associated with a new coronavirus of*
probable bat origin. Nature, 2020. **579**(7798): p. 270-273.
8.　Peng, L., et al., *SARS-CoV-2 can be detected in urine, blood, anal swabs, and*
oropharyngeal swabs specimens. J Med Virol, 2020. **92**(9): p. 1676-1680.
9.　Sun, J., et al., *Isolation of infectious SARS-CoV-2 from urine of a COVID-19*
patient. Emerg Microbes Infect, 2020. **9**(1): p. 991-993.
10.　Wang, W., et al., *Detection of SARS-CoV-2 in Different Types of Clinical*
Specimens. JAMA, 2020. **323**(18): p. 1843-1844.
11.　Hao, Z., et al., *Digestive system is a potential route of COVID-19: an analysis of*
single-cell coexpression pattern of key proteins in viral entry process. Gut,
2020. **69**(6): p. 1010.
12.　Lamers, M.M., et al., *SARS-CoV-2 productively infects human gut enterocytes.*
Science, 2020. **369**(6499): p. 50-54.

13.   Zang, R., et al., *TMPRSS2 and TMPRSS4 promote SARS-CoV-2 infection of*
641   *human small intestinal enterocytes.* Sci Immunol, 2020. **5**(47).

642   14.   Candel, S., et al., *The nasopharyngeal microbiome in COVID-19.* Emerg
643   Microbes Infect, 2023. **12**(1): p. e2165970.

644   15.   Ancona, G., et al., *Gut and airway microbiota dysbiosis and their role in*
645   *COVID-19 and long-COVID.* Front Immunol, 2023. **14**: p. 1080043.

646   16.   De Bruyn, A., et al., *Secondary infection in COVID-19 critically ill patients: a*
647   *retrospective single-center evaluation.* BMC Infect Dis, 2022. **22**(1): p. 207.

648   17.   Gu, S., et al., *Alterations of the Gut Microbiota in Patients With Coronavirus*
649   *Disease 2019 or H1N1 Influenza.* Clin Infect Dis, 2020. **71**(10): p. 2669-2678.

650   18.   Zuo, T., et al., *Alterations in Gut Microbiota of Patients With COVID-19 During*
651   *Time of Hospitalization.* Gastroenterology, 2020. **159**(3): p. 944-955 e8.

652   19.   Wu, Y., et al., *Altered oral and gut microbiota and its association with SARS-*
653   *CoV-2 viral load in COVID-19 patients during hospitalization.* NPJ Biofilms
654   Microbiomes, 2021. **7**(1): p. 61.

655   20.   Cao, J., et al., *Integrated gut virome and bacteriome dynamics in COVID-19*
656   *patients.* Gut Microbes, 2021. **13**(1): p. 1-21.

657   21.   Health., N.I.o., *https://www.covid19treatmentguidelines.nih.gov/*. 2019.

658   22.   Rodriguez-Nogales, A., et al., *Differential intestinal anti-inflammatory effects of*
659   *Lactobacillus fermentum and Lactobacillus salivarius in DSS mouse colitis:*
660   *impact on microRNAs expression and microbiota composition.* Mol Nutr Food
661   Res, 2017. **61**(11).

662   23.   .

663   24.   ., R.C.T. and *R: A language and environment for statistical*
664   *computing.* R Foundation for Statistical Computing, Vienna, Austria., 2013.

665   25.   Guo, X., et al., *Abnormal blood microbiota profiles are associated with*
666   *inflammation and immune restoration in HIV/AIDS individuals.* mSystems,
667   2023: p. e0046723.

668   26.   Mutlu, E.A., et al., *A compositional look at the human gastrointestinal*
669   *microbiome and immune activation parameters in HIV infected subjects.* PLoS
670   Pathog, 2014. **10**(2): p. e1003829.

671   27.   Yamamoto, S., et al., *The human microbiome and COVID-19: A systematic*
672   *review.* PLoS One, 2021. **16**(6): p. e0253293.

28.   Di Stadio, A., et al., *The Microbiota/Host Immune System Interaction in the Nose to Protect from COVID-19.* Life (Basel), 2020. **10**(12).

29.   Zhou, F., et al., *Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study.* Lancet, 2020. **395**(10229): p. 1054-1062.

30.   Mauvais-Jarvis, F., *Aging, Male Sex, Obesity, and Metabolic Inflammation Create the Perfect Storm for COVID-19.* Diabetes, 2020. **69**(9): p. 1857-1863.

31.   Aolymat, I., *A Cross-Sectional Study of the Impact of COVID-19 on Domestic Violence, Menstruation, Genital Tract Health, and Contraception Use among Women in Jordan.* Am J Trop Med Hyg, 2020. **104**(2): p. 519-525.

32.   Morgan, R., et al., *Using gender analysis matrixes to integrate a gender lens into infectious diseases outbreaks research.* Health Policy and Planning, 2022. **37**(7): p. 935-941.

33.   Robert, S.A., *Community-level socioeconomic status effects on adult health.* Journal of Health and Social Behavior, 1998. **39**(1): p. 18-37.

34.   Pijls, B.G., et al., *Temporal trends of sex differences for COVID-19 infection, hospitalisation, severe disease, intensive care unit (ICU) admission and death: a meta-analysis of 229 studies covering over 10M patients.* F1000Res, 2022. **11**: p. 5.

35.   Ritter, O. and G. Kararigas, *Sex-Biased Vulnerability of the Heart to COVID-19.* Mayo Clin Proc, 2020. **95**(11): p. 2332-2335.

36.   Wang, K., M. Gheblawi, and G.Y. Oudit, *Angiotensin Converting Enzyme 2: A Double-Edged Sword.* Circulation, 2020. **142**(5): p. 426-428.

37.   Pullen, A.B., et al., *Molecular and Cellular Differences in Cardiac Repair of Male and Female Mice.* J Am Heart Assoc, 2020. **9**(8): p. e015672.

38.   Wehbe, Z., et al., *Molecular and Biological Mechanisms Underlying Gender Differences in COVID-19 Severity and Mortality.* Front Immunol, 2021. **12**: p. 659339.

39.   Kermali, M., et al., *The role of biomarkers in diagnosis of COVID-19 - A systematic review.* Life Sci, 2020. **254**: p. 117788.

40.   Ranard, B.L., et al., *Identification of Endotypes of Hospitalized COVID-19 Patients.* Front Med (Lausanne), 2021. **8**: p. 770343.

41.   Rostami, M. and H. Mansouritorghabeh, *D-dimer level in COVID-19 infection: a systematic review.* Expert Rev Hematol, 2020. **13**(11): p. 1265-1275.

42.   Zhou, C., et al., *Increased Serum Levels of Hepcidin and Ferritin Are Associated*
708   *with Severity of COVID-19.* Med Sci Monit, 2020. **26**: p. e926178.

709   43.   Gomez-Pastora, J., et al., *Hyperferritinemia in critically ill COVID-19 patients -*
710   *Is ferritin the product of inflammation or a pathogenic mediator?* Clin Chim
711   Acta, 2020. **509**: p. 249-251.

712   44.   Wang, G., et al., *C-Reactive Protein Level May Predict the Risk of COVID-19*
713   *Aggravation.* Open Forum Infect Dis, 2020. **7**(5): p. ofaa153.

714   45.   Cooper, I.D., et al., *Relationships between hyperinsulinaemia, magnesium,*
715   *vitamin D, thrombosis and COVID-19: rationale for clinical management.* Open
716   Heart, 2020. **7**(2).

717   46.   Henry, B.M., et al., *Hematologic, biochemical and immune biomarker*
718   *abnormalities associated with severe illness and mortality in coronavirus*
719   *disease 2019 (COVID-19): a meta-analysis.* Clin Chem Lab Med, 2020. **58**(7):
720   p. 1021-1028.

721   47.   Honda, K. and D.R. Littman, *The microbiota in adaptive immune homeostasis*
722   *and disease.* Nature, 2016. **535**(7610): p. 75-84.

723   48.   Gupta, A., et al., *Nasopharyngeal microbiome reveals the prevalence of*
724   *opportunistic pathogens in SARS-CoV-2 infected individuals and their*
725   *association with host types.* Microbes Infect, 2022. **24**(1): p. 104880.

726   49.   Gao, M., et al., *Characterization of the Human Oropharyngeal Microbiomes in*
727   *SARS-CoV-2 Infection and Recovery Patients.* Adv Sci (Weinh), 2021. **8**(20): p.
728   e2102785.

729   50.   De Maio, F., et al., *Nasopharyngeal Microbiota Profiling of SARS-CoV-2*
730   *Infected Patients.* Biol Proced Online, 2020. **22**: p. 18.

731   51.   Braun, T., et al., *SARS-CoV-2 does not have a strong effect on the*
732   *nasopharyngeal microbial composition.* Sci Rep, 2021. **11**(1): p. 8922.

733   52.   Xu, R., et al., *Temporal association between human upper respiratory and gut*
734   *bacterial microbiomes during the course of COVID-19 in adults.* Commun Biol,
735   2021. **4**(1): p. 240.

736   53.   Wang, M., et al., *The relationship between gut microbiota and COVID-19*
737   *progression: new insights into immunopathogenesis and treatment.* Front
738   Immunol, 2023. **14**: p. 1180336.

54. Bozkurt, H.S. and O. Bilen, *Oral booster probiotic bifidobacteria in SARS-*
COV-2 patients. Int J Immunopathol Pharmacol, 2021. **35**: p.
20587384211059677.

55. Nardelli, C., et al., *Nasal Microbiome in COVID-19: A Potential Role of*
Corynebacterium in Anosmia. Curr Microbiol, 2022. **80**(1): p. 53.

56. Wei, N., et al., *Characterization of oral bacterial and fungal microbiome in*
recovered COVID-19 patients. BMC Microbiol, 2023. **23**(1): p. 123.

57. Bassis, C.M., et al., *The nasal cavity microbiota of healthy adults.* Microbiome,
2014. **2**: p. 27.

58. Perez-Losada, M., et al., *Nasal Bacteriomes of Patients with Asthma and*
Allergic Rhinitis Show Unique Composition, Structure, Function and
Interactions. Microorganisms, 2023. **11**(3).

59. Garcia-Vidal, C., et al., *Incidence of co-infections and superinfections in*
hospitalized patients with COVID-19: a retrospective cohort study. Clin
Microbiol Infect, 2021. **27**(1): p. 83-88.

60. Seibert, B., et al., *Mild and Severe SARS-CoV-2 Infection Induces Respiratory*
and Intestinal Microbiome Changes in the K18-hACE2 Transgenic Mouse
Model. Microbiol Spectr, 2021. **9**(1): p. e0053621.

61. Schippa, S., et al., *Nasal Microbiota in RSV Bronchiolitis.* Microorganisms,
2020. **8**(5).

62. Kageyama, Y., et al., *Lactobacillus plantarum induces innate cytokine responses*
that potentially provide a protective benefit against COVID-19: A single-arm,
double-blind, prospective trial combined with an in vitro cytokine response
assay. Exp Ther Med, 2022. **23**(1): p. 20.

63. Lopez-Santamarina, A., et al., *Probiotic Effects against Virus Infections: New*
Weapons for an Old War. Foods, 2021. **10**(1).

64. Ma, S., et al., *Metagenomic analysis reveals oropharyngeal microbiota*
alterations in patients with COVID-19. Signal Transduct Target Ther, 2021.
6(1): p. 191.

65. Merenstein, C., et al., *Signatures of COVID-19 Severity and Immune Response*
in the Respiratory Tract Microbiome. mBio, 2021. **12**(4): p. e0177721.

66. Mizutani, T., et al., *Correlation Analysis between Gut Microbiota Alterations*
and the Cytokine Response in Patients with Coronavirus Disease during
Hospitalization. Microbiol Spectr, 2022. **10**(2): p. e0168921.

67.   Mankowska-Wierzbicka, D., et al., *Alterations in Gut Microbiota Composition*
in Patients with COVID-19: A Pilot Study of Whole Hypervariable 16S rRNA
Gene Sequencing. Biomedicines, 2023. **11**(2).

68.   Ubeda, C., et al., *Intestinal microbiota containing Barnesiella species cures*
vancomycin-resistant Enterococcus faecium colonization. Infect Immun, 2013.
81(3): p. 965-73.

69.   Giacobbe, D.R., et al., *Enterococcal bloodstream infections in critically ill*
patients with COVID-19: a case series. Ann Med, 2021. **53**(1): p. 1779-1786.

70.   Wu, X., et al., *Coinfection with influenza virus and non-typeable Haemophilus*
influenzae aggregates inflammatory lung injury and alters gut microbiota in
COPD mice. Front Microbiol, 2023. **14**: p. 1137369.

71.   Murphy, E.C. and I.M. Frick, *Gram-positive anaerobic cocci--commensals and*
opportunistic pathogens. FEMS Microbiol Rev, 2013. **37**(4): p. 520-53.

72.   Chen, J., et al., *Comparison of the respiratory tract microbiome in hospitalized*
COVID-19 patients with different disease severity. J Med Virol, 2022. **94**(11): p.
5284-5293.

73.   Haran, J.P., et al., *Inflammation-type dysbiosis of the oral microbiome*
associates with the duration of COVID-19 symptoms and long COVID. JCI
Insight, 2021. **6**(20).

74.   Lu, S., et al., *Metatranscriptomic analysis revealed Prevotella as a potential*
biomarker of oropharyngeal microbiomes in SARS-CoV-2 infection. Front Cell
Infect Microbiol, 2023. **13**: p. 1161763.

75.   Seong, H., et al., *Clinical implications of gut microbiota and cytokine responses*
in coronavirus disease prognosis. Front Immunol, 2023. **14**: p. 1079277.

76.   Segata, N., et al., *Metagenomic biomarker discovery and explanation.* Genome
Biol, 2011. **12**(6): p. R60.

77.   Ning, Y., et al., *Comparative analysis of the gut microbiota composition*
between knee osteoarthritis and Kashin-Beck disease in Northwest China.
Arthritis Res Ther, 2022. **24**(1): p. 129.

78.   Mengyi, Z., et al., *Plasma metagenomics reveals regional variations of*
emerging and re-emerging pathogens in Chinese blood donors with an emphasis
on human parvovirus B19. One Health, 2023. **17**: p. 100602.

79. Li, S., et al., *Microbiome Profiling Using Shotgun Metagenomic Sequencing*
Identified Unique Microorganisms in COVID-19 Patients With Altered Gut
Microbiota. Front Microbiol, 2021. **12**: p. 712081.

80. Liu, Q., et al., *Gut microbiota dynamics in a prospective cohort of patients with*
post-acute COVID-19 syndrome. Gut, 2022. **71**(3): p. 544-552.

81. de Nies, L., et al., *Altered infective competence of the human gut microbiome in*
COVID-19. Microbiome, 2023. **11**(1): p. 46.

82. Xu, X., et al., *Integrated analysis of gut microbiome and host immune responses*
in COVID-19. Front Med, 2022. **16**(2): p. 263-275.

83. Lymberopoulos, E., et al., *COVID-19 severity is associated with population-*
level gut microbiome variations. Front Cell Infect Microbiol, 2022. **12**: p.
963338.

84. Thissen, J.B., et al., *Evaluation of co-circulating pathogens and microbiome*
from COVID-19 infections. PLoS One, 2022. **17**(12): p. e0278543.

85. Sulaiman, I., et al., *Microbial signatures in the lower airways of mechanically*
ventilated COVID-19 patients associated with poor clinical outcome. Nat
Microbiol, 2021. **6**(10): p. 1245-1258.

86. DeVoe, C., et al., *Increased rates of secondary bacterial infections, including*
Enterococcus bacteremia, in patients hospitalized with coronavirus disease
2019 (COVID-19). Infect Control Hosp Epidemiol, 2022. **43**(10): p. 1416-1423.

87. Devi, P., et al., *Longitudinal study across SARS-CoV-2 variants identifies*
transcriptionally active microbes (TAMs) associated with Delta severity.
iScience, 2023. **26**(10): p. 107779.

88. Moreira-Rosario, A., et al., *Gut Microbiota Diversity and C-Reactive Protein*
Are Predictors of Disease Severity in COVID-19 Patients. Front Microbiol,
2021. **12**: p. 705020.

89. Zhou, Y., et al., *Gut Microbiota Dysbiosis Correlates with Abnormal Immune*
Response in Moderate COVID-19 Patients with Fever. J Inflamm Res, 2021. **14**:
p. 2619-2631.

90. Parker, B.J., et al., *The Genus Alistipes: Gut Bacteria With Emerging*
Implications to Inflammation, Cancer, and Mental Health. Front Immunol,
2020. **11**: p. 906.

91. Li, J., et al., *Assessment of microbiota in the gut and upper respiratory tract*
associated with SARS-CoV-2 infection. Microbiome, 2023. **11**(1): p. 38.

92.   Ley, R.E., et al., *Microbial ecology: human gut microbes associated with*
obesity. Nature, 2006. **444**(7122): p. 1022-3.
841