Differential tissue deformability underlies shape divergence of the embryonic brain and spinal cord under fluid pressure

Susannah B.P. McLaren[1,2,*], Shi-Lei Xue[3], Siyuan Ding[1,2], Alex Winkel[2], Oscar Baldwin[1,2], Shreya Dwarakacherla[1,2], Kristian Franze[2,4], Edouard Hannezo[3], Fengzhu Xiong[1,2,*]

[1] Wellcome Trust / CRUK Gurdon Institute, University of Cambridge, Cambridge, UK

[2] Department of Physiology, Development and Neuroscience, University of Cambridge, Cambridge, UK

[3] Institute of Science and Technology Austria, Klosterneuburg, Austria

[4] Friedrich-Alexander-Universität Erlangen-Nürnberg, Germany; Max-Planck-Zentrum für Physik und Medizin, Germany

*Correspondence: sbm35@cam.ac.uk; fx220@cam.ac.uk

ABSTRACT

An expanded brain enables the complex behaviours of vertebrates that promote their adaptation in diverse ecological niches[1–3]. Initial morphological differences between the brain and spinal cord emerge as the antero-posteriorly patterned neural plate folds to form the neural tube[4–7] during embryonic development. Following neural tube closure, a dramatic expansion of the brain diverges its shape from the spinal cord[8], setting their distinct morphologies for further development[9,10]. How the brain and the spinal cord expand differentially remains unclear. Here, using the chicken embryo as a model, we show that the hindbrain expands through dorsal tissue thinning under a positive hydrostatic pressure from the neural tube lumen[11,12] while the dorsal spinal cord shape resists the same pressure. Using magnetic droplets and atomic force microscopy, we reveal

that the dorsal tissue in the hindbrain is more fluid than in the spinal cord. The dorsal

hindbrain harbours more migratory neural crest cells[13] and exhibits reduced apical actin

and a disorganised laminin matrix compared to the dorsal spinal cord. Blocking the activity

of neural crest-associated matrix metalloproteinases inhibited dorsal tissue thinning,

leading to abnormal brain morphology. Transplanting early dorsal hindbrain cells to the

spinal cord was sufficient to create a region with expanded brain-like morphology

including a thinned-out roof. Our findings open new questions in vertebrate head

evolution and neural tube defects, and suggest a general role of mechanical pre-pattern in

creating shape differences in epithelial tubes.

MAIN

The neural tube gives rise to the brain and spinal cord during vertebrate embryonic

development. In avian embryos, the volumetric expansion of the brain over a relatively

stable spinal cord takes place shortly after neural tube closure (Figure 1A) and depends on

the hydrostatic pressure from the neural tube lumen[14,11]. In the stages prior to brain

expansion (before Hamburg-Hamilton (HH) stage 13[8]), the neural tube has large openings at

both the anterior and posterior ends that allow fluid to move between the tube lumen and

its surroundings (Figure 1B). These openings narrow and close concomitant with the onset

of brain expansion around HH13[15], such that the neural tube lumen becomes a closed and

continuous fluid filled cavity (Figure S1A). Consistent with pioneering studies[16], we recorded

a positive internal lumen pressure of ~15Pa in the initial stages of brain expansion (HH13-

15), which increased to ~25Pa as brain expansion progressed (HH19-21. Figure 1C and S1B).

To assess how neural tube morphology changes as internal lumen pressure increases, we

measured tissue thickness around the circumference of the lumen in cross sections of the

hindbrain and spinal cord prior to and after brain expansion (HH11 and HH16, respectively.

Figure 1D-F). Dorsal hindbrain tissue approximately halved in thickness and formed a single-

cell thick epithelium concomitant with brain expansion, whilst little dorsal thinning or tissue

shape change was observed in the spinal cord (Figure S1C and D).

To test the role of lumen pressure in dorsal thinning, we intubated[14] embryos in the brain or

spinal cord just prior to brain expansion (HH11), leaving the tube lumen connected to the

embryo's surroundings to equate the pressure (Figure S1E). Although we observed an

increase in hindbrain cross-sectional area, in line with proliferation and tissue growth, dorsal

hindbrain thinning and expansion were inhibited in intubated embryos following 20 hours of

incubation (Figure 1G, I and J). Not only brain-intubated but also a spinal cord-intubated

embryo showed inhibited hindbrain thinning, confirming that the lumen is connected along

the anterior-posterior axis of the neural tube and the tissues experience the same

hydrostatic pressure. Conversely, treating embryos with β-D-xyloside (BDX) [17], which has

been found to osmotically increase lumen pressure by ~30% [12], led to an increase in

expansion and extension of the thinned dorsal roof in the hindbrain and a moderately

enlarged spinal lumen with no dorsal thinning (Figure 1H, K and L). Together these findings

demonstrate that neural tube lumen pressure drives dorsal tissue thinning in the hindbrain

but is resisted in the dorsal spinal cord, leading to brain expansion relative to the spinal cord.

The differential thinning of dorsal tissue between the hindbrain and the spinal cord may

result from a few scenarios. First, different initial tissue curvature at the onset of pressure

increase (Figure 1D) can lead to diverging dynamics of thinning and volume expansion. To

explore this possibility, we formulated a physical model describing the neural tube as a

cellularised, homogeneous material under internal pressure (Supplemental Text 1). The

model predicted that regions of the tube with a greater initial radius (lower curvature) will

expand more in the short-term than parts of the tube with an initially smaller radius (higher

curvature) as circumferential stress will be higher in low curvature regions (Figure 2A and A'),

which is consistent with the shape dynamics observed in the hindbrain and the spinal cord.

To experimentally test this possibility, we modified spinal cord lumen shape prior to brain

expansion by performing *in ovo* surgeries to remove somites and anterior presomitic

mesoderm on either side of the spinal cord (Figure 2B). This operation releases tissue

confinement from not only the somites (Figure S2A) but also the dorsal non-neural

ectoderm as it was cut through during somite ablation. After ~20 hours of incubation, these

embryos healed and exhibited reduced spinal cord dorsal curvature and in one case clear

rounding of the spinal cord cross section (Figure 2C, S2B-C). However, this did not lead to the

formation of a thinned-out roof, as was observed in the hindbrain (Figure S2D). These results

suggest that the higher curvature of the spinal cord dorsal region is not sufficient to explain

the lack of dorsal tissue thinning in the spinal cord. At the same time, this experiment ruled

out a second alternative model in which local confinement from neighbouring tissues (such

as somites and the presomitic mesoderm) might modulate neural tube shape ([18] and Figure

S2A) by constraining lumen expansion of the spinal cord relative to the hindbrain. This led us

to explore a third hypothesis: that the differential expansion of the brain and spinal cord was

due to the dorsal tissue of the hindbrain being more deformable than that of the spinal cord,

enabling local thinning despite a spatially uniform hydrostatic pressure. To test whether the

material properties of the hindbrain and spinal cord dorsal tissue differed before the onset

of increased lumen pressure, we injected ferrofluid oil droplets into the forebrain of HH11

embryos and positioned the droplets in either the hindbrain or spinal cord using a magnetic

field (Figure 2D). We then observed the resulting tissue deformation from the stress

generated by the droplets as they minimize surface tension after removal of the magnetic

field (Figures 2E). Droplets were injected either at equal volumes (2nl) or at a ratio to

account for the difference in initial lumen size (4nl in the larger hindbrain lumen, 1nl in the

smaller spinal cord lumen). In both cases the droplets showed a higher curvature at the

droplet-lumen interface in the spinal cord, indicating a higher pressure was exerted on the

spinal cord tissue than the hindbrain by this mechanical perturbation (Figure S2E). Still, the

thickness change of dorsal tissue at the droplet site was greater in the hindbrain than in the

spinal cord (Figure 2F), indicating that hindbrain dorsal tissue was more deformable. To

directly measure tissue mechanical properties, we used an atomic force microscopy (AFM)

cantilever to indent the dorsal region of the hindbrain and spinal cord (Figure 2G) in the

same embryos and measure tissue relaxation in response to a sustained force (50 or 75nN).

We found that the dorsal hindbrain had a higher fluidity than that of the spinal cord prior to

brain expansion, whilst no significant difference in instantaneous elastic modulus was

detected (Figure 2H and I). While *in vivo* AFM measurements may contain a contribution of

differences from other non-local structures such as the extracellular matrix (ECM) anchoring

the neural tube, these results further support the hypothesis that the dorsal hindbrain will

deform more than the dorsal spinal cord under the same lumen pressure.

To investigate the mechanism of dorsal tissue deformability, we explored the cellular

organisation of the tissue at stages leading up to brain expansion (HH9-12). The dorsal

neural tube tissue is occupied by neural crest cells that reorganise their actomyosin

cytoskeleton as they undergo an epithelial-to-mesenchymal transition (EMT) and delaminate

from the tube dorsal surface as the walls of the neural plate fuse dorsally [19,20,21]. The

dynamics of neural crest delamination between cranial and trunk regions is known to differ

[22,23], making them an exciting candidate for underlying tissue intrinsic differences between

the dorsal hindbrain and spinal cord. We observed clear apical organisation of actin and

myosin within the dorsal spinal cord in HH12 embryos, however this organisation was largely

absent in the neural crest domain of the hindbrain (Figure 3A-C). This change in cytoskeleton

organisation indicated that a transition away from an epithelial cell state to a more

mesenchymal cell state was occurring in the dorsal hindbrain. A key player in this process is

the zinc-finger transcription factor Snail2 which is expressed in the pre-migratory and early

migrating neural crest [24,25]. Using Snail2 as a marker, we observed that the dynamics of

neural crest delamination and migration at the dorsal surface of the neural tube differed

between the hindbrain and the spinal cord leading up to brain expansion (HH9-12), with

Snail2+ cells migrating in a cell-by-cell fashion in spinal cord regions, whilst those in the

hindbrain migrate in a collective wave (Figure 3D and S3A). Neural crest cells utilise matrix

metalloproteases (MMPs) to remodel the surrounding ECM [26,27]. ECM architecture plays an

important role in determining tissue material properties, with higher levels of ECM

organisation conferring greater stiffness (or elasticity) and more disordered matrix

organisation leading to softening (or a more viscous behaviour) [28,29], which contributes to

long-term deformability.

This led us to investigate ECM organisation in the hindbrain and the spinal cord as a potential

mediator of tissue deformability. Cross-sectional views showed that laminin was

redistributed throughout the dorsal hindbrain in the premigratory neural crest domain,

whilst comparatively little redistribution was observed in the dorsal spinal cord (Figure 3E).

This was associated with a more sparce laminin coating of the hindbrain compared to the

spinal cord when viewed from the dorsal side (Figure 3F and S3B and C). These findings

suggest that ECM is being actively reorganised and redistributed out of the plane of the

basement membrane to a greater extent in the hindbrain.

To test whether ECM remodelling plays a role in dorsal hindbrain thinning, we inhibited

MMP activity in embryos prior to brain expansion stages using an MMP-specific small

molecule inhibitor (GM6001, 1000μM). Brain expansion was reduced in inhibitor treated

embryos and a smaller surface area of the dorsal hindbrain was apparent (Figure 3G-H). In

embryos with obvious observable developmental defects (0/6 control, 4/8 inhibitor treated),

the hindbrains did not undergo dorsal thinning and in some cases aggregated a mass of

Snail2+ cells within the lumen (Figure S3D). In embryos that developed normally overall (6/6

control, 4/8 inhibitor treated), the hindbrains of the treated embryos showed thicker or

shorter dorsal roofs than those of the controls (Figure 3I, J and S3D). The laminin surface

covering the basal extent of the dorsal hindbrain appeared less disrupted in embryos with

inhibited roof thinning (Figure 3I) and the dorsal tissue was still populated with Snail2+ cells,

suggesting that neural crest specification continued in the presence of the inhibitor.

Together, these data are consistent with the idea that neural crest mesenchymal behaviour

and subsequent ECM remodelling is required for thinning and expansion of the dorsal

hindbrain.

To definitively test the intrinsic ability of early hindbrain dorsal cells to generate a thinned

roof under lumen pressure, we performed an *in ovo* graft experiment, moving a small piece

of dorsal hindbrain (or spinal cord as a control) tissue from a GFP+ embryo to the pre-cut

dorsal region of the spinal cord of a wildtype host before the onset of lumen pressure

(Figure 4A). Re-integration of the graft to reform a closed neural tube that would continue

normal morphogenesis was expected to be rare given the difficulty of positioning the graft in

such a small incision site and maintaining contact between the small graft and the host

tissue. After overnight culture, we found 5 embryos with different degrees of hindbrain

dorsal tissue integration into the spinal cord, and 2 spinal cord-to-spinal cord controls (Figure

4B, S4A and S4B). The control embryos exhibited a thick dorsal region at the grafted site

(Figure 4C, D and S4B, 2/2 embryos). Interestingly, 2 hindbrain-to-spinal cord grafted

embryos exhibited dorsal tissue thinning in the grafted region (Figure 4C, D and S4B). In the

cleanest graft where the donor tissue completely replaced the host dorsal region, we found

a distinctively thinned out dorsal roof and hindbrain-like morphology specific to the grafted

region (Figure 4C), with a thicker spinal cord-like host dorsal region present on the anterior

and posterior sides of the graft. Cells in the grafted region were Snail2+ (Figure 4E) and the

thinned-out roof coincided with loss of apical actin localisation (Figure S4C), suggesting that

the grafted region contained premigratory neural crest cells undergoing EMT. This striking

ability of Snail2+ dorsal hindbrain cells to form a thinned-out and elongated roof in the

spinal cord region suggests that hindbrain premigratory neural crest cells are sufficient to

drive dorsal tissue thinning in response to lumen pressure.

Taken together, our findings support a mechanism where deformability of the neural tube

dorsal tissue and ECM organisation underlie the distinct shape changes of the brain and the

spinal cord under a shared lumen pressure (Figure 4F). These differences likely result from

different behaviours of the neural crest cells, which are known to exhibit different signalling,

metabolism and fates between cranial and trunk levels. Our study thus implies a closely

coordinated initial developmental step coupling brain expansion and neural crest EMT, both

essential contributors to the formation and evolution of the vertebrate head [30]. We note

that these findings are in agreement with classic chimera grafting experiments in which

reduced dorsal roof thinning and elongation can be observed in hindbrain regions lacking

neural crest cells [31]. Irregularly shaped mesenchymal cells in the dorsal hindbrain have also

been observed to contribute to the formation of a thinned out hindbrain roof as the narrow

lumen of the zebrafish brain undergoes expansion [32], suggesting that neural crest mediated

dorsal tissue fluidisation may contribute to brain expansion in other species. Future work

will resolve the detailed molecular and cellular regulation of dorsal neural tissue mechanics

and identify genetic and environmental factors that may cause developmental defects

associated with aberrant brain expansion through this mechanism. Given the role of lumen

pressure in driving morphogenesis of a variety of epithelial tissues and organs [33,34,35,36] our

findings here in the neural tube imply a general strategy for creating diverse biological

shapes via a mechanical property pre-pattern before the onset of changes in fluid pressure.

ACKNOWLEDGEMENTS

We thank A. Dimitracopoulos, K. Kawaguchi, J. Vidigueira, B. Baum, I. McLaren and members

of the Buckley, Scarpa, Steventon, Kawaguchi and Xiong labs for technical assistance and

constructive feedback. We thank Ryan Greenhalgh for methods developed to obtain fluidity

values from AFM data. We thank Nicola Lawrence, Alex Sossick and Sargon Gross-Thebing

from the Gurdon Institute Imaging Facility for microscopy support. This work was supported

by a Wellcome Trust / Royal Society Sir Henry Dale Fellowship (215439/Z/19/Z) and an UKRI-

EPSRC Frontier Research Grant (EP/X023761/1, originally selected as an ERC Starting Grant)

to F.X.

AUTHOR CONTRIBUTION

S.B.P.M and F.X. conceived the project, designed experiments and wrote the manuscript.

S.B.P.M performed all experiments and analysed the data. S.X and E.H. developed the

theoretical model. S.D, O.B, S.D and F.X contributed to embryo experiments. K.F. and A.W.

designed and performed AFM experiments.

MATERIALS & METHODS

Egg lines and embryo culture. Wild type fertilised chicken eggs were obtained from MedEgg

Inc., GFP+ eggs (Cytoplasmic [37]and Membrane[38]) were obtained from the National Avian

Research Facility (NARF) at University of Edinburgh. Eggs were kept in a fridge at 14°C for

storage and in a 37.5°C humidified (40%-60%) incubator (Brinsea) during incubation. No

animal protocol under the UK Animals (Scientific Procedures) Act 1986 was required for the

chicken embryo stages investigated (incubated under 2 weeks, or 2/3 of the gestation time).

Ex ovo culture was performed using a modified EC culture[39] protocol as described[40]. For *in*

ovo operations, egg shells were windowed with surgical scissors after removing some

albumen. A small volume of Chinese calligraphy ink (YiDeGe, Black) was then injected

underneath the embryo to help with visualisation. After performing surgical, mechanical or

chemical perturbations, eggs were resealed with clear plastic tape and returned to

incubation.

Pressure measurement. Pressure measurements were performed as detailed in [41]. Briefly, a

glass microneedle connected to a pressure sensor was inserted into the neural tube lumen

of chicken embryos at varying stages of development. Pressure readings were continuously

recorded to include the pressure just outside the lumen, after insertion into the lumen, and

after retraction from the lumen for each experiment. The pressure difference between the

outside and inside of the lumen was taken as the neural tube lumen pressure.

Measurements where no stable pressure reading could be obtained inside the neural tube

lumen, likely due to needle clogging by tissue debris during the insertion, were excluded

from analysis.

Pressure modulation. Intubation experiments were performed by inserting an open-ended

short glass capillary tube into the forebrain or spinal cord region of HH10-12 stage embryos.

Control experiments were performed by either inserting a glass capillary into tissues just

neighbouring the neural tube or inserting a very narrow glass capillary and not submerging it

in fluid with the embryo. β-D-xyloside (BDX) experiments were performed by adding either

DMSO or BDX to embryo culture medium at a final concentration of 2mM in L15 media. In all

cases embryos were cultured overnight *ex ovo*. Only embryos that displayed continued

development were included in the analysis.

Assessment of lumen continuity. Fluorescently labelled Dextran was injected into the neural

tube lumen of HH11-13 stage embryos. The fluorescent signal was imaged following

injection to assess how far the dye travelled along the neural tube lumen. The dye was

observed to spread anteriorly into the brain region and posteriorly into the posterior spinal

cord when injected into the spinal cord lumen mid-way along the trunk.

Removal of confining tissues. Posterior somites and anterior presomitic mesoderm were

removed from embryos *in ovo* using a sharpened tungsten needle. The vitelline membrane

was peeled back and cuts were made at the interface between the spinal cord region of the

neural tube and posterior somites and anterior presomitic mesoderm. Somites and

presomitic mesoderm were peeled away to leave the posterior spinal cord unconfined on its

sides. Eggs were then resealed and incubated overnight. Embryos that showed a good

degree of healing and no obvious spinal cord ruptures were included in analysis.

<u>Grafting experiments.</u> Small patches of dorsal hindbrain or spinal cord cells from GFP

positive HH10-11 embryos were transplanted into the posterior spinal cord of stage-

matched wild type embryos. A sharpened tungsten needle was used to peel back the

vitelline membrane of donor and host embryos and to dissect out a small patch of dorsal

cells from either the hindbrain or spinal cord. Tissue patches were transferred using a Gilson

pipette and Ringers solution to the host embryos. A small incision was made into the

posterior host embryo dorsal neural tube and tissue patches were pushed into this incision.

Embryos were cultured *in ovo* overnight.

<u>Ferrofluid droplet experiments.</u> Ferrofluid oil droplets were injected into the forebrain region

of HH10-12 stage embryos using a Nanojet injector set to inject a volume of either 1, 2, or

4nl. Droplets were positioned in the lumen of either the hindbrain or spinal cord using a

magnet and custom-built embryo holder. Embryos were fixed approximately 1.5 hours post

droplet injection prior to immunostaining and confocal imaging or imaged live. Dorsal roof

tissue thickness was measured at the point of maximum roof deformation in the hindbrain

and spinal cord of injected embryos at the droplet location. Uninjected embryos at

equivalent stages were measured at the same anterior-posterior level as controls.

<u>Atomic force microscopy.</u> Atomic force microscopy (AFM) indentation measurements were

performed using the setup described in [42]. Briefly, HH10-12 stage chicken embryos were

cultured on agarose plates and the vitelline membrane was removed to expose the embryo's

dorsal hindbrain and spinal cord. 89.3 µm diameter polystyrene beads were glued to Arrow-

TL1 cantilevers (nominal spring constant = 0.01 N/m; NanoWorld, Neuchâtel, Switzerland) as

probes. During measurements, the cantilever was approached with 10 µm/s until an

indentation force of either 50nN or 75nN was reached. This force was maintained for 3

seconds, thus enabling the acquisition of the deformation over time ('creep').

In order to extract fluidity and elastic modulus values from the force-distance curves, [43]

introduced the following piecewise model for the force ramp:

$$
F(t) = F_C \begin{cases} 0 \; ; & t < t_C \\ \left(\dfrac{t - t_C}{\Delta t_A}\right)^{\alpha} \; ; & t_C \leq t < t_C + \Delta t_A \\ 1 \; ; & t \geq t_C + \Delta t_A \end{cases}
$$

where F_C is the magnitude of the applied force during the creep measurement reached in

time Δt_A and applied for time t_C. α describes the shape of the force ramp ($1 < \alpha < 2$), where α

= 1 corresponds to a linear force ramp and α closer to 2 to a polynomial force ramp. Fitting

this equation yields F_C, t_C, Δt_A, and α . These values then derive the following equation for

the indentation δ, here modified by substituting $E_0/(1- \nu^2)$ with the reduced instantaneous

elastic stiffness, k_0

$$
\delta(t) = \left[\frac{3}{4} \frac{1}{k_0 \sqrt{R}} F_C \frac{\alpha (t - t_C)^{\alpha + \beta}}{\Delta t_A^{\alpha} t_0^{\beta}} B\left(\frac{\Delta t_A}{t - t_C}; \alpha, \beta + 1\right) \right]^{2/3}
$$

where R is the radius of the bead, $B()$ is the incomplete beta function and the exponent, and

β is the fluidity, which can have values ranging from 0 (for an elastic solid) to 1 (for a viscous

fluid). Mean values for k_0 and β were determined from measurements taken across the

hindbrain and along the spinal cord. Values obtained for k_0 were close to values obtained for

the reduced apparent elastic modulus $K = E_0/(1- \nu^2)$ obtained by fitting the Hertz model

$F = \frac{4}{3} K \delta^{3/2} \sqrt{R}$ to the instantaneous indentation data (Figure. S2).

<u>MMP inhibitor treatment.</u> After removal of the vitelline membrane covering the brain, 20µl

of 1000µM broad-spectrum MMP inhibitor (GM6001) or DMSO diluted in Ringers solution,

was added to the dorsal surface of HH10-11 stage embryos *in ovo.* Embryos were incubated

overnight.

<u>Immunostaining.</u> After fixing embryos overnight in 500μl 4% PFA at 4°C, they were washed

3X for ~20 minutes per wash in 0.5% Triton-X- 100 (Sigma-Aldrich, X100) in PBS (PBST).

Embryos were incubated in blocking solution consisting of 4% Donkey serum in PBST for 1

hour at room temperature. Primary antibodies were diluted in blocking solution and

incubated with embryos for 2 days at 4°C. After washing embryos in PBST ~4 times over the

course of 2 hours, secondary antibodies, Hoecst and Phalloidin were diluted in blocking

solution and incubated with embryos for another 2 days at 4°C. Finally, embryos were

washed in PBST prior to imaging.

<u>Primary and secondary antibodies used.</u> Laminin (DSHB, 3H11) 1:100, Snail2 (Cell Signaling

Technology, 9585S) 1:200, pMLC2 (Cell Signaling Technology, 3671S) 1:100, Hoecst 1:1000,

Phalloidin 647 (Thermo Scientific, A22287) 1:500, Donkey Anti-Mouse 488n IgG (Abcam,

ab150105), Donkey Anti-Rabbit 594 (Abcam, ab150076)

<u>Confocal imaging.</u> Embryo cross sections and wholemounts were mounted on glass-bottom

confocal imaging dishes in VECTASHIELD (Vector Labs, H-1000-10) mounting medium. Z-

stacks were acquired at either 0.3 or 0.5μm z-steps using either an SP8 (Leica) or a SoRa

Spinning Disk (Nikon) using either a 10X, 20X or 40X objective.

<u>Data analysis.</u> All data analysis was performed in Python (www.python.org), with data stored

in Excel. Boxplots were obtained using the seaborn library and scatterplots using matplotlib.

Statistical analysis was performed using the scipy.stats Python library. Paired t-tests were

used to calculate p-values for the AFM data. Independent t-tests were used to calculate all

other p-values. Details of measurements of reported features are listed below:

1. Tissue shape analysis. The thickness profile of neural tube cross-sections was acquired by

measuring the radial tissue thickness along the lumen circumference starting from the

approximate mid-point of the dorsal hindbrain roof. For wildtype characterisation, the

distance along the lumen circumference was normalised for all samples so that tissue

thickness at distance '0' fell at the dorsal mid-point and tissue thickness at distance '1' at the

ventral midpoint of the neural tube cross section. A 4^{th} degree polynomial was fit to the

thickness profiles to get an 'average' thickness profile for the hindbrain and spinal cord at

HH11 and HH16. Images acquired by confocal microscopy were visualised using either

ImageJ[44] or napari[45]. Cross sections were either visualised directly, or by reordering the axes

of image data acquired from wholemount samples in napari. For intubation experiment

analysis the mean thickness of the first 100µm of dorsal roof tissue from the dorsal midpoint

was compared. For BDX experiment analysis the mean thickness of the first 300µm of dorsal

roof tissue from the dorsal midpoint was compared. Equivalent positions along the hindbrain

were compared for assessing hindbrain roof thickness in MMP treated and control embryos.

2. Actin intensity along the apical surface. Curves were drawn along the apical surface of

neural tube cross sections starting from the dorsal midpoint and moving ventrally. The actin

intensity along the curve was measured using the ImageJ PlotProfile function. Normalised

average actin intensities along the apical surface were obtained for each embryo and binned

into three groups starting from the dorsal-most position and moving more ventrally. The

mean actin intensity for each embryo within each bin was plotted.

3. Somite removal analysis. Z planes were selected in the region where somite removal had

been performed and in a corresponding location in controls. Roof thickness was measured at

the dorsal midpoint for z planes from each embryo.

SUPPLEMENTAL TEXT 1

In this Supplementary note, we provide detailed analysis on morphological evolution of neural tube driven by luminal pressure. The brain and the spinal cord expand differentially under lumen pressure.

1.   Elastic deformation of neural tube

The anterior and posterior ends of the neural tube get closed around Hamburg-Hamilton (HH) stage 13. After that, fluids accumulate in the enclosed lumen and tend to inflate the neural tube. To check if differences in the initial geometry and mechanical properties (such as elastic modulus) can explain the remarkable morphological differences between the hindbrain and spinal cord, especially the distinct radial expansion, after the neural tube pressurized with luminal fluids, here we extend previous model on the deformation of elastic tube after inflation[46] by including initial geometrical and mechanical inhomogeneities. The neural tube is modelled as an elastic, three-dimensional cylindrical membrane enclosed with a lumen filled with fluids. For simplicity, we consider initial shape of the neural tube is axisymmetric about A-P axis (defined as the Z-axis in Figure S5), but can be inhomogeneous along A-P axis: the initial radius $R = R(Z)$. The initial tissue thickness of the neural tube is assumed to be a constant H_0. Besides the axial and radial coordinates (R, Z) for the initial undeformed configuration, we also introduce the coordinates (r, z) for the current deformed configuration, as shown in Figure S5

Notice in both the undeformed and deformed configurations of the neural tube, the tissue surface is not necessarily parallel to the A-P axis (i.e. Z- or z-axis), but can derivate from it (see Figure S5 for schematic, where ϕ is the angle between the deformed tissue and z-axis). In this scenario, to evaluate the tissue stretching in the meridional direction, we introduce S and s to respectively track positions along the undeformed and deformed meridians.

376   First, we discuss the geometry of the deformed neural tube. The geometric variables in the

377   deformed configuration satisfy

378
$$\frac{\mathrm{d}r}{\mathrm{d}s} = \sin\phi, \quad \frac{\mathrm{d}z}{\mathrm{d}s} = \cos\phi, \tag{1}$$

379   and the two principal curvatures of the deformed tissue are given by

380
$$\kappa_1 = -\frac{\mathrm{d}\phi}{\mathrm{d}s}, \quad \kappa_2 = \frac{\cos\phi}{r}, \tag{2}$$

381   where the subscript "1" and "2" respectively represent the meridional and azimuthal

382   directions.

383   Second, we consider the mechanics of the neural tube. Pressurized with a luminal pressure,

384   the tissue of the neural tube undergoes axial stretching λ_1 and radial expansion, which can

385   be respectively quantified by the stretch ratios in the meridional and azimuthal directions:

386
$$\lambda_1 = \frac{\mathrm{d}s}{\mathrm{d}S}, \quad \lambda_2 = \frac{r}{R}. \tag{3}$$

387   The stretch ratio λ_α larger (or smaller) than 1 corresponds to tissue stretching (or

388   compression) in that direction. Besides, such information on tissue deformation allows us to

389   obtain the current thickness of the tissue H, which satisfies $H = H_0/(\lambda_1\lambda_2)$. To balance the

390   luminal pressure P, the tissue forces T_1 and T_2 yield

391
$$T_1 = \frac{1}{2}\frac{Pr}{\cos\phi}, \tag{4}$$

392
$$T_1\kappa_1 + T_2\kappa_2 = P. \tag{5}$$

The relation between the tissue forces (T_1 and T_2) and tissue deformation (λ_1 and λ_2) is an

intrinsic property of a tissue (or material). Such relation can be conveniently deduced by

considering the stored elastic energy density of the tissue W. For incompressible materials

such as soft tissue, $W = \mu f(\lambda_1, \lambda_2)$, with μ the initial shear modulus (a parameter used to

quantify tissue stiffness). To include mechanical inhomogeneity in the hindbrain and spinal

cord, here we consider $\mu = \mu(Z)$. The tissue forces are related to tissue deformation as $T_1 =$

$\frac{H_0}{\lambda_2}\frac{\partial W}{\partial \lambda_1}$, $T_2 = \frac{H_0}{\lambda_1}\frac{\partial W}{\partial \lambda_2}$. Hereafter, we use $w_\alpha = \partial W/\partial \lambda_\alpha$ for notational convenience.

The above equation system (1)-(5) can be recast as

$$\frac{d\lambda_1}{dZ} = \frac{1}{Rw_{11}}[\sin\phi\,(w_2 - w_{12}\lambda_1)S' + (w_{12}\lambda_2 - w_1)R'] - \frac{1}{w_{11}}\frac{\partial w_1}{\partial \mu}\mu',$$

$$\frac{d\lambda_2}{dZ} = \frac{1}{R}(\sin\phi\,\lambda_1 S' - \lambda_2 R'),$$

$$\frac{d\phi}{dZ} = \frac{\cos\phi}{R}\left(\frac{w_2}{w_1} - 2\frac{\lambda_1}{\lambda_2}\right)S',$$

$$\frac{dz}{dZ} = \lambda_1 \cos\phi\, S'. \tag{6}$$

where $R' = dR/dZ$, $S' = dS/dZ$, and $\mu' = d\mu/dZ$.

In the end, the elastic deformation of the pressurized neural tube is controlled by the

differential equation system (6) with four unknowns: two stretch ratios λ_1 and λ_2, the angle

ϕ, and the axial position z.

In the development and morphogenesis of the neural tube, the brain region gradually bulges

up and the spinal cord stays in a slim shape. We choose the middle point of the brain as the

starting point of our calculation, and set $z(0) = 0$. Due to the symmetry of the bulge, we

have $\phi(0) = 0$. The posterior end of the spinal cord can be approximated as $\lambda_2(L) = 1$,

which would not affect the theoretical profile of the neural tube.

Figure 1. Neural tube lumen pressure drives hindbrain and spinal cord shape divergence

(A) Schematic depicting brain expansion during chicken embryo development, Hamburger and

Hamilton stages of development are given. (B-F) Lumen pressure increase and dorsal tissue thinning

occur concomitantly with brain expansion. (B) Bright-field images of chicken embryos prior to brain

expansion (HH10) and following brain expansion (HH16) and confocal microscopy images of the early

anterior and posterior neuropore at the level of the white boxes. (C) Pressure readings obtained by

inserting a microneedle connected to a pressure sensor into the neural tube lumen in embryos at the

onset of and following brain expansion. (D) Schematic showing tissue thickness measurement

approach and confocal microscopy images of neural tube cross-sections taken at the level of the

dashed white lines in (B). (E) Apical-basal tissue thickness along the hindbrain lumen circumference,

n=6 hh11 embryos, n=3 HH16 embryos. (F) Apical-basal tissue thickness along the spinal cord lumen

circumference, n=6 hh11 embryos, n=3 HH16 embryos. (G-L) Pressure decrease and increase

experiments. (G) Bright-field images of control and intubated embryos at ~20 hours post-intubation

and confocal images of corresponding hindbrain and spinal cord cross-sections. (H) Bright-field

images of control and β-D-xyloside (BDX) treated embryos ~20h post treatment and corresponding

confocal images of hindbrain and spinal cord cross-sections. (I) Tissue thickness along the hindbrain

circumference moving from dorsal to ventral in control, anterior intubated and spinal cord intubated

embryos. (J) Hindbrain roof thickness (averaged over the first 100µm from the dorsal midpoint in

control (n=6), anterior intubated (n=2) and spinal cord intubated (n=3) embryos. (K) Tissue thickness

along the hindbrain circumference in control and BDX treated embryos. (L) Hindbrain total roof

length in control (n=5) and BDX treated (n=4) embryos.

Black scale bars are 500µm, white scale bars are 50µm unless otherwise stated.

Figure 2. The dorsal hindbrain is more deformable than the dorsal spinal cord prior to

brain expansion

(A-A') A physical model predicts a greater degree of expansion in regions with initially larger radius. A

cellularised model of an epithelial tube was used to determine how the initial radius at a point along

the tube may affect its expansion dynamics under internal pressure. The anterior neural tube is

positioned at z = 0. (B-C) Lateral tissue removal experiments. (B) Bright-field image of a pre-brain

expansion stage embryo with posterior somites and anterior presomitic mesoderm removed (n=4,

control n=3). (C) Confocal images of the dorsal surface view and cross-section view of the spinal cord

in a ~20h post somite-removal embryo. (D-F) Dorsal tissue deformation under ferrofluid droplet

injection. (D) Widefield images of HH11 stage embryos with ferrofluid droplet injections in either the

hindbrain or spinal cord lumen. (E) Confocal images showing cross-sectional views of the spinal cord

and hindbrain in HH10-11 stage uninjected embryos (n=6 spinal cord, n=6 hindbrain) and embryos

with ferrofluid droplets injected into either the hindbrain or spinal cord (n=6 hindbrain, n=8 spinal

cord). White arrows indicate thickness measurements (acquired at the thinnest dorsal point

observable). (F) Quantification of hindbrain and spinal cord roof thickness in uninjected and droplet

injected embryos. (G-I) Atomic force microscopy experiments. (G) Schematic depicting atomic force

microscopy measurement set up. (H) Mean Beta value (corresponding to fluidity) of the hindbrain

and spinal cord in HH10, 11 and 12 embryos (n=9). (I) Mean instantaneous elastic modulus of the

hindbrain and spinal cord in HH10, 11 and 12 embryos (n=9).

Black scale bars are 500µm. White scale bars are 50µm.

Figure 3. Differential neural crest cell behaviour underlies the acquisition of a thinned out and expanded dorsal hindbrain roof

(A-C) Actomyosin localisation in the dorsal hindbrain and spinal cord. (A) Confocal images of actin and phosphorylated myosin light chain kinase II in the hindbrain and spinal cord of HH12 stage embryos. (B) Quantification of apical actin signal intensity along the lumen circumference in the hindbrain of HH11 and HH12 stage embryos. (C) Quantification of apical actin signal intensity along the lumen circumference in the spinal cord of HH11 and HH12 stage embryos. (D) Premigratory neural crest in the hindbrain and spinal cord. Confocal images showing Snail2+ cell migration in the

hindbrain and spinal cord cross sections of embryos progressing towards brain-expansion stages

(HH9 to HH12). (E-F) ECM organisation in the dorsal hindbrain and spinal cord. (E) Confocal images

showing laminin and actin organisation in cross sections of the spinal cord and hindbrain of HH12

stage embryos. F) Confocal images showing laminin organisation at the dorsal surface in a 3D

rendering of the spinal cord and hindbrain of HH11 stage embryos. (G-H) ECM remodelling

perturbation experiments. (G) Brightfield images of embryos treated with DMSO or MMP inhibitor at

HH11 and incubated for 20 hours. White arrows indicate hindbrain at level of otic vesicle. (H) Dorsal

views of DMSO and MMP inhibitor treated embryos. Yellow highlight fills the brain dorsal surface. (I)

Confocal images showing the dorsal hindbrain of control (n=3) and MMP inhibitor treated embryos

(n=5), ~20 hours post treatment. Bottom panels show laminin surface with white arrows indicating

breaks in laminin continuity. (J) Hindbrain roof length obtained by measuring the length of the single-

cell thick roof in control and inhibitor treated embryos. White scale bars are 50μm.

Figure 4. Premigratory neural crest cells in the dorsal hindbrain are sufficient to generate a

thinned dorsal roof and neural tube expansion

(A-E) Grafting experiments. (A) Bright-field image of pre-brain expansion stage (HH11) embryos with

a spinal cord-to-spinal cord graft (n=2) or a hindbrain-to-spinal cord graft (n=5). (B) Confocal images

showing dorsal surface view of spinal cord-to-spinal cord and hindbrain-to-spinal cord graft

integration ~20 hours post-grafting. (C) Confocal images showing cross-sectional views of spinal cord-

to-spinal cord and hindbrain-to-spinal cord graft integration ~20 hours post-grafting. Cross section

levels correspond to dashed lines in (B). White bracket highlights the thinned-out roof in the graft

region. D) Spinal cord tissue thickness in the region with spinal cord or hindbrain grafted cells

(thickness measurements taken along lumen circumference going from dorsal midpoint and moving

ventrally). (E) Confocal images showing Snail2+ cells in graft regions. White arrows indicate Snail2+

cells within the GFP+ graft. (F) Schematic depicting proposed model of brain expansion relative to the

spinal cord. Premigratory neural crest cells in pre-brain expansion stage embryos show differences in

their behaviour between the hindbrain and spinal cord regions of the neural tube. A greater extent of

mesenchymal behaviour and corresponding extracellular matrix remodelling underlies a more

deformable dorsal roof in the hindbrain compared to the spinal cord. This allows the hindbrain roof

to deform more under internal lumen pressure, driving hindbrain expansion relative to the spinal

cord during early embryo development.

Black scale bars are 500µm. White scale bars are 25µm unless otherwise stated.

Figure S1. Neural tube characterisation and intubation experiments

(A) An HH12 stage embryo injected in the neural tube lumen with fluorescent Dextran. (B) Brightfield

images of the anterior of embryos used in pressure measurements. (C) Mean thickness of the dorsal

hindbrain in HH11 and HH16 stage embryos. (D) Mean thickness of the dorsal hindbrain in HH11 and

503   HH16 stage embryos. (E) Brightfield images of control and intubated embryos just after intubation

504   and 20 hours later. White arrows mark site of intubation. White asterisks mark the hindbrain.

505   All scale bars are 500μm.

Figure S2. Geometry and tissue mechanics experiments

(A) Confocal images of dorsal and cross-sectional views of the spinal cord region in an HH12 stage

embryo. Dashed lines indicate level of cross-sectional views. (B) A widefield image of an embryo ~20

hours after posterior somite and PSM removal with zoomed-in view of the widened spinal cord

region. (C) Confocal images of cross-sectional views of the spinal cord in embryos 20 hours after

control cuts or somite removal. (D) Dorsal spinal cord thickness in control and somite removed

embryos, measured at the dorsal midpoint. (E) Close up views of ferrofluid droplets in the hindbrain

and spinal cord and a schematic illustrating the higher curvature at the droplet-lumen interface and

thus higher pressure exerted by droplets in the spinal cord region compared to the hindbrain region.

(F) Fitting of AFM indentation data to extract fluidity 'β'. G) Mean elastic modulus in the hindbrain

and spinal cord of HH10, 11 and 12 stage embryos.

White scale bars are 50µm unless otherwise stated.

 Figure S3. Neural crest and ECM experiments

 (A) Confocal images showing Snail2+ cells at the dorsal surface of HH9 and HH11 stage brain and

 spinal cord regions. Dashed lines depict the extent of Snail2+ cell spreading away from the dorsal

 midline. (B) Total area of gaps in ECM on the dorsal surface of the hindbrain and spinal cord of pre-

 brain expansion embryos. (C) Confocal images stitched together to show a 3D rendering of the

 laminin matrix in a HH12 stage embryo. Dashed lines depict the boundaries of the hindbrain region.

 (D) Confocal images of DMSO and MMP inhibitor treated embryo hindbrains, ~20 hours post

 treatment.

 White scale bars are 100µm unless otherwise stated.

 Figure S4. Graft experiments

 (A) Widefield images of an embryo ~20 hours after introduction of a GFP+ graft into the spinal cord

 and a zoomed-in view of the graft region. (B) Confocal images of cross-sectional views of the spinal

 cord at spinal cord-spinal cord and hindbrain-spinal cord graft sites. (C) Confocal images showing

actin organisation at the apical lumen surface in a spinal cord-spinal cord graft region and a

hindbrain-spinal cord graft region. White arrow indicates loss of apical actin localisation.

White scale bars are all 50μm unless otherwise stated.

Figure S5. Theoretical model of neural tube expansion under pressure

A schematic showing the initial undeformed configuration (left) and the current deformed

configuration (right) pressurized with luminal fluids.

References:

1. Sugahara, F. *et al.* Evidence from cyclostomes for complex regionalization of the

ancestral vertebrate brain. *Nature* **531**, 97–100 (2016).

2. Kotrschal, A. *et al.* Artificial Selection on Relative Brain Size in the Guppy Reveals Costs

and Benefits of Evolving a Larger Brain. *Curr Biol* **23**, 168–171 (2013).

3. Sol, D., Duncan, R. P., Blackburn, T. M., Cassey, P. & Lefebvre, L. Big brains, enhanced

cognition, and response of birds to novel environments. *Proceedings of the National*

Academy of Sciences **102**, 5460–5465 (2005).

4. del Corral, R. D. *et al.* Opposing FGF and Retinoid Pathways Control Ventral Neural

Pattern, Neuronal Differentiation, and Segmentation during Body Axis Extension. *Neuron*

40, 65–79 (2003).

5. Nordström, U., Jessell, T. M. & Edlund, T. Progressive induction of caudal neural character

by graded Wnt signaling. *Nat Neurosci* **5**, 525–532 (2002).

6. Bel-Vialar, S., Itasaki, N. & Krumlauf, R. Initiating Hox gene expression: in the early chick

neural tube differential sensitivity to FGF and RA signaling subdivides the HoxBgenes in

two distinct groups. *Development* **129**, 5103–5115 (2002).

7. Colas, J.-F. & Schoenwolf, G. C. Towards a cellular and molecular understanding of

neurulation. *Developmental Dynamics* **221**, 117–145 (2001).

8. Hamburger, V. & Hamilton, H. L. A series of normal stages in the development of the

chick embryo. *Developmental Dynamics* **195**, 231–272 (1992).

9. Tallinen, T. *et al.* On the growth and form of cortical convolutions. *Nature Physics* **12**,

588–593 (2016).

10. Mota, B. & Herculano-Houzel, S. Cortical folding scales universally with surface area and

thickness, not number of neurons. *Science* **349**, 74–77 (2015).

11. Desmond, M. E. & Jacobson, A. G. Embryonic brain enlargement requires cerebrospinal

fluid pressure. *Developmental Biology* **57**, 188–198 (1977).

12. Garcia, K. E., Stewart, W. G., Espinosa, M. G., Gleghorn, J. P. & Taber, L. A. Molecular and

mechanical signals determine morphogenesis of the cerebral hemispheres in the chicken

embryo. *Development* **146**, dev174318 (2019).

13. Le Douarin, N. M. Cell Line Segregation During Peripheral Nervous System Ontogeny.

Science **231**, 1515-1522 (1986)

14. Garcia, K. E., Okamoto, R. J., Bayly, P. V. & Taber, L. A. Contraction and stress-dependent

growth shape the forebrain of the early chicken embryo. *Journal of the Mechanical*

Behavior of Biomedical Materials **65**, 383–397 (2017).

15. Schoenwolf, G. C. Observations on closure of the neuropores in the chick embryo.

American Journal of Anatomy **155**, 445–465 (1979).

16. Desmond, M. E., Levitan, M. L. & Haas, A. R. Internal luminal pressure during early chick

embryonic brain growth: Descriptive and empirical observations. *The Anatomical Record*

Part A: Discoveries in Molecular, Cellular, and Evolutionary Biology **285A**, 737–747

(2005).

17. Alonso, M. I., Gato, A., Moro, J. A. & Barbosa, E. Disruption of proteoglycans in neural

tube fluid by β-D-xyloside alters brain enlargement in chick embryos. *Anat. Rec.* **252**,

499–508 (1998).

18. Xiong, F., Ma, W., Bénazéraf, B., Mahadevan, L. & Pourquié, O. Mechanical Coupling

Coordinates the Co-elongation of Axial and Paraxial Tissues in Avian Embryos.

Developmental Cell **55**, 354-366.e5 (2020)

19. Shellard, A., Szabó, A., Trepat, X. & Mayor, R. Supracellular contraction at the rear of

neural crest cell groups drives collective chemotaxis. *Science* **362**, 339–343 (2018).

20. Berndt, J. D., Clay, M. R., Langenberg, T. & Halloran, M. C. Rho-kinase and myosin II affect

dynamic neural crest cell behaviors during epithelial to mesenchymal transition in vivo.

Developmental Biology **324**, 236–244 (2008).

21. Scarpa, E. *et al.* Cadherin Switch during EMT in Neural Crest Cells Leads to Contact

Inhibition of Locomotion via Repolarization of Forces. *Developmental Cell* **34**, 421–434

(2015).

22. Theveneau, E., Duband, J.-L. & Altabef, M. Ets-1 Confers Cranial Features on Neural Crest

Delamination. *PLoS ONE* **2**, e1142. (2007)

23. Rocha, M. *et al.* From head to tail: regionalization of the neural crest. *Development* **147**,

dev193888 (2020).

24. Taneyhill, L. A., Coles, E. G. & Bronner-Fraser, M. Snail2 directly represses cadherin6B

during epithelial-to-mesenchymal transitions of the neural crest. *Development* **134**,

1481–1490 (2007).

25. Nieto, M. A., Sargent, M. G., Wilkinson, D. G. & Cooke, J. Control of Cell Behavior During

Vertebrate Development by Slug, a Zinc Finger Gene. *Science* **264**, 835–839 (1994).

26. Cai, D. H., Vollberg Sr., T. M., Hahn-Dantona, E., Quigley, J. P. & Brauer, P. R. MMP-2

expression during early avian cardiac and neural crest morphogenesis. *The Anatomical*

Record **259**, 168–179 (2000).

27. Monsonego-Ornan, E. *et al.* Matrix metalloproteinase 9/gelatinase B is required for

neural crest cell migration. *Developmental Biology* **364**, 162–177 (2012).

28. Elosegui-Artola, A. *et al.* Matrix viscoelasticity controls spatiotemporal tissue

organization. *Nat. Mater.* **22**, 117–127 (2023).

29. Crest, J., Diz-Muñoz, A., Chen, D.-Y., Fletcher, D. A. & Bilder, D. Organ sculpting by

patterned extracellular matrix stiffness. *Elife* **6**, e24958 (2017).

30. Gans, C. & Northcutt, R. G. Neural Crest and the Origin of Vertebrates: A New Head.

Science **220**, 268–273 (1983).

31. Couly, G., Grapin-Botton, A., Coltey, P. & Le Douarin, N. M. The regeneration of the

cephalic neural crest, a problem revisited: the regenerating cells originate from the

contralateral or from the anterior and posterior neural fold. *Development* **122**, 3393–

3407 (1996).

32. Campo-Paysaa, F., Clarke, J. D. & Wingate, R. J. Generation of the squamous epithelial

roof of the 4th ventricle. *Elife* **8**, e38485 (2019)

33. Mosaliganti, K. R. *et al.* Size control of the inner ear via hydraulic feedback. *Elife* **8**, 2837

(2019).

34. Chan, C. J. *et al.* Hydraulic control of mammalian embryo size and cell fate. *Nature* **571**,

112–116 (2019).

35. Dumortier, J. G. *et al.* Hydraulic fracturing and active coarsening position the lumen of

the mouse blastocyst. *Science* **365**, 465–468 (2019).

36. Nelson, C. M., Gleghorn, J. P., Pang, M. F., & 2017. Microfluidic chest cavities reveal that

transmural pressure controls the rate of lung development. *Development* **144**, 4328-

4335 (2017)

37. McGrew, M. J. *et al.* Localised axial progenitor cell populations in the avian tail bud are

not committed to a posterior Hox identity. *Development* **135**, 2289–2299 (2008).

38. Rozbicki, E. *et al.* Myosin-II-mediated cell shape changes and cell intercalation contribute

to primitive streak formation. *Nat Cell Biol* **17**, 397–408 (2015).

39. Chapman, S. C., Collignon, J., Schoenwolf, G. C. & Lumsden, A. Improved method for

chick whole-embryo culture using a filter paper carrier. *Dev Dyn* **220**, 284–289 (2001).

40. Kunz, D. *et al.* Downregulation of extraembryonic tension controls body axis formation in

avian embryos. *Nat Commun* **14**, 3266 (2023).

41. McLaren, S. B. P. & Xiong, F. Lumen pressure modulation in chicken embryos. bioRxiv

569177 (2023).

42. Thompson, A. J. *et al.* Rapid changes in tissue mechanics regulate cell behaviour in the

developing embryonic brain. *Elife* **8**, e39356 (2019).

43. Schierbaum, N., Rheinlaender, J. & Schäffer, T. E. Combined atomic force microscopy

(AFM) and traction force microscopy (TFM) reveals a correlation between viscoelastic

material properties and contractile prestress of living cells. *Soft Matter* **15**, 1721–1729

(2019).

44. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat*

Methods **9**, 676–682 (2012).

45. Ahlers, J. *et al.* napari: a multi-dimensional image viewer for Python. (2023)

46. Kyriakides, S. & Yu-Chung, C. The initiation and propagation of a localized instability in an

inflated elastic tube. *International Journal of Solids and Structures* **27**, 1085–1111 (1991).