Beyond microtubules: The cellular environment at the

endoplasmic reticulum attracts proteins to the nucleus, enabling

nuclear transport

Seok Joo Chae[1,2], Dae Wook Kim[3], Oleg A. Igoshin[4,5,6,7], Seunggyu Lee[2,8] and Jae Kyoung

Kim[1,2,9,*]

[1] Department of Mathematical Sciences, KAIST, Daejeon 34141, Republic of Korea

[2] Biomedical Mathematics Group, Pioneer Research Center for Mathematical and

Computational Sciences, Institute for Basic Science, Daejeon 34126, Republic of Korea

[3] Department of Mathematics, University of Michigan, Ann Arbor, MI 48109, USA

[4] Department of Bioengineering, Rice University, Houston, TX 77005, USA

[5] Center for Theoretical Biological Physics, Rice University, Houston, TX 77005, USA

[6] Department of Chemistry, Rice University, Houston, TX 77005, USA

[7] Department of Biosciences, Rice University, Houston, TX 77005, USA

[8] Division of Applied Mathematical Sciences, Korea University, Sejong 30019, Republic of Korea

[9] Lead Contact

*Correspondence: jaekkim@kaist.ac.kr

18
19
Summary

All proteins are translated in the cytoplasm, yet many, including transcription factors,

play vital roles in the nucleus. While previous research has concentrated on molecular motors for

the transport of these proteins to the nucleus, recent observations reveal perinuclear

accumulation even in the absence of an energy source, hinting at alternative mechanisms. Here,

we propose that structural properties of the cellular environment, specifically the endoplasmic

reticulum (ER), can promote molecular transport to the perinucleus without requiring additional

energy expenditure. Specifically, physical interaction between proteins and the ER impedes their

diffusion and leads to their accumulation near the nucleus. This result explains why larger

proteins, more frequently interacting with the ER membrane, tend to accumulate at the

perinucleus. Interestingly, such diffusion in a heterogeneous environment follows Chapman's law

rather than the popular Fick's law. Our findings suggest a novel protein transport mechanism

arising solely from characteristics of the intracellular environment.

Highlights

- The interaction of proteins with ER slows down their diffusion at the perinucleus.

- This leads proteins to migrate toward the perinucleus without ATP consumption.

- Frequent ER interaction of larger proteins promotes perinuclear accumulation.

- Diffusion with the physical interaction can be described by Chapman's law.

Graphical abstract

Introduction

In eukaryotic cells, transcription factors and other proteins that function in the nucleus are translated throughout the cytoplasm but must be translocated across the nuclear membrane[1-3]. The translocation of proteins is regulated by the nuclear pore complexes, which are localized in the nuclear envelope[4,5]. To reach the vicinity of the envelope, nuclear proteins must traverse the cytoplasm. This mainly occurs via intracellular transport mechanisms facilitated by cytoskeletal filament systems, including microtubules and actin, with the assistance of cytoskeletal motors[6-10]. Moreover, actomyosin contraction can also contribute to protein transport by generating cytoplasmic flow[11]. These mechanisms consume adenosine triphosphate (ATP) to fuel motor proteins or contract myosin fibers[6-11].

Recent studies have shed light on an as-yet-unidentified alternative mechanism that operates passively (i.e., without consuming ATP). Specifically, exogenous particles like nanoparticles and particulate matter, accumulate in the perinuclear region upon entering cells[12-14]. This perinuclear accumulation was also still observed when the energy source (ATP) for the motor proteins was depleted (Figure 1(i))[14]. Moreover, recombinant EGFP-dynactin can migrate toward the nucleus and accumulate in the perinuclear region even without its microtubule binding motif (Figure 1(ii))[15]. Altogether, these studies support the existence of the transport mechanism independent of microtubule-based transport. Interestingly, the accumulation depends on temperature, i.e., decreasing as the temperature decreases (Figure 1(iii))[14]. These characteristics suggest that the unknown mechanism leading to perinuclear accumulation is likely based on diffusion. Notably, diffusion is not homogenous across the cell. In particular, the endoplasmic reticulum (ER), a cellular organelle surrounding the nucleus, is known to impede diffusion[16]. This observation raises the question of whether the heterogeneous diffusive environment in the cytoplasm leads to the migration of proteins toward the nucleus.

Our study uncovers that the structural characteristics of the intracellular environment that lead to heterogeneous diffusion of proteins also enable proteins to migrate toward the nucleus. Specifically, our agent-based model (ABM) simulation reveals that such migration occurs due to the ER, which impedes protein diffusion. Notably, our ABM also predicts that larger proteins

interact with the ER more frequently, leading to their increased accumulation at the perinucleus

compared to smaller ones. Indeed, when we analyzed all the proteins whose sizes and

subcellular localization are known simultaneously, we found that perinuclear and nuclear proteins

tend to have larger radii than cytoplasmic proteins. This passive mechanism offers an energy-

efficient alternative for transporting proteins. Furthermore, we found that the diffusion of proteins

in a heterogeneous cellular environment can be effectively described using a partial differential

equation (PDE) based on Chapman's law, rather than the conventional Fick's law that is typically

employed to describe diffusion in the cell. This result suggests that employing Fick's law to

explain diffusion in the presence of physical interactions within a cell might not provide accurate

results. Taken together, these findings shift our understanding of protein transport, demonstrating

that passive mechanisms can be remarkably efficient and that the cell's own architecture plays a

crucial role in guiding protein movement.

Results

The cellular environment at the ER leads to perinuclear accumulation

The diffusion process within the cytoplasm is not uniform due to the presence of cellular

organelles and macromolecules. Among these organelles, the ER surrounds the nucleus with a

maze-like structure (Figure 2A). A previous study suggested that due to the ER, the movement of

particles around the nucleus is hindered, and the diffusion significantly slowed down around the

nucleus[16]. Furthermore, the diffusion coefficient was decreased as the ER density increased[16].

Intriguingly, when the ER was disassembled, the diffusion coefficient near the nucleus was

comparable to that in the peripheral cytoplasm (Figure 2B)[16].

To explore how proteins move in a heterogeneous diffusive cytoplasm due to the ER, we

utilized an ABM, which has previously been used to track molecule movement in various

biological systems[17-20] (see STAR Methods for details). Protein molecules were introduced into

the cytoplasm and were set to move randomly in different directions at each time step, simulating

diffusive movement. Additionally, we incorporated the ER structure surrounding the nucleus,

which impeded protein diffusion. We modeled that this hindrance of protein diffusion stems from

protein binding to the ER membrane upon collision with it (see STAR Methods for details).

Interestingly, ABM simulation revealed that as the protein size increases, more proteins

accumulate in the perinuclear region (Figure 2C). Specifically, as the radius of protein increases,

the portion of protein accumulation at the perinucleus increases (Figure 2D). This is because

larger proteins collide more frequently with the ER membrane than smaller proteins despite lower

diffusion coefficient of larger proteins (Figure S1). Notably, proteins with a radius of ~10nm

exhibit 2.2 times more collisions in the perinuclear region, compared to proteins with a radius of ~

2nm (Figure S1). The higher collision frequency leads to a steeper decline in the diffusion

coefficient of larger proteins at the perinucleus compared to their smaller counterparts (Figure

2E), consistent with a previously observed experiment by Reisch et al.: ER impedes the diffusion

of nanoparticles more as their size increases (Figure 2F)[21]. Our findings indicate that the ER

selectively impedes the diffusion of proteins depending on their sizes, which leads to size-

dependent perinuclear accumulation.

To further validate the size-dependent perinuclear accumulation with experimental data,

we comprehensively analyzed all the proteins with known sizes and intracellular distribution

(Table S1). Specifically, we compared the Stokes radius of proteins[22-34] based on their subcellular

localization, as annotated in previous experiments[35,36]. The proteins localized in the perinucleus

or the nucleus have Stokes radii larger than proteins localized in the cytoplasm (Figure 2G),

consistent with the model prediction.

An effective PDE can recapitulate the perinuclear accumulation of proteins

Even though our ABM explains size-dependent accumulation of proteins in perinuclear

region (Figure 2), it is challenging to mathematically analyze it. To enable mathematical analysis,

we aimed to convert the ABM to a PDE that accounts for the perinuclear accumulation of proteins

without explicitly modeling ER geometry. For this, we utilized a simple microscopic model where

proteins freely diffuse in the cytoplasm but become immobile upon binding to cellular

components, such as the ER (see STAR Methods for details). Under the assumption that the

binding and unbinding is faster than the diffusion (see STAR Methods for details), we can derive

that this effective diffusion of proteins impeded by the binding can be described by Chapman's

law, i.e., the probability of finding a protein $p(x, t)$ at a given position x and time t is governed

by the following equation (Figure 3A; see STAR Methods for details):

$$\frac{\partial p(x,t)}{\partial t} = \Delta\big(D(x)p(x,t)\big),$$

where $D(x)$ is the effective diffusion coefficient at x. Note that Chapman's law is distinct from the

commonly used Fick's law (Figure 3A).

The effective diffusion coefficient, $D(x)$, accounts for how protein diffusion is impeded by

physical interactions. In the absence of such interactions, $D(x)$ remains constant throughout the

cytoplasm, resulting in a homogeneous distribution of $D(x)$ (Figure 3B). Conversely, when

protein interact with the ER near the nucleus, $D(x)$ in this region becomes smaller than that of

the peripheral cytoplasm, leading to a heterogeneous distribution of $D(x)$ across the cytoplasm.

Notably, proteins with larger radii more frequently interact with the ER (Figure S1), causing

increased hindrance of diffusion compared to proteins with smaller radii (Figure 2E). Thus, $D(x)$

for larger proteins near the nucleus decreases more than that of smaller proteins. To explore the

impact of size-dependent $D(x)$ on protein movement and accumulation near the nucleus, we

investigated moderately and highly heterogeneous $D(x)$ (Figure 3B).

In the homogeneous environment, perinuclear accumulation did not occur (Figure 3C).

On the other hand, in heterogeneous environments, proteins gather near the nucleus (Figures 3D

and 3E), resulting in a higher concentration (i.e., higher $p(x, t)$ near the nucleus). Furthermore,

$p(x, t)$ near the nucleus is higher in the highly heterogeneous environment compared to the

moderate heterogeneous environment. Notably, unlike Chapman's law, the PDE based on

conventional Fick's law cannot capture perinuclear accumulation even in a highly heterogeneous

environment (Figure 3F).

To understand why only Chapman's law, not Fick's law, leads to perinuclear

accumulation, we analyzed the flux (i.e., the amount of protein flows) of PDEs in the highly

heterogeneous environment (Figure 3G). In Fick's law, the flux ($D\nabla p$) is directly proportional to

the concentration gradient, causing proteins to migrate from regions of high concentration to

regions of low concentration. This flux leads to a steady state with a uniform concentration

distribution (Figure S2). In contrast, Chapman's law incorporates an additional drift term ($p\nabla D$)

that arises from the gradient of the effective diffusion coefficient in its flux ($D\nabla p + p\nabla D$) (Figure

3G(i))[37]. Since the drift term heads toward the nucleus, proteins begin to gather near the nucleus

from their initial homogeneous concentration distribution (Figure 3G(ii)). As proteins gather near

the nucleus, the diffusive flux ($D\nabla p$) directs proteins outward from the nucleus (Figure 3G(iii)),

opposing the drift term. Despite this opposition, the drift term ($p\nabla D$) remains stronger than the

diffusive flux ($D\nabla p$), causing the total flux ($D\nabla p + p\nabla D$) to still be directed toward the nucleus

(Figure 3G(i)). When the diffusive flux cancels out the drift term, the total flux becomes zero,

resulting in a steady state with protein accumulation near the nucleus (Figure 3G(i)). At the

steady state (i.e., $D\nabla p + p\nabla D = \nabla(Dp) = 0$), $p(x,t)$ is inversely proportional to $D(x)$, highlighting

the crucial role of physical interactions in perinuclear accumulation. Taken together, our findings

suggest that Chapman's law is more suitable than Fick's law for describing the existing

experimental results on diffusion in cellular environments[36].

An ABM, which corresponds to PDE, accurately captures the perinuclear

accumulation

Although the PDE based on Chapman's law provides a simple way to account for the

perinuclear accumulation, it can become cumbersome for complex cell shapes and intracellular

environments. In such cases, using ABM equivalent to PDE based on Chapman's law can be a

versatile alternative. Specifically, incorporating heterogenous effective diffusion in ABM without

explicitly modeling binding and unbinding reactions unlike the ABM in Figure 2 can be an efficient

way to simulate perinuclear accumulation of proteins within a heterogeneous diffusive

environment. Thus, we aimed to explore the conditions under which ABM becomes equivalent to

the PDE based on Chapman's law, enabling the reproduction of perinuclear accumulation of

proteins within a heterogeneous diffusive environment (Figure 3A).

In ABMs, the diffusion of proteins is characterized by a random walk[18,20]. The step size of

each step of random walk is proportional to the square root of the effective diffusion coefficient[38].

Since the effective diffusion coefficient varies within the cytoplasm (Figure 3B), the effective

diffusion coefficient can change even during a single step. Thus, different choices of step size can

be made based on the position where the effective diffusion coefficient is determined. For

example, determination of the effective diffusion coefficient can be based on the current position

before the step or the position after the step. The choice of where to evaluate the diffusion

coefficient results in distinct types of random walk, leading to distinct dynamics. Notably, when

the effective diffusion coefficient is evaluated at the current position, the random walk becomes

equivalent to the discretization of the Ito integral, and the dynamics of ABM follows Chapman's

law[39] (Figure 4A). Therefore, when this random walk scheme was incorporated into the ABM,

perinuclear accumulation was observed over time in a highly heterogeneous diffusion

environment (Figure 4B). Taken together, ABM utilizing the effective diffusion coefficient of

current position of the protein can effectively simulate protein diffusion in heterogeneous

intracellular environments to capture perinuclear accumulation of proteins.

Such ABM framework can depict the diffusion in various intracellular environments.

Specifically, the ABM can be easily adapted to describe more realistic but complex environments,

such as cells with the ER structure partially surrounding the nucleus (Figure 4C), or elliptical-

shaped cells (Figure 4D) (see STAR Methods for details). Remarkably, even in these complex

cell geometries, the ABM simulations replicate perinuclear accumulation (Figures 4C and 4D)

observed in idealized cells (Figure 4B).

204 Discussion

To maintain normal physiological processes within cells, it is crucial to effectively and

selectively transport to the nucleus proteins translated in the cytoplasm. Our research has shown

that proteins can passively diffuse to accumulate near the nucleus without requiring active

transport mechanisms, such as transport by motor proteins along microtubules. This

accumulation occurs because physical interactions between proteins and the ER slow the

diffusion at the perinucleus relative to the peripheral cytoplasm (Figures 2E and 3B). Larger

proteins are more likely to accumulate at the perinucleus (Figures 2C and 2D) because physical

interactions occur more frequently as the size of the molecules increases (Figure S1). Our

findings reveal that diffusion within a heterogeneous cellular environment enables the ATP-free

transport of large-sized proteins, contrasting with the conventional microtubule-based transport,

which requires ATP as an energy source[6,7].

In addition to the ER, the perinucleus is more crowded than the cell periphery because

the perinucleus contains more organelles and denser microtubule networks[40]. This molecular

crowding can also impede protein diffusion if proteins can bind to organelles or microtubules.

Since physical interactions occur more frequently as the size of the molecules increases (Figure

S1), larger proteins are more likely to accumulate at the perinucleus (Figures 2C and 2D). This

provides insight for previous experimental observations: 1) proteins destined to reach the nucleus

often form complexes before translocating to the nucleus[41,42], 2) attaching specific moieties to

anticancer drug particles, thereby increasing their size, facilitates targeting to the ER[43,44], and 3)

different types of particles, such as lipid nanoparticles and CaCO3, tend to accumulate in the

perinuclear region based on their sizes[12-14]. While we focused on proteins with sizes comparable

to the ER width (~20 nm), cells also contain significantly larger particles (>>20nm) like vesicles

and lysosomes[45,46]. These extremely large particles might interact with actin filaments, whose

mesh size is approximately 50nm[40]. Investigating the diffusion of these extremely large particles

and their potential interactions with the cytoskeleton would be an interesting avenue for future

work.

The ER consists of two main structural components: a tubular network structure in the

cell periphery and a sheet structure near the nucleus. While both structures contribute to the

heterogeneity in diffusion coefficients of proteins[16], their effects differ significantly. The tubular

network results in the compartmentalization of the diffusion in the cell periphery, and the diffusion

coefficient changes minimally due to a sparse ER density[16]. In contrast, the sheet structure is

highly dense near the nucleus, and thus large proteins slow down as they approach the nucleus

(Figure 2E)[16]. This suggests that the sheet structure is important for helping proteins move

towards the nucleus, while the tubular network likely has little effect on translocation.

Among various types of PDEs describing diffusion[47-55], it is challenging to determine a

priori the most appropriate diffusion law (e.g., Fick's law or Chapman's law) for modeling diffusion

in heterogeneous environments[56,57]. This challenge arises from the diverse physical causes

giving rise to heterogeneous diffusion, each manifested in distinct forms of diffusion equations[57].

Consequently, a thorough understanding of the underlying physical phenomena governing

heterogeneity is essential. Nevertheless, intracellular diffusion has been primarily described by

Fick's law[58-62]. Our study suggests that describing protein diffusion with binding to cellular

components like the ER (Figure 3B) is equivalent to the "trap model" in Sokolov, resulting in

Chapman's law[57]. When such cellular components impede the diffusion near the nucleus, the

PDE based on Chapman's law but not Fick's Law yields a non-uniform concentration distribution

in a steady state, and accurately describes the perinuclear accumulation (Figures 3C-3F).

However, mathematical models describing circadian clocks have adopted Fick's law to describe

the diffusion of clock proteins[18,59], although a recent study showed that a clock protein is

accumulated at the perinucleus[18]. Thus, utilizing Chapman's law to develop the circadian clock

model would be more appropriate to investigate the spatiotemporal dynamics of molecular

circadian clock[18,20,59,63-65]. Furthermore, it would be an interesting future work to investigate

dynamics of various biological oscillators based on transcriptional feedback loops requiring

nuclear transport of transcriptional factors (e.g., p53 oscillator[66-68], NF-κb oscillator[69], and

synthetic oscillator[70-73]) when Chapman's law is used rather than Fick's law.

Previous studies of diffusion in heterogeneous environments, particularly focusing on

highly crowded cellular environments, have highlighted the prevalence of subdiffusion[74-76] for

various particles, including small labeled proteins, lipid and insulin granules, and virus particle[77-

80]. This subdiffusive behavior, characterized by slower-than-linear growth in displacement

variance, suggests hindered transport and potential inefficiencies in cellular processes. On the

other hand, our study demonstrates how heterogeneous diffusion, specifically under the influence

of physical interactions with the ER, can facilitate directional protein transport toward the nucleus,

as implied by Chapman's law. Consistently, heterogeneous diffusion was known to be beneficial

for reducing the first-passage time of reaction events, even though it slows down the overall

dynamics of proteins[81]. Taken together, diffusion in heterogeneous environments can also

contribute to efficient cellular functioning.

Limitations of the study

We have comprehensively examined all the proteins for which both their sizes and

subcellular localizations (i.e., locations of proteins) are known, and this analysis supports that

larger proteins tend to accumulate near the nucleus (Figure 2G). However, the scarcity of

proteins with known sizes and subcellular localizations restricts the generalizability of our

conclusions. It would be an important topic to explore how subcellular localization and proteins

sizes are correlated across a broader range of proteins. Moreover, it is crucial to verify whether a

protein does not contain NLS when examining its subcellular localization, since proteins and

particles with NLS are reported to induce microtubule-based transport[82,83]. To further strengthen

our conclusion, it would also be helpful to investigate whether smaller proteins can accumulate

near the nucleus after fusing with fluorescent proteins, as these modifications can significantly

increase the size of protein[84].

Acknowledgements

We thank Yong Jung Kim and Ho-Youn Kim for the helpful discussion. We thank Life Science

Editors for editorial assistance. This work was supported by the Institute for Basic Science (grant

no. IBS-R029-C3) (J.K.K.) and Welch Foundation grant no. C-1995 (O.A.I.). Some parts of figure

2 were retrieved from Biorender.

Author Contributions

S.J.C., D.W.K., S.L. and J.K.K. designed the study. S.J.C. performed computational modeling

and simulation and all authors contributed to analysis. J.K.K. supervised the project. S.J.C. and

J.K.K. wrote the manuscript draft, and all authors revised the manuscript.

Declaration of interests

The authors declare no competing interests.

Main figures and legends

298
Figure 1. Perinuclear accumulation of protein occurs without consuming ATP. Proteins that
need to be transported to the nucleus typically rely on motor proteins powered by ATP for their
transport. However, surprisingly, exogenous particle can accumulate in the perinuclear region
even when the ATP is depleted (i)[14], suggesting an alternative transport mechanism. Furthermore,
disruption of microtubule binding does not impede this accumulation. Specifically, removing the
binding motif of dynactin, a protein assisting dynein in transportation, does not hinder its
movement toward the nucleus (ii)[15]. Notably, such accumulation was weakened when the
temperature was lowered (iii)[14]. This indicates that the accumulation may have a diffusive
character. Reprinted by permission from Rockefeller University Press: Journal of Cell Biology
2007 (doi: 10.1083/jcb.200608128)[15].
309

310
Figure 2. The endoplasmic reticulum (ER) hampers the diffusion of large-sized protein
near the nucleus, resulting in migration toward the nucleus. (**A**) The complex structure of ER
can impede the diffusion of protein molecules near the nucleus. (**B**) The diffusion coefficient of a
quantum dot near the nucleus was much lower compared to the peripheral cytoplasm. When the
ER was degraded, the diffusion coefficients became similar throughout the entire cell. The figure
is obtained by analyzing Figures 3 and 5 in a previous study[16]. (**C** and **D**) To mimic diffusion in the
presence of the ER, an agent-based model (ABM) describing intracellular diffusion was utilized.
In particular, we located obstacles with narrow intervals (dark grey region) near the nucleus to
describe ER (C). When a protein collided with an obstacle, the protein could bind to the obstacle
and become immobile, thereby restricting its movement. This restriction in movement affected
larger proteins more than smaller ones since larger proteins more frequently collide with the ER
(Figure S1) and thus larger proteins more likely to be accumulate near the nucleus (D). In (C),
proteins and obstacles are magnified 20 times for better visualization while biologically realistic
sizes are used in the simulation for (D) (see STAR Methods for details). (**E**) The perinuclear
diffusion coefficient relative to cytoplasm decreases with increasing protein radius. Protein
diffusion coefficients were estimated from their mean-squared displacement (   ) using the
Einstein relation (   ). (**F**) Consistent with the ABM simulation, diffusion of larger
nanoparticles is more likely to be disrupted by the ER compared to that of smaller particles[21]. (**G**)
Aligning with the ABM simulation, proteins which are localized in the perinucleus and the nucleus
(blue) have larger sizes, while proteins that are localized in the cytoplasm (red) have smaller

sizes. The protein size was determined using the Stokes radii (i.e., hydrodynamic radii) from
previous studies[27,30-33].

333
Figure 3. A partial differential equation (PDE) based on Chapman's law captures the
movement of proteins toward the nucleus by heterogeneous diffusion. (**A**) The hindered
diffusion due to the physical interaction with the ER near the nucleus can be described by the
PDE based on Chapman's law. This is different from the PDE based on Fick's law, which has
been widely utilized to describe diffusion. Here,   is the diffusion coefficient at the position
and   is the probability of finding the protein at the position   and the time . (**B**) Three
different environments of diffusion coefficient. (**C-E**) While the PDE based on Chapman's law
resulted in uniform concentration distribution in the homogeneous environment (C), proteins
gathered near the nucleus in heterogeneous environments (D and E). (**F**) Fick's law, which has
been widely used to describe diffusion[60-62], did not exhibit perinuclear accumulation even in the
highly heterogeneous environment. (**G**) The flux of Chapman's law (i) has two components: a drift
term (   ) resulting from the gradient of the diffusion coefficient (ii) and a diffusive flux (   )
resulting from the concentration gradient (iii). In the heterogeneous intracellular environment, the
drift term (ii) directs proteins toward the nucleus. This leads the diffusive flux (iii) to push proteins
outward from the nucleus. While the drift term (ii) remains stronger than the diffusive flux (iii), the
protein continues to accumulate near the nucleus (t=0, 50). Eventually, when the drift term (ii)
and the diffusive flux (iii) become similar in strength, the net flux reaches the equilibrium, meaning
there is no net movement of proteins. (t=100).

352
353

354
Figure 4. ABM scheme that corresponds to Chapman's law results in the perinuclear
accumulation of proteins. (**A**) For the ABM simulation, the step size of the protein movement is
determined by the diffusion coefficient. When the diffusion coefficient varies over the cytoplasm,
the step size can vary depending on the position at which the diffusion coefficient is obtained.
When the step size is determined by the molecule's current position, the dynamics of proteins is
equivalent to Chapman's law. (**B**) This ABM scheme was simulated in the highly heterogeneous
environment, which is used in Figure 3E. Initially (t=0), proteins were uniformly dispersed
throughout the cytoplasm. Over time, the proteins gradually accumulated near the nucleus (t=50,
100). (**C**) To simulate the ER partially surrounding the nucleus, we selectively reduced the
diffusion coefficient in specific perinuclear regions (blue). Even when the ER is partially
surrounding the nucleus, perinuclear accumulation still occurred. (**D**) Besides, ABM provided a
convenient approach for describing diffusion in cells of various shapes, including elliptical cells.
Proteins again accumulated near the nucleus in elliptical cells.
368

STAR★Methods

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the lead contact, Jae Kyoung Kim (jaekkim@kaist.ac.kr).

Materials availability

This study did not generate new unique reagents.

Data and code availability

-    This paper analyzes publicly available existing data. These accession numbers for the

378   datasets are listed in the key resources table.

379   -   The MATLAB and Netlogo codes of the computational package are available in the

380   following Database: The link will be available upon acceptance of the manuscript.

381   -   Any additional information required to analyze the data is available from the lead contact

382   upon request.

METHOD DETAILS

ABM describing diffusion in the presence of the ER

To describe the heterogeneous diffusion in a cell, we utilized an ABM which describes

the intracellular diffusion of proteins. Following previous studies[18,20], we simplified the cell

geometry as a two-dimensional circular shape with the radius of 10μm[85]. Within the cell, a circular

nucleus resided, sharing the same center and possessing a radius that is one-third of the cell

radius (i.e., 3.33μm), so that the nucleus occupies about 10% of the cell area[18]. Furthermore, we

incorporated the ER by introducing obstacles near the nucleus (Figure 2C). These obstacles were

distributed across 80 concentric layers. Starting from 3.58μm, each layer increased in radius by

0.25μm, reflecting the width of the ER sheet structure[85]. Within each layer, 1,000 uniformly

distributed obstacles were uniformly distributed to mimic increased ER density as proteins

approached the nucleus. Each obstacle was modeled as a circle with a 0.0025μm radius.

A protein molecule was represented as a circle with the radius r. In our simulation, we

varied r between $2\,\mathrm{nm}$, corresponding to the radius of carbonic anhydrase[33], to $10\,\mathrm{nm}$,

corresponding to the PER2 complex[86]. The diffusion of proteins was described by using a random

walk. During each time step of length Δt, each molecule was moved in a randomly chosen

direction with a step size δ. The angles of movement were randomly sampled from the uniform

distribution, ranging from 0° to 360°. The step size δ was chosen to be sufficiently smaller

than the protein size to ensure precise and fine-scale movements. For example, for a protein with

10nm radius, δ was set to be $1.6\mathrm{nm}$. For proteins with smaller radii, the step size was adjusted

by using the Stokes-Einstein equation (i.e., $D \propto \frac{1}{r}$, and thus $\delta \propto \frac{1}{\sqrt{r}}$). The time step length (Δt)

was determined by using the Einstein relation $\delta^2 = 4D\Delta t$[38], where D is the diffusion coefficient.

In this study, we utilized the known diffusion coefficient of PER2 protein (D=$0.2\mu\mathrm{m}^2/\mathrm{s}$)[87] and its

radius of 10nm[86], (i.e., $\delta = 1.6\mathrm{nm}$), resulting in $\Delta t = 0.0000032\mathrm{s}$.

When proteins collided to an obstacle, they were repositioned to the point of collision.

Subsequently, they bound to the obstacle with a binding probability of $1 - \exp(-k_{bind}\Delta t)$, where

k_{bind} is the binding rate. Once bound, at each time step, proteins could unbind with a probability

of $1 - \exp(-k_{unbind}\Delta t)$, where k_{unbind} is the unbinding rate. In our simulation, k_{bind} and k_{unbind}

were set to $1{,}562{,}500/s$ and $625/s$, respectively.

We implemented reflecting boundary conditions to prevent the protein from leaving the

cytoplasm. That is, we repositioned proteins as if they were elastically reflected by the boundary

(cell membrane or the nuclear membrane). Specifically, proteins exceeding the cytoplasmic

boundary (cytoplasmic membrane or nuclear membrane) are mirrored across the tangent line of

the boundary at the collision point with the boundary. Additionally, for the initial condition, protein

began diffusion from a uniform distribution of the protein throughout the cytoplasm.

Physical interpretation of Chapman's law

Since our mechanism of protein transport operates without the need for energy sources,

such as ATP hydrolysis, we expect that steady state distribution will correspond to the

thermodynamic equilibrium. This steady state with non-uniform concentration distribution should

result from the energy difference between spatial locations. We postulate that the difference in

energy level comes from the physical interaction between proteins and the cellular organelles,

including the ER.

To investigate this, we considered a one-dimensional diffusion of protein in the

cytoplasm. Specifically, we considered a protein moving along a linear chain of N sites ($i =$

$1 \sim N$), where the chain represents the cytoplasm and each site is separated by an interval h. The

protein located at the site i can interact with cellular organelles, such as the ER, with a binding

rate of $k_f^{(i)}$ and it can also detach from the organelles with an unbinding rate of $k_d^{(i)}$. If the protein

at the site i is not bound, it can diffuse to the adjacent site (e.g., $i + 1$ or $i - 1$) with a diffusion

coefficient of D. Denoting the probability of finding the bounded and unbounded protein at the site

i at time t by $p_i^b(t)$ and $p_i^u(t)$, respectively, we obtain the following ordinary differential equation

(ODE):

$$\frac{dp_i^u}{dt} = \frac{D}{h^2}(p_{i-1}^u - 2p_i^u + p_{i+1}^u) + k_d^{(i)}p_i^b - k_f^{(i)}p_i^u,$$

Eq. (1)

$$\frac{dp_i^b}{dt} = -k_d^{(i)}p_i^b + k_f^{(i)}p_i^u.$$

436   Here, we use the convention of $p_0^u = p_1^u$ and $p_{N+1}^u = p_N^u$ by imposing Neumann boundary

437   conditions. Since we are primarily interested in the probability of finding proteins at the i-th site,

438   regardless of whether they are bound or not, we introduce a new variable p_i^T, the probability of

439   finding the protein at the site i, i.e., $p_i^T = p_i^u + p_i^b$. Then, Eq. (1) can be written as shown below:

$$\frac{dp_i^T}{dt} = \frac{D}{h^2}\left[(p_{i-1}^T - p_{i-1}^b) - 2(p_i^T - p_i^b) + (p_{i+1}^T - p_{i+1}^b)\right],$$

$$\frac{dp_i^b}{dt} = -k_d^{(i)}p_i^b + k_f^{(i)}(p_i^T - p_i^b).$$

Eq. (2)

We assume that binding and unbinding reactions occur significantly faster than diffusion. Then

p_i^T is affected by only slow diffusion, whereas p_i^b is affected by fast binding and unbinding

reactions, leading to a complete time scale separation. This allows us to assume that p_i^b reaches

the quasi-equilibrium (the fast binding and unbinding equilibrium) at each site before p_i^T changes

appreciably (i.e., $-k_d^{(i)}p_i^b + k_f^{(i)}(p_i^T - p_i^b) = 0$ for each i)[88-92]. Then, p_i^b can be expressed by p_i^T

($p_i^b = \dfrac{p_i^T}{\left(1+\frac{k_f^{(i)}}{k_b^{(i)}}\right)}$). Consequently, p_i^T follows a reduced ODE shown below:

$$\frac{dp_i^T}{dt} = \frac{D}{h^2}\left(\frac{p_{i-1}^T}{1+K_{eq}^{(i-1)}} - \frac{2p_i^T}{1+K_{eq}^{(i)}} + \frac{p_{i+1}^T}{1+K_{eq}^{(i+1)}}\right),$$ Eq. (3)

446 where $K_{eq}^{(i)} = \dfrac{k_f^{(i)}}{k_d^{(i)}}$. Taking the limit of infinitesimal interval ($h \to 0$), we obtain the following

447 Chapman's law:

$$\frac{\partial p(x)}{\partial t} = \Delta\big(D_{eff}(x)p(x)\big),$$ Eq. (4)

where $D_{eff}(x) = \dfrac{D}{1+K_{eq}(x)}$. To validate the model reduction, we compared whether the original

model (Eq. (1)) and the reduced model exhibit similar behavior. As long as the binding and

unbinding occur significantly faster than the diffusion, the two models exhibit similar dynamics

(Figure S3).

Numerical simulation for the PDE

After deriving the PDE based on Chapman's law in the previous section, we solved the

PDE numerically. Depicting diffusion within a cell accurately using the standard finite difference

method on a rectangular domain presents challenges because cells usually have a non-

rectangular shape and contain a nucleus where diffusion does not occur. To address this issue,

we adopted an approach from a previous study which described the diffusion under a domain

with a complex geometry[93]. Specifically, we solved the PDE based on Chapman's law ($\frac{\partial p}{\partial t} =$

$\Delta(Dp)$) in a rectangular domain $\Omega = [-L, L] \times [-L, L]$, where, L was set to be $1.1 \cdot R_C$, where R_C

is the radius of the cell. By letting $h = \dfrac{2L}{N+1}$, we obtained a uniform mesh with a set of grid edges

as $\Omega^h = \{(x_i, y_j): 1 \le i, j \le N\}$, where $x_i = -L + (i-1)h$ and $y_j = -L + (j-1)h$. Here, we used

$N = 100$. At the edge (x_i, y_j), the boundary control function $G_{i,j}$ was defined as

$G = \begin{cases} 1\,(R_N \le \|(x_i, y_j)\| \le R_C) \\ 0\,(otherwise) \end{cases}$, where R_N is the radius of the nucleus. In the grid edge, G is

defined as $G_{i+\frac{1}{2},j} = G_{i,j}G_{i+1,j}$ and $G_{i,j+\frac{1}{2}} = G_{i,j}G_{i,j+1}$. Utilizing G allowed us to utilize the finite

difference method in a complex domain mimicking the cell by simply computing PDE in a

rectangular domain[93].

We denote the discrete approximation $p(x_i, y_j, n\Delta t)$ as $p_{i,j}^n$. Then, we can define the

discrete differential operator ∇_d as follows:

$$\nabla_{d,x} p_{i+\frac{1}{2},j}^n = \frac{p_{i+1,j} - p_{i,j}}{h},$$

Eq. (5)

$$\nabla_{d,y} p_{i,j+\frac{1}{2}}^n = \frac{p_{i,j+1} - p_{i,j}}{h}.$$

Furthermore, to calculate ∇_d at the boundary, the value of $p_{i,j}^n$ at ghost points located outside of

the domain also required ($i, j = 0 \ or \ N + 1$). To determine values out of the domain, we utilized

the Neumann boundary condition. Specifically, we set $\nabla_{d,x}\left(p_{i+\frac{1}{2},j}^n\right) = 0$ for $i = 0, N + 1$ and

$\nabla_{d,y}\left(p_{i,j+\frac{1}{2}}^n\right) = 0$ for $j = 0, N + 1$. The boundary condition and boundary control function G

ensures the conservation of $p(x,t)$ over time, i.e., $\int p(x,t)dx$ remains constant over time[94].

Finally, the PDE based on Chapman's law can be written as the following:

$$\frac{p_{i,j}^{n+1} - p_{i,j}^n}{\Delta t} = \nabla_d \cdot (G\nabla_d Dp)_{i,j}$$

$$= \frac{1}{h}\left(G_{i+\frac{1}{2},j}\nabla_{d,x}\left(D_{i+\frac{1}{2},j}p_{i+\frac{1}{2},j}^n\right) - G_{i-\frac{1}{2},j}\nabla_{d,x}\left(D_{i-\frac{1}{2},j}p_{i-\frac{1}{2},j}^n\right) + G_{i,j+\frac{1}{2}}\nabla_{d,y}\left(D_{i,j+\frac{1}{2}}p_{i,j+\frac{1}{2}}^n\right)\right.$$

Eq. (6)

$$\left. - G_{i,j-\frac{1}{2}}\nabla_{d,y}\left(D_{i,j-\frac{1}{2}}p_{i,j-\frac{1}{2}}^n\right)\right),$$

for $1 \leq i,j \leq N$, where h=0.2μm and $\Delta t = 0.00004$s were used. To determine G, the same value

of R_N and R_C as in the ABM are used. Using the PDE (Eq. (6)), we obtained the solution of

Chapman's law in Figure 3.

To describe the diffusion with the existence of physical interactions, we have utilized

various effective diffusion coefficients at (x, y), $D(x, y)$. A homogeneous case described by using

$D(x) = 0.2$μm^2/s. For moderately and highly heterogeneous cases, we employed functions that

decrease as the protein approaches the nucleus. Specifically, we used $D(x, y) = 0.2 \cdot \left(0.27 +\right.$

$0.72 \cdot \left(\frac{(\|(x,y)\|-r_N)^2}{(\|(x,y)\|-r_N)^2+r_N^2}\right)^2 \mu\text{m}^2/\text{s}$ and $D(x,y) = 0.2 \cdot \left(0.09 + 0.91\frac{(\|(x,y)\|-r_N)^2}{(\|(x,y)\|-r_N)^2+r_N^2}\right)^2 \mu\text{m}^2/\text{s}$ to describe

moderately heterogeneous and highly heterogeneous cases, respectively. Here, r_N is the radius

of the nucleus, and $\|(x,y)\|$ is the distance between the protein and the cell center.

Simulating ABM based on Chapman's law

To simulate an ABM equivalent to Chapman's law, we introduced a variation in the

protein's movement (Figures 4B-4D). Specifically, the protein molecule at position (x,y) moves

by a step size $\delta(x,y)$ in a random direction. The step size of the protein was determined by the

diffusion coefficient of the current position. We modeled the decrease in the step size by using a

Hill function: $\delta(x,y) = 0.2 \cdot \left(0.1 + \frac{(\|(x,y)\|-r_N)^2}{(\|(x,y)\|-r_N)^2+r_N^2}\right)\mu\text{m}$, where r_N is the radius of the nucleus, and

$\|(x,y)\|$ is the distance between the protein and the cell center. This choice of step size of the

same form of Hill function to PDE ($\delta^2 \propto D$) ensures that the ABM aligns with the highly

heterogeneous environment. As the protein approaches the nucleus, the step size decreases to

be 10% of the maximum value (=0.02μm). The length of time of step Δt was obtained by using

the relation $\delta_M^2 = 4D\Delta t$, where δ_M is the maximum value of the step size and D is the diffusion

coefficient of PER2 protein (D=0.2μm^2/s)[87].

In Figures 4C and 4D, we have extended the ABM to describe cells with diverse

geometries. To model the ER structure partially surrounding the nucleus in Figure 4C, we

implemented the following step size function:

$\delta(x,y) = 0.2 \cdot \left(0.1 + \frac{(\|(x,y)\|-r_N)^2}{(\|(x,y)\|-r_N)^2+r_N^2}\right) \cdot \mathbf{1}_O\left[\arctan\left(\frac{y}{x}\right)\right] \mu\text{m}$,   where

$O = \{\theta: 60° \leq \theta < 210° \text{ or } 270° \leq \theta < 360°\}$ is and $\mathbf{1}_O(x)$ is the indicator function of O. In

Figure 4D, a cell was represented by an ellipse with a semi-major axis of 15μm and a semi-minor

axis of 10μm, centered at the origin (Figure 4D). The nucleus was depicted as a circle with a

radius of $3.33\mu\text{m},$ centered at (-5μm, 0μm).

References

1.   Tsai, Y.J., Lee, H.I., and Lin, A. (2012). Ribosome distribution in HeLa cells during the
510   cell cycle. PLoS One *7*, e32820. 10.1371/journal.pone.0032820.
511   2.   Kim, Y.H., Han, M.E., and Oh, S.O. (2017). The molecular mechanism for nuclear
512   transport and its application. Anat Cell Biol *50*, 77-85. 10.5115/acb.2017.50.2.77.
513   3.   Lu, J., Wu, T., Zhang, B., Liu, S., Song, W., Qiao, J., and Ruan, H. (2021). Types of
514   nuclear localization signals and mechanisms of protein import into the nucleus. Cell
515   Communication and Signaling *19*, 60. 10.1186/s12964-021-00741-y.
516   4.   Beck, M., and Hurt, E. (2017). The nuclear pore complex: understanding its function
517   through structural insight. Nat Rev Mol Cell Biol *18*, 73-89. 10.1038/nrm.2016.147.
518   5.   Lin, D.H., and Hoelz, A. (2019). The Structure of the Nuclear Pore Complex (An Update).
519   Annu Rev Biochem *88*, 725-783. 10.1146/annurev-biochem-062917-011901.
520   6.   Reck-Peterson, S.L., Redwine, W.B., Vale, R.D., and Carter, A.P. (2018). The
521   cytoplasmic dynein transport machinery and its many cargoes. Nat Rev Mol Cell Biol *19*,
522   382-398. 10.1038/s41580-018-0004-3.
523   7.   Sweeney, H.L., and Holzbaur, E.L.F. (2018). Motor Proteins. Cold Spring Harb Perspect
524   Biol *10*. 10.1101/cshperspect.a021931.
525   8.   Olenick, M.A., and Holzbaur, E.L.F. (2019). Dynein activators and adaptors at a glance.
526   Journal of Cell Science *132*. 10.1242/jcs.227132.
527   9.   Xiang, X., and Qiu, R. (2020). Cargo-Mediated Activation of Cytoplasmic Dynein in vivo.
528   Front Cell Dev Biol *8*, 598952. 10.3389/fcell.2020.598952.
529   10.   Egan, Martin J., McClintock, Mark A., Hollyer, Ian H.L., Elliott, Hunter L., and Reck-
530   Peterson, Samara L. (2015). Cytoplasmic Dynein Is Required for the Spatial Organization
531   of Protein Aggregates in Filamentous Fungi. Cell Reports *11*, 201-209.
532   https://doi.org/10.1016/j.celrep.2015.03.028.
533   11.   Murrell, M., Oakes, P.W., Lenz, M., and Gardel, M.L. (2015). Forcing cells into shape: the
534   mechanics of actomyosin contractility. Nature Reviews Molecular Cell Biology *16*, 486-
535   498. 10.1038/nrm4012.
536   12.   Chakraborty, S.K., Fitzpatrick, J.A.J., Phillippi, J.A., Andreko, S., Waggoner, A.S.,
537   Bruchez, M.P., and Ballou, B. (2007). Cholera Toxin B Conjugated Quantum Dots for
538   Live Cell Labeling. Nano Letters *7*, 2618-2626. 10.1021/nl0709930.
539   13.   Wei, W., Ma, G.-H., Hu, G., Yu, D., McLeish, T., Su, Z.-G., and Shen, Z.-Y. (2008).
540   Preparation of Hierarchical Hollow CaCO3 Particles and the Application as Anticancer
541   Drug Carrier. Journal of the American Chemical Society *130*, 15808-15810.
542   10.1021/ja8039585.
543   14.   Rivolta, I., Panariti, A., Collini, M., Lettiero, B., D'Alfonso, L., Sironi, L., Miserocchi, G.,
544   and Chirico, G. (2011). A biophysical model of intracellular distribution and perinuclear
545   accumulation of particulate matter. Biophysical Chemistry *158*, 134-140.
546   https://doi.org/10.1016/j.bpc.2011.06.009.
547   15.   Kim , H., Ling , S.-C., Rogers , G.C., Kural , C., Selvin , P.R., Rogers , S.L., and Gelfand ,
548   V.I. (2007). Microtubule binding by dynactin is required for microtubule organization but
549   not cargo transport. Journal of Cell Biology *176*, 641-651. 10.1083/jcb.200608128.
550   16.   Li, H., Dou, S.-X., Liu, Y.-R., Li, W., Xie, P., Wang, W.-C., and Wang, P.-Y. (2015).
551   Mapping Intracellular Diffusion Distribution Using Single Quantum Dot Tracking:
552   Compartmentalized Diffusion Defined by Endoplasmic Reticulum. Journal of the
553   American Chemical Society *137*, 436-444. 10.1021/ja511273c.
554   17.   Longo, D., Peirce, S.M., Skalak, T.C., Davidson, L., Marsden, M., Dzamba, B., and
555   DeSimone, D.W. (2004). Multicellular computer simulation of morphogenesis: blastocoel
556   roof thinning and matrix assembly in Xenopus laevis. Dev Biol *271*, 210-222.
557   10.1016/j.ydbio.2004.03.021.
558   18.   Beesley, S., Kim, D.W., D'Alessandro, M., Jin, Y., Lee, K., Joo, H., Young, Y., Tomko,
559   R.J., Faulkner, J., Gamsby, J., et al. (2020). Wake-sleep cycles are severely disrupted by
560   diseases affecting cytoplasmic homeostasis. Proceedings of the National Academy of
561   Sciences *117*, 28402-28411. 10.1073/pnas.2003524117.

19. Shinde, S.B., and Kurhekar, M.P. (2020). Agent-Based Modeling of the Adaptive Immune
System Using Netlogo Simulation Tool.   held in Singapore, 2020//. K.N. Das, J.C.
Bansal, K. Deep, A.K. Nagar, P. Pathipooranam, and R.C. Naidu, eds. (Springer
Singapore), pp. 463-474.
20. Chae, S.J., Kim, D.W., Lee, S., and Kim, J.K. (2023). Spatially coordinated collective
phosphorylation filters spatiotemporal noises for precise circadian timekeeping. iScience
26, 106554. https://doi.org/10.1016/j.isci.2023.106554.
21. Reisch, A., Heimburger, D., Ernst, P., Runser, A., Didier, P., Dujardin, D., and
Klymchenko, A.S. (2018). Protein-Sized Dye-Loaded Polymer Nanoparticles for Free
Particle Diffusion in Cytosol. Advanced Functional Materials *28*, 1805157.
https://doi.org/10.1002/adfm.201805157.
22. Kushner, I., and Somerville, J.A. (1970). Estimation of the molecular size of C-reactive
protein and Cx-reactive protein in serum. Biochimica et Biophysica Acta (BBA) - Protein
Structure *207*, 105-114. https://doi.org/10.1016/0005-2795(70)90140-6.
23. Morgenstern, R., Guthenberg, C., and Depierre, J.W. (1982). Microsomal Glutathione S-
Transferase. European Journal of Biochemistry *128*, 243-248.
https://doi.org/10.1111/j.1432-1033.1982.tb06958.x.
24. Randahl, H., Slabý, I., and Holmgren, A. (1982). An Improved Purification Method and a
Physical Characterization of Phage T7 DNA Polymerase. European Journal of
Biochemistry *128*, 445-449. https://doi.org/10.1111/j.1432-1033.1982.tb06984.x.
25. Triebel, H., Bär, H., Walter, A., Osipova, T.N., Ramm, E.I., Kostyleva, E.I., and Vorob'ev,
V.I. (1989). Structural differences between histone H1 molecules from sea urchin
(Strongylocentrotus intermedius) sperm and calf thymus: hydrodynamic and c.d. studies.
International Journal of Biological Macromolecules *11*, 153-158.
https://doi.org/10.1016/0141-8130(89)90060-3.
26. Tomkinson, A.E., Lasko, D.D., Daly, G., and Lindahl, T. (1990). Mammalian DNA ligases.
Catalytic domain and size of DNA ligase I. Journal of Biological Chemistry *265*, 12611-
12617. https://doi.org/10.1016/S0021-9258(19)38387-5.
27. Forsberg, E.C., Lam, L.T., Yang, X.J., Nakatani, Y., and Bresnick, E.H. (1997). Human
histone acetyltransferase GCN5 exists in a stable macromolecular complex lacking the
adapter ADA2. Biochemistry *36*, 15918-15924. 10.1021/bi971664x.
28. Raab-Graham, K.F., and Vandenberg, C.A. (1998). Tetrameric Subunit Structure of the
Native Brain Inwardly Rectifying Potassium Channel Kir 2.2*. Journal of Biological
Chemistry *273*, 19699-19707. https://doi.org/10.1074/jbc.273.31.19699.
29. Adamczyk, Z., Cichocki, B., Ekiel-Jeżewska, M.L., Słowicka, A., Wajnryb, E., and
Wasilewska, M. (2012). Fibrinogen conformations and charge in electrolyte solutions
derived from DLS and dynamic viscosity measurements. Journal of Colloid and Interface
Science *385*, 244-257. https://doi.org/10.1016/j.jcis.2012.07.010.
30. Rajani, K.R., Pettit Kneller, E.L., McKenzie, M.O., Horita, D.A., Chou, J.W., and Lyles,
D.S. (2012). Complexes of vesicular stomatitis virus matrix protein with host Rae1 and
Nup98 involved in inhibition of host transcription. PLoS Pathog *8*, e1002929.
10.1371/journal.ppat.1002929.
31. Moor, N.A., Vasil'eva, I.A., Anarbaev, R.O., Antson, A.A., and Lavrik, O.I. (2015).
Quantitative characterization of protein-protein complexes involved in base excision DNA
repair. Nucleic Acids Res *43*, 6009-6022. 10.1093/nar/gkv569.
32. Ritterhoff, T., Das, H., Hofhaus, G., Schröder, R.R., Flotho, A., and Melchior, F. (2016).
The RanBP2/RanGAP1*SUMO1/Ubc9 SUMO E3 ligase is a disassembly machine for
Crm1-dependent nuclear export complexes. Nature Communications *7*, 11482.
10.1038/ncomms11482.
33. La Verde, V., Dominici, P., and Astegno, A. (2017). Determination of Hydrodynamic
Radius of Proteins by Size Exclusion Chromatography. Bio Protoc *7*, e2230.
10.21769/BioProtoc.2230.
34. Kilgore, H.R., Latham, A.P., Ressler, V.T., Zhang, B., and Raines, R.T. (2020). Structure
and Dynamics of N-Glycosylated Human Ribonuclease 1. Biochemistry *59*, 3148-3156.
10.1021/acs.biochem.0c00191.

35.   Rastogi, S., and Rost, B. (2011). LocDB: experimental annotations of localization for
618   Homo sapiens and Arabidopsis thaliana. Nucleic Acids Res *39*, D230-234.
619   10.1093/nar/gkq927.
620   36.   Shaiken, T.E., and Opekun, A.R. (2014). Dissecting the cell to nucleus, perinucleus and
621   cytosol. Scientific Reports *4*, 4923. 10.1038/srep04923.
622   37.   Mehrer, H. (2007). Diffusion in solids: fundamentals, methods, materials, diffusion-
623   controlled processes (Springer Science & Business Media).
624   38.   Berg, H.C. (1993). Random Walks in Biology (Princeton University Press).
625   39.   Bressloff, P.C. (2014). Stochastic Processes in Cell Biology. Interdisciplinary Applied
626   Mathematics,. 1st ed. Springer International Publishing : Imprint: Springer,.
627   40.   Jiang, C., Yang, M., Li, W., Dou, S.X., Wang, P.Y., and Li, H. (2022). Spatiotemporal
628   three-dimensional transport dynamics of endocytic cargos and their physical regulations
629   in cells. iScience *25*, 104210. 10.1016/j.isci.2022.104210.
630   41.   Takahashi, J.S. (2017). Transcriptional architecture of the mammalian circadian clock.
631   Nat Rev Genet *18*, 164-179. 10.1038/nrg.2016.150.
632   42.   Wintrebert, M., Nguyen, M.C., and Smith, G.R. (2021). Activation of meiotic
633   recombination by nuclear import of the DNA break hotspot-determining complex in fission
634   yeast. J Cell Sci *134*. 10.1242/jcs.253518.
635   43.   Ghosh, C., Nandi, A., and Basu, S. (2019). Lipid Nanoparticle-Mediated Induction of
636   Endoplasmic Reticulum Stress in Cancer Cells. ACS Applied Bio Materials *2*, 3992-4001.
637   10.1021/acsabm.9b00532.
638   44.   Liu, Y., Jia, H.-R., Han, X., and Wu, F.-G. (2021). Endoplasmic reticulum-targeting
639   nanomedicines for cancer therapy. Smart Materials in Medicine *2*, 334-349.
640   https://doi.org/10.1016/j.smaim.2021.09.001.
641   45.   Huang, C., Quinn, D., Sadovsky, Y., Suresh, S., and Hsia, K.J. (2017). Formation and
642   size distribution of self-assembled vesicles. Proceedings of the National Academy of
643   Sciences *114*, 2910-2915. doi:10.1073/pnas.1702065114.
644   46.   de Araujo, M.E.G., Liebscher, G., Hess, M.W., and Huber, L.A. (2020). Lysosomal size
645   matters. Traffic *21*, 60-75. https://doi.org/10.1111/tra.12714.
646   47.   Landsberg, P.T. (1984). D grad□v or grad(Dv)? Journal of Applied Physics *56*, 1119-
647   1122. 10.1063/1.334083.
648   48.   Van Kampen, N.G. (1988). Diffusion in inhomogeneous media. Journal of Physics and
649   Chemistry of Solids *49*, 673-677. https://doi.org/10.1016/0022-3697(88)90199-0.
650   49.   Lançon, P., Batrouni, G., Lobry, L., and Ostrowsky, N. (2002). Brownian walker in a
651   confined geometry leading to a space-dependent diffusion coefficient. Physica A:
652   Statistical Mechanics and its Applications *304*, 65-76. https://doi.org/10.1016/S0378-
653   4371(01)00510-6.
654   50.   van Milligen, B.P., Bons, P.D., Carreras, B.A., and Sánchez, R. (2005). On the
655   applicability of Fick's law to diffusion in inhomogeneous systems. European Journal of
656   Physics *26*, 913. 10.1088/0143-0807/26/5/023.
657   51.   Sattin, F. (2008). Fick's law and Fokker–Planck equation in inhomogeneous
658   environments. Physics Letters A *372*, 3941-3945.
659   https://doi.org/10.1016/j.physleta.2008.03.014.
660   52.   Huang, F., Chakraborty, P., Lundstrom, C.C., Holmden, C., Glessner, J.J., Kieffer, S.W.,
661   and Lesher, C.E. (2010). Isotope fractionation in silicate melts by thermal diffusion.
662   Nature *464*, 396-400. 10.1038/nature08840.
663   53.   Choi, B., and Kim, Y.-J. (2019). Diffusion of Biological Organisms: Fickian and Fokker--
664   Planck Type Diffusions. SIAM Journal on Applied Mathematics *79*, 1501-1527.
665   10.1137/18m1163944.
666   54.   Won, Y.-Y., and Ramkrishna, D. (2019). Revised Formulation of Fick's, Fourier's, and
667   Newton's Laws for Spatially Varying Linear Transport Coefficients. ACS Omega *4*,
668   11215-11222. 10.1021/acsomega.9b00736.
669   55.   Kim, Y.-J., and Seo, H. (2021). Model for Heterogeneous Diffusion. SIAM Journal on
670   Applied Mathematics *81*, 335-354. 10.1137/19M130087X.
671   56.   van Kampen, N.G. (1981). Itô versus Stratonovich. Journal of Statistical Physics *24*, 175-
672   187. 10.1007/BF01007642.

57.   Sokolov, I.M. (2010). Ito, Stratonovich, Hänggi and all the rest: The thermodynamics of
interpretation. Chemical Physics *375*, 359-363.
https://doi.org/10.1016/j.chemphys.2010.07.024.
58.   Bressloff, P.C., and Newby, J.M. (2013). Stochastic models of intracellular transport.
Reviews of Modern Physics *85*, 135-196. 10.1103/RevModPhys.85.135.
59.   Eliaš, J., and Clairambault, J. (2014). Reaction–diffusion systems for spatio-temporal
intracellular protein networks: A beginner's guide with two examples. Computational and
Structural Biotechnology Journal *10*, 12-22. https://doi.org/10.1016/j.csbj.2014.05.007.
60.   Yuan, J., and Sundén, B. (2014). On mechanisms and models of multi-component gas
diffusion in porous structures of fuel cell electrodes. International Journal of Heat and
Mass Transfer *69*, 358-374. https://doi.org/10.1016/j.ijheatmasstransfer.2013.10.032.
61.   Soh, S., Byrska, M., Kandere-Grzybowska, K., and Grzybowski, B.A. (2010). Reaction-
diffusion systems in intracellular molecular transport and control. Angew Chem Int Ed
Engl *49*, 4170-4198. 10.1002/anie.200905513.
62.   Aly, S., Kim, I., and Sheen, D. (2011). Turing instability for a ratio-dependent predator–
prey model with diffusion. Applied Mathematics and Computation *217*, 7265-7281.
https://doi.org/10.1016/j.amc.2011.02.018.
63.   Gotoh, T., Kim, J.K., Liu, J., Vila-Caballer, M., Stauffer, P.E., Tyson, J.J., and Finkielstein,
C.V. (2016). Model-driven experimental approach reveals the complex regulatory
distribution of p53 by the circadian factor Period 2. Proceedings of the National Academy
of Sciences *113*, 13516-13521. doi:10.1073/pnas.1607984113.
64.   Asgari-Targhi, A., and Klerman, E.B. (2019). Mathematical modeling of circadian
rhythms. WIREs Systems Biology and Medicine *11*, e1439.
https://doi.org/10.1002/wsbm.1439.
65.   Rojas, P., Piro, O., and Garcia, M. (2023). Biological rhythms generated by a single
activator-repressor loop with heterogeneity and diffusion.
66.   Purvis, J.E., Karhohs, K.W., Mock, C., Batchelor, E., Loewer, A., and Lahav, G. (2012).
p53 Dynamics Control Cell Fate. Science *336*, 1440-1444. 10.1126/science.1218351.
67.   Kim, J.K., and Jackson, T.L. (2013). Mechanisms That Enhance Sustainability of p53
Pulses. PLOS ONE *8*, e65242. 10.1371/journal.pone.0065242.
68.   Eliaš, J., Dimitrio, L., Clairambault, J., and Natalini, R. (2014). The dynamics of p53 in
single cells: physiologically based ODE and reaction–diffusion PDE models. Physical
Biology *11*, 045001. 10.1088/1478-3975/11/4/045001.
69.   Son, M., Frank, T., Holst-Hansen, T., Wang, A.G., Junkin, M., Kashaf, S.S., Trusina, A.,
and Tay, S. (2022). Spatiotemporal NF-κB dynamics encodes the position, amplitude,
and duration of local immune inputs. Science Advances *8*, eabn6240.
doi:10.1126/sciadv.abn6240.
70.   Tigges, M., Marquez-Lago, T.T., Stelling, J., and Fussenegger, M. (2009). A tunable
synthetic mammalian oscillator. Nature *457*, 309-312. 10.1038/nature07616.
71.   Purcell, O., Savery, N.J., Grierson, C.S., and di Bernardo, M. (2010). A comparative
analysis of synthetic genetic oscillators. Journal of The Royal Society Interface *7*, 1503-
1524. doi:10.1098/rsif.2010.0183.
72.   Kim, J.K., Chen, Y., Hirning, A.J., Alnahhas, R.N., Josić, K., and Bennett, M.R. (2019).
Long-range temporal coordination of gene expression in synthetic microbial consortia.
Nature Chemical Biology *15*, 1102-1109. 10.1038/s41589-019-0372-9.
73.   Macnamara, C.K., Mitchell, E.I., and Chaplain, M.A.J. (2019). Spatial-Stochastic
modelling of synthetic gene regulatory networks. Journal of Theoretical Biology *468*, 27-
44. https://doi.org/10.1016/j.jtbi.2019.02.003.
74.   Cherstvy, A.G., Chechkin, A.V., and Metzler, R. (2013). Anomalous diffusion and
ergodicity breaking in heterogeneous diffusion processes. New Journal of Physics *15*,
083039. 10.1088/1367-2630/15/8/083039.
75.   Cherstvy, A.G., and Metzler, R. (2014). Nonergodicity, fluctuations, and criticality in
heterogeneous diffusion processes. Physical Review E *90*, 012134.
10.1103/PhysRevE.90.012134.

76. Ghosh, S.K., Cherstvy, A.G., Grebenkov, D.S., and Metzler, R. (2016). Anomalous, non-Gaussian tracer diffusion in crowded two-dimensional environments. New Journal of Physics *18*, 013027. 10.1088/1367-2630/18/1/013027.

77. Wachsmuth, M., Waldeck, W., and Langowski, J. (2000). Anomalous diffusion of fluorescent probes inside living cell nuclei investigated by spatially-resolved fluorescence correlation spectroscopy11Edited by W. Baumeister. Journal of Molecular Biology *298*, 677-689. https://doi.org/10.1006/jmbi.2000.3692.

78. Seisenberger, G., Ried, M.U., Endreß, T., Büning, H., Hallek, M., and Bräuchle, C. (2001). Real-Time Single-Molecule Imaging of the Infection Pathway of an Adeno-Associated Virus. Science *294*, 1929-1932. 10.1126/science.1064103.

79. Banks, D.S., and Fradin, C. (2005). Anomalous Diffusion of Proteins Due to Molecular Crowding. Biophysical Journal *89*, 2960-2971. https://doi.org/10.1529/biophysj.104.051078.

80. Jeon, J.-H., Tejedor, V., Burov, S., Barkai, E., Selhuber-Unkel, C., Berg-Sørensen, K., Oddershede, L., and Metzler, R. (2011). In Vivo Anomalous Diffusion and Weak Ergodicity Breaking of Lipid Granules. Physical Review Letters *106*, 048103. 10.1103/PhysRevLett.106.048103.

81. Lanoiselée, Y., Moutal, N., and Grebenkov, D.S. (2018). Diffusion-limited reactions in dynamic heterogeneous media. Nature Communications *9*, 4398. 10.1038/s41467-018-06610-6.

82. Salman, H., Abu-Arish, A., Oliel, S., Loyter, A., Klafter, J., Granek, R., and Elbaum, M. (2005). Nuclear localization signal peptides induce molecular delivery along microtubules. Biophys J *89*, 2134-2145. 10.1529/biophysj.105.060160.

83. Halbi, G., Fayer, I., Aranovich, D., Gat, S., Bar, S., Erukhimovitch, V., Granek, R., and Bernheim-Groswasser, A. (2021). Nano-Particles Carried by Multiple Dynein Motors Self-Regulate Their Number of Actively Participating Motors. International Journal of Molecular Sciences *22*, 8893.

84. Snapp, E. (2005). Design and use of fluorescent fusion proteins in cell biology. Curr Protoc Cell Biol *Chapter 21*, 21 24 21-21 24 13. 10.1002/0471143030.cb2104s27.

85. Milo, R., and Phillips, R. (2015). Cell Biology by the Numbers (CRC Press).

86. Aryal, R.P., Kwak, P.B., Tamayo, A.G., Gebert, M., Chiu, P.L., Walz, T., and Weitz, C.J. (2017). Macromolecular Assemblies of the Mammalian Circadian Clock. Mol Cell *67*, 770-782 e776. 10.1016/j.molcel.2017.07.017.

87. Koch, A.A., Bagnall, J.S., Smyllie, N.J., Begley, N., Adamson, A.D., Fribourgh, J.L., Spiller, D.G., Meng, Q.J., Partch, C.L., Strimmer, K., et al. (2022). Quantification of protein abundance and interaction defines a mechanism for operation of the circadian clock. Elife *11*. 10.7554/eLife.73976.

88. Tzafriri, A.R. (2003). Michaelis–Menten kinetics at high enzyme concentrations. Bulletin of Mathematical Biology *65*, 1111-1129. https://doi.org/10.1016/S0092-8240(03)00059-4.

89. Kim, J.K., and Tyson, J.J. (2020). Misuse of the Michaelis-Menten rate law for protein interaction networks and its remedy. PLoS Comput Biol *16*, e1008258. 10.1371/journal.pcbi.1008258.

90. Song, Y.M., Hong, H., and Kim, J.K. (2021). Universally valid reduction of multiscale stochastic biochemical systems using simple non-elementary propensities. PLOS Computational Biology *17*, e1008952. 10.1371/journal.pcbi.1008952.

91. Eilertsen, J., Schnell, S., and Walcher, S. (2022). On the anti-quasi-steady-state conditions of enzyme kinetics. Mathematical Biosciences *350*, 108870. https://doi.org/10.1016/j.mbs.2022.108870.

92. Jeong, E.M., Song, Y.M., and Kim, J.K. (2022). Combined multiple transcriptional repression mechanisms generate ultrasensitivity and oscillations. Interface Focus *12*, 20210084. doi:10.1098/rsfs.2021.0084.

93. Shin, J., Choi, Y., and Kim, J. (2019). The Cahn–Hilliard Equation with Generalized Mobilities in Complex Geometries. Mathematical Problems in Engineering *2019*, 1710270. 10.1155/2019/1710270.

781   94.   Shin, J., Jeong, D., and Kim, J. (2011). A conservative numerical method for the Cahn–
782   Hilliard equation in complex domains. Journal of Computational Physics *230*, 7441-7455.
783   https://doi.org/10.1016/j.jcp.2011.06.009.