CtBP restricts DUX-dependent and -independent genetic program for the 2-cell-

like state in murine embryonic stem cells

Kazuma Yoshioka[1], Kota Sugiyama[1], Maki Ichisakino[1], Selma Alamanda Abadi[1], Nao

Hayakawa, Miyu Marutani, Ryo Masuda, Kyo Takahashi and Yoshiyuki Seki

Department of Biomedical Sciences, School of Biological and Environmental Sciences,

Kwansei Gakuin University, Hyogo, Japan

[1]Co-first author

[*]Correspondence: yseki@kwansei.ac.jp

Abstract

After fertilization, maternally deposited mRNA is cleared, and *de novo* mRNA is

transcribed from the zygotic genome through zygotic genome activation (ZGA), a

process known as maternal-to-zygotic transition (MZT) occurring in the mouse at 2-cell

(2C) stage. 2C-like cells (2CLCs) marked by MERVL expression are transcriptionally

similar to 2C embryos spontaneously emerge from mouse embryonic stem cells

(mESCs). Although the emergence of 2CLCs completely depends on DUX function, a

recent knockout study clearly showed that DUX is dispensable for mouse embryos,

suggesting that DUX-independent molecular pathways are not recapitulated in 2CLCs.

We present here that the disruption of C-terminal binding protein 1/2 (*Ctbp1/2*)

activates DUX-dependent and -independent molecular pathway associated with the

development of early mouse embryos mediated by the upregulation of Preferentially

expressed antigen of melanoma family-like 7 (PRAMEL7). Furthermore, the

abnormality of the gene expression profile caused by *Dux* KO is partially rescued by the

overexpression of PRAMEL7 in mESCs. Our study provides new insights into the

DUX-independent molecular pathway for the activation of early embryonic genes in

mESCs.

Introduction

After fertilization, maternal factors direct development and trigger zygotic genome activation (ZGA) at the around late one-cell stage in mice, called maternal-to-zygotic transition (MZT)(Eckersley-Maslin et al., 2018). The early-cleavage stage embryos acquire cellular totipotency to generate both embryonic proper and extraembryonic tissues during MZT. Although totipotency was thought to be exclusive to the early-cleavage stage embryos, a part of the molecular network is recapitulated in the totipotent-like cells (2-cell-like cells; 2CLCs) marked by the expression of endogenous retrovirus MERVL, which are an endogenous subpopulation of mouse embryonic stem cells (ESCs). (Macfarlan et al., 2012a) (Rodriguez-Terrones et al., 2018).

The transition from pluripotent stem cells into 2CLCs is a useful *in vitro* system for identifying the gene regulatory network for ZGA, and several genes that control the emergence of 2CLCs from pluripotent cells have been identified (Eckersley-Maslin et al., 2019; Hu et al., 2020; Ishiuchi et al., 2015; Percharde et al., 2018; Rodriguez-Terrones et al., 2018). The transcription factor DUX (DUX4 in humans) is a critical factor for regulating the transition from pluripotent stem cells into 2CLCs and the expression of ZGA genes (De Iaco et al., 2017; Hendrickson et al., 2017; Whiddon et al., 2017). Ectopic expression of DUX enhances the population of 2CLCs, whereas the *Dux* disruption eliminates the emergence of spontaneous 2CLCs during mESC culture. Several 2CLC regulatory genes have been identified, but they are either direct activators (DPPA2/4, p53 and NELFA) or direct repressors (CAF-1, LINE-1 and SETDB1) of *Dux* expression (Eckersley-Maslin et al., 2019; Grow et al., 2021; Ishiuchi et al., 2015; Percharde et al.,

2018; Wu et al., 2020). However, both *Dux* zygotic and maternal knockout embryos can

survive full-term (Chen and Zhang, 2019; Guo et al., 2019), suggesting that DUX-

independent molecular pathways not recapitulated in 2CLCs are critical for ZGA and the

early development of mice.

In this study, we show that disruption of Ctbp1/2 (C-terminal binding protein 1/2)

(Stankiewicz et al., 2014)activates two distinct sets of genes in ESCs: the first set is

transiently upregulated in a DUX-dependent manner during minor ZGA, and the second

set contains maternally expressed genes and the upregulated genes in a DUX-independent

manner during major ZGA. Preferentially Expressed Antigen in Melanoma-like 7

(PRAMEL7) is directly upregulated by the loss of *Ctbp1/2* without DUX function, and

upregulated PRAMEL7 activates minor ZGA genes in a DUX-dependent manner and

maternally expressed genes and major ZGA in DUX-independent manner in mESCs.

These data revealed that CtBP restricts the DUX-dependent and -independent genetic

programs for the 2-cell-like state in mESCs.

Material and Methods

Cell Culture

E14tg2a ESCs (RIKEN Cell Bank) were cultured in Glasgow minimum essential medium

(GMEM; Wako) containing 10% fetal calf serum (FCS; Invitrogen), 1 mM glutamine

(Wako), nonessential amino acids (Wako) and 500 μM Monothioglycerol Solution

(Wako). The culture was supplemented with 1000 U/ml of leukaemia inhibitory factor

(LIF) (Wako) without feeder cells. ESCs were cultured in N2B27 medium with 2i

(PD0325901, 0.4 μM: WAKO, CHIR99021, 3 μM: Sigma-Aldrich).

Generation of knockout ESCs using a CRISPR-Cas9 system

Guide RNAs for *Ctbp1/2*, *Dux* and *Pramel7* were generated in pX330-U6- Chimeric_BB-

CBh-hSpCas9 (Addgene plasmid #42230; deposited by Feng Zhang; Cong et al., 2013).

The guide RNA sequences are listed in Supplementary Table 1. The pX330 was co-

transfected with pCAG-EGFP into E14tg2a with MERVL::TurboGFP (Addgene plasmid

#69072; deposited by Maria-Elena Torres-Padilla; (Ishiuchi et al., 2015). The GFP-

positive fraction was sorted by Cell Sorter (No. SH800; Sony Biotechnology, Inc., Tokyo,

Japan). Genomic DNA deletions were sequenced (Supplementary Figure 1).

qRT-PCR and RNA-Seq

Total mRNA was purified with a PureLink™ RNA Mini Kit (Ambion, Inc.). For

quantitative RT-PCR, cDNA was synthesized by ReverTra Ace® qPCR RT master Mix

(Toyobo), and then analyzed by qPCR with FastStart SYBR Green Master Mix (Roche)

using a specific primer set for each gene (Supplementary Table 1). RNA-seq libraries

were generated using Collibri^{TM} Stranded RNA library prep for ilIumina^{TM} Systems

(ThermoFisher Scientific) and sequenced using HiSeq X^{TM} Ten (Illumina) (150-bp

paired-ends reads).

98   Data analysis of RNA-Seq and public ChIP-seq data

Total mRNA of each cell line was purified with a PureLink^{TM} RNA Mini Kit (Ambion,

Inc.) by biological duplicate. RNA-seq libraries were generated using Collibri^{TM} Stranded

RNA library prep for ilIumina^{TM} Systems (ThermoFisher Scientific) and sequenced using

HiSeq X^{TM} Ten (Illumina). Reads were aligned by Hisat2 to the Mus musculus genome

(mm10). Raw read counts of genes were calculated by R package FeatureCounts using

mm10 KnownGenes. GTF from UCSC for annotation. Normalized count data, principal

component and K-means clustering analysis were exported by the interactive Expression

and Pathway analysis (iDEP) webserver using the count data or normalized expression

data of early mouse embryos (Ge et al., 2018). Differentially expressed genes (DEGs)

were selected using a cut-off at a false discovery rate (FDR) $\leqq$ 0.1 and a $\geqq$ 2-fold

change in expression. Gene set enrichment analysis (GSEA) was used to determine

whether upregulated genes by *Ctbp1/2* knockouts were enriched for the upregulated genes

in 2-cell embryos or 2CLCs as described previously. Venn diagrams showing overlapping

genes were constructed by BioVenn, a web application (Hulsen et al., 2008).

Pubic ChIP-seq data for CtBP2 were re-analyzing using GREAT (Genomic Regions

Enrichment of Annotations tool) with the "two nearest genes" setting, which assigns each

peak to the two closest transcription start sites (TSSs) within 1 Mb. Peak visualization of

ChIP-seq peaks for CtBP2, H3K27Ac and DUX was performed using Integrative

Genomics Viewer (IGV) (Hendrickson et al., 2017; Kim et al., 2015; McLean et al., 2010;

Zhang et al., 2020).

Immunofluorescence analysis

Cells were fixed by 4% paraformaldehyde at 4℃ for 20 min, followed by incubation with

PBS containing 0.5% Triton-X at 4℃ for 5 min. Permeabilized cells were incubated with

a primary antibody solution containing anti-GFP (Nacarai Tesque, 04404-84, 1:500) with

anti-ZSCAN4 (Merk Millipore, AB4340, 1:1000) or anti-MuERVL-Gag (Epigentek, A-

2801-050, 1:250) or anti-OCT4 (Santa Cruz, Sc-8628, 1:500) or anti-CtBP2 (BE

Biosciences, 612044, 1:500) followed by secondary antibody solution containing anti-

Rat IgG Dylight^{TM} 488 antibody (Rockland, 612-141-120, 1:500) or anti-Rabbit IgG

Dylight^{TM} 549 antibody (KPL, 042-04-15-06, 1:500) or anti-Mouse IgG Dylight® 550

antibody (Abcam, ab96876, 1:500) or anti-Goat IgG Dylight^{TM} 549 antibody (Rockland,

605-742-002, 1:500) with 1 μg/ml of DAPI solution (DOJINDO, 340-07971).

Luciferase assay

The enhancer candidate region of *Pramel7* or *Dux* was cloned into pGL4.23, driving the

luciferase gene. Wild-type or *Ctbp1/2* DKO ESCs were transfected with the reported

constructs with *Ctbp2* or *Pramel7* expression vector using HilyMax (DOJINDO). Cells

were harvested 36 hours after transfection, and the luciferase activities were measured

using the Dual-luciferase Reporter Assay System (Promega).

Western blotting analysis

Cells were lysed by boiling SDS sample buffer (WAKO). Before lysed proteins were

applied to polyacrylamide-SDS gels, 2-mercaptoethanol was added to denature the

proteins. The lysed proteins were separated on polyacrylamide-SDS gels, blotted on a

polyvinylidene fluoride membrane, and probed using the following primary antibodies:

anti-UHRF1 (MBL, D289-3, 1:1000) or anti-TUBLIN (SIGMA, T5168, 1:5000).

Following the primary antibody reaction, the membrane were incubated with horseradish

peroxidase (HRP)-coupled secondary antibodies. All antibodies were detected with

Immunostar LD (WAKO).

Phylogenetic tree inference

Amino acid sequences of PRAME family proteins were obtained from Ensemble release

112. Multiple-sequence alignment was performed using the ClustalW algorithm with the

MEGA software, version 11 with default parameters. Molecular phylogenetic trees were

inferred with the maximum-likelihood method using MEGA software, version 11 with

default parameters.

Results

CtBP1/2 acts as negative regulators of the transition from pluripotent ESCs to two-cell-like cells (2CLCs)

We have previously shown that CtBP1/2 are essential for the transition from formative to naïve pluripotency in mouse embryonic stem cells (mESCs) (Yamamoto et al., 2020). To investigate their roles of CtBP1/2 beyond pluripotency maintenance, we performed RNA sequencing (RNA-seq) to compare the gene expression profiles between wild-type (WT) and *Ctbp1/2* double-knockout (DKO) ESCs. An MA-plot revealed a marked upregulation of two-cell (2C)-specific genes, including *Dux*, *Zscan4* and *Zfp352*, together with repeat elements such as MERVL, MT2 and GSAT_Mm in *Ctbp1/2* DKO ESCs (Figure 1A, B). Gene set enrichment analysis (GSEA) using a reference set of 2C embryo and 2C-like cell (2CLC)-enriched genes (n=404; (Macfarlan et al., 2012)) confirmed a strong enrichment of 2C-related transcripts among those upregulated in the absence of CtBP1/2 (Figure 1C).

To directly monitor the induction of 2CLC state, we generated *Ctbp1/2* DKO ESCs carrying the MERVL-EGFP reporter (Ishiuchi et al., 2015). The GFP-positive fraction was approximately 0.1 % in WT ESC, whereas *Ctbp1/2* DKO ESCs exhibited an over 50-fold increase (7.4%) in GFP-positive cells, indicative of robust 2CLC induction (Figure 1D). These GFP-positive cells co-expressed hallmark 2CLC markers such as ZSCAN4 and MERVL-GAG, while pluripotency-associated factors such as OCT4 were markedly reduced (Figure 1E), consistent with previous reports (Eckersley-Maslin et al., 2016; Zhu et al., 2021).

Figure 1

Figure 1. Disruption of *Ctbp1/2* expands the 2-cell-like cell (2CLC) population.

(A), (B) MA-plots showing the relationship between average gene (or repeat) expression

and log$_2$ fold change between *Ctbp1/2* DKO and wild-type (WT) ESCs. Differentially

expression was defined as adjusted $p \leq 0.05$ and fold change ≥ 2. Red and blue dots

indicate up- and down-regulated genes, respectively. (C) Gene set enrichment analysis

(GSEA) of genes upregulated in 2-cell embryos compared with oocytes and enriched in

MERVL-positives 2CLCs relative to MERVL-negative ESCs (GSE33923) (Macfarlan et

al., 2012). (D) Flow cytometry analysis of MERVL::TurboGFP reporter activity in WT

ESCs and *Ctbp1/2* DKO ESCs. (E) Immunofluorescence analysis showing that

MERVL::TurboGFP-positive cells co-express ZSCAN4 and MERVL-GAG proteins, and

are mutually exclusive with OCT4 expression. Nuclei were counterstained with DAPI.

Scale bar: 10 μm.

CtBP1/2 repress DUX-dependent minor ZGA genes and DUX-independent maternal

and major ZGA genes in ESCs

DUX, the homeobox transcription factor, is both essential and sufficient for the

emergence of spontaneous 2CLCs from ESCs (De Iaco et al., 2017; Whiddon et al., 2017).

To determine whether the 2C-like program triggered by *Ctbp1/2* deficiency depends on

DUX, we deleted the *Dux* cluster in both WT and *Ctbp1/2* DKO ESCs (Supplementary

Figure 1C). We first compared the population of MERVL-EGFP-positive cells among

WT, *Dux* KO, *Ctbp1/2* DKO and *Ctbp1/2/Dux* triple-KO (TKO) ESCs. While the strong

GFP-positive population was markedly reduced in TKO compared to DKO ESCs, a weak

GFP-positive population persisted in the TKO (Figure 2A, B). RT-qPCR analysis

confirmed that *Pramel7*, which is upregulated at the 2C stage in WT and *Dux* KO

embryos (Guo et al., 2019), remained elevated in both DKO and TKO ESCs, indicating

DUX-independent activation (Figure 2C). In contrast, *Zscan4c* upregulation upon

Ctbp1/2 depletion was completely abolished by *Dux* deletion, indicating strict DUX

dependency.

To assess transcriptional activity of the MERVL long terminal repeat (MERVL-LTR)

sequences, we performed a luciferase reporter assay (Supplemental Figure 2A). *Ctbp1/2*

loss led to robust activation of MERVL-LTR reporter, and mutation of DUX-binding

motif (DUXmut-MERVL-LTR) markedlyreduced, but did not abolish, this activity.

Reporter activity in *Ctbp1/2/Dux* TKO remained substantially higher than *Dux* KO,

indicating that CtBP1/2 also repress MERVL through DUX-independent mechanisms.

Published DUX ChIP-seq data (Hendrickson et al., 2017) revealed DUX occupancy near

both the *Zscan4c* and *Naalad2* loci, with strong binding signals overlapping MERVL-

LTR or MT2 family elements (Supplementary Figure 2B). At the *Zscan4c* locus, DUX

binds both proximal regulatory region with a strong peak and distal MERVL elements. At

the *Naalad2* locus, DUX-binding sites are exclusively overlapped with MERVL

sequences. Reporter assay confirmed that MERVL-independent elements proximal to the

Zscan4c locus were activated by *Ctbp1/2* loss in a DUX-dependent manner, whereas

MERVL-LTR-containing elements derived from the *Zscan4c* and *Naalad2* loci retained

partial activity even in *Ctbp1/2/Dux* TKO ESCs (Supplementary Figure 2C). These

luciferase assay results were consistent with our RNA-seq data (Supplementary Figure

2E).

We next performed RNA-seq to identify genes induced by *Ctbp1/2* loss that are DUX-

dependent or DUX-independent. A scatter plot comparing gene expression changes

between WT and *Ctbp1/2* DKO (Y-axis) and between *Ctbp1/2/Dux* TKO and *Dux* KO

(X-axis) revealed a moderate correlation (R = 0.52), suggesting that a substantial

population of CtBP1/2-regulated genes act independent of DUX (Figure 2D). Venn

diagram showed that more than 75% of genes upregulated in DKO were also upregulated

in TKO ESCs (Figure 2E).

To classify these regulatory patterns, we applied K-means clustering analysis (K=5) using

the 2,000 most variable genes across WT, *Dux* KO, *Ctbp1/2* DKO, and *Ctbp1/2/Dux* TKO

ESCs (Figure 2F, G). Cluster 1 genes (e.g., *Zscan4*, *Usp17*) were slightly repressed in

Dux KO ESCs but strongly upregulated in *Ctbp1/2* DKO ESCs, and this upregulation was

abolished by *Dux* KO, defining DUX-dependent minor ZGA genes. Cluster 2 genes (e.g.,

Gadd45g) were upregulated in both *Ctbp1/2* DKO and *Ctbp1/2/Dux* TKO ESCs,

representing DUX-independent target. Cluster 3 genes (e.g., *Gata1*, *Pramel7*, *Zscan5b*)

showed moderate downregulation in *Dux* KO ESCs and were modestly upregulated in

both *Ctbp1/2* DKO and *Ctbp1/2/Dux* TKO ESCs, corresponding to major ZGA and

maternal genes. Cluster 4 (e.g, *Spic*) genes were downregulated by *Dux* KO in WT ESCs.

However, this downregulation by *Dux* KO was not observed in *Ctbp1/2* DKO ESCs,

indicating that the loss of CtBP1/2 abolishes the dependency of these genes on DUX.

Cluster 5 genes were consistently downregulated in both *Ctbp1/2* DKO and *Ctbp1/2/Dux*

TKO ESCs, indicating positive regulation by CtBP1/2 in a DUX-independent manner.

To evaluate the expression dynamics of these clusters in early mouse embryos, we

compared these clusters with public RNA-seq data from early embyos (Zhang et al.,

2016), excluding genes with an FPKM average ≤ 1 (Figure 2H). Cluster 1 genes were

transient activated at the 2-4-cell stage, typical of minor ZGA genes, whereas Cluster2-4

included maternal genes downregulated post-fertilization and zygotic genes activated

after the late 2-cell stage.

Finally, to further examine the dependency of genes upregulated by *Ctbp1/2* DKO on

DUX, we measured the expression changes of genes belonging to Cluster 1 and 3 upon

inducible *Dux* expression using doxycycline-inducible *Dux* ESCs (Figure 2I). Consistent

with our RNA-seq data, Cluster 1 genes were consistently upregulated whereas Cluster 3

genes showed little or no change following short-term DUX induction. Taken together,

these results demonstrate that the *Ctbp1/2* depletion broadly upregulates DUX-dependent

minor ZGA genes and DUX-independent maternal and major ZGA genes, revealing a

dual regulatory system that links CtBP1/2-mediated repression to early embryonic

transcriptional programs.

Figure 2

Figure 2. CtBP1/2 repress both DUX-dependent and -independent transcriptional programs associated with preimplantation embryos in mouse ESCs.

(A) Flow cytometry analysis (FACS) of MERVL::TurboGFP reporter expression in wild-

type (WT), *Dux* knockout (KO), *Ctbp1/2* double knockout (DKO) and *Ctbp1/2/Dux* triple

knockout (TKO) ESCs. (B) Percentage of MERVL-EGFP-positive cells obtained by

FACS Shown are standard error of means (SEM) at biological triplicates. (C) RT-qPCR

analysis of 2C-associated genes in WT, *Dux* KO, *Ctbp1/2* DKO and *Ctbp1/2/Dux* TKO

ESCs. Values are shown as relative expression with WT set to 1. Error bar indicates the

standard error of means. ***$P < 0.001$, **$P < 0.005$, *$P < 0.05$ (Tukey-Kramer multiple

comparison test). (D) Scatter plot comparing gene expression changes induced by

Ctbp1/2 DKO in WT (Y-axis) and *Dux* KO (X-axis) backgrounds. (E) Venn diagram

showing the overlap of genes upregulated in *Ctbp1/2* DKO and TKO. (F) K-means

clustering analysis (K=5) of the top 2,000 differentially expressed genes across WT, *Dux*

KO, *Ctbp1/2* DKO and *Ctbp1/2/Dux* TKO ESCs. (G) Average expression profiles of

genes from each cluster at WT, *Dux* KO, *Ctbp1/2* DKO and *Ctbp1/2/Dux* TKO ESCs.

Grey lines indicate individual gene, and black line represents the average expression of

genes within each cluster. (H) Hierarchical clustering of genes from each cluster using

public RNA-seq data of early mouse embryo (Zhang et al., 2016). (I) RT-qPCR after *Dux*

induction (0 h vs 24 h) for selected genes (Cluster 1: *Dux, Zscan4c, Obox4, Pramel16*

and Cluster 3: Gata1, Klf17, Pramel7). The ΔCt of each gene was calculated by

subtracting the average Ct of Gapdh from that of each gene Ct. Means of ΔCt were

calculated from three biological triplicates. Error bar indicates the standard error of means

(SEM). ***$P < 0.001$, **$P < 0.005$, *$P < 0.05$ (Tukey-Kramer multiple comparison test).

Temporal regulation of ZGA-associated genes by CtBP1/2 defines rapid and delayed

response genes

To monitor the dynamics of ZGA-asscoated gene expression in response to CtBP2

induction and depletion in ESCs, we established *Ctbp1/2* DKO ESCs carrying a

doxycycline-inducible CtBP2 transgene. Using this system, we analyzed the expression

of representative ZGA-related genes by qRT-PCR. *Dux*, MERVL, *Zscan4c* and *Zfp352*

were markedly downregulated between day 3 and day 9 following CtBP2 induction, and

subsequently reactivated from day 12 to day 18 upon CtBP2 depletion (Figure 3A). In

contrast, *Pramel7* exhibited a rapid response, being strongly downregulated between

day 0 and day 3, and then upregulated between day 9 and day 12.

To distinguish between rapid and delayed transcriptional responses, we performed

RNA-seq using *Ctbp1/2* conditional KO (cKO) ESCs at defined time points. Principal

component analysis (PCA) revealed that short-term CtBP2 induction (days 0-3) and

depletion (days 9-12) drove opposing changes along the PC1 axis, whereas long-term

induction (days 3-9) and depletion (days 12-18) contributed to opposite shifts along

PC2 (Figure 3B). By overlapping differentially expressed genes under these conditions,

we identified two kinetically distinct groups: rapid-response genes, including major

ZGA genes (*Pramel7*, *Gata1*, *Phlda2*) and maternal factors (*Klf17*, *Pramel13*) (Figure

3C), and delayed-response genes, comprising minor ZGA genes such as *Zscan4*,

Usp17l, and *Gm4340* (Figure 3D). Temporal trajectory analysis showed that rapid-

response genes (N = 89) were immediately repressed upon CtBP2 induction (days 0-3)

and quickly reactivated following CtBP2 depletion (days 9-12) (Figure 3E). By

contrast, delayed-response genes (N = 48) displayed gradual downregulation during

CtBP2 expression and delayed recovery upon CtBP2 depletion (Figure 3F). Heatmap

analysis illustrated their distinct temporal expression patterns (Figure 3G).

Finally, to assess the DUX dependency of these gene sets, we compared gene

expression changes induced by *Ctbp1/2* DKO in WT and *Dux* KO backgrounds. Rapid-

response genes were equivalently upregulated in both context, whereas the induction of

delayed-response genes were significantly attenuated in the absence of DUX (Figure

3H). Collectively, these results indicate that CtBP1/2 represses two transcriptional

cascades with distinct temporal kinetics: a DUX-independent program directly targeting

major ZGA and maternal genes, and a DUX-dependent program that controls minor

ZGA genes.

Figure 3

Figure 3. Temporal regulation of ZGA genes by CtBP1/2 defines rapid and delayed

responses.

(A) RT-qPCR analysis of *Dux*, MERVL*, Zscan4c Zfp352* and *Pramel7* in *Ctbp1/2* DKO

ESCs with or without doxycycline (Dox). The ΔCt of each gene was calculated by

subtracting the average Ct of *Gapdh* from that of each gene Ct. Means of ΔCt were

calculated from three biological triplicates. Error bars indicate the standard error of

means (SEM). ***$P < 0.001$, **$P < 0.005$, *$P < 0.05$ (Tukey-Kramer multiple

comparison test). (B) Principal component analysis (PCA) of *Ctbp1/2* DKO ESCs with

or without Dox. (C) Venn diagram showing the overlap between genes downregulated 3

days after CtBP2 induction and genes upregulated 3 days after CtBP2 depletion. (D)

Venn diagram showing the overlap between genes downregulated during CtBP2

induction (day3-9) and genes upregulated during CtBP2 depletion (day 12-18). (E, F)

Temporal expression dynamics of rapid and delayed response genes upon CtBP2

induction or depletion. Individual gene trajectories (thin lines) and the mean ± SEM

(bold lines with error bars) are shown for Rapid response genes (N = 89, panel E) and

Delayed response genes (N = 48, panel F). (G) Heatmap showing the expression

profiles of selected rapid and delayed response genes. (H) Boxplot showing expression

changes of rapid and delayed response genes upon CtBP1/2 depletion in wild-type and

Dux KO backgrounds.

CtBP2 directly represses major ZGA genes Pramel7 and Gata1 in ESCs

To examine whether CtBP2 directly regulates major ZGA genes in ESCs, we analyzed

the CTBP2 occupancy at rapid- and delayed- response genes using public ChIP-seq

datasets (Kim et al., 2015). Genomic peaks were annotated to nearby genes using

Genomic Regions Enrichment of Annotations tools (GREAT) with the "two nearest

genes" setting (McLean et al., 2010), which assigns each regulatory region to the two

closest transcription start sites (TSSs) within 1 Mb (Figure 4A). Approximately half of

the rapid-response genes (40 out of 89), but only 2 of 48 delayed-response genes,

contained CtBP2-binding peaks. Genome browser views of H3K27Ac (Zhang et al.,

2020), a marker for active enhancers, DUX (Hendrickson et al., 2017), and CtBP2

occupancy demonstrated that CtBP2 was bound at regulatory regions of rapid-response

genes such as *Pramel7* and *Gata1*, whereas DUX binds to delayed-response loci such as

Duxf3 and *Zscan4c* (Figure 4B, C).

To functionally validate the repressive activity of CtBP2-bound regions, we performed

luciferase reporter assays using distal enhancer fragments derived from *Pramel7* and

Gata1 loci in wild-type and *Ctbp1/2* DKO ESCs. Each enhancer activities was elevated

in *Ctbp1/2* DKO ESCs and was suppressed upon CtBP2 re-expression (Figure 4D, E),

supporting the idea that CtBP2 directly represses *Gata1* and *Pramel7* transcription

through their distal enhancer elements in ESCs.

Immunofluorescence analysis further showed that MERVL-EGFP-positive cells

expressed GATA1 in WT ESCs, whereas this expression was abolished in *Dux* KO

ESCs (Figure 4F). Consistent with the FACS profiles described earlier (Figure 2A), we

observed both EGFP-weak and strong-populations in *Ctbp1/2* DKO ESCs, and both

populations expressed GATA1. Deletion of *Dux* in *Ctbp1/2* DKO ESCs abolished

selectively eliminated the EGFP-strong, and GATA1-positive cells, whereas the EGFP-

weak-positive cells, GATA1-positive cells persisted. These findings suggest that

Ctbp1/2 DKO ESCs comprise two transcriptionally distinct 2CLC population: (1) a

DUX-dependent, GATA1-positive subset corresponding to classical 2CLCs marked by

strong MERVL-EGFP-strong expression, and (2) a DUX-independent, GATA1-positive

subset representing a novel 2CLC state associated with major ZGA gene activation.

Figure 4

Figure 4. Major ZGA genes, *Pramel7* and *Gata1*, are direct targets of CtBP2 in ESCs.

(A) Proportion of CtBP2-bound genes in rapid- and delayed-response groups based on public CtBP2 ChIP-seq data (Kim et al., 2015). Peak-to-gene assignment performed using GREAT (McLean et al., 2010) under the "two nearest genes" rule (≤1 Mb). (B, C) Genome browser views showing H3K27ac (Zhang et al., 2020), DUX (Hendrickson et al., 2017) and CtBP2 ChIP-seq signals at representative rapid (B) and delayed (C)

response genes. (D, E) Luciferase assays using distal regulatory regions of *Pramel7* (D)

or *Gata1* (E) loci in wild-type (WT) and *Ctbp1/2* DKO ESCs with or without CtBP2

overexpression. (F) Immunofluorescence analysis of WT, *Dux* KO, *Ctbp1/2* DKO and

Ctbp1/2/Dux TKO ESCs stained with anti-GFP (green) and GATA1 (red) antibodies.

Nuclei were counterstained with DAPI. Arrowheads indicate MERVL-EGFP-positive

cells.

PRAMEL7 regulates subsets of both DUX-dependent and -independent ZGA-

associated genes

Pramel7 belongs to the PRAME (preferentially expressed antigen in melanoma) gene

family, whose expression is normally restricted to testis and cancers (Kern et al., 2021).

As the third largest gene family in mouse genome (Church et al., 2009), we examined

its regulation in wild-type (WT), *Dux* KO, *Ctbp1/2* DKO and *Ctbp1/2/Dux* TKO ESCs.

A phylogenetic tree constructed from amino acid sequences annotated in mm39

revealed that PRAME family genes are organized into six chromosomal clustres, with

high intracluster similarity (Supplementary Figure 3A, B). RNA-seq analysis showed

that most PRAME—except *Lrrc14*, *Lrrc14b*, and *Pramel48*—were markedly

upregulated in *Ctbp1/2* DKO ESCs (Figure 5A).

Based on expression patterns across WT, *Dux* KO, *Ctbp1/2* DKO and *Ctbp1/2/Dux*

TKO ESCs, PRAME genes were classified into four groups: (1) genes unaffected by

Ctbp1/2 DKO (e.g., Lrrc14); (2) genes further upregulated by *Dux* KO (e.g.,

Pramel14/30); (3) genes suppressed by *Dux* KO (e.g., *Pramel16/19/31*, DUX-

dependent); and (4) genes unaffected by *Dux* KO (e.g., *Pramel7/12*, DUX-independent).

Single-cell RNA-seq analysis of early embryonic stages (Zhang et al., 2016) revealed

that DUX-dependent PRAME genes (e.g., *Pramel16/19/31*) are transiently induced at

the mid-2-cell stage. In contrast, DUX-independent PRAME genes could be further

subdivided into two categories (Figure 5B): (i) *Pramel6/7/30*, which are upregulated

from mid-2-cell and peak at ICM before being silenced in ESCs; and (ii) *Pramel12/13*,

which are maternally expressed and downregulated by at 4-cell stage.

Given the DUX-independent function of *Pramel7* and its known importance in early

mouse development (Graf et al., 2017), we generated ESCs overexpressing *Pramel7*

(P7OE) in both WT and *Dux* KO backgrounds (Figure 5C, D). *Pramel7* overexpression

increased the fraction of MERVL-EGFP-positive 2CLCs (Figure 5E, F), although to a

lesser extent than in *Ctbp1/2* DKO. This increase was abolished in *Dux* KO

background, and the MERVL-EGFP-weak positive population was absent, indicating

that PRAMEL7-driven 2CLC partially depends on DUX.

We next compared genes upregulated by PRAMEL7 with those upregulated by

Ctbp/1/2 DKO in either DUX-dependent or -independent contexts. Venn diagram

analysis revealed that a subset of genes activated in a DUX-dependent (55 out of 390)

as well as DUX-independent manner (194 out of 1,097) upon *Ctbp1/2* DKO were also

upregulated by PRAMEL7 overexpression (Figure 5G). To further investigate the

transcriptional programs controlled by PRAMEL7, we performed K-means clustering

(K=5) of the top 1,000 most variable genes from RNA-seq of WT, *Dux* KO, P7OE and

P7OE/*Dux* KO ESCs (Figure 5H). Cluster 1 contained PRAMEL7-induced genes

unaffected by *Dux* KO, including germline-specific and DNA methylation-repressed

genes (e.g., *Xlr3/4/5*, *Hormad1*) (Escalier et al., 2002; Shin et al., 2010), which were

enriched for meiotic cell cycle pathways (Figure 5I). Cluster 3 genes, including *Zscan4*

and *Usp17l* family members, were upregulated by PRAMEL7 overexpression but

completely abolished upon *Dux* KO background (Figure 5J), indicating that PRAMEL7

indirectly promotes the expression of minor ZGA genes through *Dux* activation,

consistent with increased *Duxf3* expression (Figure 5J). Cluster 4 comprised genes

downregulated by *Dux* KO ESCs in WT ESCs but not in P7OE ESCs, suggesting that

these genes are suppressed as a consequence of *Pramel7* downregulation in the absence

of DUX. Consistent with this, *Pramel7* expression was significantly reduced in Dux KO

ESCs (Figure 5K). Together, these results identify PRAMEL7 as a key mediator that

links DUX-dependent and -independent transcriptional programs in establishing the

totipotent-like states.

Figure 5

Figure 5. PRAMEL7 controls a subset of ZGA-associated genes through DUX-dependent and -independent mechanisms.

(A) Hierarchical clustering heatmap of of PRAME family genes from RNA-seq of wild-

type (WT), *Dux* KO, *Ctbp1/2* DKO and *Ctbp1/2/Dux* KO ESCs. (B) Expression dynamics

of PRAME family genes during early mouse embryos (Zhang et al., 2016). Red letters

indicate DUX-dependent genes. (C) Western blot analysis of FLAG-tagged PRAMEL7

protein levels WT, PRAMEL7-overexpressing (P7OE), *Dux* KO and P7OE/*Dux* KOESCs.

(D) Genotype PCR confirming large deletion of *Dux* cluster locus using primer sets

flanking the gRNA target site. (E) Fluorescence images showing MERVL-EGFP signals

in WT, P7OE, *Dux* KO and P7OE/*Dux* KO ESCs. (F) FACS analysis quantifying the

percentage of MERVL-EGFP-positive cells in WT, P7OE, *Dux* KO and P7OE/*Dux* KO

ESCs. (G) Venn diagram showing the overlap of genes upregulated by *Ctbp1/2* DKO

(with or without DUX) and those induced by PRAMEL7 overexpression (H) K-means

clustering (K=5) of the top 1,000 most variable genes from RNA-seq of WT, P7OE, *Dux*

KO and P7OE/*Dux* KO ESCs. ((I) Gen ontology enrichment analysis of each gene cluster

identified in (H). (J) RT-qPCR analysis of *Duxf3* in WT and P7OE ESCs. (K) RT-qPCR

analysis of *Pramel7* in WT and <u>*Dux*</u> KO ESCs. The ΔCt values were calculated relative

to *Gapdh*, and error bar indicates the standard error of means (SEM) from three biological

replicates. Statistical significance was assessed by Tukey-Kramer multiple comparison

test (***$P < 0.001$, **$P < 0.005$, *$P < 0.05$).

Dux controls part of the ZGA transcriptional program through PRAMEL7 functions

To investigate the functional reationship between DUX and PRAMEL7, we generated

Pramel7 KO ESCs and compared transcriptomic changes between *Dux* KO and

Pramel7 KO ESCs using scatter plot analysis (Figure 6A). Global expression alterations

showed a moderate positive correlation ($R = 0.39$), particularly among downregulated

genes. Gene ontology (GO) analysis of genes downregulated in both *Pramel7* KO and

Dux KO ESCs revealed enrichment for terms related to "Nucleic acid binding" and

"Homeodomain", including *Zfp600*, *Zic2* and *Cphx*, *Pou3f1*. In contrast, genes

downregulated genes only in *Dux* KO were enriched for "Multicellular organism

development" and "Metal ion binding", including *Piwil1*, *Dppa5* and *Zsacn4c*, *Gata1/2*

(Figure 6B). The Overlap between genes upregulated in *Dux* KO and *Pramel7* KO was

modest compared with strong overlap observed for downregulated genes (Figure 6C).

Furthermore, GSEA demonstrated that the overexpression of PRAMEL7 in *Dux* KO

ESCs effectively rescued the downregulated genes resulting from *Dux* KO (Figure 6D),

indicating that downregulation of *Pramel7* expression is major cause of the

transcriptional abnormalities in *Dux* KO ESCs.

Because PRAMEL7 is known to promote UHRF1 degradation, we next examined

whether DNA methylation contributes to gene silencing in *Dux* and *Pramel7* KO ESCs.

More than half of the genes downregulated in both *Dux* KO and *Pramel7* ESCs were

upregulated by *Dnmt1* KO ESCs (Figure 6E), suggesting that the transcriptional

repression observed in these mutants is mediated, at least in part, by aberrant DNA

methylation.

We next analyzed the expression dynamics of genes downregulated in both *Dux* KO and

Pramel7 KO ESCs, as well as those specifically downregulated only in *Dux* KO ESCs,

using public single-cell RNA-seq data from early mouse embryos (Zhang et al., 2016).

Gene downregulated genes both in *Dux* KO and *Pramel7* KO ESCs were primarily

associated with maternally expressed genes, major ZGA genes and post-ZGA genes

(Figure 6F). In contrast, genes downregulated only in *Dux* KO ESCs included not only

maternally expressed genes, major ZGA genes and post-ZGA genes but also minor

ZGA genes (Figure 6G).

Taken together, these findings suggest a mechanistic model in which DUX primarily

activate minor ZGA genes, whereas PRAMEL7—one of the indirect downstream

targets of DUX—regulates maternally, major ZGA genes and post-ZGA genes through

modulation of DNA methylation mediated by UHRF1 degradation in 2CLCs.

Figure 6

Figure 6. DUX regulates part of ZGA gene program through activation or *Pramel7*

in ESCs.

(A) Scatter plot showing the correlation between expression changes in *Dux* KO and

Pramel7 KO ESCs. (B) Venn diagram showing the overlapping of the downregulated

genes by *Dux* KO in ESCs with the downregulated genes by *Pramel7* KO in ESCs. (C)

Venn diagram showing the overlapping of the upregulated genes by *Dux* KO in ESCs

with the upregulated genes by *Pramel7* KO in ESCs. (D) Gene set enrichment analysis

(GSEA) using genes downregulated by the *Dux* KO to compare gene expression between

PRMAEL7-overexpressing in *Dux* KO ESCs and *Dux* KO ESCs. (E) Venn diagram

showing the overlap among genes upregulated in *Dnmt1* KO ESCs and genes

downregulated in *Pramel7* KO and *Dux* KO ESCs. (F, G) Heatmaps showing hierarchical

clustering of genes downregulated genes in both *Dux* KO and *Pramel7* KO ESCs (F), or

exclusively in *Dux* KO ESCs (G), using public RNA-seq data of early mouse embryo

(Zhang et al., 2016).

Discussion

Figure 7

Figure 7. A schematic model for the regulation of 2C-related genes by DUX, CtBP1/2

and PRAMEL7.

In this study, we demonstrate that the loss of *Ctbp1/2* in ESCs activates both DUX-

dependent and DUX-independent transcriptional cascades to establish a totipotent-like

state (Figure 7). Most DUX-dependent genes activated upon *Ctbp1/2* depletion

correspond to minor ZGA genes transiently expressed at the two-cell (2C) stage. In

contrast, DUX-independent genes upregulated upon *Ctbp1/2* loss can be classified into

maternally expressed genes and post-major ZGA genes. Among them, *Pramel7* is directly

upregulated by *Ctbp1/2* loss in a DUX-independent manner and further modulates both

DUX-dependent and -independent pathways through the activation of DUX and its

downstream targets. Collectively, these findings uncover a dual regulatory framework in

which CtBP1/2 represses both DUX-mediated and DUX-independent transcriptional

programs, providing new insights into molecular mechanisms governing early mouse

embryogenesis.

Our data clearly demonstrate that the loss of *Ctbp1/2* in ESCs upregulates maternally

expressed and post-major ZGA genes in a DUX-independent manner, highlighting an

alternative regulatory axis operating during early embryonic development. Interestingly,

recent studies have demonstrated that OBOX proteins, a family of PRD-like homeobox

transcription factors, can compensate for DUX function by directly activating MERVL as

well as both minor and major ZGA genes (Guo et al., 2024; Ji et al., 2023; Sakamoto et

al., 2024). However, our transcriptome analysis revealed that only *Obox4* was upregulated

in *Ctbp1/2* double-knockout (DKO) ESCs, and this upregulation was completely

abolished by *Dux* KO. These results suggest that the loss of *Ctbp1/2* regulates early

developmental genes through pathways independent of both DUX and OBOX.

By integrating transcriptome data from conditional *Ctbp2* KO ESCs with public CtBP2

ChIP-seq datasets, we identified direct CtBP2 target genes (e.g., *Pramel7*, *Gata1*, and

Klf17) and distinguished them from DUX-dependent regulatory genes. Furthermore, we

found that MERVL-EGFP-weak positive cells in both *Ctbp1/2* DKO and *Ctbp1/2/Dux*

TKO ESCs expressed GATA1, a direct CtBP2 target. Importantly, GATA1-positive cells

were absent in *Dux* KO ESCs, suggesting that *Ctbp1/2* loss not only relieves DUX

dependency but also rewires the transcriptional hierarchy governing ZGA-associated

programs.

Prame is one of the most amplified gene families in mammals, and the mouse *Prame*

gene family represents the third-largest multigene family in the genome (Chang et al.,

2011). We showed that the disruption of *Ctbp1/2* activates the majority of *Prame* family

genes in ESCs, some of which are upregulated by DUX, while others are induced directly

by the loss of CtBP1/2. DUX-dependent *Prame* genes are enriched among minor ZGA

genes, whereas DUX-independent *Prame* genes are upregulated after major ZGA.

Moreover, many *Prame* genes are rapidly repressed upon the re-expression of CtBP2 in

Ctbp1/2 DKO ESCs, indicating direct repression by CtBP1/2.

The positive regulators of *Prame* family genes remain largely unknown. Notably,

published RNA-seq data indicate that OBOX3/5, but not DUX, can activate *Pramel7* and

Pramef17 (*Pramel14*), both of which are upregulated by *Ctbp1/2* loss even in the absence

of DUX. In contrast, *Pramel16*, *Gm12794* (*Pramel19*) and *A430089l19Rik* (*Pramel39*)

require DUX for activation (Sakamoto et al., 2024). Because *Obox* family genes were

silenced in *Ctbp1/2/Dux* TKO ESCs, it is possible that another positive regulator, distinct

from OBOX proteins, is activated upon *Ctbp1/2* depletion or constitutively expressed in

WT ESCs. Expression of *Prame* family genes has been observed in various cancer and

germline cells (Kern et al., 2021), and their expression pattern resembles that of *Pou5f1*

(Oct4) (Bortvin et al., 2003), suggesting that an OCT4-centered transcriptional network

may function as a positive regulatory axis for *Prame* genes.

We further demonstrated that the overexpression of PRAMEL7 partially rescues the

transcriptional abnormalities observed in *Dux* KO ESCs, in which *Pramel7* expression

was markedly reduced. PRAMEL7 activates two distinct genetic programs: one involving

minor ZGA genes through *Dux* upregulation, and another encompassing maternally

expressed, major ZGA and post-ZGA genes. Previous work showed that PRAMEL7

promotes global DNA demethylation by targeting UHRF1 for proteasomal degradation

in ESCs (Graf et al., 2017). In addition, *Dux* expression is epigenetically repressed by

DNA methylation maintained by the ZBTB24-CDCA7-HELLS axis (Guo et al., 2025).

These findings suggest that PRAMEL7 regulates DUX-dependent transcriptional

cascades by promoting DNA demethylation at the *Dux* locus. Furthermore, we found that

maternally expressed and major ZGA genes regulated by PRAMEL7 are also activated

upon *Dnmt1* depletion, indicating that PRAMEL7 controls these genes through DNA

demethylation mediated by UHRF1 degradation. Because PRAMEL7 also promotes

degradation of NuRD complex at chromatin to sustain pluripotency in ESCs (Rupasinghe

et al., 2024), it will be important to explore whether NuRD degradation by PRAMEL7

contributes to the regulation of ZGA-associated transcriptional programs.

Although DUX is a master regulator of the transition from pluripotent ESCs to 2-cell-like

cells, our results demonstrated that PRAMEL7 overexpression rescues most of the defects

observed in *Dux* KO ESCs. *Prame* family genes are upregulated from the early phase of

2CLC emergence traced by the *Zscan4* and MERVL reporter system based on scRNA-

seq (Eckersley-Maslin et al., 2016; Rodriguez-Terrones et al., 2018), and MERVL-

positive 2CLCs are eliminated by *Dux* KO (De Iaco et al., 2017). Our data further showed

that *Pramel7* expression was significantly reduced upon *Dux* KO in WT ESCs but

remained unaffected in *Ctbp1/2* DKO ESCs, indicating that the loss of *Ctbp1/2* rescues

the *Dux* KO phenotypic consequences through *Pramel7* upregulation in ESCs. Taken

together, we propose that DUX orchestrates two distinct transcriptional programs in early

mouse embryos: (1) direct activation of minor ZGA genes (e.g., MERVL, *Zscan4c* and

Obox4), and (2) PRAMEL7-mediated transcriptional cascades. While *Dux* KO ESCs

exhibit a marked reduction in *Pramel7* expression, *Pramel7* remains normally expressed

in *Dux* KO embryos at the mid-2C stage (Guo et al., 2019). OBOX5, which is maternally

expressed, represents a strong candidate regulator for the DUX-independent activation of

Pramel7 in *Dux* KO embryos.

Our findings highlight fundamental differences between *in vitro* induction of 2CLCs and

in vivo regulation of ZGA, which are likely governed by maternal factors such as OBOX

and CtBP2. To better characterize these DUX-independent cascades, dual-reporter ESC

models (e.g., MERVL-tdTomato/*Gata1* or *Pramel7*-EGFP reporters) combined with

single-cell RNA-seq represent powerful approaches to dissect the molecular networks

underlying 2CLC heterogeneity induced by *Ctbp/1/2* loss.

References

Bortvin, A., Eggan, K., Skaletsky, H., Akutsu, H., Berry, D. L., Yanagimachi, R., Page, D. C. and Jaenisch, R. (2003). Incomplete reactivation of Oct4-related genes in mouse embryos cloned from somatic nuclei. *Development* **130**, 1673–1680.

Chang, T. C., Yang, Y., Yasue, H., Bharti, A. K., Retzel, E. F. and Liu, W. S. (2011). The expansion of the PRAME gene family in Eutheria. *PLoS One* **6**,.

Chen, Z. and Zhang, Y. (2019). Loss of DUX causes minor defects in zygotic genome activation and is compatible with mouse development. *Nat Genet* **51**, 947–951.

De Iaco, A., Planet, E., Coluccio, A., Verp, S., Duc, J. and Trono, D. (2017). DUX-family transcription factors regulate zygotic genome activation in placental mammals. *Nat Genet* **49**, 941–945.

Eckersley-Maslin, M. A., Svensson, V., Krueger, C., Stubbs, T. M., Giehr, P., Krueger, F., Miragaia, R. J., Kyriakopoulos, C., Berrens, R. V., Milagre, I., et al. (2016). MERVL/Zscan4 Network Activation Results in Transient Genome-wide DNA Demethylation of mESCs. *Cell Rep* **17**, 179–192.

Eckersley-maslin, M. A., Alda-catalinas, C. and Reik, W. (2018). Dynamics of the epigenetic landscape during the maternal- to-zygotic transition. *Nat Rev Mol Cell Biol* **19**, 436–450.

Eckersley-Maslin, M., Alda-Catalinas, C., Blotenburg, M., Kreibich, E., Krueger, C. and Reik, W. (2019). Dppa2 and Dppa4 directly regulate the Dux-driven zygotic transcriptional program. *Genes Dev* **33**, 194–208.

Escalier, D., Eloy, L. and Garchon, H. J. (2002). Sex-specific gene expression during meiotic prophase I: Xlr (X linked, lymphocyte regulated), not its male homologue Xmr (Xlr related, meiosis regulated), is expressed in mouse oocytes. *Biol Reprod* **67**, 1646–1652.

Ge, S. X., Son, E. W. and Yao, R. (2018). iDEP: An integrated web application for differential expression and pathway analysis of RNA-Seq data. *BMC Bioinformatics* **19**, 1–24.

Graf, U., Casanova, E. A., Wyck, S., Dalcher, D., Gatti, M., Vollenweider, E., Okoniewski, M. J., Weber, F. A., Patel, S. S., Schmid, M. W., et al. (2017). Pramel7 mediates ground-state pluripotency through proteasomal-epigenetic combined pathways. *Nat Cell Biol* **19**, 763–773.

Grow, E. J., Weaver, B. D., Smith, C. M., Guo, J., Stein, P., Shadle, S. C.,
Hendrickson, P. G., Johnson, N. E., Butterfield, R. J., Menafra, R., et al. (2021).
p53 convergently activates Dux/DUX4 in embryonic stem cells and in
facioscapulohumeral muscular dystrophy cell models. *Nat Genet* **53**, 1207–1220.

Guo, M., Zhang, Y., Zhou, J., Bi, Y., Xu, J., Xu, C., Kou, X., Zhao, Y., Li, Y., Tu, Z.,
et al. (2019). Precise temporal regulation of Dux is important for embryo
development. *Cell Res* **7**, 1–4.

Guo, Y., Kitano, T., Inoue, K., Murano, K., Hirose, M., Li, T. D., Sakashita, A., Ishizu,
H., Ogonuki, N., Matoba, S., et al. (2024). Obox4 promotes zygotic genome
activation upon loss of dux. *Elife* **13**, 1–25.

Guo, D., Du, Z., Liu, Y., Lin, M., Lu, Y., Hardikar, S., Xue, Y., Zhang, J., Chen, T.
and Dan, J. (2025). The ZBTB24-CDCA7-HELLS axis suppresses the totipotent
2C-like reprogramming by maintaining Dux methylation and repression. *Nucleic*
Acids Res **53**,.

Hendrickson, P. G., Doráis, J. A., Grow, E. J., Whiddon, J. L., Lim, J.-W., Wike, C.
L., Weaver, B. D., Pflueger, C., Emery, B. R., Wilcox, A. L., et al. (2017).
Conserved roles of mouse DUX and human DUX4 in activating cleavage-stage
genes and MERVL/HERVL retrotransposons. *Nat Genet* **49**, 925–934.

Hu, Z., Tan, D. E. K., Chia, G., Tan, H., Leong, H. F., Chen, B. J., Lau, M. S., Tan,
K. Y. S., Bi, X., Yang, D., et al. (2020). Maternal factor NELFA drives a 2C-like
state in mouse embryonic stem cells. *Nat Cell Biol* **22**, 175–186.

Hulsen, T., de Vlieg, J. and Alkema, W. (2008). BioVenn - A web application for the
comparison and visualization of biological lists using area-proportional Venn
diagrams. *BMC Genomics* **9**, 1–6.

Ishiuchi, T., Enriquez-Gasca, R., Mizutani, E., Bošković, A., Ziegler-Birling, C.,
Rodriguez-Terrones, D., Wakayama, T., Vaquerizas, J. M. and Torres-Padilla,
M.-E. (2015). Early embryonic-like cells are induced by downregulating
replication-dependent chromatin assembly. *Nat Struct Mol Biol* **22**, 662–71.

Ji, S., Chen, F., Stein, P., Wang, J., Zhou, Z., Wang, L., Zhao, Q., Lin, Z., Liu, B., Xu,
K., et al. (2023). OBOX regulates mouse zygotic genome activation and early
development. *Nature* **620**, 1047–1053.

Kern, C. H., Yang, M. and Liu, W. S. (2021). The PRAME family of cancer testis
antigens is essential for germline development and gametogenesis. *Biol Reprod* **105**,

290–304.

Kim, T. W., Kang, B. H., Jang, H., Kwak, S., Shin, J., Kim, H., Lee, S. E., Lee, S. M.,
Lee, J. H., Kim, J. H., et al. (2015). Ctbp2 Modulates NuRD-Mediated
Deacetylation of H3K27 and Facilitates PRC2-Mediated H3K27me3 in Active
Embryonic Stem Cell Genes during Exit from Pluripotency. *Stem Cells* **33**, 2442–
2455.

Kreibich, E., Blotenburg, M., Alda-Catalinas, C., Reik, W., Krueger, C. and
Eckersley-Maslin, M. (2019). Dppa2 and Dppa4 directly regulate the Dux-driven
zygotic transcriptional program. *Genes Dev* **33**, 194–208.

Macfarlan, T. S., Gifford, W. D., Driscoll, S., Lettieri, K., Rowe, H. M., Bonanomi,
D., Firth, A., Singer, O., Trono, D. and Pfaff, S. L. (2012). Embryonic stem cell
potency fluctuates with endogenous retrovirus activity. *Nature* **487**, 57–63.

McLean, C. Y., Bristor, D., Hiller, M., Clarke, S. L., Schaar, B. T., Lowe, C. B.,
Wenger, A. M. and Bejerano, G. (2010). GREAT improves functional
interpretation of cis-regulatory regions. *Nat Biotechnol* **28**, 495–501.

Percharde, M., Lin, C., Yin, Y., Huang, B., Shen, X., Ramalho-santos, M., Percharde,
M., Lin, C., Yin, Y., Guan, J., et al. (2018). Article A LINE1-Nucleolin Partnership
Regulates Early Development and ESC Identity Article A LINE1-Nucleolin
Partnership Regulates Early Development and ESC Identity. *Cell* **174**, 1–15.

Rodriguez-Terrones, D., Gaume, X., Ishiuchi, T., Weiss, A., Kopp, A., Kruse, K.,
Penning, A., Vaquerizas, J. M., Brino, L. and Torres-Padilla, M. E. (2018). A
molecular roadmap for the emergence of early-embryonic-like cells in culture. *Nat*
Genet **50**, 106–119.

Rupasinghe, M., Bersaglieri, C., Leslie Pedrioli, D. M., Pedrioli, P. G. A., Panatta,
M., Hottiger, M. O., Cinelli, P. and Santoro, R. (2024). PRAMEL7 and CUL2
decrease NuRD stability to establish ground-state pluripotency. *EMBO Rep* **25**,
1453–1468.

Sakamoto, M., Ito, A., Wakayama, S., Sasaki, H., Wakayama, T. and Ishiuchi, T.
(2024). Detection of newly synthesized RNA reveals transcriptional reprogramming
during ZGA and a role of Obox3 in totipotency acquisition. *Cell Rep* **43**, 114118.

Shin, Y. H., Choi, Y., Erdin, S. U., Yatsenko, S. A., Kloc, M., Yang, F., Wang, P. J.,
Meistrich, M. L. and Rajkovic, A. (2010). Hormad1 mutation disrupts
synaptonemal complex formation, recombination, and chromosome segregation in

mammalian meiosis. *PLoS Genet* **6**,.

Stankiewicz, T. R., Gray, J. J., Winter, A. N. and Linseman, D. A. (2014). C-terminal

binding proteins: Central players in development and disease. *Biomol Concepts* **5**,

489–511.

Whiddon, J. L., Langford, A. T., Wong, C. J., Zhong, J. W. and Tapscott, S. J. (2017).

Conservation and innovation in the DUX4-family gene network. *Nat Genet* **49**, 935–

940.

Wu, K., Liu, H., Wang, Y., He, J., Xu, S., Chen, Y., Kuang, J., Liu, J., Guo, L., Li, D.,

et al. (2020). SETDB1-Mediated Cell Fate Transition between 2C-Like and

Pluripotent States. *Cell Rep* **30**, 25-36.e6.

Yamamoto, M., Suwa, Y., Sugiyama, K., Okashita, N., Kawaguchi, M., Tani, N.,

Matsubara, K., Nakamura, A. and Seki, Y. (2020). PRDM14-CtBP1/2-PRC2

complex regulates transcriptional repression during transition from primed to naïve

pluripotency. *J Cell Sci* **133**, 1–13.

Zhang, B., Zheng, H., Huang, B., Li, W., Xiang, Y., Peng, X., Ming, J., Wu, X., Zhang,

Y., Xu, Q., et al. (2016). Allelic reprogramming of the histone modification

H3K4me3 in early mammalian development. *Nature* **537**, 553–557.

Zhang, T., Zhang, Z., Dong, Q., Xiong, J. and Zhu, B. (2020). Histone H3K27

acetylation is dispensable for enhancer activity in mouse embryonic stem cells.

Genome Biol **21**, 1–7.

Author Contributions

Y.S. conceived the study and experimental design, performed experiments and wrote the manuscript. K.S., K.Y. and N.H. performed most experiments. R.M. M.M., A.S. and Y.M performed some experiments. K.S., K.Y. and N.H. contributed equally to this work.

Acknowledgements

We thank all members of Seki laboratory for stimulating discussions. This study was supported by JSPS KAKENHI Grant Numbers JP18H02422, JP20H05375, JP22H04682 and a Special Research Grant for Individual Researchers from Kwansei Gakuin University.

Conflict of interest

The authors declare no competing interests.

Data availability

RNA-Seq data has been deposited in Gene Expression Omnibus (GSE253944)