**The disruption of CtBP regulates DUX-dependent and -independent genetic**

**program for the 2-cell-like state in murine embryonic stem cells**

Kota Sugiyama[1], Kazuma Yoshioka[1], Nao Hayakawa[1], Miyu Marutani, Ryo Masuda,

Selma Alamanda Abadi, Maki Ichisakino and Yoshiyuki Seki

Department of Biomedical Sciences, School of Biological and Environmental Sciences,

Kwansei Gakuin University, Hyogo, Japan

[1]Co-first author

*Correspondence: yseki@kwansei.ac.jp

## Abstract

After fertilization, maternally deposited mRNA is cleared, and *de novo* mRNA is

transcribed from the zygotic genome through zygotic genome activation (ZGA), which

is called maternal-to-zygotic transition (MZT) occurring in the mouse at 2-cell (2C)

stage. 2C-like cells (2CLCs) marked by MERVL expression and transcriptionally

similar to 2C embryos spontaneously emerge from mouse embryonic stem cells

(mESCs). Although the emergence of 2CLCs completely depends on DUX function, a

recent knockout study clearly showed that DUX is dispensable for mouse embryos,

suggesting that DUX-independent molecular pathways are not recapitulated in 2CLCs.

We present here that the disruption of C-terminal binding protein 1/2 (*Ctbp1/2*)

activates DUX-dependent and -independent molecular pathway associated with the

development of early mouse embryos mediated by the upregulation of Preferentially

expressed antigen of melanoma family-like 7 (PRAMEL7). Furthermore, the

abnormality of the gene expression profile caused by *Dux* KO is partially rescued by the

overexpression of PRAMEL7 in mESCs. Our study provides new insights into the

DUX-independent molecular pathway for developing early mouse embryos.

**Introduction**

After fertilization, maternal factors direct development and trigger zygotic genome activation (ZGA) at the around late one-cell stage in mice, called maternal-to-zygotic transition (MZT)(Eckersley-Maslin et al., 2018). The early-cleavage stage embryos acquire cellular totipotency to generate both embryonic proper and extraembryonic tissues during MZT. Although totipotency was thought to be exclusive to the early-cleavage stage embryos, a part of the molecular network is recapitulated in the totipotent-like cells (2-cell-like cells; 2CLCs) marked by the expression of endogenous retrovirus MERVL, which are an endogenous subpopulation of mouse embryonic stem cells (ESCs). (Macfarlan et al., 2012a) (Rodriguez-Terrones et al., 2018).

The transition from pluripotent stem cells into 2CLCs is a useful *in vitro* system for identifying the gene regulatory network for ZGA, and several genes that control the emergence of 2CLCs from pluripotent cells have been identified (Eckersley-Maslin et al., 2019; Hu et al., 2020; Ishiuchi et al., 2015; Percharde et al., 2018; Rodriguez-Terrones et al., 2018). The transcription factor DUX (DUX4 in humans) is a critical factor for regulating the transition from pluripotent stem cells into 2CLCs and the expression of ZGA genes (De Iaco et al., 2017; Hendrickson et al., 2017; Whiddon et al., 2017). Ectopic expression of DUX enhances the population of 2CLCs, whereas the *Dux* disruption eliminates the emergence of spontaneous 2CLCs during mESC culture. Several 2CLC regulatory genes have been identified, but they are either direct activators (DPPA2/4, p53 and NELFA) or direct repressors (CAF-1, LINE-1 and SETDB1) of *Dux* expression (Eckersley-Maslin et al., 2019; Grow et al., 2021; Ishiuchi et al., 2015b; Percharde et al.,

2018; Wu et al., 2020). However, both *Dux* zygotic and maternal knockout embryos can

survive full-term (Chen and Zhang, 2019; Guo et al., 2019), suggesting that DUX-

independent molecular pathways not recapitulated in 2CLCs are critical for ZGA and the

early development of mice.

In this study, we show that disruption of Ctbp1/2 (C-terminal binding protein 1/2)

(Stankiewicz et al., 2014) activates two distinct sets of genes in ESCs: the first set is

transiently upregulated in a DUX-dependent manner during minor ZGA, and the second

set contains maternally expressed genes and the upregulated genes in a DUX-independent

manner during major ZGA. Preferentially Expressed Antigen in Melanoma (PRAME) 7

is directly upregulated by the loss of *Ctbp1/2* without DUX function, and upregulated

PRAMEL7 activates minor ZGA genes in a DUX-dependent manner and maternally

expressed genes and major ZGA in DUX-independent manner. These data revealed that

the disruption of *Ctbp1/2* recapitulates a part of DUX-independent ZGA cascades in ESCs,

and *Ctbp1/2* DKO ESCs are useful *in vitro* systems to investigate DUX-independent

molecular cascades ensuring ZGA in mice.

**Material and Methods**

**Cell Culture**

E14tg2a ESCs (RIKEN Cell Bank) were cultured in Glasgow minimum essential medium (GMEM; Wako) containing 10% fetal calf serum (FCS; Invitrogen), 1 mM glutamine (Wako), nonessential amino acids (Wako) and 500 μM Monothioglycerol Solution (Wako). The culture was supplemented with 1000 U/ml of leukaemia inhibitory factor (LIF) (Wako) without feeder cells. ESCs were cultured in N2B27 medium with 2i (PD0325901, 0.4 μM: WAKO, CHIR99021, 3 μM: Sigma-Aldrich).

**Generation of knockout ESCs using a CRISPR-Cas9 system**

Guide RNAs for *Ctbp1/2*, *Dux* and *Pramel7* were generated in pX330-U6- Chimeric_BB-CBh-hSpCas9 (Addgene plasmid #42230; deposited by Feng Zhang; Cong et al., 2013). The guide RNA sequences are listed in Supplementary Table 1. The pX330 was co-transfected with pCAG-EGFP into E14tg2a with MERVL::TurboGFP (Addgene plasmid #69072; deposited by Maria-Elena Torres-Padilla; (Ishiuchi et al., 2015a). The GFP-positive fraction was sorted by Cell Sorter (No. SH800; Sony Biotechnology, Inc., Tokyo, Japan). Genomic DNA deletions were sequenced (Supplementary Figure 1).

**qRT-PCR and RNA-Seq**

Total mRNA was purified with a PureLink™ RNA Mini Kit (Ambion, Inc.). For quantitative RT-PCR, cDNA was synthesized by ReverTra Ace® qPCR RT master Mix (Toyobo), and then analyzed by qPCR with FastStart SYBR Green Master Mix (Roche)

using a specific primer set for each gene (Supplementary Table 1). RNA-seq libraries

were generated using Collibri$^{TM}$ Stranded RNA library prep for illumina$^{TM}$ Systems

(ThermoFisher Scientific) and sequenced using HiSeq X$^{TM}$ Ten (Illumina) (150-bp

paired-ends reads).

**Data analysis of RNA-Seq**

Total mRNA was purified with a PureLink$^{TM}$ RNA Mini Kit (Ambion, Inc.). RNA-seq

libraries were generated using Collibri$^{TM}$ Stranded RNA library prep for illumina$^{TM}$

Systems (ThermoFisher Scientific) and sequenced using HiSeq X$^{TM}$ Ten (Illumina).

Reads were aligned by Hisat2 to the Mus musculus genome (mm10). Raw read counts of

genes were calculated by R package FeatureCounts using mm10 KnownGenes. GTF from

UCSC for annotation. Normalized count data, principal component and K-means

clustering analysis were exported by the interactive Expression and Pathway analysis

(iDEP) webserver using the count data or normalized expression data of early mouse

embryos (Ge et al., 2018). Gene set enrichment analysis (GSEA) was used to determine

whether upregulated genes by *Ctbp1/2* knockouts were enriched for the upregulated genes

in 2-cell embryos or 2CLCs as described previously. Venn diagrams showing overlapping

genes were constructed by BioVenn, a web application (Hulsen et al., 2008).

**Bisulfite sequencing of specific genes and RRBS**

Genomic DNA was purified using Wizard® SV Genomic DNA Purification System

(Promega). For bisulfite sequencing of specific genes, the unmethylated cytosine of the

genomic DNA was converted to uracil using EpiTect Bisulfite Kit (QIAGEN) according

to the manufacturer's instructions. Bisulfited DNA was amplified by the region-specific

primer, followed by the sequencing. According to the manufacturer's instructions, DNA

libraries for RRBS analysis were prepared using the Premium RRBS kit V2 and

sequenced using HiSeq X$^{TM}$ Ten (Illumina) (150-bp paired-ends reads). RRBS reads

were trimmed using trim galore (www.bioinformatics.babraham.ac.uk/projects/

trim_galore/) with the RRBS option invoked. Bismark was used to align reads to UCSC

version mm10 of the mouse genome. CpG methylation was extracted using

bismark_methylation_extractor and displayed by Broad Institute's Integrative Genomic

Viewer (IGV) (Robinson et al., 2011).

**Immunofluorescence analysis**

Cells were fixed by 4% paraformaldehyde at 4℃ for 20 min, followed by incubation with

PBS containing 0.5% Triton-X at 4℃ for 5 min. Permeabilized cells were incubated with

a primary antibody solution containing anti-GFP (Nacarai Tesque, 04404-84, 1:500) with

anti-ZSCAN4 (Merk Millipore, AB4340, 1:1000) or anti-MuERVL-Gag (Epigentek, A-

2801-050, 1:250) or anti-OCT4 (Santa Cruz, Sc-8628, 1:500) or anti-CtBP2 (BE

Biosciences, 612044, 1:500) followed by secondary antibody solution containing anti-

Rat IgG Dylight$^{TM}$ 488 antibody (Rockland, 612-141-120, 1:500) or anti-Rabbit IgG

Dylight$^{TM}$ 549 antibody (KPL, 042-04-15-06, 1:500) or anti-Mouse IgG Dylight® 550

antibody (Abcam, ab96876, 1:500) or anti-Goat IgG Dylight$^{TM}$ 549 antibody (Rockland,

605-742-002, 1:500) with 1 µg/ml of DAPI solution (DOJINDO, 340-07971).

**Luciferase assay**

The enhancer candidate region of *Pramel7* or *Dux* was cloned into pGL4.23, driving the

*luciferase* gene. Wild-type or *Ctbp1/2* DKO ESCs were transfected with the reported

constructs with *Ctbp2* or *Pramel7* expression vector using HilyMax (DOJINDO). Cells

were harvested 36 hours after transfection, and the luciferase activities were measured

using the Dual-luciferase Reporter Assay System (Promega).

**Western blotting analysis**

Cells were lysed by boiling SDS sample buffer (WAKO). Before lysed proteins were

applied to polyacrylamide-SDS gels, 2-mercaptoethanol was added to denature the

proteins. The lysed proteins were separated on polyacrylamide-SDS gels, blotted on a

polyvinylidene fluoride membrane, and probed using the following primary antibodies:

anti-UHRF1 (MBL, D289-3, 1:1000) or anti-TUBLIN (SIGMA, T5168, 1:5000).

Following the primary antibody reaction, the membrane were incubated with horseradish

peroxidase (HRP)-coupled secondary antibodies. All antibodies were detected with

Immunostar LD (WAKO).

**Phylogenetic tree inference**

Amino acid sequences of PRAME family proteins were obtained from Ensemble release

112. Multiple-sequence alignment was performed using the ClustalW algorithm with the

158    MEGA software, version 11 with default parameters. Molecular phylogenetic trees were

159    inferred with the maximum-likelihood method using MEGA software, version 11 with

160    default parameters.

**Results**

***Ctbp1/2 Deficiency enhances the transition from pluripotent stem cells to 2-cell-like***

***Cells***

We have previously shown that the CtBP1/2 are required to transition from formative to

naïve pluripotency in mouse embryonic stem cells (mESCs) (Yamamoto et al., 2020). To

elucidate the molecular functions of CtBP1/2 in the transition from formative to naïve

pluripotency in mESCs, we compared the gene expression profile between wild-type

(WT) and *Ctbp1/2* double knockout (DKO) ESCs using RNA-sequencing (RNA-Seq).

MA-plot comparing WT ESCs with *Ctbp1/2* DKO ESCs showed that the 2-cell (2C)-

restricted genes, including *Dux*, *Zscan4* and *Zfp352*, as well as three repeat elements

MERVL, MT2 and GSAT_Mm are significantly upregulated by *Ctbp1/2* DKO (Figure

1A, B). Gene set enrichment analysis (GSEA) using a 2-cell embryo and 2-cell-like cell

(2CLC)-enriched genes (404) (Macfarlan et al., 2012) showed that 2C-restricted genes

were significantly enriched in the upregulated genes by the loss of *Ctbp1/*2 (Figure 1C).

Next, to monitor the dynamics of the transition from pluripotent stem cells into 2C-like

cells, we established *Ctbp1/2* DKO ESCs carrying the turboGFP reporter under the

control of MERVL long terminal repeat (*Mervl*-EGFP) (Ishiuchi et al., 2015a). The GFP-

positive fraction was about 0.1 % in WT ESCs, while remarkedly, *Ctbp1/2* DKO induced

a more than 50-fold increase of 2CLC (7.37%). To confirm the critical markers of 2CLCs,

we performed immunofluorescence analysis with anti-ZSCAN4, Mervl-GAG and OCT4

antibodies (Figure 1E). *Mervl*-EGFP-positive cells emerged by *Ctbp1/2* DKO expressed

ZSCAN4 and Mervl-GAG and lacked OCT4. Our data suggested that the downregulation

of *Ctbp1/2* might be involved in the zygotic genome activation *in vivo*. Consistent with

this idea, maternal mRNA of CtBP2 was reduced between the 2-cell to 4-cell stage based

on the single-cell RNA-Seq data of mouse early embryo (Gerlinger et al., 2014) (Figure

1F). However, the expression level of CtBP2 was maintained in spontaneous 2CLCs

(Figure 1G), suggesting that the molecular pathway triggered by the loss of *Ctbp1/2* in

the 2-cell embryo is stalled in spontaneous 2CLCs.

**Figure 1**

**Figure 1. The disruption of *Ctbp1/2* expands the 2-cell-like cells**

(A), (B) MA-plot showing mean expression and fold change per gene or repeat elements in *Ctbp1/2* DKO versus WT ESCs. Cut-off values were set to $0.05 \geq P$ and over 2-fold changes. Red and blue dots indicate upregulated genes and downregulated genes, respectively. (C) Gene set enrichment analysis (GSEA) using ZGA and 2-cell enriched genes defined in (GSE33923) (Macfarlan et al., 2012)to compare *Ctbp1/2* DKO ESCs

with WT ESCs. (D) FACS analysis of MERVL::TurboGFP in WT ESCs and *Ctbp1/2* DKO ESCs. (E) MERVL::TurboGFP-positive cells were marked by ZSCAN4 and Mervl-gag proteins and mutually exclusive with OCT4-positive cells. Scale bar: 10 μm. (F) mRNA levels of *Ctbp2* in early mouse development based on the published single-cell RNA-Seq data (Gerlinger et al., 2014). (G) Immunofluorescence for CtBP2 (red) in WT ESCs with MERVL::TurboGFP (green). Scale bar: 10 μm.

### *2C program induced by Ctbp1/2 Deficiency partially depends on DUX*

DUX, the homeobox transcription factor, is essential and sufficient for the emergence of spontaneous 2CLCs from ESCs (De Iaco et al., 2017; Whiddon et al., 2017). To determine the dependency of the 2C program driven by *Ctbp1/2* deficiency on DUX, we deleted the *Dux* cluster in WT and *Ctbp1/2* DKO ESCs (Supplementary Figure 1C). Firstly, we compared the population of MERVL-EGFP-positive cells between *Ctbp1/2* DKO and *Ctbp1/2/Dux* TKO ESCs. Although *Dux* KO lost most of the MERVL-EGFP-strong positive cells in *Ctbp1/2* DKO ESCs, MERVL-EGFP-weak positive cells were found in *Ctbp1/2/Dux* TKO ESCs (Figure 2A). We have analyzed the expression levels of 2C genes in WT, *Dux* KO, *Ctbp1/2* DKO and TKO ESCs using qRT-PCR. Upregulation of Preferentially Expressed Antigen in Melanoma-like 7 (*Pramel7*) (Graf et al., 2017), which is upregulated at the 2C stage of the mouse embryo (Guo et al., 2019), by the loss of *Ctbp1/2* was unaffected by *Dux* KO, while *Dux* KO downregulated the expression of *Zscan4c* in *Ctbp1/2* DKO ESCs (Figure 2B). Next, to determine the effect of the loss of *Ctbp1/2* and *Dux* in the transcriptional activity of MERVL-LTR, we performed a luciferase assay using the reporter driven by MERVL-LTR. *Ctbp1/2* deficiency significantly upregulated the reporter driven by MERVL-LTR, and the mutation of the

DUX-recognition sequence impaired this activation, but the activity was substantially

higher than those in WT ESCs (Supplementary Figure 2A). Furthermore, the activation

of MERVL-Luciferase (Luc) by the loss of *Ctbp1/2* was significantly suppressed by the

deletion of *Dux*. However, the reporter activity of MERVL-Luc and MERVL-Luc with

the mutation of the DUX-recognition sequence (DUXm) in *Ctbp1/2/Dux* TKO ESCs was

considerably higher than those in *Dux* KO ESCs. These findings suggest that the loss of

*Ctbp1/2* activates the reporter activity of MERVL-Luc without the DUX function. qRT-

PCR analysis showed that the activation of *Zscan4c* is more strongly dependent on DUX

than the activation of *Mervl* and *Naalad2* by the loss of *Ctbp1/2* (Figure 2B). Published

ChIP-Seq data (Hendrickson et al., 2017) of DUX showed that DUX is localized at two

distinct regions around the *Zscan4c* locus (Supplementary Figure 2B). No MERVL

sequence was found at the major peak of DUX-binding, and the weak peak of DUX-

binding was localized in the MERVL sequence. In contrast, all DUX-binding peaks

around the *Naalad2* locus are co-localized with the MERVL sequence. These findings

suggested that *Zscan4c* is regulated by the DUX-binding elements with or without

MERVL, while only DUX-binding elements with MERVL regulate *Naalad2*. To confirm

these possibilities, we examined the reporter activity driven by DUX-binding elements

without MERVL and with MERVL. Reporter activity driven by the DUX-binding element

without MERVL upstream of *Zscan4c* was upregulated by the loss of *Ctbp1/2*, and this

activation was cancelled entirely by the deletion of *Dux* (Supplementary Figure 2C). The

reporter activity driven by the DUX-binding element with MERVL at *Zscan4c* and

*Naalad2* was also upregulated by the *Ctbp1/2* disruption, and this upregulation was

significantly but not entirely repressed by the loss of *Dux*. Consistent with the results of

MERVL-Luc, these reporter activities in *Ctbp1/2/Dux* TKO ESCs were higher than those

in *Dux* KO ESCs.

Next, to identify the 2C genes whose expression is upregulated by *Ctbp1/2* DKO without

DUX function, we performed RNA-Seq using WT, *Dux* KO, *Dppa2* KO, *Ctbp1/2* DKO,

*Ctbp1/2/Dux* TKO and *Ctbp1/2/Dppa2* TKO ESCs. DPPA2/DPPA4 complex is the

upstream element for the activation of *Dux* in ESCs (Eckersley-Maslin et al., 2019). To

gain insight into the dependency of DUX on the expression variation by the loss of

*Ctbp1/2*, we created a scatter plot displaying the correlation of expression variation by

the *Ctbp1/2* DKO between WT and *Dux* KO ESCs (Figure 2C). The distribution of the

scatter plots showed that the dependency of DUX on the transcriptional regulation by

*Ctbp1/2* is relatively weak, except for some genes, including *Zscan4*, *Sp110* and *Usp17*.

The minimal effect of DUX on CtBP1/2-mediated transcriptional regulation was

confirmed by gene set enrichment analysis (GSEA) comparing *Ctbp1/2/Dux* TKO and

*Dux* KO ESCs using the upregulated genes in *Ctbp1/2* DKO (Figure 2D).

We then performed K-means clustering analysis (K=5) using the top 2000 differentially

expressed genes to separate groups based on the expression alteration at each KO

condition (Figure 2E). DUX and DPPA2-dependent upregulated genes by the loss of

*Ctbp1/2,* including *Zscan4* and *Pramef25* (*Pramel16*)*,* were enriched in Cluster 2. Cluster

3, which was enriched with a group of genes whose expression is reduced by *Dux* KO

and this reduction is canceled by *Ctbp1/2* DKO, contained KRAB-ZFP cluster

(*Zfp900/981/600/534/458*) (Wolf et al., 2020). Cluster 4, in which the genes are

upregulated by *Ctbp1/2* DKO, *Ctbp1/2/Dux* TKO and *Ctbp1/2/Dppa2* TKO, contained

the genes upregulated by major ZGA, including *Pramel6/7*, *Pramef17*, *Cphx1/2/3*, *Gata1*

and *Crxos* (Supplementary Table 2). Interestingly, Cluster 5, enriched with a group of

genes whose expression was decreased by *Dppa2* KO and increased by *Ctbp1/2* DKO,

was enriched in GO regarding reproduction, including *Tex11*, *Dazl* and *Hormad1*. To

verify whether the gene expression changes in ESCs induced by *Ctbp1/2* DKO reflect the

gene expression dynamics in early embryos, we analyzed the expression changes of genes

belonging to Cluster 2 (DUX-dependent genes), Cluster 4 (DUX-independent genes) and

Cluster 5 (DPPA2-dependent genes) in early mouse embryo using public RNA-Seq data

(Gerlinger et al., 2014) (Figure 2F and Figure 3A). We analyzed the expression dynamics

of Cluster 2 genes in early embryos by removing genes with low expression in early

embryos (FPKM ≤ 1). Most genes were transiently upregulated in 2-4 cell stages. In

contrast, K-means clustering (K=6) analysis of cluster 4 showed that they were classified

into two major groups: maternal factors whose expression decreased after fertilization

(Cluster4a and b) and genes whose expression was activated after ZGA (Cluster 4c-f).

Next, we focused on the loss of *Ctbp1/2, which* appears to reverse the gene expression

alteration caused by *Dux* KO, as shown in Clusters 3 and 4 of the K-means clustering

analysis. A comparison of the expression alteration by *Dux* KO in WT ESCs versus those

by the loss of *Ctbp1/2* in *Dux* KO ESCs, as demonstrated by scatter plots, revealed a

negative correlation (Figure 2G). Furthermore, the Venn diagram overlapping the

downregulated genes by *Dux* KO and the upregulated by *Ctbp1/2* DKO in *Dux* KO

showed that most of the down-regulated genes by *Dux* KO are upregulated by *Ctbp1/2*

DKO in *Dux* KO (Figure 2H). These results suggest that the loss of *Ctbp1/2* activates the

maternal genes and major ZGA genes located downstream of DUX, either directly or

indirectly.

**Figure 2**

**Figure 2. *Ctbp1/2* disruption partially rescues the *Dux* KO phenotype in ESCs**

(A) FACS analysis of MERVL::TurboGFP in WT, *Ctbp1/2* DKO and *Ctbp1/2*/*Dux* TKO

ESCs. (B) RT-qPCR analysis of *Dux*, *Zscan4c* and *Pramel7* in WT, *Dux* KO, *Ctbp1/2*

DKO and Ctbp1/2/*Dux* TKO ESCs. The ΔCt of each gene was calculated by subtracting

the average Ct of each gene from that of *Gapdh*. Means of ΔCt were calculated from three

biological triplicates. ***$P < 0.001$, **$P < 0.005$, *$P < 0.05$ (Tukey-Kramer multiple

comparison test). (C) Scatter plot analysis of the correlation between the expression

changes by *Ctbp1/2* DKO in WT ESCs and *Ctbp1/2* DKO in *Dux* KO ESCs. (D) Gene

Set Enrichment Analysis (GSEA) of upregulated genes by *Ctbp1/2* DKO in *Ctbp1/2/Dux*

TKO and *Dux* KO ESCs. (E) Scatter plot analysis of the correlation between the

expression changes by *Dux* KO in WT ESCs and *Ctbp1/2* DKO in *Dux* KO ESCs. (F)

Gene Set Enrichment Analysis (GSEA) of the downregulated genes by *Dux* KO in

*Ctbp1/2/Dux* TKO and *Dux* KO ESCs. (G) K-means clustering analysis (K=5, N=2000)

using RNA-Seq data of WT, *Dux* KO, *Dppa2* KO, *Ctbp1/2* DKO, *Ctbp1/2/Dux* TKO and

*Ctbp1/2/Dppa2* TKO ESCs. K-means clustering analysis of the genes belonging to

Cluster 2 and Cluster 4 using published single-cell RNA-Seq data of early mouse embryos

(Gerlinger et al., 2014). (H) Venn diagram illustrating the overlap between the down-

regulated genes by *Dux* KO and the upregulated genes by *Ctbp1/2* DKO in *Dux* KO. (I)

Volcano plot showing differentially accumulated repetitive elements between WT ESCs

and *Dux* KO ESCs, WT ESCs and *Ctbp1/2* DKO ESCs, *Dux* KO ESCs and *Ctbp1/2/Dux*

TKO ESCs.

***Dppa2 KO phenotype is partially rescued by the disruption of Ctbp1/2***

K-means clustering analysis of transcriptome data using WT, *Ctbp1/2* DKO, *Dppa2* KO

and *Ctbp1/2/Dppa2* TKO ESCs revealed that down-regulated genes by *Dppa2* KO tend

to be upregulated by *Ctbp1/2* DKO (Cluster 5) (Figure 2C). These genes were categorized

into two groups based on the gene expression dynamics in early embryos: maternally

expressed genes (Cluster 1, 2) and the genes activated after major ZGA (Cluster 3-5)

(Figure 3A). Comparing the expression changes caused by *Dppa2* KO in WT ESCs with

those caused by *Ctbp1/2* DKO in *Dppa2* KO ESCs showed a clear negative correlation

(Figure 3B). The Venn diagram illustrating the overlap between the genes downregulated

by *Dppa2* KO and those upregulated by *Ctbp1/2* DKO in *Dppa2* KO showed that

approximately half of the genes downregulated by *Dppa2* KO were upregulated by

*Ctbp1/2* DKO in *Dppa2* KO (Figure 3C). It was previously reported that DPPA2

safeguard against *de novo* DNA methylation through the establishment of H3K4me3

enrichment in ESCs cultured in 2i plus LIF condition (2iL) (Gretarsson and Hackett,

2020)(Eckersley-maslin et al., 2020). We compared the DNA methylation status at around

the CpG island of *Dazl* and promoter regions of LINE1 in WT, *Dppa2* KO, *Ctbp1/2* DKO,

*Ctbp1/2/Dppa2* TKO ESCs with serum LIF (SL) and 2iL using bisulfite sequencing

analysis. DNA methylation levels were consistently elevated by *Dppa2* KO in WT ESCs

but not in *Ctbp1/2* DKO, and this tendency was more pronounced in ESCs with 2iL than

SL (Figure 3D). We compared the global DNA methylation levels in WT, *Dppa2* KO,

*Ctbp1/2* DKO and *Ctbp1/2/Dppa2* TKO ESCs under 2iL using reduced-representation

bisulfite sequencing (RRBS). Because it has been reported that DPPA2 preferentially

binds to CpG island and protects aberrant *de novo* DNA methylation (Gretarsson and

Hackett, 2020), we extracted the promoter region with CpG island in which the DNA

methylation levels were elevated by *Dppa2* KO in ESCs (Figure 3E). Interestingly, the

DNA methylation levels at almost all these promoter regions, including meiosis-

associated genes, *Dazl*, *Ddx4* and *Sycp1/2/3*, were reduced by the disruption of *Ctbp1/2*

in *Dppa2* KO ESCs. (Figure 3E, F). To determine the molecular mechanism that *Ctbp1/2*

disruption impaired the abnormal elevation of DNA methylation caused by *Dppa2* KO in

ESCs, we compared the mRNA expression of the regulators of DNA methylation and

DNA methyltransferases by RT-qPCR analysis. *Pramel7,* which is involved in degrading

UHRF1 protein (Graf et al., 2017), was significantly repressed in *Dppa2* KO ESCs, and

this repression was entirely cancelled by *Ctbp1/2* disruption (Figure 3G). Consistent with

the expression of *Pramel7*, the protein level of UHRF1 was elevated by *Dppa2* KO, and

this up-regulation was diminished by *Ctbp1/2* disruption (Figure 3H). RRBS and

published ChIP-Seq data of DPPA2 (Engelen et al., 2015) showed that DNA methylation

level was elevated upstream region of *Pramel7* associated with DPPA2-binding (Figure

3I), suggesting that DPPA2 maintain DNA hypomethylation through the activation of

*Pramel7* expression.

# Figure 3

**Figure 3. The loss of *Ctbp1/2* rescues abnormal DNA methylation by *Dppa2* KO.**

(A) K-means clustering analysis of the genes of Cluster 5 (Figure 2C) using published single-cell RNA-Seq data of early mouse embryos (Gerlinger et al., 2014). (B) A scatter plot of RNA-Seq data represents genes upregulated or downregulated by *Dppa2* KO in WT ESCs and *Ctbp1/2* DKO in *Dppa2* KO ESCs. (C) The Venn diagram shows the overlapping between the down-regulated genes by *Dppa2* KO in WT ESCs and the upregulated genes by *Ctbp1/2* DKO in *Dppa2* KO ESCs. (D) Bisulfite sequencing of the promoter region of *Dazl* and *LINE1* in WT, *Dppa2* KO, *Ctbp1/2* DKO and *Ctbp1/2/Dppa2* TKO ESCs under Serum LIF and 2i LIF. \*\*\**P* < 0.001, \*\**P* < 0.005, \**P* < 0.05 (Mann-Whitney U-test). (E) Venn diagram showing the overlapping between the region with CpG island in which DNA methylation is elevated by *Dppa2* KO in ESCs and the region with CpG island in which Ctbp1/2 DKO reduces DNA methylation in *Dppa2* KO ESCs. (F) Gene track representing the methylation levels derived from RRBS at *Dazl*, *Ddx4*, *Ric3* and *Twist2* in WT, *Dppa2* KO, *Ctbp1/2* DKO and *Ctbp1/2/Dppa2* TKO ESCs under 2iL. (G) Heat map showing the qRT-PCR data of the regulators of DNA methylation and DNA methyltransferases in WT, *Dppa2* KO, *Ctbp1/2* DKO and *Ctbp1/2/Dppa2* TKO ESCs. (H) Western blotting of UHRF1 and TUBULIN in WT, *Dppa2* KO, *Ctbp1/2* DKO and *Ctbp1/2/Dppa2* TKO ESCs. (I) Gene track representing the DPPA2-binding peaks reconstructed using published data (Engelen et al., 2015) and the methylation levels derived from RRBS upstream of *Pramel7* in WT, *Dppa2* KO, *Ctbp1/2* DKO and *Ctbp1/2/Dppa2* TKO ESCs.

***The loss of Ctbp1/2 controls two genetic cascades for ZGA: DUX-dependent and DUX-independent***

Next, to identify the direct target of CtBP1/2 controlling the emergence of MERVL-EGFP-positive cells, we established *Ctbp1/2* DKO ESCs carrying the expression unit of *Ctbp2* regulated by the doxycycline administration. We traced the expression dynamics of some critical genes associated with zygotic genome activation (ZGA) using qRT-PCR analysis. *Dux*, *Mervl*, *Zscan4c* and *Zfp352* were significantly decreased from 3 to 9

days after induction of *Ctbp2* in *Ctbp1/2* DKO ESCs, while these genes were

upregulated from 3 to 9 days (d12-d18) after the depletion of *Ctbp2* expression (Figure

4A). In contrast, *Pramel7* was sharply downregulated (d0-d3) or upregulated (d9-d12)

after *Ctbp2* induction or depletion in *Ctbp1/2* DKO ESCs. These findings suggested that

2-cell associated genes upregulated by the loss of *Ctbp1/2* are classified into two

categories: CtBP1/2-indirect target genes such as *Dux*, *Mervl*, *Zscan4c* and *Zfp352*, and

CtBP1/2-direct target genes such as *Pramel7*. To identify the direct and indirect targets

of CtBP2, we performed RNA-Seq using *Ctbp1/2* conditional KO (cKO) ESCs.

Principal component analysis (PCA) revealed that the global changes of gene

expression profiled by the short-term induction (d0-d3) or depletion (d9-d12) of *Ctbp2*

in *Ctbp1/2* DKO ESCs contributed PC1 axis in the opposite direction (Figure 4B). On

the other hand, the global gene expression changes profiled by the long-term induction

(d3-d9) or depletion of *Ctbp2* (d12-d18) in *Ctbp1/2* DKO ESCs contributed to the PC2

axis in the opposite direction. K-means clustering analysis revealed that CtBP2 rapidly

and positively regulated the genes associated with system development, including

*Ascl2*, *Zic1/4* and *Olig1/3* in ESCs (Figure 4C and Supplementary Table 1,2).

Interestingly, the downregulated genes by CtBP1/2 were separated into two groups. The

genes in cluster 2 containing minor ZGA genes, including *Dux*, *Zscan4*, *Zfp352* and

*Gm4340,* were repressed 3 to 9 days after *Ctbp2* induction (Figure 4C). Major ZGA

genes, including *Pramel6/7/14*, *Crxos* and *Gata1,* and maternal factors, including

*Pou2f3*, *Klf17* and *Pramef12* (*Pramel13*), were in cluster 3, in which the genes were

rapidly downregulated or upregulated by the loss or gain of *Ctbp2*. Next, to extract the

direct or indirect targets of CtBP2, we compared a group of genes rapidly or slowly

repressed by *Ctbp2* induction with a group of genes that are rapidly or delay activated

by *Ctbp2* depletion, respectively (Figure 4D, E). We defined overlapped genes as the

candidates for the direct target (rapid response genes) and indirect target of CtBP2

(delayed response genes), and we displayed the RNA-Seq data of parts of these genes in

the heatmap (Figure 4F). To determine the dependency of rapid response genes and

delayed response genes on DUX function, we constructed a Venn diagram illustrating

the overlap of DUX-independent upregulated genes with rapid response genes and

delayed response genes (Figure 4G). Almost all rapid response genes, but less than half

of delayed response genes, were upregulated by the loss of CtBP2 without DUX

function. Next, to identify the direct target of CtBP1/2, we overlapped the genes that

CtBP2 is localized within +/- 10 kb of the transcription start site (TSS) with rapid and

delayed response genes using the Venn diagram using public ChIP-seq data of CtBP2 in

ESCs (Kim et al., 2015) (Figure 4H). CtBP2 binding peaks were observed in

approximately half of the rapid response genes, whereas only two genes among the

delayed response genes exhibited CtBP2 binding peaks near their TSS. Next, we

displayed the ChIP-Seq track of DUX- and CtBP2-binding peaks and histone H3

Lysine-27 acetylation (H3K27Ac), a typical enhancer mark (Hendrickson et al., 2017)

(Kim et al., 2015)(Zhang et al., 2020), at around the *Pramel7*, *Pou2f3* (rapid response

genes) and *Duxf3*, *Zscan4*c (delayed response genes) (Figure 4I). CtBP2 was localized

within the H3K27Ac-enriched distal enhancer of *Pramel7* and *Pou2f3*. In contrast,

CtBP2-binding peaks were absent, and DUX-binding peaks were localized around the

TSS of *Duxf3* and *Zsacn4c*. To determine the direct effect of CtBP2 at the distal

enhancer of *Pramel7*, we analyzed the effect of CtBP2 overexpression against the

transcriptional activity driven by the distal enhancer of *Pramel7* in WT and *Ctbp1/2*

DKO ESCs (Figure 4J). Luciferase activity was activated in *Ctbp1/2* DKO ESCs

compared with those in WT ESCs, and the CtBP2 overexpression completely

diminished this activation. In summary, our findings indicate that the loss of CtBP1/2

controls two pathways regulating 2C programs: one mediated by *Dux* activation

(*Zscan4c* and *Usp17lc*) and the other directly activated by the loss of CtBP1/2 (*Prame7*

and *Pou2f3)*.

## Figure 4

**Figure 4. The loss of *Ctbp1/2* controls two genetic cascades for ZGA.**

(A) RT-qPCR analysis of *Dux*, *Mervl, Zscan4c Zfp352* and *Pramel7* in *Ctbp1/2* DKO

ESCs without Dox or with Dox. The ΔCt of each gene was calculated by subtracting the

average Ct of Gapdh from that of each gene Ct. Means of ΔCt were calculated from

three biological triplicates. \*\*\*$P < 0.001$, \*\*$P < 0.005$, \*$P < 0.05$ (Tukey-Kramer

multiple comparison test). (B) Principal component analysis (PCA) of *Ctbp1/2* DKO

ESCs without Dox or with Dox. (C) K-means clustering analysis of *Ctbp1/2* DKO ESCs

without Dox or with Dox using the top 1000 differentially expressed genes. (D) Venn

diagram shows the overlapping of down-regulated genes three days after Ctbp2

induction and upregulated genes three days after *Ctbp2* removal. (E) Venn diagram

showing the overlapping of down-regulated genes whose expression is down-regulated

from 3 days to 9 days after *Ctbp2* induction and upregulated genes whose expression is

upregulated from 3 days to 9 days after *Ctbp2* removal. (F) Heatmap showing the

selected genes from K-means clustering analysis (C). (G) Venn diagram shows the

overlapping of upregulated genes by *Ctbp1/2* DKO in *Dux* KO ESCs with rapid or

gradual responses against *Ctbp2* expression. (H) The Venn diagram demonstrates the

extent of overlap between CtBP2-binding genes and the rapid or gradual responses that

are regulated by Ctbp2 expression. (I) Gene track of ChIP-Seq data of H3K27Ac

(GSM4205678), DUX (GSM2525771) and CtBP2 (ERR532426) around *Pramel7*,

*Pou2f3*, *Duxf3* and *Zscan4c*. The red bar indicates the region used for luciferase assay in

(J). (J) The effect of CtBP2 overexpression for the enhancer activity of *Pramel7* distal

enhancer (I) in *Ctbp1/2* DKO ESCs. Relative luciferase activity was calculated with the

values of the empty vector as 1. Means of relative luciferase activity were calculated

from three biological triplicates. \*\*\*$P < 0.001$, \*\*$P < 0.005$, \*$P < 0.05$ (Tukey-Kramer

multiple comparison test).

***Prame family control genetic program of ZGA: DUX-dependent and -independent***

*Pramel7* belongs to the preferentially expressed antigen in melanoma (PRAME*)* gene

family, which is restricted to the testis and various cancers (Kern et al., 2021). Because

the PRAME family gene is one of the most amplified gene families in mammals and is

considered the third largest gene family in the mouse genome (Church et al., 2009), we

focused on the expression of the PRAME gene family in *Ctbp1/2* DKO ESCs. We

collected the amino acid sequences of the PRAME family proteins annotated on the

mm10 genome sequence and created the phylogenetic tree (Supplementary Figures 3A).

The PRAME family genes were found to be clustered on six distinct chromosomes, and

a high similarity of PRAME family proteins was observed to occur on the same

chromosome (Supplementary Figure 3B). Next, we selected the expression levels of

PRAME family genes in WT, *Dux* KO, *Ctbp1/2* DKO and *Ctbp1/2/Dux* TKO from

RNA-Seq data and displayed the hierarchical clustering data in the heatmap (Figure

5A). Interestingly, most PRAME family genes, except for *Lrrc14*, *Lrrc14b*, and

*D5Ertd577e* (*Pramel48*), were upregulated in *Ctbp1/2* DKO ESCs. According to gene

expression patterns, the genes of the PRAME family can be grouped into four

categories. In the first category, including *Lrrc14*, there are genes whose expression is

unaffected by *Ctbp1/2* DKO. In the second category, including *Pramel6* and *Pramel14*,

the genes upregulated by *Ctbp1/2* DKO were further elevated by *Dux* KO. In the third

category, including *Pramel19* and *Pramel39*, the genes upregulated by *Ctbp1/2* DKO

were completely repressed by *Dux* KO (DUX-dependent PRAME family genes). In the

fourth category, including *Pramel7* and *Pramel12*, the genes upregulated by *Ctbp1/2*

DKO were unaffected by *Dux* KO (DUX-independent *Prame* family genes). Next, we

analyzed the expression dynamics of PRAME family genes in the early mouse embryo

using public scRNA-Seq data by removing genes with low expression in early embryos

(FPKM ≤ 1) (Gerlinger et al., 2014). *Pramel19* and *Pramel39*, DUX-dependent PRAME

family genes, were transiently upregulated at the mid-2-cell stage after transient

upregulation of *Dux* at the early 2-cell stage. DUX-independent PRAME family genes

were subdivided into three categories. The first category, including *Pramel6* and

*Pramel7*, adjacent to chromosome 2, were gradually upregulated from mid-2-cell to

peak expression at ICM. The genes, including *Pramel48* and *Pramel13* in the second

category, were not upregulated by ZGA. The genes belonging to the third category,

including *Pramel14* and *Pramel31,* were activated soon after ZGA and continued to be

expressed until 4 cell embryos. Because it has been reported that *Pramel7*, which is

upregulated by the loss of *Ctbp1/2* and ZGA not depending on DUX function, is critical

for early mouse development (Graf et al., 2017), we focused on the PRAMEL7

functions on the emergence of the 2CLCs from pluripotent stem cells. We generated

ESCs with forced expression of *Pramel7* with or without *Dux* KO (Figure 5C, D).

*Pramel7* overexpression expanded 2CLCs, and this expansion was entirely cancelled by

the *Dux* KO (Figure 5E, F). Gene set enrichment analysis comparing WT and *Pramel7*-

overexpressing ESCs using the upregulated genes by the loss of *Ctbp1/2* showed that

the upregulated genes by the loss of *Ctbp1/2* tended to be upregulated by *Pramel7*

overexpression (Figure 5G). Germline-specific genes and ZGA-associated genes were

consistently upregulated by the loss of *Ctbp1/2* and *Pramel7* overexpression (Figure

5H). Some sets of ZGA-associated genes, including *Crxos*, *Gata1* and *Klf17,* were only

upregulated by the loss of *Ctbp1/2*, suggesting that the loss of *Ctbp1/2* controls two

genetic cascades: PRAMEL7-dependent and PRAMEL7-independent pathway. Next,

we separated genes into five categories using K-means clustering analysis based on the

gene expression profiles in WT, *Dux* KO, *Pramel7*-OE and *Pramel7*-OE/*Dux* KO ESCs

(Figure 5I). The first category included the minor ZGA genes whose expression was

upregulated by PRAMEL7 depending on DUX functions. Most genes belonging to the

first cluster were transiently upregulated at mid-2-cell embryos soon after the transient

upregulation of *Dux,* such as the *Zscan4* family; the rest of the genes were upregulated

after eight-cell embryos (Figure 5J). *Dux* expression was elevated in PRAMEL7-

overexpressing ESCs (Figure 5K), and enhancer activity was also upregulated by

PRAMEL7 overexpression (Figure 5L), suggesting that PRAMEL7 controls genetic

program for 2CLCs through the upregulation of *Dux* expression in ESCs. Next, we

focused on cluster 2, in which the genes were downregulated by *Dux* KO and this

downregulation was diminished by PRAMEL7 overexpression (Figure 5I). Cluster 2

was subdivided into two groups based on the gene expression dynamics in early mouse

development (Figure 5M). The first groups, clusters 2a and b, contained the maternal

factor, including *Dazl*, *Zp3,* and *Mael*. The second group consisted of clusters 2c-f, in

which genes were upregulated after major ZGA, including *Pramef17*, *Zp600* and *Sox17*.

Figure 5

**Figure 5. PRMAEL7 controls two genetic cascades for ZGA.**

(A) The heatmap showing the hierarchical clustering of the expressions of Preme family

genes using RNA-Seq data in WT, *Dux* KO, *Ctbp1/2* DKO and *Ctbp1/2/Dux* KO ESCs.

(B) The heatmap shows the hierarchical clustering of the expressions of Preme family

genes using public single-cell RNA-Seq data in early mouse embryos (Gerlinger et al.,

2014). (C) Western blotting anti-FLAG (FLAG-PRAMEL) antibody using lysates of WT,

PRAMEL7-overexpressing, *Dux* KO and PRAMEL7-overexpressing *Dux* KO ESCs. (D)

Genotype PCR using the primer sets located outside the gRNA location to confirm the

large deletion of the *Dux* cluster. (E) Dark fields image visualizing *Mervl*-EGFP signals

in WT, PRAMEL7-overexpressing, *Dux* KO and PRAMEL7-overexpressing *Dux* KO

ESCs. (F) FACS analysis measuring the percentage of *Mervl*-EGFP-positive cells in WT,

PRAMEL7-overexpressing, *Dux* KO and PRAMEL7-overexpressing *Dux* KO ESCs. (G)

Gene set enrichment analysis (GSEA) using the upregulated genes by the *Ctbp1/2*

disruption to compare the gene expression of PRMAEL7-overexpressing ESCs with WT

ESCs. (H) Venn diagram showing the overlapping of the upregulated genes by the

*Ctbp1/2* disruption in ESCs with the upregulated genes by PRAMEL7 overexpression in

ESCs. (I) K-means clustering (K=5) of the gene expression in WT, PRAMEL7-

overexpressing, *Dux* KO, and PRAMEL7-overexpressing *Dux* KO ESCs using top 2000

differentially expressed genes. (J and M) K-means clustering analysis of the genes of

Cluster 1 and Cluster 4 using published single-cell RNA-Seq data of early mouse

embryo(Gerlinger et al., 2014) (K) RT-qPCR analysis of *Dux* in WT and PRAMEL7-

overexpressing ESCs. ΔCt value was calculated by subtracting the Ct value of *Dux* from

that of *Gapdh*. The means of ΔCt was calculated from three biological triplicates. ***$P$

$< 0.001$, **$P < 0.005$, *$P < 0.05$ (Tukey-Kramer multiple comparison test). (L) The effect

of PRAMEL7 overexpression against luciferase activity regulated by the *Dux* enhancer

region. Relative luciferase activity was calculated by setting the luciferase activity of the

empty vector as 1. The means of relative luciferase activity was calculated from three

biological triplicates. ***$P < 0.001$, **$P < 0.005$, *$P < 0.05$ (Tukey-Kramer multiple

comparison test).

***Dux controls a part of genetic cascades for the ZGA program through the PRAMEL7***

***functions***

We generated *Pramel7* KO ESCs and compared the gene expression profiles of WT,

*Pramel7* KO, *Dux* KO, *Pramel7*-OE and *Pramel7*-OE/*Dux* KO ESCs (Supplementary

Figure 1D). Hierarchical clustering analysis showed that *Dux* KO ESCs and *Pramel7*

KO ESCs were in the same cluster (Figure 6A). To investigate the similarity of global

gene expression between *Dux* KO ESCs and *Pramel7* KO ESCs, we employed scatter

plot analysis to visualize the relative expression alteration induced by *Dux* KO and

*Pramel7* KO (Figure 6B). The alterations in global expression observed in *Dux* KO and

*Pramel7* exhibited a positive correlation, particularly for downregulated genes. GSEA

also supported a positive correlation of expression changes between *Dux* KO and

*Pramel7* KO (Figure 6C, D). The Venn diagram using the upregulated or

downregulated genes by *Dux* KO and *Pramel7* KO showed the shared genes and

independent genes regulated by DUX and PRAMEL7 (Figure 6E, F). The genes

associated with ZGA were separated into DUX-dependent (*Zscan4* family) and

DUX/PRAMEL7-dependent (*Cphx* and *Zfp534/600*). Since we found that the

expression of *Pramel7* was found to be significantly reduced in *Dux* KO ESCs (Figure

6G), we tested the possibility that PRAMEL7-overexpression could rescue the

phenotype of *Dux* KO ESCs. The alteration of global gene expression by *Dux* KO in

WT ESCs and PRAMEL7 overexpression in *Dux* KO ESCs demonstrated a negative

correlation (Figure 6H). Furthermore, GSEA demonstrated that the overexpression of

PRAMEL7 in *Dux* KO ESCs effectively rescued the downregulated genes resulting

from *Dux* KO (Figure 6I). These findings indicate that DUX controls two genetic

pathways of the 2C program; one is directly regulated by DUX, and the other is

regulated by activation.

**Figure 6**

**Figure 6. DUX regulates parts of the genetic program for ZGA by activating**

***Pramel7.***

(A) Hierarchical clustering analysis of WT ESCs, *Pramel7* KO ESCs, *Dux* KO ESCs, PRMAEL7-overexpressing ESCs and PRMAEL7-overexpressing *Dux* KO ESCs based on the gene expression. (B) Scatter plot analysis of the correlation between the expression changes by *Dux* KO and *Pramel7* KO in WT ESCs. (C) Gene set enrichment analysis (GSEA) using the down-regulated genes by the *Dux* KO to compare the gene expression of *Pramel7* KO ESCs with WT ESCs. (D) Gene set enrichment analysis (GSEA) using the upregulated genes by the *Dux* KO to compare the gene expression of *Pramel7* KO ESCs with WT ESCs. (E) Venn diagram showing the overlapping of the down-regulated

genes by *Dux* KO in ESCs with the down-regulated genes by *Pramel7* KO in ESCs. (F)

Venn diagram showing the overlapping of the upregulated genes by *Dux* KO in ESCs

with the upregulated genes by *Pramel7* KO in ESCs. (G) qRT-PCR analysis of *Pramel7*

in wild type and *Dux* KO ESCs. ΔCt value was calculated by subtracting the Ct value of

*Pramel7* from that of *Gapdh*. The means of ΔCt was calculated from three biological

triplicates. ***$P < 0.001$, **$P < 0.005$, *$P < 0.05$ (Tukey-Kramer multiple comparison

test). (H) Scatter plot analysis of the correlation between the expression changes by *Dux*

KO in WT ESCs and *Pramel7*-OE in *Dux* KO ESCs. (K) Gene set enrichment analysis

(GSEA) using the down-regulated genes by the *Dux* KO to compare the gene expression

of PRMAEL7-overexpressing in *Dux* KO ESCs with *Dux* KO ESCs.

**Discussion**

**Figure 7**

WT ESCs                    *Ctbp1/2* DKO ESCs            *Ctbp1/2* DKO ESCs

**Figure 7. A schematic model for the activation of 2C-related genes by the Ctbp1/2**

**disruption.**

In wild-type embryonic stem cells (ESCs), minor and major ZGA genes and germline-

specific genes are repressed. *Pramel7* and some parts of major ZGA genes are rapidly

reactivated following the disruption of *Ctbp1/2*. PRAMEL7 activates germline-specific

genes and minor ZGA genes in a manner dependent on UHRF1 degradation or DUX,

respectively.

In this study, we revealed that the disruption of *Ctbp1/2* in ESCs activates DUX-

dependent and independent molecular cascades to establish totipotent states (Figure 7).

Most DUX-dependent genes activated by *Ctbp1/2* DKO are transiently expressed at

around 2C stage embryo, belonging to the minor ZGA genes. In contrast, DUX-

independent genes activated by *Ctbp1/2* DKO are categorized into two groups:

Maternally expressed genes and the genes upregulated after major ZGA, some of which

are regulated by DPPA2. PRAMEL7, which is directly activated by the disruption of

*Ctbp1/2* in a DUX-independent manner, controls DUX-dependent genetic cascades

through DUX activation and DUX-independent genetic cascades. We present new

insights into the DUX-independent molecular pathways that ensure the early mouse

embryo's survival.

Our data clearly showed that reducing CtBP protein levels in ESCs activates a part of the

genetic program regulating the development of early mouse embryos. scRNA-seq data of

early mouse embryos shows that a large amount of *Ctbp2* mRNA is maternally deposited

in unfertilized eggs and significantly reduced after fertilization (Gerlinger et al., 2014).

In contrast, the protein level of CtBP2 was consistent between the 2CLCs that emerged

from pluripotent stem cells and pluripotent stem cells, indicating that the expression

levels of CtBP proteins are quite different between 2C embryos and 2CLCs. This

difference may explain the dependence of the emergence of 2CLCs and early mouse

development on DUX functions. However, the molecular mechanism of the clearance of

*Ctbp2* mRNA after fertilization is currently unclear. It has been reported that the inhibition

of minor ZGA inhibition with 5, 6-dichloro-1-β-D-ribofuranosyl-benzimidazole (DRB)

failed in a part of maternal mRNA clearance, including maternal *Ctbp2* mRNA (Abe et

al., 2018). Moreover, the maternal knockout of B-cell translocation gene-4 (BTG4),

which involves the maternal mRNA clearance by CCR4-NOT-dependent deadenylation,

was found to impair the maternal *Ctbp2* mRNA clearance. (Yu et al., 2016). These

findings suggest that the products of minor ZGA may work with BTG4 to destabilize the

clearance of maternal *Ctbp2* mRNA.

Transcriptome analysis of several knockout lines of ESCs provided evidence that the

activated genes by the disruption of *Ctbp1/2* are subdivided into three groups: both DUX

and DPPA2-dependent genes, DPPA2-dependent genes and both DUX and DPPA2-

independent genes (Figure 7). DUX- and DPPA2-dependent genes were enriched in the

minor ZGA genes at 2C embryo, including the *Zscan4* family. In contrast, DUX and

DPPA2-independent genes contained maternal factors, including *Dazl* and *Mael*, and the

major ZGA genes (*Gata1*, *Crxos* and *Pramef17*) in the early mouse embryo. A previous

study has shown that totipotent-like stem cells (TLSCs) induced by the chemical

inhibitors of DOT1L, KDM5B and G9a/GLP expressed a large number of minor and

major ZGA genes as well as maternal genes (Yang et al., 2022). The re-establishment of

broad H3K4me3 domains induced by the inhibition of KDM5B is thought to be involved

in the upregulation of maternal genes. However, because TLSCs were not established

using *Dux* KO ESCs, the role of DUX in the transition from ESCs to TLSCs and the

upregulation of maternal genes are largely unknown. Our data demonstrate that the

upregulation of maternal genes by *Ctbp1/2* DKO does not depend on DUX function and

is partially regulated by *Pramel7* induction.

*Prame* is one of the most amplified gene families in mammals, and the mouse *Prame*

gene family is the third largest family in the genome (Chang et al., 2011). We presented

that the disruption of *Ctbp1/2* activates the majority of *Prame* family genes in ESCs, some

of which are upregulated by DUX function and the rest directly by CtBP1/2. Interestingly,

*Prame* genes that depend on DUX are enriched in minor ZGA genes, including *Zscan4*

family genes, while *Prame* genes independent of DUX are upregulated after major ZGA.

Many *Prame* genes are repressed soon after induction of CtBP2 in *Ctbp1/2* DKO ESCs,

indicating that CtBP1/2 directly repress *Prame* family genes in ESCs. However, the

positive regulators of *Prame* family genes are largely unknown. Recent reports revealed

that the overexpression of multiple homeobox genes, oocyte-specific homeobox (Obox)

activate minor and major ZGA genes in mESCs (Guo et al., 2024; Ji et al., 2023;

Sakamoto et al., 2024). Interestingly, the overexpression of OBOX3/5 but not DUX

activate *Prame7* and *Pramef17* (*Pramel14*), which are upregulated by the loss of *Ctbp1/2*

without DUX, but not *Gm12794* (*Pramel19*) and *A430089l19Rik* (*Pramel39*), which are

upregulated in a DUX-dependent manner (Sakamoto et al., 2024). Because Obox family

genes were silent in *Ctbp1/2* DKO ESCs, another positive regulator for Prame family

genes may be activated by *Ctbp1/2* DKO in mESCs. Expression of *Prame* family genes

has been discovered in various cancer and germline cells (Kern et al., 2021), and the

expression pattern of *Prame* family genes is similar to that of *Pou5f1* (Oct4) (Bortvin et

al., 2003), suggesting that transcriptional network centred upon OCT4 is a positive

regulator of *Prame* family genes.

Although DUX is a master regulator for the transition from pluripotent stem cells into

2CLC, *Dux* KO embryos are normally born and fertile. The data presented here show that

most of the *Dux* KO phenotype is rescued by PRAMEL7 overexpression in ESCs. It has

been reported that *Prame* family genes are upregulated from the early phase of 2CLCs

emergence traced by the *Zscan4* and *Mervl* reporter system based on the scRNA-seq

(Eckersley-Maslin et al., 2016; Rodriguez-Terrones et al., 2018) and MERVL-positive

2CLCs are eliminated by *Dux* KO (De Iaco et al., 2017). Based on these findings, we

speculate that DUX controls two major genetic cascades for the development of early

mouse embryos: direct transcriptional regulation and PRAME-mediated transcriptional

regulation. *Dux* KO ESCs showed a significant reduction of *Pramel7* transcription, while

the expression of *Pramel7* is normally activated in *Dux* KO embryos at the mid-2-cell

stage (Guo et al., 2019). These findings suggest that the DUX-independent mechanism

activates the *Prame* family *in vivo*, and this mechanism might be induced in ESCs by the

disruption of *Ctbp1/2*.

Our results provide new insights into understanding DUX-independent genetic cascades,

ensuring the development of early mouse embryos. Although the visualization of 2CLCs

marked by the transcription of MERVL contributed to identifying the regulators for the

transition between pluripotent stem cells and 2CLCs, the knockout phenotypes of the

identified factors were functionally dependent on DUX. Developing the visualization

system for new 2CLCs regulating by DUX-independent genetic cascades in ESCs will be

needed.

**References**

**Abe, K., Funaya, S., Tsukioka, D., Kawamura, M., Suzuki, Y., Suzuki, M. G., Schultz,**
**R. M. and Aoki, F.** (2018). Minor zygotic gene activation is essential for mouse
preimplantation development. *Proceedings of the National Academy of Sciences* **115**,
E6780–E6788.

**Bortvin, A., Eggan, K., Skaletsky, H., Akutsu, H., Berry, D. L., Yanagimachi, R.,**
**Page, D. C. and Jaenisch, R.** (2003). Incomplete reactivation of Oct4-related genes
in mouse embryos cloned from somatic nuclei. *Development* **130**, 1673–1680.

**Chang, T. C., Yang, Y., Yasue, H., Bharti, A. K., Retzel, E. F. and Liu, W. S.** (2011).
The expansion of the PRAME gene family in Eutheria. *PLoS One* **6**,.

**Chen, Z. and Zhang, Y.** (2019). Loss of DUX causes minor defects in zygotic genome
activation and is compatible with mouse development. *Nat Genet* **51**, 947–951.

**De Iaco, A., Planet, E., Coluccio, A., Verp, S., Duc, J. and Trono, D.** (2017). DUX-
family transcription factors regulate zygotic genome activation in placental
mammals. *Nat Genet* **49**, 941–945.

**Eckersley-Maslin, M. A., Svensson, V., Krueger, C., Stubbs, T. M., Giehr, P., Krueger,**
**F., Miragaia, R. J., Kyriakopoulos, C., Berrens, R. V., Milagre, I., et al.** (2016).
MERVL/Zscan4 Network Activation Results in Transient Genome-wide DNA
Demethylation of mESCs. *Cell Rep* **17**, 179–192.

**Eckersley-maslin, M. A., Alda-catalinas, C. and Reik, W.** (2018). Dynamics of the
epigenetic landscape during the maternal- to-zygotic transition. *Nat Rev Mol Cell*
*Biol* **19**, 436–450.

**Eckersley-Maslin, M., Alda-Catalinas, C., Blotenburg, M., Kreibich, E., Krueger, C.**
**and Reik, W.** (2019). Dppa2 and Dppa4 directly regulate the Dux-driven zygotic
transcriptional program. *Genes Dev* **33**, 194–208.

**Eckersley-maslin, M. A., Parry, A., Blotenburg, M., Krueger, C., Ito, Y., Nila, V.,**
**Franklin, R., Narita, M., Santos, C. S. D. and Reik, W.** (2020). Epigenetic priming
by Dppa2 and 4 in pluripotency facilitates multi-lineage commitment. *Nat Struct*
*Mol Biol* **27**, 696–705.

**Engelen, E., Brandsma, J. H., Moen, M. J., Signorile, L., Dekkers, D. H. W.,**
**Demmers, J., Kockx, C. E. M., Ozgür, Z., Van Ijcken, W. F. J., Van Den Berg,**
**D. L. C., et al.** (2015). Proteins that bind regulatory regions identified by histone

modification chromatin immunoprecipitations and mass spectrometry. *Nat Commun*

**6**,.

**Ge, S. X., Son, E. W. and Yao, R.** (2018). iDEP: An integrated web application for

differential expression and pathway analysis of RNA-Seq data. *BMC Bioinformatics*

**19**, 1–24.

**Gerlinger, M., Swanton, C., Greaves, M. and Maley, C. C.** (2014). Single-Cell RNA-

Seq Reveals Dynamic, Random Monoallelic Gene Expression in Mammalian Cells.

**343**, 193–196.

**Graf, U., Casanova, E. A., Wyck, S., Dalcher, D., Gatti, M., Vollenweider, E.,**

**Okoniewski, M. J., Weber, F. A., Patel, S. S., Schmid, M. W., et al.** (2017).

Pramel7 mediates ground-state pluripotency through proteasomal-epigenetic

combined pathways. *Nat Cell Biol* **19**, 763–773.

**Gretarsson, K. H. and Hackett, J. A.** (2020). Dppa2 and Dppa4 counteract de novo

methylation to establish a permissive epigenome for development. *Nat Struct Mol*

*Biol* **27**, 706–716.

**Grow, E. J., Weaver, B. D., Smith, C. M., Guo, J., Stein, P., Shadle, S. C.,**

**Hendrickson, P. G., Johnson, N. E., Butterfield, R. J., Menafra, R., et al.** (2021).

p53 convergently activates Dux/DUX4 in embryonic stem cells and in

facioscapulohumeral muscular dystrophy cell models. *Nat Genet* **53**, 1207–1220.

**Guo, M., Zhang, Y., Zhou, J., Bi, Y., Xu, J., Xu, C., Kou, X., Zhao, Y., Li, Y., Tu, Z.,**

**et al.** (2019). Precise temporal regulation of Dux is important for embryo

development. *Cell Res* **7**, 1–4.

**Guo, Y., Kitano, T., Inoue, K., Murano, K., Hirose, M., Li, T. D., Sakashita, A., Ishizu,**

**H., Ogonuki, N., Matoba, S., et al.** (2024). Obox4 promotes zygotic genome

activation upon loss of dux. *Elife* **13**, 1–25.

**Hendrickson, P. G., Doráis, J. A., Grow, E. J., Whiddon, J. L., Lim, J.-W., Wike, C.**

**L., Weaver, B. D., Pflueger, C., Emery, B. R., Wilcox, A. L., et al.** (2017).

Conserved roles of mouse DUX and human DUX4 in activating cleavage-stage

genes and MERVL/HERVL retrotransposons. *Nat Genet* **49**, 925–934.

**Hu, Z., Tan, D. E. K., Chia, G., Tan, H., Leong, H. F., Chen, B. J., Lau, M. S., Tan,**

**K. Y. S., Bi, X., Yang, D., et al.** (2020). Maternal factor NELFA drives a 2C-like

state in mouse embryonic stem cells. *Nat Cell Biol* **22**, 175–186.

**Hulsen, T., de Vlieg, J. and Alkema, W.** (2008). BioVenn - A web application for the

comparison and visualization of biological lists using area-proportional Venn diagrams. *BMC Genomics* **9**, 1–6.

Ishiuchi, T., Enriquez-Gasca, R., Mizutani, E., Bošković, A., Ziegler-Birling, C., Rodriguez-Terrones, D., Wakayama, T., Vaquerizas, J. M. and Torres-Padilla, M.-E. (2015a). Early embryonic-like cells are induced by downregulating replication-dependent chromatin assembly. *Nat Struct Mol Biol* **22**, 662–71.

Ishiuchi, T., Enriquez-gasca, R., Mizutani, E., Bošković, A., Ziegler-birling, C., Rodriguez-terrones, D., Wakayama, T. and Vaquerizas, J. M. (2015b). Articles Early embryonic-like cells are induced by downregulating replication-dependent chromatin assembly. *Nature Publishing Group* **22**, 662–671.

Ji, S., Chen, F., Stein, P., Wang, J., Zhou, Z., Wang, L., Zhao, Q., Lin, Z., Liu, B., Xu, K., et al. (2023). OBOX regulates mouse zygotic genome activation and early development. *Nature* **620**, 1047–1053.

Kern, C. H., Yang, M. and Liu, W. S. (2021). The PRAME family of cancer testis antigens is essential for germline development and gametogenesis. *Biol Reprod* **105**, 290–304.

Kim, T. W., Kang, B. H., Jang, H., Kwak, S., Shin, J., Kim, H., Lee, S. E., Lee, S. M., Lee, J. H., Kim, J. H., et al. (2015). Ctbp2 Modulates NuRD-Mediated Deacetylation of H3K27 and Facilitates PRC2-Mediated H3K27me3 in Active Embryonic Stem Cell Genes during Exit from Pluripotency. *Stem Cells* **33**, 2442–2455.

Kreibich, E., Blotenburg, M., Alda-Catalinas, C., Reik, W., Krueger, C. and Eckersley-Maslin, M. (2019). Dppa2 and Dppa4 directly regulate the Dux-driven zygotic transcriptional program. *Genes Dev* **33**, 194–208.

Macfarlan, T. S., Gifford, W. D., Driscoll, S., Lettieri, K., Rowe, H. M., Bonanomi, D., Firth, A., Singer, O., Trono, D. and Pfaff, S. L. (2012). Embryonic stem cell potency fluctuates with endogenous retrovirus activity. *Nature* **487**, 57–63.

Percharde, M., Lin, C., Yin, Y., Huang, B., Shen, X., Ramalho-santos, M., Percharde, M., Lin, C., Yin, Y., Guan, J., et al. (2018). Article A LINE1-Nucleolin Partnership Regulates Early Development and ESC Identity Article A LINE1-Nucleolin Partnership Regulates Early Development and ESC Identity. *Cell* **174**, 1–15.

Robinson, J. T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E. S., Getz, G. and Mesirov, J. P. (2011). Integrative genomics viewer. *Nat Biotechnol* **29**, 24–

26.

**Rodriguez-Terrones, D., Gaume, X., Ishiuchi, T., Weiss, A., Kopp, A., Kruse, K.,**
**Penning, A., Vaquerizas, J. M., Brino, L. and Torres-Padilla, M. E.** (2018). A
molecular roadmap for the emergence of early-embryonic-like cells in culture. *Nat*
*Genet* **50**, 106–119.

**Sakamoto, M., Ito, A., Wakayama, S., Sasaki, H., Wakayama, T. and Ishiuchi, T.**
(2024). Detection of newly synthesized RNA reveals transcriptional reprogramming
during ZGA and a role of Obox3 in totipotency acquisition. *Cell Rep* **43**, 114118.

**Whiddon, J. L., Langford, A. T., Wong, C. J., Zhong, J. W. and Tapscott, S. J.** (2017).
Conservation and innovation in the DUX4-family gene network. *Nat Genet* **49**, 935–
940.

**Wolf, G., de Iaco, A., Sun, M. A., Bruno, M., Tinkham, M., Hoang, D., Mitra, A.,**
**Ralls, S., Trono, D. and Macfarlan, T. S.** (2020). Krab-zinc finger protein gene
expansion in response to active retrotransposons in the murine lineage. *Elife* **9**, 1–
22.

**Wu, K., Liu, H., Wang, Y., He, J., Xu, S., Chen, Y., Kuang, J., Liu, J., Guo, L., Li, D.,**
**et al.** (2020). SETDB1-Mediated Cell Fate Transition between 2C-Like and
Pluripotent States. *Cell Rep* **30**, 25-36.e6.

**Yamamoto, M., Suwa, Y., Sugiyama, K., Okashita, N., Kawaguchi, M., Tani, N.,**
**Matsubara, K., Nakamura, A. and Seki, Y.** (2020). PRDM14-CtBP1/2-PRC2
complex regulates transcriptional repression during transition from primed to naïve
pluripotency. *J Cell Sci* **133**, 1–13.

**Yang, M., Yu, H., Yu, X., Liang, S., Hu, Y., Luo, Y., Izsvák, Z., Sun, C. and Wang, J.**
(2022). Chemical-induced chromatin remodeling reprograms mouse ESCs to
totipotent-like stem cells. *Cell Stem Cell* **29**, 400-418.e13.

**Yu, C., Ji, S. Y., Sha, Q. Q., Dang, Y., Zhou, J. J., Zhang, Y. L., Liu, Y., Wang, Z. W.,**
**Hu, B., Sun, Q. Y., et al.** (2016). BTG4 is a meiotic cell cycle-coupled maternal-
zygotic-transition licensing factor in oocytes. *Nat Struct Mol Biol* **23**, 387–394.

**Zhang, T., Zhang, Z., Dong, Q., Xiong, J. and Zhu, B.** (2020). Histone H3K27
acetylation is dispensable for enhancer activity in mouse embryonic stem cells.
*Genome Biol* **21**, 1–7.

**Author Contributions**

Y.S. conceived the study and experimental design, performed experiments and wrote the

manuscript. K.S., K.Y. and N.H. performed most experiments. R.M. M.M., A.S. and Y.M

performed some experiments. K.S., K.Y. and N.H. contributed equally to this work.

**Acknowledgements**

We thank all members of Seki laboratory for stimulating discussions. This study was

supported by JSPS KAKENHI Grant Numbers JP18H02422, JP20H05375 and

JP22H04682.

**Conflict of interest**

The authors declare no competing interests.

**Data availability**

RNA-Seq data has been deposited in Gene Expression Omnibus (GSE253944)