Circatidal control of gene expression in the deep-sea hot vent shrimp

Rimicaris leurokolos

Hongyin Zhang[1,2,†], Takuya Yahagi[3,†], Norio Miyamoto[4], Chong Chen[4], Qingqiu

Jiang[1,2], Pei-Yuan Qian[5,6], and Jin Sun[1,2,#]

[1]Key Laboratory of Evolution & Marine Biodiversity (Ministry of Education) and

Institute of Evolution & Marine Biodiversity, Ocean University of China, Qingdao

266003, China

[2]Laboratory for Marine Biology and Biotechnology, Qingdao Marine Science and

Technology Center, Laoshan Laboratory, Qingdao 266237, China

[3]Department of Marine Ecosystems Science, Atmosphere and Ocean Research Institute,

The University of Tokyo, 5-1-5 Kashiwanoha, Kashiwa, Chiba 277-8564, Japan

[4]X-STAR, Japan Agency for Marine-Earth Science and Technology (JAMSTEC), 2-15

Natsushima-cho, Yokosuka, Kanagawa 237-0061, Japan

[5]Department of Ocean Science, Hong Kong University of Science and Technology,

Clear Water Bay Road, Hong Kong SAR, China

[6]Southern Marine Science and Engineering Guangdong Laboratory (Guangzhou),

Guangzhou, China

Correspondence:

Jin Sun, jin_sun@ouc.edu.cn

[†]These authors contributed equally to this work

Abstract

Biological clocks are a ubiquitous feature of all life, enabling the use of natural environmental cycles to track time. Although studies on circadian rhythms have contributed greatly to the knowledge of chronobiology, biological rhythms in dark biospheres such as the deep sea remain poorly understood. Here, based on a laboratory free-running experiment, we reveal potentially endogenous rhythms in gene expression of the deep-sea hydrothermal vent shrimp *Rimicaris leurokolos*. Oscillations with ~12-hour periods, likely reflecting tidal influence, greatly prevail over others in the temporal transcriptome, indicating *R. leurokolos* likely depends on a circatidal clock consisting of at least some components independent of the circadian clocks. The tidal transcripts exhibit an antiphased expression pattern divided into two internally synchronized clusters, correlated with wide-ranging biological processes that occur in the nucleus and cytoplasm, respectively. In addition, comparing the tidal transcripts with the ~12-hour ultradian rhythms genes in fruit flies and mice shows large similarity, indicating the likely scenario of broad impact of the tide on the ~12-hour oscillations across the metazoan. These findings not only provide new insights into the temporal adaptations in deep-sea organisms but also highlight deep-sea hydrothermal vent organisms as intriguing models for chronobiological, particularly 12-hour ultradian rhythms, studies.

Keywords: biological rhythm, deep sea, hydrothermal vent, adaptation, *Rimicaris leurokolos*, transcriptomics

1. Introduction

As an adaptation to natural periodic environmental changes, biological clocks have been evolving in all kingdoms of life to keep track of time [1-3]. Numerous organisms exhibit endogenous rhythms dominated by circadian periodicity of ~24 hours at behavioral, physiological, and molecular levels, which help them anticipate daily changes such as light, temperature, and food availability [4-6]. To achieve this, animals rely on their internal circadian clock systems like transcription-translation feedback loops (TTFLs) that can be entrained to match the environmental cycles [7]. In dark biospheres such as the underground and deep-sea habitats, however, the absence of sunlight veils organisms there from the direct effects of everlasting day-night transitions. How organisms living in the dark schedule their activities through time, as an adaptation to darkness, remains obscure [8, 9].

The deep ocean, the largest but the least studied habitat on Earth [10], is of great interest for chronobiological research due to its distinct ecological environment [11]. With increasing depth, the sunlight intensity dramatically decreases and the visible light spectrum rapidly narrows to result in zero sunlight penetration below 1000 m even in the clearest oceanic waters [12]. From the ecological perspective, it is more appropriate for deep-sea organisms to synchronize their biological rhythms with other zeitgebers (e.g., tidal cycles) instead of the light cycle [13]. Indeed, various environmental fluctuations related to tidal cycles have been observed in the deep, especially in hydrothermal vent ecosystems [14-17]. Vent faunas rely strongly on their local environment, and survive on organic compounds produced by chemosynthetic microbes utilizing dissolved chemicals in the vent fluids [18]. Therefore, vent animals are likely to be affected by spatial and temporal variations in environmental factors modulated by the tide. Several previous studies have reported potential tide-related rhythms of behaviour and community dynamics in deep-sea organisms from chemosynthetic ecosystems [19-21], and a recent study revealed tidal cycles dominated the gene expression of the deep-sea vent mussel *Bathymodiolus azoricus* through periodic *in situ* sampling across 24.8 hours [22]. Although various evidence suggests that vent organisms exhibit biological rhythms in the wild, whether they need endogenous rhythms or clocks to accommodate and anticipate local environmental changes to maximize their fitness is still poorly understood.

In the Iheya North hydrothermal vent field of the Okinawa Trough (figure 1*a*), tidal modulation on the deep-sea current dynamics was observed previously [23]. According

to a published long-term temperature measurement (Mean ± SD = 4.4 ± 0.1□) recorded
at the periphery of the vent field where bathymodioline mussels typically occur, a strong
semidiurnal tidal cycle (12.4-hour periodicity) was found in the environmental
fluctuations as well (figure 1*b*). Unlike mussels at the vent periphery, however, the
alvinocaridid shrimp *Rimicaris leurokolos* (formerly *Shinkaicaris leurokolos* [24]) lives
in close proximity to the vigorously venting orifice (0.2–0.8 m and occasionally less
than 0.2 m from the vent [25]) where the hot fluid and cold surrounding seawater mix
spontaneously and dramatically [26] (figure 1*c*). In the *R. leurokolos* colony,
environmental factors fluctuate frequently and dramatically at the seconds-minutes scale,
which adds unpredictable noises to the tidal cycles. In fact, the temperature could
rapidly fluctuate between 7.63 to 17.74°C (Mean ± SD = 10.16 ± 1.71°C) within
approximately 2.5 minutes (figure 1*d*, electronic supplementary material, data S1). As
such, *R. leurokolos* is likely unable to directly use the unsteady, highly variable
environmental factors as zeitgebers to entrain its biological rhythm. However, patterns of
venting activity across a longer time-span is known to be linked to tidal cycles [27].
Therefore, we hypothesised that an internal clock may integrate useful periodic cues
whether they are clear (e.g., pressure) or noisy (e.g., water chemistry), and thus help hot
vent shrimps adapt and keep track of periodic environmental changes.
100
Although circadian patterns are dominant in terrestrial and coastal animals, recent
studies revealed robust ultradian rhythms (e.g., ~12-hour cycling gene expression) in
model organisms such as mice and fruit flies [28, 29]. These rhythms were found to be
cell- autonomous, driven by a distinct 12-hour pacemaker largely independent from the
circadian clock. As many homologous genes were found rhythmic with a period of ~12
hours both in terrestrial and marine animals, a hypothesis was then proposed that the
circatidal rhythms in marine animals and ~12-hour rhythms in terrestrial animals are
evolutionarily conserved [30]. However, given the dominant effects of the ~24-hour
circadian system, it is challenging to distinguish the ultradian rhythms from circadian
rhythms in terrestrial or coastal animals. The hot vent shrimp studied herein lives in the
deep sea environment lacking day-night cycles, and thus provides an opportunity to
examine this hypothesis.
113
Here, through a laboratory free-running experiment across 72 hours, we addressed the
biological rhythms of gene expression in the vent shrimp *R. leurokolos* and tested if it
can maintain rhythmicity under constant conditions. Our work reveals free-running
rhythms dominated by 12-hour tidal patterns in the transcriptome of *R. leurokolos*,

which are likely driven by an unidentified endogenous clock including but not limited to
circatidal components. The tidal transcripts exhibit a bimodal gene expression pattern,
and influence broad cellular processes such as DNA replication, DNA repair, regulation
of transcription, transport, cell cycle, and responses to environmental stresses. In
addition, we found *R. leurokolos* shares many common genes and biological processes
in the tidal rhythms with the ~12-hour rhythms in sea anemones, fruit flies, and
mammals, suggesting the evolutionary conservation of ~12-hour gene expression across
a broad range of phyla. These findings highlight the prospect for deep-sea organisms in
chronobiology, especially in the evolution and molecular mechanisms of ~12-hour
ultradian rhythms.
128

2. Materials and methods

(a) Environmental temperature monitoring
The local fluctuations in the temperature of a *Rimicaris leurokolos* colony on the NBC
chimney (126°53.80'E, 27°47.45'N; 982 m deep) of the Original Site, Iheya North vent
field were measured using a RINKO dissolved oxygen and temperature sensor (JFE
Advantech Co., Ltd.) during dive #1610 of the remotely operated vehicle (ROV)
Hyper-Dolphin on-board R/V *KAIYO* cruise KY14-01 for 162 seconds, and data were
recorded every second (electronic supplementary material, data S1). Published
temperature data in a peripheral area of the original site, Iheya North vent field far away
from active chimneys, which were recorded continuously every 10 min from the 4th of
September 2010 to the 20th of August 2011, were re-analyzed [31]. We used the
Lomb-Scargle periodogram to detect periodicity in the temperature time series.
141

(b) Sampling procedure and experiment design
A total of 184 adult individuals of *Rimicaris leurokolos* were collected by a suction
sampler from the top of the NBC chimney at the Original Site, Iheya North hydrothermal
field during dive #1613 of ROV *Hyper-Dolphin* on-board R/V *KAIYO* in the cruise
KY14-01. Specimens of *R. leurokolos* were kept in canister bottles on the ROV prior to
recovery on-board the research vessel. After sampling, it took about 2 hours
(15:17–17:07 JST) for the ROV to be recovered. Once on-board the ship, under natural
light, 154 live shrimps, excluding 25 dead individuals and 5 ovigerous females, were
immediately transferred into a lightproof seawater container. This container was placed
in a dark cold room at 4°C by 19:30 JST for the subsequent free-running experiment.
152

The free-running experiment began at 22:00 JST, aiming to investigate the gene

expression of *R. leurokolos* in a constant environment over time. During the entire
experiment light was turned off, although a faint orange light from the power strip
remained lit. As this was very dim and with a wavelength close to red light, the impact
of this should be minimal. At each sampling timepoint, shrimps were collected by a
meshed scoop, dissected with scissors without anesthesia into cephalothorax and
abdomen, and fixed every 4 hours for a total duration of 72 hours. The cephalothorax of
each individual were preserved in RNAlater (Invitrogen) and used for RNA-Seq in this
study. During the sampling, we did not provide any food source as the previous
laboratory study on behaviours of vent shrimps revealed that the introduction of food
largely increased the activities of these shrimps [32]. As a possible zeitgeber [33], an
external food source may give an opportunity for shrimps to synchronize their activities
and even gene expression at a certain time, which should be avoided in the free-running
experiment. Despite that, *R. leurokolos* may feed on bacteria attached to other
individuals. We did not find any dead shrimp during the free-running experiment, which
might reflect the health conditions of *R. leurokolos*.
169
(c) RNA extraction, sequencing and assembly
Total RNA was extracted from individuals using TRIzol reagent (ThermoFisher)
following the manufacturer's protocol. The mRNA from each shrimp cephalothorax
was enriched by Oligo-dT probes and further reverse transcribed to first-strand cDNA
for library construction. All cDNA libraries were sequenced on an Illumina NovaSeq
6000 platform with 150 bp paired-end mode. A total of 57 RNA-Seq samples were
generated from *R. leurokolos* during 72 hours of sampling at 4-hour intervals with three
replicates per time point. Raw reads were firstly trimmed using Trimmomatic version
0.39 [34], and then potentially associated bacteria contamination reads were removed
with Kraken2 version 2.0.8-beta [35]. *De novo* assembly was performed with Trinity
version 2.11.0 [36] (set jaccard clip option). Salmon version 1.4.0 [37] was used to
quantify the expression of each assembled transcript.
182
(d) Gene quantification and annotation
The TMM cross-sample normalization method [38] was applied to remove the
sequencing variations and potential "batch effects" among samples using the
abundance_estimates_to_matrix.pl script in the Trinity toolkit. In addition, lowly
expressed transcripts with 0 counts in at least 38 samples were filtered out, leaving 23,905
transcripts for subsequent analyses. The predicted proteins were annotated by searching
the candidate protein sequences against the NCBI non-redundant (NR) database using

BLASTP with DIAMOND version 2.0.13 [39] with ultra-sensitive mode and E-value
threshold of 1×10^{-5}, the Kyoto Encyclopedia of Genes and Genomes (KEGG) database
via KEGG Automatic Annotation Server (KAAS), and the Gene Ontology (GO) via
BLAST2GO version 6.0 [40]. The gene enrichment analyses were performed with
ClusterProfiler version 4.2.2 [41].
195

(e) Gene expression and detrend analyses
Principle components analysis (PCA) was performed using the plotPCA function in R
package DESeq2 version 1.34.0 [42]. Permutational multivariate analysis of variance
(PERMANOVA) and analysis of similarities (ANOSIM) were applied to test the
statistical significance of the PCA results. The potential monotonic trends in the raw gene
expression of all transcripts were determined using the Mann-Kendall trend test, and we
pinpointed two overall clusters of transcripts that had either an upward trend (S score > 0
and *P*-value < 0.05) or a downward trend (S score < 0 and *P*-value < 0.05).
204

To remove the trend and produce a steady baseline for subsequent rhythm analyses, we
used CEEMDAN [43], an adaptive decomposition method designed for non-stationary
and non-linear time series signals, which minimizes the impact of distorting the period,
phase or amplitude change of the oscillations. By this method, each gene expression
data could be decomposed into three intrinsic mode functions (IMFs), and a residual
that represents the trend (electronic supplementary material, figure S1). In addition to
the potential periodic patterns, the three IMFs are also likely to contain noises. We then
selected the relevant IMFs and reconstructed the detrended gene expression data under
the criterion suggested by Ayenu-Prah & Attoh-Okine [44]. To verify the effectiveness
of CEEMDAN detrend analysis, we compared the rhythm detection results using gene
expression data before and after detrending. Rhythmic datasets identified using raw and
detrended data share many common transcripts, and we detected more rhythmic
transcripts through the detrended data (electronic supplementary material, figure S2). In
addition, using the detrended gene expression, we detected extra periodic patterns in
some transcripts that had a significant monotonic trend within the raw gene expression,
which were hard to recognize using the raw data (electronic supplementary material,
figure S3). These findings suggest that the detrend analysis increases the power of
rhythmicity detection methods and thus helps us to get a clearer picture of the periodic
patterns within the gene expression data. The ceemdan detrend was performed with the R
package Rlibeemd version 1.4.2 [45] (set parameters: num_imfs $= 0$; noise_strength $= 0.4$;
S_number $= 0$; num_siftings $= 50$; ensemble_size $= 250$)

(f) Rhythm detection analyses

For *R. leurokolos*, we knew nothing about its biological rhythm, and no golden-standard

circadian or non-circadian genes were available for evaluating the detection power (such

as AUC values and ROC curves) of different algorithms [46, 47]. Thus, we assessed

various rhythm detection methods based on our experiment design (for details see

electronic supplementary material, table S1). Overall, we chose four rhythm detection

algorithms in this study, the mathematic method eigenvalue/pencil [28] (carried out with

customized scripts), the Extended Circadian Harmonic Oscillator application version

4.0.1 [48] (ECHO), and two non-parametric statistical methods RAIN version 1.28.0 [49]

and empirical JTK_CYCLE version 1.0 [50] (eJTK), which were designed to detect

asymmetric waveforms. For eigenvalue/pencil and ECHO, rhythmic transcripts with the

periodicity of 10-14 hours and 22-26 hours were considered as tidal and daily oscillated

[51]. Oscillations with the largest amplitude in one transcript for eigenvalue/pencil were

considered the dominant oscillation [52]. For RAIN and eJTK, rhythmic transcripts with

a period of 12 hours and 24 hours were considered tidal and daily cycling.

Multiple-testing was carried out to adjust *P*-values using the Benjamini-Hochberg

method for RAIN and ECHO, and an empirical method based on the gamma distribution

for combining dependent *P*-values for eJTK. Transcripts with an FDR $\leq$ 0.05 were

considered statistically rhythmic. The root mean square (RMS) amplitude and

Rhythmicity Index (RI) of rhythmic transcripts were calculated by LimoRhyde2 version

0.1.0 [53] and autocorrelation analysis, respectively. We also used ECHO to fit data with

an extended fixed amplitude model and categorize the oscillation type using amplitude

change coefficient (γ) as forced (-0.15 $< \gamma <$ -0.03; amplified amplitude), harmonic (-0.03

$\leq \gamma \leq$ 0.03; steady amplitude), damped (0.03 $< \gamma <$ 0.15; reduced amplitude),

overexpressed ($\gamma \leq$ -0.15), and repressed ($\gamma >$ 0.15), of which the latter two are considered

arrhythmic [48]. The results of rhythm detection analyses are listed in electronic

supplementary material, data S2.

(g) Identification of putative circadian clock genes

We searched for putative circadian clock genes in our *de novo* assembled transcriptome

using the protein Basic Local Alignment Search Tool (BLASTP) carried out by

DIAMOND with a series of known core clock and clock-associated genes as queries

(electronic supplementary material, data S3). For candidate circadian clock transcripts,

sequences were scanned against the coding regions and conserved functional domains

using the InterProScan tools [54]. To determine the sequence homology of candidate

transcripts, we further conducted the phylogenetic analysis. The amino acid sequences
were first aligned using MAFFT version 7.490 [55] and then trimmed using trimAl
version 1.4.1 with the '-automated1' option [56]. The maximum-likelihood trees were
then constructed using the IQ-Tree version 2.1.3 [57] with the ultrafast bootstrap
approximation (UFBoot) and ModelFinder for examining the best-fit model. No
Bmal/*Cycle* transcript was identified in *R. leurokolos* and the *Clock* transcript exhibited
extremely low expression that rendered it unfeasible for rhythm analysis. Overall, we
reconstructed four phylogenetic trees using the Q.insect+Γ4 model for the Clock and
Bmal/Cycle tree, the Q.insect+I+Γ4 model for the Period tree and the Timeless/Timeout
tree, plus the LG+R4 model for the Photolyase/Cryptochrome tree. In the identified
circadian clock genes, the *Clock* transcript exhibited extremely low expression and was
thus excluded from the rhythm detection. Visualization of all phylogenetic trees was
performed with ggtree version 3.8.2 [58].
275

(h) Conserved rhythmic orthologs analysis
The 12-hour rhythmic gene expression data of mice liver and *Drosophila* S2 cell used in
this study were published previously [28, 29]. Together with the *de novo* transcriptome
of *R. leurokolos* assembled in this study, the protein sequences from mouse genome
assembly GCF_000001635 and fruit fly genome assembly GCF_000001215 were used
to construct orthogroups using Orthofinder version 2.5.4 [59] with default settings. The
conserved rhythmic analysis was performed with the single-copy orthologous genes.
Pathway network enrichment analysis of conserved single-copy orthologous genes was
performed with aPEAR version 1.0.0 [60].
285

3. Results and discussion

(a) Transcriptome analysis of deep-sea vent shrimp *R. leurokolos*
To study the potential endogenous rhythms in *R. leurokolos*, we carried out a 72-hour
free-running experiment with 4-hour sampling intervals under constant conditions.
According to previous studies on crustaceans, we worked on the whole cephalothorax
that contained the nervous and visual system where the central clock likely lies, since
the exact location in crustaceans remains unclear [61]. The reference *de novo*
transcriptome contained 90,755 predicted open reading frames (ORFs), which showed a
92% complete BUSCO score. Candidate protein-coding transcripts with low expression
were filtered out, leaving 23,905 transcripts with a 60.4% annotation rate for rhythmic
analyses. A principal components analysis (PCA) of normalized raw gene expression
showed a rough trend from the start of sampling (0 hours) to the end (72 hours, electronic

supplementary material, figure S4). Performing the Mann-Kendall trend test on the raw
dataset, we found that 10,273 transcripts (i.e., 42.97% of the transcriptome) had a
significant monotonic trend (P-value < 0.05). The trend was then removed using
CEEMDAN [43] to produce a steady baseline and enhance the strength of rhythm
detection methods (electronic supplementary material, figure S5). In the PCA results
plotted using detrended gene expression, the trend disappeared and sample points from
different sampling times clustered together (figure 2*a*), indicating the shrimps were
seemingly experiencing the same conditions.
306
(b) General biological rhythms in the temporal transcriptome of *R. leurokolos*
We first used the eigenvalue/pencil [28] and ECHO [48] algorithms to carry out
unbiased identification of any potential periodic oscillations in the gene expression of *R.*
leurokolos. The results uncovered similar rhythmicity that most cycling transcripts were
identified as ultradian rhythms, which oscillated in the tidal range of 10-14 hours, while
much fewer cycling transcripts oscillated in the daily range of 22-26 hours (figure 2*b*).
The eigenvalue/pencil algorithm was able to identify superimposed oscillations from
one cycling transcript. For cycling transcripts with multiple components composed of
tidal and daily oscillations, the amplitude of the tidal component is significantly larger,
suggesting stronger rhythmicity (electronic supplementary material, figure S6). Despite
the high sensitivity of detecting weak ultradian rhythms, the eigenvalue/pencil method
could have high positive false rates [52]. To validate the results, we then detected
cycling transcripts with two robust nonparametric algorithms, RAIN [49] and empirical
JTK_CYCLE [50] (eJTK). Searching for cycling transcripts (FDR < 0.05) with periods
in the range of 8-28 hours at regular intervals of 4 hours, we found the number of
rhythmic transcripts identified by RAIN and eJTK both reached a peak at the 12-hour
period that corresponded to a tidal cycle (electronic supplementary material, figure S7).
Interestingly, ECHO shared many common cycling transcripts with RAIN (60.8% of
transcripts) and eJTK (65.8% of transcripts) but detected numerous extra tidal
transcripts, likely due to the inclusiveness of identifying different harmonics of
oscillating elements [62]. Merging transcripts that were identified as rhythmic by at
least one method, we identified a total of 8,122 rhythmic transcripts (33.98% of the
transcriptome) in the tidal range (figure 2*c*), versus 1,230 rhythmic transcripts (5.15% of
the transcriptome) in the daily range (figure 2*d*). Then we calculated and compared the
root mean square (RMS) amplitude between tidal and daily transcripts, and found the
tidal transcripts had a significantly larger amplitude than daily ones, indicating tidal
cycles contributed more biologically relevant effects (figure 2*e*). The list of rhythmic

transcripts is provided in electronic supplementary material, data S2.

These results demonstrate sustainable endogenous rhythmic gene expression in *R.*
leurokolos under constant conditions. The temporal transcriptome is dominated by the
approximately 12-hour tidal rhythms, which greatly prevail over other oscillations with
different periods, especially the 24-hour daily transcripts that are likely unrelated to the
circadian rhythms.

(c) Patterns of tidal gene expression in *R. leurokolos*
To uncover the biological rhythms ticking in the deep, we then focused on the dominant
tidal rhythms in the temporal transcriptome of *R. leurokolos*. Our 72-hour duration of
sampling covered approximately six tidal cycles, allowing us to test the stability and
robustness of rhythmicity in the tidal gene expression [63]. To capture the amplitude
changes over time, we fitted the gene expression to an extended fixed-amplitude model
using ECHO [48] and categorized the results into five types of oscillation groups (details
see Materials and methods). Near half of the tidal transcripts (n = 3,294) were identified
as damped type, showing a reduced amplitude (figure 3*a*).

Although there was a significantly amplified or reduced amplitude in at least 75% of
tidal rhythmic transcripts, the period and phase of most tidal cycles were seemingly
fixed. We divided the tidal transcriptome into two clusters by gene expression using
K-Means clustering (electronic supplementary material, figures S8*a* and S8*b*). The two
clusters showed a largely antithetical gene expression pattern over time, and transcripts
within each cluster were peaking at the same time, resulting in an almost
complementary phase distribution in one tidal cycle for the two clusters (figures 3*b*).
This bimodal expression pattern is widely observed in the gene expression of
endogenous rhythms. For example, terrestrial organisms such as mice and baboons
experience transcriptional 'rush hours' preceding dawn and dusk, which is considered to
assist in preparing for the impending change in lighting [5, 6]. In the hydrothermal vent,
tidal cycles could strongly affect variations in environmental factors such as
temperature, water chemistry and dissolved oxygen. Vent fauna would thus experience
two opposite environmental conditions in a tidal cycle that one is high temperature, high
sulfide, low oxygen, etc., and another is comparatively low temperature, low sulfide,
high oxygen, etc. In this case, the bimodal tidal expression pattern could help *R.*
leurokolos save energy during the transition periods between acrophase and trough in
the tidal cycle, and make preparations for the upcoming significant environmental

changes [51, 64, 65].

The rhythmicity of 3,837 transcripts in cluster 1 was relatively stable with almost
equivalent amounts of modest forced, harmonic, and damped oscillations, whereas the
2,524 transcripts in cluster 2 were nearly occupied by damped oscillations that
decreased obviously in amplitude, particularly during the second half of experiment
(figure 3*c*). Indeed, there was seemingly a strong correlation between trend directions
and oscillation types (electronic supplementary material, figure S8*c*). In other words, a
tidal transcript with a decreasing expression trend was more likely to have a reduced
amplitude, and vice versa. This likelihood was extremely high in cluster 2 where over
94% of tidal transcripts with a downward trend were also damped. Since we isolated *R.*
leurokolos shrimps from the hydrothermal vent and cultivated them in the lab under
constant conditions for 72 hours, there were unsurprisingly considerable changes in the
gene expression over time corresponding to transcriptional responses to potential
stresses such as decompression. Therefore, the amplitude likely decreases as a
byproduct of declining gene expression levels, especially for most transcripts in cluster
2. Although desynchronization between different tissues of *R. leurokolos* could lead to
arrhythmic gene expression in our free-running study, we still detected a considerable
number of rhythmic transcripts with highly synchronized expression, suggesting the
effect of desynchronization was likely minor.

(d) Wide range of biological processes related to tidal cycles in *R. leurokolos*
The tidal rhythmic transcripts could be categorized by gene expression, oscillation types,
and original trends, all of which included several subgroups that likely reflected similar
or different biological relevance. To search for potential biological functions modulated
by tidal rhythms, we performed functional enrichment analyses over the tidal
transcriptome based on these three classification schemes using clusterProfiler [41].
Gene ontology (GO) and KEGG pathway analyses revealed that tidal rhythms generally
affected various biological processes such as DNA repair, regulation of transcription,
translation, transport, organelle organization, and citrate cycle (figure 3*d*, electronic
supplementary material, figure S9). There was an expected high correlation between the
biological functions of classified subgroups. For example, terms enriched in cluster 2
were more or less the same as those in the damped or downward subgroup. Notably,
rhythmic processes related to the nucleus such as chromatin organization, DNA
biosynthetic process, and transcription were uniquely enriched in cluster 1; whereas
rhythmic processes supported by the cytoplasm such as translation, protein folding,

vesicle-mediated transport, and oxidative phosphorylation were highly enriched in cluster 2. Thus, it is likely that *R. leurokolos* shrimps can organize various rhythmic biological processes in terms of spatial and temporal expression patterns to coordinate their internal physiology, regardless of the presence of clear zeitgebers.

In hydrothermal vents, particularly near the fluid orifice where *R. leurokolos* inhabits, hot, anoxic fluids with abundant potentially toxic chemical substances (e.g., hydrogen sulfide) mix dramatically with the cold surrounding seawater, resulting in an extreme and variable environment. Vent organisms have evolved various adaptive strategies at morphological, physiological, and biochemical levels [18]. We hypothesized that intrinsic rhythms could help vent animals schedule their activities to cope with the harsh environment. As expected, we found many tidally oscillating transcripts were related to various cellular stress response processes such as sulfur metabolism, reactive oxygen species (ROS) homeostasis, cellular detoxification, heat and hypoxia response and innate immune response (electronic supplementary material, figures S10*a-d*). The representative tidal genes related to stress responses include *Sqor*, *Cat*, *Sod*, *Gst*, *Gpx*, *Egln1*, *Dnaja1*, *Hsp90*, *Ctsl*, among others. Most of these transcripts showed damped rhythmic expression over time under constant conditions, suggesting that *in situ* environmental cues were necessary for the maintenance and synchronization of rhythmic processes related to stress responses. The results reveal broad tidal regulation of gene expression in the cellular response to various environmental stresses, suggesting an adaptive advantage for *R. leurokolos* living in harsh, fluctuating environments.

Mounting evidence suggests the presence of cross-talking between circadian clocks and the cell cycle [66, 67]. Here, we found the annotation of many tidal transcripts was related to the cell cycle (electronic supplementary material, figures S11*a-c*). The transition between the G1 and S phase is controlled and regulated by key genes involved in the checkpoint pathways [67-69], of which several transcripts were tidally oscillating, including *Ccnd2*, *TP53*, *Mycbp2*, *Rbl1*, *E2f2*, *Tipin*, *Mdm2*, among others. In addition, genes that encoded major components of the RAD9-RAD1-HUS1 clamp complex (e.g., *Rad9a*, *Rad1*, *Hus1*), which play pivotal roles in DNA repair, also oscillated in the tidal range [70]. As crucial elements of the spindle assembly checkpoint (SAC), genes such as *Mad2*, *Bub1*, and *Bub3* that regulate chromosome separation were also found to be tidally cycling [71]. Taken together, tidal rhythms appear to generally regulate the progression of the cell cycle in *R. leurokolos* to cope with DNA damage derived from periodic intrinsic and environmental stresses.

(e) Possible molecular basis and zeitgebers of biological rhythms in *R. leurokolos*

We searched for and identified *R. leurokolos* homologs of the canonical circadian clock genes that form the molecular oscillator in the fruitfly *Drosophila melanogaster*, including *Clock (Clk)*, *Period (Per)*, *Timeless (Tim)*, *Cryptochrome 2 (Cry2)*, *Clockwork orange (Cwo)*, *Doubletime (Dbt)*, *Vrille (Vri)*, and *Pdp1ε* (electronic supplementary material, figures S12*a-d* and data S3). Indeed, all identified *R. leurokolos* circadian clock gene homologs lacked 24-hour rhythmicity (figure 4). This may provide intrinsic fitness for these hydrothermal vent shrimps that are likely completely isolated from the surface as adults [72]. Moreover, most clock genes showed a lack of rhythmicity in the tidal range as well. Although the transcriptional repressor *Period* and *Doubletime*, which encodes a kinase that phosphorylates the PER protein [73], were statistically oscillated in the tidal range, the Period transcript showed an opposite expression with the Timeless transcript in the second half of the free-running experiment, which was abnormal for the circadian clock regulated by the TTFL mechanism. Since a robust endogenous clock is required to be free running with a self-sustained period for maintaining the rhythmicity of tidal oscillations in *R. leurokolos*, we hypothesize that there is a circatidal clock with at least part of independent molecular components from the circadian clocks. Separate circadian and circatidal clocks were also reported in the intertidal isopod *Eurydice pulchra* [74] and mangrove cricket *Apteronemobius asahinai* [75]. Nevertheless, the core clock genes still encode functional proteins with conserved domains (electronic supplementary material, figure S12*e*), and we cannot exclude the possibility that some core circadian clock genes are necessary for tidal rhythms [76, 77], even though they lose free-running transcription rhythmicity. These circadian clock genes may also be useful during the dispersal of planktotrophic larvae, which may take place in the photic zone [78].

To precisely synchronize with the tidal cycle, we consider *R. leurokolos* shrimps depend on an endogenous circatidal clock to integrate all potential periodic cues. In the microhabitat where *R. leurokolos* lives, crucial environmental factors such as temperature, water chemistry, and dissolved oxygen can vary irregularly and dramatically in a very short time, generating noisy signals. Therefore, although these environmental factors are crucial for the survival of hot vent shrimps, they are likely poor periodic cues. Hydrostatic pressure and tidal currents, however, are largely in line with the standard of a synchronization cue [79, 80]. In addition, they show significant coherence with temperature, water chemistry, and dissolved oxygen [81]. The closer the

distance from hot vents, the stronger of modulation of tidal pressure on environmental
factors [17]. Therefore, we propose hydrostatic pressure as the most likely
synchronization cue for *R. leurokolos*. Over longer time scales the overall levels of
venting activity is known to be modulated by the tidal cycle. As such, by synchronizing
to pressure, it is possible for hot vent shrimps to predict major environmental
fluctuations. In addition to pressure and currents, light emissions from vents [82], food
sources [32] or even soundscape cues from vents [83] could be potentially useful
periodic cues for *R. leurokolos*. Nevertheless, how exactly hot vent shrimps synchronize
their endogenous rhythms with environmental cycles remains a question for the future.
487
(f) Rhythmicity comparison of two vent species living in different
microenvironments
To compare the tidal rhythms in the deep-sea hydrothermal vent organisms, we applied
our transcriptome analysis and rhythm detection pipeline to the published *in situ* gene
expression dataset of the Atlantic deep-sea vent mussel *B. azoricus* [22]. Despite the
different geographic localities of the two species, *B. azoricus* and *R. leurokolos* both
show similar biological rhythms predominated by a tidal pattern that is related to the
responses to environmental stresses such as DNA damage, and thermal, immune, and
oxidative stress, suggesting a generally temporal adaptation to the harsh deep-sea
hydrothermal environments.
498
Compared to *R. leurokolos*, however, the proportion of tidal transcripts in the
transcriptome was lower in *B. azoricus* regardless of the detection algorithm used
(electronic supplementary material, figure S13*a* and S13*b*). Indeed, considering that
rhythmic gene expression was organ-specific [5], the result was disputable as we
detected overall rhythmicity within mixed tissues of *R. leurokolos*, whereas the study on
B. azoricus only worked on gills. We further examined and compared the strength of
tidal rhythms in the two species by calculating the rhythmicity index (RI) using the
autocorrelation analysis, and the results suggested the general rhythmicity of tidal gene
expression was greater in *R. leurokolos* (electronic supplementary material, figure S13*c*).
This is unexpected since it should be easier for *B. azoricus*, living in a relatively distant
area from the vent orifice with stable diffuse flow, to detect environmental factors
influenced by tidal cycles. A possible explanation is that *R. leurokolos* needs to depend
on a strong pacemaker to integrate all the periodic cues for predicting environmental
fluctuations and preparing for transcriptional regulation, whereas *B. azoricus* could
schedule the transcription by either relying on an endogenous clock or responding

directly to the tidal-modulated environmental cues [22]. Why *R. leurokolos* inhabiting a more dynamic environment shows more robust rhythmicity compared to *B. azoricus* is an intriguing topic warranting future studies, although this could also be related to intrinsic differences between mussels and shrimps, as well as the influence of chemosymbiosis on the host [84].

(g) The potential evolutionary conservation between marine tidal rhythms and terrestrial 12-hour rhythms

We show that the deep-sea vent shrimp *R. leurokolos* exhibits robust endogenous rhythms related to tidal effects instead of daily cycles, making it a useful model for testing the hypothesis that marine tidal rhythms and 12-hour rhythms of terrestrial animals share conserved features. To investigate this, we compared the tidal gene expression of *R. leurokolos* with published ~12-hour rhythms in the *Bmal1* knockout mice liver [28] and *Drosophila* S2 cells [29], two systems without canonical circadian clock gene expressions. A total of 406 single-copy orthologous genes were identified as showing ~12-hour rhythms in these three species. The vent shrimp *Rimicaris* shared more genes with the fruit fly compared with the mouse (54 genes vs 36 genes), suggesting a phylogenetic relationship. We also found 21 genes with a period of ~12-hour rhythms in all three species (figure 5*a*). These genes were enriched in the processes of DNA replication, regulation of transcription, ribosome biogenesis, protein folding, and trans-Golgi network, which are also highly enriched in the ~12-hour rhythms of human beings [85] (figure 5*b*). The heat shock protein 70 cochaperone *Dnaja1*, which was also reported in 12-hour rhythms of mammals [85] and tidal rhythms of the sea anemone *Aiptasia diaphana* [84], also showed a robust free-running tidal rhythm in *R. leurokolos*. These results support the evolutionary conservation of ~12-hour gene expression related to the processes of central dogma information flow in both terrestrial and marine animals across a broad range of phyla, indicating the ~12-hour rhythms in animals overall likely initially evolved for adjusting to the tidal cycle.

4. Conclusion

In this study, our analyses provide evidence for endogenous biological rhythms in the deep-sea hydrothermal vent shrimps *R. leurokolos* through a 72-hour free-running experiment and constrain the possible zeitgeber for biological clocks ticking in the deep, shedding light on biological rhythms in the light-absent biosphere (figure 6). We propose that rhythms with a period of 24 hours driven by the conventional circadian

clock largely decrease or even disappear in adult deep-sea hydrothermal vent fauna.
Instead, rhythmic gene expression coordinated with approximately 12-hour tidal cycles,
which are likely controlled by an independent circatidal clock, evolved as a temporal
adaptation to the environmental fluctuations in hydrothermal ecosystems. Independent
of solar energy but being influenced by tidal cycles, organisms in the deep-sea
hydrothermal vent are likely powerful models for chronobiologists to decipher the
evolution and molecular mechanisms of circatidal clocks and ~12-hour rhythms,
important across a wide range of living organisms.
558

Data accessibility

560 Raw RNA-Seq data are available on NCBI as BioProject PRJNA993859. The *de novo*
561 reference transcriptome of *Rimicaris leurokolos* is available in figshare
562 (https://doi.org/10.6084/m9.figshare.23667327). All original codes used in this work are
563 available at GitHub (https://github.com/belphe/Shinkaicaris).
564

Author's contributions

566 Conceptualization, J.S., T.Y., N.M., and C.C.; Methodology, T.Y., N.M., C.C., and
567 H.Y.Z.; Formal Analysis, H.Y.Z. and J.S.; Investigation, T.Y., C.C.; Writing – Original
568 Draft, H.Y.Z.; Writing – Review & Editing, J.S., C.C.; Visualization, H.Y.Z. and Q.Q.J.;
569 Supervision, J.S. and T.Y.; Project Administration, J.S., P-Y.Q. and T.Y.; Funding
570 Acquisition, J.S. and P-Y.Q.
571

Competing interests

573 The authors declare that they have no competing interests.
574

Funding

This work was supported by: Science and Technology Innovation Project of Laoshan
Laboratory (LSKJ202203104); Fundamental Research Funds for the Central
Universities (202172002 and 202241002); Young Taishan Scholars Program of
Shandong Province (tsqn202103036); Natural Science Foundation of Shandong
Province (ZR2023JQ014); Principal investigator projects of the Southern Marine
Science and Engineering Guangdong Laboratory (Guangzhou) (2021HJ01); Major
Basic and Applied Research Projects of Guangdong Province (2019B030302004-04);

587 Acknowledgements

We are grateful to the ship captain and crews of R/V *KAIYO* as well as pilots and the
technical team of ROV *Hyper-Dolphin* for their great efforts during sample collection
during cruise KY14-01 led by Ken Takai (JAMSTEC). Hiroyuki Yamamoto, Shinsuke
Kawagucci, and Hiromi K. Watanabe (JAMSTEC) are thanked for providing raw data
of the temperature probe recordings published in a previous study [31]. We also thank
Bokai Zhu (University of Pittsburgh) for his insightful advice on an earlier version of
this manuscript.
595

References

1. Kondo, T., Mori, T., Lebedeva, N. V., Aoki, S., Ishiura, M., Golden, S. S. 1997 Circadian rhythms in rapidly dividing cyanobacteria. *Science*. **275**, 224-227. (doi:10.1126/science.275.5297.224)

2. Panda, S., Hogenesch, J. B., Kay, S. A. 2002 Circadian rhythms from flies to human. *Nature*. **417**, 329-335. (doi:10.1038/417329a)

3. Edgar, R. S., Green, E. W., Zhao, Y., van Ooijen, G., Olmedo, M., Qin, X., Xu, Y., Pan, M., Valekunja, U. K., Feeney, K. A*., et al.* 2012 Peroxiredoxins are conserved markers of circadian rhythms. *Nature*. **485**, 459-464. (doi:10.1038/nature11088)

4. Stoleru, D., Peng, Y., Agosto, J., Rosbash, M. 2004 Coupled oscillators control morning and evening locomotor behaviour of *Drosophila*. *Nature*. **431**, 862-868. (doi:10.1038/nature02926)

5. Zhang, R., Lahens, N. F., Ballance, H. I., Hughes, M. E., Hogenesch, J. B. 2014 A circadian gene expression atlas in mammals: Implications for biology and medicine. *Proc. Natl Acad. Sci. USA*. **111**, 16219-16224. (doi:10.1073/pnas.1408886111)

6. Mure, L. S., Le, H. D., Benegiamo, G., Chang, M. W., Rios, L., Jillani, N., Ngotho, M., Kariuki, T., Dkhissi-Benyahya, O., Cooper, H. M*., et al.* 2018 Diurnal transcriptome atlas of a primate across major neural and peripheral tissues. *Science*. **359**, eaao0318. (doi:10.1126/science.aao0318)

7. Patke, A., Young, M. W., Axelrod, S. 2020 Molecular mechanisms and physiological importance of circadian rhythms. *Nat. Rev. Mol. Cell Biol.* **21**, 67-84. (doi:10.1038/s41580-019-0179-2)

8. Beale, A. D., Whitmore, D., Moran, D. 2016 Life in a dark biosphere: a review of circadian physiology in "arrhythmic" environments. *J. Comp. Physiol. B*. **186**, 947-968. (doi:10.1007/s00360-016-1000-6)

9. Abhilash, L., Shindey, R., Sharma, V. K. 2017 To be or not to be rhythmic? A review of studies on organisms inhabiting constant environments. *Biol. Rhythm Res.* **48**, 677-691. (doi:10.1080/09291016.2017.1345426)

10. Ramirez-Llodra, E., Brandt, A., Danovaro, R., De Mol, B., Escobar, E., German, C. R., Levin, L. A., Martinez Arbizu, P., Menot, L., Buhl-Mortensen, P*., et al.* 2010 Deep, diverse and definitely different: unique attributes of the world's largest ecosystem. *Biogeosciences*. **7**, 2851-2899. (doi:10.5194/bg-7-2851-2010)

11. Aguzzi, J., Costa, C., Menesatti, P., Garcia, J. A., Bahamon, N., Puig, P., Sarda, F. 2011 Activity rhythms in the deep-sea: a chronobiological approach. *Front. Biosci. (Landmark Ed)*. **16**, 131-150. (doi:10.2741/3680)

12. Warrant, E. J., Locket, N. A. 2004 Vision in the deep sea. *Biol. Rev. Camb. Philos.*
Soc. **79**, 671-712. (doi:10.1017/S1464793103006420)

13. Aguzzi, J., Costa, C., Furushima, Y., Chiesa, J. J., Company, J. B., Menesatti, P.,
Iwase, R., Fujiwara, Y. 2010 Behavioral rhythms of hydrocarbon seep fauna in
relation to internal tides. *Mar. Ecol. Prog. Ser.* **418**, 47-56. (doi:10.3354/meps08835)

14. Chevaldonné, P., Desbruyères, D., Le Haître, M. 1991 Time-series of temperature
from three deep-sea hydrothermal vent sites. *Deep Sea Res. Part A Oceanogr. Res.*
Pap. **38**, 1417-1430. (doi:10.1016/0198-0149(91)90014-7)

15. Tivey, M. K., Bradley, A. M., Joyce, T. M., Kadko, D. 2002 Insights into tide-related
variability at seafloor hydrothermal vents from time-series temperature
measurements. *Earth. Planet. Sci. Lett.* **202**, 693-707.
(doi:10.1016/S0012-821X(02)00801-4)

16. Crone, T. J., Wilcock, W. S. D. 2005 Modeling the effects of tidal loading on
mid-ocean ridge hydrothermal systems. *Geochem. Geophys. Geosyst.* **6**,
(doi:10.1029/2004GC000905)

17. Barreyre, T., Escartín, J., Sohn, R. A., Cannat, M., Ballu, V., Crawford, W. C. 2014
Temporal variability and tidal modulation of hydrothermal exit-fluid temperatures at
the Lucky Strike deep-sea vent field, Mid-Atlantic Ridge. *J. Geophys. Res. Solid*
Earth. **119**, 2543-2566. (doi:10.1002/2013JB010478)

18. Childress, J. J., Fisher, C. R. 1992 The biology of hydrothermal vent animals:
physiology, biochemistry, and autotrophic symbioses. *Oceanogr. Mar. Biol. Annu.*
Rev. **30**, 337-441.

19. Nedoncelle, K., Lartaud, F., de Rafelis, M., Boulila, S., Le Bris, N. 2013 A new
method for high-resolution bivalve growth rate studies in hydrothermal environments.
Mar. Biol. **160**, 1427-1439. (doi:10.1007/s00227-013-2195-7)

20. Cuvelier, D., Legendre, P., Laes, A., Sarradin, P.-M., Sarrazin, J. 2014 Rhythms and
community dynamics of a hydrothermal tubeworm assemblage at main Endeavour
field–a multidisciplinary deep-sea observatory approach. *PLoS One.* **9**, e96924.
(doi:10.1371/journal.pone.0096924)

21. Cuvelier, D., Legendre, P., Laës-Huon, A., Sarradin, P. M., Sarrazin, J. 2017
Biological and environmental rhythms in (dark) deep-sea hydrothermal ecosystems.
Biogeosciences. **14**, 2955-2977. (doi:10.5194/bg-14-2955-2017)

22. Mat, A. M., Sarrazin, J., Markov, G. V., Apremont, V., Dubreuil, C., Eché, C.,
Fabioux, C., Klopp, C., Sarradin, P.-M., Tanguy, A., *et al.* 2020 Biological rhythms
in the deep-sea hydrothermal mussel *Bathymodiolus azoricus*. *Nat. Commun.* **11**,
3454. (doi:10.1038/s41467-020-17284-4)

23. Furushima, Y., Higashi, H., Koshikawa, H., Furuichi, N., Yamamoto, H. Year
Periodic behaviors of deep-ocean flow and turbulent mixing in hydrothermal field on
the Okinawa Trough, Japan. OCEANS 2016 MTS/IEEE Monterey; 2016 19-23
Sept. 2016; 2016. p. 1-5.

24. Methou, P., Chen, C., Komai, T. 2024 Revision of the alvinocaridid shrimp genus
Rimicaris Williams & Rona, 1986 (Decapoda: Caridea) with description of a new
species from the Mariana Arc hydrothermal vents. *Zootaxa*. **5406**, 501-518.
(doi:10.11646/ZOOTAXA.5406.4.1)

25. Watanabe, H., Kojima, S. 2015 Vent fauna in the Okinawa Trough. In *Subseafloor*
biosphere linked to hydrothermal systems: TAIGA concept. (ed.^eds. pp. 449-459

26. Yahagi, T., Watanabe, H., Ishibashi, J., Kojima, S. 2015 Genetic population structure
of four hydrothermal vent shrimp species (Alvinocarididae) in the Okinawa Trough,
Northwest Pacific. *Mar. Ecol. Prog. Ser.* **529**, 159-169. (doi:10.3354/meps11267)

27. Van Audenhaege, L., Matabos, M., Brind'Amour, A., Drugmand, J., Laës-Huon, A.,
Sarradin, P.-M., Sarrazin, J. 2022 Long-term monitoring reveals unprecedented
stability of a vent mussel assemblage on the Mid-Atlantic Ridge. *Prog. Oceanogr.*
204, 102791. (https://doi.org/10.1016/j.pocean.2022.102791)

28. Zhu, B., Zhang, Q., Pan, Y., Mace, E. M., York, B., Antoulas, A. C., Dacso, C. C.,
O'Malley, B. W. 2017 A cell-autonomous mammalian 12 hr clock coordinates
metabolic and stress rhythms. *Cell Metab.* **25**, 1305-1319.e1309.
(doi:10.1016/j.cmet.2017.05.004)

29. Zhu, B., Liu, S. 2023 Preservation of ~12-h ultradian rhythms of gene expression of
mRNA and protein metabolism in the absence of canonical circadian clock. *Front.*
Physiol. **14**, (doi:10.3389/fphys.2023.1195001)

30. Zhu, B., Dacso, C. C., O'Malley, B. W. 2018 Unveiling "Musica Universalis" of the
cell: a brief history of biological 12-hour rhythms. *J. Endocr. Soc.* **2**, 727-752.
(doi:10.1210/js.2018-00113)

31. Kawagucci, S., Miyazaki, J., Nakajima, R., Nozaki, T., Takaya, Y., Kato, Y., Shibuya,
T., Konno, U., Nakaguchi, Y., Hatada, K.*, et al.* 2013 Post-drilling changes in fluid
discharge pattern, mineral deposition, and fluid chemistry in the Iheya North
hydrothermal field, Okinawa Trough. *Geochem. Geophys. Geosyst.* **14**, 4774-4790.
(doi:10.1002/2013GC004895)

32. Matabos, M., Cuvelier, D., Brouard, J., Shillito, B., Ravaux, J., Zbinden, M.,
Barthelemy, D., Sarradin, P. M., Sarrazin, J. 2015 Behavioural study of two
hydrothermal crustacean decapods: *Mirocaris fortunata* and *Segonzacia mesatlantica*,
from the Lucky Strike vent field (Mid-Atlantic Ridge). *Deep Sea Res. II: Top. Stud.*
Oceanogr. **121**, 146-158. (doi:10.1016/j.dsr2.2015.04.008)

33. Stephan, F. K. 2002 The "other" circadian system: food as a Zeitgeber. *J. Biol.*
Rhythms. **17**, 284-292. (doi:10.1177/074873040201700402)

34. Bolger, A. M., Lohse, M., Usadel, B. 2014 Trimmomatic: a flexible trimmer for
Illumina sequence data. *Bioinformatics*. **30**, 2114-2120.
(doi:10.1093/bioinformatics/btu170)

35. Wood, D. E., Lu, J., Langmead, B. 2019 Improved metagenomic analysis with
Kraken 2. *Genome Biol.* **20**, 257. (doi:10.1186/s13059-019-1891-0)

36. Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I.,
Adiconis, X., Fan, L., Raychowdhury, R., Zeng, Q.*, et al.* 2011 Full-length
transcriptome assembly from RNA-Seq data without a reference genome. *Nat.*
Biotechnol. **29**, 644-652. (doi:10.1038/nbt.1883)

37. Patro, R., Duggal, G., Love, M. I., Irizarry, R. A., Kingsford, C. 2017 Salmon
provides fast and bias-aware quantification of transcript expression. *Nat. Methods*. **14**,
417-419. (doi:10.1038/nmeth.4197)

38. Robinson, M. D., Oshlack, A. 2010 A scaling normalization method for differential
expression analysis of RNA-seq data. *Genome Biol.* **11**, R25.
(doi:10.1186/gb-2010-11-3-r25)

39. Buchfink, B., Reuter, K., Drost, H.-G. 2021 Sensitive protein alignments at
tree-of-life scale using DIAMOND. *Nat. Methods*. **18**, 366-368.
(doi:10.1038/s41592-021-01101-x)

40. Conesa, A., Götz, S., García-Gómez, J. M., Terol, J., Talón, M., Robles, M. 2005
Blast2GO: a universal tool for annotation, visualization and analysis in functional
genomics research. *Bioinformatics*. **21**, 3674-3676.
(doi:10.1093/bioinformatics/bti610)

41. Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., Zhan,
L.*, et al.* 2021 clusterProfiler 4.0: A universal enrichment tool for interpreting omics
data. *Innovation*. **2**, 100141. (doi:10.1016/j.xinn.2021.100141)

42. Love, M. I., Huber, W., Anders, S. 2014 Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550.
(doi:10.1186/s13059-014-0550-8)

43. Torres, M. E., Colominas, M. A., Schlotthauer, G., Flandrin, P. Year A complete
ensemble empirical mode decomposition with adaptive noise. 2011 IEEE
international conference on acoustics, speech and signal processing (ICASSP); 2011:
IEEE; 2011. p. 4144-4147.

44. Ayenu-Prah, A., Attoh-Okine, N. 2010 A criterion for selecting relevant intrinsic
mode functions in empirical mode decomposition. *Adv. Adapt. Data Anal.* **2**, 1-24.

(doi:10.1142/S1793536910000367)

45. Luukko, P. J. J., Helske, J., Räsänen, E. 2016 Introducing libeemd: a program
package for performing the ensemble empirical mode decomposition. *Comput. Stat.*
31, 545-557. (doi:10.1007/s00180-015-0603-9)

46. Laloum, D., Robinson-Rechavi, M. 2020 Methods detecting rhythmic gene
expression are biologically relevant only for strong signal. *PLoS Comput. Biol.* **16**,
e1007666. (doi:10.1371/journal.pcbi.1007666)

47. Mei, W., Jiang, Z., Chen, Y., Chen, L., Sancar, A., Jiang, Y. 2020 Genome-wide
circadian rhythm detection methods: systematic evaluations and practical guidelines.
Brief. Bioinform. **22**, bbaa135. (doi:10.1093/bib/bbaa135)

48. De los Santos, H., Collins, E. J., Mann, C., Sagan, A. W., Jankowski, M. S., Bennett,
K. P., Hurley, J. M. 2019 ECHO: an application for detection and analysis of
oscillators identifies metabolic regulation on genome-wide circadian output.
Bioinformatics. **36**, 773-781. (doi:10.1093/bioinformatics/btz617)

49. Thaben, P. F., Westermark, P. O. 2014 Detecting rhythms in time series with RAIN. *J.*
Biol. Rhythms. **29**, 391-400. (doi:10.1177/0748730414553029)

50. Hutchison, A. L., Maienschein-Cline, M., Chiang, A. H., Tabei, S. M. A., Gudjonson,
H., Bahroos, N., Allada, R., Dinner, A. R. 2015 Improved statistical methods enable
greater sensitivity in rhythm detection for genome-wide data. *PLoS Comput. Biol.* **11**,
e1004094. (doi:10.1371/journal.pcbi.1004094)

51. Connor, K. M., Gracey, A. Y. 2011 Circadian cycles are the dominant transcriptional
rhythm in the intertidal mussel *Mytilus californianus*. *Proc. Natl Acad. Sci. USA.* **108**,
16110. (doi:10.1073/pnas.1111076108)

52. Pan, Y., Ballance, H., Meng, H., Gonzalez, N., Kim, S.-M., Abdurehman, L., York, B.,
Chen, X., Schnytzer, Y., Levy, O.*, et al.* 2020 12-h clock regulation of genetic
information flow by XBP1s. *PLoS Biol.* **18**, e3000580.
(doi:10.1371/journal.pbio.3000580)

53. Obodo, D., Outland, E. H., Hughey, J. J. 2023 LimoRhyde2: genomic analysis of
biological rhythms based on effect sizes. *PLoS One.* **18**, e0292089.
(doi:10.1371/journal.pone.0292089)

54. Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H.,
Maslen, J., Mitchell, A., Nuka, G.*, et al.* 2014 InterProScan 5: genome-scale protein
function classification. *Bioinformatics.* **30**, 1236-1240.
(doi:10.1093/bioinformatics/btu031)

55. Katoh, K., Standley, D. M. 2013 MAFFT multiple sequence alignment software
version 7: improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772-780.

(doi:10.1093/molbev/mst010)

56. Capella-Gutiérrez, S., Silla-Martínez, J. M., Gabaldón, T. 2009 trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*. **25**, 1972-1973. (doi:10.1093/bioinformatics/btp348)

57. Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., von Haeseler, A., Lanfear, R. 2020 IQ-TREE 2: new models and efficient methods for phylogenetic inference in the genomic era. *Mol. Biol. Evol.* **37**, 1530-1534. (doi:10.1093/molbev/msaa015)

58. Yu, G., Smith, D. K., Zhu, H., Guan, Y., Lam, T. T.-Y. 2017 ggtree: an r package for visualization and annotation of phylogenetic trees with their covariates and other associated data. *Methods Ecol. Evol.* **8**, 28-36. (doi:10.1111/2041-210X.12628)

59. Emms, D. M., Kelly, S. 2019 OrthoFinder: phylogenetic orthology inference for comparative genomics. *Genome Biol.* **20**, 238. (doi:10.1186/s13059-019-1832-y)

60. Kerseviciute, I., Gordevicius, J. 2023 aPEAR: an R package for autonomous visualization of pathway enrichment networks. *Bioinformatics*. **39**, (doi:10.1093/bioinformatics/btad672)

61. Strauss, J., Dircksen, H. 2010 Circadian clocks in crustaceans: identified neuronal and cellular systems. *Front. Biosci.* **15**, 1040-1074. (doi:10.2741/3661)

62. De los Santos, H., Collins, E. J., Hurley, J. M., Bennett, K. P. Year Circadian rhythms in *Neurospora* exhibit biologically relevant driven and damped harmonic oscillations. Proceedings of the 8th ACM International Conference on Bioinformatics, Computational Biology, and Health Informatics; 2017; 2017. p. 455-463.

63. Yamazaki, S., Numano, R., Abe, M., Hida, A., Takahashi, R.-i., Ueda, M., Block, G. D., Sakaki, Y., Menaker, M., Tei, H. 2000 Resetting central and peripheral circadian oscillators in transgenic rats. *Science*. **288**, 682-685. (doi:10.1126/science.288.5466.682)

64. Gracey, A. Y., Chaney, M. L., Boomhower, J. P., Tyburczy, W. R., Connor, K., Somero, G. N. 2008 Rhythms of gene expression in a fluctuating intertidal environment. *Curr. Biol.* **18**, 1501-1507. (doi:10.1016/j.cub.2008.08.049)

65. Satoh, A., Terai, Y. 2019 Circatidal gene expression in the mangrove cricket *Apteronemobius asahinai*. *Sci. Rep.* **9**, 3719. (doi:10.1038/s41598-019-40197-2)

66. Hunt, T., Sassone-Corsi, P. 2007 Riding tandem: circadian clocks and the cell cycle. *Cell*. **129**, 461-464. (doi:10.1016/j.cell.2007.04.015)

67. Farshadi, E., van der Horst, G. T. J., Chaves, I. 2020 Molecular links between the circadian clock and the cell cycle. *J. Mol. Biol.* **432**, 3515-3524. (doi:10.1016/j.jmb.2020.04.003)

68. Yoshizawa-Sugata, N., Masai, H. 2007 Human Tim/Timeless-interacting protein,
Tipin, is required for efficient progression of S phase and DNA replication
checkpoint. *J. Biol. Chem.* **282**, 2729-2740. (doi:10.1074/jbc.M605596200)

69. Masri, S., Cervantes, M., Sassone-Corsi, P. 2013 The circadian clock and cell cycle:
interconnected biological circuits. *Curr. Opin. Cell Biol.* **25**, 730-734.
(doi:10.1016/j.ceb.2013.07.013)

70. Bermudez, V. P., Lindsey-Boltz, L. A., Cesare, A. J., Maniwa, Y., Griffith, J. D.,
Hurwitz, J., Sancar, A. 2003 Loading of the human 9-1-1 checkpoint complex onto
DNA by the checkpoint clamp loader hRad17-replication factor C complex *in vitro*.
Proc. Natl Acad. Sci. USA. **100**, 1633-1638. (doi:10.1073/pnas.0437927100)

71. Klebig, C., Korinth, D., Meraldi, P. 2009 Bub1 regulates chromosome segregation in
a kinetochore-independent manner. *J. Cell Biol.* **185**, 841-858.
(doi:10.1083/jcb.200902128)

72. Methou, P., Chen, C., Kayama Watanabe, H., Cambon, M.-A., Pradillon, F. 2022
Reproduction in deep-sea vent shrimps is influenced by diet, with rhythms apparently
unlinked to surface production. *Ecol. Evol.* **12**, e9076. (doi:10.1002/ece3.9076)

73. Parnell, A. A., De Nobrega, A. K., Lyons, L. C. 2021 Translating around the clock:
Multi-level regulation of post-transcriptional processes by the circadian clock. *Cell.*
Signal. **80**, 109904. (doi:10.1016/j.cellsig.2020.109904)

74. Zhang, L., Hastings, Michael H., Green, Edward W., Tauber, E., Sladek, M., Webster,
Simon G., Kyriacou, Charalambos P., Wilcockson, David C. 2013 Dissociation of
circadian and circatidal timekeeping in the marine crustacean *Eurydice pulchra*. *Curr.*
Biol. **23**, 1863-1873. (doi:10.1016/j.cub.2013.08.038)

75. Takekata, H., Numata, H., Shiga, S., Goto, S. G. 2014 Silencing the circadian clock
gene Clock using RNAi reveals dissociation of the circatidal clock from the circadian
clock in the mangrove cricket. *J. Insect Physiol.* **68**, 16-22.
(doi:10.1016/j.jinsphys.2014.06.012)

76. Lin, Z., Green, E. W., Webster, S. G., Hastings, M. H., Wilcockson, D. C., Kyriacou,
C. P. 2023 The circadian clock gene *bmal1* is necessary for co-ordinated circatidal
rhythms in the marine isopod *Eurydice pulchra* (Leach). *PLoS Genet.* **19**, e1011011.
(doi:10.1371/journal.pgen.1011011)

77. Kwiatkowski, E. R., Schnytzer, Y., Rosenthal, J. J., Emery, P. 2023 Behavioral
circatidal rhythms require *Bmal1* in *Parhyale hawaiensis*. *Curr. Biol.* **33**, 1867-1882.
e1865. (doi:10.1016/j.cub.2023.03.015)

78. Yahagi, T., Kayama Watanabe, H., Kojima, S., Kano, Y. 2017 Do larvae from
deep-sea hydrothermal vents disperse in surface waters? *Ecology*. **98**, 1524-1534.
(doi:10.1002/ecy.1800)

79. Reid, D. G., Naylor, E. 1990 Entrainment of bimodal circatidal rhythms in the shore
crab *Carcinus maenas*. *J. Biol. Rhythms*. **5**, 333-347.
(doi:10.1177/074873049000500405)

80. Akiyama, T. 2004 Entrainment of the circatidal swimming activity rhythm in the
cumacean Dimorphostylis asiatica (Crustacea) to 12.5-hour hydrostatic pressure
cycles. *Zoolog. Sci.* **21**, 29-38, 10.
(doi:10.2108/0289-0003(2004)21[29:EOTCSA]2.0.CO;2)

81. Mittelstaedt, E., Fornari, D. J., Crone, T. J., Kinsey, J., Kelley, D., Elend, M. 2016
Diffuse venting at the ASHES hydrothermal field: Heat flux and tidally modulated
flow variability derived from in situ time-series measurements. *Geochem. Geophys.*
Geosyst. **17**, 1435-1453. (doi:10.1002/2015GC006144)

82. Van Dover, C. L., Reynolds, G. T., Chave, A. D., Tyson, J. A. 1996 Light at deep-sea
hydrothermal vents. *Geophys. Res. Lett.* **23**, 2049-2052. (doi:10.1029/96GL02151)

83. Crone, T. J., Wilcock, W. S. D., Barclay, A. H., Parsons, J. D. 2006 The sound
generated by mid-ocean ridge black smoker hydrothermal vents. *PLoS One*. **1**, e133.
(doi:10.1371/journal.pone.0000133)

84. Sorek, M., Schnytzer, Y., Waldman Ben-Asher, H., Caspi, V. C., Chen, C.-S., Miller,
D. J., Levy, O. 2018 Setting the pace: host rhythmic behaviour and gene expression
patterns in the facultatively symbiotic cnidarian *Aiptasia* are determined largely by
Symbiodinium. *Microbiome*. **6**, 83. (doi:10.1186/s40168-018-0465-9)

85. Zhu, B., Liu, S., David, N. L., Dion, W., Doshi, N. K., Siegel, L. B., Amorim, T.,
Andrews, R. E., Kumar, G. N., Li, H*., et al.* 2023 Evidence for conservation of
primordial ~12-hour ultradian gene programs in humans under free-living conditions.
bioRxiv. 2023.2005.2002.539021. (doi:10.1101/2023.05.02.539021)

86. Zhu, B., Liu, S., David, N. L., Dion, W., Doshi, N. K., Siegel, L. B., Amorim, T.,
Andrews, R. E., Kumar, G. V. N., Li, H*., et al.* 2024 Evidence for ~12-h ultradian
gene programs in humans. *npj Biological Timing and Sleep*. **1**, 4.
(doi:10.1038/s44323-024-00005-1)

87. Häfker, N. S., Meyer, B., Last, K. S., Pond, D. W., Hüppe, L., Teschke, M. 2017
Circadian clock involvement in zooplankton diel vertical migration. *Curr. Biol.* **27**,
2194-2201.e2193. (doi:10.1016/j.cub.2017.06.025)

88. Oosthuizen, M. K., Bennett, N. C. 2022 Clocks ticking in the dark: a review of
biological rhythms in subterranean African mole-rats. *Front. Ecol. Evol.* **10**,
(doi:10.3389/fevo.2022.878533)

Figure Legends

Figure 1. Deep-sea chronobiology study on the hydrothermal vent shrimp *Rimicaris leurokolos*. (*a*) This work was focused on the Iheya North hydrothermal vent field, Okinawa Trough. (*b*) Time series of published temperature data covering 351 days recorded at the peripheral area of the active NBC mound vent. Lomb-Scargle (LS) periodogram analysis reveals a strong semidiurnal tidal cycle dominated by the M2 tidal constituent. M2, principal lunar semidiurnal constituent; S2, principal solar semidiurnal constituent; N2, larger lunar elliptic semidiurnal constituent; K1, luni-solar declinational diurnal constituent; O1, Lunar declinational diurnal constituent. (*c*) *R. leurokolos* shrimps (in the yellow circle) live in a hot zone near vent orifice where hydrothermal fluids mix with cold surrounding seawater spontaneously. (*d*) Time series of *in situ* temperature and dissolved oxygen records during 162 seconds when the sensor was placed in the *R. leurokolos* colony showing frequent environmental fluctuations with high variability.

Figure 2. The *Rimicaris leurokolos* temporal transcriptome was dominated by tidal oscillations under constant conditions. (*a*) Principal components analysis (PCA) plot showing the detrended transcriptomic data from the free-running experiment samples (n = 57). Significance of inter-group differences was examined by the PERMANOVA (*P*-value = 0.376) and ANOSIM (*P*-value = 0.958) analysis both on Bray-Curtis dissimilarities. (*b*) Distribution of the periods of oscillations identified by the eigenvalue/pencil method and the ECHO application. Oscillations with a period range of 10-14 hours and 22-26 hours are regarded as tidal and daily cycling, respectively. (*c,d*) Venn diagrams detailing the number of rhythmic transcripts (FDR < 0.05) identified by ECHO, RAIN, and eJTK in the (*c*) tidal range and (*d*) daily range. (*e*) Raincloud plot of the root mean square (RMS) amplitude for 12-hour tidal transcripts and 24-hour daily transcripts. The stars indicated the level of significance: **** *P*-value < 0.0001.

Figure 3. Free-running tidal transcriptomic patterns of *Rimicaris leurokolos*. (*a*) Density distribution based on amplitude change coefficient (γ) of tidal transcripts detected by ECHO. Oscillation types are categorized as forced (-0.15 < γ < -0.03; amplified amplitude), harmonic (-0.03 ≤ γ ≤ 0.03; steady amplitude), damped (0.03 < γ < 0.15; reduced amplitude), overexpressed (γ ≤ -0.15), and repressed (γ > 0.15), and the latter two are considered arrhythmic. (*b*) Rose chart showing phase distribution comparison of tidal transcripts in cluster 1 and cluster 2. For each oscillation, the phase

is divided by period length and then times 360 (degrees) to normalize phases of transcripts with different period lengths. (*c*) Heatmap showing highly synchronized gene expression of tidal transcripts (ECHO FDR < 0.05). Each row represents a transcript with three individuals per time point. Transcripts are clustered and ordered by phases, oscillation type, and monotonic trend. The expression of each gene is Z-score normalized. (*d*) Bubble diagram showing significantly enriched gene ontology (GO) terms related to biological processes of tidal subgroups categorized by three classification schemes (from left to right: cluster, oscillation type, trend). Cross symbols represent a lack of enriched terms in the corresponding subgroup.

Figure 4. Expression profiles of *Rimicaris leurokolos* homologs of candidate circadian clock genes. Data points are displayed as mean values ± SEM (n = 3 biological replicates per time point). Minimum *P*-values of results from tidal or daily detection of each transcript are presented in the plot. Significant levels of oscillations are indicated with stars: * FDR < 0.05.

Figure 5. Homologous genes with a period of 12-hour rhythms between *Rimicaris leurokolos*, *Mus musculus*, and *Drosophila melanogaster*. (*a*) Venn diagram showing the comparison of ~12-hour rhythms of single-copy orthologous genes between three species. (*b*) Pathway enrichment network showing biological functions of 21 conserved ~12-hour genes in three species.

Figure 6. A new perspective of biological rhythms in the dark biosphere. Terrestrial and shallow-water organisms largely rely on circadian clocks for anticipating periodic environmental changes such as the day-night transition. Terrestrial organisms such as human beings exhibit robust ~24-hour rhythms driven by the circadian clock [2], and potential ~12-hour ultradian rhythms (with asterisk) that are assumed to be homologous with circatidal rhythms [86], though the biological functions and synchronization cues are still unclear. The intertidal mussel *Mytilus californianus* shows both circatidal and circadian rhythms, while the latter is the dominant one [51]. The copepod *Calanus finmarchicus* also shows behavioral rhythms in diel vertical migration (DVM) [87]. However, a different scenario can be found in the dark biosphere. Living away from direct sunlight, subterranean mole rats scarcely depend on circadian rhythms for survival, and the endogenous circadian system is understood to be in the process of regression, even though still present [88]. In this work, we found that in addition to the vent mussel *B. azoricus* [22], the vent shrimp *Rimicaris leurokolos* shows strong

956   endogenous 12-hour rhythms in the gene expression, likely reflecting important effects

957   of tidal cycles on deep-sea hot vent organisms.

a Permanova R²: 0.91; P-value: 0.376 Anosim R: -0.09; P-value: 0.958