Beta$_2$-Adrenergic Agonists in Treatment for Parkinsonism, with Implications for Neurodegenerative and Neoplastic Disorders.

Mario A. Inchiosa, Jr.

Department of Pharmacology, New York Medical College, Valhalla, NY 10595

Short title: **Beta$_2$-Adrenergic Agonists: In Parkinsonism, Neurodegenerative Disease and Cancer.**

Abstract: There is a long record of observations suggesting that β_2-adrenergic agonists may have therapeutic value in Parkinson's disease. Recent studies have focused on the possible role of β_2-receptor agonists in suppressing the formation of α-synuclein protein, the component of Lewy bodies. Levalbuterol, the pure levo isomer of the β2 selective agonist, albuterol, has been found to possess significant anti-inflammatory activity, a property that may have the potential to suppress cytokine mediated degeneration of dopaminergic neurons and progression of Parkinsonism. All of the β2 agonist and anti-inflammatory activities of albuterol reside in the levo isomer. The dextro isomer of albuterol substantially negates the efficacies of the levo form. Epinephrine, the prototypical β2 agonist and certain other adrenergic agents, when modeled in the Harvard/MIT Broad Institute genomic database, CLUE, demonstrated strong associations with the gene-expression signatures of drugs possessing glucocorticoid receptor agonist activity. Gene-expression signatures generated by the interaction of the adrenergic drugs of interest in 8 human tumor cell lines were compared with the entire CLUE database of more than 8,000 agents. The signatures were summarized for their consistency (connectivity) across all 8 cell lines and ranked for their relative degree of similarity to the agents in the database. Possible associations with anti-inflammatory activity of glucocorticoids prompted *in vivo* biological confirmation for levalbuterol and related agonists in the Jackson Laboratory human peripheral blood mononuclear cell (PBMC)-engrafted mouse. Levalbuterol inhibited the release of the eosinophil attractant chemokine, eotaxin-1 (specifically, CCL11), when the mice were challenged with mononuclear antibodies known to provoke cytokine release. Eotaxin is implicated in CNS and peripheral inflammatory disorders. Also, elaboration of the broad tumor-promoting angiogenic factor, VEGFa, and the pro-inflammatory cytokine, IL-13, from activated PBMCs were also inhibited by levalbuterol. These observations suggest possible translation to Parkinson's disease, other neurodegenerative syndromes, and malignancies, by several mechanisms.

Keywords: beta2-adrenergics; anti-inflammatory effects of levalbuterol; inhibition of eotaxin elaboration; inhibition of VEGF elaboration; cortisol-epinephrine interaction

1. Introduction

Magistrelli and Comi [1] have reviewed the earlier history of reports of an association between treatment with β_2-adrenergic receptor (β2AR) agonists and a therapeutic benefit in patients with Parkinson's disease (PD). Three open-label studies were noted, dating from 1992, 1994 and 2003, with the use of salbutamol as an adjunct to levodopa in a total of 25 patients. In the earliest study (9 PD patients), there were significant increases in "on time" and shortening of response latency [2]. The 1994 study with 8 PD patients [3] showed statistically significant improvement in PD tests for manual dexterity and time to arise and walk 20 feet and return and sit again; the 2003 study with 8 PD patients also found improvement in manual dexterity and in the Unified Parkinson's Disease Rating Scale (UPDRS) scores. [4]

1.1. α- Synuclein gene (SNCA) and PD

The observations in these earlier studies have advanced considerably with the recognition that β2AR ligands modulate the transcription of the α-synuclein gene (*SNCA*); and, the gene product, α-synuclein (α-syn), is implicated in the development of PD [1,5]. PD pathology is clinically correlated with the extent of excessive expression of *SNCA* and synthesis of α-syn, which accumulates intracellularly in Lewy bodies in the brains of PD patients.

In addition to PD, synucleinopathies include the neurodegenerative diseases Lewy bodies dementia and multiple system atrophy. All three demonstrate cellular accumulation of α-syn and loss of dopaminergic neurons [6].

Mittal et al. [5] screened 1126 FDA approved drugs and a variety of health supplements in cultured human neuroblastoma cells (SK-N-MC) for their ability to reduce transcription of *SNCA* mRNA in a four-stage gene expression assay. The highest efficacies were found in 3 β2AR agonists,

metaproterenol, clenbuterol and salbutamol and, interestingly, in the drug, riluzole, which has been 67

approved by the FDA for intervention in amyotrophic lateral sclerosis, where it was found to 68

decrease dopaminergic degeneration in a PD model in the rat.   All three β2AR agonists caused 69

statistically significant relative reductions for *SNCA* mRNA abundance and α-syn protein abundance 70

in SK-N-MC cells [5].   Of particular relevance to PD pathology and loss of dopaminergic neurons, 71

these investigators also found that intraperitoneal clenbuterol treatment for 24 hours reduced 72

expression of *SNCA* in the substantia nigra of C57BL/6J wild type mice, with decreases in nigral 73

mRNA and α-syn protein. Additionally, higher cellular levels of α-syn produced mitochondrial 74

dysfunction and production of superoxide anions and other reactive oxygen species (ROS); 75

clenbuterol treatment in neuronal precursor cells (iPSC-derived) from a PD patient suppressed these 76

changes [5]. 77

1.2. PD causation from neuroinflammation 78

A number of studies have focused on the importance of inflammatory processes in 79

neurodegenerative diseases, including PD.   In this connection, there is considerable interest in the 80

apparent role of β2AR signaling on immune cells in suppressing the inflammatory potential of 81

microglia and macrophages, and in T cells and B cells. Signaling through this receptor can influence 82

the inflammatory response of these cells [1,5,7–13]. 83

The role of neuroinflammation in the initiation and progression of PD is a major focus of these 84

present investigations. The functional role of the β2AR in biological responses has been further 85

elucidated by the development of "short-acting" and "long-acting" relatively selective β2AR agonists 86

that have little β1AR efficacy. These drugs have been used primarily for the treatment of bronchial 87

asthma and chronic obstructive pulmonary disease (COPD).   The nomenclature for these agents in 88

the literature includes either the "United States Adopted Name" (USAN) or the "International 89

Nonproprietary Name" (INN), usually depending upon the country of origin of the research or the 90

date of the publication. The designated names of the principal drugs that were investigated are as 91

follows: Albuterol sulfate is the USAN for a short acting bronchodilator that has an INN of 92

salbutamol sulfate. Albuterol is a racemic mixture of equal quantities of the S and R isomers; all of 93

the β2-adrenergic activity resides in the R isomer. Some proprietary names for albuterol sulfate are 94

Ventolin, Proventil and ProAir. The pure R-isomer has the USAN of levalbuterol HCl (a proprietary 95

formulation as levalbuterol tartrate is named Xopenex); its INN is levosalbutamol (it is also referred 96

to as R-salbutamol). Arformoterol is the USAN for the pure R isomer of a long-acting β2AR agonist; 97

the racemic mixture of the drug is named formoterol. The bronchodilator drug theophylline was also 98

studied; it does not have β2AR agonist activity but inhibits the metabolism of cyclic AMP (cAMP) 99

by phosphodiesterase enzymes thereby preserving intracellular levels; cAMP is the initial signaling 100

messenger that results from activation of both β1- and β2-ARs. 101

It is important to note the differences in the expected anti-inflammatory activity of β2AR 102

agonists regarding whether they are racemic mixtures, such as albuterol, or the pure R-isomer, such 103

as levalbuterol. In addition to the fact that the R-isomer has all the bronchodilator activity of 104

albuterol, it also has all the anti-inflammatory activity. The S-isomer of albuterol has pro- 105

inflammatory activity and partially, completely, or overwhelmingly negates the anti-inflammatory 106

activity of the R-isomer. Mazzoni et al. [14] demonstrated a hyperresponsiveness to histamine in 107

guinea pig airways following pretreatment with S-albuterol. This is apparently related to the capacity 108

of that isomer to increase smooth muscle proliferation, as observed in human bronchial cell cultures 109

[15]; levalbuterol decreased cell proliferation compared with the control cultures enriched with 5% 110

fetal bovine serum. Baramki et al. [16], Chorley et al. [17] and Wang et al. [7] have all confirmed the 111

anti-inflammatory effects of levalbuterol (R-albuterol) and the pro-inflammatory and compromising 112

effects of S-albuterol on those of levalbuterol. 113

1.3. Epidemiological associations between beta2-adrenergic effects and PD 114

Several large epidemiological studies have revealed the finding that the clinical use of β2AR 115

agonists (primarily for patients with asthma) had a decreased incidence of PD compared to the 116

general population and that the use of a β2AR antagonist (primarily for the treatment of

hypertension) had an increased incidence. Mittal et al. [5] studied the Norwegian Prescription

Database over an 11-year period for the entire population alive on January 1, 2004 (n = 4.6 million).

Patients treated with salbutamol (albuterol) showed a decreased PD risk; the rate ratio (incidence

with salbutamol/incidence without exposure to salbutamol) was 0.66 (95% confidence interval, CI of

0.58 to 0.76). Patients treated with propranolol, a β1-,β2AR antagonist had a marked increased rate

ratio of PD to 2.20 (95% CI of 1.62 to 3.00).

Using health records from the major Israeli health provider, Gronich et al. [18] studied medical

records of 1,762,164 individuals who did not have a PD diagnosis on January 1, 2004 and followed

their health histories until June 30, 2017. Each participant who was treated with salbutamol was

compared with 10 cohorts as controls and showed a decreased rate ratio for development of PD of

0.89 (0.82-0.96); p = 0.004. Participants treated with propranolol showed an increased PD rate ratio

of 2.60 (2.40-2.81); p <0.001. A special feature of this study [18] was that it included a number of

participants that were treated with a group of β1 selective antagonists instead of the β1-, β2AR

antagonist, propranolol. A total of 3,032 participants received either metoprolol, atenolol or

bisoprolol. Since these drugs leave β2ARs largely available, it would be expected that they would be

chronically stimulated by endogenous epinephrine that was released during daily activities and

exercise. These participants showed no change in PD rate ratio as compared with their cohort

controls; 1.00 (0.95-1.05); p = 0.94. All the above epidemiological studies consistently

demonstrated that drugs that stimulated the β2AR receptor, or interventions that left it available for

activation, decreased the incidence of PD.

Chen et al. [19] conducted a meta analysis on the association between β-adrenoceptor drugs and

Parkinson's disease and distilled 640 studies down to 8 that were appropriate for meta-analysis.

Salbutamol yielded a decreased rate ratio for PD of 0.888 (0.822-0.960); p = 0.003. When the studies

that used one or more of the β2AR agonists were analyzed together, the rate ratio for PD was still

lowered: 0.840 (0.714-0.987); p = 0.035. This included the β2 selective agonists salbutamol, formoterol, salmeterol, and terbutaline. It must be noted that all these drugs are the racemic mixtures; in the studies noted above, only the levo (R) isomers have been shown to have anti-inflammatory activity while the S-isomer portion is actually proinflammatory and compromises the effects of the R-isomer. Thus, the decreased incidence of PD in participants that received β2AR agonists may have been greater if the single R-isomer agents had been studied.

1.4. Life-style factors that may influence the incidence or progression of PD

There is considerable evidence that certain lifestyle practices delay the initiation of PD and the rate of its progression. These include physical exercise, nicotine exposure from smoking, and intake of caffeine from consumption of coffee or tea. In a potential connection with the present study, these life-style interventions are associated with an increased outflow of the sympathetic nervous system and elevations in circulating levels of epinephrine, with its prominent β2AR agonist activity.

Exercise programs are a major therapeutic recommendation on the websites of the Parkinson's Foundation and the Michael J. Fox Foundation to forestall the progression of disease and to ameliorate the symptoms of PD. Moore et al. [20] reviewed extensive literature showing the beneficial effects of physical exercise in human and animal models of PD. They also presented the projected design of a Phase II trial, "Study in Parkinson Disease of Exercise," (SPARX). It was intended to serve as a pilot study to determine whether a larger study on graded intensity of exercise was warranted. It included patients who were at an early stage of PD and were not on dopaminergic therapy or, if so, for only a short period. There were 3 groups: High intensity treadmill exercise for 30 min, 4 times/week at 80-85% of their maximum heart rate; moderate intensity exercise for 30 min, 4 times/week at 60-65% of maximum heart rate; and usual care for a wait-list group who were eligible later for inclusion in the study. The ability to identify treatment effects was complicated by the fact that the usual care group were permitted to continue their current exercise programs since, as noted above, the value of exercise in PD was already appreciated and it would not have been ethical

to prevent the control group from possible benefits of their existing programs. The groups were to be maintained and monitored for a period of 6 months.

Schenkman et al. [21] reported the results of the Phase II pilot trial that included a total of 128 participants. There were improvements in the Movement Disorder Society revised Unified Parkinson Disease Rating Scale (MDS-UPDRS) in both treadmill groups, but statistical significance was only reached at the high intensity level. Recruitment is now ongoing for the SPARX3 trial that will be conducted at 29 sites in the United States and Canada (Study No. NCT04284436; sparx3pd.com) with 370 participants. This study will have considerably more statistical power since there will only be two groups divided at the two intensity levels; the changes in MDS-UPDRS scores will be evaluated for each participant between their initial values and at the end of the study period. The SPARX3 trial has been delayed by the COVID-19 pandemic and completion of recruitment is now estimated to be July, 2025, with the results available approximately 18 months after close of the study.

The studies of Kjaer et al. [22] on epinephrine release with exercise are of relevance in relation to the evolving focus on the intensity of exercise as therapy in PD. Their findings in a series of studies showed a direct relationship between exercise intensity in healthy adults and epinephrine arterial plasma concentrations. Four levels of exercise intensity were studied based on the individual subjects' maximal oxygen uptake (VO_2MAX). Resting epinephrine concentration averaged 0.5 nmol/l. Plasma epinephrine concentration after 60 min at 40-56% VO_2MAX averaged 1.6 nmol/l; after 25 min at 60-70% VO_2MAX, 2.2 nmol/l; after 25 min at 82-89% VO_2MAX, 4.5 nmol/l; and after 4 min at 100-110% VO_2MAX, 8.7 nmol/l.

Tobacco smoking has a long and established negative association with the onset and progression of PD [23–27]. Meta-analysis by Hernan et al. [27] that included 44 case-controlled studies and 4 cohort studies showed a relative risk of PD between current smokers and individuals that had never smoked of 0.39 (95% confidence interval of 0.32-0.47). The pharmacological effects of nicotine

inhalation on the sympathetic nervous system would appear to be a convincing explanation for this protective effect. In addition to stimulating outflow of the sympathetic nervous system from the brain, nicotine activates nicotinic receptors in sympathetic ganglia peripherally to release norepinephrine from sympathetic nerve endings and also generates release of epinephrine by stimulation of nicotine receptors in the adrenal medulla.

Cryer et al. [28] studied the effects on catecholamine plasma concentrations in 10 male subjects in conjunction with two trials of smoking 5.5 cm of a cigarette in a 10 min period. The men were chronic smokers but had fasted overnight and had not smoked until the time of the experiments. Plasma concentrations of norepinephrine and epinephrine rose rapidly during the smoking period to maxima at about the 10-min time point. The average epinephrine concentration before smoking of 44 +/- 4 (mean; S.E.) pg/ml rose to 113 +/- 27 pg/ml (p < 0.05) at 10 min; it remained noticeably above the basal level for the next 20 min until measurements were halted. Grassi et al. [29,30] also studied the acute effects of cigarette smoking on epinephrine plasma levels. Their protocol required 48 hours of abstinence from smoking and completion of the cigarette in a 5 min period. Although the smoking period was half that in the previous study [28], epinephrine concentrations increased from 25 +/- 8.6 (mean; S.E.) pg/ml before smoking to 47 +/- 11.5 pg/ml (p <0.01) at the end of the period.

The relationship of caffeine consumption to a decreased incidence of PD has a long and consistent documentation [27,31–36]. Tan et al. [32], in a completely Chinese population, found that 10 years of coffee consumption at 3 cups/day was associated with a decreased odds ratio for PD of 0.78 (95% CI 0.66-0.93) p = 0.006, and for 10 years of tea drinking at 3 cups/day odds ratio of 0.72 (95% CI 0.56-0.94) p = 0.014). Ren and Chen [36] have reviewed the several large prospective and meta-analyses that show strong evidence of the negative association between caffeine consumption and PD. Notably, a meta-analysis by Qi and Li [35] that involved a total of 901,764 participants from 13 studies showed a maximum benefit in PD incidence reduction at approximately 3 cups of coffee/day: relative risk of 0.72 (95% CI 0.65-0.81). This same review [36] presented

evidence from animal studies that caffeine may reduce neuroinflammation and its attendant loss of

dopaminergic neurons. Also, in α-syn animal models of PD, caffeine administration was associated

with a decreased aggregation of α-syn, the component of Lewy bodies.

Finally, consistent with the above observations with physical exercise and cigarette smoking,

caffeine consumption results in increases in plasma epinephrine. In a precisely controlled clinical

study, Robertson et al. [37] demonstrated that 250 mg of caffeine in non-coffee drinkers resulted in a

significant increase in plasma epinephrine concentration one hour after consumption from 36 +/- 5

pg/ml (mean, S.E.) for a control beverage to 89 +/- pg/ml with caffeine; $p < 0.001$. Plasma

epinephrine concentrations were still statistically different from controls in the final measurements 3

hrs. after caffeine consumption.

Benowitz et al. [38] observed similar findings in caffeine consumption of 2 and 4 mg/kg with a

cross-over design in coffee drinkers who had abstained for 3 days before the study. The results,

expressed as maximum changes in epinephrine concentrations were 35 +/- 19 (mean, S.E.) pg/ml for

the 2 mg/kg dose and 63 +/-18 pg/ml for 4 mg/kg. Both doses were statistically different from

controls ($p < 0.05$) and the 4 mg/kg dose was statistically greater than the 2 mg/kg dose ($p < 0.05$).

Measurements were made over a 180 min period and the areas under the time x plasma

concentrations (AUCs) were also statistically and markedly different from the controls, with the 4

mg/kg dose again statistically different from the 2 mg/kg dose (differences of $p < 0.05$).

1.5. Genomic modeling of the potential anti-inflammatory effects of epinephrine

The evidence outlined above on the possible role of β2AR agonist activity on: 1, Reducing the

expression of *SNCA* mRNA and its protein product α syn; 2, The reduction in neuroinflammation

and loss of dopaminergic neurons; 3, Epidemiological associations with decreased incidence of PD

in their routine clinical use; and 4, Life-style factors such as physical exercise, cigarette smoking and

caffeine consumption that are negatively associated with PD incidence, and which all result in

increased plasma epinephrine concentration, prompted an attempt to explore a possible link between 241

the gene-expression characteristics of epinephrine and an anti-inflammatory potential. 242

Epinephrine was modeled in the Harvard/MIT Broad Institute genomic database, CLUE [39]. It 243

was found to have a remarkably coherent similarity in gene-expression activity to the human 244

glucocorticoid, cortisol (hydrocortisone), which has profound anti-inflammatory activity. Of 245

particular interest, this property was not shared with norepinephrine, which is known to have 246

minimal β2AR agonist activity. These observations led to the investigation of the potential anti- 247

inflammatory effects of a series of β2AR agonists in the Jackson Laboratory human peripheral blood 248

mononuclear cell (PBMC)-engrafted mouse. 249

1.6. Anti-inflammatory studies in the Cytokine Release Syndrome (CRS) assay 250

The CRS assay of Jackson Laboratories in humanized mice was utilized to evaluate the anti- 251

inflammatory potential of several of the β2AR agonists and related compounds noted above, 252

primarily to evaluate their capacity to protect against and modulate neuroinflammation. This assay 253

employs immunodeficient mice that have been engrafted with human peripheral blood mononuclear 254

cell cultures (PBMCs) that are then challenged with several monoclonal antibodies (mAbs) that are 255

known to potently induce clinically injurious cytokine release [40,41]. The mice can be pre- and 256

chronically treated with agents of interest for anti-inflammatory effects in conjunction with the mAb 257

challenges. The Jackson Laboratories, Bar Harbor, Maine website provides additional details of the 258

CRS assay. 259

The direct β2AR agonists, albuterol, levalbuterol, and arformoterol, and the indirect agonist 260

theophylline, all noted above to increase cellular levels of cAMP, were studied in the mouse CRS 261

assays. Phenoxybenzamine, a noncompetitive antagonist of alpha2-adrenergic agonist activity that 262

results in increased cellular levels of cAMP [53], was also assayed. All the compounds except 263

albuterol, the racemic mixture of S and R albuterol, showed anti-inflammatory potential in the CRS 264

assay. 265

The possible anti-inflammatory activity of beta2-adrenergic agonists, as suggested by their gene-expression similarity to that of glucocorticoid receptor agonists, became a principal aim of these investigations. This hypothesis was tentatively confirmed by the inhibitory effects of beta2-adrenergic drugs on the release of several cytokines and chemokines in the activated human PBMC-engrafted mouse.

2. Materials and Methods

2.1. Analyses of gene-expression signatures of β2-adrenergic agonists and related compounds in the Harvard/MIT Broad Institute genomic database, CLUE

Both the earlier online Broad Institute CMap genomic dataset and the new cloud-based CLUE platform are based on the gene-expression signatures that result from the perturbation of actively proliferating cells that are malignant. The earlier online platform showed predictions of antitumor and histone deacetylase activity of phenoxybenzamine [56]. The strength of these predictions was extended when phenoxybenzamine was profiled on the cloud-based platform [57]. Both references provide additional details of the methodology.

The database and associated software are accessible at https://clue.io. The drugs, chemical agents and genetic inputs are referred to as, "perturbagens" [39, 42].   The present analyses focused on gene-expression signatures located in the "TOUCHSTONE" dataset (accessed in the "Tools" menu) that was selected because it represents a set of thoroughly annotated small molecule perturbagens that would be expected to be relevant to comparisons with the possible effects of β2-adrenergic agonists; see: https://clue.io/connectopedia/tag/TOUCHSTONE.   The Touchstone dataset contains a total of approximately 8,400 perturbagens that have produced gene signatures that were generated from testing on a panel of 9 proliferating malignant cell lines; those cell lines are detailed in Table 1.

Table 1. Cell Lines Profiled in the TOUCHSTONE Database.

Cell Line	Description
A375	Human malignant melanoma
A549	Human non-small cell carcinoma
HA1E	Human kidney epithelial immortalized
HCC515	Human non-small cell lung adenocarcinoma
HEPG2	Human hepatocellular carcinoma cell line
MCF7	Human breast adenocarcinoma
PC3	Human prostate adenocarcinoma
VCAP	Human metastatic prostate cancer
HT29	Human colorectal adenocarcinoma

.   In almost all instances, the perturbagens were tested at a concentration of 10 μM, allowing for the

comparison of the strength of the connectivity of gene-expression between agents at equimolar

concentrations.

It should be noted that the dataset in the earlier online platform of CMap was based almost

entirely on perturbations in only 3 tumor cell lines, MCF7 (human breast adenocarcinoma), PC3

(human prostate adenocarcinoma) and HL60 (human promyeloblast); the first two cell lines are

among the 9 in the current CLUE platform and consistency among the gene-expression signatures for

the different cell lines is strengthened by the inclusion of additional cell lines.

The scoring value obtained in the gene expression analyses is termed "tau;" it ranges from 100

to -100 and is a measure of the connectivity between the gene expression-signature of the

perturbagen of interest and those of the other 8400 perturbagens in the database.   A positive tau

indicates a relative similarity between two perturbagens or group of perturbagens, while a negative

score indicates relative opposing gene signatures.   Thus, for example, a tau score of 95 indicates that

only 5% of the signatures in the database had connectivity that was higher than that of the

perturbagen being profiled; see: https://clue.io/connectopedia/connectivity_scores.   Positive scores

above 90 are generally considered worthy of consideration as representing possibly important

similarities in gene expression. The gene-expression data outputs from the analyses places emphasis

on this range.

The Touchstone database returns two data formats in response to a query about the similarities

or differences between a perturbagen of interest and the other perturbagens in the database.   The

compound being searched against all other entries in the Touchstone database is termed the

"INDEX."   A "heatmap" format presents the connectivity score between the query perturbagen and

the reference perturbagens in the database for each of the 9 cells lines that have been studied.

(Seven to 9 cell lines may have been chosen for a particular INDEX analysis.)   The second format is

the "detailed list" output.   In our experience, this output provided the most convenient access to 316

information that relates to the primary focus of the present studies.   The detailed list format also 317

includes the protein targets of the individual perturbagens.   Both formats provide comparison of the 318

gene signatures of the INDEX compound with members of the four perturbagen classes in the 319

Touchstone database.   Those classes are identified as: Chemical compound/pharmacologic agent 320

(CP); Gene knock-down (KD); Gene over-expression (OE); and perturbagen class (PCL).   In the 321

heat map format, INDEX signatures are compared with the entire Touchstone database of 322

perturbagens (approximately 8400).   Since our primary interest was to search for drugs and 323

chemicals that had recognized mechanisms of action that might strengthen previous findings and 324

support hypotheses for the repurposing of drugs therapeutically, we focused our searches in the 325

detailed list format primarily to the CP agents. 326

The heatmap output that is presented gives the calculated *median* tau score (connectivity) for 327

the 9 cell lines and a "summarization" measure.   For all other results, the connectivity scores that 328

are presented are the summarization measure.   This measure is calculated across the 9 cell lines. It 329

is valuable since it provides a measure of the consistency of perturbations from one cell line to 330

another.   The algorithms for the calculation of tau and the summarization measure, and for the 331

associated formulas, are presented at https://clue.io/connectopedia/cmap_algorithms. Gene- 332

expression connectivity scores are also associated with a "rank" measure for that score; the rank 333

score has an almost perfect non-linear inverse correlation with the gene expression 334

similarity/connectivity score. These measures are detailed further in S1 Figure. 335

The results represent the status of the CLUE platform when accessed primarily between May, 336

2020 and July, 2023; the database is dynamic and changes slightly if new perturbagens are added. 337

2.2. CRS assay methodology 341

Female NSG-SGM3 mice [*NOD.Cg-Prkdc^scid Il2rg^tmlWjl* Tg(CMV-IL3,CSF2,KITLG) 342
lEav/MloySzJ;stock #013062] were used for the study. The mice were ear notched for 343
identification and housed in individually ventilated polysulfonate cages with HEPA filtered air at 344
a density of up to 5 mice per cage. Cages were changed every two weeks. The animal room was 345
lit entirely with artificial fluorescent lighting, with a controlled 12 h light/dark cycle (6 am to 6 346
pm light). The normal temperature and relative humidity ranges in the animal rooms were 20-26°C 347
and 30-70%, respectively. The animal rooms are set to have up to 15 air exchanges per hour. A 1% 348
sucrose solution of filtered tap water, acidified to a pH of 2.5 to 3.0, and standard lab chow were 349
provided ad libitum. 350

<div align="center">Study Design</div> 351

1. Mice were irradiated and injected with PBMCs from a single donor on Day 0. 352

2. Animals were monitored for body weight and clinical observations once daily. 353

3. Animals were euthanized before study end if they showed >20% body weight loss or a body 354
condition score of ≥2. 355

4. Treatment compounds were delivered in drinking water starting on Day 4 for the duration of 356
study. 357

5. Mice were dosed on Day 6 with monoclonal antibodies OKT3 or anti-CD28, to induce 358
cytokine release [41], or with PBS as control. 359

6. Mice were bled via retro-orbital (RO) method at 6 h post dose on Day 6. Blood was 360
processed and measured for human cytokines and chemokines using MILLIPEX Human 361
Cytokine/Chemokine/Growth Panel A Magnetic Bead Panel, Cat# HCYTOMAG-60K. 362

7. Bodyweight and clinical CRS score were assessed daily starting Day 6. Mice were euthanized by CO2 asphyxiation at 72 h post dose Day 9.

The experimental design of the study is summarized by Study Day (SD) as follows:

The clinical condition of the mice was rank-scored as follows:

Clinical CRS score:
0: normal activity
1: normal activity, piloerection, tiptoe gait
2: hunched, reduced activity but still mobile
3: hypomobile but mobile when prompted
4: moribund (non-responsive to touch)

Dosing schedule and interventions

Group	N	Drinking Water* (D4)	Compound (D6)	Dose (mg/kg)	Dosing Route	Dosing Frequency	Bleed** RO	Terminal
1	5		PBS	N/A	IV	Single dose	D6	D9
2	5	No pre-treatment	OKT3	0.25	IV	Single dose	D6	D9
3	5		Anti-CD28	1	IV	Single dose	D6	D9
4	5		PBS	N/A	IV	Single dose	D6	D9
5	5	Albuterol 8 ug/mL	OKT3	0.25	IV	Single dose	D6	D9
6	5		Anti-CD28	1	IV	Single dose	D6	D9
7	5		PBS	N/A	IV	Single dose	D6	D9
8	5	Levalbuterol 4 ug/mL	OKT3	0.25	IV	Single dose	D6	D9
9	5		Anti-CD28	1	IV	Single dose	D6	D9
10	5	Phenoxy-	PBS	N/A	IV	Single dose	D6	D9
11	5	benzamine	OKT3	0.25	IV	Single dose	D6	D9
12	5	5 ug/mL	Anti-CD28	1	IV	Single dose	D6	D9
13	5		PBS	N/A	IV	Single dose	D6	D9
14	5	Theophylline 400 ug/mL	OKT3	0.25	IV	Single dose	D6	D9
15	5		Anti-CD28	1	IV	Single dose	D6	D9
16	5	Arformoterol	PBS	N/A	IV	Single dose	D6	D9
17	5	Tartrate	OKT3	0.25	IV	Single dose	D6	D9
18	5	0.008 ug/mL	Anti-CD28	1	IV	Single dose	D6	D9
Total	90							

*Freshly prepared drinking water was provided daily. **Mice were bled via retro-orbital (RO) on Day 6 and cardiac puncture on Day 9

Levalbuterol Hydrochloride, Albuterol sulfate, and Phenoxybenzamine Hydrochloride were

purchased as reference standards from the United States Pharmacopeia. Arformoterol Tartrate and

Theophylline (anhydrous powder) were purchased from Sigma-Aldrich Chemical Company.

2.3. Statistics

Group sizes of these valued experimental animals were kept as small as possible and were

consistent with the spirit of FDA Amendment 2.0. The cytokine and chemokine data from these

small groups largely failed normality distributions. Thus, all analyses were conducted with the

Kruskal-Wallis multiple comparative approach for nonparametric data followed by post-hoc

adjustment of p values with the Dunnett's test. Each drug treatment effect was only compared with

its sucrose control. A p value of 0.05 (*) or less was considered statistically significant; ** p <0.01; ***p <0.001; ****p <0.0001.

3. Results and Discussion

3.1. Analyses of gene expression signatures of β2-adrenergic agonists and related compounds in the Harvard/MIT Broad Institute genomic database, CLUE

In preliminary analyses, salbutamol (albuterol), the racemic mixture of the drug that was prominent in the Norwegian and Israeli epidemiological studies in relation to PD frequency [5,18] was profiled in CLUE.   As might be expected, salbutamol showed high connectivity with other beta-adrenergic receptor agonists; in this CMap class profiled in CLUE, 8 had gene-expression connectivity scores that were higher than 90.0 in relation to salbutamol (Table 2).

Table 2. Gene-expression connectivity scores of adrenergic receptor agonists in relation to salbutamol.

Rank	Score	Name	Description	Target
3	99.93	cimaterol	Adrenergic receptor agonist	ADRB3, ADRB1, ADRB2
27	99.37	procaterol	Adrenergic receptor agonist	ADRB2
118	97.54	dobutamine	Adrenergic receptor agonist	ADRB1, ADRB2, ADRA1A 3
183	96.72	orciprenaline	Adrenergic receptor agonist	ADRB2
255	95.52	buphenine	Adrenergic receptor agonist	ADRB2
288	95.21	fenoterol	Adrenergic receptor agonist	ADRB2, ADRB1, ADRB3
515	92.56	isoxsuprine	Adrenergic receptor agonist	ADRB2
553	92.25	salmeterol	Adrenergic receptor agonist	ADRB2

ADRB2, beta2-adrenergic receptor; ADRB1, beta1-adrenergic receptor; ADRA1A, alpha1a-adrenergic receptor; ADRB3, beta3-adrenergic receptor.

Also, when CLUE was queried for connectivity with dopamine receptor agonists, 5 had high scores in relation to salbutamol; this included dopamine itself (Table 3).   In view of the apparent associations between epinephrine and PD, it would appear to be important that there is an overlap of agonist activity of epinephrine on β2AR and dopamine receptors [43,44].

Table 3. Gene-expression connectivity scores of dopamine receptor agonists in relation to salbutamol.

Rank	Score	Name	Description	Target
8	99.65	SKF-77434	Dopamine receptor agonist	DRD1
51	98.70	fenoldopam	Dopamine receptor agonist	DRD1, ADRA1A, ADRA1B, ADRA1D, ADRA2A, ADRA2B, ADRA2C, DRD4, DRD5
52	98.66	pramipexole	Dopamine receptor agonist	DRD3, DRD2, ADRA2A, ADRA2B, ADRA2C, DRD1, DRD4, DRD5, HTR1A, HTR1B, HTR1D, HTR2A, HTR2B, HTR2C
335	93.45	RO-10-5824	Dopamine receptor agonist	DRD4
385	92.63	dopamine	Dopamine receptor agonist	DRD2, DRD1, DRD3, DRD4, DRD5, DBH, TR1A, HTR7, SLC6A2, SLC6A3, SLC6A4

DRDI, dopamine receptor D1; DRD2, dopamine receptor D2; DRD3, dopamine receptor D3; DRD4, dopamine receptor D4; DRD5, dopamine receptor D5; ADRA1A, alpha1a-adrenergic receptor; ADRA1B, alpha1b-adrenergic receptor; ADRA1D, alpha1d-adrenergic receptor; ADRA2A, alpha2a-adrenergic receptor; ADRA2B, alpha2b-adrenergic receptor; ADRA2C, alpha2c-adrenergic receptor; HTR1A, HTR1B, HTR1D, HTR2A, HTR2B, HTR2C, serotonin receptors 1A, 1B, 1D, 2A, 2B, 2C; DBH, dopamine beta-hydroxylase; TR1A, T regulatory Type 1 cells; HTR7, serotonin 7 receptor; SLC6A2, norepinephrine transporter; SLC6A3, sodium dependent dopamine transporter; SLC6A4. serotonin transporter.

The potential anti-inflammatory importance of β2-adrenergic agonists was first observed in the current study with the observation of the marked similarity of the gene expression signatures of epinephrine to those of glucocorticoid receptor (GR) agonists. A portion of the heat map with the highest tau scores when epinephrine was modeled in CLUE already showed the predominance of GR agonists with high tau scores for the top matches in the database; the protein targets of the matching compounds were also replete with the codes, NRC31, glucocorticoid receptor, and NRC32, mineralocorticoid receptor (Fig. 1). In comparison, norepinephrine, which has approximately equal β1-adrenergic activity as epinephrine but has little or no β2 activity, showed essentially no

connectivity to GR agonists in a comparable portion of its heatmap (Fig. 2). More than 60% of the

gene expression signatures with connectivity to epinephrine in Fig. 1 were GR agonists while there

was only 1 (0.025%) GR agonist that matched with a high tau score with norepinephrine (Fig. 2.).

Fig 1. Epinephrine HeatMap.
HeatMap of association strengths (median tau score) between gene expression signatures of
epinephrine in 8 malignant cell cultures and those for all entries in the Touchstone database, with
targets. Chemical compound (CP); Gene knock-down (KD); Gene over-expression (OE); and
perturbagen class (PCL).

Fig 2. Norepinephrine HeatMap.
HeatMap of association strengths (median tau score) between gene expression signatures of
norepinephrine in 8 malignant cell cultures and those for all entries in the Touchstone database, with
targets. Chemical compound (CP); Gene knock-down (KD); Gene over-expression (OE); and
perturbagen class (PCL).

The detailed analysis outputs from the CLUE platform added further evidence of the strong

association of epinephrine with drugs that have GR receptor agonist activity. A table is produced

that shows the gene enrichment scores for CMap classes in the Touchstone database (Table 4). A

high gene enrichment score is noted for the 44 GR agonists in this CMap class. As would be

expected, there is also a high enrichment score for the 11 beta-adrenergic receptor agonists CMap

class (Table 4). And, consistent with the almost complete lack of GR receptor agonists in the

norepinephrine heatmap (Fig. 2), no connections with the GR CMap class are found in the

comparable CMap class output for norepinephrine (Table 5). However, the general category of

beta-adrenergic receptor agonists is represented (Table 5) because epinephrine and norepinephrine

have equal $\beta1$-adrenergic potency.

Still further evidence that an association exists between GR receptor agonist activity and the

neurotransmitter, epinephrine, and that this is unique to this agent that has $\beta2$-adrenergic activity and

not shared by norepinephrine, is seen in a comparison of the detailed analyses associated with the

CMap class score differences. The strength of associations in gene signatures with GR receptor

agonists for epinephrine and norepinephrine are presented in Tables 6 and 7, respectively.   Of the 44

glucocorticoids in its CMap class, all have gene signature connectivity scores above 90.0 in relation

to epinephrine (Table 6), while only 7 have scores above 90.0 in connectivity with norepinephrine

and some have negative (opposing) connectivity (Table 7).   These rankings are presented

graphically in Fig. 3.

Table 4. CMap Class Connectivity: EPINEPHRINE.

Perturbagen Class (PCL)	Group Enrichment Score
Progesterone receptor agonist x 12	100.00
Glucocorticoid receptor agonist x 44	99.67
Aromatase inhibitor x 3	97.00
Beta-adrenergic receptor agonist x 11	96.12
FLT3 inhibitor x 6	95.46
Progesterone receptor antagonist x 5	95.13
Leucine rich repeat kinase inhibitor x 3	94.15
Bromodomain Inhibitor x 6	93.59
DNA synthesis inhibitor x 3	92.13
VEGFR inhibitor x 13	91.84
RAF inhibitor x 9	91.19

"Sets of compound perturbagens with enrichment scores above 90 (similar) and below -90
(opposing)" in relation to their correspondence with the gene expression signatures of epinephrine.

Table 5. CMap Class Connectivity: NOREPINEPHRINE.

Perturbagen Class (PCL)	Group Enrichment Score
Beta-adrenergic receptor agonist x 11	99.95
Adenosine receptor agonist x 6	99.58
Reverse transcriptase inhibitor x 4	99.27
Bacterial 30S ribosomal subunit inhibitor x 5	99.26
GABA receptor antagonist x 5	95.90
Sigma receptor antagonist x 3	95.58
Bile acid x 4	93.38
PPAR receptor agonist x 16	92.04
Benzodiazepine receptor agonist x 4	90.55
FXR antagonist x 3	90.27
Estrogen receptor agonist x 17	90.13

"Sets of compound perturbagens with enrichment scores above 90 (similar) and below -90 (opposing)" in relation to their correspondence with the gene expression signatures of norepinephrine.

Table 6.   Gene Expression-Connectivity Scores for EPINEPHRINE vs GR Agonists.

Rank	Score	Name	Description
4	99.89	Hydrocortisone	Glucocorticoid receptor agonist
5	99.89	Betamethasone	Glucocorticoid receptor agonist
7	99.79	Budesonide	Glucocorticoid receptor agonist
8	99.79	Flunisolide	Cytochrome P450 inhibitor
9	99.75	Mometasone	Glucocorticoid receptor agonist
11	99.75	Fluocinonide	Glucocorticoid receptor agonist
13	99.68	Triamcinolone	Glucocorticoid receptor agonist
14	99.68	Fludroxycortide	Glucocorticoid receptor agonist
19	99.62	Clobetasol	Glucocorticoid receptor agonist
21	99.61	Triamcinolone	Glucocorticoid receptor agonist
22	99.61	Flumetasone	Glucocorticoid receptor agonist
23	99.58	Fluticasone	Glucocorticoid receptor agonist
24	99.58	fluorometholone	Glucocorticoid receptor agonist
26	99.54	Dexamethasone	Glucocorticoid receptor agonist
27	99.52	Prednicarbate	Phospholipase activator
28	99.51	Isoflupredone	Glucocorticoid receptor agonist
29	99.47	Hydrocortisone	Glucocorticoid receptor agonist
30	99.44	Diflorasone	Corticosteroid agonist
31	99.44	Dexamethasone	Glucocorticoid receptor agonist
32	99.4	Medrysone	Glucocorticoid receptor agonist
33	99.37	Prednisolone	Glucocorticoid receptor agonist
34	99.33	Halcinonide	Glucocorticoid receptor agonist
37	99.12	Betamethasone	Glucocorticoid receptor agonist

39	99.05	Beclomethasone	Glucocorticoid receptor agonist
40	99.05	Fluocinolone	Glucocorticoid receptor agonist
42	99.05	Hydrocortisone	Glucocorticoid receptor agonist
43	99.01	Hydrocortisone	Glucocorticoid receptor agonist
45	98.77	Clocortolone	Glucocorticoid receptor agonist
46	98.77	Beclomethasone	Glucocorticoid receptor agonist
48	98.6	Westcort	Glucocorticoid receptor agonist
51	98.4	Prednisolone	Glucocorticoid receptor agonist
52	98.38	Fluticasone	Glucocorticoid receptor agonist
55	98.34	Hydrocortisone	Glucocorticoid receptor agonist
56	98.32	Rimexolone	Glucocorticoid receptor agonist
58	98.17	Fludrocortisone	Glucocorticoid receptor agonist
59	98.17	Amcinonide	Glucocorticoid receptor agonist
61	98.03	methylprednisolone	Glucocorticoid receptor agonist
67	97.71	desoximetasone	Glucocorticoid receptor agonist
75	97.67	Halometasone	Glucocorticoid receptor agonist
88	97.53	Loteprednol	Glucocorticoid receptor agonist
96	97.5	Depomedrol	Glucocorticoid receptor agonist
117	97.39	Prednisolone	Glucocorticoid receptor agonist
137	97.15	Fluocinonide	Glucocorticoid receptor agonist
632	93.24	Alclometasone	Glucocorticoid receptor agonist

Table 7. Gene-Expression Connectivity Scores for NOREPINEPHRINE vs GR agonists.

Rank	**Score**	**Name**	**Description**
23	99.55	fluocinonide	Glucocorticoid receptor agonist
244	97.47	depomedrol	Glucocorticoid receptor agonist
376	96.44	beclometasone	Glucocorticoid receptor agonist
471	95.46	fluocinolone	Glucocorticoid receptor agonist
615	94.13	prednisolone	Glucocorticoid receptor agonist
890	91.89	triamcinolone	Glucocorticoid receptor agonist
943	91.4	Amcinonide	Glucocorticoid receptor agonist
1479	86.61	Rimexolone	Glucocorticoid receptor agonist
1810	83.17	hydrocortisone	Glucocorticoid receptor agonist
1959	82.28	halometasone	Glucocorticoid receptor agonist
2159	80.37	beclometasone	Glucocorticoid receptor agonist
2349	78.38	triamcinolone	Glucocorticoid receptor agonist
2445	77.7	prednisolone	Glucocorticoid receptor agonist
2855	73.57	betamethasone	Glucocorticoid receptor agonist
2864	73.36	hydrocortisone	Glucocorticoid receptor agonist

2877	73.18	alclometasone	Glucocorticoid receptor agonist
3084	71.41	Westcort	Glucocorticoid receptor agonist
3155	70.7	Fluticasone	Glucocorticoid receptor agonist
3160	70.64	Loteprednol	Glucocorticoid receptor agonist
3348	68.45	dexamethasone	Glucocorticoid receptor agonist
3561	66.22	fluocinonide	Glucocorticoid receptor agonist
3591	65.9	desoximetasone	Glucocorticoid receptor agonist
4051	61.03	betamethasone	Glucocorticoid receptor agonist
4147	60	prednicarbate	Phospholipase activator
4393	56.96	clocortolone	Glucocorticoid receptor agonist
4404	56.86	hydrocortisone	Glucocorticoid receptor agonist
4465	56.37	fluorometholone	Glucocorticoid receptor agonist
4489	56.05	Flunisolide	Cytochrome P450 inhibitor
4701	52.82	mometasone	Glucocorticoid receptor agonist
5218	45.35	dexamethasone	Glucocorticoid receptor agonist
5512	41.08	prednisolone	Glucocorticoid receptor agonist
5594	39.97	fludroxycortide	Glucocorticoid receptor agonist
5618	39.72	methylprednisolone	Glucocorticoid receptor agonist
5696	38.68	Fluticasone	Glucocorticoid receptor agonist
5837	36.6	hydrocortisone	Glucocorticoid receptor agonist
6223	31.46	Medrysone	Glucocorticoid receptor agonist
6329	29.69	Budesonide	Glucocorticoid receptor agonist
6560	25.98	isoflupredone	Glucocorticoid receptor agonist
6962	19.87	Clobetasol	Glucocorticoid receptor agonist
7001	19.45	fludrocortisone	Glucocorticoid receptor agonist
7129	16.81	Halcinonide	Glucocorticoid receptor agonist
8126	-10.56	flumetasone	Glucocorticoid receptor agonist
8351	-40.7	hydrocortisone	Glucocorticoid receptor agonist
8433	-53.42	Diflorasone	Corticosteroid agonist

Fig 3. Comparison of epinephrine and norepinephrine gene-expression scores as glucocorticoid
receptor agonists.
The horizontal lines are the median values for the individual distributions. Mann-Whitney analysis of
the distributions showed a statistically significant difference (p <0.0001), with strong association for
epinephrine with gene-expression similarity to glucocorticoid agonists.

It is of particular interest, and unexpected, that the perturbagen class output for epinephrine

(Table 4) included leucine rich repeat kinase (LRRK) inhibitors as showing gene-expression

connectivity to epinephrine with a strong positive score.   Among the agents in the CLUE database,

499
500
501

LRRK2 inhibitors, XMD-1150, Broad Institute ID (BRD)- K01436366 and XMD-885 (BRD-K64857848), ranked with connectivity scores to epinephrine of 90.78 and 90.47, respectively. LRRK2 inhibitors are currently a major focus as potential treatments for PD [45-48]. Table 4 also includes a positive score for gene-expression connectivity with epinephrine for vascular endothelial growth factor receptor (VEGFR) inhibitors; this will be commented upon below. Quite remarkably, the CLUE output in Table 4 also includes a positive score for connectivity of epinephrine with rapidly accelerated fibrosarcoma (RAF) inhibitors. The proto-oncogene, *B-RAF*, is part of the pathway of *RAF* activation of many cancers [49]. Although this could be of considerable interest in relation to possible anti-neoplastic activity of β2AR agonists, it was not pursued in the current study.

It can be noted that the positive rankings for LRRK2, VEGFR, and RAF inhibitors with epinephrine were not identified with gene expression connectivity to norepinephrine (Table 5), which is consistent with the association of these properties with β2- adrenergic activity.

The CLUE analyses of a marked difference between the associations of epinephrine and norepinephrine with gene signatures that are predictive of GR agonist activity are parallel with the established biological differences between the agonist activities of epinephrine and norepinephrine (Table 8). Epinephrine and norepinephrine have essentially equal α1-, α2- and β1-adrenergic agonist activity, but epinephrine is substantially more active than norepinephrine as a β2-adrenergic agonist. The evidence outlined above that supports an anti-inflammatory activity of β2-adrenergic agonists, and a potentially therapeutic effect in relation to the neuroinflammation aspect of PD, is a major focus of these investigations. Also, the fact that GR agonists are widely recognized for their anti-inflammatory activity adds to the plausibility of these associations with β2-adrenergic activity.

Table 8. Modified from: Goodman and Gilman's The Pharmacological Basis of Therapeutics, 9[th] Edition, 1996. Joel G. Hardman and Lee E. Limbird, Eds. McGraw-Hill, New York. Chapter 6, Neurotransmission, Page 125 (Table 6-3).

Characteristics of Subtypes of Adrenergic Receptors[1]

Receptor	Agonists	Antagonists	Tissue	Responses
Alpha1	Epi $\geq$NE>>Iso Drug example, phenylephrine	Prazosin	Vascular smooth muscle	Contraction
			Genitourinary smooth muscle	Contraction
			Liver	Glycogenolysis; gluconeogenesis
			Intestinal smooth muscle	Hyperpolarization and relaxation
			Heart	Increased contractile force, arrhythmias
Alpha2	Epi $\geq$NE>>Iso Drug example, Clonidine	Yohimbine	Pancreatic islet (β cells)	Decreased insulin secretion
			Platelets	Aggregation
			Nerve terminals	Decreased release of NE
			Vascular Smooth Muscle	Contraction
Beta1	Iso>>Epi=NE Drug example, Dobutamine	Metoprolol	Heart	Increased force and rate of contraction and AV nodal conduction velocity
			Juxtaglomerular cell	Increased renin secretion
Beta2	Iso>>Epi >>>NE Drug example, Terbutaline	ICI 118551 (Inverse agonist)	Smooth muscle (vascular, bronchial, gastrointestinal, and genitourinary)	Relaxation
			Skeletal muscle	Glycogenolysis; uptake of K+
			Liver	Glycogenolysis; gluconeogenesis
Beta3	Iso=NE>Epi	ICI 118551	Adipose tissue	Lipolysis

[1]This table provides examples of drugs that act on adrenergic receptors and of the location of subtypes of adrenergic receptors. Abbreviations: epinephrine (Epi); norepinephrine (NE); isoproterenol (Iso). Epi $\geq$NE>>Iso (Epi equal to/slightly greater than NE, which is greater than Iso); Iso>>Epi=NE (Iso greater than Epi, which is equal to NE); Iso>>Epi>>>NE (Iso greater than Epi, which is much greater than NE); Iso=NE>Epi (Iso equal to NE, which is slightly greater than Epi).

When the natural endogenous human glucocorticoid, cortisol (hydrocortisone), was profiled in 531

CLUE, as expected, it showed strong gene signature connectivity with the synthetic glucocorticoid 532

drugs in the CLUE CMap class (Table 9).   Interestingly, it also ranked the beta-adrenergic receptor 533

agonist class with a high enrichment score, similar to that for epinephrine (Table 4). The detailed

analyses also demonstrated that the 44 synthetic glucocorticoids in the CMap class all had gene

signature connectivity scores above 90.0 with hydrocortisone (Table 10); again, this is similar to the

finding with epinephrine (Table 6).

To summarize, Fig. 4 presents the comparison between the gene expression signatures when

CLUE is probed with epinephrine and hydrocortisone as INDEX compounds (i.e., Table 6 vs. Table

10 results).

Table 9. CMap Class Connectivity: HYDROCORTISONE.

Perturbagen Class (PCL)	Group Enrichment Score
Glucocorticoid receptor agonist x 44	99.89
Cyclooxygenase inhibitor x 6	97.34
Beta-adrenergic receptor agonist x 11	95.88
Adenosine receptor agonist x 6	95.49
Na-K-Cl transporter inhibitor x 3	94.89
Rho associated kinase inhibitor x 4	93.79
MEK inhibitor x 8	92.61

"Sets of compound perturbagens with enrichment scores above 90 (similar) and below -90 (opposing)" in relation to their correspondence with the gene expression signatures of hydrocortisone.

Table 10. Gene-expression connectivity scores for HYDROCORTISONE vs GR agonists.

Rank	Score	Name	Description
1	99.99	hydrocortisone	Glucocorticoid receptor agonist
2	99.93	triamcinolone	Glucocorticoid receptor agonist
3	99.93	budesonide	Glucocorticoid receptor agonist
4	99.93	fluocinonide	Glucocorticoid receptor agonist
5	99.93	fluorometholone	Glucocorticoid receptor agonist
7	99.72	mometasone	Glucocorticoid receptor agonist
8	99.72	fluticasone	Glucocorticoid receptor agonist
9	99.72	fludrocortisone	Glucocorticoid receptor agonist
11	99.61	betamethasone	Glucocorticoid receptor agonist
12	99.61	betamethasone	Glucocorticoid receptor agonist
13	99.58	clocortolone	Glucocorticoid receptor agonist
14	99.58	flunisolide	Cytochrome P450 inhibitor
15	99.58	dexamethasone	Glucocorticoid receptor agonist

16	99.58	clobetasol	Glucocorticoid receptor agonist
17	99.54	isoflupredone	Glucocorticoid receptor agonist
20	99.37	fludroxycortide	Glucocorticoid receptor agonist
21	99.33	prednisolone	Glucocorticoid receptor agonist
22	99.3	beclomethasone	Glucocorticoid receptor agonist
23	99.3	fluticasone	Glucocorticoid receptor agonist
24	99.3	alclometasone	Glucocorticoid receptor agonist
26	99.3	hydrocortisone	Glucocorticoid receptor agonist
27	99.26	fluocinolone	Glucocorticoid receptor agonist
28	99.25	flumetasone	Glucocorticoid receptor agonist
30	99.16	rimexolone	Glucocorticoid receptor agonist
32	99.06	loteprednol	Glucocorticoid receptor agonist
34	98.91	hydrocortisone	Glucocorticoid receptor agonist
35	98.8	halcinonide	Glucocorticoid receptor agonist
36	98.66	triamcinolone	Glucocorticoid receptor agonist
37	98.63	dexamethasone	Glucocorticoid receptor agonist
38	98.59	beclomethasone	Glucocorticoid receptor agonist
39	98.52	prednicarbate	Phospholipase activator
44	98.48	prednisolone	Glucocorticoid receptor agonist
45	98.48	amcinonide	Glucocorticoid receptor agonist
46	98.1	halometasone	Glucocorticoid receptor agonist
48	97.82	methylprednisolone	Glucocorticoid receptor agonist
50	97.68	diflorasone	Corticosteroid agonist
52	97.5	hydrocortisone	Glucocorticoid receptor agonist
53	97.5	fluocinonide	Glucocorticoid receptor agonist
55	97.36	desoximetasone	Glucocorticoid receptor agonist
58	97.33	prednisolone	Glucocorticoid receptor agonist
66	97.15	depomedrol	Glucocorticoid receptor agonist
83	96.69	medrysone	Glucocorticoid receptor agonist
141	95.45	westcort	Glucocorticoid receptor agonist
187	95.1	hydrocortisone	Glucocorticoid receptor agonist

Fig 4. Comparison of epinephrine and hydrocortisone gene-expression scores as glucocorticoid receptor agonists.
The horizontal lines are the median values for the individual distributions, which are essentially identical. Mann-Whitney analysis of the distributions showed no difference in gene-expression characteristics for epinephrine and hydrocortisone as glucocorticoid receptor agonists (p = 0.805).

The gene-product targets for the 44 glucocorticoid analyses in this CMap class are presented in

S1 Table. All members of this class have the glucocorticoid receptor, NR3C1, as a target; some also

have the mineralocorticoid receptor, NR3C2, as a target. Only the natural human glucocorticoid   558

cortisol (hydrocortisone) and the synthetic drugs dexamethasone and amcinonide have the gene   559

product ANXA1 as a target. Annexin A1, also termed lipocortin I, is the product of the gene   560

ANXA1, and appears to primarily serve the anti-inflammatory effect of several glucocorticoids. [50]   561

In addition to the gene-product targets for the glucocorticoids, S1 Table includes the Broad Institute   562

ID numbers for each analysis of gene-expression in the database.   563

It should be noted that the CMap class output for hydrocortisone (Table 9) included a strong   564

group score (92.61) for gene-expression connectivity to mitogen-activated protein kinase kinase   565

(MEK) inhibitors. This kinase is on the pathway to signaling of malignant proliferation and   566

survival, i.e., RAF > MEK > extracellular signal-related kinases (ERK). This is similar to the   567

observations noted above for epinephrine (Table 4) where gene-expression connectivity was   568

observed for inhibitory mediators of the anti-proliferative/anti-neoplastic effects of LRRK2, VEGFR,   569

and RAF. In the case of the MEK inhibitors, the drug with the highest score for the class when   570

probed separately in CLUE was selumetinib (96.51), which has proven to have a broad role as an   571

anti-neoplastic agent [51, 52].   572

The striking similarities in the gene-expression signature connectivity for epinephrine and   573

hydrocortisone drew attention to their known interrelationships in regard to synthesis of cAMP [53-   574

55]. Perez [53] noted that α2 agonist activity mediated a decreased synthesis of cAMP. When the   575

classical non-competitive, irreversible α2 antagonist, phenoxybenzamine, was modeled in CLUE, it   576

also showed considerable gene-expression signature connectivity with GR receptor agonists. The   577

gene enrichment score for phenoxybenzamine with the GR agonist CMap class is shown in Table 11.   578

Table 11. CMap Class Connectivity: PHENOXYBENZAMINE

Perturbagen Class (PCL)	Group Enrichment Score
Progesterone receptor agonist x 12	99.50
Glucocorticoid receptor agonist x 44	98.78
Aromatase inhibitor x 3	97.98
Glycogen synthase kinase inhibitor x 11	97.63
HSP inhibitor x 7	96.76
EGFR inhibitor x 14	96.33
PPAR receptor agonist x 16	95.40
DNA synthesis inhibitor x 3	95.02
PKC inhibitor x 7	93.75
JAK inhibitor x 5	92.59
HIV protease inhibitor x 4	92.34
Cannabinoid receptor agonist x 8	91.13

"Sets of compound perturbagens with enrichment scores above 90 (similar) and below -90 (opposing)" in relation to their correspondence with the gene expression signatures of phenoxybenzamine.

Detailed analysis of the gene-expression connectivity scores for phenoxybenzamine with GR receptor agonists is listed in Table 12.

Table 12. Gene-Expression Connectivity Scores for PHENOXYBENZAMINE vs GR agonists.

Rank	Score	Name	Description
4	99.93	Depomedrol	Glucocorticoid receptor agonist
6	99.93	Halcinonide	Glucocorticoid receptor agonist
8	99.93	Clocortolone	Glucocorticoid receptor agonist
11	99.93	Budesonide	Glucocorticoid receptor agonist
12	99.93	Prednisolone	Glucocorticoid receptor agonist
13	99.93	Fludroxycortide	Glucocorticoid receptor agonist
15	99.93	Fludrocortisone	Glucocorticoid receptor agonist
18	99.93	Amcinonide	Glucocorticoid receptor agonist
20	99.89	Beclomethasone	Glucocorticoid receptor agonist
23	99.82	Fluocinonide	Glucocorticoid receptor agonist
33	99.72	Mometasone	Glucocorticoid receptor agonist
34	99.72	Triamcinolone	Glucocorticoid receptor agonist
46	99.51	Beclomethasone	Glucocorticoid receptor agonist
55	99.44	Prednisolone	Glucocorticoid receptor agonist
113	98.27	Fluticasone	Glucocorticoid receptor agonist
114	98.27	Isoflupredone	Glucocorticoid receptor agonist
127	98.13	Westcort	Glucocorticoid receptor agonist
141	97.74	Hydrocortisone	Glucocorticoid receptor agonist
163	97.53	flunisolide	Cytochrome P450 inhibitor
167	97.5	Flumetasone	Glucocorticoid receptor agonist

180	97.38	Clobetasol	Glucocorticoid receptor agonist
221	96.45	Rimexolone	Glucocorticoid receptor agonist
291	95.26	Loteprednol	Glucocorticoid receptor agonist
347	93.94	Fluocinolone	Glucocorticoid receptor agonist
598	89.54	Prednisolone	Glucocorticoid receptor agonist
769	87.05	Alclometasone	Glucocorticoid receptor agonist
779	86.97	Diflorasone	Corticosteroid agonist
814	86.66	Betamethasone	Glucocorticoid receptor agonist
882	85.73	Prednicarbate	Phospholipase activator
953	84.66	Hydrocortisone	Glucocorticoid receptor agonist
1310	79.4	Betamethasone	Glucocorticoid receptor agonist
1637	74.67	Hydrocortisone	Glucocorticoid receptor agonist
1655	74.17	Fluorometholone	Glucocorticoid receptor agonist
1702	73.53	Hydrocortisone	Glucocorticoid receptor agonist
1719	73.16	Triamcinolone	Glucocorticoid receptor agonist
1836	71.31	Desoximetasone	Glucocorticoid receptor agonist
1930	69.94	Dexamethasone	Glucocorticoid receptor agonist
2543	61.53	fluticasone	Glucocorticoid receptor agonist
2665	59.91	Halometasone	Glucocorticoid receptor agonist
3383	47.82	Dexamethasone	Glucocorticoid receptor agonist
3419	47.37	Fluocinonide	Glucocorticoid receptor agonist
4447	33.18	Hydrocortisone	Glucocorticoid receptor agonist
4539	31.79	methylprednisolone	Glucocorticoid receptor agonist
4566	31.68	Medrysone	Glucocorticoid receptor agonist

The strong gene-expression connectivity of phenoxybenzamine with GR agonist activity and the 589

potential anti-neoplastic properties of phenoxybenzamine (Table 11) have been presented in previous 590

reports [56, 57]. 591

The associations between agents that increase cAMP levels and GR receptor agonist activity 592

were investigated further by modeling two methylxanthine drugs, theophylline and caffeine, in 593

CLUE. Theophylline has a long therapeutic history as a bronchodilator in the treatment of 594

respiratory diseases. As discussed above, it does not directly increase synthesis of cAMP, but 595

results in higher levels of cellular cAMP by inhibition of phosphodiesterase enzymes that metabolize 596

cAMP. Caffeine has similar activity but is a much more potent CNS stimulant than a peripheral 597

agonist. 598

The CMap class output scores from CLUE for theophylline are presented in Table 13.   Gene-expression connectivity scores vs GR agonists are presented in S2 Table.

Table 13. CMap Class Connectivity: THEOPHYLLINE

Perturbagen Class (PCL)	Group Enrichment Score
Dihydrofolate reductase inhibitor x 4	99.78
Reverse transcriptase inhibitor x 4	99.40
Glucocorticoid receptor agonist x 44	98.37
Vitamin D receptor agonist x 6	97.36
Nucleoside reverse transcriptase inhibitor x 3	97.07
IMPDH inhibitor x 3	95.65
Bacterial 30S ribosomal subunit inhibitor x 5	94.42
FXR antagonist x 3	93.95
Calmodulin antagonist x 3	93.51
Bacterial cell wall synthesis inhibitor x 9	90.77

"Sets of compound perturbagens with enrichment scores above 90 (similar) and below -90 (opposing)" in relation to their correspondence with the gene expression signatures of theophylline.

The CMap class output scores from CLUE for caffeine are presented in Table 14. The GR agonist class scored substantially lower for caffeine than for theophylline.   Gene-expression connectivity scores vs GR agonist are presented in S3 Table.

Table 14. CMap Class Connectivity: CAFFEINE.   613

Perturbagen Class (PCL)	Group Enrichment Score
Bacterial DNA gyrase inhibitor x 5	95.54
Aromatase inhibitor x 3	93.32
Dihydrofolate reductase inhibitor x 4	91.76
Cyclooxygenase inhibitor x 6	90.65
Glucocorticoid receptor agonist x 44	90.24

"Sets of compound perturbagens with enrichment scores above 90 (similar) and below -90   614
(opposing)" in relation to their correspondence with the gene expression signatures of caffeine.   615

A summary of agents that have been investigated in relation to their gene-expression signatures   618

with connectivity to GR agonists is presented in Figure 5.   619

Fig 5. Comparison of beta2-adrenergic related agents and hydrocortisone for gene-expression   621
connectivity scores as glucocorticoid receptor agonists.   622
The horizontal lines are the median values for the individual distributions. The relative differences   623
among the gene-expression scores can be appreciated visually. Caffeine was not studied further.   624
625

Epinephrine, a potent β2AR agonist has been presented as a prototype for drugs of this class in   627

these studies because levalbuterol and arformoterol have not been profiled for gene-expression   628

connectivity in CLUE.   In July 2020, as our interest in levalbuterol advanced, we formally   629

nominated it to the Broad Institute for gene-expression connectivity study in the CLUE database   630

based on its apparent anti-inflammatory potency [7]; it remained on the queue for possible testing for   631

several months but was never screened.   632

It would appear to be significant that the drugs noted in Figure 5 that have gene-expression   633

signatures with connectivity to GR agonists are all capable of mediating an increase in cellular   634

cAMP. Since increases in cAMP are a characteristic of drugs that mediate anti-inflammatory effects,   635

it appears that this may represent a unifying hypothesis for this study.   636

3.2. CRS assays of agents with β2-adrenergic and related activities

The CRS assay was used to test the efficacy of the several β2-adrenergic related drugs discussed above for possible relationships between predictions generated in the CLUE genomic analyses and a biological test of anti-inflammatory activity in the humanized mouse assay. All animal assays were carried out in the Jackson Laboratories facility in Sacramento, California.

Several of the drugs of interest were tested for their ability to modulate cytokine release *in vivo* after the mice were challenged with injections of the monoclonal antibodies OKT3 or aCD28. The application of an *in vivo* model to simulate a clinical translation was considered a feature of these investigations. The drugs were administered in the drinking water. They are all water soluble and are known to have good bioavailability by the oral route. Doses were calculated from the standard human clinical range of mg/kg body weight per day. The FDA factor of 12.3 was applied for conversion from human to mouse mg/kg dose. A mouse body weight of 20 grams and an average intake of drinking water of 4 ml/day were assumed. The resulting doses were as follows: Albuterol sulfate, 8 µg/ml drinking water; levalbuterol HCl, 4 µg/ml; phenoxybenzamine HCl, 5 µg/ml; theophylline, 400 µg/ml; and arformoterol tartrate, 0.008 µg/ml.

Mice treated with levalbuterol showed a statistically significant lower body weight loss than the control group (1% sucrose) when CRS was induced with OKT3 (Figure 6).

Fig 6. Percent body weight loss calculated from each animal's initial body weight at Study Day 0.
Body weight loss was statistically significantly less for the group that was pretreated with levalbuterol and challenged with the monoclonal antibody, OKT3.

There were no statistically significant differences in the clinical scoring between any of the groups and the control group (Figure 7). All 90 animals included in the study survived to the time of blood collection by cardiac puncture on Study Day 9, 72 hours after toxic induction with OKT3 or aCD28. This would suggest that the doses translated from human to mouse were reasonably appropriate.

Fig 7. **CRS clinical scores for the period following toxic induction of cytokine release with OKT3 or aCD28.**

There were no statistically significant differences in the clinical scores that were attributable to the pretreatments; the doses of pretreatments were calculated from typical human doses that were translated using FDA factors.

Water consumption in both the OKT3 and anti-CD28 groups peaked on Study Day 6 before toxicity was induced and declined progressively to Study Day 9 (Figure 8). Analysis of the data was limited since water consumption was made by the cage for each group of five mice. However, one individual comparison (Wilcoxon signed rank test) for water consumption for levalbuterol vs. sucrose control in the OKT3 groups did trend for a statistical advantage favoring more water intake for levalbuterol ($p = 0.068$). (Data for theophylline groups were accidentally not recorded.)

Fig 8. Water consumption recorded per cage of 5 mice as a percentage of the intake on Study Day 6 prior to the induction challenges with OKT3 and anti-CD28.

The toxic antibody challenges to induce cytokine release caused marked decreases in water consumption in all of the groups. Since the pretreatment drugs were administered in the drinking water, it would be expected that drug delivery was compromised.

Analyses of the effects of the beta2 adrenergic agonists and related compounds on the release of cytokines and chemokines 6 hours after induction with OKT3 in the CRS assay demonstrated the following primary outcomes: Figure 9 presents the results for the inhibition of the release of the eosinophil-selective chemokine, eotaxin, which is responsible for a number of eosinophilic disorders, and the angiogenic proto-oncogene, vascular endothelial growth factor (VEGF), by several of the β2-adrenergic related drugs. Albuterol, the racemic mixture of the levo and dextro isomers, was not different from the sucrose control in any of the assays. It should be noted that albuterol was administered at twice the concentration of levalbuterol, therefore the mice received the same dose of the levo isomer as those receiving the pure levalbuterol. However, as noted above, the dextro isomer is pro-inflammatory and interferes with the anti-inflammatory activity of the levo isomer. Levalbuterol also inhibited the release of the pro-inflammatory cytokine, IL-13 with OKT3 induction (Figure 9). The relationships between the treatment drugs and their gene-expression connectivity to glucocorticoid receptor agonist activity is also included in Figure 9.

Fig 9. The effects of β2-adrenergic related drugs on the release of eotaxin and VEGF.

Elaboration of Eotaxin (A), VEGF (B), and IL-13 (C), in the CRS assay. D, comparisons are illustrated in the same order for the gene-expression connectivity of the agents as glucocorticoid receptor agonists in the CLUE database. As noted, the gene-expression connectivity for epinephrine, the prototypical beta2-adrenergic agonist, is substituted for levalbuterol and arfomoterol since these drugs were never profiled in the CLUE database. Albuterol, the racemic mixture of R and S levalbuterol, showed no inhibition of cytokine/chemokine release in any of the assays (A, B, C).

The changes seen in cytokine and chemokine inhibition at 6 hours after OKT3 induction did not persist until assays on Day 9, 72 hours after induction. A possible explanation may be related to the fact that water intake containing the adrenergic-related pretreatments was reduced by approximately 55 to 70 percent among the treatments by Day 9 (Figure 8).

The results for eotaxin and VEGF with levalbuterol (Fig. 9) showed that some of the other beta2-adrenergic related drugs were quite close to statistical differences from their controls, so a simple "power analysis" was done by duplicating the data to double the sample size to 10 mice/group; the variance was kept the same. All the drug treatments except albuterol showed significantly less elaboration of VEGF and eotaxin with these increases in sample size (Figure 10).

Fig 10. "Power" analysis of the results for the VEGF and eotaxin assays.
Duplication of data from 5-animal CRS assays but variance unchanged.

Possible trends toward statistical significance were examined for all the cytokines and chemokines that were measured. Unpaired t-tests, without correction for multiple comparisons, identified the following inhibitory effects for levalbuterol on cytokine elaboration after OKT3 induction: TNFα, $p = 0.0321$; TNFβ, $p = 0.0373$; M-CSF1, $p = 0.0342$; IL-1α, 0.0399; and IL-18, $p = 0.0003$.

The MILLIPLEX panel used to quantify cytokines in the above assays identified eotaxin-1 (CCL11), which is the most extensively characterized and studied of the eotaxins. When eotaxin links to its most specific receptor, CCR3, in organs and the brain it recruits eosinophils that upon degranulation release eosinophil granule proteins, growth factors and cytokines causing cellular damage; eotaxin plays a central role in mediating a number of eosinophilic disorders, including

atopic dermatitis, eosinophilia in asthma, chronic rhinosinusitis and eosinophilic esophagitis [58, 59]. ₇₃₆

Teixeira et al. [60] have reviewed the association of increased blood levels of eotaxin-1/CCL11 in ₇₃₇

patients with major psychiatric disorders including schizophrenia, major depression, bipolar disease, ₇₃₈

autism spectrum disorder, dysthymia, and Alzheimer disease; they also discussed the presence of the ₇₃₉

CCR3 receptor on microglia, and Peterson et al. [11] have proposed a pro-inflammatory role of ₇₄₀

activated microglia in destruction of dopaminergic neurons in PD. Chandra et al. demonstrated that ₇₄₁

eotaxin is markedly elevated in the substantia nigra pars compacta in post-mortem brains of PD ₇₄₂

patients compared to non-affected controls [61]. ₇₄₃

The finding that IL-13 release was also inhibited by levalbuterol (Figure 9) is of interest because ₇₄₄

IL-13 is one of the cytokines that is central to cellular trafficking; blocking its ligands is a feature of ₇₄₅

dupilumab in the treatment of chronic obstructive pulmonary disease (COPD) and other eosinophilic ₇₄₆

disorders [59, 62]. ₇₄₇

Figure 9 also presents the results for the inhibition of the release of VEGF by levalbuterol and ₇₄₈

theophylline. There is a vast literature on the importance of VEGF in neoplastic disease including an ₇₄₉

earlier comprehensive review by Goel and Mercurio [63], and most recent efforts to block certain ₇₅₀

particularly aggressive cancers that depend on binding of VEGF to a co-factor, neuropillin-2 [64]. ₇₅₁

The MILLIPLEX panel that was used to quantify this cytokine in the CRS assays was specific for ₇₅₂

the $VEGF_{165}$ variant. Although it is many decades since the recognition of the involvement of VEGF ₇₅₃

in tumor angiogenesis, treatment complexity persists. Mabeta and Steenkamp [65] have presented a ₇₅₄

recent review of VEGF variants and $VEGF_{165}$ is the prototypical cytokine and is upregulated in ₇₅₅

approximately 70% of tumors including breast cancer, glioblastoma multiforme, ovarian cancer, ₇₅₆

melanoma, esophageal cancer, gastrointestinal tract cancers, lung cancer, renal carcinomas and ₇₅₇

others. $VEGF_{165}$ promotes sprouting of existing vessels as well as splitting of vessels as part of its ₇₅₈

angiogenic effects; these processes may be part of an effort by tumors to escape therapeutic ₇₅₉

interventions that produce conditions jeopardizing tumor expansion [65]. It is interesting to speculate ₇₆₀

that inhibition of release of VEGF, as seen in Figure 9, may be superior to the complexities of 761

control of angiogenesis at the tumor level. 762

3.3. Relevant observations 763

The marked similarity between gene-expression connectivity for epinephrine (and presumably 764

other β2AR agonists) and glucocorticoids, as presented in these investigations, raises the question as 765

to why β2AR agonists have shown absolutely no evidence of the classical adverse effects of 766

glucocorticoids. This is true with chronic treatment of β2AR agonists at pharmacological doses in 767

their primary role as bronchodilators. Adverse side effects of glucocorticoids include hypertension, 768

hyperglycemia, osteoporosis, glaucoma, cataract formation, peptic ulcer, gastrointestinal bleeding, 769

and others [66]. These effects are dose dependent and reduction in dosages leads to loss of 770

therapeutic effectiveness. 771

Dose limitations represent a major disadvantage in the clinical use of glucocorticoids. In 772

addition to their efficacy as anti-inflammatory/immunomodulatory agents, these drugs have major 773

anti-neoplastic effects in the treatment of several hematopoietic malignancies with lymphatic lineage, 774

including chronic lymphocytic leukemia, acute lymphoblastic leukemia, multiple myeloma, 775

Hodgkin's lymphoma, and non-Hodgkin's lymphoma [67, 68].   Induction of apoptosis appears to be 776

a primary mechanism in the treatment of these malignancies [69]. 777

There has been considerable effort directed in attempts to separate the desired effects of 778

glucocorticoids from the initiation of adverse events.   Agents of this type are non-steroidal 779

chemically and have been variously termed as "dissociated glucocorticoid receptor ligands" [70], 780

"selective glucocorticoid receptor agonists" (SEGRAs) [71], or examples of "biased signaling" [72]. 781

Kleiman and Tuckerman have summarized the results with some of the biological trials with 782

SEGRAs and they have shown limited success [73]. 783

The fact that epinephrine, the prime example of a β2AR agonist, showed exactly the same gene- 784

expression profile to hydrocortisone in the CLUE analyses (Tables 6 and 10; Fig. 4) but is not 785

associated with any of the adverse effects of glucocorticoids is now known with confidence to be related to the gene-signaling pathways from different receptor ligands. The biological result is related to which gene-signaling pathways are favored [74].

An updated comprehensive review of the glucocorticoid receptor by Nicolaides et al. [75] discusses the two major classifications of the signaling of gene expression in relation to the potential for adverse events with longer treatment and higher doses versus the pathway for therapeutic predominance. The contrasting pathways are termed "transactivation" and "transrepression." Transactivation correlates with downstream transcription of protein sequences that mediate the toxic manifestations of glucocorticoid treatment (diabetes mellitus, osteoporosis, hypertension, etc.). Transrepression is primarily associated with therapeutic effects of glucocorticoids, i.e., their anti-inflammatory, immunomodulatory, and certain anti-neoplastic effects. The mechanisms of transactivation and transrepression are complex and are discussed in extensive detail by Lesovaya et al [76]. In view of the potential anti-inflammatory effects of β2AR-related drugs observed in the present study, it would be expected that they mediate via a transrepression effect at the GR.

The chapter by Nicolaides et al. [75] also discusses the evolution of the glucocorticoid receptor, which apparently represents an important step in the survival advantage of a branch that ultimately led to mammalian species. That step, which included specificity of GR for cortisol, took place about 450 million years ago with the first appearance of the GR in ray-finned fish and then to land vertebrates. Of interest to the present study, the evolution of the sympathetic nervous system took place in that same period of geologic time, and it is presumed that a sympathetic nervous system represented a level of functional sophistication that contributed to survival to further evolution [77].

The adrenal steroids and sympathetic nervous systems are critical parts of the integrated response to stress in humans today and this may be the reason that the CLUE platform of the Harvard/MIT Broad Institute database showed essentially the same gene-expression signatures for hydrocortisone and epinephrine, as noted above. In these integrated systems, glucocorticoids have

been demonstrated to promptly increase elaboration of cAMP by several mechanisms that would be expected to inhibit the release of pro-inflammatory cytokines and chemokines: *a*) Hydrocortisone (cortisol) directly activates adenylyl cyclase to elaborate cAMP; *b*) it increases the density of β2ARs on cell membranes of elements of the innate immune system; *c*) it facilitates linking of those receptors to adenylyl cyclase to increase cAMP; *d*) it preserves responsiveness (i.e. reverses tachyphylaxis) that may develop with continuous occupancy of β2ARs [54, 55, 78-82]; *e*) and finally, there is the added fact that, as part of the stress response, hydrocortisone selectively activates the phenylethanolamine N-methyltransferease (PNMT) enzyme in the adrenal medulla to increase the synthesis of epinephrine [75, 83, 84]. It is interesting to speculate that the selective beta2-adrenergic-related agents studied in this investigation may extend the clinically therapeutic applications of the glucocorticoid-sympathetic stress response without incurring the adverse effects of higher glucocorticoid doses.

In particularly interesting observations by Flydal et al. [85] levalbuterol was identified as a small molecular "chaperone" that stabilized the enzyme, tyrosine hydroxylase. This enzyme catalyzes the rate-limiting step in the synthesis of dopamine. It is depleted or dysfunctional in the tyrosine hydroxylase deficiency syndrome and in PD. In an effort to repurpose treatments for these deficiencies, these investigators screened 1280 recognized drugs for their abilities to prevent excessive aggregation and/or misfolding of the tyrosine hydroxylase molecule, and to reduce the feedback inhibitory effects of dopamine on the activity of the enzyme. Through active site modeling, levalbuterol was proposed to bind to the same iron moiety as dopamine at the active site. However, binding by levalbuterol did not inhibit activity but did competitively reduce the feedback inhibitory effects of dopamine. These properties of levalbuterol may have value in reducing the dyskinesias associated with the chronic treatment of PD. Nineteen drugs from the initial screening showed promise but validation studies eliminated all but 4. Of those, levalbuterol demonstrated dose-related efficacy most consistently [85].

3.4. Clinical translation of the findings

The drugs studied in the present work are all FDA-approved medications, which could facilitate their possible translation to therapies that have relevance to neurodegenerative and neoplastic disorders. There is considerable clinical data regarding the use of levalbuterol and other beta2-adrenergic selective agents for the treatment of bronchial asthma in children and adults. The doses that were used in the present studies to test their effects on cytokine release in the human PBMC-engrafted mice were translated from human adult doses that relieve asthmatic attacks. Heart rate changes were not monitored in the mice but there were no untoward effects noted. In a clinical comparison between levalbuterol and the racemic drug albuterol in children aged 4 to 11 years in age, levalbuterol appeared to be superior in terms of improvement in respiratory function indices [86]. Children who received the recommended dose of levalbuterol for this age group (0.31 mg by nebulized inhalation) showed no mean change in heart rate; those that required double this dose for a satisfactory treatment of an asthmatic attack had a mean increase in heart rate of approximately 7 beats per minute. It must be noted that this route of administration essentially delivers the entire dose to the lungs where there is rapid systemic absorption that impacts the heart; this would not be the case if the drug was administered episodically by the oral route. It was administered continuously in drinking water in the mouse studies. A recent meta-analysis covering the use of this class of drugs from 1996 – 2021 demonstrated their safety [87].

Clinical translation would also be aided by the fact that there are no formulation issues for levalbuterol. There are three FDA-approved dose levels that could be used for episodic administration orally.

3.5. Limitations

An unfortunate limitation in the studies, as noted above, was the fact that levalbuterol and arformoterol have never been profiled for gene-expression connectivity to glucocorticoids by the

Harvard/MIT Broad Institute. We did formally nominate it for assay in CLUE, but this has not taken place. This remains to be addressed in future studies.

Another limitation was that we were not able to assay the proto-typical beta2-adrenergic agonist, epinephrine, in the mouse CRS assay. The assay was designed to provide a low-stress delivery of pretreatments to the mice by providing the drugs in the drinking water. Epinephrine would undergo autoxidation under these conditions.

4. Conclusions

Advances in the development of beta2-selective adrenergic agonists has led to their recognition as potent anti-inflammatory agents with potential for treatment in Parkinson's disease and inflammatory disorders. This potential was reinforced in the present study by observations that epinephrine, the prototype beta2-adrenergic agonist, showed an almost identical gene expression signature as cortisol (hydrocortisone) and other glucocorticoids.

Levalbuterol and other beta2-adrenergic-related agents inhibited the release of eotaxin, VEGFa and IL-13 from mice engrafted with human peripheral blood monocyte cultures in a Jackson Laboratory assay, supporting the predictions of similar gene expression mechanisms and potential translation to inflammatory pathologies and neoplastic syndromes. The observations with the group of drugs that were studied suggested a unifying mechanism of action of maintenance or elevation in cellular levels of cAMP. This aspect requires further investigation.

While therapeutic efforts in neurodegenerative, inflammatory, and neoplastic diseases that block downstream pathways and receptors have advanced considerably, there remain the issues of development of tolerance, escape mechanisms and off-target adverse effects. There may be a philosophical advantage to inhibition of the release of inciting elements in these pathologies, as suggested by the present studies.

The agents studied are FDA-approved drugs with minimal adverse effect profiles. The fact that they showed no toxicity in the animal studies at doses translated from human experience, and that

there are existing formulations that can be administered orally, may facilitate their repurposing and 885

ultimate translation to appropriate pathologies. 886

Supporting Information 888

S1 Table: Glucocorticoid receptor agonist list with targets. 889

S2 Table**:** Gene expression connectivity scores for THEOPHYLLINE vs GR agonists. 890

S3 Table: Gene expression connectivity scores for CAFFEINE vs GR agonists 891

S1 Figure: Association between rank score and gene-expression connectivity score. 892

Author Contributions: The author conceived the investigation, carried out bioinformatics 893

studies, contributed to the design of the animal assays, and wrote the manuscript. 894

Funding: The study was part of an academic collaboration between New York Medical College, 895

Valhalla, NY and Jackson Laboratories, Bar Harbor, ME. Some costs were covered by New York 896

Medical College with funds contributed by a private donor. 897

Data availability: Essentially all the data are included in the manuscript. Body weights for 898

individual animals and water consumption data by cage are available upon request. 899

Acknowledgments: The author is grateful to Ms. Jennifer Brown for editorial assistance in the 900

preparation of the manuscript, and to Ms. Brown and Ms. Gail Anderson of the Department of 901

Pharmacology, New York Medical College, for facilitating the administrative and financial 902

procedures required for the studies. The author also recognizes the contributions of Ms. Maribeth 903

Balingit and associates of the Phillip Capozzi, M.D. Library, New York Medical College, for 904

assistance in assembling the bibliography of the manuscript. 905

Conflicts of Interest: The author declares no conflicts of interest. 906

References

1. Magistrelli L, Comi C. Beta2-Adrenoceptor Agonists in Parkinson's Disease and Other Synucleinopathies. J Neuroimmune Pharmacol. 2020;15(1):74-81.

2. Hishida R, Kurahashi K, Narita S, Baba T, Matsunaga M. "Wearing-off" and beta 2-adrenoceptor agonist in Parkinson's disease. Lancet. 1992;339(8797):870.

3. Alexander GM, Schwartzman RJ, Nukes TA, Grothusen JR, Hooker MD. Beta 2-adrenergic agonist as adjunct therapy to levodopa in Parkinson's disease. Neurology. 1994;44(8):1511-3.

4. Uc EY, Lambert CP, Harik SI, Rodnitzky RL, Evans WJ. Albuterol improves response to levodopa and increases skeletal muscle mass in patients with fluctuating Parkinson disease. Clin Neuropharmacol. 2003;26(4):207-12.

5. Mittal S, Bjornevik K, Im DS, Flierl A, Dong X, Locascio JJ, et al. beta2-Adrenoreceptor is a regulator of the alpha-synuclein gene driving risk of Parkinson's disease. Science. 2017;357(6354):891-8.

6. Devine MJ, Gwinn K, Singleton A, Hardy J. Parkinson's disease and alpha-synuclein expression. Mov Disord. 2011;26(12):2160-8.

7. Wang S, Liu F, Tan KS, Ser HL, Tan LT, Lee LH, et al. Effect of (R)-salbutamol on the switch of phenotype and metabolic pattern in LPS-induced macrophage cells. J Cell Mol Med. 2020;24(1):722-36.

8. Friere M, Pergolizzi R, Millon C, Dominguez PJ. Cytokine, chemokine, and nitric oxide release in stimulated small airway epithelial cells treated with β2 agonist enantiomers of salbutamol. J Allergy Clin Immunolo. 2000;105.

9. Severn A, Rapson NT, Hunter CA, Liew FY. Regulation of tumor necrosis factor production by adrenaline and beta-adrenergic agonists. J Immunol. 1992;148(11):3441-5.

10. Elenkov IJ, Chrousos GP. Stress hormones, proinflammatory and antiinflammatory cytokines, and autoimmunity. Ann N Y Acad Sci. 2002;966:290-303.

11. Peterson L, Ismond KP, Chapman E, Flood P. Potential benefits of therapeutic use of beta2-adrenergic receptor agonists in neuroprotection and Parkinsonmus disease. J Immunol Res. 2014;2014:103780.

12. Elenkov IJ, Wilder RL, Chrousos GP, Vizi ES. The sympathetic nerve--an integrative interface between two supersystems: the brain and the immune system. Pharmacol Rev. 2000;52(4):595-638.

13. Sharma M, Flood PM. beta-arrestin2 regulates the anti-inflammatory effects of Salmeterol in lipopolysaccharide-stimulated BV2 cells. J Neuroimmunol. 2018;325:10-9.

14. Mazzoni L, Naef R, Chapman ID, Morley J. Hyperresponsiveness of the airways to histamine following exposure of guinea pigs to racemic mixtures and distomers of β2-selective sympathomimetics. Pulm Pharmacol. 1994;7.

15. Ibe BO, Abdallah MF, Raj JU. Mechanisms by which S-albuterol induces human bronchial smooth muscle cell proliferation. Int Arch Allergy Immunol. 2008;146(4):321-33.

16. Baramki D, Koester J, Anderson AJ, Borish L. Modulation of T-cell function by (R)- and (S)-isomers of albuterol: anti-inflammatory influences of (R)-isomers are negated in the presence of the (S)-isomer. J Allergy Clin Immunol. 2002;109(3):449-54.

17. Chorley BN, Li Y, Fang S, Park JA, Adler KB. (R)-albuterol elicits antiinflammatory effects in human airway epithelial cells via iNOS. Am J Respir Cell Mol Biol. 2006;34(1):119-27.

18. Gronich N, Abernethy DR, Auriel E, Lavi I, Rennert G, Saliba W. beta2-adrenoceptor agonists and antagonists and risk of Parkinson's disease. Mov Disord. 2018;33(9):1465-71.

19. Chen CL, Wang SY, Chen TC, Chuang CS. Association between beta2-Adrenoreceptor Medications and Risk of Parkinson's Disease: A Meta-Analysis. Medicina (Kaunas). 2021;57(10).

20. Moore CG, Schenkman M, Kohrt WM, Delitto A, Hall DA, Corcos D. Study in Parkinson disease of exercise (SPARX): translating high-intensity exercise from animals to humans. Contemp Clin Trials. 2013;36(1):90-8.

21. Schenkman M, Moore CG, Kohrt WM, Hall DA, Delitto A, Comella CL, et al. Effect of High-Intensity Treadmill Exercise on Motor Symptoms in Patients With De Novo Parkinson Disease: A Phase 2 Randomized Clinical Trial. JAMA Neurol. 2018;75(2):219-26.

22. Kjaer M. Epinephrine and some other hormonal responses to exercise in man: with special reference to physical training. Int J Sports Med. 1989;10(1):2-15.

23. Baron JA. Cigarette smoking and Parkinson's disease. Neurology. 1986;36(11):1490-6.

24. Mappin-Kasirer B, Pan H, Lewington S, Kizza J, Gray R, Clarke R, et al. Tobacco smoking and the risk of Parkinson disease: A 65-year follow-up of 30,000 male British doctors. Neurology. 2020;94(20):e2132-e8.

25. Fratiglioni L, Wang HX. Smoking and Parkinson's and Alzheimer's disease: review of the epidemiological studies. Behav Brain Res. 2000;113(1-2):117-20.

26. Allam MF, Campbell MJ, Hofman A, Del Castillo AS, Fernandez-Crehuet Navajas R. Smoking and Parkinson's disease: systematic review of prospective studies. Mov Disord. 2004;19(6):614-21.

27. Hernan MA, Takkouche B, Caamano-Isorna F, Gestal-Otero JJ. A meta-analysis of coffee drinking, cigarette smoking, and the risk of Parkinson's disease. Ann Neurol. 2002;52(3):276-84.

28. Cryer PE, Haymond MW, Santiago JV, Shah SD. Norepinephrine and epinephrine release and adrenergic mediation of smoking-associated hemodynamic and metabolic events. N Engl J Med. 1976;295(11):573-7.

29. Grassi G, Seravalle G, Calhoun DA, Bolla G, Mancia G. Cigarette smoking and the adrenergic nervous system. Clin Exp Hypertens A. 1992;14(1-2):251-60.

30. Grassi G, Seravalle G, Calhoun DA, Bolla GB, Giannattasio C, Marabini M, et al. Mechanisms responsible for sympathetic activation by cigarette smoking in humans. Circulation. 1994;90(1):248-53.

31. Ragonese P, Salemi G, Morgante L, Aridon P, Epifanio A, Buffa D, et al. A case-control study on cigarette, alcohol, and coffee consumption preceding Parkinson's disease. Neuroepidemiology. 2003;22(5):297-304.

32. Tan EK, Tan C, Fook-Chong SM, Lum SY, Chai A, Chung H, et al. Dose-dependent protective effect of coffee, tea, and smoking in Parkinson's disease: a study in ethnic Chinese. J Neurol Sci. 2003;216(1):163-7.

33. Ascherio A, Zhang SM, Hernan MA, Kawachi I, Colditz GA, Speizer FE, et al. Prospective study of caffeine consumption and risk of Parkinson's disease in men and women. Ann Neurol. 2001;50(1):56-63.

34. Palacios N, Gao X, McCullough ML, Schwarzschild MA, Shah R, Gapstur S, et al. Caffeine and risk of Parkinson's disease in a large cohort of men and women. Mov Disord. 2012;27(10):1276-82.

35. Qi H, Li S. Dose-response meta-analysis on coffee, tea and caffeine consumption with risk of Parkinson's disease. Geriatr Gerontol Int. 2014;14(2):430-9.

36. Ren X, Chen JF. Caffeine and Parkinson's Disease: Multiple Benefits and Emerging Mechanisms. Front Neurosci. 2020;14:602697.

37. Robertson D, Frolich JC, Carr RK, Watson JT, Hollifield JW, Shand DG, et al. Effects of caffeine on plasma renin activity, catecholamines and blood pressure. N Engl J Med. 1978;298(4):181-6.

38. Benowitz NL, Jacob P, 3rd, Mayan H, Denaro C. Sympathomimetic effects of paraxanthine and caffeine in humans. Clin Pharmacol Ther. 1995;58(6):684-91.

39. Subramanian A, Narayan R, Corsello SM, Peck DD, Natoli TE, Lu X, et al. A Next Generation Connectivity Map: L1000 Platform and the First 1,000,000 Profiles. Cell. 2017;171(6):1437-52 e17.

40. Shultz LD, Keck J, Burzenski L, Jangalwe S, Vaidya S, Greiner DL, et al. Humanized mouse models of immunological diseases and precision medicine. Mamm Genome. 2019;30(5-6):123-42.

41. Ye C, Yang H, Cheng M, Shultz LD, Greiner DL, Brehm MA, et al. A rapid, sensitive, and reproducible in vivo PBMC humanized murine model for determining therapeutic-related cytokine release syndrome. FASEB J. 2020;34(9):12963-75.

42. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. Proc Natl Acad Sci U S A. 2005;102(43):15545-50.

43. Lanau F, Zenner MT, Civelli O, Hartman DS. Epinephrine and norepinephrine act as potent agonists at the recombinant human dopamine D4 receptor. J Neurochem. 1997;68(2):804-12.

44. Wu Y, Zeng L, Zhao S. Ligands of Adrenergic Receptors: A Structural Point of View. Biomolecules. 2021;11(7).

45. Xiao B, Tan EK. Targeting LRRK2 in Parkinson's disease. Cell Rep Med. 2022;3(10):100778.

46. Cresto N, Gardier C, Gubinelli F, Gaillard MC, Liot G, West AB, et al. The unlikely partnership between LRRK2 and alpha-synuclein in Parkinson's disease. Eur J Neurosci. 2019;49(3):339-63.

47. Zhao Y, Dzamko N. Recent Developments in LRRK2-Targeted Therapy for Parkinson's Disease. Drugs. 2019;79(10):1037-51.

48. Jennings D, Huntwork-Rodriguez S, Henry AG, Sasaki JC, Meisner R, Diaz D, et al. Preclinical and clinical evaluation of the LRRK2 inhibitor DNL201 for Parkinson's disease. Sci Transl Med. 2022;14(648):eabj2658.

49. Davies H, Bignell GR, Cox C, Stephens P, Edkins S, Clegg S, et al. Mutations of the BRAF gene in human cancer. Nature. 2002;417(6892):949-54.

50. Purvis GSD, Solito E, Thiemermann C. Annexin-A1: Therapeutic Potential in Microvascular Disease. Front Immunol. 2019;10:938.

51. Gross AM, Wolters PL, Dombi E, Baldwin A, Whitcomb P, Fisher MJ, et al. Selumetinib in Children with Inoperable Plexiform Neurofibromas. N Engl J Med. 2020;382(15):1430-42.

52. Ciombor KK, Bekaii-Saab T. Selumetinib for the treatment of cancer. Expert Opin Investig Drugs. 2015;24(1):111-23.

53. Perez DM. alpha1-Adrenergic Receptors in Neurotransmission, Synaptic Plasticity, and Cognition. Front Pharmacol. 2020;11:581098.

54. Davies AO, Lefkowitz RJ. Regulation of beta-adrenergic receptors by steroid hormones. Annu Rev Physiol. 1984;46:119-30.

55. Aksoy MO, Mardini IA, Yang Y, Bin W, Zhou S, Kelsen SG. Glucocorticoid effects on the beta-adrenergic receptor-adenylyl cyclase system of human airway epithelium. J Allergy Clin Immunol. 2002;109(3):491-7.

56. Inchiosa MA, Jr. Anti-tumor activity of phenoxybenzamine and its inhibition of histone deacetylases. PLoS One. 2018;13(6):e0198514.

57. Inchiosa MA, Jr. Further investigation of the potential anti-neoplastic, anti-inflammatory and immunomodulatory actions of phenoxybenzamine using the Broad Institute CLUE platform [Internet]. The Preprint Server for Biology; Cold Spring Harbor Laboratory 2019. doi.org/10.1101/767392.

58. Charo IF, Ransohoff RM. The many roles of chemokines and chemokine receptors in inflammation. N Engl J Med. 2006;354(6):610-21.

59. Doran E, Cai F, Holweg CTJ, Wong K, Brumm J, Arron JR. Interleukin-13 in Asthma and Other Eosinophilic Disorders. Front Med (Lausanne). 2017;4:139.

60. Teixeira AL, Gama CS, Rocha NP, Teixeira MM. Revisiting the Role of Eotaxin-1/CCL11 in Psychiatric Disorders. Front Psychiatry. 2018;9:241.

61. Chandra G, Rangasamy SB, Roy A, Kordower JH, Pahan K. Neutralization of RANTES and Eotaxin Prevents the Loss of Dopaminergic Neurons in a Mouse Model of Parkinson Disease. J Biol Chem. 2016;291(29):15267-81.

62. Bhatt SP, Rabe KF, Hanania NA, Vogelmeier CF, Cole J, Bafadhel M, et al. Dupilumab for COPD with Type 2 Inflammation Indicated by Eosinophil Counts. N Engl J Med. 2023.

63. Goel HL, Mercurio AM. VEGF targets the tumour cell. Nat Rev Cancer. 2013;13(12):871-82.

64. Xu Z, Goel HL, Burkart C, Burman L, Chong YE, Barber AG, et al. Inhibition of VEGF binding to neuropilin-2 enhances chemosensitivity and inhibits metastasis in triple-negative breast cancer. Sci Transl Med. 2023;15(694):eadf1128.

65. Mabeta P, Steenkamp V. The VEGF/VEGFR Axis Revisited: Implications for Cancer Therapy. Int J Mol Sci. 2022;23(24).

66. Schacke H, Docke WD, Asadullah K. Mechanisms involved in the side effects of glucocorticoids. Pharmacol Ther. 2002;96(1):23-43.

67. Isaacs C, Wellstein A, Riegel AT. Hormones and Related Agents in the Therapy of Cancer. In: Brunton LL, editor. Goodman & Gilman's The Pharmacological Basis of Therapeutics. New York: McGraw Hill; 2018. p. 1237-47.

68. Lin KT, Wang LH. New dimension of glucocorticoids in cancer treatment. Steroids. 2016;111:84-8.

69. Greenstein S, Ghias K, Krett NL, Rosen ST. Mechanisms of glucocorticoid-mediated apoptosis in hematological malignancies. Clinical cancer research : an official journal of the American Association for Cancer Research. 2002;8(6):1681-94.

70. Schacke H, Rehwinkel H. Dissociated glucocorticoid receptor ligands. Curr Opin Investig Drugs. 2004;5(5):524-8.

71. Schacke H, Berger M, Rehwinkel H, Asadullah K. Selective glucocorticoid receptor agonists (SEGRAs): novel ligands with an improved therapeutic index. Mol Cell Endocrinol. 2007;275(1-2):109-17.

72. Keenan CR, Lew MJ, Stewart AG. Biased signaling from the glucocorticoid receptor: Renewed opportunity for tailoring glucocorticoid activity. Biochem Pharmacol. 2016;112:6-12.

73. Kleiman A, Tuckermann JP. Glucocorticoid receptor action in beneficial and side effects of steroid therapy: lessons from conditional knockout mice. Mol Cell Endocrinol. 2007;275(1-2):98-108.

74. De Bosscher K, Haegeman G, Elewaut D. Targeting inflammation using selective glucocorticoid receptor modulators. Curr Opin Pharmacol. 2010;10(4):497-504.

75. Nicolaides NC, Chrousos G, Kino T. Glucocorticoid Receptor. In: Feingold KR, Anawalt B, Blackman MR, Boyce A, Chrousos G, Corpas E, et al., editors. Endotext. South Dartmouth (MA)2000.

76. Lesovaya EA, Chudakova D, Baida G, Zhidkova EM, Kirsanov KI, Yakubovskaya MG, et al. The long winding road to the safer glucocorticoid receptor (GR) targeting therapies. Oncotarget. 2022;13:408-24.

77. Haming D, Simoes-Costa M, Uy B, Valencia J, Sauka-Spengler T, Bronner-Fraser M. Expression of sympathetic nervous system genes in Lamprey suggests their recruitment for specification of a new vertebrate feature. PLoS One. 2011;6(10):e26543.

78. Davies AO, Lefkowitz RJ. Agonist-promoted high affinity state of the beta-adrenergic receptor in human neutrophils: modulation by corticosteroids. J Clin Endocrinol Metab. 1981;53(4):703-8.

79. Lefkowitz RJ. Clinical physiology of adrenergic receptor regulation. Am J Physiol. 1982;243(1):E43-7.

80. Logsdon PJ, Middleton E, Jr., Coffey RG. Stimulation of leukocyte adenyl cyclase by hydrocortisone and isoproterenol in asthmatic and no asthmatic subjects. J Allergy Clin Immunol. 1972;50(1):45-56.

81. Marone G, Lichtenstein LM, Plaut M. Hydrocortisone and human lymphocytes: increases in cyclic adenosine 3':5'-monophosphate and potentiation of adenylate cyclase-activating agents. J Pharmacol Exp Ther. 1980;215(2):469-78.

82. Parker CW, Huber MG, Baumann ML. Alterations in cyclic AMP metabolism in human bronchial asthma. 3. Leukocyte and lymphocyte responses to steroids. J Clin Invest. 1973;52(6):1342-8.

83. Adams M, Meijer OC, Wang J, Bhargava A, Pearce D. Homodimerization of the glucocorticoid receptor is not essential for response element binding: activation of the phenylethanolamine N-methyltransferase gene by dimerization-defective mutants. Mol Endocrinol. 2003;17(12):2583-92.

84. Tai TC, Claycomb R, Her S, Bloom AK, Wong DL. Glucocorticoid responsiveness of the rat phenylethanolamine N-methyltransferase gene. Mol Pharmacol. 2002;61(6):1385-92.

85. Flydal MI, Krakenes T-A, Tai MDS, Tran MPA, Teigen K, Martinez A. Levalbuterol lowers the feedback inhibition by dopamine and delays misfolding and aggregation in tyrosine hydroxylase. Biochimie. 2021;183:126-132.

86. Milgrom H, Skoner DP, Bensch G, Kim KT, Claus R, Baumgartner RA. Low-dose levalbuterol in children with asthma: Safety and efficacy in comparison with placebo and racemic albuterol. J Allergy Clin Immunol. 2001;108:938-45.

87. Sriprasart T, Waterer G, Garcia G, Rubin A, Andrade MAL, Roguska A, et al. Safety of SABA monotherapy in asthma management: a systematic review and meta-analysis. Adv Ther. 2023;40:133-158.

Figure 1. HeatMap of association strengths (median tau score) between gene expression signatures of epinephrine in 8 malignant cell cultures and those for all entries in the Touchstone database, with targets. Chemical compound (CP); Gene knock-down (KD); Gene over-expression (OE); and perturbagen class (PCL).

type	median_tau_score	name	description	target
	99.99	norepinephrine	adrenergic receptor agonist, DNA inhibitor, free radical scavenger	ADRA1A, ADRA1B, ADRA1D, ADRA2A, ADRA2B, ADRA2C, ADRB1,
	99.87	terbutaline	adrenergic receptor agonist	ADRB2
	98.46	tetracycline	30S ribosomal protein inhibitor, 30S ribosomal subunit inhibitor, protein argi	PRNP
	97.99	dopamine	dopamine receptor agonist	DRD2, DRD1, DRD3, DRD4, DRD5, DBH, HTR1A, HTR7, SLC6A2, SL
	97.75	forskolin	adenylyl cyclase activator, Adenylate cyclase stimulant, growth hormone re	ADCY2, ADCY5, GNAS
	97.21	orciprenaline	adrenergic receptor agonist	ADRB2
	96.84	phylloquinone	antifibrinolytic agent, gamma carboxylase enzyme	BGLAP, GGCX
	96.43	acebutolol	adrenergic receptor antagonist	ADRB1, ADRB2
	96.23	dobutamine	adrenergic receptor agonist	ADRB1, ADRB2, ADRA1A
	96.10	PROKR1	GPCR / Class A : Prokineticin receptors, prokineticin receptor 1	
	95.69	fenoterol	adrenergic receptor agonist	ADRB2, ADRB1, ADRB3
	95.52	mupirocin	isoleucyl-tRNA synthetase inhibitor	
	95.49	Beta-Adrenergic Rece		
	95.49	PSMB9	Proteasome (prosome, macropain) subunits, proteasome (prosome, macro	
	95.47	fluocinonide	corticosteroid agonist, corticosteroid hormone receptor agonist	NR3C1, SERPINA6, SMO
	95.24	VU-0400193-3	glutamate receptor modulator	GRM4
	95.07	procaterol	adrenergic receptor agonist	ADRB2
	94.56	nevirapine	non-nucleoside reverse transcriptase inhibitor, reverse transcriptase inhibit:	CYP2B6, CYP2C19, CYP3A4
	94.54	17-beta-estradiol	estrogen receptor agonist	ESR1
	94.37	selegiline	monoamine oxidase inhibitor, superoxide dismutase stimulant	MAOB, MAOA, CYP2B6, CYP2C19
	94.22	ARID5B	AT rich interactive domain 5B (MRF1-like)	
	94.21	danazol	estrogen receptor antagonist, luteinizing hormone releasing hormone antag	ESR1, GNRHR, AR, CCL2, CYP2C8, GNRHR2, PGR, TNF
	94.10	pyrazinamide	fatty acid synthase inhibitor	FASN
	94.10	EIF5A	eukaryotic translation initiation factor 5A	
	94.08	CNNM2	cyclin M2	
	93.98	mirtazapine	adrenergic receptor antagonist, serotonin receptor antagonist	ADRA2A, HTR2A, HTR2C, ADRA2C, HTR3A, ADRA1A, ADRA1B, ADI
	93.81	labetalol	adrenergic receptor antagonist	ADRB1, ADRB2, ADRA1A, ADRA1D, ADRA1B
	93.63	TMEM127	transmembrane protein 127	
	93.27	POLR1E	RNA polymerase subunits, polymerase (RNA) I polypeptide E, 53kDa	
	93.17	R-96544	serotonin receptor antagonist	HTR2A
	93.03	acadesine	adenosine release stimulant, AMPK activator, AMPK stimulant	
	92.76	GR-79236	adenosine receptor agonist	ADORA1, ADORA2A
	92.70	kynuramine	aryl hydrocarbon receptor activator	
	92.68	cimaterol	adrenergic receptor agonist	
	92.48	tetrahydrobiopterin	nitric oxide stimulant, nitric oxide synthase stimulant, phenylalanine 4-hydr(	PAH, TH, DBH, NOS3, TPH1
	92.43	BRD-K14844937	GABA benzodiazepine site receptor inverse agonist	GABRA1, GABRA2, GABRA3, GABRA5
	92.41	etilefrine	adrenergic receptor agonist	ADRA1A, ADRB1
	92.41	PAQR7	progestin and adipoQ receptor family member VII	

Figure 2. HeatMap of association strengths (median tau score) between gene expression signatures of norepinephrine in 8 malignant cell cultures and those for all entries in the Touchstone database. Chemical/pharmacologic agent (CP); Gene knock-down (KD); Gene over-expression (OE); and perturbagen class (PCL).

Fig 3. Comparison of epinephrine and norepinephrine gene-expression scores as glucocorticoid receptor agonists.
The horizontal lines are the median values for the individual distributions. Mann-Whitney analysis of the distributions showed a statistically significant difference (p <0.0001), with strong association for epinephrine with gene-expression similarity to glucocorticoid agonists.

Fig 4. Comparison of epinephrine and hydrocortisone gene-expression scores as glucocorticoid receptor agonists.
The horizontal lines are the median values for the individual distributions, which are essentially identical. Mann-Whitney analysis of the distributions showed no difference in gene-expression characteristics for epinephrine and hydrocortisone as glucocorticoid receptor agonists (p = 0.805).

Fig 5. Comparison of beta2-adrenergic related agents and hydrocortisone for gene expression connectivity scores as glucocorticoid receptor agonists.
The horizontal lines are the median values for the individual distributions. The relative differences among the gene-expression scores can be appreciated visually. Caffeine was not studied further.

Fig 6. Percent body weight loss calculated from each animal's initial body weight at Study Day 0.
Body weight loss was statistically significantly less for the group that was pretreated with levalbuterol and challenged with the monoclonal antibody, OKT3.

Fig 7. CRS clinical scores for the period following toxic induction of cytokine release with OKT3 or aCD28.

There were no statistically significant differences in the clinical scores that were attributable to the pretreatments; the doses of pretreatments were calculated from typical human doses that were translated using FDA factors.

Fig 8. Water consumption recorded per cage of 5 mice as a percentage of the intake on Study Day 6 prior to the induction challenges with OKT3 and anti-CD28.

The toxic antibody challenges to induce cytokine release caused marked decreases in water consumption in all of the groups. Since the pretreatment drugs were administered in the drinking water, it would be expected that drug delivery was compromised.

Fig 9. The effects of β2-adrenergic related drugs on the release of eotaxin and VEGF.

Elaboration of Eotaxin (A), VEGF (B), and IL-13 (C), in the CRS assay. D, comparisons are illustrated in the same order for the gene-expression connectivity of the agents as glucocorticoid receptor agonists in the CLUE database. As noted, the gene-expression connectivity for epinephrine, the prototypical beta2-adrenergic agonist, is substituted for levalbuterol and arfomoterol since these drugs were never profiled in the CLUE database. Albuterol, the racemic mixture of R and S levalbuterol, showed no inhibition of cytokine/chemokine release in any of the assays (A, B, C).

Fig 10. "Power" analysis of the results for the VEGF and eotaxin assays.

Duplication of data from 5-animal CRS assays but variance unchanged.