# Genome and tissue-specific transcriptome of the tropical milkweed (*Asclepias curassavica*)

Honglin Feng[1,5], Jing Zhang[1], Adrian F. Powell[1], Gretta L. Buttelmann[4], Lily Yang[1], Ethan Yan[1], Fumin Wang[1], Steven B. Broyles[2], Georg Jander[1*], and Susan R. Strickler[3,4*]

[1]Boyce Thompson Institute, Ithaca, NY 14853, USA
[2]Biological Sciences, State University of New York, Cortland, NY 13045, USA
[3]Negaunee Institute for Plant Conservation Science and Action, Chicago Botanic Garden, Glencoe, IL 60022, USA
[4]Plant Biology and Conservation Program, Northwestern University, Evanston, IL 60208, USA
[5]Present address: Department of Entomology, Louisiana State University, Baton Rouge, LA 70803, USA

*To whom correspondence should be addressed: sstrickler@chicagobotanic.org, gj32@cornell.edu

## Abstract

Tropical milkweed (*Asclepias curassavica*) serves as a host plant for monarch butterflies (*Danaus plexippus*) and other insect herbivores that can tolerate the abundant cardiac glycosides that are characteristic of this species. Cardiac glycosides, along with additional specialized metabolites, also contribute to the ethnobotanical uses of *A. curassavica*. To facilitate further research on milkweed metabolism, we assembled the 241 Mbp genome of a fifth-generation inbred line of *A. curassavica* into 851 contigs, with an N50 of 2.4 Mbp. Scaffolding resulted in 97.9% of the assembly being anchored to 11 chromosomes, which are colinear with the previously assembled common milkweed (*A. syriaca*) genome. Assembly completeness evaluations showed that 97.9% of the BUSCO gene set is present in the *A. curassavica* assembly. The transcriptomes of six tissue types (young leaves, mature leaves, stems, flowers, buds, and roots), with and without methyl jasmonate treatment, showed both tissue-specific gene expression and induced expression of genes that may be involved in cardiac glycoside biosynthesis. Together, this genome sequence and transcriptome analysis provide important resources for further investigation of the ecological and medicinal uses of *A. curassavica*.

**Introduction**

Plants in the genus *Asclepias* (milkweeds) are known for their production of cardiac glycosides (Figure 1A), defensive compounds that are inhibitors of essential $Na^+/K^+$-ATPases in animal cells. Specialized herbivores from at least six insect orders have developed target-site resistance to the cardiac glycosides in their *Asclepias* host plants (Dobler et al., 2012; Petschenka et al., 2017; Karageorgi et al., 2019). In particular, the iconic interactions between milkweeds and *Danaus plexippus* (monarch butterflies), which sequester toxic cardiac glycosides from their milkweed host plants for their own protection, have been the subject of extensive ecological research and hundreds of publications. Among the more than 200 known milkweed species, *Asclepias syriaca* (common milkweed) and *Asclepias curassavica* (tropical milkweed; Figure 1B) are the most common host plants for monarch caterpillars (Agrawal, 2017).

Whereas *A. syriaca* is found in temperate areas of North America, the native range of *A. curassavica* (Figure 1B) comprises tropical North and South America, including the Caribbean islands (Woodson, 1954). A shrubby perennial, *A. curassavica* has been introduced as a garden ornamental or a naturalized weed in other tropical areas of the world. In areas where *A. syriaca* does not grow, *A. curassavica* is the main host plant for monarch butterflies. Caterpillars from non-migratory monarch populations in southern Florida were reported to feed primarily on introduced *A. curassavica* (Knight and Brower, 2009).

Due to its ethnobotanical uses, Carl Linnaeus named the genus *Asclepias* after Asclepius, the Greek god of medicine (Linnaeus, 1754; Quattrocchi, 1999). The numerous reported medical applications of *A. curassavica* include the antimicrobial (Navarro Garcia et al., 2003; Peraza-Sanchez et al., 2007; Jacobo-Salcedo Mdel et al., 2011; Alonso-Castro et al., 2021) and anticancer (Yuan et al., 2016; Zheng et al., 2019) properties of plant extracts. More broadly, cardiac glycosides produced by milkweed and other plants, in particular foxglove (*Digitalis* spp.), have been used for centuries to treat congestive heart failure (Tomov, 1981; Rehman and Hai, 2023). The first enzyme of the cardiac glycoside biosynthesis pathway, which catalyzes sterol side chain cleavage, has been confirmed in species from three plant families: Plantaginaceae, woolly foxglove (*Digitalis lanata*) and common foxglove (*D. purpurea*); Apocynaceae, Sodom apple (*Calotropis procera*); and Brassicaceae, wormseed wallflower (*Erysimum cheiranthoides*) (Carroll et al., 2023; Kunert et al., 2023; Younkin et al., 2024).

Although *A. syriaca* genome assemblies have been described previously (Straub et al., 2011; Weitemier et al., 2019; Boyle et al., 2022), this species is not very tractable as a laboratory model system. In contrast to *A. syriaca*, *A. curassavica* grows well in pots in the greenhouse, does not require a winter diapause, can be propagated from cuttings, and is self-fertile (Wyatt and Broyles, 1997). To facilitate molecular genetic research on the extensive ecological interactions and medical applications of *Asclepias*, we initiated an *A. curassavica* genome sequencing project. Here we present the assembly and annotation of a high-quality genome, as well as the transcriptomes of six tissue types, from a fifth-generation inbred lineage of *A. curassavica*.

## Methods

### *Plants and growth conditions*

*Asclepias curassavica* seeds were initially purchased from Butterfly Encounters (www.butterflyencounters.com/). Plants were grown in a greenhouse under ambient light and were self-pollinated manually for five generations. To produce a larger number of seeds in the sixth generation, a flowering plant was placed in a cage with a bumblebee colony (*Bombus terrestris*; https://www.biobestgroup.com) for more efficient pollination. Seeds from the sixth generation of inbreeding were submitted to the USDA-ARS Ornamental Plant Germplasm Center (https://opgc.osu.edu/ ; https://www.ars-grin.gov/ ; accession number OPGC 7586). The fifth-generation inbred line of *A. curassavica* was used to prepare material for genomic DNA and cDNA sequencing.

For plant growth, seeds were washed three times with 0.6% sodium hypochlorite in deionized water with one drop of dish soap. After the washes, seeds were transferred to a 90 mm Petri dish containing a layer of moist brown paper towel. Sealed Petri dishes with seeds were incubated at 30°C for 3-6 days, until germination. After the seeds had germinated, seedlings were then moved to pots containing Cornell Mix (by weight 56% peat moss, 35% vermiculite, 4% lime, 4% Osmocote slow-release fertilizer [Scotts, Marysville, OH], and 1% Unimix [Scotts]) in Conviron (Winnipeg, Canada) growth chambers with a 16:8 photoperiod, 180 µM m$^{-2}$ s$^{-1}$ photosynthetic photon flux density, 60% humidity, and constant 23 °C temperature.

### *Methyl jasmonate treatment*

Synchronized flowering *A. curassavica* plants were treated with methyl jasmonic acid (MeJA) before tissue collection. Each plant was sprayed with 0.01% MeJA (dissolved in 0.1% ethanol) to cover the whole plant, both sides of the leaves and the stem, to the point of liquid run-off, in a separated spray room. Control plants were sprayed with 0.1% ethanol. After the spray treatment, MeJA-treated and control-treated plants were separated in two different greenhouses with the same temperature and light settings. After 24 hours, all plants were sprayed again. Six hours after the booster spray, different plant tissues, including young leaves (the top 4 newly emerged leaves), mature leaves, stems, flowers, buds, and roots, were collected, flash-frozen in liquid nitrogen, and stored at -80°C until use. The six tissue types were collected in eight-fold replication, with and without MeJA treatment, for a total of 96 samples.

## *DNA extraction, library preparation, and sequencing*

To extract intact DNA for long-read sequencing, *A. curassavica* leaf tissue was harvested and placed into 2 ml screw-cap vials, 3 three-mm steel balls were added to each tube, tubes were flash-frozen in liquid nitrogen, and samples were homogenized on a paint shaker (Harbil, Wheeling, IL, USA). DNA was extracted from the homogenized tissue using a modified cetyl trimethylammonium bromide (CTAB) protocol using chloroform:isoamyl alcohol 24:1 (Fulton et al., 1995). To ensure that DNA fragments were of the correct size for Nanopore sequencing, the DNA was then size-selected using the Circulomics SRE kit v2.0 (SS-100-101-01, PacBio, Menlo Park, CA, USA). A concentration of 150 ng/µL of DNA was initially used, and 55 µL of Buffer EB was added prior to the incubation step. To prepare the library, the Oxford Nanopore Genomic DNA by Ligation kit (SQK-LSK109, Oxford Nanopore Technologies, New York, NY, USA) was used with the undiluted ~2.5 mg of SRE-treated DNA from the previous step. Sequencing was done using a Nanopore MinION flow cell (flow cell number FAL75881-1282) on a MinION sequencing device. To maximize the available runtime of the flow cell and generate more data, the library was split in two and each half was sequenced successively on the same flow cell, one before and one after completing the Nanopore Nuclease flush protocol (Oxford Nanopore Technologies, New York, NY, USA).

For Illumina sequencing, DNA was extracted from *A. curassavica* leaves using the Wizard® Genomic DNA Purification Kit (Promega, Madison WI, USA). The quantity and quality of genomic DNA was assessed using a Qubit 3 fluorometer (Thermo Fisher, Waltham, MA, USA) and a Bioanalyzer DNA12000 kit (Agilent, Santa Clara, CA, USA). The PCR-free

TruSeq DNA (Illumina, San Diego, CA) was used to prepare a library. Samples were sequenced on an Illumina MiSeq instrument (paired-end 2×150 bp) at the Cornell University Biotechnology Resource Center (Ithaca, NY, USA). A Hi-C library was prepared from *A. curassavica* leaves using the Proximo Hi-C Plant Kit (Phase Genomics, Seattle, WA, USA) and was sequenced on an Illumina MiSeq instrument (paired-end 2×75bp) at the Cornell University Biotechnology Resource Center (Ithaca, NY, USA).

### *RNA extraction, cDNA library preparation, and sequencing*

RNA samples were prepared using the RNeasy® Plant Mini Kit (Qiagen, Hilden, Germany) according to the manufacturer's protocol. Briefly, tissue samples were ground using a 1600 MiniG® Automated Tissue Homogenizer and Cell Lyser (Cole-Parmer, Vernon, IL, USA) and ~100 mg of each sample were used for RNA extraction. RNA concentrations were measured using a Qubit® RNA BR kit (ThermoFisher Scientific, Waltham, MA, USA) on a Qubit® 3.0 Fluorometer (ThermoFisher Scientific), and 10 samples were randomly selected and evaluated using the Agilent 2100 bioanalyzer (Agilent, Santa Clara, CA, USA) to ensure high RNA quality (RNA Integrity Number > 7).

High-quality RNA samples were used for cDNA library preparation with the Illumina Stranded mRNA Prep kit following a ligation protocol (Illumina, San Diego, CA, USA). Briefly, from the total RNAs (~ 1 ug), we first captured and enriched mRNA with polyA tails using oligo (dT) magnetic beads. Purified mRNA samples were then fragmented and denatured for first-strand cDNA synthesis using the first-strand synthesis master mix with reverse transcriptase. Then, the mRNA templates were removed and second-strand cDNA was synthesized to generate blunt-ended double-stranded cDNA fragments. In place of deoxythymidine triphosphate (dTTP), deoxyuridine triphosphate, (dUTP) was incorporated to quench the second strand during amplification to achieve strand specificity. The cDNA fragments were captured and purified using the Agencourt AMPure XP magnetic beads (Beckman Coulter, Brea, CA, USA). The blunt-ended double-stranded cDNA was adenylated at the 3' end to prevent self-ligation and facilitate ligation to the adapters with corresponding thymine (T) nucleotides on the 3' end in a later step. Adenylated cDNA fragments were further ligated with pre-index RNA anchors for dual indexing. To amplify cDNA libraries, the anchor-ligated DNA fragments were selectively amplified and dual-indexed using unique indexes and primer sequences with 10 cycles of PCR. Finally, the cDNA libraries were cleaned up using AMPure XP magnetic beads and

concentrations were measured using the Qubit® dsDNA BR Assay kit (ThermoFisher Scientific) using a Qubit® 3.0 Fluorometer.

RNAseq data were generated using Illumina sequencing at the Cornell University Biotechnology Resource Center (Ithaca, NY). The 96 uniquely dual-indexed cDNA libraries were pooled based on their concentrations measured using a Qubit® 3.0 Fluorometer. The pooled libraries were first evaluated using the Agilent 2100 Bioanalyzer for quality check and then sequenced with the Illumina MiSeq platform (Illumina) to ensure approximately equal amounts input of each individual library for Illumina HiSeq. Finally, about 5 million 2x75 bp pair-ended RNAseq reads were generated for each of the 96 libraries using the Illumina NextSeq500 platform (Illumina).

### *Genome assembly and annotation*

Genome size was estimated by k-mer analysis of Illumina reads using Jellyfish v2.2.7 (Marçais and Kingsford, 2011) and GenomeScope.v1.0 (Vurture et al., 2017). Genomic *A. curassavica* nanopore and Illumina sequences were assembled using MaSurCA v3.4.1 (Zimin et al., 2013). The assembly was corrected using Illumina sequences and Pilon v1.23 (Walker et al., 2014). Redundancy was removed using Purge Hapolotigs (Roach et al., 2018). Hi-C was used to scaffold the contigs using 3D-DNA v180922 (Dudchenko et al., 2017). The assembly was manually adjusted in Juicebox v2.15.07 (Durand et al., 2016). Assembly gaps were filled using the nanopore reads and draft assembly as input to LR_gapcloser v3 (Xu et al., 2019).

LTR_retriever (Ou and Jiang, 2018) was used with outputs from LTRharvest (Ellinghaus et al., 2008) and LTR_FINDER (Xu and Wang, 2007) to identify long terminal repeat retrotransposons (LTRs). The LTR library was then used to hard mask the genome, and RepeatModeler version: open-1.0.11 (Smit and Hubley, 2015) was used to identify additional repetitive elements in the remaining unmasked segments of the genome. Protein-coding sequences were excluded using blastx v2.8.1+ (Altschul et al., 1990; Ellinghaus et al., 2008) results in conjunction with the ProtExcluder.pl script from the ProtExcluder v1.2 package (Campbell et al., 2014). The libraries from RepeatModeler and LTR_retriever were then combined and used with RepeatMasker v4.0.9 (Smit et al., 2015) to produce the final masked version of the genome.

RNAseq reads were mapped to the genome with HISAT2 v2.1.0 (Kim et al., 2015). Portcullis v1.2.0 (Mapleson et al., 2018) and Mikado v2.3.2 (Venturini et al., 2018) were used to

process and filter the resulting bam files. Augustus v3.3.3 (Stanke et al., 2008) and Snap v2006-07-28 (Korf, 2004) were trained and implemented through the Maker v2.31.10 pipeline (Cantarel et al., 2008), with proteins from Swiss-Prot (Boutet et al., 2007) and processed RNAseq added as evidence. Gene models were filtered with the following criteria: 1) at least one match found in the Trembl database (4-17-19) (Boutet et al., 2007) with an E-value less than 1e-20, 2) InterProScan matches to repeats were removed, 3) genes with an AED score of 1 and no InterPro domain were removed, and 4) single-exon genes with no InterPro domain were removed. Functional annotation and classification were performed using BLASTx v2.8.1+ (Altschul et al., 1990) and InterProScan v5.36-75.0 (Jones et al., 2014). Both genome and annotation completeness were assessed by BUSCO v3.1.0 (Waterhouse et al., 2017) using the embryophyta lineage set.

### *Synteny, gene family, and phylogenetic analysis*

A synteny plot between the *A. curassavica* and *A. syriaca* pseudomolecules was created using sequence matches calculated with nucmer v4.0.0beta2 (Kurtz et al., 2004) and plotted using Circos v0.69-6 (Krzywinski et al., 2009). OrthoFinder v2.5.2 (Emms & Kelly, 2015) was used to identify gene families using protein sequences from genome-sequenced close relatives to *A. curassavica* with genome annotations. The following species were included: *A. curassavica* v1.0, *Asclepias syriaca* v1.0 (Boyle et al., 2022), *Calotropis gigantea* (Hoopes et al., 2018), *Catharanthus roseus* (Kellner et al., 2015), *Rhazya stricta* (Sabir et al., 2016), *Eustoma grandiflorum* (Liang et al., 2022), *Gelsemium sempervirens* (Frank et al., 2015), *Neolamarckia cadamba* (Zhao et al., 2022), *Chiococca alba* (Lau et al., 2020), *Coffea canephora* v1.0 (Denoeud et al., 2014). These results were used to create an Upset plot using the UpSetR package v1.4.0 (Conway et al., 2017) depicting shared and unique gene clusters. Maximum likelihood methods were used to identify significantly expanded and contracted gene families in *Asclepias*. Gene families from Orthofinder and the species tree were input into CAFÉ v.3.0 (Han et al., 2013). Before inputting, the tree was ultrametricized with penalized likelihood using the chronopl() function in the R package APE (Paradis et al., 2004). An error model and a single lambda value was estimated for the tree. For the gene families showing significant expansion and contraction ($p<0.05$) in *A curassavica*, we retrieved the list of associated gene ontology (GO) terms from our annotation. The resulting list of expanded or contracted *A. curassavica* orthogroups and their associated GO terms was input to topGO (Alexa and Rahnenfuhrer, 2023)

for enrichment analyses. We searched for enrichment in GO terms associated with biological functions, molecular process, and cellular component and used Fisher's exact test to determine significance.

For phylogenetic analysis of the CPY87A4 gene family, the *Digitalis lanata* CPY87A4 protein sequence (Accession ID OR134564) was retrieved from Genbank. The gene family that best matched this protein based on sequence identity was identified by BLAST queries and the alignment was retrieved from the OrthoFinder multiple sequence alignment (MSA) results. A gene tree was constructed in iQ-Tree2 2.2.2.6 (Minh et al., 2020) using the MSA and the following parameters `--alrt 1000 -B 1000`, the JTT+I+R3 model, and 1000 bootstrap replicates. The tree was visualized with TreeViewer v2.2.0 .(Bianchini and Sánchez-Baracaldo, 2023).

### *Differential expression analysis*

The paired-end sequencing reads from cDNA libraries were aligned to gene models from the reference *A. curassavica* genome using bowtie v2.2.5 with default settings (Langmead and Salzberg, 2012). The raw read counts for each gene model were extracted using the pileup.sh script from BBMap v35.85 (https://sourceforge.net/projects/bbmap/). Raw read counts were then used to generate a data matrix for differential expression analyses using the Bioconductor package edgeR v3.36.0 (Robinson et al., 2010) in Rstudio v2022.02.3 (https://www.rstudio.com). For differential expression analyses, the raw read counts for each gene were first normalized as counts per million (CPM) to the total read counts of all genes in each sample. Only genes with > 2 CPMs of expression and detected in at least 8 samples were retained for further analyses. The filtered normalized reads were then investigated using principal component analysis (PCA) to exclude the possibility of outliers or batch effects. Then, differential expression analyses were performed between the MeJA and control treatments, independently within each tissue type. To identify differentially expressed genes (MeJA vs control), we applied the exactTest function followed by the decideTestsDGE function in edgeR, where we retained genes with a significant differential expression value of $p \leq 0.05$, and a Benjamini & Hochberg adjusted false discovery rate (FDR) $\leq 0.05$ (Robinson et al., 2010).

To facilitate visualization of variations in gene expression, the common differentially expressed genes across different tissue types were identified using a web-based Venn diagram generator from Bioinformatics & Evolutionary Genomics (https://bioinformatics.psb.ugent.be)

and the result was plotted with UpSetR v1.4.0 (Conway et al., 2017). To understand the function of those genes, in addition to using the existing genome annotation, we further annotated the genes by blasting their protein sequences against Uniprot database, using PANNZER2 (Törönen et al., 2018) and local BLAST (Altschul et al., 1997; Schäffer et al., 2001), as well as KEGG pathway annotation (https://www.genome.jp/kegg/pathway.html). To visualize the expression results, we generated a heatmap for all the log2 fold changes of the common differentially expressed genes using the R package pheatmap v1.0.12 (Kolde, 2018).

## Results

### A. curassavica *genome assembly and annotation*

We sequenced the *A. curassavica* genome, which was estimated to be 639 Mbp by flow cytometry (Lewis and Ruter, 2023), using paired-end Illumina and Oxford Nanopore sequencing, with 93X and 45X coverage, respectively. K-mer analysis suggested that the genome size is only 161 Mbp, significantly smaller than the previous estimates, with a heterozygosity 0.2% (Figure S1). The Illumina and nanopore sequencing reads were assembled into 851 contigs with an N50 of 2.4 Mbp and a total length of 241.1 Mbp. Scaffolding resulted in 97.9% of the assembly being anchored to 11 chromosomes. Assembly completeness evaluations using BUSCO showed that 97.9% of the BUSCO set is present in the assembly, and only 1.1% of the genes were duplicated (Table S1).

Repeat analysis indicates that 51.5% of the genome is composed of repetitive elements, with LTR elements being the most abundant (Figure 2). Gene prediction resulted in 26,035 gene models, which tended to be located toward distal regions of the chromosomes. The annotation captures 95.2% of the BUSCO set (Table S1). Altogether, 21,221 genes were functionally annotated and 13,196 were assigned GO terms.

### *Evolution of the* A. curassavica *genome*

Although smaller in total length, the 11 pseudomolecules of the *A. curassavica* genome assembly are highly colinear with the *A. syriaca* genome assembly (Boyle et al., 2022) (Figure 3). The *A. syriaca* genome was found to have a greater expansion of LTR retroelements relative to *A. curassavica*, contributing to its larger size (Figure 2). Gene family analysis involving closely related species with sequenced genomes, *A. curassavica, A. syriaca*, *C. gigantea*, *C. roseus*, *C.*

*alba*, *C. canephora, E. grandiflorum*, *G. sempervirens*, *M. speciosa, N. cadamba*, and *R. stricta*, assigned 92.5% of detected genes (336,215/ 363,631) to 26,866 orthogroups. A total of 9,052 gene families containing 168,416 genes were shared between all species (Figure 4). To identify gene families that had expanded or contracted we first estimated a lambda rate with CAFE5 of 0.638. A total of 406 gene families have expanded in *A. curassavica* and 564 have contracted (Figure 5). We then focused on rapidly evolving gene families that had changed significantly along the *A. curassavica* lineage. Rapidly expanding gene families in *A. curassavica* include genes involved in amine metabolic processes and defense responses, whereas rapidly contracting families included genes involved in magnesium ion transport and manganese ion homeostasis (Table S2).

## *Differentially expressed genes upon jasmonic acid treatment*

We collected samples from six tissue types (buds, roots, stems, flowers, young leaves, and mature leaves), with and without MeJA treatment, in 8-fold replication for analysis of gene expression. Illumina sequencing (RNAseq) generated approximately 5 million paired-end reads for each of the samples. Approximately 75% of RNAseq reads of each sample were mapped to the coding sequences of gene models annotated in the genome. Principal component analysis (PCA) showed that the overall gene expression patterns from both control samples (Figure 6A) and MeJA-treated samples (Figure 6B) were distinct in each tissue type. Similarly, PCA clustering based on tissue types demonstrated clear separation between control and MeJA-treated samples (Figure 6C-H). It is worth noting that, compared to other tissue types, samples from buds showed less distinct separation (Figure 6C), possibly due to their rapid growth and dynamic changes in gene expression in this tissue type.

Differential gene expression analyses were conducted independently for each tissue type comparing MeJA-treated samples to controls. In buds, roots, stems, flowers, young leaves, and mature leaves, we identified 2465, 4329, 5864, 6262, 7769, and 9608 differentially expressed genes, respectively (Figure 7). While many of these differentially expressed genes were specific to individual tissue types (blue bars shown in Figure 4), 349 were differentially expressed in a similar manner across all tissue types (red bar shown in Figure 7). Genes that were significantly upregulated by MeJA include *GGT1* (glutathione hydrolase 1), *GGPS* (geranylgeranyl pyrophosphate synthase), and *OMR1* (threonine dehydratase 1). Genes that were consistently

downregulated include *AMT1-2* (ammonium transporter 1 member 2), *CYP450-98A2* (cytochrome P450 98A2), and *ECR26* (eceriferum 26) (Figure 8). Functional annotation resulted in 89% (*i.e.*, 310/349) of the differentially expressed genes being assigned putative functions (Figure 8 and Table S3).

Analysis of gene expression changes indicated that MeJA treatment elicited a robust defense response in *A. syriaca.* Across all six tissue types, 18 genes associated with jasmonic acid biosynthesis and signaling were consistently upregulated upon MeJA treatment (Figure 8 and Table S3) (Ruan et al., 2019; Li et al., 2022; Hewedy et al., 2023). These genes included five *JAZ* (jasmonate-ZIM domain) genes, which play a crucial role in the jasmonate signaling cascade, as well as *ACOX1* (acyl-coezyme A oxidase 1), *AOC* (allene oxide cyclase), *AOS* (allene oxide synthase), two *LOX* (linoleate 13S-lipoxygenases) genes, *KAT2* (3-ketoacyl-CoA thiolase 2), *4CLL5* (4-coumarate-CoA ligase-like 5), *NINJA* (novel interactor of JAZ), and *MYC2* (a jasmonate-responsive transcription factor) (Figure 8 and Table S3). In addition, MeJA treatment induced other genes associated with plant defense responses, including *ITH* (inhibitor of trypsin and hageman factor), a proteinase inhibitor that is potentially involved in plant resistance to herbivory (Chen et al., 1992), as well as *PAMP-A70* (pathogen-associated molecular patterns-induced protein A70) and *R1A-10* (late blight resistance protein homolog), which are both associated with plant disease resistance (Figure 8 and Table S3).

In addition to triggering a general plant resistance response, MeJA treatment elicited expression changes in genes potentially involved in cardiac glycoside biosynthesis. Among those genes, two were annotated as *SMO* (squalene monooxygenase) and *SS1* (squalene synthase 1), which are involved in the final steps of terpenoid backbone biosynthesis (Kreis et al., 1998; Kreis and Müller-Uri, 2013; Zheng et al., 2014; Pandey et al., 2016; Zhang et al., 2017). One gene was annotated as *DWF1* or *DHCR24* (delta(24)-sterol reductase), which is associated with sterol biosynthesis to generate cholesterol for cardiac glycoside biosynthesis (Zerenturk et al., 2013; Tsukagoshi et al., 2016; Knoch et al., 2018). In addition, we also found 14 other cytochrome P450 genes from families that are involved in specialized metabolite biosynthesis in other species, including *CYP711A1, CYP714A1, CYP72A219, CYP81E8, CYP90B1, CYP94C1, CYP98A2, CYP450-AOS*, and *CYP450-HPL*. While homologs of *CYP94C1, CYP450-AOS* and *CYP450-HPL* were annotated to be involved in JA biosynthesis and signaling (Ruan et al., 2019; Li et al., 2022), other cytochrome P450s are potentially involved in cardiac glycoside

biosynthesis. For example, in *Digitalis* spp., *C. gigantea*, and *E. cheiranthoides*, enzymes in the CYP87A family catalyze the first step in cardiac glycoside biosynthesis from phytosterols (Carroll et al., 2023; Kunert et al., 2023; Younkin et al., 2024), and there is likely a homolog in *A. curassavica* (Figures S4 and S5).

Beyond enzymes involved in the biosynthesis of the steroid core, modifying enzymes can generate cardiac glycoside diversity, For example, a permissive glycosyltransferase, UGT74AN1, catalyzes steroid C3 glycosylation to produce diverse glycosides and other bioactive molecules in *A. curassavica* (Wen et al., 2018). Our differential expression analysis revealed different groups of enzymes that may contribute to secondary modifications that generate the extensive cardiac glycoside diversity found in milkweed (Figure 8 and Table S3). These enzymatic activities encoded by these genes include kinase, synthase, acyltransferase, glycosyl/glucosyltransferase, reductase, oxidase, dehydrogenase, ligase, hydrogenase, phosphatase, epimerase, esterase, cyclase, and more. Many identified enzymes in cardiac glycoside biosynthesis from other species belong to those enzymatic families, such as progesterone 5β-reductase (Herl et al., 2008; Herl et al., 2008), 3β-hydroxysteroid dehydrogenase (Finsterbusch et al., 1999; Munkert et al., 2014), and UDPG glucosyltransferase (Yoshikawa and Furuya, 1979). However, while confirming homology between our candidates to published sequences is possible, their functions in milkweed remain to be tested.

**Discussion**

We have assembled and annotated a high-quality *A. curassavica* reference genome for use in ecological and molecular studies. By also sequencing the transcriptomes of six tissue types, with and without MeJA elicitation, we have identified both tissue-specific gene expression (Figure 7) and induced expression of genes that are likely to be involved in plant defenses (Figure 8). Previous milkweed research demonstrated tissue-specific accumulation of cardiac glycosides and resistance to insect herbivory (López-Goldar et al., 2022; Agrawal and Hastings, 2023). The tissue-specific milkweed transcriptomes described here will facilitate future research to identify the genetic basis of this within-plant natural variation in milkweed defenses.

Although the *A. curassavica* genome is smaller and has fewer LTR elements than the previously assembled *A. syriaca* genome (Figure 2), the two genomes are almost completely colinear (Figure 3). Therefore, it is likely that results from functional genomic analyses with *A.*

*curassavica* can be directly applied for research with *A. syriaca* and possibly other milkweed species for which genome sequences are not yet available. Comparative genomics studies will make it possible to identify genes that are present in one species but not another, *e.g.*, genes that are required for species-specific differences in the production of cardiac glycosides and other defense-related metabolites.

Prior to sequencing the *A. curassavica* genome, we conducted five generations of inbreeding by hand-pollinating individual plants. The resulting low level of DNA sequence heterozygosity facilitated our *A. curassavica* genome assembly. The availability of this inbred line will increase the reproducibility of future experiments to study the function of individual milkweed genes in ecological interactions and the biosynthesis of medically relevant specialized metabolites. To make our genome-sequenced *A. curassavica* lineage a permanent resource for milkweed researchers, we have submitted sixth-generation inbred seeds to a public repository, the USDA-ARS Ornamental Plant Germplasm Center (https://opgc.osu.edu/ ; https://www.ars-grin.gov/ ; accession number OPGC 7586).

**Data Availability**

DNA and RNA sequences produced for this research have been deposited in GenBank, (PRJNA1055959). Seeds of the inbred line of *A. curassavica* that was used for genome sequencing are available from the USDA-ARS Ornamental Plant Germplasm Center, accession number OPGC 7586.

**Supplementary Materials**

**Figure S1.** Genome heterozygosity, repeat content, and size estimate
**Figure S2.** Hi-C contact heat map of assembled pseudomolecules
**Figure S3.** Feature density along the A. curassavica pseudomolecules
**Figure S4.** Alignment of CYP87A4 proteins
**Figure S5.** Phylogenetic tree of CYP87A4 proteins
**Table S1.** Genome assembly metrics
**Table S2.** Analysis of *A. curassavica* gene families
**Table S3.** Differentially expressed genes

**Competing Interests**

The authors declare that they have no competing interests.

**Acknowledgements**

This study was funded by US National Science Foundation awards 1645256 and 1850796, United States Department of Agriculture – National Institute of Food and Agriculture award 2020-67013-30896, and the Triad Foundation. We thank Arielle Johnson for assistance with the DNA sample preparation.

**Authors' Contributions**

SBB, FW, and GJ created an inbred line of *A. curassavica*; HF, AFP, JZ, LY, EY, and SRS assembled and annotated the genome; GLB conducted gene family analysis, HF generated and analyzed transcriptome data; and HF, SRS, and GJ wrote the manuscript.

**References**

**Agrawal AA** (2017) Monarchs and Milkweed: A Migrating Butterfly, a Poisonous Plant, and Their Remarkable Story of Coevolution. Princeton University Press, Princeton, NJ

**Agrawal AA, Hastings AP** (2023) Tissue-specific plant toxins and adaptation in a specialist root herbivore. Proc Natl Acad Sci U S A **120**: e2302251120

**Alexa A, Rahnenfuhrer J** (2023) topGO: Enrichment Analysis for Gene Ontology.

**Alonso-Castro AJ, Arana-Argaez V, Yanez-Barrientos E, Torres-Romero JC, Chable-Cetz RJ, Worbel K, Euan-Canto AJ, Wrobel K, Gonzalez-Ibarra A, Solorio-Alvarado CR, et al** (2021) Pharmacological activities of *Asclepias curassavica* L. (Apocynaceae) aerial parts. J Ethnopharmacol **281**: 114554

**Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ** (1990) Basic local alignment search tool. J Mol Biol **215**: 403–410

**Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ** (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. Nucleic Acids Res **25**: 3389–402

**Bianchini G, Sánchez-Baracaldo P** (2023) TreeViewer - Cross-platform software to draw phylogenetic trees. doi: 10.5281/zenodo.10044388

**Boutet E, Lieberherr D, Tognolli M, Schneider M, Bairoch A** (2007) UniProtKB/Swiss-Prot. *In* D Edwards, ed, Plant Bioinformatics: Methods and Protocols. Humana Press, Totowa, NJ, pp 89–112

**Boyle JH, Strickler SR, Twyford A, Ricono A, Powell AF, Zhang J, Xu H, Dalgleish HJ, Jander G, Agrawal AA, et al** (2022) Temporal matches and mismatches between monarch butterfly and milkweed population changes over the past 12,000 years. bioRxiv https://doi.org/10.1101/2022.02.25.481796:

**Campbell MS, Law M, Holt C, Stein JC, Moghe GD, Hufnagel DE, Lei J, Achawanantakun R, Jiao D, Lawrence CJ, et al** (2014) MAKER-P: a tool kit for the rapid creation, management, and quality control of plant genome annotations. Plant Physiol **164**: 513–524

**Cantarel BL, Korf I, Robb SMC, Parra G, Ross E, Moore B, Holt C, Sánchez Alvarado A, Yandell M** (2008) MAKER: an easy-to-use annotation pipeline designed for emerging model organism genomes. Genome Res **18**: 188–196

**Carroll E, Ravi Gopal B, Raghavan I, Mukherjee M, Wang ZQ** (2023) A cytochrome P450 CYP87A4 imparts sterol side-chain cleavage in digoxin biosynthesis. Nat Commun **14**: 4042

**Chen M-S, Feng G, Zen KC, Richardson M, Valdes-Rodriguez S, Reeck GR, Kramer KJ** (1992) α-amylases from three species of stored grain coleoptera and their inhibition by wheat and corn proteinaceous inhibitors. Insect Biochemistry and Molecular Biology **22**: 261–268

**Conway JR, Lex A, Gehlenborg N** (2017) UpSetR: an R package for the visualization of intersecting sets and their properties. Bioinformatics **33**: 2938–2940

**Denoeud F, Carretero-Paulet L, Dereeper A, Droc G, Guyot R, Pietrella M, Zheng C, Alberti A, Anthony F, Aprea G, et al** (2014) The coffee genome provides insight into the convergent evolution of caffeine biosynthesis. Science **345**: 1181–1184

**Dobler S, Dalla S, Wagschal V, Agrawal AA** (2012) Community-wide convergent evolution in insect adaptation to toxic cardenolides by substitutions in the Na,K-ATPase. Proc Natl Acad Sci U S A **109**: 13040–13045

**Dudchenko O, Batra SS, Omer AD, Nyquist SK, Hoeger M, Durand NC, Shamim MS, Machol I, Lander ES, Aiden AP, et al** (2017) *De novo* assembly of the *Aedes aegypti* genome using Hi-C yields chromosome-length scaffolds. Science **356**: 92–95

**Durand NC, Robinson JT, Shamim MS, Machol I, Mesirov JP, Lander ES, Aiden EL** (2016) Juicebox provides a visualization system for Hi-C contact maps with unlimited zoom. Cell Syst **3**: 99–101

**Ellinghaus D, Kurtz S, Willhoeft U** (2008) LTRharvest, an efficient and flexible software for de novo detection of LTR retrotransposons. BMC Bioinformatics **9**: 18

**Finsterbusch A, Lindemann P, Grimm R, Eckerskorn C, Luckner M** (1999) Δ5-3β-Hydroxysteroid dehydrogenase from *Digitalis lanata* Ehrh. – a multifunctional enzyme in steroid metabolism? Planta **209**: 478–486

**Frank MH, Edwards MB, Schultz ER, McKain MR, Fei Z, Sorensen I, Rose JK, Scanlon MJ** (2015) Dissecting the molecular signatures of apical cell-type shoot meristems from two ancient land plant lineages. New Phytol **207**: 893–904

**Fulton M, Chunwongse J, Tanksley SD** (1995) Microprep protocol for extraction of DNA from tomato and other herbaceous plants. Plant Mol Biol Rep **13**: 207–209

**Han MV, Thomas GWC, Lugo-Martinez J, Hahn MW** (2013) Estimating gene gain and loss rates in the presence of error in genome assembly and annotation using CAFE 3. Mol Biol Evol **30**: 1987–1997

**Herl V, Albach DC, Müller-Uri F, Bräuchler C, Heubl G, Kreis W** (2008) Using progesterone 5β-reductase, a gene encoding a key enzyme in the cardenolide biosynthesis, to infer the phylogeny of the genus *Digitalis*. Plant Syst Evol **271**: 65–78

**Hewedy OA, Elsheery NI, Karkour AM, Elhamouly N, Arafa RA, Mahmoud GA-E, Dawood MF-A, Hussein WE, Mansour A, Amin DH, et al** (2023) Jasmonic acid regulates plant development and orchestrates stress response during tough times. Environmental and Experimental Botany **208**: 105260

**Hoopes GM, Hamilton JP, Kim J, Zhao D, Wiegert-Rininger K, Crisovan E, Buell CR** (2018) Genome assembly annotation of the medicinal plant *Calotropis gigantea*, a producer of anticancer and antimalarial cardenolides. G3 **8**: 385–391

**Jacobo-Salcedo Mdel R, Alonso-Castro AJ, Salazar-Olivo LA, Carranza-Alvarez C, Gonzalez-Espindola LA, Dominguez F, Maciel-Torres SP, Garcia-Lujan C, Gonzalez-Martinez Mdel R, Gomez-Sanchez M, et al** (2011) Antimicrobial and cytotoxic effects of Mexican medicinal plants. Nat Prod Commun **6**: 1925–8

Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J, Mitchell A, Nuka G, et al (2014) InterProScan 5: genome-scale protein function classification. Bioinformatics **30**: 1236–1240

Karageorgi M, Groen SC, Sumbul F, Pelaez JN, Verster KI, Aguilar JM, Hastings AP, Bernstein SL, Matsunaga T, Astourian M, et al (2019) Genome editing retraces the evolution of toxin resistance in the monarch butterfly. Nature **574**: 409–412

Kellner F, Kim J, Clavijo BJ, Hamilton JP, Childs KL, Vaillancourt B, Cepela J, Habermann M, Steuernagel B, Clissold L, et al (2015) Genome-guided investigation of plant natural product biosynthesis. Plant J **82**: 680–692

Kim D, Langmead B, Salzberg SL (2015) HISAT: a fast spliced aligner with low memory requirements. Nat Methods **12**: 357–360

Knight A, Brower LP (2009) The influence of eastern North American autumnal migrant monarch butterflies (*Danaus plexippus* L.) on continuously breeding resident monarch populations in southern Florida. J Chem Ecol **35**: 816–23

Knoch E, Sugawara S, Mori T, Poulsen C, Fukushima A, Harholt J, Fujimoto Y, Umemoto N, Saito K (2018) Third DWF1 paralog in Solanaceae, sterol Δ24-isomerase, branches withanolide biosynthesis from the general phytosterol pathway. Proc Natl Acad Sci U S A **115**: E8096–E8103

Kolde R (2018) Pheatmap: Pretty heatmaps, R package version 1.0.10. https://cran.microsoft.com/snapshot/2018-06-22/web/packages/pheatmap/pheatmap.pdf

Korf I (2004) Gene finding in novel genomes. BMC Bioinformatics **5**: 59

Kreis W, Hensel A, Stuhlemmer U (1998) Cardenolide biosynthesis in foxglove. Planta Med **64**: 491–499

Kreis W, Müller-Uri F (2013) Cardenolide Aglycone Formation in Digitalis. *In* TJ Bach, M Rohmer, eds, Isoprenoid Synthesis in Plants and Microorganisms: New Concepts and Experimental Approaches. Springer, New York, NY, pp 425–438

Krzywinski M, Schein J, Birol İ, Connors J, Gascoyne R, Horsman D, Jones SJ, Marra MA (2009) Circos: An information aesthetic for comparative genomics. Genome Res **19**: 1639–1645

Kunert M, Langley C, Lucier R, Ploss K, Rodríguez López CE, Serna Guerrero DA, Rothe E, O'Connor SE, Sonawane PD (2023) Promiscuous CYP87A enzyme activity initiates cardenolide biosynthesis in plants. Nat Plants **9**: 1607–1617

**Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg SL** (2004) Versatile and open software for comparing large genomes. Genome Biology **5**: R12

**Langmead B, Salzberg SL** (2012) Fast gapped-read alignment with Bowtie 2. Nature methods **9**: 357–9

**Lau KH, Bhat WW, Hamilton JP, Wood JC, Vaillancourt B, Wiegert-Rininger K, Newton L, Hamberger B, Holmes D, Hamberger B, et al** (2020) Genome assembly of *Chiococca alba* uncovers key enzymes involved in the biosynthesis of unusual terpenoids. DNA Res. doi: 10.1093/dnares/dsaa013

**Lewis M, Ruter JM** (2023) Use of flow cytometry to assess total genomic content for *Asclepias* spp. HortScience **58**: 583–587

**Li C, Xu M, Cai X, Han Z, Si J, Chen D** (2022) Jasmonate signaling pathway modulates plant defense, growth, and their trade-offs. International Journal of Molecular Sciences **23**: 3945

**Liang Y, Li F, Gao Q, Jin C, Dong L, Wang Q, Xu M, Sun F, Bi B, Zhao P, et al** (2022) The genome of *Eustoma grandiflorum* reveals the whole-genome triplication event contributing to ornamental traits in cultivated lisianthus. Plant Biotechnol J **20**: 1856–1858

**Linnaeus C** (1754) Genera Plantarum, ed. 5. Impensis Laurentii Salvii, Stockholm

**López-Goldar X, Hastings A, Züst T, Agrawal A** (2022) Evidence for tissue-specific defence-offence interactions between milkweed and its community of specialized herbivores. Mol Ecol **31**: 3254–3265

**Mapleson D, Venturini L, Kaithakottil G, Swarbreck D** (2018) Efficient and accurate detection of splice junctions from RNA-seq with Portcullis. Gigascience. doi: 10.1093/gigascience/giy131

**Marçais G, Kingsford C** (2011) A fast, lock-free approach for efficient parallel counting of occurrences of k-mers. Bioinformatics **27**: 764–770

**Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, Lanfear R** (2020) IQ-TREE 2: New models and efficient methods for phylogenetic inference in the genomic era. Molecular Biology and Evolution **37**: 1530–1534

**Munkert J, Ernst M, Müller-Uri F, Kreis W** (2014) Identification and stress-induced expression of three 3β-hydroxysteroid dehydrogenases from *Erysimum crepidifolium* Rchb. and their putative role in cardenolide biosynthesis. Phytochemistry **100**: 26–33

**Navarro Garcia VM, Gonzalez A, Fuentes M, Aviles M, Rios MY, Zepeda G, Rojas MG** (2003) Antifungal activities of nine traditional Mexican medicinal plants. J Ethnopharmacol **87**: 85–8

**Ou S, Jiang N** (2018) LTR_retriever: A highly accurate and sensitive program for identification of long terminal repeat retrotransposons. Plant Physiol **176**: 1410–1422

**Pandey A, Swarnkar V, Pandey T, Srivastava P, Kanojiya S, Mishra DK, Tripathi V** (2016) Transcriptome and metabolite analysis reveal candidate genes of the cardiac glycoside biosynthetic pathway from Calotropis procera. Sci Rep **6**: 34464

**Paradis E, Claude J, Strimmer K** (2004) APE: Analyses of Phylogenetics and Evolution in R language. Bioinformatics **20**: 289–290

**Peraza-Sanchez SR, Cen-Pacheco F, Noh-Chimal A, May-Pat F, Sima-Polanco P, Dumonteil E, Garcia-Miss MR, Mut-Martin M** (2007) Leishmanicidal evaluation of extracts from native plants of the Yucatan peninsula. Fitoterapia **78**: 315–8

**Petschenka G, Wagschal V, von Tschirnhaus M, Donath A, Dobler S** (2017) Convergently evolved toxic secondary metabolites in plants drive the parallel molecular evolution of insect resistance. Am Nat **190**: S29–S43

**Quattrocchi U** (1999) CRC World Dictionary of Plant Names: Common Names, Scientific Names, Eponyms, Synonyms, and Etymology. CRC Press, Boca Raton, FL

**Rehman R, Hai O** (2023) Digitalis Toxicity. StatPearls

**Roach MJ, Schmidt SA, Borneman AR** (2018) Purge Haplotigs: allelic contig reassignment for third-gen diploid genome assemblies. BMC Bioinformatics **19**: 460

**Robinson MD, McCarthy DJ, Smyth GK** (2010) edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics **26**: 139–140

**Ruan J, Zhou Y, Zhou M, Yan J, Khurshid M, Weng W, Cheng J, Zhang K** (2019) Jasmonic acid signaling pathway in plants. International Journal of Molecular Sciences **20**: 2479

**Sabir JSM, Jansen RK, Arasappan D, Calderon V, Noutahi E, Zheng C, Park S, Sabir MJ, Baeshen MN, Hajrah NH, et al** (2016) The nuclear genome of *Rhazya stricta* and the evolution of alkaloid diversity in a medically relevant clade of Apocynaceae. Sci Rep **6**: 33782

**Schäffer AA, Aravind L, Madden TL, Shavirin S, Spouge JL, Wolf YI, Koonin EV, Altschul SF** (2001) Improving the accuracy of PSI-BLAST protein database searches with composition-based statistics and other refinements. Nucleic Acids Res **29**: 2994–3005

**Smit AFA, Hubley R** (2015) RepeatModeler Open-1.0.

**Smit AFA, Hubley R, Green P** (2015) RepeatMasker Open-4.0. 2013-2015. http://www.repeatmasker.org

**Stanke M, Diekhans M, Baertsch R, Haussler D** (2008) Using native and syntenically mapped cDNA alignments to improve de novo gene finding. Bioinformatics **24**: 637–644

**Straub SC, Fishbein M, Livshultz T, Foster Z, Parks M, Weitemier K, Cronn RC, Liston A** (2011) Building a model: developing genomic resources for common milkweed (*Asclepias syriaca*) with low coverage genome sequencing. BMC Genomics **12**: 211

**Tomov I** (1981) Cardiac glycoside therapy of congestive heart failure. II. Vutr Boles **20**: 1–9

**Törönen P, Medlar A, Holm L** (2018) PANNZER2: a rapid functional annotation web server. Nucleic Acids Res **46**: W84–W88

**Tsukagoshi Y, Suzuki H, Seki H, Muranaka T, Ohyama K, Fujimoto Y** (2016) Ajuga Δ24-sterol reductase catalyzes the direct reductive conversion of 24-methylenecholesterol to campesterol. J Biol Chem **291**: 8189–8198

**Venturini L, Caim S, Kaithakottil GG, Mapleson DL, Swarbreck D** (2018) Leveraging multiple transcriptome assembly methods for improved gene structure annotation. Gigascience. doi: 10.1093/gigascience/giy093

**Vurture GW, Sedlazeck FJ, Nattestad M, Underwood CJ, Fang H, Gurtowski J, Schatz MC** (2017) GenomeScope: fast reference-free genome profiling from short reads. Bioinformatics **33**: 2202–2204

**Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q, Wortman J, Young SK, et al** (2014) Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. PLoS One **9**: e112963

**Waterhouse RM, Seppey M, Simão FA, Manni M, Ioannidis P, Klioutchnikov G, Kriventseva EV, Zdobnov EM** (2017) BUSCO applications from quality assessments to gene prediction and phylogenomics. Mol Biol Evol. doi: 10.1093/molbev/msx319

**Weitemier K, Straub SCK, Fishbein M, Bailey CD, Cronn RC, Liston A** (2019) A draft genome and transcriptome of common milkweed (*Asclepias syriaca*) as resources for evolutionary, ecological, and molecular studies in milkweeds and Apocynaceae. PeerJ **7**: e7649

**Wen C, Huang W, Zhu X-L, Li X-S, Zhang F, Jiang R-W** (2018) UGT74AN1, a Permissive Glycosyltransferase from Asclepias curassavica for the Regiospecific Steroid 3-O-Glycosylation. Org Lett **20**: 534–537

**Woodson RE** (1954) The North American species of *Asclepias*. Ann Miss Bot Gard **41**: 1–211

**Wyatt R, Broyles SB** (1997) The weedy tropical milkweeds *Asclepias curassavica* and *A. fruticosa* are self-compatible. Biotropica **29**: 232–234

**Xu G-C, Xu T-J, Zhu R, Zhang Y, Li S-Q, Wang H-W, Li J-T** (2019) LR_Gapcloser: a tiling path-based gap closer that uses long reads to complete genome assembly. Gigascience. doi: 10.1093/gigascience/giy157

**Xu Z, Wang H** (2007) LTR_FINDER: an efficient tool for the prediction of full-length LTR retrotransposons. Nucleic Acids Res **35**: W265-8

**Yoshikawa T, Furuya T** (1979) Purification and properties of sterol: UDPG glucosyltransferase in cell culture of *Digitalis purpurea*. Phytochemistry **18**: 239–241

**Younkin GC, Alani ML, Capador AP, Fischer HD, Mirzaei M, Hastings AP, Agrawal AA, Jander G** (2024) Cardiac glycosides protect wormseed wallflower (*Erysimum cheiranthoides*) against some, but not all, glucosinolate-adapted herbivores. New Phytologist in press:

**Yuan WQ, Zhang RR, Wang J, Ma Y, Li WX, Jiang RW, Cai SH** (2016) Asclepiasterol, a novel C21 steroidal glycoside derived from *Asclepias curassavica*, reverses tumor multidrug resistance by down-regulating P-glycoprotein expression. Oncotarget **7**: 31466–83

**Zerenturk EJ, Sharpe LJ, Ikonen E, Brown AJ** (2013) Desmosterol and DHCR24: unexpected new directions for a terminal step in cholesterol synthesis. Prog Lipid Res **52**: 666–680

**Zhang J, Li X, Lu F, Wang S, An Y, Su X, Li X, Ma L, Han G** (2017) *De novo* sequencing and transcriptome analysis reveal key genes regulating steroid metabolism in leaves, roots, adventitious roots and calli of *Periploca sepium* Bunge. Frontiers in Plant Science 8:594

**Zhao X, Hu X, OuYang K, Yang J, Que Q, Long J, Zhang J, Zhang T, Wang X, Gao J, et al** (2022) Chromosome-level assembly of the *Neolamarckia cadamba* genome provides insights into the evolution of cadambine biosynthesis. Plant J **109**: 891–908

**Zheng K, Zhang G, Jiang N, Yang S, Li C, Meng Z, Guo Q, Long G** (2014) Analysis of the transcriptome of *Marsdenia tenacissima* discovers putative polyoxypregnane glycoside biosynthetic genes and genetic markers. Genomics **104**: 186–193

**Zheng X, Xu Y, Liu B, Qi Y, Zhang Y, Li X, Zhang X, Pu X, Li S, Chen Z, et al** (2019) Ethyl acetate extract of *Asclepias curassavica* induced apoptosis in human cancer cells via activating p38 and JNK MAPK signaling pathways. Evid Based Complement Alternat Med **2019**: 9076269

**Zimin AV, Marçais G, Puiu D, Roberts M, Salzberg SL, Yorke JA** (2013) The MaSuRCA genome assembler. Bioinformatics **29**: 2669–2677

A.

 B.

**Figure 1.** *Asclepias curassavica*, **tropical milkweed.** A. Structure of voruscharin, the most abundant cardiac glycoside in *A. curassavica*. B. Flowers of the greenhouse-grown *A. curassavica* lineage that was used for the current study.

**Figure 2: Pie charts showing percentages of genome features in the** *Asclepias syriaca and Asclepias curassavica* **pseudomolecules.** The size of each pie is scaled to the genome size of the species.

**Figure 3: Synteny plot showing regions of collinearity between the *Asclepias curassavica* and *Asclepias syriaca* pseudomolecules.** Sequence matches were calculated with nucmer and plotted using Circos.

**Figure 4: Upset plot showing the intersection of gene families among *Asclepias curassavica* and sequenced near relatives.** The numbers of gene families are indicated for each species and species intersection.

**Figure 5: Species phylogeny showing gene family expansions and contractions.**
Expansions are show in green and contractions are shown in red.

**Figure 6. Principal component analyses (PCA) of RNAseq data.** (A) PCA of control samples treated with 0.1% ethanol (B) PCA of methyl jasmonic acid (MeJA)-treated samples. (C-H) comparison of control samples (green dots) and MeJA-treated samples (blue dots) of (C) buds, (D) flowers, (E) mature leaves, (F) roots, (G) stems, and (H) young leaves.

**Figure 7. Number of differentially expressed genes across tissue types after methyl jasmonic acid (MeJA) treatment.** The horizontal bar plot in the inset box shows the number of differentially expressed genes (MeJA-treated *vs*. control) in each of the six tested tissue types. The intersection plot shows the number of differentially expressed genes across different tissue type combinations. The red bar represents the number of core differentially expressed genes across all tissue types, blue bars represent the number of differentially expressed genes that are specific to each tissue type, and black bars represent other tissue intersections indicated by the dot connections under the bar plot. This plot was generated using UpSetR v1.4.0.

**Figure 8. Heatmap of $\log_2$ fold changes of commonly differentially expressed genes across different tissues.** $\log_2$ fold changes were calculated with methyl jasmonate (MeJA)-treated samples, comparing to control samples within each tissue type. Red represents upregulation, while blue represents downregulation. Tissue (in columns) and gene (in rows) clusters were calculated using default method in pheatmap to facilitate pattern identification. Gene identifications are listed in Supplemental Table S2.