Investigating the molecular basis of cleptobiosis in eusocial stingless bees (Apidae: Hymenoptera)

Paulo Cseri Ricardo[a]*, Natalia De Souza Araujo[b] and Maria Cristina Arias[a]

[a]*Departamento de Genética e Biologia Evolutiva – Instituto de Biociências, Universidade de São Paulo, São Paulo, Brazil*
[b] *Unit of Evolutionary Biology & Ecology – Université libre de Bruxelles, Brussels, Belgium.*
*Corresponding author: P. C. Ricardo - cseri.bio@gmail.com

Abstract

Cleptobiosis, the act of raiding other species to obtain food or resources, is widespread among animals but rarely obligatory. The eusocial stingless bee *Lestrimelitta limao* is one of the few species that depend entirely on cleptobiosis, raiding other stingless bee colonies for survival. To investigate the molecular basis and evolutionary adaptations associated with this specialized lifestyle, we compared the transcriptomes of foraging workers of *L. limao* and three non-robber stingless bees – *Nannotrigona testaceicornis*, *Scaptotrigona* aff. *depilis*, and *Tetragonisca angustula*.

Our analysis revealed that differentially expressed orthologs were predominantly downregulated in *L. limao* workers, suggesting reduced transcriptional activity during foraging in this species. These downregulated genes fall into three major functional categories potentially linked to cleptobiotic adaptations: (1) detoxification and chemoreception genes, including cytochrome P450s and odorant receptors, indicating decreased exposure to phytochemicals; (2) neuronal and synaptic genes, such as *para* and *Dys*, possibly reflecting neurophysiological modifications; and (3) mitochondrial and carbohydrate metabolism genes, suggesting lower energetic demands. These findings provide novel insights into the molecular mechanisms shaping cleptobiotic behavior in eusocial bees.

Keywords

Lestrimelitta; robber bees; Meliponini; Neotropical bees; parasitism

1. Introduction

Cleptobiosis is an ecological relationship in which an individual or a group of individuals from the same species loot food or other valuable resources from another

(Breed et al., 2012). This behavior is widespread in vertebrates and invertebrates and can be facultative or obligatory, although obligate cleptobiosis is far less common (reviewed in Iyengar 2008). Often the term cleptobiosis is associated with species of social insects, especially ants and bees (Breed et al., 2012). It was first proposed as a classification for the ecological interactions of thief ants (Forel, 1901; Wheeler, 1910), where several cases of inter and intraspecific food theft are known (Hölldobler, 1986; Perfecto and Vandermeer, 1993; Yamaguchi, 1995; Richard et al., 2004; Ronque et al., 2018; Yamaguchi, 1995).

Among eusocial bees, facultative cleptobiosis is common in species from the family Apidae encompassing the tribes Meliponini (stingless bees) and Apini (honeybees), especially when species are facing food shortage (Bohart, 1970; Costa et al., 2018; Michener, 2007; Wcislo, 1987). However, obligate cleptobiosis has only been confirmed in two genera of stingless bees: *Lestrimelitta* and *Cleptotrigona*. Species of *Trichotrigona* are also likely to show obligate cleptobiosis based on their nest architecture, but no direct observations of this behavior have been recorded so far (Camargo and Pedro, 2007; Pedro and Cordeiro, 2015). *Lestrimelitta* is a Neotropical genus comprising 26 described species, all exhibiting obligatory cleptobiotic behavior (Engel et al., 2023). In contrast, the Afrotropical genus *Cleptotrigona* includes only one extant species widely distributed across sub-Saharan Africa (Eardley, 2004), *Cleptotrigona cubiceps* (Lepeco et al., 2024; Portugal-Araújo, 1958). *Cleptotrigona* belongs to the Afrotropical Meliponini clade, while *Lestrimelitta* belongs to the Neotropical Meliponini clade (Rasmussen and Cameron, 2009). These two clades diverged over 70 million years ago (Rasmussen and Cameron, 2009). Thus, obligatory cleptobiosis in bees arose independently from flower-collecting ancestors at least twice.

Lestrimelitta and Cleptotrigona species attack nests of other stingless bee species to rob larval food, pollen, honey, and cerumen (Nogueira-Neto, 1997; Portugal-Araújo, 1958; Sakagami et al., 1993; Wittmann et al., 1990). Accordingly, these species have morphological and behavioral adaptations likely driven by their cleptobiotic behavior. Maybe the most striking of these adaptations is the absence of a functional corbicula, a morphological feature common to all corbiculate bees (Eardley, 2004; Michener, 2007; Shôichi F Sakagami and Laroca, 1963; Wille, 1983). Indeed, the reduction or complete loss of pollen-collecting structures seems to be a convergent adaptation in parasitic bees, as it is also observed in cleptoparasitic (*e.g.,* Nomadinae bees brood parasites) and social parasitic species (*e.g.*, *Bombus Psithyrus* subgenus) (Michener, 2000). Queens of *Lestrimelitta* have a higher number of ovarioles per ovary (10-15) than most stingless bees (~4) (Sakagami, 1982), enabling the production of more workers. Moreover, the robust mandibles of the *Lestrimelitta* workers (James et al., 2022; Nogueira-Neto, 1997) help them overcome the colony defenses of host species.

An interesting behavioral adaptation in *Lestrimelitta* and *Cleptotrigona* species is their foraging strategy, as they carry out organized robbing raids in which several individuals simultaneously attack colonies of other species; in *C. cubiceps* the number of individuals participating in these raids may reach a few dozen (Portugal-Araújo, 1958), while in *Lestrimelitta* it could exceed several hundred (Sakagami et al., 1993). Raiding recruitment in *Lestrimelitta* has been reported to involve the use of semiochemicals (Blum, 1966; Sakagami et al., 1993) that have toxin functions (James et al., 2022), and may also play a role in the host response to raids (James et al., 2022; von Zuben et al., 2016). Likewise, in an apparent strategy of chemical deception, it has been

suggested that *Lestrimelitta* may share cuticular hydrocarbons with their preferred host species to better conceal their presence (Vázquez et al., 2022).

Among all robber bees, *Lestrimelitta limao* is the best-known and most studied species (Rech et al., 2013). They are named after the citral volatile compound produced and released by workers, a compound common to all *Lestrimelitta* species (Blum et al., 1970; Francke et al., 2000). Like other species of *Lestrimelitta*, *L. limao* is specialized in raiding stingless bee nests (Wittmann et al., 1990), although records of pillaging in honeybee colonies exist (Sakagami et al., 1993). There are records of *L. limao* raids in more than a dozen stingless bee species from different genera, including *Friesella*, *Frieseomelitta*, *Melipona*, *Nannotrigona*, *Plebeia*, *Nannotrigona*, *Scaptotrigona*, *Tetragonisca* and *Trigonisca* (reviewed in Grüter et al. 2016). Nevertheless, *L. limao* host preference is apparently opportunistic depending on the location and the composition of the local stingless bee fauna (Sakagami and Laroca, 1963). *L. limao* raid attacks are common enough for this species to be considered a relevant commercial threat to bee-raising activities (meliponiculture), as it frequently attacks nests of species of commercial interest, killing many individuals and potentially leading to colony destruction (Sakagami et al., 1993; Sakagami and Laroca, 1963). In natural environments, it has been hypothesized that this species may control the size of stingless bee populations (Nogueira-Neto, 1997). In turn, evidence suggests that cleptobiotic interactions between *L. limao* and other stingless bee species may have driven the evolution of a soldier sub-caste in the assaulted species, which would be specialized in colony defense (Grüter et al., 2017, 2012). The number and size of these defensive soldiers are affected by the colony exposure to cleptobionts (Segers et al., 2016), corroborating the hypothesis that cleptobiosis would represent a relevant selective pressure in driving the evolution of host species.

Despite the evolutionary, ecological, and economic relevance of cleptobiosis in stingless bees, the mechanisms involved in the foraging strategy of *L. limao* and how they diverge from other non-parasitic bees are poorly understood. Likewise, the adaptations at the molecular level related to the evolution and maintenance of obligatory cleptobiosis in eusocial bees are unknown. Herein, we investigated the broad molecular mechanisms involved in stingless bees cleptobiosis using comparative transcriptomics. We generated transcriptomic data from foragers of the robber bee *L. limao* and for three pollen-collecting (non-parasitic) stingless bees, *Nannotrigona testaceicornis*, *Scaptotrigona* aff. *depilis*, and *Tetragonisca angustula*. These data were comparatively analyzed to identify and describe the molecular pathways associated with the cleptobiotic strategy of this robber bee during foraging.

2. Methods

2.1. Sampling and RNA sequencing

Workers were collected while performing foraging activities between 11:00 am and 1:00 pm to control for clock gene differences. We used foragers for the transcriptome comparative analyses as they are responsible for collecting food resources. The workers of *L. limao* were collected from three natural colonies in the city of São Paulo (Brazil); these workers were sampled when returning to their nests. Foragers of *N. testaceicornis* and *S.* aff. *depilis* were obtained from three rational colonies housed at the Bee Laboratory at the Universidade de São Paulo. For these species, workers were marked at their emergency and returned to their nests. After 30 days, when workers of several stingless bees are known to perform forager tasks outside the colony (see Grüter 2020), the marked bees were collected. For *T. angustula*, we used RNA-Seq data from two samples generated in a previous study, each consisting of

a pool of three adult foragers collected using a similar sampling approach (Araujo and Arias, 2021). To account for potential sequencing batch effects, we additionally sequenced one batch sample from three *T. angustula* workers collected from a nest at the Instituto de Biociências, Universidade de São Paulo. These workers were collected while returning to their colony from foraging trips. Total RNA was extracted from each bee whole body using the RNeasy® Mini Kit (Qiagen). RNA extractions from individuals of the same nest were then pooled (3 individuals/pool) to reduce individual variation expression within the colonies. These pools are hereafter referred to as samples. In summary, three samples (one per colony) of *L. limao*, *N. testaceicornis,* and *S. depilis*; and one sample of *T. angustula* were sequenced. Library construction and sequencing were performed by Macrogen (South Korea) using the Illumina® NovaSeq 6000 platform and Illumina® suggested protocols to obtain 100 bp paired-end reads.

2.2.   *Transcriptome assembly*

Read qualities were evaluated with FastQC 0.11.5 (Andrews, 2018). Raw reads were then quality-trimmed using Trimmomatic 0.38 (Bolger et al., 2014) (options: SLIDINGWINDOW:4:30 LEADING:3 TRAILING:3 MINLEN:80 AVGQUAL:30 HEADCROP:14). *De novo* transcriptome assembly was performed independently for each species using Trinity 2.10.0 (Grabherr et al., 2013; Haas et al., 2013), following default parameters, except for the *in silico* read normalization option (set up to 20x maximum coverage). The transcripts resulting from these assemblies were then clustered into SuperTranscripts (Davidson et al., 2017) using the Trinity gene splice modeler script implemented in the trinityrnaseq toolbox (Grabherr et al., 2013; Haas et al., 2013). Downstream analyses were performed using the SuperTranscripts datasets to reduce misassembly biases associated with the *de novo* approach. Quality parameters

from SuperTranscripts were analyzed with Transrate 1.0.3 (Smith-Unna et al., 2016) and BUSCO 5.2.2 (Simão et al., 2015).

2.3. *Annotation and orthologs identification*

SuperTranscripts were annotated with Annocript 2.0.1 (Musacchia et al., 2015) using the UniProt Reference Clusters (UniRef90) (Suzek et al., 2015) and UniProtKB/Swiss-Prot (Bairoch, 1996) databases from February 2021. SuperTranscripts identified as possible contaminants (*i.e.*, those annotated to Acari, Alveolata, Archaea, Bacteria, Fungi, Rhizaria, Rhodophyta, Viridiplantae and Viruses) were removed from the final data set using custom scripts (https://github.com/PauloCseri/Annotation.git). To identify orthologs among the four species, we followed the methodology from Ricardo et al. (2024). First, amino acid sequences derived from SuperTranscripts open reading frames (ORFs) generated by Annocript were used to identify orthogroups among all species using Orthofinder 2.1.1 (Emms and Kelly, 2019, 2015). Subsequently, for each orthogroup inferred by Orthofinder, only a one-to-one ortholog sequence per species was selected based on the best BLASTp alignment (BLAST 2.9.0+) (Altschul, 1997; Camacho et al., 2009) using default parameters.

2.4. Differential expression analyses

Differential expression (DE) analyses were performed using the data set of one-to-one orthologs of each species. To estimate the expression levels, the paired reads were mapped to the species correspondent set of ortholog sequences using Bowtie 2.3.5.1 (Langmead and Salzberg, 2012) and further processed with RSEM 1.3.3 (Li and Dewey, 2011). The identification of differentially expressed orthologs between *L. limao* and all other stingless bee species was pair-wise performed using two methods, as

suggested by Ricardo et al. (2024): (1) using the TMM normalized counts (Robinson and Oshlack, 2010) compared in edgeR 3.34.0 (Robinson et al., 2010); and (2) using the scale based normalized (SCBN) counts (Zhou et al., 2019) and NOISeq 2.36.0 (Tarazona et al., 2015) for comparing expression levels. The SCBN method is a count normalization strategy optimized for cross-species DE analyses (Zhou et al., 2019) implemented in the Bioconductor package SCBN 1.10.0 (Zhou, 2021). Only significant differentially expressed transcripts (at a cut-off of $|Log2FC| \geq 2$, adjusted p-value < 0.05) identified using both methods were filtered as DE orthologs. Lastly, we compared all pairwise differential expression results to identify orthologs consistently differently expressed between *L. limao* and all pollen-collecting species.

2.5. *GO functional analysis*

Among the DE orthologs identified between *L. limao* and all the other species, we identified GO-enriched terms using the Bioconductor package TopGO 2.20.0 (Alexa and Rahnenfuhrer, 2020). Redundant enriched terms for Biological Process (BP) were summarized in a two-level hierarchical GO set using REVIGO web server (Supek et al., 2011) and these hierarchical sets were represented in a chart generated using the CirGO 2.0 software (Kuznetsova et al., 2019).

3. Results

3.1. *Transcriptome assembly and annotation*

The RNA sequencing generated 254,713,432 raw reads for *L. limao*, 261,049,812 for *N. testaceicornis*, and 259,553,460 for *S.* aff. *depilis*. The number of reads decreased to 182,096,230, 187,198,886, and 185,326.996, respectively, after cleaning, For *T. angustula*, 89,081,896 new raw reads were sequenced additionally to 118,318,606 cleaned reads from a previous study (Araujo and Arias, 2021), resulting in

a final set of 182,707,804 cleaned reads. The main quality parameters associated with each transcriptome assembly are summarized in Table 1.

Table 1. Main quality parameters for the transcriptome assemblies of *Lestrimelitta limao*, *Nannotrigona testaceicornis*, *Scaptotrigona* aff. *depilis*, and *Tetragonisca angustula*.

Metric	*L. limao*	*N. testaceicornis*	*S.* **aff.** *depilis*	*T. angustula*
Total transcripts assembled	97,316	127,496	137,488	152,927
Number of clustered SuperTranscripts	57,125	82,622	91,872	85,001
SuperTranscripts ≥ 1,000 nt	11,023	13,839	16,366	20,346
p fragments mapped[a]	0.92	0.91	0.92	0.91
p bases uncovered[b]	0.02	0.01	0.01	0.02
Transrate optimal score[c]	0.48	0.48	0.50	0.52
Complete BUSCOs (%)	88.6	89.2	90.6	92.6
Missing BUSCOs (%)	7.6	6.6	6.2	4.6

[a] proportion of read pairs that mapped successfully.
[b] proportion of bases that were not covered by any read.
[c] this discrete score represents the accuracy and completeness of the assembly based on the assessment of alignments between contigs and reads. The minimum score is 0, while the maximum is 1.0. For a description of the components used in the calculation of this metric, see Smith-Unna et al. (2016).

The number of transcripts assembled for *L. limao* was lower in comparison to the other species even though the amount of input RNA-seq reads was similar for all of them. Accordingly, after the annotation step and the exclusion of potential contaminants (Supplementary Material 1), *L. limao* was the species with the lowest number of SuperTranscripts returning significant blast hits with protein-coding gene sequences from the UniRef90 database, 24,859 (43,5% of total SuperTranscripts). For *N. testaceicornis*, *S.* aff. *depilis*, and *T. angustula* this numbers were 32,452 (39,2%), 38,273 (41,65%), and 33,596 (39,5%), respectively. Moreover, *L. limao* also presented a considerably lower number of potentially contaminating sequences (379) against *N. testaceicornis* (13,802), *S.* aff. *depilis* (14,414), and *T. angustula* (9,096). Most of the potential contaminants obtained for the non-cleptobiont bees, i.e., *N. testaceicornis*, *S.* aff. *depilis*, and *T. angustula*, resulted in significant blast hits against plant sequences (13,588, 14,033, and 8,467, respectively), particularly with plants of Fabaceae and Myrtaceae families, which is likely explained by their pollen-collecting behavior. *L. limao*, on the other hand, presented only 379 SuperTranscripts identified as potential

contaminant sequences, of which 87 were from plants, and the remaining were attributed mainly to fungi and bacteria.

3.2.   *Differential expression of orthologs*

Orthofinder identified 44,968 orthogroups, of which 8,213 were common to all four species (Figure 1A). *L. limao* showed the smallest number of species-specific orthogroups (140), and only 49 of them included at least one annotated transcript. We retrieved 7,482 one-to-one orthologs from all species. This set of orthologs was used for differential expression analyses. In the pair-wise comparisons, the most significant difference in expression was between *T. angustula* and *N. testaceicornis* (total of 1,229 DE orthologs), while the least amount of differentially expressed orthologs reported was between *S.* aff. *depilis* and *L. limao* (total of 734 DE orthologs). Finally, we identified 175 orthologs commonly differentially expressed between *L. limao* and all the other three bee species studied (Figure 1B, Supplementary Material 2), most of which (108) were consistently downregulated in *L. limao*. Among these, we find an odorant receptor (with significant BLAST hits to *Or1* from *Apis cerana* and LOC100741725 from *Bombus impatiens*), and several genes of the cytochrome P450 monooxygenase superfamily (*CYP* genes).

Figure 1. Analysis of the orthologs identified for the species *Lestrimelitta limao*, *Nannotrigona testaceicornis*, *Scaptotrigona* aff. *depilis* and *Tetragonisca angustula*. (A) Venn diagram representing the sets of orthogroups shared between/among species. (B) Venn diagram representing the sets of orthologs differentially expressed (DE) between/among species. Color scales represent count numbers ranging from lowest (dark gray) to highest (white).

The enrichment analysis of GO terms returned 26 enriched Biological Processes (BP) among the differentially expressed orthologs (Supplementary Material 3). The redundant BP terms were summarized in a two-level hierarchical list for visualization (Figure 2). The orthologs annotated to these enriched terms are shown in Table 2. Top enriched BP terms were related to *transmission of nerve impulse*, *maintenance of presynaptic active zone structure*, and *positive regulation of calcium ion transport*.

Figure 2. Summarized GO Biological Process terms enriched in the differentially expressed orthologs between the cleptobiotic *L. limao* and the pollen-collecting Meliponini species *Nannotrigona testaceicornis*, *Scaptotrigona* aff. *depilis* and *Tetragonisca angustula*. The legend next to the chart represents the color, name, and proportion of the GO parents for each two-level hierarchical set of child terms that were enriched.

Table 2. Differentially expressed orthologs between the cleptobiotic *L. limao* and the pollen-collecting Meliponini species *Nannotrigona testaceicornis*, *Scaptotrigona* aff. *depilis,* and *Tetragonisca angustula* annotated to the top enriched Biological Process GO term. Orthologs are represented by the ID of their respective orthogroups. Genes and respective species from the database hit are also shown (based on the Swiss-Prot database annotation) for each orthogroup. Expression patterns of orthologs are indicated by arrows: ↑ (upregulated) and ↓ (downregulated) as well as their function.

Orthogroup	Swiss-Prot homologs (organism)	Expression	Putative function
Peroxidase			
OG0001295	*Duox* (*Drosophila melanogaster*)	*L. limao* ↑ Pollen-collecting ↓	Production of reactive oxygen species (ROS) and potential role in defense against pathogens (Azzouz-Olden et al., 2018; Kawahara et al., 2007).
Peptidase			
OG0003146	*PAN2* (*D. melanogaster*)	*L. limao* ↑ Pollen-collecting ↓	Involved in mRNA degradation, and in hencing the regulation of transcript abundance (Parker and Song, 2004; Wahle

and Winkler, 2013).

Uracil-DNA glicosilase			
OG0010914	*TDG* (*Homo sapiens*)	*L. limao* ↑ Pollen-collecting ↓	Repair of damaged DNA by removing thymines and uracils that have been mispaired with guanines (Hardeland et al., 2001).
DNA helicase			
OG0009618	*MCM3* (*H. sapiens, Pongo abelli*)	*L. limao* ↑ Pollen-collecting ↓	Essential in the initiation and elongation phases of chromosome replication (helicase function) (Li et al., 2011).
Sodium channel			
OG0001240	*para* (*D. melanogaster*)	*L. limao* ↓ Pollen-collecting ↑	Postsynaptic excitatory potential regulation; Potential role in mechanosensory behavior (Engel and Wu, 1994; Xiao et al., 2017).
Distrofin			
OG0003765	*Dys* (*D. melanogaster*)	*L. limao* ↓ Pollen-collecting ↑	Control of neuromuscular synaptic homeostasis (van der Plas et al., 2006); modulation of the microRNA profile (Marrone et al., 2012).
Glucose-6-phosphate dehydrogenase			
OG0000574	*G6PD/Zw* (*H. sapiens, D. melanogaster*)	*L. limao* ↓ Pollen-collecting ↑	Role in pentose phosphate pathway and redox balance (Legan et al., 2008; Stincone et al., 2015).
Serine-threonine phosphatase			
OG0000759	*Ppp3ca/Pp2B-14D* (*Rattus norvegicus, H.* sapiens, *Bos taurus, D. melanogaster*)	*L. limao* ↓ Pollen-collecting ↑	Regulates diverse processes like immunity, sleep, circadian rhythms, and female meiosis (Dijkers and O'Farrell, 2007; Kweon et al., 2018; Nakai et al., 2011; Takeo et al., 2012).
ATPase			
OG0000773	*ATAD3A/bor* (*H. sapiens,* *Xenopus tropicalis,* *D. melanogaster*)	*L. limao* ↓ Pollen-collecting ↑	Regulation of the activity and morphology of mitochondria (Gilquin et al., 2010; Harel et al., 2016).
RING-box protein			
OG0000879	*rbx1/RNF7*/Roc1a (*Salmo salar, Mus musculus, H. sapiens, D. melanogaster,*)	*L. limao* ↓ Pollen-collecting ↑	Involved in protein degradation via ubiquitination. Roles in *Drosophila* development and neural remodeling (Arama et al., 2007; Noureddine et al., 2002; Wong et al., 2013).
Transcriptional elongation factor			
OG0000899	*ELP4* (*H. sapiens, Xenopus laevis, Danio rerio*)	*L. limao* ↓ Pollen-collecting ↑	Implicated in several processes including transcriptional control, α-tubulin acetylation, and tRNA modifications; related to translational efficiency and

			fidelity (Dalwadi and Yip, 2018; Johansson et al., 2018).
OG0002245	*Iws1* (*M. musculus*, *R. norvegicus*)	*L. limao* ↓ Pollen-collecting ↑	Involved in the elongation, processing, and nuclear exportation of mRNA; important for ensuring the efficient translation of mRNA into protein (Brès et al., 2008; Pujari et al., 2010; Yoh et al., 2007).
Type IV collagen protein			
OG0001022	*COL4A1/COL4A2/emb-9/Col4a1* (*H. sapiens*, *B. taurus*, *Caenorhabditis elegans*, *D. melanogaster*)	*L. limao* ↓ Pollen-collecting ↑	Encodes collagen IV chains, crucial components of the basement membrane (Morrissey and Sherwood, 2015). Role in the adequate development of the structure and function of cardiac, skeletal, and visceral muscle cells during morphogenesis (Kelemen-Valkony et al., 2012; Kiss et al., 2016).
DNA-binding protein			
OG0001379	*SMARCD1/Bap60* (*H. sapiens*, *D. melanogaster*)	*L. limao* ↓ Pollen-collecting ↑	*Bap60*/*SMARCD1* homologs mediate interactions between the Brahma complex and species-specific transcription factors, implicating them in a variety of regulatory processes, e.g., immune response to microbial infections, and neurodevelopmental genes regulation (Liu et al., 2016; Nixon et al., 2019).
Carboxylase			
OG0002309	*PCK2/Pepck1* (*H. sapiens*, *D. melanogaster*)	*L. limao* ↓ Pollen-collecting ↑	Encodes an enzyme crucial for gluconeogenesis (Hanson and Patel, 2006).

4. Discussion

Our comparisons revealed that the cleptobiotic species exhibits lower expression levels for most differentially expressed orthologs. Among these downregulated orthologs, we highlight three groups associated with genes encoding for proteins that might play a role in the ecological differences among *L. limao* and non-robber bees. The first group comprises orthologs involved in detoxification and chemoreception pathways, including genes homologous to the cytochrome P450 monooxygenase superfamily and a putative odorant receptor (*Or1*). We hypothesize that a reduced expression of these genes could be associated with non-floral foraging lifestyle adaptations. The second group contains orthologs from the top enriched GO terms, *i.e.*,

transmission of nerve impulses, maintenance of presynaptic active zone structure, and positive regulation of calcium ion transport. Although the exact functional consequences of these processes in cleptobiosis are unclear, the critical role of these genes in synaptic transmission, neuromuscular function, and sensory processing suggests they may be associated with neurophysiological adaptations of cleptoparasitism. Finally, the third group of orthologs is involved in mitochondrial activity and carbohydrate metabolism. We hypothesize that the reduced expression of these genes in *L. limao* workers suggests a shift in metabolic pathways leading to a low-energy-demanding worker state compared to pollen-collecting bees.

4.1. *Group I of downregulated orthologs: detoxification and chemoreception*

The cytochrome P450 superfamily (CYP genes) is widely conserved across living organisms (Werck-Reichhart and Feyereisen, 2000). Genes in this superfamily encode enzymes that catalyze a broad range of oxidative processes, including both endogenous and xenobiotic detoxification (Feyereisen, 2012). In arthropods, six major CYP clans have been identified, with the most prominent being the CYP2, CYP3, CYP4, and mitochondrial clans (Dermauw et al., 2020). Our results indicate that *L. limao* exhibits a reduced expression of orthologs from the CYP2 (*Cyp304a1*) and CYP3 (*Cyp6a14* and *Cyp6a2*) clans. The CYP2 clan is evolutionarily conserved and participates in key biological pathways, such as juvenile hormone and ecdysteroid biosynthesis (Bassett et al., 1997; Li et al., 2014; Namiki et al., 2005; Rewitz et al., 2006), as well as in the development of sensory structures (Willingham and Keil, 2004). The CYP3 clan is the most diverse among arthropods (Dermauw et al., 2020) and includes the CYP6 family, which comprises *Cyp6a14*, *Cyp6a2*, and several other genes involved in the metabolism of xenobiotics and phytochemicals (Feyereisen, 2012). Functional studies have demonstrated that *Cyp6a14* variants in *Tribolium castaneum*

enhance detoxification capacity against plant-derived toxins (Ercan et al., 2020), while in *Aphis glycines*, increased expression of a *CYP6A14*-like gene confers resistance to pyrethroid insecticides (Xi et al., 2015). Similarly, *Cyp6a2* has been linked to resistance against organophosphates (Dunkov et al., 1997) and DDT (Pedra et al., 2004) in *Drosophila*.

In bees, Johnson et al. (2018) found evidence of positive selection in the genes of the CYP6AS subfamily (part of the CYP6A family, CYP3 clan). Based on these findings, the authors hypothesized that the role of these genes in metabolizing and detoxifying phytochemicals present in plant nectar, pollen, and resins (compounds that are toxic if not degraded) has enabled honeybees to consume a flavonoid-rich diet (Johnson et al., 2018; Liu et al., 2005; Mao et al., 2017). Since *L. limao* does not directly harvest floral resources but instead obtains them already processed through nest robbing (Michener, 1946; Shoichi F Sakagami and Laroca, 1963),— a behavior corroborated by the significantly lower number of plant-associated contaminant sequences in *L. limao* compared to pollen-collecting species in our dataset —we hypothesize that the downregulation of these CYP genes may reflect reduced exposure of this species to phytochemicals, as this behavioral shift likely decreases the need for flower detoxification mechanisms.

Another example of a downregulated ortholog in *L. limao*, potentially linked to the loss of floral foraging, comes from a putative odorant receptor (significant blast hits against *Or1* from *Apis cerana* and LOC100741725 from *Bombus impatiens*). *AcerOr1* has been implicated in detecting volatile floral scents (Zhao et al., 2014), and its downregulation agrees with a non-floral foraging lifestyle as *L. limao* workers likely do not rely on floral scents to obtain food. Still, the odorant receptor (OR) gene family is large and highly diverse in Hymenoptera (Obiero et al., 2021). Thus, it is worth

mentioning that the identified orthologs may not have retained their functions across species. Chemosensory-related genes, including ORs, likely play crucial roles in intraspecific communication during nest-robbing events (Sakagami et al., 1993; von Zuben et al., 2016) and the specific cues guiding *L. limao* workers in robbing nests of diverse hosts remain unknown, as well as the impact of cleptobiotic behavior on the evolution of these gene families.

4.2. *Group II of downregulated orthologs: synaptic transmission and neurosensory processing.*

Among the orthologs associated with the top enriched GO terms (*i.e.*, transmission of nerve impulses, maintenance of presynaptic active zone structure, and positive regulation of calcium ion transport) we found the genes *para* and *Dys*. In *Drosophila*, *para* encodes voltage-activated sodium channels (Loughney et al., 1989), which are essential for regulating postsynaptic excitatory potential (Xiao et al., 2017) and are widely expressed in the central nervous system of adult flies (Ravenscroft et al., 2020). Alternative splicing in this gene leads to channel variants of distinct properties (Lin et al., 2012, 2009; O'Dowd and Smith, 1996; Thackeray and Ganetzky, 1995), and mutations in *para* have been linked to seizure-like behavior, paralysis, and altered mechanosensory responses (Engel and Wu, 1994; Parker et al., 2011; Takai et al., 2020). The gene *Dys* encodes the protein Dystrophin, a conserved component of the dystrophin-glycoprotein complex (DGC) (Greener and Roberts, 2000; Neuman et al., 2005). In *Drosophila*, *Dys* is involved in neuromuscular synaptic homeostasis (Jantrapirom et al., 2019; van der Plas et al., 2006), photoreceptor axon migration (Marrone et al., 2011; Shcherbata et al., 2007; Zhan et al., 2010), and microRNA profile modulation (Marrone et al., 2012). The loss of functional *Dys* has been associated with age-dependent muscle degeneration and movement impairments (Shcherbata et al.,

2007). The critical roles of *para* and *Dys* in synaptic transmission, neuromuscular function, and sensory processing suggest that these orthologs may contribute to neurophysiological adaptations associated with cleptoparasitism. However, their precise functional consequences remain unclear. Future studies integrating neurophysiological and behavioral analyses will be necessary to elucidate the impact of these transcriptional differences in robber bees.

4.3. Group III of downregulated orthologs: mitochondrial activity and carbohydrate metabolism.

Processes associated with mitochondrial activity and carbohydrate metabolism were also enriched in our GO term analysis and include downregulated orthologs in *L. limao* homologs to the vertebrate *ATAD3A*, to the human *PCK2*, and the *Drosophila bor*, *Pepck1*, *G6PD*, and *Zw*. *ATAD3A* plays a critical role in mitochondrial dynamics, particularly at the interface of the inner and outer mitochondrial membranes, where it regulates essential processes such as mitochondrial fusion and mitochondrial DNA organization (Gilquin et al., 2010; Harel et al., 2016; He et al., 2007; Hoffmann et al., 2009; Wang and Bogenhagen, 2006). Overexpression of *ATAD3A* enhances mitochondrial ATP production efficiency, a key driver of cellular metabolic activity (Gomes et al., 2011; Harel et al., 2016). *PCK2/Pepck1*, in turn, encodes an enzyme essential for gluconeogenesis (Hanson and Patel, 2006), playing a key role in glucose homeostasis in insects and initiating trehalose synthesis (Miyamoto and Amrein, 2017). Trehalose, the main sugar in insect hemolymph, is a crucial energy source for ATP production during flight (Becker et al., 1996; Matsuda et al., 2015). *G6PD* encodes glucose-6-phosphate 1-dehydrogenase, an enzyme critical for carbohydrate metabolism (reviewed in Stincone et al. 2015) and cellular redox balance (Kletzien et al., 1994; Legan et al., 2008). In this regard, it has been proposed that PPP may reduce the

oxidative stress brought on by flying activity, particularly in species whose diets are rich in sugars, as is the case of insects that feed on nectar (Levin et al., 2017).

Given that these orthologs are linked to energic metabolism and protection against oxidative damage and that their expression levels are reduced in *L. limao* compared to non-robber bees, we conclude that the *L. limao* workers sampled in this study were not engaged in high-energy-demanding activities. We took special care to sample *L. limao* workers when returning to their colonies, this choice was based on the behavior of foragers from pollen-collecting species and ensured the sampling of workers at equivalent life stages across species for transcriptomic comparisons. However, *L. limao* workers are also known to raid other stingless bee colonies, and we did not sample individuals engaged in this activity. We hypothesize the expression pattern of workers may shift significantly during robbing raids since the metabolic demand should increase as worker activity intensifies. Accordingly, two interpretations are possible to explain our findings: either *L. limao* workers have a reduced global energetic demand compared to other stingless bees, or an alternative foraging state exists in this species. In this case, foraging workers would alternately engage in two distinct activities, robbing raids (a high-energy-demanding activity) and low-energy-demanding tasks, such as short-distance recognition flights. Further investigation comparing these two states is required to confirm these hypotheses.

5. Conclusions

Lestrimelitta species constitute one of the most notable and well-known examples of obligate cleptobiosis. Despite this, there is still a large gap in knowledge about the genetic factors underlying this behavior. Herein, we provide a starting point for these investigations. Our results demonstrated that most DE orthologs are downregulated in *L. limao* workers, and some of them may reflect specific aspects of

obligate cleptobiosis in bees, such as the downregulation of *CYP* genes, which is associated with the non-floral foraging lifestyle. The enriched GO terms indicate that several DE orthologs potentially regulate neuronal and synaptic activities in addition to other important cellular processes. Moreover, we found evidence that robber foragers have reduced energetic demands when compared to other eusocial bee workers, which could suggest the existence of two foraging states in *L. limao,* one more active and another less active. In summary, the distinctive expression pattern of genes in *L. limao* provides valuable insights into the molecular mechanisms associated with its cleptobiotic behavior during foraging, offering avenues for further investigation into the adaptation and ecological dynamics of this unique species.

Credit authorship contribution statement

Paulo C. Ricardo: Writing – review & editing, Writing – original draft, Conceptualization, Methodology, Sampling, Investigation, Formal analysis. **Natalia de S. Araujo:** Writing – review & editing, Conceptualization, Methodology, Supervision, Funding acquisition. **Maria C. Arias:** Writing – review & editing, Conceptualization, Supervision, Funding acquisition, Project administration.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The authors acknowledge Susy Coelho Oliveira and Thiago Geronimo Pires Alegria for technical assistance, and Isabel Alves dos-Santos, Sheina Koffler, Larissa Nunes do Prado and Sergio Dias Hilário for support in specimen sampling.

Funding

This study was supported in part by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – CAPES, Brasil [scholarship to PCR, Proc. 1783961 and Proc. 88882.377416/2019-01, Finance Code 001]; Conselho Nacional de Desenvolvimento Científico e Tecnológico –CNPq, Brasil [research sponsorship to MCA - 306932/2016-4], and Fundação de Amparo à Pesquisa do Estado de São Paulo – FAPESP [Proc. 2013/12530-4, 2016/24669-5, 2019/23186-9 and 2022/09046-2]. NSA was funded by the Fonds De La Recherche Scientifique (FNRS) [grant number 40005980].

Data availability

The raw sequence reads have been deposited in the NCBI Sequence Read Archive (SRA) under the following accession numbers: SRR29109009, SRR29109008, SRR29109007, SRR29109006, SRR29109005, SRR29109004, SRR29109003, SRR29109002, SRR29109001, SRR29109000. The corresponding BioProject accession number is PRJNA1113344. The transcriptome assemblies, as well as the sets of SuperTranscripts, are available on FigShare (https://doi.org/10.6084/m9.figshare.26064586.v1).

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work the authors used ChatGPT (GPT-4o mini) and DeepSeek (DeepThink R1) in order to improve the clarity and coherence of the text and to assist in language editing and grammar correction. After using these tools, the authors reviewed and edited the content as needed, taking full responsibility for the content of the published article.

References

Alexa, A., Rahnenfuhrer, J., 2020. topGO: Enrichment Analysis for Gene Ontology. R package version 2.20.0.

Altschul, S., 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. Nucleic Acids Res 25, 3389–3402. https://doi.org/10.1093/nar/25.17.3389

Andrews, S., 2018. FastQC: a quality control tool for high throughput sequence data.

Arama, E., Bader, M., Rieckhof, G.E., Steller, H., 2007. A Ubiquitin Ligase Complex Regulates Caspase Activation During Sperm Differentiation in *Drosophila*. PLoS Biol 5, e251. https://doi.org/10.1371/journal.pbio.0050251

Araujo, N.S., Arias, M.C., 2021. Gene expression and epigenetics reveal species-specific mechanisms acting upon common molecular pathways in the evolution of task division in bees. Sci Rep 11, 3654. https://doi.org/10.1038/s41598-020-75432-8

Azzouz-Olden, F., Hunt, A., DeGrandi-Hoffman, G., 2018. Transcriptional response of honey bee (*Apis mellifera*) to differential nutritional status and *Nosema* infection. BMC Genomics 19, 628. https://doi.org/10.1186/s12864-018-5007-0

Bairoch, A., 1996. The SWISS-PROT protein sequence data bank and its new supplement TREMBL. Nucleic Acids Res 24, 21–25. https://doi.org/10.1093/nar/24.1.21

Bassett, M.H., McCarthy, J.L., Waterman, M.R., Sliter, T.J., 1997. Sequence and developmental expression of Cyp18, a member of a new cytochrome P450 family from *Drosophila*. Mol Cell Endocrinol 131, 39–49. https://doi.org/10.1016/S0303-7207(97)00093-2

Becker, A., Schlöder, P., Steele, J.E., Wegener, G., 1996. The regulation of trehalose metabolism in insects. Experientia 52, 433–439. https://doi.org/10.1007/BF01919312

Blum, M.S., 1966. Chemical Releasers of Social Behavior. VIII. Citral in the Mandibular Gland Secretion of Lestrimelitta limao (Hymenoptera: Apoidea: Melittidae)1. Ann Entomol Soc Am 59, 962–964. https://doi.org/10.1093/aesa/59.5.962

Blum, M.S., Crewe, R., Kerr, L.H., Keith, L.H., Garrison, A.W., Walker, M.M., 1970. Citral in stingless bees: Isolation and functions in trail- laying and robbing. J Insect Physiol 16, 1637–1648.

Bohart, G.E., 1970. The Evolution of Parasitism among bees, 41 st. ed, Faculty Honor Lectures. Utah State University.

Bolger, A.M., Lohse, M., Usadel, B., 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics 30, 2114–2120. https://doi.org/10.1093/bioinformatics/btu170

Breed, M.D., Cook, C., Krasnec, M.O., 2012. Cleptobiosis in social insects. Psyche (Camb Mass) 2012, 1–7. https://doi.org/10.1155/2012/484765

Brès, V., Yoh, S.M., Jones, K.A., 2008. The multi-tasking P-TEFb complex. Curr Opin Cell Biol 20, 334–340. https://doi.org/10.1016/j.ceb.2008.04.008

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., Madden, T.L., 2009. BLAST+: architecture and applications. BMC Bioinformatics 10, 421. https://doi.org/10.1186/1471-2105-10-421

Camargo, J.M.F., Pedro, S.R.M., 2007. Notas sobre a bionomia de *Trichotrigona extranea* Camargo & Moure (Hymenoptera, Apidae, Meliponini). Rev Bras Entomol 51, 72–81. https://doi.org/10.1590/S0085-56262007000100013

Costa, L., Galaschi-Teixeira, J.S., Maia, U.M., Imperatriz-Fonseca, V.L., 2018. Plasticity of stingless bee *Melipona fuliginosa* Lepeletier to obtain food resources in Amazonia. Sociobiology 65, 744. https://doi.org/10.13102/sociobiology.v65i4.3490

Dalwadi, U., Yip, C.K., 2018. Structural insights into the function of Elongator. Cellular and Molecular Life Sciences 75, 1613–1622. https://doi.org/10.1007/s00018-018-2747-6

Davidson, N.M., Hawkins, A.D.K., Oshlack, A., 2017. SuperTranscripts: a data driven reference for analysis and visualisation of transcriptomes. Genome Biol 18, 148. https://doi.org/10.1186/s13059-017-1284-1

Dermauw, W., Van Leeuwen, T., Feyereisen, R., 2020. Diversity and evolution of the P450 family in arthropods. Insect Biochem Mol Biol 127, 103490. https://doi.org/10.1016/j.ibmb.2020.103490

Dijkers, P.F., O'Farrell, P.H., 2007. *Drosophila* Calcineurin Promotes Induction of Innate Immune Responses. Current Biology 17, 2087–2093. https://doi.org/10.1016/j.cub.2007.11.001

Dunkov, B.C., Guzov, V.M., Mocelin, G., Shotkoski, F., Brun, A., Amichot, M., Ffrench-Constant, R.H., Feyereisen, R., 1997. The *Drosophila* Cytochrome P450 Gene *Cyp6a2*□: Structure, Localization, Heterologous Expression, and Induction by Phenobarbital. DNA Cell Biol 16, 1345–1356. https://doi.org/10.1089/dna.1997.16.1345

Eardley, C.D., 2004. Taxonomic revision of the African stingless bees (Apoidea: Apidae: Apinae: Meliponini). Afr Plant Prot 10, 63–96.

Emms, D.M., Kelly, S., 2019. OrthoFinder: phylogenetic orthology inference for comparative genomics. Genome Biol 20, 238. https://doi.org/10.1186/s13059-019-1832-y

Emms, D.M., Kelly, S., 2015. OrthoFinder: solving fundamental biases in whole genome comparisons dramatically improves orthogroup inference accuracy. Genome Biol 16, 157. https://doi.org/10.1186/s13059-015-0721-2

Engel, J.E., Wu, C.-F., 1994. Altered mechanoreceptor response in *Drosophila* bang-sensitive mutants. Journal of Comparative Physiology A 175, 267–278. https://doi.org/10.1007/BF00192986

Engel, M.S., Rasmussen, C., Ayala, R., de Oliveira, F.F., 2023. Stingless bee classification and biology (Hymenoptera, Apidae): a review, with an updated key to genera and subgenera. Zookeys 1172, 239–312. https://doi.org/10.3897/zookeys.1172.104944

Ercan, F.S., Azarkan, S.Y., Ercan, N., Koc, M., 2020. Sequence variants of *CYP345a1* and *CYP6a14<i/> gene regions in Tribolium castaneum (Coleoptera: Tenebrionidae) adults treated with the novel characterized Bolanthus turcicus (Caryophyllaceae) extract. Mol Biol Res Commun 9, 105–110. https://doi.org/10.22099/mbrc.2020.35861.1472*

Feyereisen, R., 2012. Insect CYP Genes and P450 Enzymes, in: Insect Molecular Biology and Biochemistry. Elsevier, pp. 236–316. https://doi.org/10.1016/B978-0-12-384747-8.10008-X

Forel, A., 1901. Fourmis termitophages, lestobiose, Atta tardigrada, sous-genres d'Euponera. Annales de la Société Entomologique de Belgique 45, 389–398.

Francke, W., Lübke, G., Schröder, W., Reckziegel, A., Imperatriz-Fonseca, V., Kleinert, A., Engels, E., Hartfelder, K., Radtke, R., Engels, W., 2000. Identification of oxygen containing volatiles in cephalic secretions of workers of Brazilian stingless bees. J Braz Chem Soc 11, 562–571. https://doi.org/10.1590/S0103-50532000000600003

Gilquin, B., Taillebourg, E., Cherradi, N., Hubstenberger, A., Gay, O., Merle, N., Assard, N., Fauvarque, M.-O., Tomohiro, S., Kuge, O., Baudier, J., 2010. The AAA⁺ ATPase ATAD3A Controls Mitochondrial Dynamics at the Interface of the Inner and Outer Membranes. Mol Cell Biol 30, 1984–1996. https://doi.org/10.1128/MCB.00007-10

Gomes, L.C., Benedetto, G. Di, Scorrano, L., 2011. During autophagy mitochondria elongate, are spared from degradation and sustain cell viability. Nat Cell Biol 13, 589–598. https://doi.org/10.1038/ncb2220

Grabherr, M.G., Haas, B.J., Yassour, M., Levin, J.Z., Thompson, D.A., Amit, I., Adiconis, X., Fan, L., Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, E., Hacohen, N., Gnirke, A., Rhind, N., di Palma, F., Birren, B.W., Nusbaum, C., Lindblad-Toh, K., Friedman, N., Regev, A., 2013. Trinity: reconstructing a full-length transcriptome assembly from RNA-Seq data without a reference genome. Nat Biotechnol 29, 644–652. https://doi.org/10.1038/nbt.1883

Greener, M.J., Roberts, R.G., 2000. Conservation of components of the dystrophin complex in Drosophila [1]. FEBS Lett 482, 13–18. https://doi.org/10.1016/S0014-5793(00)02018-4

Grüter, C., 2020. Colony Organisation and Division of Labour. pp. 203–232. https://doi.org/10.1007/978-3-030-60090-7_6

Grüter, C., Menezes, C., Imperatriz-Fonseca, V.L., Ratnieks, F.L.W., 2012. A morphologically specialized soldier caste improves colony defense in a neotropical

eusocial bee. Proceedings of the National Academy of Sciences 109, 1182–1186. https://doi.org/10.1073/pnas.1113398109

Grüter, C., Segers, F.H.I.D., Menezes, C., Vollet-Neto, A., Falcón, T., von Zuben, L., Bitondi, M.M.G., Nascimento, F.S., Almeida, E.A.B., 2017. Repeated evolution of soldier sub-castes suggests parasitism drives social complexity in stingless bees. Nat Commun 8, 4. https://doi.org/10.1038/s41467-016-0012-y

Grüter, C., von Zuben, L.G., Segers, F.H.I.D., Cunningham, J.P., 2016. Warfare in stingless bees. Insectes Soc 63, 223–236. https://doi.org/10.1007/s00040-016-0468-0

Haas, B.J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P.D., Bowden, J., Couger, M.B., Eccles, D., Li, B., Lieber, M., MacManes, M.D., Ott, M., Orvis, J., Pochet, N., Strozzi, F., Weeks, N., Westerman, R., William, T., Dewey, C.N., Henschel, R., LeDuc, R.D., Friedman, N., Regev, A., 2013. De novo transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. Nat Protoc 8, 1494–1512. https://doi.org/10.1038/nprot.2013.084

Hanson, R.W., Patel, Y.M., 2006. Phosphoenolpyruvate Carboxykinase (GTP): the Gene and the Enzyme. pp. 203–281. https://doi.org/10.1002/9780470123157.ch6

Hardeland, U., Bentele, M., Lettieri, T., Steinacher, R., Jiricny, J., Schär, P., 2001. Thymine DNA Glycosylase. Prog Nucleic Acid Res Mol Biol 68, 235–253. https://doi.org/10.1016/S0079-6603(01)68103-0

Harel, T., Yoon, W.H., Garone, C., Gu, S., Coban-Akdemir, Z., Eldomery, M.K., Posey, J.E., Jhangiani, S.N., Rosenfeld, J.A., Cho, M.T., Fox, S., Withers, M., Brooks, S.M., Chiang, T., Duraine, L., Erdin, S., Yuan, B., Shao, Y., Moussallem, E., Lamperti, C., Donati, M.A., Smith, J.D., McLaughlin, H.M., Eng, C.M., Walkiewicz, M., Xia, F., Pippucci, T., Magini, P., Seri, M., Zeviani, M., Hirano, M., Hunter, J.V., Srour, M., Zanigni, S., Lewis, R.A., Muzny, D.M., Lotze, T.E., Boerwinkle, E., Gibbs, R.A., Hickey, S.E., Graham, B.H., Yang, Y., Buhas, D., Martin, D.M., Potocki, L., Graziano, C., Bellen, H.J., Lupski, J.R., 2016. Recurrent De Novo and Biallelic Variation of ATAD3A , Encoding a Mitochondrial Membrane Protein, Results in Distinct Neurological Syndromes. The American Journal of Human Genetics 99, 831–845. https://doi.org/10.1016/j.ajhg.2016.08.007

He, J., Mao, C.-C., Reyes, A., Sembongi, H., Di Re, M., Granycome, C., Clippingdale, A.B., Fearnley, I.M., Harbour, M., Robinson, A.J., Reichelt, S., Spelbrink, J.N., Walker, J.E., Holt, I.J., 2007. The AAA$^+$ protein ATAD3 has displacement loop binding properties and is involved in mitochondrial nucleoid organization. Journal of Cell Biology 176, 141–146. https://doi.org/10.1083/jcb.200609158

Hoffmann, M., Bellance, N., Rossignol, R., Koopman, W.J.H., Willems, P.H.G.M., Mayatepek, E., Bossinger, O., Distelmaier, F., 2009. C. elegans ATAD-3 Is Essential for Mitochondrial Activity and Development. PLoS One 4, e7644. https://doi.org/10.1371/journal.pone.0007644

Hölldobler, B., 1986. *Food robbing in ants, a form of interference competition.* Oecologia 69, 12–15. https://doi.org/10.1007/BF00399031

IYENGAR, E. v., 2008. *Kleptoparasitic interactions throughout the animal kingdom and a re-evaluation, based on participant mobility, of the conditions promoting the evolution of kleptoparasitism.* Biological Journal of the Linnean Society 93, 745–762. https://doi.org/10.1111/j.1095-8312.2008.00954.x

James, C.C., Sánchez, D., Cruz-López, L., Nieh, J., 2022. *Fighting ability and the toxicity of raiding pheromone in an obligate kleptoparasite, the stingless bee Lestrimelitta niitkib.* Behav Ecol Sociobiol 76, 38. https://doi.org/10.1007/s00265-022-03129-1

Jantrapirom, S., Nimlamool, W., Temviriyanukul, P., Ahmadian, S., Locke, C.J., Davis, G.W., Yamaguchi, M., Noordermeer, J.N., Fradkin, L.G., Potikanond, S., 2019. *Dystrobrevin is required postsynaptically for homeostatic potentiation at the Drosophila NMJ.* Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease 1865, 1579–1591. https://doi.org/10.1016/j.bbadis.2019.03.008

Johansson, M.J.O., Xu, F., Byström, A.S., 2018. *Elongator—a tRNA modifying complex that promotes efficient translational decoding.* Biochimica et Biophysica Acta (BBA) - Gene Regulatory Mechanisms 1861, 401–408. https://doi.org/10.1016/j.bbagrm.2017.11.006

Johnson, R.M., Harpur, B.A., Dogantzis, K.A., Zayed, A., Berenbaum, M.R., 2018. *Genomic footprint of evolution of eusociality in bees: floral food use and CYPome "blooms."* Insectes Soc 65, 445–454. https://doi.org/10.1007/s00040-018-0631-x

Kawahara, T., Quinn, M.T., Lambeth, J.D., 2007. *Molecular evolution of the reactive oxygen-generating NADPH oxidase (Nox/Duox) family of enzymes.* BMC Evol Biol 7, 109. https://doi.org/10.1186/1471-2148-7-109

Kelemen-Valkony, I., Kiss, M., Csiha, J., Kiss, A., Bircher, U., Szidonya, J., Maróy, P., Juhász, G., Komonyi, O., Csiszár, K., Mink, M., 2012. *Drosophila basement membrane collagen col4a1 mutations cause severe myopathy.* Matrix Biology 31, 29–37. https://doi.org/10.1016/j.matbio.2011.09.004

Kiss, M., Kiss, A.A., Radics, M., Popovics, N., Hermesz, E., Csiszár, K., Mink, M., 2016. *Drosophila type IV collagen mutation associates with immune system activation and intestinal dysfunction.* Matrix Biology 49, 120–131. https://doi.org/10.1016/j.matbio.2015.09.002

Kletzien, R.F., Harris, P.K.W., Foellmi, L.A., 1994. *Glucose□6□phosphate dehydrogenase: a "housekeeping" enzyme subject to tissue□specific regulation by hormones, nutrients, and oxidant stress.* The FASEB Journal 8, 174–181. https://doi.org/10.1096/fasebj.8.2.8119488

Kuznetsova, I., Lugmayr, A., Siira, S.J., Rackham, O., Filipovska, A., 2019. *CirGO: an alternative circular way of visualising gene ontology terms.* BMC Bioinformatics 20, 84. https://doi.org/10.1186/s12859-019-2671-2

Kweon, S.H., Lee, J., Lim, C., Choe, J., 2018. High-Amplitude Circadian Rhythms in Drosophila Driven by Calcineurin-Mediated Post-translational Control of sarah. Genetics 209, 815–828. https://doi.org/10.1534/genetics.118.300808

Langmead, B., Salzberg, S.L., 2012. Fast gapped-read alignment with Bowtie 2. Nat Methods 9, 357–359. https://doi.org/10.1038/nmeth.1923

Legan, S.K., Rebrin, I., Mockett, R.J., Radyuk, S.N., Klichko, V.I., Sohal, R.S., Orr, W.C., 2008. Overexpression of Glucose-6-phosphate Dehydrogenase Extends the Life Span of Drosophila melanogaster. Journal of Biological Chemistry 283, 32492–32499. https://doi.org/10.1074/jbc.M805832200

Lepeco, A., Branstetter, M.G., Melo, G.A.R., Freitas, F. V., Tobin, K.B., Gan, J., Jensen, J., Almeida, E.A.B., 2024. Phylogenomic insights into the worldwide evolutionary relationships of the stingless bees (Apidae, Meliponini). Mol Phylogenet Evol 201, 108219. https://doi.org/10.1016/j.ympev.2024.108219

Levin, E., Lopez-Martinez, G., Fane, B., Davidowitz, G., 2017. Hawkmoths use nectar sugar to reduce oxidative damage from flight. Science (1979) 355, 733–735. https://doi.org/10.1126/science.aah4634

Li, B., Dewey, C.N., 2011. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. BMC Bioinformatics 12, 323–333. https://doi.org/10.1186/1471-2105-12-323

Li, J., Deng, M., Wei, Q., Liu, T., Tong, X., Ye, X., 2011. Phosphorylation of MCM3 Protein by Cyclin E/Cyclin-dependent Kinase 2 (Cdk2) Regulates Its Function in Cell Cycle. Journal of Biological Chemistry 286, 39776–39785. https://doi.org/10.1074/jbc.M111.226464

Li, Z., Ge, X., Ling, L., Zeng, B., Xu, J., Aslam, A.F.M., You, L., Palli, S.R., Huang, Y., Tan, A., 2014. CYP18A1 regulates tissue-specific steroid hormone inactivation in Bombyx mori. Insect Biochem Mol Biol 54, 33–41. https://doi.org/10.1016/j.ibmb.2014.08.007

Lin, W.-H., Günay, C., Marley, R., Prinz, A.A., Baines, R.A., 2012. Activity-Dependent Alternative Splicing Increases Persistent Sodium Current and Promotes Seizure. The Journal of Neuroscience 32, 7267–7277. https://doi.org/10.1523/JNEUROSCI.6042-11.2012

Lin, W.-H., Wright, D.E., Muraro, N.I., Baines, R.A., 2009. Alternative Splicing in the Voltage-Gated Sodium Channel DmNa $_v$ Regulates Activation, Inactivation, and Persistent Current. J Neurophysiol 102, 1994–2006. https://doi.org/10.1152/jn.00613.2009

Liu, F., He, J., Fu, W., 2005. Highly controlled nest homeostasis of honey bees helps deactivate phenolics in nectar. Naturwissenschaften 92, 297–299. https://doi.org/10.1007/s00114-005-0629-x

Liu, N., Wang, X.-W., Sun, J.-J., Wang, L., Zhang, H.-W., Zhao, X.-F., Wang, J.-X., 2016. Akirin interacts with Bap60 and 14-3-3 proteins to regulate the expression of

antimicrobial peptides in the kuruma shrimp (Marsupenaeus japonicus). Dev Comp Immunol 55, 80–89. https://doi.org/10.1016/j.dci.2015.10.015

Loughney, K., Kreber, R., Ganetzky, B., 1989. Molecular analysis of the para locus, a sodium channel gene in Drosophila. Cell 58, 1143–1154. https://doi.org/10.1016/0092-8674(89)90512-6

Mao, W., Schuler, M.A., Berenbaum, M.R., 2017. Disruption of quercetin metabolism by fungicide affects energy production in honey bees (Apis mellifera). Proceedings of the National Academy of Sciences 114, 2538–2543. https://doi.org/10.1073/pnas.1614864114

Marrone, A.K., Edeleva, E. V, Kucherenko, M.M., Hsiao, N.-H., Shcherbata, H.R., 2012. Dg-Dys-Syn1 signaling in Drosophila regulates the microRNA profile. BMC Cell Biol 13, 26. https://doi.org/10.1186/1471-2121-13-26

Marrone, A.K., Kucherenko, M.M., Rishko, V.M., Shcherbata, H.R., 2011. New Dystrophin/Dystroglycan interactors control neuron behavior in Drosophila eye. BMC Neurosci 12, 93. https://doi.org/10.1186/1471-2202-12-93

Matsuda, H., Yamada, T., Yoshida, M., Nishimura, T., 2015. Flies without Trehalose. Journal of Biological Chemistry 290, 1244–1255. https://doi.org/10.1074/jbc.M114.619411

Michener, C.D., 2007. The bees of the world, 2nd ed. The Johns Hopkins University Press, Baltimore, Maryland.

Michener, C.D., 2000. The Bees of the World. The Johns Hopkins University Press, Baltimore and London.

Michener, C.D., 1946. Notes on the Habits of Some Panamanian Stingless Bees (Hymenoptera, Apidae). Journal of the New York Entomological Society 54, 179–197.

Miyamoto, T., Amrein, H., 2017. Gluconeogenesis: An ancient biochemical pathway with a new twist. Fly (Austin) 11, 218–223. https://doi.org/10.1080/19336934.2017.1283081

Morrissey, M.A., Sherwood, D.R., 2015. An active role for basement membrane assembly and modification in tissue sculpting. J Cell Sci 128, 1661–1668. https://doi.org/10.1242/jcs.168021

Musacchia, F., Basu, S., Petrosino, G., Salvemini, M., Sanges, R., 2015. Annocript: a flexible pipeline for the annotation of transcriptomes able to identify putative long noncoding RNAs. Bioinformatics 31, 2199–2201. https://doi.org/10.1093/bioinformatics/btv106

Nakai, Y., Horiuchi, J., Tsuda, M., Takeo, S., Akahori, S., Matsuo, T., Kume, K., Aigaki, T., 2011. Calcineurin and Its Regulator Sra/DSCR1 Are Essential for Sleep in Drosophila. The Journal of Neuroscience 31, 12759–12766. https://doi.org/10.1523/JNEUROSCI.1337-11.2011

Namiki, T., Niwa, R., Sakudoh, T., Shirai, K., Takeuchi, H., Kataoka, H., 2005. Cytochrome P450 CYP307A1/Spook: A regulator for ecdysone synthesis in insects.

Biochem Biophys Res Commun 337, 367–374.
https://doi.org/10.1016/j.bbrc.2005.09.043

Neuman, S., Kovalio, M., Yaffe, D., Nudel, U., 2005. The Drosophila homologue of the dystrophin gene - Introns containing promoters are the major contributors to the large size of the gene. FEBS Lett 579, 5365–5371. https://doi.org/10.1016/j.febslet.2005.08.073

Nixon, K.C.J., Rousseau, J., Stone, M.H., Sarikahya, M., Ehresmann, S., Mizuno, S., Matsumoto, N., Miyake, N., Baralle, D., McKee, S., Izumi, K., Ritter, A.L., Heide, S., Héron, D., Depienne, C., Titheradge, H., Kramer, J.M., Campeau, P.M., 2019. A Syndromic Neurodevelopmental Disorder Caused by Mutations in SMARCD1, a Core SWI/SNF Subunit Needed for Context-Dependent Neuronal Gene Regulation in Flies. The American Journal of Human Genetics 104, 596–610. https://doi.org/10.1016/j.ajhg.2019.02.001

Nogueira-Neto, P., 1997. Vida e criação de abelhas indígenas sem ferrão. Editora Nogueirapis, São Paulo.

Noureddine, M.A., Donaldson, T.D., Thacker, S.A., Duronio, R.J., 2002. Drosophila Roc1a Encodes a RING-H2 Protein with a Unique Function in Processing the Hh Signal Transducer Ci by the SCF E3 Ubiquitin Ligase. Dev Cell 2, 757–770. https://doi.org/10.1016/S1534-5807(02)00164-8

Obiero, G.F., Pauli, T., Geuverink, E., Veenendaal, R., Niehuis, O., Große-Wilde, E., 2021. Chemoreceptor Diversity in Apoid Wasps and Its Reduction during the Evolution of the Pollen-Collecting Lifestyle of Bees (Hymenoptera: Apoidea). Genome Biol Evol 13. https://doi.org/10.1093/gbe/evaa269

O'Dowd, D.K., Smith, M.A., 1996. Single-cell analysis of gene expression in the nervous system. Mol Neurobiol 13, 199–211. https://doi.org/10.1007/BF02740623

Parker, L., Padilla, M., Du, Y., Dong, K., Tanouye, M.A., 2011. Drosophila as a Model for Epilepsy: bss Is a Gain-of-Function Mutation in the Para Sodium Channel Gene That Leads to Seizures. Genetics 187, 523–534. https://doi.org/10.1534/genetics.110.123299

Parker, R., Song, H., 2004. The enzymes and control of eukaryotic mRNA turnover. Nat Struct Mol Biol 11, 121–127. https://doi.org/10.1038/nsmb724

Pedra, J.H.F., McIntyre, L.M., Scharf, M.E., Pittendrigh, B.R., 2004. Genome-wide transcription profile of field- and laboratory-selected dichlorodiphenyltrichloroethane (DDT)-resistant Drosophila. Proceedings of the National Academy of Sciences 101, 7034–7039. https://doi.org/10.1073/pnas.0400580101

Pedro, S.R.M., Cordeiro, G.D., 2015. A new species of the stingless bee Trichotrigona (Hymenoptera: Apidae, Meliponini). Zootaxa 3956, 389. https://doi.org/10.11646/zootaxa.3956.3.4

Perfecto, I., Vandermeer, J.H., 1993. Cleptobiosis in the ant Ectatomma ruidum in Nicaragua. Insectes Soc 40, 295–299. https://doi.org/10.1007/BF01242365

Portugal-Araújo, V., 1958. A Contribution to the Bionomics of Lestrimelitta Cubiceps (Hymenoptera , Apidae). J Kans Entomol Soc 31, 203–211.

Pujari, V., Radebaugh, C.A., Chodaparambil, J. v., Muthurajan, U.M., Almeida, A.R., Fischbeck, J.A., Luger, K., Stargell, L.A., 2010. The Transcription Factor Spn1 Regulates Gene Expression via a Highly Conserved Novel Structural Motif. J Mol Biol 404, 1–15. https://doi.org/10.1016/j.jmb.2010.09.040

Rasmussen, C., Cameron, S.A., 2009. Global stingless bee phylogeny supports ancient divergence, vicariance, and long distance dispersal. Biological Journal of the Linnean Society 99, 206–232. https://doi.org/10.1111/j.1095-8312.2009.01341.x

Ravenscroft, T.A., Janssens, J., Lee, P.-T., Tepe, B., Marcogliese, P.C., Makhzami, S., Holmes, T.C., Aerts, S., Bellen, H.J., 2020. Drosophila Voltage-Gated Sodium Channels Are Only Expressed in Active Neurons and Are Localized to Distal Axonal Initial Segment-like Domains. The Journal of Neuroscience 40, 7999–8024. https://doi.org/10.1523/JNEUROSCI.0142-20.2020

Rech, A.R., Schwade, M.A., Schwade, M.R.M., 2013. Amazonian stingless bees protect meliponaries against robber bees. Acta Amazon 43, 389–393. https://doi.org/10.1590/S0044-59672013000300016

Rewitz, K.F., Rybczynski, R., Warren, J.T., Gilbert, L.I., 2006. The Halloween genes code for cytochrome P450 enzymes mediating synthesis of the insect moulting hormone. Biochem Soc Trans 34, 1256–1260. https://doi.org/10.1042/BST0341256

Ricardo, P.C., Arias, M.C., de Souza Araujo, N., 2024. Decoding bee cleptoparasitism through comparative transcriptomics of Coelioxoides waltheriae and its host Tetrapedia diversipes. Sci Rep 14, 12361. https://doi.org/10.1038/s41598-024-56261-5

Richard, F.-J., Dejean, A., Lachaud, J.-P., 2004. Sugary food robbing in ants: a case of temporal cleptobiosis. C R Biol 327, 509–517. https://doi.org/10.1016/j.crvi.2004.03.002

Robinson, M.D., McCarthy, D.J., Smyth, G.K., 2010. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics 26, 139–140. https://doi.org/10.1093/bioinformatics/btp616

Robinson, M.D., Oshlack, A., 2010. A scaling normalization method for differential expression analysis of RNA-seq data. Genome Biol 11, R25. https://doi.org/10.1186/gb-2010-11-3-r25

Ronque, M.U. v., Migliorini, G.H., Oliveira, P.S., 2018. Thievery in rainforest fungus-growing ants: interspecific assault on culturing material at nest entrance. Insectes Soc 65, 507–510. https://doi.org/10.1007/s00040-018-0632-9

Sakagami, S.F., 1982. Stingless Bees, in: Social Insects. Elsevier, pp. 361–423. https://doi.org/10.1016/B978-0-12-342203-3.50011-4

Sakagami, Shôichi F, Laroca, S., 1963. Additional Observations on the Habits of the Cleptobiotic Stingless Bees, the Genus Lestrimelitta Friese (Hymenoptera,

Apoidea). Journal of the Faculty of Science Hokkaido University Series VI. Zoology 15, 319–339. https://doi.org/10.18960/seitai.14.6_259_5

Sakagami, Shoichi F, Laroca, S., 1963. Additional observations on the habits of the cleptobiotic stingless bees , the genus Lestrimelitta Friese (Hymenoptera, Apoidea). Journal of the Faculty of Science, Hokkaido University: Zoology. Series VI 15, 319–339.

Sakagami, S.F., Roubik, D.W., Zucchi, R., 1993. Ethology of the robber stingless bee, Lestrimelitta limao (Hymenoptera: Apidae). Sociobiology 21, 237–277.

Segers, F.H.I.D., von Zuben, L., Grüter, C., 2016. Local differences in parasitism and competition shape defensive investment in a polymorphic eusocial bee. Ecology 97, 417–426. https://doi.org/10.1890/15-0793.1

Shcherbata, H.R., Yatsenko, A.S., Patterson, L., Sood, V.D., Nudel, U., Yaffe, D., Baker, D., Ruohola-Baker, H., 2007. Dissecting muscle and neuronal disorders in a Drosophila model of muscular dystrophy. EMBO J 26, 481–493. https://doi.org/10.1038/sj.emboj.7601503

Simão, F.A., Waterhouse, R.M., Ioannidis, P., Kriventseva, E. v., Zdobnov, E.M., 2015. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. Bioinformatics 31, 3210–3212. https://doi.org/10.1093/bioinformatics/btv351

Smith-Unna, R., Boursnell, C., Patro, R., Hibberd, J.M., Kelly, S., 2016. TransRate: reference-free quality assessment of de novo transcriptome assemblies. Genome Res 26, 1134–1144. https://doi.org/10.1101/gr.196469.115

Stincone, A., Prigione, A., Cramer, T., Wamelink, M.M.C., Campbell, K., Cheung, E., Olin□Sandoval, V., Grüning, N., Krüger, A., Tauqeer Alam, M., Keller, M.A., Breitenbach, M., Brindle, K.M., Rabinowitz, J.D., Ralser, M., 2015. The return of metabolism: biochemistry and physiology of the pentose phosphate pathway. Biological Reviews 90, 927–963. https://doi.org/10.1111/brv.12140

Supek, F., Bošnjak, M., Škunca, N., Šmuc, T., 2011. REVIGO Summarizes and Visualizes Long Lists of Gene Ontology Terms. PLoS One 6, e21800. https://doi.org/10.1371/journal.pone.0021800

Suzek, B.E., Wang, Y., Huang, H., McGarvey, P.B., Wu, C.H., 2015. UniRef clusters: a comprehensive and scalable alternative for improving sequence similarity searches. Bioinformatics 31, 926–932. https://doi.org/10.1093/bioinformatics/btu739

Takai, A., Yamaguchi, M., Yoshida, H., Chiyonobu, T., 2020. Investigating Developmental and Epileptic Encephalopathy Using Drosophila melanogaster. Int J Mol Sci 21, 6442. https://doi.org/10.3390/ijms21176442

Takeo, S., Swanson, S.K., Nandanan, K., Nakai, Y., Aigaki, T., Washburn, M.P., Florens, L., Hawley, R.S., 2012. Shaggy/glycogen synthase kinase 3β and phosphorylation of Sarah/regulator of calcineurin are essential for completion of Drosophila female meiosis. Proceedings of the National Academy of Sciences 109, 6382–6389. https://doi.org/10.1073/pnas.1120367109

Tarazona, S., Furió-Tarí, P., Turrà, D., Pietro, A. di, Nueda, M.J., Ferrer, A., Conesa, A., 2015. Data quality aware analysis of differential expression in RNA-seq with NOISeq R/Bioc package. Nucleic Acids Res 43, gkv711. https://doi.org/10.1093/nar/gkv711

Thackeray, J.R., Ganetzky, B., 1995. Conserved alternative splicing patterns and splicing signals in the Drosophila sodium channel gene para. Genetics 141, 203–214. https://doi.org/10.1093/genetics/141.1.203

van der Plas, M.C., Pilgram, G.S.K., Plomp, J.J., de Jong, A., Fradkin, L.G., Noordermeer, J.N., 2006. Dystrophin Is Required for Appropriate Retrograde Control of Neurotransmitter Release at the Drosophila Neuromuscular Junction. The Journal of Neuroscience 26, 333–344. https://doi.org/10.1523/JNEUROSCI.4069-05.2006

Vázquez, M., Muñoz, D., Medina, R., Paxton, R.J., de Oliveira, F.F., Quezada-Euán, J.J.G., 2022. Sympatric cleptobiotic stingless bees have species-specific cuticular profiles that resemble their hosts. Sci Rep 12, 2621. https://doi.org/10.1038/s41598-022-06683-w

von Zuben, L.G., Schorkopf, D.L.P., Elias, L.G., Vaz, A.L.L., Favaris, A.P., Clososki, G.C., Bento, J.M.S., Nunes, T.M., 2016. Interspecific chemical communication in raids of the robber bee Lestrimelitta limao. Insectes Soc 63, 339–347. https://doi.org/10.1007/s00040-016-0474-2

Wahle, E., Winkler, G.S., 2013. RNA decay machines: Deadenylation by the Ccr4–Not and Pan2–Pan3 complexes. Biochimica et Biophysica Acta (BBA) - Gene Regulatory Mechanisms 1829, 561–570. https://doi.org/10.1016/j.bbagrm.2013.01.003

Wang, Y., Bogenhagen, D.F., 2006. Human Mitochondrial DNA Nucleoids Are Linked to Protein Folding Machinery and Metabolic Enzymes at the Mitochondrial Inner Membrane. Journal of Biological Chemistry 281, 25791–25802. https://doi.org/10.1074/jbc.M604501200

Wcislo, Wi.T., 1987. The roles of seasonality, host synchrony, and behaviour in the evolutions and distributions of nest parasites in Hymenoptera (Insecta), with special reference to bees (Apoidea). Biological Reviews 62, 515–542. https://doi.org/10.1111/j.1469-185X.1987.tb01640.x

Werck-Reichhart, D., Feyereisen, R., 2000. Cytochromes P450: a success story. Genome Biol 1. https://doi.org/10.1186/gb-2000-1-6-reviews3003

Wheeler, W.M., 1910. Ants: Their Structure, Development, and Behavior. The Columbia University Press, New York.

Wille, A., 1983. Biology of the Stingless Bees. Annu Rev Entomol 28, 41–64. https://doi.org/10.1146/annurev.en.28.010183.000353

Willingham, A.T., Keil, T., 2004. A tissue specific cytochrome P450 required for the structure and function of Drosophila sensory organs. Mech Dev 121, 1289–1297. https://doi.org/10.1016/j.mod.2004.04.017

Wittmann, D., Radtke, R., Zeil, J., Lübke, G., Francke, W., 1990. Robber bees (Lestrimelitta limao) and their host chemical and visual cues in nest defense by Trigona (Tetragonisca) angustula (Apidae: Meliponinae). J Chem Ecol 16, 631–641. https://doi.org/10.1007/BF01021793

Wong, J.J.L., Li, S., Lim, E.K.H., Wang, Y., Wang, C., Zhang, H., Kirilly, D., Wu, C., Liou, Y.-C., Wang, H., Yu, F., 2013. A Cullin1-Based SCF E3 Ubiquitin Ligase Targets the InR/PI3K/TOR Pathway to Regulate Neuronal Pruning. PLoS Biol 11, e1001657. https://doi.org/10.1371/journal.pbio.1001657

Xi, J., Pan, Y., Bi, R., Gao, X., Chen, X., Peng, T., Zhang, M., Zhang, H., Hu, X., Shang, Q., 2015. Elevated expression of esterase and cytochrome P450 are related with lambda–cyhalothrin resistance and lead to cross resistance in Aphis glycines Matsumura. Pestic Biochem Physiol 118, 77–81. https://doi.org/10.1016/j.pestbp.2014.12.002

Xiao, X., Chen, C., Yu, T.-M., Ou, J., Rui, M., Zhai, Y., He, Y., Xue, L., Ho, M.S., 2017. Molecular Chaperone Calnexin Regulates the Function of Drosophila Sodium Channel Paralytic. Front Mol Neurosci 10. https://doi.org/10.3389/fnmol.2017.00057

Yamaguchi, T., 1995. Intraspecific competition through food robbing in the harvester ant, Messor aciculatus (Fr. Smith), and its consequences on colony survival. Insectes Soc 42, 89–101. https://doi.org/10.1007/BF01245701

Yoh, S.M., Cho, H., Pickle, L., Evans, R.M., Jones, K.A., 2007. The Spt6 SH2 domain binds Ser2-P RNAPII to direct Iws1-dependent mRNA splicing and export. Genes Dev 21, 160–174. https://doi.org/10.1101/gad.1503107

Zhan, Y., Melian, N.Y., Pantoja, M., Haines, N., Ruohola-Baker, H., Bourque, C.W., Rao, Y., Carbonetto, S., 2010. Dystroglycan and Mitochondrial Ribosomal Protein L34 Regulate Differentiation in the Drosophila Eye. PLoS One 5, e10488. https://doi.org/10.1371/journal.pone.0010488

Zhao, H., Gao, P., Du, H., Ma, W., Tian, S., Jiang, Y., 2014. Molecular characterization and differential expression of two duplicated odorant receptor genes, AcerOr1 and AcerOr3, in Apis cerana cerana. J Genet 93, 53–61. https://doi.org/10.1007/s12041-014-0332-9

Zhou, Y., 2021. SCBN: A statistical normalization method and differential expression analysis for RNA-seq data between different species. R package version 1.10.0. https://doi.org/10.18129/B9.bioc.SCBN

Zhou, Y., Zhu, J., Tong, T., Wang, J., Lin, B., Zhang, J., 2019. A statistical normalization method and differential expression analysis for RNA-seq data between different species. BMC Bioinformatics 20, 163. https://doi.org/10.1186/s12859-019-2745-1