Pupil size and eye movements differently index effort
in both younger and older adults

Björn Herrmann[1,2], Jennifer D. Ryan[1,2,3]

[1]Rotman Research Institute,
Baycrest Academy for Research and Education, M6A 2E1, North York, ON, Canada

[2]Department of Psychology,
University of Toronto, M5S 1A1, Toronto, ON, Canada

[3]Department of Psychiatry,
University of Toronto, M5S 1A1, Toronto, ON, Canada

Correspondence concerning this article should be addressed to Björn Herrmann, Rotman Research Institute, Baycrest Academy for Research and Education, 3560 Bathurst St, North York, ON, M6A 2E1, Canada. E-mail: bherrmann@research.baycrest.org

Keywords: Effort, speech perception, pupillometry, eye movements, aging

Word count: Abstract (249), Significance statement (120), Introduction (648), Discussion (1593)

Page, figure, table count: Number of pages (27), Number of figures (5), Number of tables (0)

Conflict of interest statement: The authors declare no competing interests.

Acknowledgments: The research was supported by the Canada Research Chair program (CRC-2019-00156, 232733), the Natural Sciences and Engineering Research Council of Canada (NSERC Discovery Grant: RGPIN-2021-02602), the Canadian Institutes of Health Research (CIHR: 186236), and the William Demant Foundation. We thank Tazeen Atif and Priya Pandey for help during data collection. We thank Ingrid Johnsrude and Lauren Fink for fruitful discussions.

Author contributions: BH: Conceptualization, methodology, validation, formal analysis, investigation, data curation, writing - original draft, writing - review & editing, visualization, supervision, project management, funding acquisition. JDR: Conceptualization, methodology, investigation, writing - review & editing.

39   Abstract

The assessment of mental effort is increasingly relevant in neuro-cognitive and lifespan domains.

Pupillometry – the measure of the pupil size – is often used to assess effort but has disadvantages.

Analysis of eye movements may provide an alternative, but research has been limited to easy and

difficult task demands in younger adults. An effort measure must be sensitive to the whole effort

profile, including 'giving-up' effort investment, and capture effort in different age groups. The current

study comprised three experiments in which younger and older adults (both sexes) listened to speech

masked by background babble at different signal-to-noise ratios associated with easy, difficult, and

impossible speech comprehension. We expected individuals to invest little effort for easy and

impossible speech ('giving up') but to exert effort for difficult speech. Indeed, pupil size was largest for

difficult, but lower for the easy and impossible speech conditions. In contrast, gaze dispersion

decreased with increasing speech masking in both age groups. Critically, gaze dispersion during

difficult speech returned to levels similar to easy speech after sentence offset, where acoustic

stimulation was similar across conditions, whereas gaze dispersion during impossible speech continued

to be reduced. These findings show that a reduction in eye movements is not a byproduct of acoustic

factors, but instead suggest that neuro-cognitive processes, different from arousal-related systems

regulating the pupil size, drive reduced eye movements during high task demands. The current data

thus show that effort in one sensory domain (audition) differentially impacts distinct functional

properties in another sensory domain (vision).

Significance statement

The assessment of mental effort is increasingly relevant in many domains. Here, we investigated the sensitivity of a novel measure of effort – the spatial dispersion of eye movements – and compared it to pupillometry, which is a more common effort-assessment tool. Individuals listened to speech masked by background babble at levels associated with easy, difficult, and impossible speech comprehension. Pupil size was largest for difficult, but lower for easy and impossible speech conditions (giving up listening). In contrast, the spatial dispersion of eye movements decreased with increasing speech masking, but this effect was independent of acoustic factors. Instead, the current results suggest that neuro-cognitive processes, different from arousal-related systems regulating the pupil size, drive reduced eye movements during high task demands.

Introduction

The concept of 'mental workload', or effort, is relevant in many neuro-cognitive domains (Brehm and
Self, 1989; Hockey, 1997; Kool and Botvinick, 2018). For example, older adults often invest substantial
effort to comprehend speech in noise, and this effort is considered an early sign of age-related hearing
loss (Eckert et al., 2016; Pichora-Fuller et al., 2016; Peelle, 2018; Herrmann and Johnsrude, 2020).
Research is thus increasingly dedicated to quantifying effort (Zekveld et al., 2010; McGarrigle et al.,
2014; Moran et al., 2016; Francis and Love, 2020; Shechter and Share, 2020; Slade et al., 2021; Kraus
et al., 2023b), but existing effort measures have limitations.

Pupillometry – the measurement of the pupil size – is commonly used to assess effort (van der
Wel and van Steenbergen, 2018; Winn et al., 2018; Zekveld et al., 2018). However, pupil size can vary
due to factors unrelated to task demands, such as changes in light (Knapen et al., 2016; Suzuki et al.,
2019; Thurman et al., 2021) or in the angle of the eye relative to the eye-tracking camera (Brisson et
al., 2013; Hayes and Petrov, 2016; Fink et al., 2021). To address the latter, individuals typically maintain
fixation on a point on a computer monitor (Zekveld et al., 2018; Winn and Teece, 2021; Kraus et al.,
2023a), but restriction of gaze can reduce memory and mental imagery (Johansson et al., 2012).
Pupillometry has been used to investigate effort in older adults (Kuchinsky et al., 2013; Ayasse and
Wingfield, 2018), but aging reduces the dilation range (Piquado et al., 2010; Guillon et al., 2016; Zhao
et al., 2019), which could lead to effort misestimations. Moreover, effort-related changes in pupil size
are driven by arousal systems in the brain (Mathôt, 2018; Joshi and Gold, 2020; Strauch et al., 2022),
but arousal can differ between and within populations, adding unwanted variability; for example,
arousal can be higher for individuals with hearing loss (Ayasse and Wingfield, 2020).

Recent work suggests that the quantification of eye movements may provide an alternative
effort metric. Eye movements decrease during periods of high compared to low memory load (Kosch
et al., 2018; Walter and Bex, 2021) and with increasing speech masking (Contadini-Wright et al., 2023;
Cui and Herrmann, 2023). This research has focused mainly on increases in effort as task demands
increase. Yet, disengagement or 'giving up' effort investment is common for tasks that feel too
effortful, when the reward does not justify the required mental costs (Braver et al., 2014; Westbrook
and Braver, 2015; Herrmann and Johnsrude, 2020). Any measure of effort must be sensitive to
disengagement, not only because it is a critical, cognitive transition point (Richter, 2013; Westgate and
Wilson, 2018), but also to ensure that the measure is indeed sensitive to variations in effort. The pupil
size shows an inverted u-shape: it is largest for difficult listening tasks, but lower during easy and
impossible listening ('giving up'; Zekveld and Kramer, 2014; Ohlenforst et al., 2018; Wendt et al.,
2018). Whether measures of eye movements also exhibit this u-shaped effort profile associated with
easy to difficult to impossible task demands is unclear. Functional differences between pupil and eye
movement patterns may suggest distinct neural mechanisms that reflect effort.

In contrast to measurements of pupil size, measuring reductions in eye movements may be less
affected by age-related changes (Sharpe and Sylvester, 1978; Spooner et al., 1980; Zackon and Sharpe,
1987; Bruenech, 2018) as the dynamics related to executing eye movements appear little affected by
aging (Abrams et al., 1998). Further, older adults tend to make more eye movements than younger
adults (Ryan et al., 2007; Liu et al., 2018; Mazloum-Farzaghi et al., 2022), which may facilitate

sensitivity to eye-movement reductions related to effort. Therefore, in the current study, measures of
pupil size and the spatial extent of eye movement exploration (i.e., gaze dispersion) were contrasted
for the full effort profile in three listening experiments for younger and older adults.

General methods

Participants

113
Younger adults and older adults participated in three experiments. Participants were native English
speakers or learned English before the age of 5 years. They reported having normal hearing and did
not wear hearing aids or were prescribed one (with the exception of one person whose data were
excluded). Participants gave written informed consent prior to the experiment and were paid $7.5 CAD
per half-hour for their participation. The study was conducted in accordance with the Declaration of
Helsinki, the Canadian Tri-Council Policy Statement on Ethical Conduct for Research Involving Humans
(TCPS2-2014), and was approved by the Research Ethics Board of the Rotman Research Institute at
Baycrest.
In Experiment 1, 23 younger adults (median age: 25 years, age range: 18–40 years; 4 male, 17
female, 1 non-binary, 1 trans male) and 20 older adults (median age: 66 years, age range: 56–78 years;
7 male, 13 female) participated. Data from four additional participants were recorded, but not
analyzed because they wore hearing aids (n=1), reported having a neurological disease (n=1), their
proportion of correct responses for the easiest condition in the behavioral task was equal to or lower
than 0.7 (n=1), or more than 50% of trials of their physiological recordings contained more than 50% of
missing data (n=1). One person's pupil data were not included in the analyses, because the person's
pupil data were an extreme outlier (note that inclusion of this data point did not change the
significance or absence thereof of statistical tests).
In Experiment 2, 20 younger adults (median age: 23 years, age range: 20–35 years; 10 male, 9
female, 1 non-binary) and 22 older adults (median age: 67 years, age range: 56–77 years; 9 male, 13
female) participated. Data from six additional participants were recorded, but not analyzed because
they were a non-native English speaker (n=1), their proportion of correct responses for the easiest
condition in the behavioral task was equal to or lower than 0.7 (n=2), or more than 50% of trials of
their physiological recordings contained more than 50% of missing data (n=3). One other person
indicated having migraines sometimes, but this did not affect their participation. The data of this
person were included.
In Experiment 3, 23 younger adults participated (median age: 25 years, age range: 18–33 years;
11 male, 10 female, 2 non-binary). Data from 2 additional participants were recorded but not analyzed
because more than 50% of trials contained more than 50% of missing data (n=2). Since younger and
older adults showed similar results in the first two experiments, only younger adults were recruited in
the last experiment.

Hearing assessment

144
Participants sat in a comfortable chair within a sound booth during all experimental procedures. In
Experiments 1 and 2, pure-tone audiometric thresholds were obtained for frequencies ranging from
0.25 to 8 kHz using a Maico M28 audiometer. Audiometry was recorded for all older adults, but only
for 17 and 9 younger adults for Experiment 1 and 2, respectively, due to misunderstandings related to
the execution of the experimental protocols. No audiometry was recorded in Experiment 3, because
no older adults were recruited. The pure-tone average threshold was calculated as the mean threshold
across 0.5 to 4 kHz frequencies. An independent-samples t-test was used to analyze age-group
differences in the pure-tone average threshold. Despite participants self-reporting normal hearing,
some adults showed clinically relevant hearing loss (see below). Data from all adults were included
nevertheless, because a) our sample is representative of community-dwelling adults who vary in their
degree of hearing loss despite self-describing their hearing as normal (Moore, 2007; Plack, 2014;
Presacco et al., 2016; Herrmann et al., 2018, 2023), b) they were able to do the behavioral speech-in-
noise task, and c) our study seeks to investigate whether eye movements are indicative of listening
effort in younger and older adults more generally, rather than investigating differences in people with
versus those without hearing loss.

Experimental setup

160
Sounds were presented via Sony Dynamic Stereo MDR-7506 headphones and a Steinberg UR22 mkII
(Steinberg Media Technologies) external sound card. Experimental procedures were run using
Psychtoolbox (v3.0.14) in MATLAB (MathWorks Inc.) on a Lenovo T450s laptop with Microsoft
Windows XP. The laptop screen was mirrored to a ViewSonic monitor with a refresh rate of 60 Hz. All
sounds were presented at a comfortable listening level that was fixed across participants (~70-75 dB
SPL).

Stimulus materials and procedure

167
Stimulus materials consist of sentences from the Harvard sentence lists spoken by a male native
English speaker (mean duration: 2.5 s; range of durations: 2–3.1 s) (IEEE, 1969). Speech recordings
were taken from a publicly available resource (speaker ID: 'PNM055';
https://depts.washington.edu/phonlab/projects/uwnu.php; Panfili et al., 2017; McCloy et al., 2018).
Sentences were embedded in a 6.3-second 12-talker babble noise (Bilger, 1984). A sentence started
2.8 s after babble onset (Zhang et al., 2022; Cui and Herrmann, 2023; Figure 1A). For younger adults,
sentences were presented at -17 dB (impossible), -4 dB (difficult), or +9 dB SNR (easy). For older adults,
SNR levels were increased by 2 dB (-15, -2, +11 dB SNR) relative to younger adults to account for
known challenges of older people with the perception of speech in noise (Gustafsson and Arlinger,
1994; Helfer and Freyman, 2008; Presacco et al., 2019; Irsik et al., 2022; Herrmann, 2023). We
expected listening effort to be low for impossible and easy SNRs, because participants would not invest
cognitively for the former and required little effort to comprehend sentences for the latter. Highest
listening effort was expected for the difficult SNR. Different SNRs were achieved by adjusting the
sound level of the sentence, while keeping the babble level constant across trials (Ohlenforst et al.,

2017; Kadem et al., 2020; Cui and Herrmann, 2023). This ensured that stimuli with different SNRs did
not differ prior to sentence onset (Ohlenforst et al., 2017; Kadem et al., 2020; Cui and Herrmann,
2023). After each stimulus, a probe word occurred on the screen. Participants were asked to indicate
whether the probe word was semantically related or unrelated to the sentence (Rodd et al., 2010a;
Rodd et al., 2010b; Kadem et al., 2020; Cui and Herrmann, 2023).

Participants listened to 60 sentences per SNR condition across 5 experimental blocks. Within
each block, participants listened to 12 sentences per SNR condition. The order of SNR conditions was
randomized such that each SNR level occurred maximally two times in direct succession (Experiment 1
and 2) or was blocked with 12 sentences of the same SNR in direct succession (Experiment 3).
Assignment of SNR levels to specific sentences was randomized across participants to avoid
confounding specific sentences with SNR levels. Participants underwent a short 12-trials training block
to familiarize them with the eye-tracking calibration procedures and the semantic-relatedness task.
Different sentences from the Harvard sentence lists were used in Experiments 1, 2, and 3.

Pupillometry studies typically require a participant to fixate centrally on a fixation point on a
computer screen to reduce eye movements (Zekveld et al., 2014; Zekveld et al., 2018; Zhao et al.,
2019; Farahani et al., 2020; Kadem et al., 2020; Winn and Teece, 2021). However, an investigation into
the degree to which eye movements reflect listening effort may be impeded by restricting eye
movements (Cui and Herrmann, 2023) and can reduce behavioral speech comprehension effects (Cui
and Herrmann, 2023). Here we presented an incidental multi-object movement display to facilitate eye
movements, similar to our recent work (Cui and Herrmann, 2023; see other work using object tracking,
Cavanagh and Alvarez, 2005; Alvarez and Franconeri, 2007; Scholl, 2009; Herrmann and Johnsrude,
2018a, b). To this end, 16 white dots [dot diameter: 1.2 cm (0.9°)] were presented on gray background
and moved on the screen during the presentation of the auditory stimuli. Presentation of dots was
constrained to a display frame of 20.6 cm width (15.6°) and 19.4 cm height (14.7°) centered on the
screen and highlighted to the participants by a light gray frame (Figure 1A). Dots never moved outside
of the display frame and never overlapped during movements; dots moved ~3.7 cm/s (2.8°/s). The dot
display occurred 0.7 s prior to babble onset and remained stationary for the 0.7 s duration. Dot
movements started with the onset of the babble and continued throughout the 6.3 s duration of the
auditory stimulus, at which point the dots disappeared. Critically, participants were instructed that
their task was to comprehend the sentence and perform the semantic-relatedness task. No task was
required for the dot display. Participants were instead instructed to look at the screen in whatever way
they wanted (Johansson et al., 2006; Johansson et al., 2011; Johansson et al., 2012; Cui and Herrmann,
2023).

The three experiments of the current study differed in the degree to which individuals could
anticipate the comprehension difficulty of an upcoming sentence. In Experiment 1, no cues were given
to participants, and they therefore only knew about the difficulty of a sentence upon sentence onset.
Importantly, anticipation of the level of difficulty is a critical component in effort frameworks to help
an individual to know when to disengage to preserve effort and reduce fatigue (Brehm and Self, 1989;
Richter, 2013; Richter et al., 2016; Herrmann and Johnsrude, 2020). In Experiment 2, a visual cue
indicated the difficulty of an upcoming sentence. To this end, the 16 dots of the 0.7-s stationary dot-
display were color-coded, with green dots indicating easy comprehension, yellow dots indicating

difficult comprehension, and red dots indicating impossible comprehension. Upon babble onset, the
color of the dots turned white and all dots started to move. The color to difficulty assignment was not
counter-balanced across participants, because we wanted to avoid confusing color-difficulty
associations (i.e., red indicating easy) and we reasoned the cue color is unlikely to influence ocular
responses 2.8 s later (sentence onset). We anticipated that color cues would enable participants to
regulate effort investment on a sentence-by-sentence basis. In Experiment 3, SNR conditions (easy,
difficult, impossible) were presented in mini-blocks of 12 sentences during an experimental block. Prior
to each mini-block, participants were also visually presented with the text "Comprehension of the next
few sentences will likely be: …" completed by either "easy", "difficult", or "impossible" depending on
the SNR condition. The dots were white throughout and did not cue SNR conditions. By blocking SNR
conditions, we reasoned that participants would more easily be able to regulate effort investment than
in Experiment 2.

Behavioral data analysis
The proportion of correct responses in the semantic-relatedness task was calculated for each SNR
condition. Behavioral and physiological data from participants who had a proportion of correct
responses equal to or lower than 0.7 in the easiest condition were excluded, based on the assumption
that, in such cases, participants either did not pay sufficient attention or experienced undue
difficulties. A repeated measures analysis of variance (rmANOVA) was calculated using the proportion
of correct responses as dependent measure. SNR (easy, difficult, impossible) was included as a within-
participants factor and age group (younger, older) was included as a between-participants factor (with
the exception of Experiment 3, which only included younger adults). Post hoc tests were calculated for
significant main effects and interactions. Holm's methods was used to correct for multiple
comparisons (Holm, 1979).

Pupillometry and eye-movement recordings
During the experiments, participants placed their head on a chin and forehead rest facing the
computer monitor at a distance of about 70 cm. Pupil area and eye movements were recorded
continuously from the right eye (or the left eye if the right eye could not be tracked accurately) using
an integrated infrared camera (EyeLink 1000 eye tracker; SR Research Ltd.) at a sampling rate of 500
Hz. Nine-point fixation was used for eye-tracker calibration prior to each block (McIntire et al., 2014).

Processing of pupil-size and eye-movement data
The main metrics of the current study were pupil size and gaze dispersion (Cui and Herrmann, 2023).
Preprocessing of pupil-size and eye-movement data involved removing eye blinks and other artifacts.
For each eye blink indicated by the eye tracker, all data points of the pupil time course and the x and y
eye-coordinate time courses between 100 ms before and 200 ms after a blink were set to NaN ('not a
number' in MATLAB). In addition, pupil-size values that differed from the mean pupil size by more than
3 times the standard deviation were classified as outliers and the corresponding data points of the

pupil time course and the x and y eye-coordinate time courses were set to NaN. Missing pupil data
(coded as NaN) resulting from artifact rejections and outlier removal were interpolated using
MATLAB's 'pchip' method. Pupil-size time courses were filtered with a 5-Hz low-pass filter (51 points,
Kaiser window). X and y eye-coordinate time courses were not interpolated. That is, missing data
points (NaNs) were ignored in the calculation of the eye-movement metric (i.e., gaze dispersion).

Changes in eye movements were investigated using spatial gaze dispersion (Cui and Herrmann,
2023). Spatial gaze dispersion is a measure of the general tendency for the eyes to move around. It
was calculated as the standard deviation in gaze across time points, averaged across x- and y-
coordinates, and transformed to logarithmic values. Smaller values indicate less gaze dispersion. To
obtain time courses for gaze dispersion, it was calculated for 1-s sliding time windows centered
sequentially on each time point. If more than 90% of data were unavailable within a 1-s time window
(that is, 450 or more samples were NaN-coded), gaze dispersion for the corresponding time point was
set to NaN and ignored during averaging (Cui and Herrmann, 2023). Gaze dispersion is a broad
measure that captures any eye movements. Analyses of saccades (including microsaccades) require
setting a velocity threshold to determine what counts as a saccade (Engbert and Kliegl, 2003; Engbert,
2006; Widmann et al., 2014; Kadem et al., 2020; Contadini-Wright et al., 2023; Cui and Herrmann,
2023), but the threshold can influence whether or not effects of cognition on saccades can be
observed (Cui and Herrmann, 2023). Nonetheless, we explored the sensitivity of saccades to changes
in SNR and results were very similar to the results for gaze dispersion.

Continuous pupil-size and gaze-dispersion data were divided into single-trial time courses
ranging from -1.7 to 6.3 s time-locked to babble onset. Data for an entire trial were excluded from
analysis if the percentage of NaN data entries made up more than 50% of the trial. Pupil size and gaze
dispersion were averaged across trials, separately for each SNR condition. Pupil size was also
normalized relative to the -1.7 s to -0.7 s baseline period (i.e., the period prior to the onset of the
visual dot display), by subtracting the mean baseline pupil size from the pupil size at each time point.
Gaze dispersion does not require baseline correction.

For statistical analyses, gaze dispersion was averaged across the 2.8–5.3-s time window during
which sentences were presented (sentences started at 2.8 s after babble onset; average sentence
offset was at 5.3 s). Mean pupil size was calculated for a time window that was delayed by 0.5 s to 3.3–
5.8-s, because the pupil size is known to change relatively slowly and peak late during sentence
listening (Knapen et al., 2016; Winn and Moore, 2018; Winn et al., 2018; Kadem et al., 2020; Zhang et
al., 2022). We also analyzed data after sentence offset, focusing on the 5.3 s to 6 s time window, to
investigate whether gaze dispersion is sensitive to decreased listening effort when speech has ended
(we ignored the last 0.3 s prior to babble offset to avoid impacting the analyses by back smearing of
eye movement activity following babble offset). Separate rmANOVAs were calculated for different
measures (pupil size, gaze dispersion) and time windows, using the within-participants factor SNR
(easy, difficult, impossible) and the between-participants factor Age Group (younger, older). Age Group
was not included in Experiment 3, because only younger adults were recruited. Post hoc tests were
calculated for significant main effects and interactions. Holm's methods was used to correct for
multiple comparisons (Holm, 1979). We also examined linear and quadratic SNR effects. A quadratic
effect in the absence of a linear effect would be consistent with the expected inverted u-shaped effort

profile. Statistical analyses were carried out in JASP software (v0.18.1; JASP, 2023). Note that JASP uses
pooled error terms and degrees of freedom for post hoc, linear, and quadratics effects of an
rmANOVA.

Results for Experiment 1

Younger and older adults participated in Experiment 1. Pure-tone audiometric thresholds were higher
in older compared to younger adults (t_{35} = 3.099, p = 0.004, d = 1.022), as expected, and some adults
showed clinically relevant hearing loss (Figure 1B). Data from these individuals were included in the
analyses nonetheless as described above.
Participants listened to sentences in background babble at three different SNR levels that were
selected to make sentence comprehension easy, difficult, or impossible. No prior cues were provided
to participants about whether comprehension for an upcoming sentence would be easy, difficult, or
impossible. Participants judged whether the probe word that followed a sentence was semantically
related to the sentence. Performance in the semantic relatedness task was higher for easy than
difficult (t_{82} = 10.974, p_{Holm} = 9.1 · 10^{-18}, d = 2.248), easy than impossible (t_{82} = 35.119, p_{Holm} = 1 · 10^{-50},
d = 7.193), and difficult than impossible SNRs (t_{82} = 24.144, p_{Holm} = 1 · 10^{-38}, d = 4.945; effect of SNR:
$F_{2,82}$ = 645.578, p = 6.6 · 10^{-51}, ω^2 = 0.902), as expected. Performance was higher in older than younger
adults for the difficult condition (t_{82} = 2.942, p_{Holm} = 0.012, d = 0.899; SNR × Age Group interaction:
$F_{2,82}$ = 4.590, p = 0.013, ω^2 = 0.049), but not for the other SNR conditions (for all p_{Holm} > 0.5).
Differences between SNR conditions were significant for both age groups (for all p_{Holm} < 0.001). Hence,
the SNR manipulation influenced behavior as intended, but older adults performed better than
younger adults on the difficult condition (Figure 1C).

Figure 1: Experimental design, pure-tone audiometric thresholds, and behavioral data. A: Stimulation paradigm. A sentence was presented in a 6.3-s babble noise. The sentence onset was 2.8 s and the averaged sentence offset was 5.3 s. A moving-dot display was presented while participants listened to the auditory stimulus. Following a 0.7-s stationary display prior to babble onset, sixteen dots smoothly moved on the screen throughout the entire duration of the auditory stimulus. No task was required using the moving-dot display. B: Left: Pure-tone audiometric thresholds: The thin lines reflect data from each individual participant. Thick lines

reflect the mean across participants. **Right:** Pure-tone average hearing threshold (mean across 0.5, 1, 2, and 4 kHz). Gray dots reflect the pure-tone average threshold for individual participants. **C:** Proportion of correct responses in the semantic-relatedness task for younger (left) and older adults (right). Gray dots reflect data from individual participants. The dashed horizontal line at 0.5 reflects chance level. Error bars reflect that standard error of the mean.

The pupil size was larger during the difficult than the impossible SNR (t_{80} = 3.107, p_{Holm} = 0.008,
d = 0.276) and numerically, but non-significantly larger for the difficult than the easy SNR (t_{80} = 1.981,
p_{Holm} = 0.102, d = 0.176), whereas easy and impossible conditions did not differ (p_{Holm} = 0.264; effect of
SNR: $F_{2,80}$ = 4.947, p = 0.009, ω^2 = 0.011; Figure 2). The quadratic (t_{80} = -2.937, p = 0.004) but not the
linear (p = 0.264) SNR effect was also significant. There was no effect of Age Group and no Age Group ×
SNR interaction (for both p > 0.05). The lower pupil size for the impossible SNR relative to the difficult
SNR suggests that individuals did not invest cognitively, and perhaps gave up listening, when
comprehension was not possible (cf. Zekveld and Kramer, 2014; Ohlenforst et al., 2017; Ohlenforst et
al., 2018; Wendt et al., 2018). That the pupil size did not differ between difficult and easy SNRs might
indicate that some cognitive investment was needed for comprehending speech under the easy
condition, although the quadratic SNR effect shows the expected inverted u-shaped effort profile.
There was also a transient increase in the pupil size for the easy SNR following sentence onset (Figure
2A). This transient increase may have been due to increased arousal following the unpredictable
sound-level increase associated with the high SNR of the sentence (+9 dB SNR and +11 dB SNR for
younger and older adults, respectively), and may have made observing a difference between the
difficult and easy SNRs more challenging.

Figure 2: Pupil size and gaze dispersion for Experiment 1. A: Pupil size time courses. The shaded area reflects the standard error of the mean (between-participant variances removed; Masson and Loftus, 2003). The dashed vertical lines indicate, from left to right, the onset of the dot display (stationary), onset of the babble noise and dot movements, sentence onset. **B:** Mean pupil size across the 3.3–5.8 s time window (sentence onset at 2.8 s and average sentence offset at 5.3 s). Error bars reflect the standard error of the mean (removal of between-participant variance; Masson and Loftus, 2003). **C:** Difference in pupil size between different SNR conditions. Violin plots reflect the histogram of individual data points. Dots reflect the mean. **D-F:** The same types of data plots as in panels A-C are depicted for gaze dispersion. Panel E and F reflect mean gaze dispersion across the 2.8-5.3 s time window.

Gaze dispersion was also sensitive to SNR ($F_{2,82}$ = 15.240, p = 9.5 · 10^{-6}, ω^2 = 0.016), but
differently than the pupil size. Gaze dispersion decreased when speech comprehension was difficult
(t_{82} = 3.465, p_{Holm} = 0.002, d = 0.204) and impossible (t_{82} = 5.455, p_{Holm} = 1.5 · 10^{-6}, d = 0.321) relative to
when comprehension was easy. Eye movements were further reduced under the impossible compared
to the difficult condition (t_{82} = 1.99, p_{Holm} = 0.05, d = 0.117). This was also reflected in the linear (t_{82} =
7.537, p = 5.7 · 10^{-11}) and quadratic (t_{82} = -4.426, p = 2.9 · 10^{-5}) SNR effects. There was no effect of Age
Group nor an Age Group × SNR interaction (for both p > 0.05). Critically, the reduction in eye
movements associated with SNR conditions is not related to SNR per se. That is, eye movements
remained reduced for the impossible relative to the difficult SNR (t_{82} = 7.601, p_{Holm} = 1.3 · 10^{-10}, d =

0.751) and the easy SNR (t_{82} = 7.537, p_{Holm} = 1.3 · 10^{-10}, d = 0.745) even after a sentence ended (5.3 to
6 s) and no acoustic differences between the three SNR conditions were present, whereas eye
movements did not differ between the easy and the difficult conditions after sentence offset (t_{82} =
0.064, p_{Holm} = 0.949, d = 0.006; effect of SNR: $F_{2,82}$ = 38.193, p = 1.9 × 10^{-12}, ω^2 = 0.110). This suggests
that a cognitive process, rather than acoustic differences, led to reduced eye movements for the
impossible condition and the dissociation from the pattern of results for the pupil size.

The results of Experiment 1 show that pupil size indexes listening effort as expected. That is,
pupil size increased for difficult speech comprehension, but not when comprehension was impossible
(Zekveld and Kramer, 2014; Ohlenforst et al., 2017; Ohlenforst et al., 2018; Wendt et al., 2018). In
contrast, eye movements decreased for both, when comprehension was difficult and when it was
impossible, suggesting different processes underlie changes in pupil size and eye movements during
listening. In Experiment 1, participants were unable to anticipate whether comprehension of an
upcoming sentence will be easy, difficult, or impossible. However, anticipating the difficulty of a task is
a critical component in frameworks that are concerned with motivation and cognitive control (Brehm
and Self, 1989; Richter, 2013; Richter et al., 2016). Anticipating when listening is challenging can help
regulate how much effort is invested. In Experiment 1, listeners may have continued attentive listening
even under impossible conditions, affecting their eye movements, but not arousal-related changes in
pupil size, because they were unable to predict ahead of time whether a sentence could be
understood. Experiment 2 was designed to investigate whether prior knowledge about the
comprehension difficulty changes the influence of SNR on pupil responses and eye movements.

Experiment 2

Younger and older adults participated in Experiment 2. Pure-tone audiometric thresholds were higher
in older compared to younger adults (t_{29} = 4.04, p = 3.6 · 10^{-4}, d = 1.599; Figure 3A,B), as expected.
Participants listened to sentences in background babble at three different SNR levels that made
comprehension easy, difficult, or impossible. Critically, in Experiment 2, a visual cue prior to babble
onset indicated whether comprehension of the upcoming sentence would be easy, difficult, or
impossible. As for Experiment 1, performance decreased with SNR (effect of SNR: $F_{2,80}$ = 646.899, p = 4
· 10^{-50}, ω^2 = 0.908). Older adults exhibited higher performance than younger adults for the difficult SNR
(t_{80} = 2.899, p_{Holm} = 0.013, d = 0.896), but not the easy or impossible SNRs (for both p_{Holm} > 0.2; Age
Group × SNR interaction: $F_{2,80}$ = 5.719, p = 0.005, ω^2 = 0.067; Figure 3C). There was no effect of Age
Group (p = 0.607).

Figure 3: Pure-tone audiometric thresholds and behavioral data for Experiment 2. A: Pure-tone audiometric thresholds: The thin lines reflect data from individual participants. Thick lines reflect the mean across participants. Note that in Experiment 2, audiograms were recorded for all older adults, but only for 9 younger adults. **B:** Pure-tone average hearing threshold (mean across 0.5, 1, 2, and 4 kHz). Gray dots reflect the pure-tone average threshold for individual participants. **C:** Proportion of correct responses in the semantic-relatedness task for younger (left) and older adults (right). Gray dots reflect data from individual participants. The dashed horizontal line at 0.5 reflects chance level. Error bars reflect that standard error of the mean.

Pupil size was modulated by SNR ($F_{2,80}$ = 8.542, p = 4.3 · 10^{-4}, ω^2 = 0.037), such that pupil size was larger under difficult than easy (t_{80} = 2.918, p_{Holm} = 0.009, d = 0.359) and impossible conditions (t_{80} = 3.994, p_{Holm} = 4.3 · 10^{-4}, d = 0.491), whereas the pupil size for easy and impossible SNRs did not differ (t_{80} = 1.077, p_{Holm} = 0.285, d = 0.132). The quadratic (t_{80} = -3.991, p = 1.5 · 10^{-4}) but not the linear (p = 0.285) SNR effect was also significant. Hence, pupillometry showed the expected inverted u-shaped profile, indicating sensitivity to listening effort (Figure 4; Zekveld and Kramer, 2014; Ohlenforst et al., 2017; Ohlenforst et al., 2018; Wendt et al., 2018). There was no age-group difference nor any interaction involving age group (ps > 0.5).

Figure 4: Pupil size and gaze dispersion for Experiment 2. A: Pupil size time courses. The shaded area reflects the standard error of the mean (between-participant variances removed; Masson and Loftus, 2003). The dashed vertical lines indicate, from left to right, the onset of the dot display (stationary), onset of the babble noise and dot movements, sentence onset. **B:** Mean pupil size across the 3.3–5.8 s time window (sentence onset at 2.8 s and average sentence offset at 5.3 s). Error bars reflect the standard error of the mean (removal of between-participant variance; Masson and Loftus, 2003). **C:** Difference in pupil size between different SNR conditions. Violin plots reflect the histogram of individual data points. Dots reflect the mean. **D-F:** The same types of data plots as in panels A-C are depicted for gaze dispersion. Panel E and F reflect mean gaze dispersion across the 2.8-5.3 s time window.

Gaze dispersion also showed the same pattern as in Experiment 1. During sentence
presentation (2.8 to 5.3 s), gaze dispersion decreased with decreasing SNR ($F_{2,80} = 33.267$, $p = 3 \cdot 10^{-11}$,
$\omega^2 = 0.030$), and there was no age-group difference or interaction (ps > 0.05). Specifically, gaze
dispersion was reduced for difficult ($t_{80} = 5.738$, $p_{Holm} = 3.3 \cdot 10^{-7}$, d = 0.303) and impossible conditions
($t_{80} = 7.890$, $p_{Holm} = 4 \cdot 10^{-11}$, d = 0.416) compared to the easy condition, and gaze dispersion was
further reduced for the impossible compared to the difficult condition ($t_{80} = 2.151$, $p_{Holm} = 0.034$, d =
0.114). This was also reflected in the linear ($t_{80} = 8.218$, $p = 3 \cdot 10^{-12}$) and quadratic ($t_{80} = -3.682$, $p = 4.2$
$\cdot 10^{-4}$) SNR effects. Moreover, gaze dispersion remained lower when comprehension was impossible,

compared to both the easy SNR (t_{80} = 8.218, p_{Holm} = 9.1 · 10^{-12}, d = 0.690) and the difficult SNR (t_{80} =
7.298, p_{Holm} = 3.8 · 10^{-10}, d = 0.612), even after sentence offset (5.3–6 s) when no acoustic differences
between conditions were present, whereas eye movements did not differ between easy and difficult
SNRs after sentence offset (p_{Holm} = 0.36; effect of SNR: $F_{2,80}$ = 40.545, p = 6.9 · 10^{-13}, ω^2 = 0.087). This
suggests, similar to Experiment 1, that cognitive processes, rather than acoustic differences, underlie
the pattern of eye movements.
In Experiment 2, participants could anticipate the level of speech comprehension difficulty
prior to sentence onset, but SNRs associated with easy, difficult, and impossible comprehension
randomly varied from trial to trial. This trial-to-trial change in SNR may reduce individuals' ability to
fully make use of anticipatory processes. In some previous pupillometry work that investigated the
relation between SNR and pupil size, trials with different SNRs were presented in separate blocks
(Wendt et al., 2018). This may facilitate the anticipation of upcoming comprehension difficulties and
may better enable individuals to give up listening or ready themselves for sustained effort. Experiment
3 was designed to test whether a blocked presentation of SNRs changes the influence of SNR on eye
movements.

Experiment 3

408
In Experiment 3, younger adults listened to sentences in background babble at three different SNR
levels that made comprehension easy, difficult, or impossible. SNR conditions were blocked such that
12 trials of the same SNR condition occurred in direct succession. Each of the 12-trial blocked
presentations was preceded by written information about the comprehension difficulty of the
upcoming sentences. Performance in the semantic-relatedness task decreased with decreasing SNR
($F_{2,44}$ = 272.306, p = 1.7 · 10^{-25}, ω^2 = 0.865; all post hoc contrasts p_{Holm} < 0.001; Figure 5A).

Figure 5: Behavioral data, pupil area, and gaze dispersion for Experiment 3. A: Proportion of correct responses in the semantic-relatedness task for younger adults. Gray dots reflect data from individual participants. The dashed horizontal line at 0.5 reflects chance level. **B:** Pupil size data. Left: time courses. The shaded area reflects the standard error of the mean (between-participant variances removed; Masson and Loftus, 2003). The dashed

vertical lines indicate, from left to right, the onset of the dot display (stationary), onset of the babble noise and dot movements, sentence onset. Middle: Mean pupil size across the 3.3–5.8 s time window (sentence onset at 2.8 s and average sentence offset at 5.3 s). Error bars reflect the standard error of the mean (removal of between-participant variance; Masson and Loftus, 2003). Right: Difference in pupil size between different SNR conditions. Violin plots reflect the histogram of individual data points. Dots reflect the mean. **C:** The same data plots as in panel B are depicted for gaze dispersion (mean across the 2.8-5.3 s time window).

There was no effect of SNR on pupil size ($F_{2,44}$ = 1.032, p = 0.365, ω^2 < 0.001; Figure 5B), nor a
linear or quadratic SNR effect (both ps > 0.25). In contrast, gaze dispersion was smaller for the difficult
(t_{44} = 3.352, p_{Holm} = 0.003, d = 0.313) and the impossible SNR (t_{44} = 3.392, p_{Holm} = 0.002, d = 0.345)
compared to the easy SNR, whereas the difficult and the impossible SNRs did not differ (p_{Holm} > 0.7;
effect of SNR: $F_{2,44}$ = 8.327, p = 8.6 · 10^{-4}, ω^2 = 0.021; Figure 5C). This was also reflected in the linear
SNR effect (t_{44} = 3.939, p = 2.9 · 10^{-4}; quadratic: p = 0.088). As for Experiments 1 and 2, the reduction in
gaze dispersion cannot be due to acoustic differences, because gaze dispersion remained reduced for
the impossible SNR compared to both the easy (t_{44} = 3.939, p_{Holm} = 8.7 · 10^{-4}, d = 0.610) and the
difficult SNR (t_{44} = 3.481, p_{Holm} = 0.002, d = 0.539) even after sentence offset (5.3–6 s; effect of SNR:
$F_{2,44}$ = 9.281, p = 4.3 · 10^{-4}, ω^2 = 0.062). Gaze dispersion did not differ between the difficult and the
easy SNR conditions after sentence offset (p_{Holm} > 0.6). These results replicate the results from
Experiments 1 and 2 and suggest that eye movements are influenced differently by listening challenges
than pupil size.

428   **Discussion**

Pupillometry is commonly used to assess effort, but it has disadvantages. Here, we investigated the
degree to which a new measure of effort – that is, eye movements – reflect effort across three
listening experiments. Younger and older adults listened to speech masked by background babble at
easy, difficult, and impossible speech comprehension levels. We expected a u-shaped pattern in pupil
size and gaze dispersion indicative of high effort for difficult comprehension, but low for easy and
impossible speech comprehension. In the latter case, we predicted that individuals would give up
investing effort when comprehension was impossible. The pupil-size data showed the expected u-
shaped profile, whereas gaze dispersion decreased with increasing speech masking. The decrease in
gaze dispersion for the impossible speech comprehension was independent of stimulus acoustics, thus
suggesting that gaze dispersion is governed by a different neuro-cognitive mechanism than pupil size.

Pupil size reflects effort
Measurements of pupil size are frequently used to assess effort during task engagement, such as
during listening to masked or degraded speech (Zekveld et al., 2010; Zekveld et al., 2014; Zekveld and
Kramer, 2014; Wendt et al., 2016; Koelewijn et al., 2018; Winn and Moore, 2018; Winn et al., 2018;
Ayasse and Wingfield, 2020; Kadem et al., 2020). Our results are consistent with this previous work,
showing that the pupil size is largest under difficult but manageable speech-comprehension

conditions, whereas it is smaller when speech comprehension is easy or impossible (Zekveld and Kramer, 2014; Ohlenforst et al., 2017; Ohlenforst et al., 2018; Wendt et al., 2018). Although the current pupil-size results show the u-shaped profile, this profile appears a bit variable; for example, it was not significant for younger adults in Experiment 3 (note that pupil-size variability generally appeared larger in younger than older adults; error bars in Figures 2, 4, 5). The variability in the u-shaped profile of the pupil size may be partly related to the moving dot display that was presented during sound presentation to facilitate sensitivity to changes in eye movements; pupillometry is typically recorded while individuals maintain fixation on a point on a computer monitor during speech listening (Zekveld et al., 2014; Zhao et al., 2019; Farahani et al., 2020; Winn and Teece, 2021), but this can alter speech processes related to mental imagery and comprehension (Johansson et al., 2012; Cui and Herrmann, 2023). Nonetheless, pupil size was associated with cognitive investment in the current study, reaffirming the utility of this marker in assessing listening effort.

Eye movements reflect listening challenges differently than the pupil size

Recent works suggest that reductions in eye movements may index effort. Eye movements decrease when individuals perform a high-load compared to a low-load memory task (Lipton et al., 1980; Hutton and Tegally, 2005; Kosch et al., 2018; Walter and Bex, 2021) and when speech masking increases (Contadini-Wright et al., 2023; Cui and Herrmann, 2023). We also found that gaze dispersion decreased with decreasing SNR (Figures 2, 4, and 5), and extend this previous work by showing that the sensitivity of eye movements to changes in speech masking does not differ between younger and older adults. The absence of an age-group difference suggests that measuring gaze dispersion may be useful across different populations.

Previous studies that investigated changes in eye movements related to effort only included easy and difficult task conditions, but a signature of effort must also be sensitive to disengagement that occurs under conditions that exceed a person's mental resources. We thus predicted a u-shaped effort profile when the range of easy to impossible listening conditions were used (Yerkes and Dodson, 1908; Zekveld and Kramer, 2014; Ohlenforst et al., 2017; Ohlenforst et al., 2018; Wendt et al., 2018). The results of the current study provide no indication of a u-shaped profile for gaze dispersion (Figures 2, 4, and 5). Instead, gaze dispersion decreased with decreasing SNR. On first glance, this may indicate that changes in eye movements are coupled to changes in the acoustic properties of the auditory environment rather than to effort per se. That is, sound intensity during sentence presentation increased with increasing SNR, because the babble was played at the same level across trials to which a sentence was added at different levels to create the three SNRs (cf. Ohlenforst et al., 2017; Kadem et al., 2020; Cui and Herrmann, 2023). Critically, acoustic differences ceased at sentence offset, after which only the babble continued. Gaze dispersion did not differ between easy and difficult speech-comprehension conditions after sentence offset, suggesting that, in the difficult condition, individuals stopped investing cognitively after a sentence ended. Yet, gaze dispersion during impossible speech comprehension remained reduced after sentence offset relative to easy and difficult comprehension. Acoustic differences thus do not appear to drive the reduction in the spatial extent of eye movements. Likewise, prior knowledge about the upcoming challenge in speech comprehension did not affect this

pattern of results. The difference in the data patterns for pupil size and gaze dispersion may therefore
reflect the underlying influence of distinct cognitive processes during challenging listening conditions.

The role of neurophysiological mechanisms in the different changes in pupil size and eye movements
A wide network of brain regions underlies oculomotor functions, including the execution of eye
movements and pupil dilation/constriction. Brain regions associated with eye movements include the
superior colliculus, thalamus, basal ganglia, cerebellum, visual cortex, frontal and supplementary eye
fields, anterior cingulate cortex, posterior parietal cortex, and the prefrontal cortex (Sparks, 2002;
Pierrot-Deseilligny et al., 2004; Lencer and Trillenberg, 2008; Pierce et al., 2019). Pupil size is regulated
by some of these same brain regions (Wang et al., 2012; Joshi and Gold, 2020; Wang and Munoz,
2021; Burlingham et al., 2022), but is also driven by regions that do not appear to play a prominent
role in regulating eye movements, such as the locus coeruleus (Mathôt, 2018; Joshi and Gold, 2020;
Strauch et al., 2022).
Critically, changes in pupil size have been linked to changes in arousal associated with locus
coeruleus function (Bradley et al., 2008; Mathôt, 2018; Wang et al., 2018; Ross and Van Bockstaele,
2021; Burlingham et al., 2022). Here, too, the arousal system may underlie the observed pattern of
pupil responses. In Experiment 1, in which the comprehension difficulty could not be predicted for a
given trial, high SNR sentences elicited a transient pupil response (despite being highly intelligible;
Figure 2). This transient pupil response was less prominent in Experiment 2 when individuals could
predict, and thus were less surprised by, the SNR of a sentence (Figure 4). Moreover, there was a
transient increase in pupil size following the visual cue that indexed the impossible SNR condition,
possibly reflecting surprisal which, in turn, elicited arousal (Figure 4).
Eye movements can be automatically triggered or volitionally initiated (Pierce et al., 2019). Eye-
movement reductions with increasing task load may include involuntary eye movements, as suggested
by observations of task-difficulty related eye-movement reductions during fixation on a point on a
computer monitor (Contadini-Wright et al., 2023; Cui and Herrmann, 2023). However, individuals may
also voluntarily reduce eye movements under high cognitive demands, for example, to free resources
associated with eye movements and to avoid processing irrelevant information. A voluntary reduction
in eye movements may involve cognitive control regions, such as the prefrontal, cingulate, and parietal
cortices (Cole and Schneider, 2007; Braver, 2012; Niendam et al., 2012), that drive structures
associated with eye movement initiation, such as the frontal and supplementary eye fields and the
superior colliculus (Pierce et al., 2019). In the current study, individuals may have tried to free
resources to gather relevant speech information by reducing eye movements during unfavorable SNRs,
independently of the engagement of arousal or other pupil-related systems. Individuals appeared to
have detected when relevant speech information discontinued, because eye movements in the
difficult condition increased to similar levels as in the easy condition after sentence offset (Figures 2, 3,
and 5). For the impossible condition, listeners were unable to hear when a sentence ended, possibly
leading them to continue to reduce eye movements throughout the trial. We would predict that in a
study in which the babble noise continued for longer following sentence offset, visual exploration in
the impossible condition would eventually increase and be similar to the easy condition. We thus

speculate that the pupil size indicates effort through the involvement of arousal systems, whereas
changes in the spatial dispersion of eye movements reflect the influence of cognitive processes related
to mental effort.

526   Conclusions
The current study investigated the degree to which changes in eye movements reflect mental effort
associated with listening in younger and older adults. Individuals listened to speech masked by
background babble at signal-to-noise ratios associated with easy, difficult, and impossible speech
comprehension. We expected individuals would invest little effort for easy and impossible speech
('giving up') but that they would exert higher effort for difficult speech. Consistent with this effort
profile, a u-shaped pattern was observed for the pupil size, such that pupil size was largest for difficult
comprehension, but lower for easy and impossible comprehension. In contrast, gaze dispersion
decreased with increasing speech masking. However, gaze dispersion during difficult comprehension
returned to levels similar to easy comprehension after sentence offset, during which acoustic
stimulation was similar across conditions, whereas the spatial dispersion of eye movements during
impossible comprehension continued to decrease. This speaks against acoustic factors driving reduced
eye movements. Our data instead suggest that a cognitive process, different from the arousal-related
systems regulating the pupil size, is involved in reduced eye movements during high cognitive load.
The current study highlights how effort in one sensory modality (audition) impacts the functional
properties in another sensory modality (vision).

542   Acknowledgements
543   The research was supported by the Canada Research Chair program (CRC-2019-00156, 232733), the
544   Natural Sciences and Engineering Research Council of Canada (NSERC Discovery Grant: RGPIN-2021-
545   02602), the Canadian Institutes of Health Research (CIHR: 186236), and the William Demant
546   Foundation. We thank Tazeen Atif and Priya Pandey for help during data collection. We thank Ingrid
547   Johnsrude and Lauren Fink for fruitful discussions.

548   Author Contributions
549   **BH:** Conceptualization, methodology, validation, formal analysis, investigation, data curation, writing -
550   original draft, writing - review & editing, visualization, supervision, project management, funding
551   acquisition. **JDR:** Conceptualization, methodology, investigation, writing - review & editing.

552   Statements and Declarations
553   The author has no conflicts or competing interests.

554   References

Abrams RA, Pratt J, Chasteen AL (1998) Aging and movement: Variability of force pulses for saccadic
eye movements. Psychology and Aging 13:387-395.

Alvarez GA, Franconeri SL (2007) How many objects can you track?: Evidence for a resource-limited
attentive tracking mechanism. Journal of Vision 7:1-10.

Ayasse ND, Wingfield A (2018) A Tipping Point in Listening Effort: Effects of Linguistic Complexity and
Age-Related Hearing Loss on Sentence Comprehension. Trends in Hearing 22:2331216518790907.

Ayasse ND, Wingfield A (2020) Anticipatory Baseline Pupil Diameter Is Sensitive to Differences in
Hearing Thresholds. Frontiers in Psychology 10.

Bilger RC (1984) Manual for the clinical use of the revised SPIN Test. Champaign, IL, USA: The
University of Illinois.

Bradley MM, Miccoli L, Escrig MA, Lang PJ (2008) The pupil as a measure of emotional arousal and
autonomic activation. Psychophysiology 45:602-607.

Braver TS (2012) The variable nature of cognitive control: a dual mechanisms framework. Trends in
Cognitive Sciences 16:106-113.

Braver TS et al. (2014) Mechanisms of motivation-cognition interaction: challenges and opportunities.
Cogn Affect Behav Neurosci 14:443-472.

Brehm JW, Self EA (1989) The Intensity of Motivation. Annual Review of Psychology 40:109-131.

Brisson J, Mainville M, Mailloux D, Beaulieu C, Serres J, Sirois S (2013) Pupil diameter measurement
errors as a function of gaze direction in corneal reflection eyetrackers. Behavior Research
Methods 45:1322-1331.

Bruenech JR (2018) Age-Related Changes in the Oculomotor System. In: Age-Related Changes of the
Human Eye (Cavallotti CAP, Cerulli L, eds), pp 343-373: Humana Press.

Burlingham CS, Mirbagheri S, Heeger DJ (2022) A unified model of the task-evoked pupil response.
Science Advances 8:eabi9979.

Cavanagh P, Alvarez GA (2005) Tracking multiple targets with multifocal attention. Trends in Cognitive
Sciences 9:349-354.

Cole MW, Schneider W (2007) The cognitive control network: Integrated cortical regions with
dissociable functions. Neuroimage 37:343-360.

Contadini-Wright C, Magami K, Mehta N, Chait M (2023) Pupil dilation and microsaccades provide
complementary insights into the dynamics of arousal and instantaneous attention during effortful
listening. The Journal of Neuroscience 43:4856-4866.

Cui ME, Herrmann B (2023) Eye Movements Decrease during Effortful Speech Listening. The Journal of
Neuroscience 43:5856-5869.

Eckert MA, Teubner-Rhodes S, Vaden Jr. KI (2016) Is Listening in Noise Worth It? The Neurobiology of
Speech Recognition in Challenging Listening Conditions. Ear & Hearing 37:101S-110S.

Engbert R (2006) A microcosm for research on oculomotor control, attention, and visual perception.
Progress in Brain Research 154:177-192.

Engbert R, Kliegl R (2003) Microsaccades uncover the orientation of covert attention. Vision Research
43:1035-1045.

Farahani M, Parsa V, Herrmann B, Kadem M, Johnsrude IS, Doyle PC (2020) An Auditory-Perceptual
and Pupillometric Study of Vocal Strain and Listening Eort in Adductor Spasmodic Dysphonia.
Applied Sciences 10:5907.
Fink L, Simola J, Tavano A, Lange E, Wallot S, Laeng B (2021) From pre-processing to advanced dynamic
modeling of pupil data. PsyArXiv.
Francis AL, Love J (2020) Listening effort: Are we measuring cognition or affect, or both? WIREs
Cognitive Science 11:e1514.
Guillon M, Dumbleton K, Theodoratos P, Gobbe M, Wooley CB, Moody K (2016) The Effects of Age,
Refractive Status, and Luminance on Pupil Size. Optometry and Vision Science 93:1093-1100.
Gustafsson HÅ, Arlinger SD (1994) Masking of speech by amplitude-modulated noise. The Journal of
the Acoustical Society of America 95:518-529.
Hayes TR, Petrov AA (2016) Mapping and correcting the influence of gaze position on pupil size
measurements. Behavior Research Methods 48:510-527.
Helfer KS, Freyman RL (2008) Aging and Speech-on-Speech Masking. Ear & Hearing 29:87-98.
Herrmann B (2023) The perception of artificial-intelligence (AI) based synthesized speech in younger
and older adults. International Journal of Speech Technology 26:395-415.
Herrmann B, Johnsrude IS (2018a) Attentional State Modulates the Effect of an Irrelevant Stimulus
Dimension on Perception. Journal of Experimental Psychology: Human Perception and
Performance 44:89-105.
Herrmann B, Johnsrude IS (2018b) Neural signatures of the processing of temporal patterns in sound.
The Journal of Neuroscience 38:5466-5477.
Herrmann B, Johnsrude IS (2020) A Model of Listening Engagement (MoLE). Hearing Research
397:108016.
Herrmann B, Maess B, Johnsrude IS (2018) Aging Affects Adaptation to Sound-Level Statistics in Human
Auditory Cortex. The Journal of Neuroscience 38:1989-1999.
Herrmann B, Maess B, Johnsrude IS (2023) Age-related changes in neural synchronization and the
sustained response to amplitude and frequency modulation of sound. Hearing Research
428:108677.
Hockey GRJ (1997) Compensatory control in the regulation of human performance under stress and
high workload: A cognitive-energetical framework. Biological Psychology 45:73-93.
Holm S (1979) A Simple Sequentially Rejective Multiple Test Procedure. Scandinavian Journal of
Statistics 6:65-70.
Hutton SB, Tegally D (2005) The effects of dividing attention on smooth pursuit eye tracking.
Experimental Brain Research 163:306-313.
IEEE (1969) IEEE Recommended Practice for Speech Quality Measurements. IEEE Transactions on
Audio and Electroacoustics 17:225-246.
Irsik VC, Johnsrude IS, Herrmann B (2022) Age-related deficits in dip-listening evident for isolated
sentences but not for spoken stories. Scientific Reports 12:5898.
JASP (2023) JASP [Computer software]. In: https://jasp-stats.org/.

Johansson R, Holsanova J, Holmqvist K (2006) Pictures and Spoken Descriptions Elicit Similar Eye Movements During Mental Imagery, Both in Light and in Complete Darkness. Cognitive Science 30:1053-1079.

Johansson R, Holsanova J, Homqvist K (2011) The Dispersion of Eye Movements During Visual Imagery is Related to Individual Differences in Spatial Imagery Ability. In: Annual Meeting of the Cognitive Science Society, pp 1200-1205: Proceedings of the Annual Meeting of the Cognitive Science Society.

Johansson R, Holsanova J, Dewhurst R, Holmqvist K (2012) Eye movements during scene recollection have a functional role, but they are not reinstatements of those produced during encoding. Journal of Experimental Psychology: Human Perception and Performance 38:1289-1314.

Joshi S, Gold JI (2020) Pupil Size as a Window on Neural Substrates of Cognition. Trends in Cognitive Sciences 24:466-480.

Kadem M, Herrmann B, Rodd JM, Johnsrude IS (2020) Pupil dilation is sensitive to semantic ambiguity and noise. Trends in Hearing 24:1-16.

Knapen T, de Gee JW, Brascamp J, Nuiten S, Hoppenbrouwers S, Theeuwes J (2016) Cognitive and Ocular Factors Jointly Determine Pupil Responses under Equiluminance. PLoS ONE 11:e0155574.

Koelewijn T, Zekveld AA, Lunner T, Kramer SE (2018) The effect of reward on listening effort as reflected by the pupil dilation response. Hearing Research 367:106-112.

Kool W, Botvinick M (2018) Mental labour. Nature Human Behavior 2:899-908.

Kosch T, Hassib M, Woźniak PW, Buschek D, Alt F (2018) Your Eyes Tell: Leveraging Smooth Pursuit for Assessing Cognitive Workload. In: CHI Conference on Human Factors in Computing Systems, p Paper 436. Montreal, QC, Canada.

Kraus F, Obleser J, Herrmann B (2023a) Pupil-size sensitivity to listening demand depends on motivational state. eNeuro.

Kraus F, Tune S, Obleser J, Herrmann B (2023b) Neural alpha oscillations and pupil size differentially index cognitive demand under competing audio-visual task conditions. The Journal of Neuroscience 43:4352-4364.

Kuchinsky SE, Ahlstrom JB, Vaden Jr. KI, Cute SL, Humes LE, Dubno JR, Eckert MA (2013) Pupil size varies with word listening and response selection difficulty in older adults with hearing loss. Psychophysiology 50:23-34.

Lencer R, Trillenberg P (2008) Neurophysiology and neuroanatomy of smooth pursuit in humans. Brain and Cognition 68:219-228.

Lipton RB, Frost LA, Holzman PS (1980) Smooth Pursuit Eye Movements, Schizophrenia, and Distraction. Perceptual and Motor Skills 50:159-167.

Liu Z-X, Shen K, Olsen RK, Ryan JD (2018) Age-related changes in the relationship between visual exploration and hippocampal activity. Neuropsychologia 119:81-91.

Masson MEJ, Loftus GR (2003) Using confidence intervals for graphically based data interpretation. Canadian Journal of Experimental Psychology and Aging 57:203-220.

Mathôt S (2018) Pupillometry: Psychology, physiology, and function. Journal of Cognition 1.

Mazloum-Farzaghi N, Shing N, Mendoza L, Barense MD, Ryan JD, Olsen RK (2022) The impact of aging and repetition on eye movements and recognition memory. Aging, Neuropsychology, and Cognition:1-27.

McCloy DR, Panfili L, John C, Winn M, Wright RA (2018) Gender, the individual, and intelligibility. In: 176th Meeting of the Acoustical Society of America. Victoria, BC, Canada.

McGarrigle R, Munro KJ, Dawes P, Stewart AJ, Moore DR, Barry JG, Amitay S (2014) Listening effort and fatigue: What exactly are we measuring? A British Society of Audiology Cognition in Hearing Special Interest Group 'white paper'. International Journal of Audiology 53:433-440.

McIntire LK, McIntire JP, McKinley RA, Goodyear C (2014) Detection of vigilance performance with pupillometry. In: Symposium on Eye Tracking Research and Applications, pp 167-174. Safety Harbor, Florida, USA: ACM.

Moore BCJ (2007) Cochlear Hearing Loss: Physiological, Psychological and Technical Issues. West Sussex, Engand: John Wiley & Sons, Ltd.

Moran A, Quinn A, Campbell M, Rooney B, Brady N, Burke C (2016) Using pupillometry to evaluate attentional effort in quiet eye: A preliminary investigation. Sport, Exercise, and Performance Psychology 5:365-376.

Niendam TA, Laird AR, Ray KL, Dean YM, Glahn DC, Carter CS (2012) Meta-analytic evidence for a superordinate cognitive control network subserving diverse executive functions. Cognitive, Affective, & Behavioral Neuroscience 12:241-268.

Ohlenforst B, Wendt D, Kramer SE, Naylor G, Zekveld AA, Lunner T (2018) Impact of SNR, masker type and noise reduction processing on sentence recognition performance and listening effort as indicated by the pupil dilation response. Hearing Research.

Ohlenforst B, Zekveld AA, Lunner T, Wendt D, Naylor G, Wang Y, Versfeld NJ, Kramer SE (2017) Impact of stimulus-related factors and hearing impairment on listening effort as indicated by pupil dilation. Hearing Research 351:68-79.

Panfili LM, Haywood J, McCloy DR, Souza PE, Wright RA (2017) The UW/NU Corpus, Version 2.0. In: https://depts.washington.edu/phonlab/projects/uwnu.php.

Peelle JE (2018) Listening Effort: How the Cognitive Consequences of Acoustic Challenge Are Reflected in Brain and Behavior. Ear & Hearing 39:204-214.

Pichora-Fuller MK, Kramer SE, Eckert MA, Edwards B, Hornsby BWY, Humes LE, Lemke U, Lunner T, Matthen M, Mackersie CL, Naylor G, Phillips NA, Richter M, Rudner M, Sommers MS, Tremblay KL, Wingfield A (2016) Hearing Impairment and Cognitive Energy: The Framework for Understanding Effortful Listening (FUEL). Ear & Hearing 37 Suppl 1:5S-27S.

Pierce JE, Clementz BA, McDowell JE (2019) Saccades: Fundamentals and Neural Mechanisms. In: Eye Movement Research: An Introduction to its Scientific Foundations and Applications (Klein C, Ettinger U, eds), pp 11-71. Cham: Springer International Publishing.

Pierrot-Deseilligny C, Milea D, Müri RM (2004) Eye movement control by the cerebral cortex. Current Opinion in Neurology 17.

Piquado T, Isaacowitz D, Wingfield A (2010) Pupillometry as a measure of cognitive effort in younger and older adults. Psychophysiology 47:560-569.

Plack CJ (2014) The sense of hearing. New York, USA: Psychology Press.

Presacco A, Simon JZ, Anderson S (2016) Evidence of degraded representation of speech in noise, in
the aging midbrain and cortex. Journal of Neurophysiology 116:2346-2355.
Presacco A, Simon JZ, Anderson S (2019) Speech-in-noise representation in the aging midbrain and
cortex: Effects of hearing loss. PLoS ONE 14:e0213899.
Richter M (2013) A Closer Look Into the Multi-Layer Structure of Motivational Intensity Theory. Social
and Personality Psychology Compass 7:1-12.
Richter M, Gendolla GHE, Wright RA (2016) Three Decades of Research on Motivational Intensity
Theory: What We Have Learned About Effort and What We Still Don't Know. In: Advances in
Motivation Science (Elliot AJ, ed), pp 149-186. Cambridge, MA, USA: Academic Press.
Rodd JM, Johnsrude IS, Davis MH (2010a) The role of domain-general frontal systems in language
comprehension: Evidence from dual-task interference and semantic ambiguity. Brain and
Language 115:182-188.
Rodd JM, Longe OA, Randall B, Tyler LK (2010b) The functional organisation of the fronto-temporal
language system: Evidence from syntactic and semantic ambiguity. Neuropsychologia 448:1324-
1335.
Ross JA, Van Bockstaele EJ (2021) The Locus Coeruleus- Norepinephrine System in Stress and Arousal:
Unraveling Historical, Current, and Future Perspectives. Frontiers in Psychiatry 11.
Ryan JD, Leung G, Turk-Browne NB, Hasher L (2007) Assessment of age-related changes in inhibition
and binding using eye movement monitoring. Psychology and Aging 22:239-250.
Scholl BJ (2009) What have we learned about attention from multiple object tracking (and vice versa)?
In: Computation, cognition, and Pylyshyn (Dedrick D, Trick L, eds), pp 49-78. Cambridge, MA, USA:
MIT Press.
Sharpe JA, Sylvester TO (1978) Effect of aging on horizontal smooth pursuit. Investigative
Ophthalmology & Visual Science 17:465-468.
Shechter A, Share DL (2020) Keeping an Eye on Effort: A Pupillometric Investigation of Effort and
Effortlessness in Visual Word Recognition. Psychological Science 32:80-95.
Slade K, Kramer SE, Fairclough S, Richter M (2021) Effortful listening: Sympathetic activity varies as a
function of listening demand but parasympathetic activity does not. Hearing Research
410:108348.
Sparks DL (2002) The brainstem control of saccadic eye movements. Nature Reviews Neuroscience
3:952-964.
Spooner JW, Sakala SM, Baloh RW (1980) Effect of Aging on Eye Tracking. Archives of Neurology
37:575-576.
Strauch C, Wang C-A, Einhäuser W, Van der Stigchel S, Naber M (2022) Pupillometry as an integrated
readout of distinct attentional networks. Trends in Neurosciences 45:635-647.
Suzuki Y, Minami T, Laeng B, Nakauchi S (2019) Colorful glares: Effects of colors on brightness illusions
measured with pupillometry. Acta Psychologica 198:102882.
Thurman SM, Cohen Hoffing RA, Madison A, Ries AJ, Gordon SM, Touryan J (2021) "Blue Sky Effect":
Contextual Influences on Pupil Size During Naturalistic Visual Search. Frontiers in Psychology 12.
van der Wel P, van Steenbergen H (2018) Pupil dilation as an index of effort in cognitive control tasks:
A review. Psychonomic Bulletin & Review 25:2005-2015.

Walter K, Bex P (2021) Cognitive load influences oculomotor behavior in natural scenes. Scientific Reports 11:12405.

Wang C-A, Munoz DP (2021) Coordination of Pupil and Saccade Responses by the Superior Colliculus. Journal of Cognitive Neuroscience 33:919-932.

Wang C-A, Boehnke SE, White BJ, Munoz DP (2012) Microstimulation of the Monkey Superior Colliculus Induces Pupil Dilation Without Evoking Saccades. The Journal of Neuroscience 32:3629.

Wang C-A, Baird T, Huang J, Coutinho JD, Brien DC, Munoz DP (2018) Arousal Effects on Pupil Size, Heart Rate, and Skin Conductance in an Emotional Face Task. Frontiers in Neurology 9.

Wendt D, Dau T, Hjortkjær J (2016) Impact of Background Noise and Sentence Complexity on Processing Demands during Sentence Comprehension. Frontiers in Psychology 7:Article 345.

Wendt D, Koelewijn T, Książek P, Kramer SE, Lunner T (2018) Toward a more comprehensive understanding of the impact of masker type and signal-to-noise ratio on the pupillary response while performing a speech-in-noise test. Hearing Research 369:67-78.

Westbrook A, Braver TS (2015) Cognitive effort: A neuroeconomic approach. Cognitive, Affective & Behavioral Neuroscience 15:395-415.

Westgate EC, Wilson TD (2018) Boring Thoughts and Bored Minds: The MAC Model of Boredom and Cognitive Engagement. Psychological Review 125:689-713.

Widmann A, Engbert R, Schröger E (2014) Microsaccadic Responses Indicate Fast Categorization of Sounds: A Novel Approach to Study Auditory Cognition. The Journal of Neuroscience 34:11152-11158.

Winn MB, Moore AN (2018) Pupillometry Reveals That Context Benefit in Speech Perception Can Be Disrupted by Later-Occurring Sounds, Especially in Listeners With Cochlear Implants. Trends in Hearing 22:2331216518808962.

Winn MB, Teece KH (2021) Listening Effort Is Not the Same as Speech Intelligibility Score. Trends in Hearing 25:1-26.

Winn MB, Wendt D, Koelewijn T, Kuchinsky SE (2018) Best Practices and Advice for Using Pupillometry to Measure Listening Effort: An Introduction for Those Who Want to Get Started. Trends in Hearing 22:2331216518800869.

Yerkes RM, Dodson JD (1908) The relation of strength of stimulus to rapidity of habit-formation. Journal of Comparative Neurology and Psychology 18:459-482.

Zackon DH, Sharpe JA (1987) Smooth pursuit in senescence: Effects of Target Acceleration and Velocity. Acta Oto-Laryngologica 104:290-297.

Zekveld AA, Kramer SE (2014) Cognitive processing load across a wide range of listening conditions: Insights from pupillometry. Psychophysiology 51:277-284.

Zekveld AA, Kramer SE, Festen JM (2010) Pupil response as an indication of effortful listening: the influence of sentence intelligibility. Ear & Hearing 31.

Zekveld AA, Koelewijn T, Kramer SE (2018) The Pupil Dilation Response to Auditory Stimuli: Current State of Knowledge. Trends in Hearing 22:1-25.

Zekveld AA, Heslenfeld DJ, Johnsrude IS, Versfeld NJ, Kramer SE (2014) The eye as a window to the listening brain: Neural correlates of pupil size as a measure of cognitive listening load. NeuroImage 101:76-86.

795   Zhang Y, Malaval F, Lehmann A, Deroche MLD (2022) Luminance effects on pupil dilation in speech-in-
796   noise recognition. PLOS ONE 17:e0278506.
797   Zhao S, Bury G, Milne A, Chait M (2019) Pupillometry as an Objective Measure of Sustained Attention
798   in Young and Older Listeners. Trends in Hearing 23:1-21.
799