Neonatal sensitivity to vocal emotions: A milestone at 37 weeks of gestational age

Xinlin Hou[1]#, Peng Zhang[1,2]#, Licheng Mo[3], Cheng Peng[1], Dandan Zhang[3,4,5,6]*

Department of Pediatrics, Peking University First Hospital, Beijing 100034, China

Department of Pediatrics, Miyun Country Maternal and Child Health Hospital, Beijing 101599, China

Institute of Brain and Psychological Sciences, Sichuan Normal University, Chengdu 610066, China

Shenzhen-Hong Kong Institute of Brain Science, Shenzhen, 518060, China

China Center for Behavioral Economics and Finance, Southwestern University of Finance and Economics, Chengdu 611130, China

School of Psychology, Chengdu Medical College, Chengdu 610500, China

* Corresponding author. Jing'an Road #5, Jinjiang District, Chengdu, 610066, China. Tel/Fax: 86-28-84760580. E-mail: zhangdd05@gmail.com

# Both authors contributed equally to the study

Running title: Early emotional milestone in neonates at 37 weeks GA

Conflict of interest disclosure. The authors declare that there are no conflicts of interest in relation to this study.

Emails:
Xinlin Hou - houxinlin66@sina.com,
Peng Zhang - jupiterzp@163.com,
Licheng Mo - molicheng1021@163.com,
Cheng Peng - pch_20082008@outlook.com,
Dandan Zhang - zhangdd05@gmail.com

Subject consent statement. Prior to data collection, written consent was obtained from the parents or legal guardians of all participating neonates for access to clinical information and EEG data collection for scientific purposes.

Funding statement. This study was funded by the National High Level Hospital Clinical Research Funding (High Quality Clinical Research Project of Peking University First Hospital, 2022CR68), the National Natural Science Foundation of China (32271102; 31920103009), the Major Project of National Social Science Foundation (20&ZD153), Shenzhen-Hong Kong Institute of Brain Science (2024SHIBS0004), and the National Key Research and Development Program of China (2021YFC2700700).

Abstract

Emotional responsiveness in neonates, particularly their ability to discern vocal emotions, plays an evolutionarily adaptive role in human communication and adaptive behaviors. The developmental trajectory of emotional sensitivity in neonates is a crucial area of inquiry for understanding the foundations of early social-emotional functioning. However, the precise onset of this sensitivity in neonates and its relationship with gestational age (GA) remain subjects of investigation. In a study involving 120 healthy neonates categorized into six groups based on their GA (ranging from 35 and 40 weeks), we delved into their emotional responses to vocal stimuli. These stimuli encompassed disyllables with happy and neutral prosodies, alongside acoustically matched nonvocal control sounds. The assessments occurred during natural sleep states in neonates, utilizing the odd-ball paradigm and event-related potentials. The results unveil a distinct developmental milestone at 37 weeks GA, marking the point at which neonates exhibit heightened perceptual acuity for emotional vocal expressions. This newfound ability is substantiated by the presence of the mismatch response, akin to an initial form of adult mismatch negativity, elicited in response to positive emotional vocal prosody. Notably, this perceptual shift's specificity becomes evident when no such discrimination is observed in acoustically matched control sounds. Neonates born before 37 weeks GA do not display this level of discrimination ability. This critical developmental milestone carries significant implications for our understanding of early social-emotional development, shedding light on the role of gestational age in shaping early perceptual abilities. Moreover, it introduces the potential for a valuable screening tool in the context of autism, which is characterized by atypical social-emotional functions. This study makes a substantial contribution to the broader field of developmental neuroscience and holds promise for early intervention in neurodevelopmental disorders.

Keywords: vocal emotion, neonate, preterm, gestational age, mismatch negativity

Significance statement

This study illuminates a key developmental milestone, pinpointing the emergence of heightened emotional perceptual acuity at 37 weeks of gestational age. Employing rigorous methods, we reveal that neonates at this stage exhibit remarkable discrimination abilities for emotional vocal prosody, a vital turning point in early social-emotional functioning. These findings emphasize the pivotal role of gestational age in shaping neonatal perception and provides a pathway for early screening of neurodevelopmental disorders, particularly autism. This insight holds profound implications for understanding the foundations of early social-emotional development in humans, offering a potential tool for early intervention in neurodevelopmental disorders, thereby enhancing child health and well-being.

Introduction

Emotions represent a fundamental aspect of human social interaction, serving as a compelling subject of inquiry within the disciplines of neuroscience, psychology, and psychiatry. Over the course of evolution, the human brain has evolved to possess a heightened sensitivity to the emotional expressions of others (Lindquist et al., 2012). Remarkably, even prior to the full maturation of their visual system, human infants exhibit a remarkable ability to discern vocal emotions (Blasi et al., 2011; Soderstrom et al., 2017; Vaish & Striano, 2004). Prosodic elements of speech, including pitch, intensity, and rhythm, function as universal and non-linguistic channels for emotional communication (Latinus & Belin, 2011). Numerous studies have established that infants, including those who have not yet acquired language, exhibit differentiated responses to emotional prosody conveying happiness, fear, anger, and sadness within the age range of 2 to 12 months (e.g., Caron et al., 1988; Fernald, 1993; Graham et al., 2013; Grossmann et al., 2010; Singh et al., 2002; Walker-Andrews & Grolnick, 1983; Zhao et al., 2021).

More specifically, during the very early stages of postnatal life, often termed the neonatal period (encompassing infants under four weeks of age), compelling evidence points to the presence of emotion-specific responses to emotional cues conveyed through vocal prosody. These responses have been identified through various measurement methods, including assessments of eye-opening scores (Mastropieri & Turkewitz, 1999), event-related potentials (Cheng et al., 2012), and near-infrared spectroscopy (Zhang et al., 2019). However, prior research has primarily focused on the perception and discrimination of emotions among traditionally defined term neonates, a group that includes infants born within a five-week span (37 to 41 weeks) of gestational age (GA), treating them as a homogenous cohort. This raises a crucial question: when does emotional sensitivity begin to manifest in newborns? Does it exist in preterm neonates (GA < 37 weeks)? And does it vary among neonates born at early term (GA = 37-38 weeks) and full term (GA = 39-40 weeks), as defined by the refined 'term' classification (Spong, 2013)? Surprisingly, to date, no study has explored emotion processing in neonates with varying GAs. The discovery of this developmental milestone not only advances our understanding of the cognitive mechanisms underlying human social-emotional functioning but also provides valuable insights for early diagnosis of neurodevelopmental disorders, such as autism (Jones et al., 2014; Molnar-Szakacs et al., 2021).

The principal objective of this study is to investigate emotional responses in neonates across a range of GAs, spanning from 35 to 40 weeks, and to determine whether their heightened sensitivity to emotional voices is influenced by GA. To achieve this, we utilized the odd-ball paradigm in conjunction with an event-related potential (ERP) component known as mismatch negativity (MMN) to probe the neurobiological encoding of emotional voices in the neonatal brain. MMN is an auditory ERP component that demonstrates a negative shift in response to deviant sounds when compared to standard sounds (Näätänen et al., 2007). Importantly, it can be elicited without requiring the subject's attention, making it particularly suitable for recording in young infants (Cheour et al., 1998, 2002). It is worth noting that in neonates, this ERP component often manifests as a positive response rather than the traditional MMN (e.g., Cheng et al., 2012; Chládková et al., 2021; Kostilainen et al., 2020; Richard et al., 2022; Thiede et al., 2019; Virtala et

167 al., 2022; Winkler et al., 2003, 2009), leading many researchers to refer to it as the mismatch
168 response (MMR) in the neonatal brain.
169
In our study, we exposed neonates to speech samples characterized by positive (i.e., happy) and
neutral prosodies. Our selection of positive emotions over negative ones (e.g., fear, sadness, or
anger) was guided not only by ethical considerations but also by previous research indicating an
early preference for positive emotions in neonates (Farroni et al., 2007; Mastropieri & Turkewitz,
1999; Zhang et al., 2019; with the exception of Cheng et al., 2012). Additionally, to eliminate the
possibility of neonates distinguishing emotional voices solely based on their low-level acoustic
features, we included another set of control sounds. These nonvocal stimuli were meticulously
matched with their vocal prosodic counterparts in terms of mean intensity and fundamental
frequency (Cheng et al., 2012). Consequently, our primary objective is to pinpoint the
developmental stage (i.e., the GA group) at which the discrimination between happy and neutral
stimuli becomes apparent for emotional voices while remaining absent for acoustically matched
control sounds.
182

Results

The MMR was extracted using ERP difference waves, computed by subtracting the ERP evoked
by the standard stimulus (neutral sound) from the ERP evoked by the deviant stimulus (happy
sound) (Näätänen et al., 2007). Brain electrical activity was recorded from the F3, F4, C3, C4, P3,
and P4 sites following the international 10/20 system. However, this study primarily focused on
data from electrodes F3 and F4, as the neonatal MMR exhibits a frontal distribution (Cheng et al.,
2012; Cheour et al., 2002). Figure 1 displays MMR waveforms recorded from all six electrodes.
190
Initially, we conducted a three-way repeated measures ANOVA on the mean MMR amplitudes
(time window: 150 ms to 400 ms after sound onset) with factors including condition
(vocal/nonvocal), hemisphere (left/right frontal, i.e., F3/F4) as within-subjects factors, and
neonatal group (GA = 35, 36, 37, 38, 39, and 40 weeks) as the between-subjects factor. However,
neither the main effect nor the interaction effects involving the hemisphere factor were statistically
significant (for detailed statistics, please refer to the supplemental file subtitled "Result of the
three-way ANOVA"). Consequently, we removed the hemisphere factor and averaged the MMR
waveforms recorded at the F3 and F4 electrodes.
199
Subsequently, we performed a two-way repeated measures ANOVA with condition and group as
the two factors. The main effect of stimuli was significant, $F(1,114) = 38.827$, $p < 0.001$, $\eta_p^2 =$
0.254. Specifically, vocal stimuli elicited larger MMRs (mean ± standard deviation: 3.839 ± 4.855
µV) compared to nonvocal stimuli (0.496 ± 4.779 µV). The main effect of group was also
significant, $F(5,114) = 3.228$, $p = 0.009$, $\eta_p^2 = 0.124$. In general, MMR amplitudes were smaller
in the GA35 (0.590 ± 4.579 µV) and GA36 (0.141 ± 4.807 µV) groups compared to the GA37
(2.801 ± 5.585 µV), GA38 (2.760 ± 4.382 µV), GA39 (3.401 ± 4.871 µV), and GA40 groups
(3.311 ± 5.491 µV). However, no significant differences were found in pairwise comparisons after
Bonferroni adjustment for multiple comparisons.

The interaction between stimuli and group was significant, $F(5,114) = 3.127$, $p = 0.011$, $\eta_p^2 =$

0.121 (as shown in Figure 2). Simple effect analysis revealed that MMR amplitudes were larger in

the vocal condition compared to the nonvocal condition in the GA37 ($F(1,114) = 15.254$, $p <$

0.001, $\eta_p^2 = 0.118$; vocal = 5.367 ± 5.165 μV, nonvocal = 0.235 ± 4.847 μV), GA38 ($F(1,114) =$

16.072, $p < 0.001$, $\eta_p^2 = 0.124$; vocal = 5.394 ± 3.145 μV, nonvocal = 0.126 ± 3.861 μV), GA39

($F(1,114) = 8.393$, $p = 0.005$, $\eta_p^2 = 0.069$; vocal = 5.305 ± 4.011 μV, nonvocal = 1.498 ± 4.998

μV), and GA40 groups ($F(1,114) = 14.482$, $p < 0.001$, $\eta_p^2 = 0.113$; vocal = 5.811 ± 5.298 μV,

nonvocal = 0.811 ± 4.546 μV). However, there were no significant differences in MMR

amplitudes between the two kinds of stimuli in the GA35 ($F(1,114) = 0.026$, $p = 0.873$, $\eta_p^2 <$

0.001; vocal = 0.695 ± 4.031 μV, nonvocal = 0.485 ± 5.173 μV) and GA36 groups ($F(1,114) =$

0.236, $p = 0.628$, $\eta_p^2 = 0.002$; vocal = 0.460 ± 4.104 μV, nonvocal = -0.179 ± 5.511 μV).

Further analysis revealed that vocal stimuli evoked varying MMR amplitudes across groups,

$F(5,114) = 6.768$, $p < 0.001$, $\eta_p^2 = 0.229$. Specifically, the MMRs evoked by vocal stimuli were

smaller in the GA35 group compared to GA37 ($p = 0.014$), GA38 ($p = 0.013$), GA39 ($p = 0.017$),

and GA40 groups ($p = 0.005$). Similarly, the MMRs evoked by vocal stimuli were smaller in the

GA36 group compared to GA37 ($p = 0.008$), GA38 ($p = 0.008$), GA39 ($p = 0.009$), and GA40

groups ($p = 0.003$). However, nonvocal stimuli did not elicit significantly different MMR

amplitudes across groups, $F(5,114) = 0.300$, $p = 0.912$, $\eta_p^2 = 0.013$.

Figure 1. Mismatch response (MMR) waveforms recorded across six electrodes in different
gestational age (GA) groups. The MMR is extracted by subtracting the event-related potential
(ERP) elicited by the standard stimulus (neutral condition) from the ERP evoked by the deviant
stimulus (happy condition).

Figure 2. Primary MMR results. A, ERP waveforms (averaged at the F3 and F4 electrodes) across
the six GA groups. The time window for assessing MMR amplitude is indicated between the two
vertical gray lines. B, Violin plot illustrating MMR amplitudes in the six GA groups. Simple effect
analysis was performed with pairwise comparisons, corrected using the Bonferroni method: ** p <
0.010, *** p < 0.001.

Discussion

The current study elucidates a pivotal developmental milestone in neonatal emotional
responsiveness by investigating their ability to perceive vocal emotions. The findings illuminate a
distinct turning point at 37 weeks of gestational age (GA), representing the onset of heightened
perceptual acuity for emotional vocal expressions. This milestone is particularly evident in the
robust MMR to positive emotional vocal prosody. Significantly, the absence of this discrimination
ability when acoustically matched control sounds were presented underscores the specificity of
this developmental shift towards emotional voice processing. Our identification of the 37-week
GA mark aligns with previous research, which indicated emotional sensitivity in term neonates
born at or after 37 weeks of gestation (Cheng et al., 2012; Farroni et al., 2007; Mastropieri &
Turkewitz, 1999; Zhang et al., 2019) and in preterm neonates (GA < 37 weeks) tested at term age
(Kostilainen et al., 2020). Notably, our findings also reveal that neonates born before 37 weeks
GA do not exhibit these emotional discrimination abilities.

In the final trimester of pregnancy, the human brain undergoes a period of rapid and continuous changes in neural structure and cognitive functions (Bayer et al., 1993; Clancy et al., 2007). Although there is no direct evidence to support the notion that preterm neonates can decode vocal emotions, they have displayed an aptitude for processing speech stimuli with social contexts. For example, neonates born at or after 29 weeks GA have shown a preference for infant-directed speech, characterized by a high pitch, exaggerated pitch modifications, and a slow rate (Butler et al., 2014). This preference has been associated with increased visual attention, heightened alertness (Eckerman et al., 1994), reduced heart rate (White-Traut et al., 1997), and enhanced speech differentiation in premature babies (Richard et al., 2022). Additionally, neonates born at or after 30 weeks GA have demonstrated an age-related increase in sensitivity to maternal voices (D Chorna et al., 2018; A. P. F. Key et al., 2012), leading to beneficial effects on cognitive and neurobehavioral development (Caskey et al., 2011; Picciolini et al., 2014; see Provenzi et al., 2018 for a review). These effects encompass improved feeding behaviors, heightened responsiveness (Katz, 1971; Krueger et al., 2010), enhanced weight gain (Zimmerman et al., 2013), and activated auditory cortical plasticity (Webb et al., 2015). Both infant-directed speech and maternal voices feature extensive pitch modulation, and the preference for these emotionally prosodic-like voices during the preterm stage may prepare the developing brain to discriminate vocal emotions at 37 weeks GA, as demonstrated in this study.

Traditionally, 37 weeks of gestation served as the benchmark for fetal maturity, and term infants born within the 37 to 41 weeks GA range were generally considered healthy, forming a homogenous group. Recent insights, however, have unveiled variations in physical and cognitive maturation within this 5-week span of full-term pregnancy. Research indicates that neonates born at 37-38 weeks GA face increased risks of neonatal mortality and pediatric respiratory, neurologic, and endocrine morbidities compared to those born at 39-41 weeks GA (Cahen-Peretz et al., 2022; Clark et al., 2009; Edwards et al., 2013; Ghartey et al., 2012; Paz Levy et al., 2017; Sengupta et al., 2013; Tita et al., 2009). Furthermore, a dose-response relationship inversely linking GA to the risk of developmental delay has been identified in infants from preterm to full-term births (Rose et al., 2013; Schonhaut et al., 2015). Early birth (34-38 weeks GA) has also been found to have a detrimental impact on child development and academic achievement during school age (Bentley et al., 2016; Chan et al., 2016; Dong et al., 2012; Hedges et al., 2021; Murray et al., 2017; Nielsen et al., 2019; Noble et al., 2012). Consequently, the definition of a full-term pregnancy has been narrowed to a two-week window starting at 39 weeks (Spong, 2013), with nonmedically indicated deliveries between 37 and 38 weeks of gestation discouraged (ACOG Committee Opinion, 2019). While accumulating evidence underscores the adverse effects of the traditional 37-week threshold, our findings contribute to a limited body of research suggesting that neonatal social-emotional functioning reaches a development milestone at 37 weeks GA.

A more comprehensive understanding of the developmental trajectory of emotional sensitivity has the potential to revolutionize decision-making in the final weeks of pregnancy and the identification of newborns at risk of emotional and neurodevelopmental disorders, particularly autism. Individuals with autism often exhibit atypical perceptual and neural processing of emotional information, including emotional prosodic voices (Kuhl et al., 2005; Lindström et al.,

2018; Van Lancker et al., 1989; Wang et al., 2007; for comprehensive reviews, see Frühholz &
Staib, 2017; Yeung, 2022). While previous studies have indicated that social-emotional behavioral
indicators typically begin to demonstrate predictive power for autism from the second year of life
(Gliga et al., 2014; Jones et al., 2014), brain functional indicators of emotional processing during
infancy, especially within the first year of life, have already shown their predictive value (Ayoub
et al., 2022; Clairmont et al., 2021; Molnar-Szakacs et al., 2021). For instance, infants
subsequently diagnosed with autism displayed a smaller amplitude and shorter duration of the
negative central (Nc) component at six months of age when viewing smiling faces compared to
toys, a pattern not observed in infants who were subsequently undiagnosed (Jones et al., 2016).
Additionally, while the Nc and P400 components were able to distinguish between smiling, fearful,
and neutral facial expressions in typically developing 9-to-10-month-old infants, these EEG
indicators failed to differentiate emotional faces in infants at high risk for autism (Di Lorenzo et
al., 2021; A. P. Key et al., 2015). Moreover, it has been observed that infants at high risk for
autism exhibit diminished activation in the fusiform gyrus and hippocampus compared to healthy
controls when exposed to sad cries between the ages of 4 and 7 months (Blasi et al., 2015). The
fusiform gyrus, a region crucial for face perception and memory, and the hippocampus, which
plays a significant role in general learning and memory processes, are both implicated in this
phenomenon (Lisman et al., 2017; Rossion et al., 2024). Consequently, the hippocampus-fusiform
network, essential for the development of social cognitive skills, may serve as a predictive
indicator for the onset of autism. Building upon these existing studies, our research introduces a
promising early screening indicator for autism: the neonatal MMR in response to emotional voices.
We recommend future longitudinal studies to further elucidate the predictive role of this
neurophysiological indicator, thereby facilitating early diagnosis and intervention for
social-emotional disorders.
324
When interpretating the current findings, it is important to consider that the nonvocal control
sounds utilized in this study may not have adequately eliminated all low-level acoustic properties
that could aid neonatal discrimination. Specifically, while the nonvocal control counterparts
retained the fundamental frequency (f0) of the emotional prosodic voices, they did not replicate
the burst of energy associated with consonants. Consequently, it cannot be ruled out that neonates
utilized consonant characteristics to discriminate emotional prosodies conveyed by disyllables.
Additionally, the nonvocal sounds were generated using a simple filtering method, resulting in
certain vocal-like components persisting in these control sounds. The limitations of the control
sound materials should be given greater consideration in future replication or further research.
334
Furthermore, there is a compelling need for future investigations to expand upon the present
findings by incorporating a broader array of emotional stimuli. Non-speech emotional
vocalizations, such as laughter, crying, or retching, as well as natural emotional auditory cues like
thunder, flowing water, hissing snakes, and bird calls, offer a rich spectrum of emotional materials
that have been shown to engage the perceptual faculties of neonates and infants (Blasi et al., 2011;
Erlich et al., 2013). This multifaceted approach could illuminate whether the developmental
milestone observed at 37 weeks GA is specific to the processing of emotional prosodic speech and
vocal expressions, or if it extends to encompass a broader range of both artificial and natural
emotional auditory cues. Moreover, the use of non-speech emotional stimuli aids in resolving the

debate between nature and nurture concerning the onset of emotional sensitivity at 37 weeks GA. It cannot solely attribute the current finding of discrimination to the innate maturational explanation, given that the auditory system becomes functional at the end of the second trimester of pregnancy, allowing exposure to spoken language in utero to influence the development of speech perception (DeCasper & Spence, 1986; Moon et al., 2013; Partanen et al., 2013). The ability to discriminate prosodic emotions starting at 37 weeks GA could stem from additional exposure in utero to speech. Future exploration is need to definitively investigate prenatal learning by utilizing emotional sounds that are infrequently encountered in the prenatal environment. Finally, the inclusion of non-speech emotional materials may offer insights into the potential right lateralization of emotional processing in the neonatal brain. While prior studies (including some cited herein) have identified right lateralization for emotional processing in full-term neonates (Cheng et al., 2012; Zhang et al., 2019; see Bisiacchi & Cainelli, 2022 for a comprehensive review), the introduction of non-speech materials can help disentangle the confounding effects of left lateralization, which is associated with language processing and has been identified in both preterm (Mahmoudzadeh et al., 2013) and full-term neonates (Kotilahti et al., 2010; May et al., 2018; Peña et al., 2003; Sato et al., 2012; Vannasing et al., 2016; Wu et al., 2022).

In summary, this study has illuminated a pivotal developmental milestone – the emergence of heightened perceptual acuity for emotional vocal expressions at 37 weeks GA. It is important to note that neonates' perceptual sensitivity at this stage is unlikely to be associated with a profound conceptual understanding of the meaning of emotions. Nevertheless, this unique discrimination ability in early life may serve as a foundational building block for the subsequent development of emotional and social cognition. Beyond its scientific significance, our findings underscore the critical role of gestational age in shaping early perceptual abilities and offer a promising avenue for early screening and intervention in neurodevelopmental disorders, particularly autism, where early detection is of paramount importance. This work not only deepens our understanding of neonatal social and emotional development but also provides a potential tool to support early diagnosis and intervention in this critical realm of child health and well-being.

Materials and methods

Subjects

The research received approval from both the Ethical Committee of Peking University First Hospital and the Chinese Clinical Trial Registry (ChiCTR2300069898). Initially, we planned to include 120 healthy neonates, with 60 being boys, in the data analysis. These participants were categorized into six groups based on their GA, specifically 35, 36, 37, 38, 39, and 40 weeks, with each group comprising twenty subjects. For instance, the GA35 group comprised neonates with GA ranging from 35 weeks plus 0 day to 6 days. However, we ultimately recruited 198 neonates to obtain 120 valid datasets due to non-cooperation of newborns (n = 75) or technical issues (n = 3). Specially, 11, 12, 11, 14, 13, and 14 neonates were excluded from data analysis in the GA35, GA36, GA37, GA38, GA39, and GA40 groups, respectively, due to crying or irritable movements during EEG device preparation and EEG recording.

385 The mothers of these neonates were monolingual and nurtured their babies in a native language
386 environment. All neonates participated the experiment within the first 24 hours after birth, with a
387 mean ± standard deviation of 17.8 ± 0.4 hours for the 120 valid data.
388
389 Prior to data collection, written consent was obtained from the parents or legal guardians of all
390 participating neonates for access to clinical information and EEG data collection for scientific
391 purposes. While sample sizes were not statistically predetermined, including twenty subjects per
392 GA group represented the maximum feasible number within a two-year period at Peking
393 University First Hospital.
394
All subjects met the following inclusion criteria: 1) normal birth weight for their GA; 2) absence
of clinical symptoms at the time of EEG recording; 3) no previous sedation or medication prior to
EEG recording; and 4) normal hearing results in an evoked otoacoustic emissions test (ILO88 Dpi,
Otodynamics Ltd, Hatfield, UK). Additionally, subjects did not exhibit any of the following
neurological or metabolic disorders: 1) hypoxic-ischemic encephalopathy, 2) intraventricular
hemorrhage or white matter damage detected by cranial ultrasound, 3) congenital malformation, 4)
central nervous system infection, 5) metabolic disorder, 6) clinical evidence of seizures, and 7)
signs of asphyxia.
403

Stimuli

A total of 85 possible combinations of consonants and vowels, which are standard in Chinese
(Lee & Zee, 2003) and common to most human languages (e.g., 'dada' and 'keke'), were
recorded by a native Chinese-speaking adult woman with the Peking dialect. Each disyllable was
recorded with four repetitions, two using a happy prosody and two with a neutral prosody,
resulting in a total of 340 disyllables (85 × 4). Twenty Chinese undergraduate students (10 men,
mean age 20.1 ± 1.2 years) performed a discrimination task, distinguishing between happy and
neutral stimuli, and rated the affective content of these stimuli.
412
In the affective rating task, participants assessed the intensity of happiness (on a 9-point scale
ranging from 1 being the least happy to 9 being the happiest) and the valence (on a 9-point scale
ranging from 1 being the most negative, 5 being neutral, to 9 being the most positive) of the 340
stimuli. This study selected five pairs of happy and neutral disyllables that shared the same
consonant-monophthong combinations and achieved 100% discrimination accuracy in the
discrimination task (i.e., 'dada', 'dudu', 'gege', 'keke', and 'tutu' in Chinese Pinyin).
Paired-samples t-tests demonstrated that the happy disyllables were rated as significantly happier
($t(4) = 24.70$, $p < 0.001$; happy intensity: 7.49 ± 0.20 versus 3.53 ± 0.19) and had a more positive
valence ($t(4) = 18.55$, $p < 0.001$; valence: 7.11 ± 0.12 versus 4.91 ± 0.24) than their neutral
counterparts. These ten disyllables were then standardized to have the same mean intensity and
had duration of 400 ms using Adobe Audition (v.2022; Adobe Systems Inc., San Jose, CA).
424
To ensure that neonates were discriminating based on prosodic cues containing emotional content
rather than low-level acoustic properties, we employed a method similar to Cheng et al. (2012)
and created a separate set of nonvocal control sounds. We hypothesized that the fundamental

frequency (f0) alone does not convey emotional content in voices and that neonates require
multiple other prosodic cues embedded in the high-frequent component to discern emotions, as
suggested by Cheng et al. (2012) and Zhang et al. (2014). As a result, ten nonvocal sounds were
generated to match the f0 contours and temporal envelopes of their corresponding vocal sounds.
This matching process was carried out using Matlab (v.2021b; MathWorks, Inc., South Natick,
MA). Specifically, we initially applied a zero-phase filter with a bandpass of mean $f0 \pm 150$ Hz to
obtain f0-matched sounds of prosodic voices. Subsequently, a normalization procedure was
implemented to ensure that the intensity of each pair of vocal and nonvocal sounds was equal.
Oscillograms and spectrograms of the auditory stimuli utilized in this study are presented in
Figure 3, generated using Praat (v.6.3.17, www.praat.org). All auditory stimuli, along with their
pronunciations and rating scores, are available in the supplemental material labeled "experimental
sounds".

To optimize the diversity of our material and increase the generalizability of our results, we
utilized ten sets of sounds. Each set included both positive and neutral prosodic voices, along with
their respective nonvocal counterparts. These auditory materials were distributed randomly and
evenly within each neonatal GA group, ensuring that each set was presented twice (to two
individuals) in each GA group.

Procedure

The sound stimuli were presented in two blocks: the vocal and nonvocal conditions, utilizing the
odd-ball paradigm. The standard stimulus was either a vocal or nonvocal neutral sound, while the
deviant stimulus was either a vocal or nonvocal happy sound. Each block consisted of 240
standard stimuli (80%) and 60 deviant stimuli (20%). The standard and deviant stimuli were
presented randomly, ensuring that each deviant stimulus was followed by at least two standard
stimuli. Each sound had a duration of 400 ms, and the inter-trial interval was silent, with varying
durations ranging from 500 to 700 ms. Each block lasted for 5 minutes, and the order of the vocal
and nonvocal blocks was counterbalanced across participants. A 5-minute break separated the two
blocks, resulting in a total EEG recording duration of 15 minutes.

The experiment took place in the neonatal ward of Peking University First Hospital. Neonates
were transported to a designated testing room for EEG recording as soon as their condition
stabilized after birth. In this room, they were separated from their mothers to minimize any
natural exposure to speech or speech stimuli other than those utilized in the experiment. Auditory
stimuli were presented through a pair of loudspeakers positioned approximately 30 cm away from
the neonates' left and right ears, at a sound pressure level of 55 to 60 dB, with an average
background noise intensity level of 30 dB. EEG recording was conducted while the neonates were
in a natural sleep state (Cheour et al., 2002; Wu et al., 2022).

Data recording and analysis

We recorded brain electrical activity using an electrical amplifier (NeuSen.W32, Neuracle,

Changzhou, China) at a sampling frequency of 1000 Hz. Initially, the data were recorded online
with reference to the left mastoid and subsequently re-referenced offline to the average of the left
and right mastoids. The ground electrode was positioned on the forehead. For the recording of
vertical eye movements, an electrooculogram (EOG) electrode was positioned beneath the left eye,
while another was placed at the left external canthi for recording horizontal eye movements.
Throughout the recording process, electrode impedances were meticulously maintained below 10
kΩ.
476
We eliminated ocular artifacts from the EEG data using a regression procedure implemented in
NeuroScan software (Scan 4.3, NeuroScan, Herndon, VA). Subsequently, we employed Matlab
(v.2021b; Mathworks, Inc., Sherborn, MA) for data processing and result presentation. The
EOG-corrected EEG data were then offline filtered with a half-amplitude cutoff range of 0.01~30
Hz and segmented from 200 ms before sound presentation until 1000 ms after sound onset.
Epochs were baseline-corrected relative to the mean voltage during the 200 ms preceding sound
presentation. Any epochs containing artifacts with peak deflections exceeding ±200 µV were
rejected (see also Biro et al., 2021; Di Lorenzo et al., 2021; Kumaravel et al., 2022), followed by
averaging for each experimental condition. The number of valid epochs did not exhibit a
significant difference across neonatal groups (please refer to the supplemental file subtitled
"Epoch number"). The time window for the MMR component was pre-defined as 150 ms to 400
ms after sound onset, based on prior knowledge (Cheour et al., 2002), and utilized throughout the
data analysis.
490
We performed statistical analyses using SPSS Statistics (v. 20.0; IBM, Somers, USA). Descriptive
data are reported as mean ± standard deviation. The significance level was set at 0.05. We applied
the Greenhouse-Geisser correction for ANOVA tests when deemed appropriate. Post-hoc tests for
significant main effects were conducted using the Bonferroni method. Significant interactions
were explored through simple effects models. We reported partial eta-squared (η_p^2) as a measure of
effect size in ANOVA tests.
497

Figure 3. Oscillograms and spectrograms displaying vocal and nonvocal auditory stimuli for the syllable 'dada' (international phonetic symbol: [ta ta]).

Acknowledgments

This study was funded by the National High Level Hospital Clinical Research Funding (High Quality Clinical Research Project of Peking University First Hospital, 2022CR68), the National Natural Science Foundation of China (32271102; 31920103009), the Major Project of National Social Science Foundation (20&ZD153), Shenzhen-Hong Kong Institute of Brain Science (2024SHIBS0004), and the National Key Research and Development Program of China (2021YFC2700700).

Data availability

The experimental materials are accessible as supplementary files for download. EEG epochs from all 120 datasets can be downloaded from https://osf.io/a3xzy/. The data and materials from this study are available for academic purpose free of charge, on the condition that proper citation to this article is provided.

Author contributions

D. Zhang and X. Hou conceptualized and designed the research. P. Zhang, X. Hou, and C. Peng conducted the experiments, while P. Zhang, L. Mo, and D. Zhang were responsible for data analysis. D. Zhang authored the paper, and X. Hou revised and provided critical input to the manuscript.

References

American College of Obstetricians and Gynecologists, & Society for Maternal-Fetal Medicine. (2019). ACOG committee opinion no. 765: Avoidance of nonmedically indicated early-term deliveries and associated neonatal morbidities. *Obstetrics & Gynecology*, *133*, e156-e163. https://doi.org/10.1097/AOG.0000000000003076

Ayoub, M. J., Keegan, L., Tager-Flusberg, H., & Gill, S. V. (2022). Neuroimaging techniques as descriptive and diagnostic tools for infants at risk for autism spectrum disorder: A systematic review. *Brain Sciences*, *12*(5), 602. https://doi.org/10.3390/brainsci12050602

Bayer, S. A., Altman, J., Russo, R. J., & Zhang, X. (1993). Timetables of neurogenesis in the human brain based on experimentally determined patterns in the rat. *Neurotoxicology*, *14*(1), 83–144.

Bentley, J. P., Roberts, C. L., Bowen, J. R., Martin, A. J., Morris, J. M., & Nassar, N. (2016). Planned birth before 39 weeks and child development: A population-based study. *Pediatrics*, *138*(6), e20162002. https://doi.org/10.1542/peds.2016-2002

Biro, S., Peltola, M. J., Huffmeijer, R., Alink, L. R. A., Bakermans-Kranenburg, M. J., & van IJzendoorn, M. H. (2021) Frontal EEG asymmetry in infants observing separation and comforting events: The role of infants' attachment relationship. *Developmental Cognitive Neuroscience*, 48, 100941. https://doi.org/10.1016/j.dcn.2021.100941.

Bisiacchi, P., & Cainelli, E. (2022). Structural and functional brain asymmetries in the early phases of life: A scoping review. *Brain Structure and Function*, *227*(2), 479–496. https://doi.org/10.1007/s00429-021-02256-1

Blasi, A., Lloyd-Fox, S., Sethna, V., Brammer, M. J., Mercure, E., Murray, L., Williams, S. C. R., Simmons, A., Murphy, D. G. M., & Johnson, M. H. (2015). Atypical processing of voice sounds in infants at risk for autism spectrum disorder. *Cortex*, *71*, 122–133. https://doi.org/10.1016/j.cortex.2015.06.015

Blasi, A., Mercure, E., Lloyd-Fox, S., Thomson, A., Brammer, M., Sauter, D., Deeley, Q., Barker, G. J., Renvall, V., Deoni, S., Gasston, D., Williams, S. C. R., Johnson, M. H., Simmons, A., & Murphy, D. G. M. (2011). Early specialization for voice and emotion processing in the infant brain. *Current Biology*, *21*(14), 1220–1224. https://doi.org/10.1016/j.cub.2011.06.009

Butler, S. C., O'Sullivan, L. P., Shah, B. L., & Berthier, N. E. (2014). Preference for infant-directed speech in preterm infants. *Infant Behavior and Development*, *37*(4), 505–511. https://doi.org/10.1016/j.infbeh.2014.06.007

Cahen-Peretz, A., Tsaitlin-Mor, L., Abu-Ahmad, W., Ben-Shushan, M. T., Levine, H., & Walfisch, A. (2022). Long-term respiratory outcomes in early-term born offspring: A systematic review

and meta-analysis. *American Journal of Obstetrics and Gynecology MFM*, *4*(3), 100570.

https://doi.org/10.1016/j.ajogmf.2022.100570

Caron, A. J., Caron, R. F., & MacLean, D. J. (1988). Infant discrimination of naturalistic

emotional expressions: The role of face and voice. *Child Development*, *59*(3), 604–616.

Caskey, M., Stephens, B., Tucker, R., & Vohr, B. (2011). Importance of parent talk on the

development of preterm infant vocalizations. *Pediatrics*, *128*(5), 910–916.

https://doi.org/10.1542/peds.2011-0609

Chan, E., Leong, P., Malouf, R., & Quigley, M. A. (2016). Long-term cognitive and school

outcomes of late-preterm and early-term births: A systematic review. *Child: Care, Health and*

Development, *42*(3), 297–312. https://doi.org/10.1111/cch.12320

Cheng, Y., Lee, S.-Y., Chen, H.-Y., Wang, P.-Y., & Decety, J. (2012). Voice and emotion

processing in the human neonatal brain. *Journal of Cognitive Neuroscience*, *24*(6),

1411–1419. https://doi.org/10.1162/jocn_a_00214

Cheour, M., Ceponiene, R., Lehtokoski, A., Luuk, A., Allik, J., Alho, K., & Näätänen, R. (1998).

Development of language-specific phoneme representations in the infant brain. *Nature*

Neuroscience, *1*(5), 351–353. https://doi.org/10.1038/1561

Cheour, M., Martynova, O., Näätänen, R., Erkkola, R., Sillanpää, M., Kero, P., Raz, A., Kaipio,

M.-L., Hiltunen, J., Aaltonen, O., Savela, J., & Hämäläinen, H. (2002). Speech sounds

learned by sleeping newborns. *Nature*, *415*(6872), 599–600. https://doi.org/10.1038/415599b

Chládková, K., Urbanec, J., Skálová, S., & Kremláček, J. (2021). Newborns' neural processing of

native vowels reveals directional asymmetries. *Developmental Cognitive Neuroscience*, *52*,

101023. https://doi.org/10.1016/j.dcn.2021.101023

Clairmont, C., Wang, J., Tariq, S., Sherman, H. T., Zhao, M., & Kong, X.-J. (2021). The value of

brain imaging and electrophysiological testing for early screening of autism spectrum

disorder: A systematic review. *Frontiers in Neuroscience*, *15*, 812946.

https://doi.org/10.3389/fnins.2021.812946

Clancy, B., Finlay, B. L., Darlington, R. B., & Anand, K. J. S. (2007). Extrapolating brain

development from experimental species to humans. *Neurotoxicology*, *28*(5), 931–937.

https://doi.org/10.1016/j.neuro.2007.01.014

Clark, S. L., Miller, D. D., Belfort, M. A., Dildy, G. A., Frye, D. K., & Meyers, J. A. (2009).

Neonatal and maternal outcomes associated with elective term delivery. *American Journal of*

Obstetrics and Gynecology, *200*(2), 156.e1-4. https://doi.org/10.1016/j.ajog.2008.08.068

D Chorna, O., L Hamm, E., Shrivastava, H., & Maitre, N. L. (2018). Feasibility of event-related

potential (ERP) biomarker use to study effects of mother's voice exposure on speech sound

differentiation of preterm infants. *Developmental Neuropsychology*, *43*(2), 123–134.

https://doi.org/10.1080/87565641.2018.1433671

DeCasper, A. J. & Spence, M. J. (1986). Prenatal maternal speech influences newborns' perception

of speech sounds. *Infant Behavior and Development*, 9, 133–150.

https://doi.org/10.1016/0163-6383(86)90025-1

Di Lorenzo, R., Munsters, N. M., Ward, E. K., de Jonge, M., Kemner, C., & van den Boomen, C.

(2021). Is it fear? Similar brain responses to fearful and neutral faces in infants with a

heightened likelihood for autism spectrum disorder. *Journal of Autism and Developmental*

Disorders, *51*(3), 961–972. https://doi.org/10.1007/s10803-020-04560-x

Dong, Y., Chen, S., & Yu, J. (2012). A systematic review and meta-analysis of long-term

development of early term infants. *Neonatology*, *102*(3), 212–221. https://doi.org/10.1159/000338099

Eckerman, C. O., Oehler, J. M., Medvin, M. B., & Hannan, T. E. (1994). Premature newborns as social partners before term age. *Infant Behavior and Development*, *17*(1), 55–70. https://doi.org/10.1016/0163-6383(94)90022-1

Edwards, M. O., Kotecha, S. J., & Kotecha, S. (2013). Respiratory distress of the term newborn infant. *Paediatric Respiratory Reviews*, *14*(1), 29–36; quiz 36–37. https://doi.org/10.1016/j.prrv.2012.02.002

Farroni, T., Menon, E., Rigato, S., & Johnson, M. H. (2007). The perception of facial expressions in newborns. *The European Journal of Developmental Psychology*, *4*(1), 2–13. https://doi.org/10.1080/17405620601046832

Fernald, A. (1993). Approval and disapproval: Infant responsiveness to vocal affect in familiar and unfamiliar languages. *Child Development*, *64*(3), 657–674.

Frühholz, S., & Staib, M. (2017). Neurocircuitry of impaired affective sound processing: A clinical disorders perspective. *Neuroscience and Biobehavioral Reviews*, *83*, 516–524. https://doi.org/10.1016/j.neubiorev.2017.09.009

Ghartey, K., Coletta, J., Lizarraga, L., Murphy, E., Ananth, C. V., & Gyamfi-Bannerman, C. (2012). Neonatal respiratory morbidity in the early term delivery. *American Journal of Obstetrics and Gynecology*, *207*(4), 292.e1-4. https://doi.org/10.1016/j.ajog.2012.07.022

Gliga, T., Jones, E. J. H., Bedford, R., Charman, T., & Johnson, M. H. (2014). From early markers to neuro-developmental mechanisms of autism. *Developmental Review*, *34*(3), 189–207. https://doi.org/10.1016/j.dr.2014.05.003

Graham, A. M., Fisher, P. A., & Pfeifer, J. H. (2013). What sleeping babies hear: A functional MRI study of interparental conflict and infants' emotion processing. *Psychological Science*, *24*(5), 782–789. https://doi.org/10.1177/0956797612458803

Grossmann, T., Oberecker, R., Koch, S. P., & Friederici, A. D. (2010). The developmental origins of voice processing in the human brain. *Neuron*, *65*(6), 852–858. https://doi.org/10.1016/j.neuron.2010.03.001

Hedges, A., Corman, H., Noonan, K., & Reichman, N. E. (2021). Gestational age at term and educational outcomes at age nine. *Pediatrics*, *148*(2), e2020021287. https://doi.org/10.1542/peds.2020-021287

Jones, E. J. H., Gliga, T., Bedford, R., Charman, T., & Johnson, M. H. (2014). Developmental pathways to autism: A review of prospective studies of infants at risk. *Neuroscience and Biobehavioral Reviews*, *39*(100), 1–33. https://doi.org/10.1016/j.neubiorev.2013.12.001

Jones, E. J. H., Venema, K., Earl, R., Lowy, R., Barnes, K., Estes, A., Dawson, G., & Webb, S. J. (2016). Reduced engagement with social stimuli in 6-month-old infants with later autism spectrum disorder: A longitudinal prospective study of infants at high familial risk. *Journal of Neurodevelopmental Disorders*, *8*, 7. https://doi.org/10.1186/s11689-016-9139-8

Katz, V. (1971). Auditory stimulation and developmental behavior of the premature infant. *Nursing Research*, *20*(3), 196.

Key, A. P. F., Lambert, E. W., Aschner, J. L., & Maitre, N. L. (2012). Influence of gestational age and postnatal age on speech sound processing in NICU infants. *Psychophysiology*, *49*(5), 720–731. https://doi.org/10.1111/j.1469-8986.2011.01353.x

Key, A. P., Ibanez, L. V., Henderson, H. A., Warren, Z., Messinger, D. S., & Stone, W. L. (2015).

Positive affect processing and joint attention in infants at high risk for autism: An exploratory study. *Journal of Autism and Developmental Disorders*, *45*(12), 4051–4062. https://doi.org/10.1007/s10803-014-2191-x

Kostilainen, K., Partanen, E., Mikkola, K., Wikström, V., Pakarinen, S., Fellman, V., & Huotilainen, M. (2020). Neural processing of changes in phonetic and emotional speech sounds and tones in preterm infants at term age. *International Journal of Psychophysiology*, *148*, 111–118. https://doi.org/10.1016/j.ijpsycho.2019.10.009

Kotilahti, K., Nissilä, I., Näsi, T., Lipiäinen, L., Noponen, T., Meriläinen, P., Huotilainen, M., & Fellman, V. (2010). Hemodynamic responses to speech and music in newborn infants. *Human Brain Mapping*, *31*(4), 595–603. https://doi.org/10.1002/hbm.20890

Krueger, C., Parker, L., Chiu, S.-H., & Theriaque, D. (2010). Maternal voice and short-term outcomes in preterm infants. *Developmental Psychobiology*, *52*(2), 205–212. https://doi.org/10.1002/dev.20426

Kuhl, P. K., Coffey-Corina, S., Padden, D., & Dawson, G. (2005). Links between social and linguistic processing of speech in preschool children with autism: Behavioral and electrophysiological measures. *Developmental Science*, *8*(1), F1–F12. https://doi.org/10.1111/j.1467-7687.2004.00384.x

Kumaravel, V. P., Farella, E., Parise, E., & Buiatti, M. (2022). NEAR: An artifact removal pipeline for human newborn EEG data. *Developmental Cognitive Neuroscience*, 54, 101068. https://doi.org/10.1016/j.dcn.2022.101068.

Latinus, M., & Belin, P. (2011). Human voice perception. *Current Biology*, *21*(4), R143-145. https://doi.org/10.1016/j.cub.2010.12.033

Lee, W.-S., & Zee, E. (2003). Standard Chinese (Beijing). *Journal of the International Phonetic Association*, *33*(1), 109–112. https://doi.org/10.1017/S0025100303001208

Lindquist, K. A., Wager, T. D., Kober, H., Bliss-Moreau, E., & Barrett, L. F. (2012). The brain basis of emotion: A meta-analytic review. *The Behavioral and Brain Sciences*, *35*(3), 121–143. https://doi.org/10.1017/S0140525X11000446

Lindström, R., Lepistö-Paisley, T., Makkonen, T., Reinvall, O., Nieminen-von Wendt, T., Alén, R., & Kujala, T. (2018). Atypical perceptual and neural processing of emotional prosodic changes in children with autism spectrum disorders. *Clinical Neurophysiology*, *129*(11), 2411–2420. https://doi.org/10.1016/j.clinph.2018.08.018

Lisman, J., Buzsáki, G., Eichenbaum, H., Nadel, L., Ranganath, C., & Redish, A. D. (2017). Viewpoints: how the hippocampus contributes to memory, navigation and cognition. *Nature Neuroscience*, 20(11), 1434-1447. https://doi.org/10.1038/nn.4661

Mahmoudzadeh, M., Dehaene-Lambertz, G., Fournier, M., Kongolo, G., Goudjil, S., Dubois, J., Grebe, R., & Wallois, F. (2013). Syllabic discrimination in premature human infants prior to complete formation of cortical layers. *Proceedings of the National Academy of Sciences of the United States of America*, *110*(12), 4846–4851. https://doi.org/10.1073/pnas.1212220110

Mastropieri, D., & Turkewitz, G. (1999). Prenatal experience and neonatal responsiveness to vocal expressions of emotion. *Developmental Psychobiology*, *35*(3), 204–214. https://doi.org/10.1002/(sici)1098-2302(199911)35:3<204::aid-dev5>3.0.co;2-v

May, L., Gervain, J., Carreiras, M., & Werker, J. F. (2018). The specificity of the neural response to speech at birth. *Developmental Science*, *21*(3), e12564. https://doi.org/10.1111/desc.12564

Molnar-Szakacs, I., Kupis, L., & Uddin, L. Q. (2021). Neuroimaging markers of risk and

pathways to resilience in autism spectrum disorder. *Biological Psychiatry. Cognitive Neuroscience and Neuroimaging*, *6*(2), 200–210. https://doi.org/10.1016/j.bpsc.2020.06.017

Moon, C., Lagercrantz, H. & Kuhl, P. K. (2013). Language experienced in utero affects vowel perception after birth: a two-country study. *Acta Paediatrica*, 102, 156–160. https://doi.org/10.1111/apa.12098

Murray, S. R., Shenkin, S. D., McIntosh, K., Lim, J., Grove, B., Pell, J. P., Norman, J. E., & Stock, S. J. (2017). Long term cognitive outcomes of early term (37-38 weeks) and late preterm (34-36 weeks) births: A systematic review. *Wellcome Open Research*, *2*, 101. https://doi.org/10.12688/wellcomeopenres.12783.1

Näätänen, R., Paavilainen, P., Rinne, T., & Alho, K. (2007). The mismatch negativity (MMN) in basic research of central auditory processing: A review. *Clinical Neurophysiology*, *118*(12), 2544–2590. https://doi.org/10.1016/j.clinph.2007.04.026

Nielsen, T. M., Pedersen, M. V., Milidou, I., Glavind, J., & Henriksen, T. B. (2019). Long-term cognition and behavior in children born at early term gestation: A systematic review. *Acta Obstetricia Et Gynecologica Scandinavica*, *98*(10), 1227–1234. https://doi.org/10.1111/aogs.13644

Noble, K. G., Fifer, W. P., Rauh, V. A., Nomura, Y., & Andrews, H. F. (2012). Academic achievement varies with gestational age among children born at term. *Pediatrics*, *130*(2), e257-264. https://doi.org/10.1542/peds.2011-2157

Partanen, E., Kujala, T., Näätänen, R., Liitola, A., Sambeth, A., & Huotilainen, M. (2013). Learning-induced neural plasticity of speech processing before birth. *Proceedings of the National Academy of Sciences of the United States of America*, 110(37), 15145–50. https://doi.org/10.1073/pnas.1302159110.

Paz Levy, D., Sheiner, E., Wainstock, T., Sergienko, R., Landau, D., & Walfisch, A. (2017). Evidence that children born at early term (37-38 6/7 weeks) are at increased risk for diabetes and obesity-related disorders. *American Journal of Obstetrics and Gynecology*, *217*(5), 588.e1-588.e11. https://doi.org/10.1016/j.ajog.2017.07.015

Peña, M., Maki, A., Kovacić, D., Dehaene-Lambertz, G., Koizumi, H., Bouquet, F., & Mehler, J. (2003). Sounds and silence: An optical topography study of language recognition at birth. *Proceedings of the National Academy of Sciences of the United States of America*, *100*(20), 11702–11705. https://doi.org/10.1073/pnas.1934290100

Picciolini, O., Porro, M., Meazza, A., Giannì, M. L., Rivoli, C., Lucco, G., Barretta, F., Bonzini, M., & Mosca, F. (2014). Early exposure to maternal voice: Effects on preterm infants development. *Early Human Development*, *90*(6), 287–292. https://doi.org/10.1016/j.earlhumdev.2014.03.003

Provenzi, L., Broso, S., & Montirosso, R. (2018). Do mothers sound good? A systematic review of the effects of maternal voice exposure on preterm infants' development. *Neuroscience and Biobehavioral Reviews*, *88*, 42–50. https://doi.org/10.1016/j.neubiorev.2018.03.009

Richard, C., Jeanvoine, A., Stark, A. R., Hague, K., Kjeldsen, C., & Maitre, N. L. (2022). Randomized trial to increase speech sound differentiation in infants born preterm. *The Journal of Pediatrics*, *241*, 103-108.e3. https://doi.org/10.1016/j.jpeds.2021.10.035

Rose, O., Blanco, E., Martinez, S. M., Sim, E. K., Castillo, M., Lozoff, B., Vaucher, Y. E., & Gahagan, S. (2013). Developmental scores at 1 year with increasing gestational age, 37-41 weeks. *Pediatrics*, *131*(5), e1475-1481. https://doi.org/10.1542/peds.2012-3215

Rossion, B., Jacques, C., & Jonas, J. (2024). The anterior fusiform gyrus: The ghost in the cortical face machine. *Neuroscience & Biobehavioral Reviews*, 158, 105535. https://doi.org/10.1016/j.neubiorev.2024.105535.

Sato, H., Hirabayashi, Y., Tsubokura, H., Kanai, M., Ashida, T., Konishi, I., Uchida-Ota, M., Konishi, Y., & Maki, A. (2012). Cerebral hemodynamics in newborn infants exposed to speech sounds: A whole-head optical topography study. *Human Brain Mapping*, *33*(9), 2092–2103. https://doi.org/10.1002/hbm.21350

Schonhaut, L., Armijo, I., & Pérez, M. (2015). Gestational age and developmental risk in moderately and late preterm and early term infants. *Pediatrics*, *135*(4), e835-841. https://doi.org/10.1542/peds.2014-1957

Sengupta, S., Carrion, V., Shelton, J., Wynn, R. J., Ryan, R. M., Singhal, K., & Lakshminrusimha, S. (2013). Adverse neonatal outcomes associated with early-term birth. *JAMA Pediatrics*, *167*(11), 1053–1059. https://doi.org/10.1001/jamapediatrics.2013.2581

Singh, L., Morgan, J. L., & Best, C. T. (2002). Infants' listening preferences: Baby talk or happy talk? *Infancy*, *3*(3), 365–394. https://doi.org/10.1207/S15327078IN0303_5

Soderstrom, M., Reimchen, M., Sauter, D., & Morgan, J. L. (2017). Do infants discriminate non-linguistic vocal expressions of positive emotions? *Cognition and Emotion*, *31*(2), 298–311. https://doi.org/10.1080/02699931.2015.1108904

Spong, C. Y. (2013). Defining "term" pregnancy: Recommendations from the defining "term" pregnancy workgroup. *JAMA*, *309*(23), 2445–2446. https://doi.org/10.1001/jama.2013.6235

Thiede, A., Virtala, P., Ala-Kurikka, I., Partanen, E., Huotilainen, M., Mikkola, K., Leppänen, P. H. T., & Kujala, T. (2019). An extensive pattern of atypical neural speech-sound discrimination in newborns at risk of dyslexia. *Clinical Neurophysiology*, *130*(5), 634–646. https://doi.org/10.1016/j.clinph.2019.01.019

Tita, A. T. N., Landon, M. B., Spong, C. Y., Lai, Y., Leveno, K. J., Varner, M. W., Moawad, A. H., Caritis, S. N., Meis, P. J., Wapner, R. J., Sorokin, Y., Miodovnik, M., Carpenter, M., Peaceman, A. M., O'Sullivan, M. J., Sibai, B. M., Langer, O., Thorp, J. M., Ramin, S. M., … Eunice Kennedy Shriver NICHD Maternal-Fetal Medicine Units Network. (2009). Timing of elective repeat cesarean delivery at term and neonatal outcomes. *The New England Journal of Medicine*, *360*(2), 111–120. https://doi.org/10.1056/NEJMoa0803267

Vaish, A., & Striano, T. (2004). Is visual reference necessary? Contributions of facial versus vocal cues in 12-month-olds' social referencing behavior. *Developmental Science*, *7*(3), 261–269. https://doi.org/10.1111/j.1467-7687.2004.00344.x

Van Lancker, D., Cornelius, C., & Kreiman, J. (1989). Recognition of emotional prosodic meanings in speech by autistic, schizophrenic, and normal children. *Developmental Neuropsychology*, *5*(2–3), 207–226. https://doi.org/10.1080/87565648909540433

Vannasing, P., Florea, O., González-Frankenberger, B., Tremblay, J., Paquette, N., Safi, D., Wallois, F., Lepore, F., Béland, R., Lassonde, M., & Gallagher, A. (2016). Distinct hemispheric specializations for native and non-native languages in one-day-old newborns identified by fNIRS. *Neuropsychologia*, *84*, 63–69. https://doi.org/10.1016/j.neuropsychologia.2016.01.038

Virtala, P., Putkinen, V., Kailaheimo-Lönnqvist, L., Thiede, A., Partanen, E., & Kujala, T. (2022). Infancy and early childhood maturation of neural auditory change detection and its associations to familial dyslexia risk. *Clinical Neurophysiology*, *137*, 159–176.

https://doi.org/10.1016/j.clinph.2022.03.005

Walker-Andrews, A. S., & Grolnick, W. (1983). Discrimination of vocal expressions by young infants. *Infant Behavior and Development*, *6*(4), 491–498. https://doi.org/10.1016/S0163-6383(83)90331-4

Wang, A. T., Lee, S. S., Sigman, M., & Dapretto, M. (2007). Reading affect in the face and voice: Neural correlates of interpreting communicative intent in children and adolescents with autism spectrum disorders. *Archives of General Psychiatry*, *64*(6), 698–708. https://doi.org/10.1001/archpsyc.64.6.698

Webb, A. R., Heller, H. T., Benson, C. B., & Lahav, A. (2015). Mother's voice and heartbeat sounds elicit auditory plasticity in the human brain before full gestation. *Proceedings of the National Academy of Sciences of the United States of America*, *112*(10), 3152–3157. https://doi.org/10.1073/pnas.1414924112

White-Traut, R. C., Nelson, M. N., Silvestri, J. M., Cunningham, N., & Patel, M. (1997). Responses of preterm infants to unimodal and multimodal sensory intervention. *Pediatric Nursing*, *23*(2), 169–175, 193.

Winkler, I., Háden, G. P., Ladinig, O., Sziller, I., & Honing, H. (2009). Newborn infants detect the beat in music. *Proceedings of the National Academy of Sciences of the United States of America*, *106*(7), 2468–2471. https://doi.org/10.1073/pnas.0809035106

Winkler, I., Kushnerenko, E., Horváth, J., Ceponiene, R., Fellman, V., Huotilainen, M., Näätänen, R., & Sussman, E. (2003). Newborn infants can organize the auditory world. *Proceedings of the National Academy of Sciences of the United States of America*, *100*(20), 11812–11815. https://doi.org/10.1073/pnas.2031891100

Wu, Y. J., Hou, X., Peng, C., Yu, W., Oppenheim, G. M., Thierry, G., & Zhang, D. (2022). Rapid learning of a phonemic discrimination in the first hours of life. *Nature Human Behaviour*, *6*(8), Article 8. https://doi.org/10.1038/s41562-022-01355-1

Yeung, M. K. (2022). A systematic review and meta-analysis of facial emotion recognition in autism spectrum disorder: The specificity of deficits and the role of task characteristics. *Neuroscience and Biobehavioral Reviews*, *133*, 104518. https://doi.org/10.1016/j.neubiorev.2021.104518

Zhang, D., Chen, Y., Hou, X., & Wu, Y. J. (2019). Near-infrared spectroscopy reveals neural perception of vocal emotions in human neonates. *Human Brain Mapping*, *40*(8), Article 8. https://doi.org/10.1002/hbm.24534

Zhang, D., Liu, Y., Hou, X., Sun, G., Cheng, Y., & Luo, Y. (2014). Discrimination of fearful and angry emotional voices in sleeping human neonates: A study of the mismatch brain responses. *Frontiers in Behavioral Neuroscience*, *8*, 422. https://doi.org/10.3389/fnbeh.2014.00422

Zhao, C., Schiessl, I., Wan, M. W., Chronaki, G., & Abel, K. M. (2021). Development of the neural processing of vocal emotion during the first year of life. *Child Neuropsychology*, *27*(3), 333–350. https://doi.org/10.1080/09297049.2020.1853090

Zimmerman, E., Keunen, K., Norton, M., & Lahav, A. (2013). Weight gain velocity in very low-birth-weight infants: Effects of exposure to biological maternal sounds. *American Journal of Perinatology*, *30*(10), 863–870. https://doi.org/10.1055/s-0033-1333669

GA = 35 weeks   GA = 36 weeks   GA = 37 weeks

GA = 38 weeks   GA = 39 weeks   GA = 40 weeks

F3   F4
C3   C4
P3   P4

μV 8 +
0   800 ms
-

vocal MMR
nonvocal MMR