1 *A novel human pluripotent stem cell-based gene activation system identifies IGFBP2*
2 *as a mediator in the production of hematopoietic progenitors in vitro.*
3

Paolo Petazzi[1], Telma Ventura[6], Francesca Paola Luongo[6], Heather McClafferty[7], Alisha

May[6], Helen Alice Taylor[6], Micheal J Shipston[7,8], Nicola Romanò[7,8], Lesley M. Forrester[6],

Pablo Menéndez[1-5], Antonella Fidanza[6,9*].

1. Josep Carreras Leukemia Research Institute, Barcelona, Spain.

2. Red Española de Terapias Avanzadas (TERAV)-Instituto de Salud Carlos III (ISCIII),

Madrid, Spain.

3. CIBER-ONC, ISCIII, Barcelona, Spain.

4. Institució Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, Spain.

5. Department of Biomedicine. School of Medicine, University of Barcelona, Barcelona, Spain.

6. Centre for Regenerative Medicine, Institute for Regeneration and Repair, University of

Edinburgh, Edinburgh, UK

7. Centre for Discovery Brain Sciences, Edinburgh Medical School, Biomedical Sciences,

University of Edinburgh, Edinburgh, UK

8. Zhejiang University-University of Edinburgh Joint Institute, Zhejiang University School of

Medicine, Zhejiang University, Haining 310058, China

9. Edinburgh Medical School, Biomedical Sciences, University of Edinburgh, Edinburgh, UK

* Correspondence to afidanza@ed.ac.uk

Abstract

A major challenge in the stem cell biology field is the ability to produce fully functional cells

from induced pluripotent stem cells (iPSCs) that are a valuable resource for cell therapy,

drug screening and disease modelling. Here we developed a novel inducible CRISPR-

mediated activation strategy (iCRISPRa) to drive the expression of multiple endogenous

transcription factors important for *in vitro* cell fate and differentiation of iPSCs to

haematopoietic progenitor cells. This work has identified a key role for IGFBP2 in the

development of hematopoietic progenitors. We first identified nine candidate transcription

factors that we predicted to be involved in blood cell emergence during development, then

generated tagged gRNAs directed to the transcriptional start site of these transcription

factors that could also be detected during scRNAseq. iCRISPRa activation of these

endogenous transcription factors resulted in a significant expansion of arterial-fated

endothelial cells expressing high levels of IGFBP2 and our analysis indicated that IGFBP2 is

involved in the remodeling of metabolic activity during *in vitro* endothelial to hematopoietic

transition. As well as providing fundamental new insights into the mechanisms of

haematopoietic cell fate and differentiation, the broader applicability of iCRISPRa provides a

valuable tool for studying dynamic processes controlling developmental events and for

recapitulating abnormal phenotypes characterised by ectopic activation of specific

endogenous gene expression in a wide range of systems.

Introduction

The haematopoietic system develops early during gestation through so called "waves" of

hematopoietic progenitors that arise in different anatomical regions and result in the

production of various progenitor and stem cells [1–5]. The precise signalling pathways leading

to the production of hematopoietic stem and progenitor cells (HSPCs) during embryonic

development are yet to be completely defined, posing a limitation on how to recapitulate the

process *in vitro* from pluripotent stem cells.

During development, HSPCs are generated by a subset of endothelial cells, known as

hemogenic endothelium [6–9], via the endothelial to hematopoietic transition (EHT) [10]. During

the EHT, endothelial cells undergo profound transcriptional remodelling whereby the

expression of endothelial genes is gradually downregulated, and the transcription of the

hematopoietic program is initiated [11] while the cells round up and eventually detach to enter

the circulation [12,13].

To explore the molecular control on the development of the hematopoietic system and to

address the differences with the *in vitro* differentiation of induced pluripotent stem cells

(iPSCs), we compared our single cell transcriptomics analysis of *in vitro* derived hemogenic

endothelium and early progenitors [14] to that of the *in vivo* HSC-primed human hemogenic

endothelium [15]. We then developed a novel DOX-inducible CRISPR gene activation system

to assess the role of the genes that were expressed at a lower level within *in vitro*-derived

cells compared to their *in vivo* counterparts. We employed single-cell RNA sequencing to

track the presence of guide RNAs and we monitored the phenotypic effects of gene

activation. With this experimental pipeline, we identified and functionally validated a novel

role for IGFBP2, IGF Binding protein 2, in the development of *in vitro* progenitors and

showed that IGBP2 remodels the metabolic activity during *in vitro* EHT.

Results

Comparison of iPSC-derived endothelial cells with AGM AGM endothelial cells dataset

identifies 9 differentially expressed transcription factors.

We, and others, have shown that *in vitro* differentiation of human iPSCs is a powerful tool to

model intraembryonic hematopoiesis [14,16–18]. To understand the molecular basis underlying

the challenges associated with the *in vitro* production of blood stem and progenitor cells *in*

vitro from differentiating iPSCs, we compared our scRNAseq dataset from differentiating

iPSCs[14] to that of cells derived *in vivo* from the human aorta-gonad-mesonephros (AGM)

region [15]. We integrated the transcriptomic data of *in vitro* derived endothelial (IVD_Endo)

and hematopoietic cells (IVD_HPC) with that of arterial endothelial cells (aEC), arterial

hemogenic cells (aHEC) and venous endothelial cells (vEC) derived from human embryos

collected between Carnegie stage 12 and 14 (Figure 1A-B). To identify possible target

genes that could be manipulated *in vitro* to improve iPSCs differentiation, we first determined

the transcription factors marking the aHEC *in vivo* (Table 1). Then, we filtered the

transcription factor genes that were expressed at lower levels in the IVD_Endo and

IVD_HPC. This strategy identified 9 target transcription factors *RUNX1T1*, *NR4A1*, *GATA2*,

SMAD7, *ZNF124*, *SOX6*, *ZNF33A*, *NFAT5* and *TFDP2* (Figure 1C), whose expression was

detected across other AGM datasets in the hemogenic endothelium[17] (Supplementary

Figure 1).

Development of a DOX-inducible dCAS9-SAM activation system in human iPSCs

We previously developed an all-in-one Synergistic Activator Mediator system, UniSAM, that

mediates the transcriptional activation of endogenous gene expression[19]. To activate the

nine target genes identified in this study, we developed a novel doxycycline-inducible SAM

(iSAM) cassette targeted into the *AAVS1* locus of human iPSCs (Figure 2A). We

demonstrated this strategy could activate of *RUNX1C* expression in HeLa cells and that this

was correlated with the doxycycline (DOX) concentration in a linear manner (Figure S2A-B).

To verify gene activation in human PSCs at single-cell resolution, we employed a RUNX1C-

GFP human embryonic stem cell (hESC) reporter cell line (Figure S2C-G). As predicted, the

level of expression of the mCherry tag, the fluorescent mCherry reporter tag within the iSAM

cassette, was proportional to the concentration of DOX (Figure S2D-E) and to the number of

cells in which RUNX1C-GFP was activated (Figure S2 D-F). Furthermore, the RUNX1C

expression level, measured by the mean fluorescence intensity (MFI) of the RUNX1C-GFP

reporter, also correlated with the concentration of DOX (Figure S2G). We then tested the

iSAM cassette in the iPSC line (SFCi55) (Figure 2B-D). Only when iPSCs were transfected

with both iSAM and the gRNA directed to *RUNX1C* and treated with DOX, was the

expression of the *RUNX1C* gene and RUNX1 protein detected (Figure 2B-C).

We then targeted the iSAM cassette into the *AAVS1* locus using a Zinc Finger Nuclease

(ZFN) strategy [14,20,21]. iPSC clones that had specifically integrated the iSAM cassette into the

AAVS1 locus were validated by genomic PCR screening (Figure S2A-B) and sequencing.

The *AAVS1* locus has been reported to be a "safe harbour" site that is resistant to epigenetic

silencing and indeed, we had previously demonstrated that transgenes inserted into the

AAVS1 locus under the control of the constitutively active CAG promoter were efficiently

expressed both in undifferentiated and in differentiated iPSCs [20,22–24]. However, after the

iSAM line had been established and cultured under self-renewal conditions, we noted a

dramatic reduction in the number of mCherry+ cells in undifferentiated iSAM iPSCs upon

DOX induction. We predicted this to be due to transgene-silencing of the rTTA DOX-

inducible cassette (Figure S3C). To overcome this problem, we treated the iSAM iPSC line

with an inhibitor of histone deacetylases (HDACs), sodium butyrate (SB), reported to have

no adverse effect on iPSC maintenance [25,26] and we also confirmed that the treatment with

SB had no effect on viability and cell proliferation in our culture conditions (Figure S3F-G). A

short 48-hour treatment significantly increased the number of mCherry+ cells upon DOX

induction, proportional to the SB concentration (Figure S3D-E). We, therefore, maintained

the iSAM iPSCs in the presence of SB and this fully restored the inducibility of the transgene

with virtually all cells expressing mCherry in the presence of DOX (Figure S3G). We did not

treat the cells with SB during the differentiation since we detected robust expression of the

iSAM cassette during the differentiation (Figure S4F). Furthermore, since the HDACs are

also involved in the chromatin remodelling during the EHT process[27] we predicted that SB

treatment would negatively affect the differentiation.

To test the effect of activating the 9 target genes on the transcriptomes of differentiating

iPSC cells, we engineered the gRNAs to allow their detection within the single cell RNA

sequencing pipeline [28]. We inserted a capture sequence prior to the termination signal to

avoid any alteration in the secondary structure of the loops thus preserving the binding of the

synergistic activators of the SAM system to the stem loops of the gRNAs. Of the two capture

sequences available[28], we decided to use the one that was predicted to result in fewer

secondary structure alterations and this new gRNA was named 2.1 (Figure 2E). We

compared the activation level achieved with the new 2.1 gRNA to that of the original 2.0

backbone using various gRNAs targeting *RUNX1C* (Figure 2F). These results convincingly

demonstrated that the addition of the capture sequence in the gRNA 2.1 does not alter the

level of endogenous gene activation that could be achieved (Figure 2G). Altogether, these

results show that the iSAM system is able to induce gene expression that predictably

translates into an increased protein expression and thus provides a platform to steer

phenotypical changes in cell identity.

To activate our target genes, we designed 5-7 gRNAs in the 200bp upstream of the
transcriptional start sites of each of the 9 target genes. We subcloned a total of 49 gRNAs
(Table S1) into the gRNA 2.1 backbone and packaged them into lentiviral particles, (herein
referred to as the AGM library) as well as a non-targeting (NT) gRNA that was used as
control. The iSAM iPSC line was transduced with the targeting gRNAs or the control non-
targeting gRNA to generate the iSAM_AGM and iSAM_NT iPSCs line, respectively (Figure
S4A). After puromycin selection, their integration in the genome was also confirmed by PCR
and sequencing (Figure S3H-I).
154
155   **Single Cell RNA sequencing in combination with CRISPR activation identifies arterial**
156   **cell type expansion in association with higher hematopoietic progenitor potential**
157
To assess the transcriptional changes in response to the activation of the target genes, we
differentiated the iSAM iPSCs, induced with DOX and subjected them to single-cell RNA
sequencing using the 10X pipeline. After 10 days of differentiation, in the presence or
absence of DOX, we FAC-sorted live CD34+ cells from iSAM_AGM and the iSAM_NT
iPSCs (Figure S4B). Following data filtering, we selected cells in which the gRNAs
expression was detected and showed that our approach activated all the target genes,
except for ZFN124. A higher level of expression of *RUNX1T1*, *NR4A1*, *GATA2*, *SMAD7*,
SOX6, *ZNF33A*, *NFAT5*, *TFDP2* was observed following DOX treatment of cells of
iSAM_AGM cells compared to the iSAM_NT cells (Figures 3A, S3G). To study the effect of
gene activation on transcriptional and cellular phenotype, we performed clustering analysis
and detected a total of 7 clusters (Figure 3B-E). A high level of *GJA4* and *DLL4* expression
was used to annotate the arterial-like cell cluster, while high levels of hemogenic markers,
such as *RUNX1* and *CD44*, were used to annotate the hemogenic clusters, EHT_1 and
EHT_2 (Figure 3B, C-D). All the clusters expressed pan-endothelial markers such as
PECAM1 and *CDH5*, coding for CD31 and VECAD, respectively, which get progressively
downregulated in EHT_1 and EHT_2, as expected for cells undergoing the EHT process.
To assess the effect of the activation on cell identity, we analysed the proportion of the cell
clusters between the libraries and detected a significant expansion of the arterial cluster in
the DOX-induced iSAM-AGM compared to the iSAM_NT cells (Figures 3E, S4D, S4I). To
validate the effect of the activation on the expansion of arterial cell population, we analysed
their prevalence by flow cytometry. Although DOX treatment resulted in an average 1.63 ±
0.13 fold increase of CD34$^+$DLL4$^+$ cells in the control iSAM_NT sample (Figure S3A), the
increase observed in the iSAM-AGM cells was significantly greater, with an expansion of
3.33 ± 0.73 fold increase of CD34$^+$DLL4$^+$ immunophenotypic arterial cells identified by flow
cytometry (Figure 3F). To assess the effect of the different cell composition in the CD34+

compartment upon activation on the emergence of colony forming progenitors, we isolated

CD34$^+$ cells using magnetic beads and cocultured 20,000 cells on OP9 supportive stromal

cells for 7 days in the presence of hematopoietic differentiation cytokines. After one week,

the progenitor cells were assessed by colony-forming unit (CFU) assays and scored 14 days

later. We detected an increased number of CFU-E (colony-forming unit erythroid) and CFU-

GM (colony-forming unit granulocyte/macrophage) and a reduction of CFU-M (colony-

forming unit macrophage) in iSAM-AGM samples cultured in the presence of DOX compared

to the absence of DOX but no significant effect of DOX in SAM_NT samples (Figure 3G).

These data indicate that activation of target transcription factors using our novel CRISPR

strategy results in transcriptional remodelling and a steer in cell identity that we detected as

a functional difference in the hematopoietic progenitor profile.

The addition of IGFBP2 to the *in vitro* differentiation leads to a higher number of

functional hematopoietic progenitor cells.

To better understand the molecular mechanism behind the increased progenitor

development, we compared the expression profile of the arterial cells between the different

activation libraries, and we obtained a list of genes upregulated upon activation of the

targets (Table 1). The most upregulated gene, *IGFBP2,* was expressed at significantly

higher levels in the iSAM_AGM library in the presence of DOX compared to the others

(Figure 4A, Table 1). Although the arterial cells expressed *IGFBP2* at the highest level

compared to other cell types, the activation of the gene was not cell-type specific and it was

detected across the various clusters (Figure S4H). We then compared the gRNA distribution

in these arterial cells from the iSAM_AGM treated with DOX to that of arterial cells without

activation. We observed a significant enrichment of the *RUNX1T1-specific* gRNAs (Table1),

indicating that the increased *IGFBP2* expression could be downstream of *RUNX1T1*

activation. IGF Binding Protein 2 is a member of the family of IGF binding proteins and is

thought to be secreted from cells where it then binds IGF1, IGF2, and other extracellular

matrix proteins, modulating their function. IGF1 and IGF2 are commonly used in

differentiation protocols[29–31], including ours[32,33], due to their direct role in blood development.

To test if the increased frequency of functional hematopoietic progenitors was due to

IGFBP2 signalling, presumably derived from the arterial cells, we supplemented the media

with IGFBP2 at 100ng/ml after the induction of endothelial cell differentiation. To explore the

role of IGFBP2 we employed the parental iPSCs line, SFCi55 from which the iSAM line was

derived. We isolated CD34+ cells at day 8 and co-cultured them on OP9 cells in presence of

IGFBP2 for one week, then tested for their hematopoietic clonogenic potential using CFU

assays (Figure 4B). Cells treated with IGFBP2 showed a significant increase in the total

number of haematopoietic CFU colonies compared to cells in control cultures. To assess
whether IGFBP2 also affected the production of arterial cells themselves, we analysed the
proportion of DLL4+ cells, but no difference was detected in the presence of IGFBP2 (Figure
4C). This implies that the mechanism of action of IGFBP2 is different from that mediated by
the gene activation that led to an expansion of the arterial population. To further understand
the relationship between colonies' potential with the arterial identity, we sortedCD34+ into
DLL4$^+$ and DLL4$^-$ and cultured them on OP9 co-culture for a week prior to methylcellulose
assay. CD34$^+$DLL4$^+$ and CD34$^+$DLL4$^-$ plated on OP9 showed different capacity to generate
suspension cells (Figure S5D), in line with the colonies' formation results showing that the
DLL4$^-$ contains the largest progenitor activity. This aligns with the observation that the EHT
process coincides with the downregulation of arterial markers such as *DLL4* and the
upregulation of hemogenic markers such as *RUNX1* (Figure 3C).
This observation supports our hypothesis that the change in hematopoietic progenitor
production following activation in the iSAM_AGM line is associated with both differential
gene expression within the arterial cells rather and their expansion. We then focused on the
characterization of the cells derived from the CD34+ cells after coculture with the OP9 in the
presence of IGFBP2. Our results showed a significant expansion of the CD34+ and CD43+
cell populations, further supporting our hypothesis (Figure 4D). To address the potential role
of *IGFBP2 in vivo*, we analysed single cell sequencing data from the human AGM at
Carnegie stages 14 and 15. *IGFBP2* is highly expressed within the AGM niche by stromal
and epithelial cells and, most importantly, highest in endothelial cells (Figure 4E). Together
these data show that IGFBP2 addition results in increased hematopoietic blood production *in*
vitro and that the endothelial compartment is the most likely source of IGFBP2 both *in vitro*
and *in vivo*.
244
245 **IGFBP2 enhances metabolic dependency on oxidative phosphorylation of**
246 **differentiating endothelial cells.**
247
Following the observation that IGFBP2 supports haematopoietic progenitor differentiation *in*
vitro from human iPSCs we performed a time course single-cell RNA experiment of SFCi55
iPSCs differentiated in its presence. We isolated CD34$^+$ cells and plated them in EHT culture
on laminin[34], rather than on OP9 support, to assess exclusively the specific effects of
IGFBP2- on differentiating iPSCs. FAC-sorted single/live adherent cells from day 10 and 13
in the presence and absence of IGFBP2 were subjected to single-cell RNAseq (Figure S4C).
Our time course transcriptomic analyses showed that IGFBP2 induced a change in the
transcriptome, specifically on day 13 (Figure 5A). A cluster of endothelial cells enriched for
genes associated with the KEGG pathway of growth factor binding, was detected almost

exclusively at day 13 (Figure S5F); these cells displayed a different transcriptional signature

(as indicated by a shift in their position in the UMAP embedding) in the presence of IGFBP2,

and were enriched upon treatment (Figure S5F). Interestingly, this cluster showed

expression of *GJA4* across the cells, indicating their broad arterial identity, with a specific

reduction of *DLL4* expression level concomitantly to an upregulation of *RUNX1* in the region

of the cluster induced by IGFBP2 (Figure 5B). This increase in hemogenic identity within the

endothelial cell compartment is consistent with the observation of increased functional

progenitor production. This suggests that IGFBP2 supports arterial cells in the acquisition of

hemogenic capacity.

We compared the transcriptome of cells at day 13 in the presence and absence of IGFBP2

and, using KEGG enrichment analysis, we observed that the genes that were upregulated

by IGFBP2 were highly enriched in the oxidative phosphorylation term (Figure 5C-D). Since

the metabolic switch between glycolytic to mitochondrial metabolism has been previously

reported in definitive hematopoiesis[35,36], we tested whether the addition of IGFBP2 could

result in a different ATP production profile. We analysed the ATP production at day 13, when

RUNX1 expression was induced by IGFBP2 addition, by using specific inhibitors of the

complex I and II and complex V to quantify the intracellular ATP and mitochondrial

synthesis, respectively. We detected a reduction in glycolytic-derived ATP (Figure 5E-F),

which translates to a higher contribution of mitochondria metabolism for IGFBP2-treated

cells (Figure 5G). We then looked at the expression levels of genes encoding the enzymes

of the glycolytic pathway and its checkpoints, glucose and lactate transporters,

monocarboxylate transporters and the enzymes associated with hexokinase and

phosphofructose reactions. We observed a general downregulation of the glycolytic

enzymes and its checkpoints, with the exception of *PFKM* and *SLC16A1*, (Figures S5A-B),

which were expressed at lower levels in the IGFBP2-treated cells at day 13 compared to the

control on the same day. These differences were not observed on day 10 cells, which is

consistent with our previous observations that the IGFBP2-induced remodelling of the

endothelial cells transcriptome happens exclusively on day 13.

To assess whether the higher clonogenic potential and effect on the bioenergetic profile of

the cells treated with IGFBP2 was the result of an increase in proliferation, we analysed the

cell cycle profile of suspension hematopoietic cells obtained from the OP9 cocultured in the

presence of IGFBP2. No differences in the cell cycle distribution of hematopoietic

progenitors (Figure S3C) was observed, indicating that the detected increase in

hematopoietic progenitors is not likely to be a consequence of their increased cycling. We,

therefore, concluded the effect of IGBP2 occurs prior to, or during their emergence via the

EHT process.

In summary, our results show that the activation of the 9 target transcription factors,
expressed at higher levels in vivo within the arterial cells in the AGM region, leads to the
expansion of the arterial cells and, consequently, to an increase in progenitor activity (Figure
6). IGFBP2 was identified as the most upregulated gene in the arterial cell upon gene
activation. The addition of IGFBP2 to the differentiation culture induced the upregulation of
RUNX1 and OxPhos genes and accelerated the metabolic dependency on OXPHOS in
association with improved progenitor activity (Figure 6).
300
301 **Discussion**
302
The complexity and dynamism of developmental hematopoiesis *in vivo* have imposed
challenges in accurately reproducing the process *in vitro*. We hypothesized that this could, in
part, be due to the inability to recapitulate the appropriate transcriptional programme in
cultured cells. Here, we developed a novel CRISPR-activation system to induce the
expression of genes that are expressed at low levels in cells that are generated *in vitro*, to
explore the downstream consequences of their activation and to assess the effects on
emerging hematopoietic progenitor cells.
When we compared endothelial cells derived *in vitro* from hiPSCs to those in the AGM
region of the human embryo, at the time point when early hematopoietic commitment takes
place, we identified nine transcription factors that were expressed at a lower level in the
cultured cells. Some of these had been associated with blood cell development, including
GATA2[37–39]*, SMAD7* [40]*, NR4A1*[40]*, SOX6*[41] and *RUNX1T1*[15,17]*. Other genes such as,
ZNF124, ZNF33A, NFAT5 and TFDP2 had not been previously associated with
hematopoiesis and could provide early evidence of their possible role in hematopoiesis that
will require further studies. We were particularly interested in *RUNX1T1* (also known as
ETO) which is associated with the leukemic fusion protein, AML1/ETO resulting from the
t(8;21) chromosomal translocation[42]. Furthermore, *RUNX1T1* expression has been recently
detected in transcriptomic analyses of the human AGM region[15,17], but its precise role during
the ontogeny of the blood system has not been elucidated. The addition of a capture
sequence to the gRNA backbone enabled their detection coincidentally with the single-cell
transcriptome, and this allowed us to demonstrate that RUNX1T1 gRNAs were significantly
enriched in cells within an expanded arterial cluster and their presence was associated with
the highest expression of *IGFBP2* across clusters. *IGFBP2* KO mice show increased
expression of cell-cycle inhibitors and HSC apoptosis, implicating IGFBP2 as a modulator of
HSCs cell cycle and survival [43]. More recently *IGFBP2* was reported as being highly
expressed in the human AGM region at CS14 when HSCs are emerging [17], supporting a

possible role during developmental hematopoiesis but the precise molecular process was

unclear. In this study, we show that the addition of IGFBP2 recombinant protein in our *in*

vitro model of EHT results in the emergence of an increased number of functional

hematopoietic progenitors, and we provide an early observation to suggest that RUNX1T1

could be involved in the regulation its expression. Because RUNX1T1 lacks a DNA binding

domain, its direct involvement in the regulation of IGFBP2 expression, or in the development

of the hematopoietic system in general, must require association with other cofactors that

are yet to be identified.

Our data, together with *in vivo* data[17], show that IGFBP2 is expressed predominantly by

endothelial cells. The supportive role of the endothelial niche in the development of HSCs

has been studied *in vivo* [44–46] and exploited *in vitro* to support HSCs emergence [45,47].

We describe here a novel role for IGFBP2 in the remodelling of the metabolism of iPSC-

derived endothelial cells and that this is associated with the induction of RUNX1, a hallmark

of hemogenic fate. The addition of IGFBP2 in the culture induces upregulation of the genes

of the oxidative phosphorylation pathway in association with an increased relative

mitochondrial contribution to cellular ATP production. This is due to reduced glycolysis in

association with the downregulation of genes coding for glycolytic enzymes and their

checkpoints. The switch between glycolytic to mitochondrial metabolism has been shown to

be essential for the EHT in definitive hematopoiesis[36,48]. This switch is induced *in vivo* by

mechanical cues downstream of the establishment of circulation, and it is required for

functional HSCs development[35], while *in vitro*, the switch is driven by pyruvate mitochondrial

catabolism, leading to definitive EHT as opposed to primitive[36]. Further *in vivo* studies are

required to characterize the role of IGFBP2 during AGM hematopoiesis to overcome the

limitation of using an *in vitro* model such as iPSCs differentiation.

In conclusion, we detected transcriptional differences between *in vivo* and *in vitro*

developmental hematopoiesis and developed a novel inducible gene activation system to

identify novel molecular players during the endothelial-to-hematopoietic transition. Our

multidisciplinary approach identified IGFBP2 as novel signalling molecules that support

human blood progenitor development *in vitro*, inducing a metabolic switch from cytoplasmic

glycolysis to mitochondrial respiration. This study demonstrates that combining CRISPR-

mediated activation of target genes with single-cell transcriptomic analysis in differentiating

hPSCs can be a powerful approach to alter cell fate, providing a tool for gene function

studies during human development. The fine epigenetic manipulation of the transcription can

be readily applied to any cell lineage simply by adding specific gRNAs and it will be

instrumental in exploring other developmental processes that can be, at least partially, be

mimicked *in vitro* with human iPSCs. Some limitations remain in applying CRISPR-mediated

gene activation strategies in long differentiation protocols due to the challenge of detecting

high copy numbers of the gRNAs, limiting the possibility of providing the fine statistical

correlation needed to predict downstream target genes. Testing different promoters driving

the gRNA expression or structural modifications of the gRNA scaffold could result in a more

robust expression and allow for correlation analysis.

Finally, using this approach, we have identified the supportive role of IGFBP2, predominantly

produced by endothelial cells, which induces transcriptional and metabolic remodelling in

association with the induction of *RUNX1* expression, and result in higher hematopoietic

progenitors' activity.

Figure 1 - Comparison of *in vitro* iPSC-derived and *in vivo* AGM-derived endothelial

cells identifies 9 differentially expressed transcription factors.

A - Schematic of the analytic pipeline used to identify the target genes. **B** - Integrative

analysis of single cell transcriptome of *in vitro* derived endothelial (IVD_Endo) and

hematopoietic cells (IVD_HPCs) with *in vivo* developed endothelial cells (venous, vEC;

arterial, aEC; arterial hemogenic, HECs) from human embryos (CS12-CS14) visualised on

UMAP dimensions. **C** - Target genes expression level showing higher expression in arterial

hemogenic endothelium *in vivo* than *in vitro* derived cells.

Figure 2 - The inducible iSAM cassette successfully mediates activation of

endogenous gene expression upon DOX induction.

A - Schematic of the iSAM cassette containing the TET-on system under the control of EF1α

and dCAS9-P2A-MS2-p65-HSF1-T2A-mCherry under the rTTA responsive elements,

separated by genetic silencer and flanked by AAVS1 specific homology arms. **B** - *RUNX1C*

gene expression activation after transient transfection of the iSAM plasmid and gRNAs in

presence or absence of DOX in human iPSC line (n=3 from independent transfections). **C** –

RUNX1 protein expression upon iSAM activation after transient transfection of the iSAM

plasmid and gRNAs with DOX in human iPSC line detected by immunostaining. **D**

Expression of the iSAM cassette reported by mCherry tag during the differentiation protocol,

the representative images (bright field – BF, and fluorescence) show embryoid bodies at day

3 of differentiation. **E** - Schematic of the gRNA 2.1 containing the capture sequence for

detection during the scRNAseq pipeline. **F** - *RUNX1C* gene activation level obtained using

either the gRNA 2.0 or 2.1 backbone (n=3 from independent transfections of the 4 different

gRNAs). **G** - Statistical analysis of the gRNAs activation level showing no significant

variation following addition of the capture sequence (n=3 for each of the 4 different gRNAs).

Figure 3 - Single Cell RNA sequencing in combination with CRISPR activation

identifies arterial cell type and functional hematopoietic expansion in association with

activation of the 9 target genes.

A - Gene expression profile of target genes following target genes' activation, heatmap

shows the expression level of the target genes in the iSAM_NT and iSAM_AGM treated with

DOX following normalisation on the -DOX control. **B** - Dimension reduction and clustering

analysis of the scRNAseq data following activation, filtered on cells where the gRNA

expression was detected. **C** - Arterial (*GJA4*, *DLL4*), venous (*NRP2*, *APLNR*) and

hemogenic marker (*CD44*, *RUNX1*) expression distribution in the clusters indicated by the

colour. **D** - Expression distribution visualised on the UMAP plot showing the location of

arterial cells marked by *DLL4*, and hemogenic endothelium marked by *CD44* and *RUNX1*.

E - Heatmap of the top 15 marker genes for each of the clusters. **F** - Contribution of the

different libraries to the clusters showing that arterial cell cluster is overrepresented in the

iSAM_AGM treated with DOX, compared to the other libraries. **G** - Expansion of the arterial

population assessed by the membrane marker expression of DLL4+ following targets'

activation, quantified by flow cytometry at day 8 of differentiation (Data are normalised on

the iSAM_NT + DOX sample, n=5 independent differentiations, * p = 0.0417 paired t-test). **H**

- Colony forming potential of the suspension progenitor cells derived from the two lines

treated with or without DOX following OP9 coculture activation, data show the colony

obtained for 104 CD34+ input equivalent (n=3 from independent differentiations * p<0.05,

Tukey's two-way ANOVA).

Figure 4 - IGFBP2 addition to the *in vitro* differentiation leads to a higher number of

functional hematopoietic progenitor cells. A - Violin plot of *IGFBP2* expression profile in

the arterial cells obtained from the different conditions, in the presence or absence of gRNAs

and DOX. **B** -Number of hematopoietic colonies obtained after coculture on OP9 in presence

or absence of IGFBP2 (n=3-4 from independent differentiations, ** p=0.0080, Sidak's Two

way ANOVA). **C** – Percentage of DLL4[+] arterial cells differentiation within the CD34[+]

compartment analysed by flow cytometry in day 8 EBs (n=4 from independent

differentiations, Two-way Anova, ns =p>0.99). **D** – Expansion of hematopoietic progenitors

analysed using markers' expression on suspension progenitors derived after coculture of

CD34+ cells onto OP9 support (data are expressed as fold over the CTR in the absence of

IGFBP2 (n=4 from independent differentiations, * p<0.02, Sidak's Two way ANOVA. **E** -

Single-cell transcriptomic analysis of developing AGM collected from human embryos at

Carnegie Stages14 and 15 enriched for CD31[+] and CD34[+] showing the *IGFBP2* expression

profile *in vivo* in the AGM.

Figure 5 - IGFBP2 alters cell metabolism by inducing a reduction in glycolytic ATP production.

A – Clustering analysis of the single cell transcriptomic time course analysis of differentiating cells at day 10 and day 13 in the absence (CTR) or presence of IGFBP2. Arrows indicate the difference in the clustering due to the addition of IGFBP2 compared to control. **B** – Expression profile of arterial markers, *GJA4* and *DLL4*, and hemogenic marker *RUNX1* (top - the dashed line shows the location of the shift in gene expression of cells treated with IGFBP2) and their expression profile in the endothelial cells cluster marked by Growth factor binding in absence (CTR) and in presence of IGFBP2 (*GJA4* p=1E^{-54}, *DLL4* p=1.2E^{-119}, *RUNX1* p=8.2E^{-163}). **C –** KEGG enrichment analysis of the genes upregulated at day 13 upon IGFBP2 treatment. The arrow shows the ranking of the Oxidative Phosphorylation pathway. **D** - Dot Plot showing the expression profile of the genes coding for the enzyme of the Oxidative Phosphorylation pathway. **E** -Oxygen Consumption Rate (OCR) and Extracellular Acidification Rate (ECAR) profile in cells at day 13 of differentiation reporting mitochondrial respiration and glycolysis, respectively. **F** – ATP production rate divided by that deriving from glycolysis and from mitochondrial respiration, in cells treated with IGFBP2 and controls at day 13. **G** – Ratio of the ATP production between glycolysis and mitochondrial respiration in cells treated with IGFBP2 and controls at day 13.

471
472

Figure 6 - Model summarising the results.
During development, some endothelial cells undergo arterialisation, as identified by their
arterial genes' expression profile (e.g. *DLL4*). Arterial cells, characterised by high
dependency on glycolysis, are expanded by our CRISPR activation approach, resulting in
more blood production since these cells are the cell-of-origin of the hemogenic endothelium.
Once arterial cells commit to hemogenic endothelium fate, they start to express
hematopoietic genes (e.g. *RUNX1*); this process is enhanced by IGFBP2 via induction of
both *RUNX1* expression and increased dependency on Oxidative Phosphorylation, known to
be important for the progression of the endothelial to hematopoietic transition.

Methods

Resource availability

R code is available at https://github.com/afidanza/CRISPRa. Raw and processed data have

been deposited to ArrayExpress (E-MTAB-12748 – currently on hold). The AAVS1-iSAM are

available on Addgene (Addgene #211495) and the gRNA 2.1 plasmids (Addgene #211496)

(both currently on hold). Further information and requests for resources and reagents should

be directed to and will be fulfilled by the corresponding author.

Pluripotent Stem Cells maintenance

hPSCs were maintained *in vitro* in StemPro hESC SFM (Gibco) with bFGF (R&D) at 20

ng/ml. Wells were coated with Vitronectin (ThermoFisher Scientific) at least 1 hour before

plating and cells were passaged using the StemPro EZPassage tool (ThermoFisher

Scientific). Media change was performed every day and cells passaged every 3–4 days at a

ratio of 1:4.

Transfection

iPSCs SFCi55 and hESCs RUNX1-GFP were plated at 3×10^5 cells per a well of a 6 well

plate and reverse transfected with 2 µg of DNA using the Xfect Transfection reagent

(Clontech) and analyzed 2 days later.

HeLa cells were cultured in Dulbecco's Modified Eagle Medium/Nutrient Mixture F-12

(DMEM/F12) with Glutmax and 5% FCS (Gibco) and passaged every few days, at a ratio of

1:6. HEL were cultured in Iscove's Modified Dulbecco's Medium (IMDM) with 10% FCS

(Gibco) and passaged every few days, at a ratio of 1:4. 2×10^5 cells were plated, transfected

at 6–8 hours with 0.75 µg of DNA using Xfect Transfection reagent (Clontech) and then

analysed 2 days after.

Immunocytochemistry

Cells were fixed in 4% PFA in PBS at room temperature for 10', permeabilized in PBS-T

(0.4% Triton-X100) for 20' and blocked in PBS-T with 1% BSA and 3% goat serum for

1 hour. Primary antibodies were incubated in blocking solution over night at 4 °C (RUNX1

1:200 - ab92336, Abcam). Cells were then washed in PBS-T and incubated with secondary

antibodies for 1 hour at room temperature (donkey α-rabbit 1:200 - A-11008 - Thermo

Scientific). Cells were washed in PBS-T and counterstained with DAPI. Images were

generated using the Zeiss Observer microscope.

Gene expression analysis

Total RNA was purified using the RNAeasy Mini Kit (Qiagen) and cDNA synthesized from

500 ng of total RNA using the High Capacity cDNA synthesis Kit (Applied Biosystem). 2 ng of

cDNA were amplified per reaction and each reaction was performed in triplicate using the

LightCycler 384 (Roche) with SYBR Green Master Mix II (Roche). A melting curve was

performed and analyzed for each gene to ensure the specificity of the amplification. *β-*

Actin was used as reference genes to normalize the data [19].

Pluripotent Stem Cells differentiation to hematopoietic progenitors

hPSCs were differentiated in a xeno-free composition of SFD medium [14], BSA was

substituted with human serum albumin, HSA (Irvine-Scientific). Day 0 differentiation medium,

containing 10 ng/ml BMP4 was added to the colonies prior cutting. Cut colonies were

transferred to a Cell Repellent 6 wells Plates (Greniner) to form embryoid bodies and

cultured for two days. At day 2 media was changed and supplemented with 3 μM CHIR

(StemMacs). At day 3, EBs were transferred into fresh media supplemented with 5 ng/ml

bFGF and 15 ng/ml VEGF. At day 6 media was changed for final haematopoietic induction in

SFD medium supplemented with 5 ng/ml bFGF, 15 ng/ml VEGF, 30 ng/ml IL3, 10 ng/ml IL6,

5 ng/ml IL11, 50 ng/ml SCF, 2 U/ml EPO, 30 ng/ml TPO, 10 ng/ml FLT3L and 25 ng/ml

IGF1. From day 6 onward, cytokines were replaced every two days.

CD34 isolation

CD34+ cells were isolated using CD34 Magnetic Microbeads from Miltenyi Biotec, according

to their manufacturing protocol. Briefly, Embryoid bodies were dissociated using Accutase

(Life Techonologies) at 37°C for 30'. Cells were centrifuged and resuspended in 150 μl of

PBS + 0.5% BSA + 2mM EDTA with 50 μl Fcr blocker and 50 μl of magnetic anti-CD34 at

4°C for 30'. Cells were washed using the same buffer and transferred to pre-equilibrated

columns, washed three times and eluted. After centrifugation, cells were resuspend in SFD

media, counted and plated for OP9 coculture.

OP9 coculture and colony assay

OP9 cells were maintained in α-MEM supplemented with 20% serum (Gibco) and sodium

bicarbonate (Gibco) and passaged with Trypsin every 3-4 days. The day before the co-

culture, 45.000 OP9 cells were plated for each 12 well plates' well in SFD media. The day of

the co-culture the 20.000 iSAM cells or 25.000 SFCi55 or H9 were plated in each well and

culture in SFD media supplemented with 5 ng/ml bFGF, 15 ng/ml VEGF, 30 ng/ml IL3, 10

ng/ml IL6, 5 ng/ml IL11, 50 ng/ml SCF, 2 U/ml EPO, 30 ng/ml TPO, 10 ng/ml FLT3L and 25

ng/ml IGF1 and 100ng/ml IGFBP2. Cytokines were replaced twice during one week of

coculture. At the end of the coculture, cells were collected by Trypsin and half of the well

equivalent was plated in 2 ml of methylcellulose medium (Human enriched H4435, Stemcell

Technologies). Cells were incubated in the assay for 14 days and then scored.

Laminin EHT culture

Laminin EHT culture was performed as previously described[34]. Briefly, 24 well plates were

coated for at least 2 hours with recombinant human Laminin-521 (Thermo-Fisher). Following

CD34+ isolation at day 8, 400.000 CD34+ cells were seeded in each well of a pre-coated

24-well plate in SFD media supplemented with 5 ng/ml bFGF, 15 ng/ml VEGF, 30 ng/ml IL3,

10 ng/ml IL6, 5 ng/ml IL11, 50 ng/ml SCF, 2 U/ml EPO, 30 ng/ml TPO, 10 ng/ml FLT3L and

25 ng/ml IGF1 and 100ng/ml IGFBP2.

Flow cytometry staining and cell sorting

Embryoid bodies were dissociated using Accutase (Life Techonologies) at 37°C for 30'.

Cells were centrifuged and resuspended in PBS + 0.5% BSA + 2mM EDTA, counted and

stained at 10^5 cells for a single tube. Cells were stained with antibodies for 30' at room

temperature gently shaking. Flow cytometry data were collected using DIVA software (BD).

For the sorting experiments, the cells were prepared as above and stained at 10^7 cells/ml in

presence of the specific antibodies. Sorting was performed using FACSAria Fusion (BD) and

cells were collected in PBS + 1% BSA. Data were analysed using FlowJo version 10.4.2.

Flow cytometry antibodies

For flow cytometry 10^5 cells per test were stained in 50 µl of staining solution with the

following antibodies: CD34 Percp-Efluor710 (4H11 eBioscience, 1:100), CD34 Pe (4H11

eBioscience, 1:200), CD43 APC (eBio84-3C1, 1:100), CD45 FITC (2D1 ebioscience, 1:100),

DLL4 Pe (MHD4-46 Biolegend, 1:200), CD41 PE (HIP8 Biolegend, 1:200), CD144 APC (16B1

eBioscience, 1:100), CD235a FITC (HIR2 BD Bioscience, 1:250).

iSAM plasmid generation

The iSAM plasmid was obtained by Gibson assembly of four fragments. The first fragment,

the backbone, was a DOX-inducible AAVS1 targeted plasmid expressing an E6-E7-IRES-

ZsGreen which was excised by BstBI and NdeI. The second fragment, one of the adapters,

was derived from the UniSAM plasmid that we previously generated (Addgene #99866) by

PCR with the following primer sets

FW_aggggacccggttcgagaaggggctcttcatcactagggccgctagctctagagagcgtcgaatt,

RV_ttcgggtcccaattgccgtcgtgctggcggctcttcccacctttctcttcttcttgggggctcatggtggcc. The UniSAM

cassette was obtained also from the UniSAM plasmid via digestion with BsrGI and BsiWI.

Finally, the last fragment consisting of another adapter for the Gibson was custom

synthetised and contained overlapping sequences flanking a chicken b-globin insulator that

we inserted to prevent cross-activation of the EF1α-promoter and the TRE-GV promoter

driving the iSAM. Correct assembly was verified by Sanger sequencing.

iSAM cell lines derivation

The iSAM plasmid was used together with ZNFs specific for the AAVS1 locus to mediate

specific integration in SFCi55 human iPSCs line [14,20]. Briefly, 10 μg of AAVS1-iSAM with 2.5

μg of each ZNFs, left and right, using Xfect (Takara) according to the manufacturer protocol.

Cells were selected using Neomycin. Single clones were picked, amplified, and initially

screened by mCherry expression upon DOX addition. Clones that expressed the fluorescent

tag were screened for specific integration using PCR followed by Sanger sequencing for the

correctly integrated clones. 100 ng of genomic DNA was amplified using the EmeraldAmp®

MAX HS Takara and specific primer sets (Table 1). For the specific AAVS1 integration site,

Sigma_AAVS1 - CGG AAC TCT GCC CTC TAA CG and NeoR -GAT ATT GCT GAA GAG

CTT GGC GG were used with the PCR conditions of 95 °C for 7 min, 32 cycles of 95 °C for

15 s, 57 °C for 30 s, and 72 °C for 1 min, with the final elongation step at 72 °C for 7 min. For

the Wild type locus screening, Sigma_AAVS1 - CGG AAC TCT GCC CTC TAA CG and

AAVS1_EXT3_RV – ACA CCC AGA CCT GAC CCA AA were used with the PCR cycling

conditions of 95 °C for 7 min, 30 cycles of 95 °C for 15 s, 57 °C for 30 s, and 72 °C for 2 min,

with the final elongation step at 72 °C for 6 min.

Capture sequencing addition to the gRNA backbone

The Capture sequence 2 was added to the gRNA_Puro_Backbone (Addgene #73797) by

PCR. Briefly, the capture sequence was added before the termination signal of the gRNA

followed by a BamHI site using the following PCR primers:

gRNA_FW gagggcctatttcccatgattcct,

gRNA_Cap_RV aaaaaaggatccaaaaaaCCTTAGCCGCTAATAGGTGAGCgcaccgactcggtgcc.

The gRNA backobone was replaced from the original plasmid via NdeI and BamHI digestion,

followed by ligation of the PCR produced following the same digestion. Correct integration of

the insert was verified by Sanger sequencing.

Target genes identification

Candidate genes were identified by comparing *in vivo* hemogenic endothelium [15]

(GSE135202) and *in vitro* iPSCs-derived endothelial cells that we previously generated [14]

(E-MTAB-9295). Briefly, the two datasets were merged and normalised using the R

package Seurat. Specific markers for hemogenic endothelium were identified and

transcription factors were sorted based on their GO annotation. Within those genes we

filtered those detected in more than 50% of the *in vivo* hemogenic endothelium and

expressed in less than 25% of the *in vitro* derived endothelial cells. This pipeline identified 9

target genes *RUNX1T1*, *NR4A1*, *GATA2*, *SMAD7*, *ZNF124*, *SOX6*, *ZNF33A*, *NFAT5*, *TFDP*.

AGM gRNAs library preparation

sgRNA design was performed by selecting the top candidates for on-target and off-target

score. Between 5 and 7 guides per gene were designed for *RUNX1T1*, *NR4A1*, *GATA2*,

SMAD7, *ZNF124*, *SOX6*, *ZNF33A*, *NFAT5*, *TFDP2* using the CRISPRpick tool from the

Broad Institute (https://portals.broadinstitute.org/gppx/crispick/public) (Table S1). All the

guide variants were Golden Gate cloned with the gRNA 2.1 backbone according to the

established protocol [49]. The 49 plasmids were pooled together in an equimolar ratio and the

library prep was subsequently used to produce lentiviral particles with a second-generation

production system. Briefly, the psPAX2 packaging plasmid, pMD2.G envelope, and the AGM

vector library were co-transfected using polyethyleneimine (PEI) (Polysciences, Warrington,

PA, USA) as previously detailed [50], Lentiviral particles-containing supernatants were

harvested 48–72 h post-transfection, concentrated by ultracentrifugation and titered in

hiPSCs cells.

iSAM_AGM and iSAM_NT cell line derivation

The selected iSAM clone (3.13 internal coding) was infected with viral particles containing

either the AGM library or the non targeting gRNA (NT) at a MOI of 10. The iSAM cells were

plated the afternoon before at $7x10^6$ cells into a T125 in the presence of 10 μM Rock

Inhibitor (Merk) which was maintained until the day following the infection. Cells were

infected in presence of 8μg/ml of Polybrene (Merk). Puromycin selection was initiated 36

hours post-infection and maintained during their culture until the beginning of the

differentiation. Both lines were tested for integration of the gRNAs. Briefly, 100 ng of isolated

gDNA was amplified using the PrimeSTAR MAX PCR mix (Takara) using the primers

gRNA_screening_FW and gRNA_screening_RV (Table1). Purified amplicons were

subjected to Sanger sequencing.

Single Cell RNA sequencing

For the iSAM_AGM and iSAM_NT single cell RNA sequencing experiment, embryoid bodies obtained from day 10 of differentiation were dissociated using Accutase (Life Technologies') at 37°C for 30'. For the IGFBP2 experiment, day 10 and day 13 cells from the Laminin EHT culture were detached from the adherent layer using Accutase (Life Technologies') at 37°C for 5'. From both experiments, cells were centrifuged and resuspended in CD34-Pe staining solution at a density of 10^7/ml. CD34+/live/single cells were FAC-sorted in PBS + 0.1 % BSA. Cell viability was also confirmed by Trypan blue stain for an accurate count. Around 15000 cells per sample were loaded into the 10X Chromium Controller, and single cell libraries were obtained using the Chromium single cell 3' Reagent Kits v3 (10XGenomics) according to manufacturer protocol. The four libraries were indexed using SI PCR primers with different i7 indexes to allow for demultiplexing of the sequencing data. RNA concentration was obtained using Quibit RNA HS (Thermo-Fisher). Quality of the obtained libraries were verified using LabChip GX (PerkinElmer). Libraries were sequenced using NextSeq 2000 technology (Illumina) at 50.000 reads/cell. Data were aligned to GRCh38 using the Cell Ranger dedicated pipeline (10XGenomics). Data filtering, dimension reduction, clustering analysis, differentially expressed genes and cell cycle analysis were obtained using Seurat R package (version 4.1.0)[51]. Cells were subjected to QC and filtering using both the number of genes (1000-7500) and the percentage of mitochondrial genes detected (1-15%), resulting in 9025, 10942, 9468, 13073 cells respectively for the samples iSAM_NT, iSAM_NT+DOX, iSAM_AGM and iSAM_AGM+DOX.Pseudotemporal ordering was performed using Monocle 3 R package [52]. KEGG pathways was performed using ShinyGo[53]. The gRNAs' expression matrix was used to select cells in which the expression of the gRNAs was detected. Briefly, for the libraries derived from the iSAM_NT control containing only the non-targeting gRNA, the filter was set for cells expressing one gRNA, while for the iSAM_AGM libraries was set to more than one. The code is available on GitHub at https://github.com/afidanza/CRISPRa, the raw data have been submitted to Array Express (E-MTAB-12748), and the browsable processed data will be made available at the time of publication on our website containing previous sequencing data at
https://lab.antonellafidanza.com.

ATP production analysis

At day 8 of differentiation, 20.000 CD34+ were plated for each precoated well of the Seahorse XFp mini Cell Culture Plates (Agilent) precoated with rhLaminin-521 (Thermo-Fisher). On day 13, the cells' ATP production rate was analysed with the Seahorse XF Real-

Time ATP Rate Assay kit (Agilent) according to manufacturer protocol, following

confirmation of comparable cell densities across replicates and conditions. Briefly, the media

was changed prior to the assay for the XF DMEM medium, pH 7.4, supplemented with 10

mM Seahorse XF glucose, 1mM Seahorse XF pyruvate and 2mM Seahorse XF Glutamine

(Agilent) and incubated for 45-60 minutes in a non-CO_2 incubator. Oligomycin and

Rotenone/AA solutions were prepared and added in the cartridge and finally loaded together

with the cells in the Seahorse XF Mini Analyzer using the dedicated software. Data were

collected at the end of the run, and the values of ATP production were calculated according

to the ATP Production Rate Calculation provided by Agilent. Briefly, OCR ATP was

calculated as OCR_{basal}-OCR_{oligo} averaged across the three reads for each well. Mitochondrial

ATP was calculated as OCR_{ATP} multiplied by the molecular oxygen consumption rate of 2

and by the P/O value of 2.75. For the glycolytic ATP production, we calculated the MitoPER

as the OCR_{basal}-OCR_{rot} times the CO_2 contribution Factor of 0.5. The PER was then

calculated as the ECAR times the Buffer Factor of 2.6, the volume of reaction of 2.28 μl and

the Kvol value of 1.1. The GlycoATP Production Rate was obtained by removing the

MitoPER from the PER, and the TotalATP was calculated by adding the GlycoATP and the

MytoATP. Mito/Glyco ratio was obtained by dividing their ATP production value.

Cell cycle analysis

DAPI staining and flow cytometry analysis were performed to verify the proliferation rate of

the cells. Briefly, cells were collected from the supernatant by aspiration using a pastette,

washed in PBS + 0.5% BSA + 2mM EDTA, counted and stained at 10^5 cells for a single

tube. Cells were stained 1:1 v:v with a solution of 1% NP40 and 5 μg/ml DAPI for 2 minutes

and acquired using the DIVA software (BD), and analysed using FlowJo version 10.4.2.

Author contribution

AF designed the study, performed experiments and bioinformatic analysis, wrote the paper

and led the research. PP, TV, FPL, HM performed experiments. AM, HT, MS provided

support to the experiments. NR performed bioinformatic analysis. LF and PM helped the

design of the study and the research. All authors provided essential feedback on the

experiments and the manuscript.

Acknowledgment

AF and LF acknowledge financial support from the Biotechnology and Biological Sciences

Research Council; Grant S002219/1. AF was supported by a European Hematology

Association Advanced Research Grant (EHA RAG 2021), and by the American Society for

Hematology (Research Global Award). TV and AM were supported by PhD studentships

from the Medical Research Council (Precision Medicine) and the College of Medicine and

Veterinary Medicine, respectively. FPL was supported by an Erasmus+ Traineeship Program

2016/2017. PM acknowledges financial support from a PERIS program from the Catalan

Government and a Retos collaboration project from the MINECO (RTC-2018-4603-1)

Bibliography

1.   Medvinsky, A. & Dzierzak, E. Definitive hematopoiesis is autonomously initiated by
the AGM region. *Cell* **86**, 897–906 (1996).
2.   Palis, J., Robertson, S., Kennedy, M., Wall, C. & Keller, G. Development of erythroid
and myeloid progenitors in the yolk sac and embryo proper of the mouse. *Development*
126, 5073–84 (1999).
3.   Patel, S. H. *et al.* Lifelong multilineage contribution by embryonic-born blood
progenitors. *Nature 2022 606:7915* **606**, 747–753 (2022).
4.   Böiers, C. *et al.* Lymphomyeloid Contribution of an Immune-Restricted Progenitor
Emerging Prior to Definitive Hematopoietic Stem Cells. *Cell Stem Cell* **13**, 535–548
(2013).
5.   Hoeffel, G. *et al.* C-Myb(+) erythro-myeloid progenitor-derived fetal monocytes give
rise to adult tissue-resident macrophages. *Immunity* **42**, 665–78 (2015).
6.   Jaffredo, T., Gautier, R., Eichmann, A. & Dieterlen-Lièvre, F. Intraaortic hemopoietic
cells are derived from endothelial cells during ontogeny. *Development* **125**, 4575–83
(1998).
7.   Zovein, A. C. *et al.* Fate Tracing Reveals the Endothelial Origin of Hematopoietic
Stem Cells. *Cell Stem Cell* **3**, 625–636 (2008).
8.   Bertrand, J. Y. *et al.* Haematopoietic stem cells derive directly from aortic
endothelium during development. *Nature* **464**, 108–111 (2010).
9.   Boisset, J. C. *et al.* In vivo imaging of haematopoietic cells emerging from the mouse
aortic endothelium. *Nature* **464**, 116–120 (2010).
10. Ottersbach, K. Endothelial-to-haematopoietic transition: an update on the process of
making blood. (2019) doi:10.1042/BST20180320.
11. Swiers, G. *et al.* Early dynamic fate changes in haemogenic endothelium characterized
at the single-cell level. *Nature Communications 2013 4:1* **4**, 1–10 (2013).
12. Kissa, K. & Herbomel, P. Blood stem cells emerge from aortic endothelium by a novel
type of cell transition. *Nature* **464**, 112–115 (2010).
13. Eilken, H. M., Nishikawa, S. I. & Schroeder, T. Continuous single-cell imaging of
blood generation from haemogenic endothelium. *Nature* **457**, 896–900 (2009).
14. Fidanza, A. *et al.* Single cell analyses and machine learning define hematopoietic
progenitor and HSC-like cells derived from human PSCs. *Blood* **136**, 2893–2904
(2020).
15. Zeng, Y. *et al.* Tracing the first hematopoietic stem cell generation in human embryo
by single-cell RNA sequencing. *Cell Res* **29**, 881–894 (2019).
16. Ng, E. S. *et al.* Differentiation of human embryonic stem cells to HOXA+ hemogenic
vasculature that resembles the aorta-gonad-mesonephros. *Nat Biotechnol* **34**, 1168–
1179 (2016).
17. Calvanese, V. *et al.* Mapping human haematopoietic stem cells from haemogenic
endothelium to birth. *Nature 2022 604:7906* **604**, 534–540 (2022).
18. Sturgeon, C. M., Ditadi, A., Awong, G., Kennedy, M. & Keller, G. Wnt signaling
controls the specification of definitive and primitive hematopoiesis from human
pluripotent stem cells. *Nat Biotechnol* **32**, 554–561 (2014).
19. Fidanza, A. *et al.* An all-in-one UniSam vector system for efficient gene activation. *Sci*
Rep **7**, 6394 (2017).
20. Yang, C.-T. *et al.* Activation of KLF1 Enhances the Differentiation and Maturation of
Red Blood Cells from Human Pluripotent Stem Cells. *Stem Cells* **35**, 886–897 (2017).
21. Lopez-Yrigoyen, M. *et al.* A human iPSC line capable of differentiating into
functional macrophages expressing ZsGreen: A tool for the study and in vivo tracking

of therapeutic cells. *Philosophical Transactions of the Royal Society B: Biological*
Sciences **373**, (2018).
22. Lopez-Yrigoyen, M. *et al.* Genetic programming of macrophages generates an in vitro
model for the human erythroid island niche. *Nat Commun* **10**, 881 (2019).
23. Lopez-Yrigoyen, M. *et al.* A human iPSC line capable of differentiating into
functional macrophages expressing ZsGreen: A tool for the study and in vivo tracking
of therapeutic cells. *Philosophical Transactions of the Royal Society B: Biological*
Sciences **373**, (2018).
24. May, A. *et al.* Modelling the erythroblastic island niche of dyserythropoietic anaemia
type IV patients using induced pluripotent stem cells. *Front Cell Dev Biol* **11**, (2023).
25. Kang, S. J., Park, Y. Il, So, B. & Kang, H. G. Sodium butyrate efficiently converts
fully reprogrammed induced pluripotent stem cells from mouse partially
reprogrammed cells. *Cell Reprogram* **16**, 345–354 (2014).
26. Zhang, Z., Xiang, D. & Wu, W. S. Sodium Butyrate Facilitates Reprogramming by
Derepressing OCT4 Transactivity at the Promoter of Embryonic Stem Cell–Specific
miR-302/367 Cluster. *Cell Reprogram* **16**, 130 (2014).
27. Thambyrajah, R. *et al.* HDAC1 and HDAC2 Modulate TGF-β Signaling during
Endothelial-to-Hematopoietic Transition. *Stem Cell Reports* **10**, 1369–1383 (2018).
28. Replogle, J. M. *et al.* Combinatorial single-cell CRISPR screens by direct guide RNA
capture and targeted sequencing. *Nat Biotechnol* (2020) doi:10.1038/s41587-020-
0470-y.
29. Ng, E. S. *et al.* Differentiation of human embryonic stem cells to HOXA+ hemogenic
vasculature that resembles the aorta-gonad-mesonephros. *Nat Biotechnol* **34**, 1168–
1179 (2016).
30. Sturgeon, C. M., Ditadi, A., Awong, G., Kennedy, M. & Keller, G. Wnt signaling
controls the specification of definitive and primitive hematopoiesis from human
pluripotent stem cells. *Nat Biotechnol* **32**, 554–561 (2014).
31. Ditadi, A. *et al.* Human definitive haemogenic endothelium and arterial vascular
endothelium represent distinct lineages. *Nat Cell Biol* **17**, 580–591 (2015).
32. Ventura, T., Egan, E. J., Romanò, N. & Fidanza, A. In Vitro Model of Fetal Human
Vessel On-chip to Study Developmental Mechanobiology. *J Vis Exp* **2023**, (2023).
33. Fidanza, A. *et al.* Single cell analyses and machine learning define hematopoietic
progenitor and HSC-like cells derived from human PSCs. *Blood* **136**, 2893–2904
(2020).
34. Ventura, T., Egan, E. J., Romanò, N. & Fidanza, A. In Vitro Model of Fetal Human
Vessel On-chip to Study Developmental Mechanobiology. *J Vis Exp* **2023**, (2023).
35. Azzoni, E. *et al.* The onset of circulation triggers a metabolic switch required for
endothelial to hematopoietic transition. *Cell Rep* **37**, (2021).
36. Oburoglu, L. *et al.* Pyruvate metabolism guides definitive lineage specification during
hematopoietic emergence. *EMBO Rep* **23**, (2022).
37. Castaño, J. *et al.* GATA2 Promotes Hematopoietic Development and Represses
Cardiac Differentiation of Human Mesoderm. *Stem Cell Reports* **13**, 515–529 (2019).
38. de Pater, E. *et al.* Gata2 is required for HSC generation and survival. *Journal of*
Experimental Medicine **210**, 2843–2850 (2013).
39. Ling, K. W. *et al.* GATA-2 Plays Two Functionally Distinct Roles during the
Ontogeny of Hematopoietic Stem Cells. *J Exp Med* **200**, 871 (2004).
40. McGarvey, A. C. *et al.* A molecular roadmap of the AGM region reveals BMPER as a
novel regulator of HSC maturation. *J Exp Med* **214**, 3731–3751 (2017).
41. McGrath, K. E. *et al.* A transient definitive erythroid lineage with unique regulation of
the β-globin locus in the mammalian embryo. *Blood* **117**, 4600 (2011).

42.   Rejeski, K., Duque-Afonso, J. & Lübbert, M. AML1/ETO and its function as a regulator of gene transcription via epigenetic mechanisms. *Oncogene* **40**, 5665–5676 (2021).

43.   Huynh, H. D. *et al.* IGF binding protein 2 supports the survival and cycling of hematopoietic stem cells. *Blood* **118**, 3236 (2011).

44.   Crosse, E. I. *et al.* Multi-layered Spatial Transcriptomics Identify Secretory Factors Promoting Human Hematopoietic Stem Cell Development. *Cell Stem Cell* **27**, 822 (2020).

45.   Hadland, B. *et al.* Engineering a niche supporting hematopoietic stem cell development using integrated single-cell transcriptomics. *Nature Communications 2022 13:1* **13**, 1–17 (2022).

46.   Hadland, B. K. *et al.* Endothelium and NOTCH specify and amplify aorta-gonad-mesonephros-derived hematopoietic stem cells. *J Clin Invest* **125**, 2032–2045 (2015).

47.   Sandler, V. M. *et al.* Reprogramming human endothelial cells to haematopoietic cells requires vascular induction. *Nature* **511**, 312–318 (2014).

48.   Azzoni, E. *et al.* The onset of circulation triggers a metabolic switch required for endothelial to hematopoietic transition. *Cell Rep* **37**, (2021).

49.   Konermann, S. *et al.* Genome-scale transcriptional activation by an engineered CRISPR-Cas9 complex. *Nature* **517**, 583–588 (2015).

50.   Petazzi, P. *et al.* Robustness of Catalytically Dead Cas9 Activators in Human Pluripotent and Mesenchymal Stem Cells. *Mol Ther Nucleic Acids* **20**, 196–204 (2020).

51.   Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29 (2021).

52.   Cao, J. *et al.* The single-cell transcriptional landscape of mammalian organogenesis. *Nature 2019 566:7745* **566**, 496–502 (2019).

53.   Ge, S. X., Jung, D. & Yao, R. ShinyGO: a graphical gene-set enrichment tool for animals and plants. *Bioinformatics* **36**, 2628–2629 (2020).

874