1   *A novel human pluripotent stem cell-based gene activation system identifies IGFBP2*
2   *as a mediator in the production of hematopoietic progenitors in vitro.*
3

Paolo Petazzi[1], Telma Ventura[6], Francesca Paola Luongo[6], Heather McClafferty[7], Alisha

May[6], Helen Alice Taylor[6], Micheal J Shipston[7,8], Nicola Romanò[7,8], Lesley M. Forrester[6],

Pablo Menéndez[1-5], Antonella Fidanza[6,9*].

1. Josep Carreras Leukemia Research Institute, Barcelona, Spain.

2. Red Española de Terapias Avanzadas (TERAV)-Instituto de Salud Carlos III (ISCIII),

Madrid, Spain.

3. CIBER-ONC, ISCIII, Barcelona, Spain.

4. Institució Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, Spain.

5. Department of Biomedicine. School of Medicine, University of Barcelona, Barcelona,

Spain.

6. Centre for Regenerative Medicine, Institute for Regeneration and Repair, University of

Edinburgh, Edinburgh, UK

7. Centre for Discovery Brain Sciences, Edinburgh Medical School, Biomedical Sciences,

University of Edinburgh, Edinburgh, UK

8. Zhejiang University-University of Edinburgh Joint Institute, Zhejiang University School of

Medicine, Zhejiang University, Haining 310058, China

9. Edinburgh Medical School, Biomedical Sciences, University of Edinburgh, Edinburgh, UK

* Correspondence to afidanza@ed.ac.uk

Abstract

A major challenge in the stem cell biology field is the ability to produce fully functional cells

from induced pluripotent stem cells (iPSCs) that are a valuable resource for cell therapy,

drug screening and disease modelling. Here we developed a novel inducible CRISPR-

mediated activation strategy (iCRISPRa) to drive the expression of multiple endogenous

transcription factors important for *in vitro* cell fate and differentiation of iPSCs to

haematopoietic progenitor cells and used this to identify a key paracrine role for IGFBP2 in

the development of hematopoietic progenitors. We first identified nine candidate

transcription factors that we predicted to be involved in blood cell emergence during

development, then generated novel gRNAs directed to the transcriptional start site of these

transcription factors that could also be detected during scRNAseq. iCRISPRa activation of

these endogenous transcription factors resulted in a significant expansion of arterial-fated

endothelial cells expressing high levels of IGFBP2 and showed that IGFBP2 remodels the

metabolic activity during *in vitro* endothelial to hematopoietic transition. As well as providing

fundamental new insights into the mechanisms of haematopoietic cell fate and

differentiation, the broader applicability of iCRISPRa provides a valuable tool for studying

dynamic processes controlling developmental events as well as for recapitulating abnormal

phenotypes characterised by ectopic activation of specific endogenous gene expression in a

wide range of systems.

Introduction

The haematopoietic system develops early during gestation through so called "waves" of

hematopoietic progenitors that arise in different anatomical regions and result in the

production of various progenitor and stem cells [1–5]. The precise signalling pathways leading

to the production of hematopoietic stem and progenitor cells (HSPCs) during embryonic

development are yet to be completely defined, posing a limitation on how to recapitulate the

process *in vitro* from pluripotent stem cells.

During development, HSPCs are generated by a subset of endothelial cells, known as

hemogenic endothelium [6–9], via the endothelial to hematopoietic transition (EHT) [10]. During

the EHT, endothelial cells undergo profound transcriptional remodelling whereby the

expression of endothelial genes is gradually downregulated, and the transcription of the

hematopoietic program is initiated [11] while the cells round up and eventually detach to enter

the circulation [12,13].

To explore the molecular control on the development of the hematopoietic system and to

address the differences with the *in vitro* differentiation of induced pluripotent stem cells

(iPSCs), we compared our single cell transcriptomics analysis of *in vitro* derived hemogenic

endothelium and early progenitors [14] to that of the *in vivo* HSC-primed human hemogenic

endothelium [15]. We then developed a novel DOX-inducible CRISPR gene activation system

to assess the role of the genes that were expressed at a lower level within *in vitro*-derived

cells compared to their *in vivo* counterparts. We employed single-cell RNA sequencing to

track the presence of guide RNAs and we monitored the phenotypic effects of gene

activation. With this experimental pipeline, we identified and functionally validated a novel

role for IGFBP2, IGF Binding protein 2, in the development of *in vitro* progenitors and

showed that IGBP2 remodels the metabolic activity during *in vitro* EHT.

Results

Comparison of iPSC-derived endothelial cells with AGM AGM endothelial cells dataset

identifies 9 differentially expressed transcription factors.

We, and others, have shown that *in vitro* differentiation of human iPSCs is a powerful tool to

model intraembryonic hematopoiesis [14,16–18]. To understand the molecular basis underlying

the challenges associated with the *in vitro* production of blood stem and progenitor cells *in*

vitro from differentiating iPSCs, we compared our scRNAseq dataset from differentiating

iPSCs[14] to that of cells derived *in vivo* from the human aorta-gonad-mesonephros (AGM)

region [15]. We integrated the transcriptomic data of *in vitro* derived endothelial (IVD_Endo)

and hematopoietic cells (IVD_HPC) with that of arterial endothelial cells (aEC), arterial

hemogenic cells (aHEC) and venous endothelial cells (vEC) derived from human embryos

collected between Carnegie stage 12 and 14 (Figure 1A). To identify possible target genes

that could be manipulated *in vitro* to improve iPSCs differentiation, we first determined the

transcription factors marking the aHEC *in vivo* (Table 1). Then, we filtered the transcription

factor genes that were expressed at lower level in the IVD_Endo and IVD_HPC. This

strategy identified 9 target transcription factors *RUNX1T1*, *NR4A1*, *GATA2*, *SMAD7*,

ZNF124, *SOX6*, *ZNF33A*, *NFAT5* and *TFDP2* (Figure 1B), whose expression was detected

across other AGM datasets in the hemogenic endothelium[17] (Supplementary Figure 1).

Development of a DOX-inducible dCAS9-SAM activation system in human iPSCs

We previously developed an all-in-one Synergistic Activator Mediator system, UniSAM, that

mediates the transcriptional activation of endogenous gene expression[19]. To activate the

nine target genes identified in this study, we developed a novel doxycycline-inducible SAM

(iSAM) cassette targeted into the *AAVS1* locus of human iPSCs (Figure 2A). We

demonstrated the activation of *RUNX1C* expression in HeLa cells and that this was

correlated with the doxycycline (DOX) concentration in a linear manner (Figure S2A-B). To

verify the activation in human PSCs at single-cell resolution, we employed a RUNX1C-GFP

human embryonic stem cell (hESC) reporter cell line (Figure S2C-G). As predicted, the level

of expression of the mCherry tag, the fluorescent reporter within the iSAM cassette, was

proportional to the concentration of DOX (Figure S2D-E) and to the number of RUNX1C-

GFP+ cells (Figure S2 D-F). Furthermore, the RUNX1C expression level, measured by the

mean fluorescence intensity (MFI) of the RUNX1C-GFP reporter, also correlated with the

concentration of DOX (Figure S2G). We then tested the iSAM cassette in the iPSC line

(SFCi55) (Figure 2B-D). Only when iPSCs were transfected with iSAM and the gRNA for

RUNX1C, in the presence of DOX, was the expression of the *RUNX1C* gene and RUNX1

protein detected (Figure 2B-C).

We then targeted the iSAM cassette into the *AAVS1* locus using a Zinc Finger Nuclease

(ZFN) strategy [14,20,21]. iPSC clones that had specifically integrated the iSAM cassette into the

AAVS1 locus were validated by genomic PCR and sequencing (Figure S2A-B). The *AAVS1*

locus has been reported to be a "safe harbour" site that is resistant to epigenetic silencing
and indeed, we had previously demonstrated that transgenes inserted into the *AAVS1* locus
under the control of the constitutively active CAG promoter were efficiently expressed both in
undifferentiated and in differentiated iPSCs [20,22–24]. However, after the iSAM line had been
established and cultured under self-renewal conditions, we noted a dramatic reduction in the
number of mCherry+ cells in undifferentiated iSAM iPSCs upon DOX induction, that we
predicted to be due to transgene-silencing of the rTTA DOX-inducible cassette (Figure S3C).
To overcome this problem, we treated the iSAM iPSC line with an inhibitor of histone
deacetylases (HDACs), sodium butyrate (SB), reported to have no adverse effect on iPSC
maintenance [25,26] and we also confirmed that the treatment with SB had no effect on viability
and cell proliferation in our culture conditions (Figure S3D-E). A short 48-hour treatment
significantly increased the number of mCherry+ cells upon DOX induction, proportional to
the SB concentration (Figure S3F). We, therefore, maintained the iSAM iPSCs in the
presence of SB and this fully restored the inducibility of the transgene with virtually all cells
expressing mCherry in the presence of DOX (Figure S3G).
To test the effect of activating the 9 target genes on the transcriptomes of differentiating
iPSC cells, we engineered the gRNAs to allow their detection within the single cell RNA
sequencing pipeline [27]. We inserted a capture sequence prior to the termination signal to
avoid any alteration in the secondary structure of the loops thus preserving the binding of the
synergistic activators of the SAM system to the stem loops of the gRNAs. Of the two capture
sequences available[27], we decided to use the one that was predicted to result in fewer
secondary structure alterations and this new gRNA was named 2.1 (Figure 2E). We
compared the activation level achieved with the new 2.1 gRNA to that of the original 2.0
backbone using various gRNAs targeting *RUNX1C* (Figure 2F). These results convincingly
demonstrated that the addition of the capture sequence in the gRNA 2.1 does not alter the
level of endogenous gene activation that could be achieved (Figure 2G). Altogether, these
results show that the iSAM system is able to induce gene expression that translates into an
increased protein expression and thus provides a platform to steer phenotypical changes in
cell identity.
To activate our target genes, we designed 5-7 gRNAs in the 200bp upstream of the
transcriptional start sites of each of the 9 target genes. We subcloned a total of 49 gRNAs
(Table S1) into the gRNA 2.1 backbone and packaged them into lentiviral particles, (herein
referred to as the AGM library) as well as a non-targeting (NT) gRNA that was used as
control. The iSAM iPSC line was transduced with the targeting gRNAs or the control non-
targeting gRNA to generate the iSAM_AGM and iSAM_NT iPSCs line, respectively. After
puromycin selection, their integration in the genome was also confirmed by PCR and
sequencing (Figure S3H-I).

Single Cell RNA sequencing in combination with CRISPR activation identifies arterial
cell type expansion in association with higher hematopoietic progenitor potential

To assess the transcriptional changes in response to the activation of the target genes, we
differentiated the iSAM iPSCs, induced with DOX and subjected them to single-cell RNA
sequencing using the 10X pipeline. After 10 days of differentiation, in the presence or
absence of DOX, we FAC-sorted live CD34+ cells from iSAM_AGM and the iSAM_NT
iPSCs. Following data filtering, we selected cells in which the gRNAs expression was
detected and showed that our approach activated all the target genes, except for ZFN124. A
higher level of expression of *RUNX1T1*, *NR4A1*, *GATA2*, *SMAD7*, *SOX6*, *ZNF33A*, *NFAT5*,
TFDP2 was observed following DOX treatment of cells of iSAM_AGM cells compared to the
iSAM_NT cells (Figure 3A). To study the effect of gene activation on transcriptional and
cellular phenotype, we performed clustering analysis and detected a total of 7 clusters
(Figure 3B). A high level of *GJA4* and *DLL4* expression was used to annotate the arterial-like
cell cluster, while high levels of hemogenic markers, such as *RUNX1* and *CD44*, were used
to annotate the hemogenic clusters, EHT_1 and EHT_2 (Figure 3B, C-D). To assess the
effect of the activation on cell identity, we analysed the proportion of the cell clusters
between the libraries and detected a significant expansion of the arterial cluster in the DOX-
induced iSAM-AGM compared to the iSAM_NT cells (Figure 3E). To validate the effect of
the activation on the expansion of arterial cell population, we analysed their prevalence by
flow cytometry. Although DOX treatment resulted in an average 1.63 ± 0.13 fold increase of
$CD34^+DLL4^+$ cells in the control iSAM_NT sample (Figure S3A), the increase observed in
the iSAM-AGM cells was significantly greater, with an expansion of 3.33 ± 0.73 fold increase
of $CD34^+DLL4^+$ immunophenotypic arterial cells identified by flow cytometry (Figure 3F). To
assess the effect of this predicted arterial cell expansion on the emergence of functional
progenitors, we isolated $CD34^+$ cells using magnetic beads and cocultured 20,000 cells on
OP9 supportive stromal cells for 7 days, in the presence of hematopoietic differentiation
cytokines. After one week, the progenitor cells were assessed by colony-forming unit (CFU)
assays and scored 14 days later. We detected an increased number of CFU-E (colony-
forming unit erythroid) and CFU-GM (colony-forming unit granulocyte/macrophage) and a
reduction of CFU-M (colony-forming unit macrophage) in iSAM-AGM samples cultured in the
presence of DOX compared to the absence of DOX but no significant effect of DOX in
SAM_NT samples (Figure 3G). These data indicate that activation of target transcription
factors using our novel CRISPR strategy results in transcriptional remodelling and a steer in
cell identity that we detected as a functional difference in the hematopoietic progenitor
profile.

The addition of IGFBP2 to the *in vitro* differentiation leads to a higher number of functional hematopoietic progenitor cells.

To better understand the molecular mechanism behind the increased progenitor development, we compared the expression profile of the arterial cells between the different activation libraries, and we obtained a list of genes upregulated upon activation of the targets (Table 1). The most upregulated gene, *IGFBP2,* was expressed at significantly higher levels in the iSAM_AGM library in the presence of DOX compared to the others (Figure 4A, Table 1). We then compared the gRNA distribution in these arterial cells from the iSAM_AGM treated with DOX to that of arterial cells without activation. We observed a significant enrichment of the *RUNX1T1-specific* gRNAs (Table1), indicating that the increased *IGFBP2* expression could be downstream of *RUNX1T1* activation. IGF Binding Protein 2 is a member of the family of IGF binding proteins and is thought to be secreted from cells where it then binds IGF1, IGF2, and other extracellular matrix proteins, modulating their function. To test if the increased frequency of functional hematopoietic progenitors was due to IGFBP2 paracrine signalling, presumably derived from the arterial cells, we supplemented the media with IGFBP2 at 100ng/ml after the induction of endothelial cell differentiation. To explore the role of IGFBP2 we employed the parental iPSCs line, SFCi55 from which the iSAM line was derived. We isolated CD34+ cells at day 8 and co-cultured them on OP9 cells in presence of IGFBP2 for one week, then tested for their hematopoietic clonogenic potential using CFU assays (Figure 4B). Cells treated with IGFBP2 showed a significant increase in the total number of haematopoietic CFU colonies compared to cells in control cultures. To assess whether IGFBP2 also had a paracrine effect on the production of arterial cells themselves, we analyzed the proportion of DLL4+ cells, but no difference was detected in the presence of IGFBP2 (Figure 4C). This observation supports our hypothesis that the change in hematopoietic progenitor production following activation in the iSAM_AGM line is associated with differential gene expression within the arterial cells rather than expansion of that cell population. We then focused on the characterization of the cells derived from the CD34+ cells after coculture with the OP9 in the presence of IGFBP2. Our results showed a significant expansion of the CD34+ and CD43+ cell populations further supporting our hypothesis (Figure 4D). To address the potential role of *IGFBP2 in vivo*, we analysed single cell sequencing data from the human AGM at Carnegie stages 14 and 15. *IGFBP2* is highly expressed within the AGM niche by stromal and epithelial cells and, most importantly, highest in endothelial cells (Figure 4E). Together these data show that IGFBP2 addition results in increased hematopoietic blood production *in vitro* and that the endothelial compartment is the most likely source of IGFBP2 both *in vitro* and *in vivo*.

IGFBP2 remodels the metabolic profile of endothelial cells.

Following the observation that IGFBP2 supports haematopoietic progenitor differentiation *in vitro* from human iPSCs we performed a time course single-cell RNA experiment of SFCi55 iPSCs differentiated in its presence. We isolated CD34[+] cells and plated them in EHT culture on laminin[28], rather than on OP9 support, to assess exclusively the specific effects of IGFBP2- on differentiating iPSCs. FAC-sorted single/live cells from day 10 and 13 in the presence and absence of IGFBP2 were subjected to single-cell RNAseq. Our time course transcriptomic analyses showed that IGFBP2 induced a change in the transcriptome, specifically on day 13 (Figure 5A). A cluster of endothelial cells enriched for genes associated with the KEGG pathway of growth factor binding, was detected almost exclusively at day 13; these cells displayed a different transcriptional signature (as indicated by a shift in their position in the UMAP embedding) in the presence of IGFBP2..
Interestingly, this cluster showed expression of *GJA4* across the cells, indicating their broad arterial identity, with a specific reduction of *DLL4* expression level concomitantly to an upregulation of *RUNX1* in the region of the cluster induced by IGFBP2 (Figure 5B). This increase in hemogenic identity within the endothelial cell compartment is consistent with the observation of increased functional progenitor production. We compared the transcriptome of cells at day 13 in the presence and absence of IGFBP2 and, using KEGG enrichment analysis, we observed that the genes that were upregulated by IGFBP2 were highly enriched in the oxidative phosphorylation term (Figure 5C-D). Since the metabolic switch between glycolytic to mitochondrial metabolism has been previously reported in definitive hematopoiesis[29,30], we tested whether the addition of IGFBP2 could result in a different ATP production profile. We analysed the ATP production at day 13, when RUNX1 expression was induced by IGFBP2 addition, by using specific inhibitors of the complex I and II and complex V to quantify the intracellular ATP and mitochondrial synthesis, respectively. We detected a reduction in glycolytic-derived ATP (Figure 5E-F), which translates to a higher contribution of mitochondria metabolism for IGFBP2-treated cells (Figure 5G). We then looked at the expression levels of genes encoding the enzymes of the glycolytic pathway and its checkpoints, glucose and lactate transporters, monocarboxylate transporters and the enzymes associated with hexokinase and phosphofructose reactions. We observed a general downregulation of the glycolytic enzymes and its checkpoints, with the exception of *PFKM* and *SLC16A1*, (Figure S4), which were expressed at lower levels in the IGFBP2-treated cells at day 13 compared to the control on the same day. These differences were not observed on day 10 cells, which is consistent with our previous observations that the IGFBP2-induced remodelling of the transcriptome happens exclusively on day 13. These

results indicate that the presence of IGFBP2 induces the upregulation of *RUNX1* and

OxPhos genes and the metabolic remodelling of endothelial cells.

Discussion

The complexity and dynamism of developmental hematopoiesis *in vivo* have imposed

challenges in accurately reproducing the process *in vitro*. We hypothesized that this could, in

part, be due to the inability to recapitulate the appropriate transcriptional programme in

cultured cells. Here, we developed a novel CRISPR-activation system to induce the

expression of genes that are expressed at low levels in cells that are generated *in vitro*, to

explore the downstream consequences of their activation and to assess the effects on

emerging hematopoietic progenitor cells.

When we compared endothelial cells derived *in vitro* from hiPSCs to those in the AGM

region of the human embryo, at the time point when early hematopoietic commitment takes

place, we identified nine transcription factors that were expressed at a lower level in the

cultured cells. Some of these had been associated with blood cell development, including

GATA2[31–33]*, SMAD7* [34]*, NR4A1*[34]*, SOX6*[35] and *RUNX1T1*[15,17]. Other genes such as,

ZNF124, ZNF33A, NFAT5 and TFDP2 had not been previously associated with

hematopoiesis and could provide early evidence of their possible role in hematopoiesis that

will require further studies. We were particularly interested in *RUNX1T1* (also known as

ETO) which is associated with the leukemic fusion protein, AML1/ETO resulting from the

t(8;21) chromosomal translocation[36]. Furthermore, *RUNX1T1* expression has been recently

detected in transcriptomic analyses of the human AGM region[15,17], but its precise role during

the ontogeny of the blood system has not been elucidated. The addition of a capture

sequence to the gRNA backbone enabled their detection coincidentally with the single-cell

transcriptome, and this allowed us to demonstrate that RUNX1T1 gRNAs were significantly

enriched in cells within an expanded arterial cluster and their presence was associated with

a high level of expression of *IGFBP2*. *IGFBP2* KO mice show increased expression of cell-

cycle inhibitors and HSC apoptosis, implicating IGFBP2 as a modulator of HSCs cell cycle

and survival [37]. More recently *IGFBP2* was reported as being highly expressed in the human

AGM region at CS14 when HSCs are emerging [17], supporting a possible role during

developmental hematopoiesis but the precise molecular process was unclear. In this study,

we show that the addition of IGFBP2 recombinant protein in our *in vitro* model of EHT results

in the emergence of an increased number of functional hematopoietic progenitors and we

provide evidence to support the hypothesis that RUNX1T1 could regulated its expression.

Because RUNX1T1 lacks a DNA binding domain, its direct involvement in the regulation of

IGFBP2 expression must require association with other cofactors that are yet to be
identified.
Our data, together with *in vivo* data[17], show that IGFBP2 is expressed predominantly by
endothelial cells. The supportive role of the endothelial niche in the development of HSCs
has been studied *in vivo* [38–40] and exploited *in vitro* to support HSCs emergence [39,41].
We describe here a novel role for IGFBP2 in the remodelling of the metabolism of iPSC-
derived endothelial cells and that this is associated with the induction of RUNX1, a hallmark
of hemogenic fate. The addition of IGFBP2 in the culture induces upregulation of the genes
of the oxidative phosphorylation pathway in association with an increased relative
mitochondrial contribution to cellular ATP production. This is due to reduced glycolysis in
association with the downregulation of genes coding for glycolytic enzymes and their
checkpoints. The switch between glycolytic to mitochondrial metabolism has been shown to
be essential for the EHT in definitive hematopoiesis[29,30]. This switch is induced *in vivo* by
mechanical cues downstream of the establishment of circulation, and it is required for
functional HSCs development[29], while *in vitro*, the switch is driven by pyruvate mitochondrial
catabolism, leading to definitive EHT as opposed to primitive[30]. Further *in vivo* studies are
required to characterize the role of IGFBP2 during AGM hematopoiesis to overcome the
limitation of using an *in vitro* model such as iPSCs differentiation.
In conclusion, we detected transcriptional differences between *in vivo* and *in vitro*
developmental hematopoiesis and developed a novel inducible gene activation system to
identify novel molecular players during the endothelial-to-hematopoietic transition. Our
multidisciplinary approach identified *RUNX1T1* and IGFBP2 as novel signalling molecules
that support human blood progenitor development *in vitro*, the latter inducing a metabolic
switch from cytoplasmic glycolysis to mitochondrial respiration. This study demonstrates that
the combination of CRISPR-mediated activation of target genes with single-cell
transcriptomic analysis in differentiating hPSCs can be a powerful approach to alter cell fate,
providing a tool for gene function studies during human development. The fine epigenetic
manipulation of the transcription can be readily applied to any cell lineage simply by adding
specific gRNAs and it will be instrumental in exploring other developmental processes that
can be, at least partially, be mimicked *in vitro* with human iPSCs.
327
328 Using this approach, we have identified the supportive role of paracrine IGFBP2,
329 predominantly produced by endothelial cells, which induces transcriptional and metabolic
330 remodelling in association with the induction of *RUNX1* expression.
331
332

333
334 **Figures**

Figure 1 - Comparison of *in vitro* iPSC-derived and *in vivo* AGM-derived endothelial

cells identifies 9 differentially expressed transcription factors.

A - Integrative analysis of single cell transcriptome of *in vitro* derived endothelial (IVD_Endo)

and hematopoietic cells (IVD_HPCs) with *in vivo* developed endothelial cells (venous, vEC;

arterial, aEC; arterial hemogenic, HECs) from human embryos (CS12-CS14) visualised on

UMAP dimensions. **B** - Target genes expression level showing higher expression in arterial

hemogenic endothelium *in vivo* compared to *in vitro* derived cells.

Figure 2 - The inducible iSAM cassette successfully mediates activation of endogenous gene expression upon DOX induction.

A - Schematic of the iSAM cassette containing the TET-on system under the control of EF1α and dCAS9-P2A-MS2-p65-HSF1-T2A-mCherry under the rTTA responsive elements, separated by genetic silencer and flanked by AAVS1 specific homology arms. **B** - *RUNX1C* gene expression activation after transient transfection of the iSAM plasmid and gRNAs in presence or absence of DOX in human iPSC line (n=3 from independent transfections). **C** – RUNX1 protein expression upon iSAM activation after transient transfection of the iSAM plasmid and gRNAs with DOX in human iPSC line detected by immunostaining. **D** Expression of the iSAM cassette reported by mCherry tag during the differentiation protocol, the representative images (bright field – BF, and fluorescence) show embryoid bodies at day 3 of differentiation. **E** - Schematic of the gRNA 2.1 containing the capture sequence for detection during the scRNAseq pipeline. **F** - *RUNX1C* gene activation level obtained using either the gRNA 2.0 or 2.1 backbone (n=3 from independent transfections of the 4 different gRNAs). **G** - Statistical analysis of the gRNAs activation level showing no significant variation following addition of the capture sequence (n=3 for each of the 4 different gRNAs).

Figure 3 - Single Cell RNA sequencing in combination with CRISPR activation

identifies arterial cell type and functional hematopoietic expansion in association with

activation of the 9 target genes.

A - Gene expression profile of target genes following target genes' activation, heatmap

shows the expression level of the target genes in the iSAM_NT and iSAM_AGM treated with

DOX following normalisation on the -DOX control. **B** - Dimension reduction and clustering

analysis of the scRNAseq data following activation, filtered on cells where the gRNA

expression was detected. **C** - Arterial (*GJA4*, *DLL4*), venous (*NRP2*, *APLNR*) and

hemogenic marker (*CD44*, *RUNX1*) expression distribution in the clusters indicated by the

colour. **D** - Expression distribution visualised on the UMAP plot showing the location of

arterial cells marked by *DLL4*, and hemogenic endothelium marked by *CD44* and *RUNX1*. **E**

- Contribution of the different libraries to the clusters showing that arterial cell cluster is

overrepresented in the iSAM_AGM treated with DOX, compared to the other libraries. **F** -

Expansion of the arterial population assessed by the membrane marker expression of

DLL4+ following targets' activation, quantified by flow cytometry at day 8 of differentiation

(Data are normalised on the iSAM_NT + DOX sample, n=5 independent differentiations, * p

397 = 0.0417 paired t-test). **G** - Colony forming potential of the suspension progenitor cells

398 derived from the two lines treated with or without DOX following OP9 coculture activation,

399 data show the colony obtained for 104 CD34+ input equivalent (n=3 from independent

400 differentiations * p<0.05, Tukey's two-way ANOVA).

Figure 4 - IGFBP2 addition to the *in vitro* differentiation leads to a higher number of functional hematopoietic progenitor cells. A - Violin plot of *IGFBP2* expression profile in the arterial cells obtained from the different conditions, in the presence or absence of gRNAs and DOX. **B** -Number of hematopoietic colonies obtained after coculture on OP9 in presence or absence of IGFBP2 (n=3-4 from independent differentiations, ** p=0.0080, Sidak's Two way ANOVA). **C** – Percentage of DLL4[+] arterial cells differentiation within the CD34[+] compartment analysed by flow cytometry in day 8 EBs (n=4 from independent differentiations, Two-way Anova, ns =p>0.99). **D** – Expansion of hematopoietic progenitors analysed using markers' expression on suspension progenitors derived after coculture of CD34+ cells onto OP9 support (data are expressed as fold over the CTR in the absence of IGFBP2 (n=4 from independent differentiations, * p<0.02, Sidak's Two way ANOVA. **E** - Single-cell transcriptomic analysis of developing AGM collected from human embryos at Carnegie Stages14 and 15 enriched for CD31[+] and CD34[+] showing the *IGFBP2* expression profile *in vivo* in the AGM.

Figure 5 - IGFBP2 alters cell metabolism by inducing a reduction in glycolytic ATP

production.

A – Clustering analysis of the single cell transcriptomic time course analysis of differentiating

cells at day 10 and day 13 in the absence (CTR) or presence of IGFBP2. Arrows indicate the

difference in the clustering due to the addition of IGFBP2 compared to control. **B** –

Expression profile of arterial markers, *GJA4* and *DLL4*, and hemogenic marker *RUNX1* (top

- the dashed line shows the location of the shift in gene expression of cells treated with

IGFBP2) and their expression profile in the endothelial cells cluster marked by Growth factor

binding in absence (CTR) and in presence of IGFBP2. **C –** KEGG enrichment analysis of the

genes upregulated at day 13 upon IGFBP2 treatment. The arrow shows the ranking of the

Oxidative Phosphorylation pathway. **D** - Dot Plot showing the expression profile of the

genes coding for the enzyme of the Oxidative Phosphorylation pathway. **E** -Oxygen

Consumption Rate (OCR) and Extracellular Acidification Rate (ECAR) profile in cells at day

13 of differentiation reporting mitochondrial respiration and glycolysis, respectively. **F** – ATP

production rate divided by that deriving from glycolysis and from mitochondrial respiration, in

cells treated with IGFBP2 and controls at day 13. **G** – Ratio of the ATP production between

glycolysis and mitochondrial respiration in cells treated with IGFBP2 and controls at day 13.

437
Methods

Resource availability

R code is available at https://github.com/afidanza/CRISPRa. Raw and processed data have

been deposited to ArrayExpress (E-MTAB-12748 – currently on hold). The AAVS1-iSAM are

available on Addgene (Addgene #211495) and the gRNA 2.1 plasmids (Addgene #211496)

(both currently on hold). Further information and requests for resources and reagents should

be directed to and will be fulfilled by the corresponding author.

Pluripotent Stem Cells maintenance

hPSCs were maintained *in vitro* in StemPro hESC SFM (Gibco) with bFGF (R&D) at 20

ng/ml. Wells were coated with Vitronectin (ThermoFisher Scientific) at least 1 hour before

plating and cells were passaged using the StemPro EZPassage tool (ThermoFisher

Scientific). Media change was performed every day and cells passaged every 3–4 days at a

ratio of 1:4.

Transfection

iPSCs SFCi55 and hESCs RUNX1-GFP were plated at 3×10^5 cells per a well of a 6 well

plate and reverse transfected with 2 µg of DNA using the Xfect Transfection reagent

(Clontech) and analyzed 2 days later.

HeLa cells were cultured in Dulbecco's Modified Eagle Medium/Nutrient Mixture F-12

(DMEM/F12) with Glutmax and 5% FCS (Gibco) and passaged every few days, at a ratio of

1:6. HEL were cultured in Iscove's Modified Dulbecco's Medium (IMDM) with 10% FCS

(Gibco) and passaged every few days, at a ratio of 1:4. 2×10^5 cells were plated, transfected

at 6–8 hours with 0.75 µg of DNA using Xfect Transfection reagent (Clontech) and then

analysed 2 days after.

Immunocytochemistry

Cells were fixed in 4% PFA in PBS at room temperature for 10', permeabilized in PBS-T

(0.4% Triton-X100) for 20' and blocked in PBS-T with 1% BSA and 3% goat serum for

1 hour. Primary antibodies were incubated in blocking solution over night at 4 °C (RUNX1

1:200 - ab92336, Abcam). Cells were then washed in PBS-T and incubated with secondary

antibodies for 1 hour at room temperature (donkey α-rabbit 1:200 - A-11008 - Thermo

Scientific). Cells were washed in PBS-T and counterstained with DAPI. Images were

generated using the Zeiss Observer microscope.

Gene expression analysis

Total RNA was purified using the RNAeasy Mini Kit (Qiagen) and cDNA synthesized from 500 ng of total RNA using the High Capacity cDNA synthesis Kit (Applied Biosystem). 2 ng of cDNA were amplified per reaction and each reaction was performed in triplicate using the LightCycler 384 (Roche) with SYBR Green Master Mix II (Roche). A melting curve was performed and analyzed for each gene to ensure the specificity of the amplification. *β-Actin* was used as reference genes to normalize the data [19].

Pluripotent Stem Cells differentiation to hematopoietic progenitors

hPSCs were differentiated in a xeno-free composition of SFD medium [14], BSA was substituted with human serum albumin, HSA (Irvine-Scientific). Day 0 differentiation medium, containing 10 ng/ml BMP4 was added to the colonies prior cutting. Cut colonies were transferred to a Cell Repellent 6 wells Plates (Greniner) to form embryoid bodies and cultured for two days. At day 2 media was changed and supplemented with 3 μM CHIR (StemMacs). At day 3, EBs were transferred into fresh media supplemented with 5 ng/ml bFGF and 15 ng/ml VEGF. At day 6 media was changed for final haematopoietic induction in SFD medium supplemented with 5 ng/ml bFGF, 15 ng/ml VEGF, 30 ng/ml IL3, 10 ng/ml IL6, 5 ng/ml IL11, 50 ng/ml SCF, 2 U/ml EPO, 30 ng/ml TPO, 10 ng/ml FLT3L and 25 ng/ml IGF1. From day 6 onward, cytokines were replaced every two days.

CD34 isolation

CD34+ cells were isolated using CD34 Magnetic Microbeads from Miltenyi Biotec, according to their manufacturing protocol. Briefly, Embryoid bodies were dissociated using Accutase (Life Techonologies) at 37°C for 30'. Cells were centrifuged and resuspended in 150 μl of PBS + 0.5% BSA + 2mM EDTA with 50 μl Fcr blocker and 50 μl of magnetic anti-CD34 at 4°C for 30'. Cells were washed using the same buffer and transferred to pre-equilibrated columns, washed three times and eluted. After centrifugation, cells were resuspend in SFD media, counted and plated for OP9 coculture.

OP9 coculture and colony assay

OP9 cells were maintained in α-MEM supplemented with 20% serum (Gibco) and sodium bicarbonate (Gibco) and passaged with Trypsin every 3-4 days. The day before the co-culture, 45.000 OP9 cells were plated for each 12 well plates' well in SFD media. The day of the co-culture the 20.000 iSAM cells or 25.000 SFCi55 or H9 were plated in each well and culture in SFD media supplemented with 5 ng/ml bFGF, 15 ng/ml VEGF, 30 ng/ml IL3, 10 ng/ml IL6, 5 ng/ml IL11, 50 ng/ml SCF, 2 U/ml EPO, 30 ng/ml TPO, 10 ng/ml FLT3L and 25

ng/ml IGF1 and 100ng/ml IGFBP2. Cytokines were replaced twice during one week of

coculture. At the end of the coculture, cells were collected by Trypsin and half of the well

equivalent was plated in 2 ml of methylcellulose medium (Human enriched H4435, Stemcell

Technologies). Cells were incubated in the assay for 14 days and then scored.

Laminin EHT culture

Laminin EHT culture was performed as previously described[28]. Briefly, 24 well plates were

coated for at least 2 hours with recombinant human Laminin-521 (Thermo-Fisher). Following

CD34+ isolation at day 8, 400.000 CD34+ cells were seeded in each well of a pre-coated

24-well plate in SFD media supplemented with 5 ng/ml bFGF, 15 ng/ml VEGF, 30 ng/ml IL3,

10 ng/ml IL6, 5 ng/ml IL11, 50 ng/ml SCF, 2 U/ml EPO, 30 ng/ml TPO, 10 ng/ml FLT3L and

25 ng/ml IGF1 and 100ng/ml IGFBP2.

Flow cytometry staining and cell sorting

Embryoid bodies were dissociated using Accutase (Life Techonologies) at 37°C for 30'.

Cells were centrifuged and resuspended in PBS + 0.5% BSA + 2mM EDTA, counted and

stained at 10^5 cells for a single tube. Cells were stained with antibodies for 30' at room

temperature gently shaking. Flow cytometry data were collected using DIVA software (BD).

For the sorting experiments, cells were stained at 10^7cells/ml in presence of the specific

antibodies. Sorting was performed using FACSAria Fusion (BD) and cells were collected in

PBS + 1% BSA. Data were analysed using FlowJo version 10.4.2.

Flow cytometry antibodies

For flow cytometry 10^5 cells per test were stained in 50 μl of staining solution with the

following antibodies: CD34 Percp-Efluor710 (4H11 eBioscience, 1:100), CD34 Pe (4H11

eBioscience, 1:200), CD43 APC (eBio84-3C1, 1:100), CD45 FITC (2D1 ebioscience, 1:100),

DLL4 Pe (MHD4-46 Biolegend, 1:200)

iSAM plasmid generation

The iSAM plasmid was obtained by Gibson assembly of four fragments. The first fragment,

the backbone, was a DOX-inducible AAVS1 targeted plasmid expressing an E6-E7-IRES-

ZsGreen which was excised by BstBI and NdeI. The second fragment, one of the adapters,

was derived from the UniSAM plasmid that we previously generated (Addgene #99866) by

PCR with the following primer sets

FW_aggggacccggttcgagaaggggctcttcatcactagggccgctagctctagagagcgtcgaatt,

RV_ttcgggtcccaattgccgtcgtgctggcggctcttccacctttctcttcttcttggggctcatggtggcc. The UniSAM

cassette was obtained also from the UniSAM plasmid via digestion with BsrGI and BsiWI.

Finally, the last fragment consisting of another adapter for the Gibson was custom

synthetised and contained overlapping sequences flanking a chicken b-globin insulator that

we inserted to prevent cross-activation of the EF1α-promoter and the TRE-GV promoter

driving the iSAM. Correct assembly was verified by Sanger sequencing.

iSAM cell lines derivation

The iSAM plasmid was used together with ZNFs specific for the AAVS1 locus to mediate

specific integration in SFCi55 human iPSCs line [14,20]. Briefly, 10 μg of AAVS1-iSAM with 2.5

μg of each ZNFs, left and right, using Xfect (Takara) according to the manufacturer protocol.

Cells were selected using Neomycin. Single clones were picked, amplified, and initially

screened by mCherry expression upon DOX addition. Clones that expressed the fluorescent

tag were screened for specific integration using PCR followed by Sanger sequencing for the

correctly integrated clones. 100 ng of genomic DNA was amplified using the EmeraldAmp®

MAX HS Takara and specific primer sets (Table 1). For the specific AAVS1 integration site,

Sigma_AAVS1 - CGG AAC TCT GCC CTC TAA CG and NeoR -GAT ATT GCT GAA GAG

CTT GGC GG were used with the PCR conditions of 95 °C for 7 min, 32 cycles of 95 °C for

15 s, 57 °C for 30 s, and 72 °C for 1 min, with the final elongation step at 72 °C for 7 min. For

the Wild type locus screening, Sigma_AAVS1 - CGG AAC TCT GCC CTC TAA CG and

AAVS1_EXT3_RV – ACA CCC AGA CCT GAC CCA AA were used with the PCR cycling

conditions of 95 °C for 7 min, 30 cycles of 95 °C for 15 s, 57 °C for 30 s, and 72 °C for 2 min,

with the final elongation step at 72 °C for 6 min.

Capture sequencing addition to the gRNA backbone

The Capture sequence 2 was added to the gRNA_Puro_Backbone (Addgene #73797) by

PCR. Briefly, the capture sequence was added before the termination signal of the gRNA

followed by a BamHI site using the following PCR primers:

gRNA_FW gagggcctatttcccatgattcct,

gRNA_Cap_RV aaaaaaggatccaaaaaaCCTTAGCCGCTAATAGGTGAGCgcaccgactcggtgcc.

The gRNA backobone was replaced from the original plasmid via NdeI and BamHI digestion,

followed by ligation of the PCR produced following the same digestion. Correct integration of

the insert was verified by Sanger sequencing.

Target genes identification

Candidate genes were identified by comparing *in vivo* hemogenic endothelium [15]

(GSE135202) and *in vitro* iPSCs-derived endothelial cells that we previously generated [14]

(E-MTAB-9295). Briefly, the two datasets were merged and normalised using the R

package Seurat. Specific markers for hemogenic endothelium were identified and

transcription factors were sorted based on their GO annotation. Within those genes we

filtered those detected in more than 50% of the *in vivo* hemogenic endothelium and

expressed in less than 25% of the *in vitro* derived endothelial cells. This pipeline identified 9

target genes *RUNX1T1, NR4A1, GATA2, SMAD7, ZNF124, SOX6, ZNF33A, NFAT5, TFDP.*

AGM gRNAs library preparation

sgRNA design was performed by selecting the top candidates for on-target and off-target

score. Between 5 and 7 guides per gene were designed for *RUNX1T1, NR4A1, GATA2,*

SMAD7, ZNF124, SOX6, ZNF33A, NFAT5, TFDP2 using the CRISPRpick tool from the

Broad Institute (https://portals.broadinstitute.org/gppx/crispick/public) (Table S1). All the

guide variants were Golden Gate cloned with the gRNA 2.1 backbone according to the

established protocol [42]. The 49 plasmids were pooled together in an equimolar ratio and the

library prep was subsequently used to produce lentiviral particles with a second-generation

production system. Briefly, the psPAX2 packaging plasmid, pMD2.G envelope, and the AGM

vector library were co-transfected using polyethyleneimine (PEI) (Polysciences, Warrington,

PA, USA) as previously detailed [43], Lentiviral particles-containing supernatants were

harvested 48–72 h post-transfection, concentrated by ultracentrifugation and titered in

hiPSCs cells.

iSAM_AGM and iSAM_NT cell line derivation

The selected iSAM clone (3.13 internal coding) was infected with viral particles containing

either the AGM library or the non targeting gRNA (NT) at a MOI of 10. The iSAM cells were

plated the afternoon before at $7x10^6$ cells into a T125 in the presence of 10 μM Rock

Inhibitor (Merk) which was maintained until the day following the infection. Cells were

infected in presence of 8μg/ml of Polybrene (Merk). Puromycin selection was initiated 36

hours post-infection and maintained during their culture until the beginning of the

differentiation. Both lines were tested for integration of the gRNAs. Briefly, 100 ng of isolated

gDNA was amplified using the PrimeSTAR MAX PCR mix (Takara) using the primers

gRNA_screening_FW and gRNA_screening_RV (Table1). Purified amplicons were

subjected to Sanger sequencing.

Single Cell RNA sequencing

For the iSAM_AGM and iSAM_NT single cell RNA sequencing experiment, embryoid bodies

obtained from day 10 of differentiation were dissociated using Accutase (Life Technologies')

at 37°C for 30'. For the IGFBP2 experiment, day 10 and day 13 cells from the Laminin EHT

culture were detached from the adherent layer using Accutase (Life Technologies') at 37°C

for 5'. From both experiments, cells were centrifuged and resuspended in CD34-Pe staining

solution at a density of 10^7/ml. CD34+/live/single cells were FAC-sorted in PBS + 0.1 %

BSA. Cell viability was also confirmed by Trypan blue stain for an accurate count. Around

15000 cells per sample were loaded into the 10X Chromium Controller, and single cell

libraries were obtained using the Chromium single cell 3' Reagent Kits v3 (10XGenomics)

according to manufacturer protocol. The four libraries were indexed using SI PCR primers

with different i7 indexes to allow for demultiplexing of the sequencing data. RNA

concentration was obtained using Quibit RNA HS (Thermo-Fisher). Quality of the obtained

libraries were verified using LabChip GX (PerkinElmer). Libraries were sequenced using

NextSeq 2000 technology (Illumina) at 50.000 reads/cell. Data were aligned to GRCh38

using the Cell Ranger dedicated pipeline (10XGenomics). Data filtering, dimension

reduction, clustering analysis, differentially expressed genes and cell cycle analysis were

obtained using Seurat R package (version 4.1.0)[44]. Pseudotemporal ordering was performed

using Monocle 3 R package [45]. KEGG pathways was performed using ShinyGo[46]. The code

is available on Github at https://github.com/afidanza/CRISPRa, the raw data have been

submitted to Array Express (E-MTAB-12748), and the browsable processed data will be

made available at the time of publication on our website containing previous sequencing

data at https://lab.antonellafidanza.com.

ATP production analysis

At day 8 of differentiation, 20.000 CD34+ were plated for each precoated well of the

Seahorse XFp mini Cell Culture Plates (Agilent) precoated with rhLaminin-521 (Thermo-

Fisher). At day 13, the cells' ATP production rate was analysed with the Seahorse XF Real-

Time ATP Rate Assay kit (Agilent) according to manufacturer protocol. Briefly, the media

was changed prior to the assay for the XF DMEM medium, pH 7.4, supplemented with 10

mM Seahorse XF glucose, 1mM Seahorse XF pyruvate and 2mM Seahorse XF Glutamine

(Agilent) and incubated for 45-60 minutes in a non-CO_2 incubator. Oligomycin and

Rotenone/AA solutions were prepared and added in the cartridge and finally loaded together

with the cells in the Seahorse XF Mini Analyzer using the dedicated software. Data were

collected at the end of the run, and the values of ATP production were calculated according

to the ATP Production Rate Calculation provided by Agilent.

Author contribution

AF designed the study, performed experiments and bioinformatic analysis, wrote the paper

and led the research. PP, TV, FPL, HM performed experiments. AM, HT, MS provided

support to the experiments. NR performed bioinformatic analysis. LF and PM helped the

design of the study and the research. All authors provided essential feedback on the

experiments and the manuscript.

Acknowledgment

AF and LF acknowledge financial support from the Biotechnology and Biological Sciences

Research Council; Grant S002219/1. AF was supported by a European Hematology

Association Advanced Research Grant (EHA RAG 2021), and by the American Society for

Hematology (Research Global Award). TV and AM were supported by PhD studentships

from the Medical Research Council (Precision Medicine) and the College of Medicine and

Veterinary Medicine, respectively. FPL was supported by an Erasmus+ Traineeship Program

2016/2017. PM acknowledges financial support from a PERIS program from the Catalan

Government and a Retos collaboration project from the MINECO (RTC-2018-4603-1)

Bibliography

1. Medvinsky, A. & Dzierzak, E. Definitive hematopoiesis is autonomously initiated by the AGM region. *Cell* **86**, 897–906 (1996).

2. Palis, J., Robertson, S., Kennedy, M., Wall, C. & Keller, G. Development of erythroid and myeloid progenitors in the yolk sac and embryo proper of the mouse. *Development* **126**, 5073–84 (1999).

3. Patel, S. H. *et al.* Lifelong multilineage contribution by embryonic-born blood progenitors. *Nat. 2022 6067915* **606**, 747–753 (2022).

4. Böiers, C. *et al.* Lymphomyeloid Contribution of an Immune-Restricted Progenitor Emerging Prior to Definitive Hematopoietic Stem Cells. *Cell Stem Cell* **13**, 535–548 (2013).

5. Hoeffel, G. *et al.* C-Myb(+) erythro-myeloid progenitor-derived fetal monocytes give rise to adult tissue-resident macrophages. *Immunity* **42**, 665–78 (2015).

6. Jaffredo, T., Gautier, R., Eichmann, A. & Dieterlen-Lièvre, F. Intraaortic hemopoietic cells are derived from endothelial cells during ontogeny. *Development* **125**, 4575–83 (1998).

7. Zovein, A. C. *et al.* Fate Tracing Reveals the Endothelial Origin of Hematopoietic Stem Cells. *Cell Stem Cell* **3**, 625–636 (2008).

8. Bertrand, J. Y. *et al.* Haematopoietic stem cells derive directly from aortic endothelium during development. *Nature* **464**, 108–111 (2010).

9. Boisset, J. C. *et al.* In vivo imaging of haematopoietic cells emerging from the mouse aortic endothelium. *Nature* **464**, 116–120 (2010).

10. Ottersbach, K. Endothelial-to-haematopoietic transition: an update on the process of making blood. (2019). doi:10.1042/BST20180320

11. Swiers, G. *et al.* Early dynamic fate changes in haemogenic endothelium characterized at the single-cell level. *Nat. Commun. 2013 41* **4**, 1–10 (2013).

12. Kissa, K. & Herbomel, P. Blood stem cells emerge from aortic endothelium by a novel type of cell transition. *Nature* **464**, 112–115 (2010).

13. Eilken, H. M., Nishikawa, S. I. & Schroeder, T. Continuous single-cell imaging of blood generation from haemogenic endothelium. *Nature* **457**, 896–900 (2009).

14. Fidanza, A. *et al.* Single cell analyses and machine learning define hematopoietic progenitor and HSC-like cells derived from human PSCs. *Blood* **136**, 2893–2904 (2020).

15. Zeng, Y. *et al.* Tracing the first hematopoietic stem cell generation in human embryo by single-cell RNA sequencing. *Cell Res.* **29**, 881–894 (2019).

16. Ng, E. S. *et al.* Differentiation of human embryonic stem cells to HOXA+ hemogenic

vasculature that resembles the aorta-gonad-mesonephros. *Nat. Biotechnol.* **34**, 1168–

1179 (2016).

17.   Calvanese, V. *et al.* Mapping human haematopoietic stem cells from haemogenic

endothelium to birth. *Nat. 2022 6047906* **604**, 534–540 (2022).

18.   Sturgeon, C. M., Ditadi, A., Awong, G., Kennedy, M. & Keller, G. Wnt signaling

controls the specification of definitive and primitive hematopoiesis from human

pluripotent stem cells. *Nat. Biotechnol.* **32**, 554–561 (2014).

19.   Fidanza, A. *et al.* An all-in-one UniSam vector system for efficient gene activation. *Sci.*

Rep. **7**, 6394 (2017).

20.   Yang, C.-T. *et al.* Activation of KLF1 Enhances the Differentiation and Maturation of

Red Blood Cells from Human Pluripotent Stem Cells. *Stem Cells* **35**, 886–897 (2017).

21.   Lopez-Yrigoyen, M. *et al.* A human iPSC line capable of differentiating into functional

macrophages expressing ZsGreen: A tool for the study and in vivo tracking of

therapeutic cells. *Philos. Trans. R. Soc. B Biol. Sci.* **373**, (2018).

22.   Lopez-Yrigoyen, M. *et al.* Genetic programming of macrophages generates an in vitro

model for the human erythroid island niche. *Nat. Commun.* **10**, 881 (2019).

23.   Lopez-Yrigoyen, M. *et al.* A human iPSC line capable of differentiating into functional

macrophages expressing ZsGreen: A tool for the study and in vivo tracking of

therapeutic cells. *Philos. Trans. R. Soc. B Biol. Sci.* **373**, (2018).

24.   May, A. *et al.* Modelling the erythroblastic island niche of dyserythropoietic anaemia

type IV patients using induced pluripotent stem cells. *Front. cell Dev. Biol.* **11**, (2023).

25.   Kang, S. J., Park, Y. Il, So, B. & Kang, H. G. Sodium butyrate efficiently converts fully

reprogrammed induced pluripotent stem cells from mouse partially reprogrammed

cells. *Cell. Reprogram.* **16**, 345–354 (2014).

26.   Zhang, Z., Xiang, D. & Wu, W. S. Sodium Butyrate Facilitates Reprogramming by

Derepressing OCT4 Transactivity at the Promoter of Embryonic Stem Cell–Specific

miR-302/367 Cluster. *Cell. Reprogram.* **16**, 130 (2014).

27.   Replogle, J. M. *et al.* Combinatorial single-cell CRISPR screens by direct guide RNA

capture and targeted sequencing. *Nat. Biotechnol.* (2020). doi:10.1038/s41587-020-

0470-y

28.   Ventura, T., Egan, E. J., Romanò, N. & Fidanza, A. In Vitro Model of Fetal Human

Vessel On-chip to Study Developmental Mechanobiology. *J. Vis. Exp.* **2023**, (2023).

29.   Azzoni, E. *et al.* The onset of circulation triggers a metabolic switch required for

endothelial to hematopoietic transition. *Cell Rep.* **37**, (2021).

30.   Oburoglu, L. *et al.* Pyruvate metabolism guides definitive lineage specification during

hematopoietic emergence. *EMBO Rep.* **23**, (2022).

31.   Castaño, J. *et al.* GATA2 Promotes Hematopoietic Development and Represses

Cardiac Differentiation of Human Mesoderm. *Stem Cell Reports* **13**, 515–529 (2019).

32.   de Pater, E. *et al.* Gata2 is required for HSC generation and survival. *J. Exp. Med.*

210, 2843–2850 (2013).

33.   Ling, K. W. *et al.* GATA-2 Plays Two Functionally Distinct Roles during the Ontogeny

of Hematopoietic Stem Cells. *J. Exp. Med.* **200**, 871 (2004).

34.   McGarvey, A. C. *et al.* A molecular roadmap of the AGM region reveals BMPER as a

novel regulator of HSC maturation. *J. Exp. Med.* **214**, 3731–3751 (2017).

35.   McGrath, K. E. *et al.* A transient definitive erythroid lineage with unique regulation of

the β-globin locus in the mammalian embryo. *Blood* **117**, 4600 (2011).

36.   Rejeski, K., Duque-Afonso, J. & Lübbert, M. AML1/ETO and its function as a regulator

of gene transcription via epigenetic mechanisms. *Oncogene* **40**, 5665–5676 (2021).

37.   Huynh, H. D. *et al.* IGF binding protein 2 supports the survival and cycling of

hematopoietic stem cells. *Blood* **118**, 3236 (2011).

38.   Crosse, E. I. *et al.* Multi-layered Spatial Transcriptomics Identify Secretory Factors

Promoting Human Hematopoietic Stem Cell Development. *Cell Stem Cell* **27**, 822

(2020).

39.   Hadland, B. *et al.* Engineering a niche supporting hematopoietic stem cell

development using integrated single-cell transcriptomics. *Nat. Commun. 2022 131* **13**,

1–17 (2022).

40.   Hadland, B. K. *et al.* Endothelium and NOTCH specify and amplify aorta-gonad-

mesonephros-derived hematopoietic stem cells. *J. Clin. Invest.* **125**, 2032–2045

(2015).

41.   Sandler, V. M. *et al.* Reprogramming human endothelial cells to haematopoietic cells

requires vascular induction. *Nature* **511**, 312–318 (2014).

42.   Konermann, S. *et al.* Genome-scale transcriptional activation by an engineered

CRISPR-Cas9 complex. *Nature* **517**, 583–588 (2015).

43.   Petazzi, P. *et al.* Robustness of Catalytically Dead Cas9 Activators in Human

Pluripotent and Mesenchymal Stem Cells. *Mol. Ther. - Nucleic Acids* **20**, 196–204

(2020).

44.   Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-

3587.e29 (2021).

45.   Cao, J. *et al.* The single-cell transcriptional landscape of mammalian organogenesis.

Nat. 2019 5667745 **566**, 496–502 (2019).

46.   Ge, S. X., Jung, D. & Yao, R. ShinyGO: a graphical gene-set enrichment tool for

animals and plants. *Bioinformatics* **36**, 2628–2629 (2020).

785