Can social interactions rescue "lost" behavioural defences in an allopatric cuckoo host?

Deryk Tolman[1,2,#], Katja Ronkä[1,2], Edward Kluen[1,2], Daniela Campobello[3], Rose Thorogood[1,2]

[1]HiLIFE Helsinki Institute of Life Science, University of Helsinki, Helsinki, Finland

[2]Research Programme in Organismal and Evolutionary Biology, Faculty of Biological and

Environmental Sciences, University of Helsinki, Helsinki, Finland

[3]Dipartimento STEBICEF, Università degli Studi di Palermo, Palermo, Italy

[#]Corresponding author. Contact email: deryk.tolman@helsinki.fi

ABSTRACT

Social information use by cuckoo-hosts provides an excellent example of how social interactions within species can influence selection on another. Neighbours observe each other's mobbing behaviour towards cuckoos, and then adjust costly defences back at their own nest in response. This may even allow naïve juveniles to 'learn' to recognise cuckoos, suggesting that 'social responsiveness' must play a key part in determining how hosts respond to changing rates of brood parasitism. However, if the 'cultural memory' of cuckoos is lost, can this plasticity in defences be rescued? This is fundamental for cuckoo-host coevolution, as the rates at which host defences are lost, gained, or retained determines evolutionary dynamics. Using populations of reed warblers at the northern- and southernmost extents of their range, we designed a field experiment to test whether populations allopatric to cuckoos (i) still recognise cuckoos as a threat, (ii) respond to social cues to upregulate defences, and (iii) vary in their strength of social responsiveness. We found defences and social responsiveness to be weaker in the south, where there is a much longer history of allopatry. In the north, only birds that retained recognition of cuckoos showed social responsiveness: social information about cuckoos did not increase hosts' propensity to attack (whereas it doubled in the range core), but it did increase the strength of mobbing in the 30% of birds that recognised the cuckoo at the northern range front. Most importantly, however, experimental cuckoo eggs remained in the nest, whereas in previous experiments elsewhere the same social information increased egg rejection ten-fold. This suggests that social responsiveness is more limited at the range edges, under relaxed selection, and this would incur little immediate cost to cuckoos' fitness if they (re)invade. Our study provides a rare example of geographic variation in social information use and highlights how current range expansions may influence the outcome of future species interactions.

INTRODUCTION

When their distributions change, species are likely to experience new selection pressures, but they may also be released from existing ones (Lahti et al., 2009). Relaxed selection can lead to the reduction or loss of previously adaptive traits (Lahti et al., 2009), contributing prominently to patterns of evolutionary change (Fong et al., 1995; Porter and Crandall, 2003; Rayner et al., 2022). In a changing world, past selection pressures can re-emerge, making long unexpressed traits adaptive again. Social learning and information-use could play a key role in determining whether behaviours are lost or retained (Foster, 2013). For example, social information is used by prey and hosts to assess their predation and parasitism risks (Gil et al., 2018) and predators also use information transmitted by other predators to find suitable prey, feeding back to alter prey defences (Hämäläinen et al., 2022). If and how individuals and populations respond to social information (their 'social responsiveness') therefore affects their own fitness and the strength of selection on interacting parties (Dall et al., 2005). Social behaviour may also determine whether populations are resilient to change, as information from others can rescue populations from extinction (Thorogood et al., 2018; Gil et al., 2019). However, behaviour and behavioural plasticity are relatively understudied in the context of relaxed selection, despite the expectation that they persist longer than morphological traits after selection for them ceases (Rayner et al., 2022).

Social information use by hosts of brood parasitic cuckoos provides an excellent example of how social interactions within a species can influence selection on another. After observing their neighbours mobbing a common cuckoo (*Cuculus canorus*), reed warblers (*Acrocephalus scirpaceus*) increase mobbing at their own nests (Davies and Welbergen, 2009). This eavesdropping may even allow naïve juveniles to 'learn' to recognise cuckoos (Feeney and Langmore, 2013), suggesting that social responsiveness must play a key part in determining how hosts respond to changing rates of brood parasitism. This is likely a specific response to information about cuckoos, rather than nest predators or novel stimuli, because hosts must discriminate between cuckoos and the deadly hawks they mimic (Welbergen and Davies, 2011). In optimising their discrimination thresholds, social interactions of hosts cause frequency-dependent selection of a plumage polymorphism in cuckoos, where the less common colour morph benefits from being less detectable (Thorogood and Davies, 2012). However, learning from your neighbours requires that your neighbours have experience, so what happens when this social memory of cuckoos is lost?

Host populations may be currently parasitised, previously but no longer parasitised, or naïve to parasitism. It is thought that gene flow from populations under varying degrees of coevolutionary

pressure determines how quickly a population responds to parasitism and thus the strength of selection on the parasite (Nuismer et al., 2003; Thompson, 2005). The movement of experienced individuals between populations creates an analogous information landscape (Bergman and Beehner, 2023), though population-scale changes in behaviour can be seen rapidly, even within the same generation, through social learning. Behaviour therefore varies geographically according to the local information and adaptive landscapes, but "cold spots" where selection is relaxed have received relatively less research attention (Lahti et al., 2009).

Defences are costly, so host populations that find themselves allopatric from cuckoos, such as through range expansion, tend to plastically reduce their expression of anti-parasite adaptations (Brooke et al., 1998; Lindholm and Thomas, 2000), or lose them altogether (Lahti, 2006; Lawson et al., 2020). Parasitised reed warblers in England, for instance, have reduced their defences proportional to the decline of cuckoos in the country (Thorogood and Davies, 2013), and hosts introduced to New Zealand do not mob cuckoo mounts (Hale and V. Briskie, 2007). Naïve populations may represent opportunities for rapid invasion or host switching by cuckoos. However, some hosts have also been seen to increase defences very rapidly upon the (re)invasion of brood parasites. When common cuckoos expanded their range in Japan and began parasitising new populations of azure-winged magpies, the magpies increased their rejection rates by 2-3% per year (Nakamura, 1991). Such rapid behavioural changes in a population implicate plasticity and social learning as their mechanisms rather than genotypic changes. It is unknown whether a host population allopatric to cuckoos can (re-)learn defences immediately from a few informed individuals, or if the mechanisms to do so are lost altogether. In fact, these outcomes may not be mutually exclusive and a population's ability to use social information may depend on how long ago it was parasitised, as an evolutionary loss of these mechanisms may occur over a long enough time.

Here we use allopatric populations of reed warblers from the northern- and southernmost limits of their range to test by experiment how social responsiveness varies with geography and time since last being parasitised. Reed warblers colonised Finland only about 100 years ago (Järvinen and Ulfstrand, 1980), escaping parasitism by their cuckoo gens which has yet to expand as far north. At the southern range limit in Sicily, reed warblers are unlikely to have been parasitised – their nearest known parasitised conspecifics are > 850 km away (Quaglierini, 2004) and reed warblers tend to disperse < 100 km (Ceresa et al., 2016) (but adults are highly philopatric (Ceresa et al., 2016; Halupka et al., 2017), limiting the influence of experienced individuals from parasitised populations on allopatric populations). There is further evidence of isolation in the Sicilian population in that they differ in

body size from Sardinian and mainland populations (Ientile et al., 2023). Cuckoos are present in Sicily only on mountainsides, while reed warblers are restricted to the island's coast and singular inland lake. This separation between the two species is likely to have existed since antiquity as cuckoos and reed warblers are mentioned in historical records but not in association with each other (Arnott, 2007).

First, we test whether reed warblers in these range-edge populations already recognise and respond to cuckoos. Second, whether they upregulate their defences in response to social information about cuckoos, as in core populations. Finally, we test if reed warblers in Finland, with a more recent history of parasitism, respond more to social information about cuckoos than those in Sicily.

METHODS

Study areas and host populations

We studied reed warblers at 30 reed-line bays near Helsinki, along 100km of southern coast of Finland, from 60.19848N 24.07305E to 60.34039N 25.71162E, and in the Fiume Ciane nature reserve, Sicily, Italy 37.04285N, 15.24273E, EPSG:3857. These represent the northern- and southernmost extents of the reed warbler's range. Throughout this range, they nest almost exclusively in *Phragmites* reeds. On the coast of Finland, reed beds are often expansive, and reed warblers nest in relatively low densities (distances to closest nest measured using GPS coordinates, median (lower quartile, upper quartile) 2019: 57 m (33, 95), 2020: 49 m (27, 77); range, 2019: 15–573, 2020: 9–398). In Fiume Ciane, reeds are concentrated along the river, its floodplains, and irrigation ditches, as well as a disused salt work (Saline di Siracusa). Reed warblers here nest more densely (median distance to closest nest: 36 m (24, 53)). Neither of the reed warbler populations, in Finland nor Sicily, are parasitised by cuckoos (see Introduction).

Measuring host mobbing of cuckoos before and after social information

In Finland, we conducted experiments at reed warbler nests in May-July in 2019 and 2020 and in Sicily in May-July 2022. We began on the day the fourth egg was laid, the usual day of clutch completion for this species. However, reed warblers typically lay 3-5 eggs, so some nests were tested either the day before or the day after clutch completion. Since incubation usually begins with the penultimate egg (Cramp and Brooks, 1992), any differences in nest attendance were expected to be minimal. The experimental design was essentially the same between the two locations: at each focal nest we first measured the base-line mobbing response to a model cuckoo, then presented social information of either a cuckoo or control model, and finally measured their mobbing response to a

cuckoo again at their nest. We waited at least 1.5 h between presentations and limited them to between

the hours of 10:00 and 18:00 local time as cuckoos tend to lay in the afternoon. As such, presentations

were split over 2 days for each nest.

The models used as controls were different in Finland and Sicily, as was the number of control model

presentations at the nest. In Finland, there were 3 presentations of models at the nest and 1 social

presentation: the first and last nest presentations were always a cuckoo, and the social presentation

was either a cuckoo or a magpie, while the second presentation at the nest was either a cuckoo or

magpie depending on treatment so that each nest saw the same models the same number of times. In

Sicily, each nest had 4 presentations of models at the nest - one cuckoo and one teal both before and

after a social presentation of either cuckoo or teal – with model order being randomised to control for

order effects. This slight difference in design was so that we could test an additional question: would

any threat to the nest elicit the same increase in nest defence or is this behaviour specific to cuckoos?

Previous studies have not experimentally considered this. Hypothesising that reed warblers in Sicily

would express a reduced response to cuckoos, we decided to replicate the methods of previous studies

(e.g. Davies and Welbergen, 2009) that used a novel, harmless intruder, so that we could even more

directly compare social responsiveness to populations in the range core. To compare how reed

warblers at each range edge responded to a benign intruder, we additionally presented teal models to

56 nests in Finland in 2018-2020. These were not the same nests used in the social information

experiment in Finland, and all teal models used were of females.

We used 3D-printed ABS plastic cuckoo models (3D QuickPrinting United Kingdom) painted with

acrylics by an artist, and the magpie and teal models were commercially available hunting decoys

(Live Decoys and Flambeau Outdoors, respectively). A similar range of mobbing behaviours were

elicited by our cuckoo models as studies using taxidermy models (e.g. Welbergen and Davies, 2008)

and they were produced with the same 3D templates as other, independent researchers (e.g. Marton

et al., 2019).

To measure mobbing at the nest, we placed models adjacent to the rim of the focal nest and upon the

arrival of the first bird within 3 m, we counted bill snaps and rasp vocalisations for 5 min (intensity).

Bill snaps and rasp calls are correlated with close approach, threat postures and direct attack of cuckoo

models (Welbergen and Davies, 2008), and differ according to the threat posed by the model

(Duckworth, 1991; Welbergen and Davies, 2008; Campobello and Sealy, 2010). Bill snaps and rasp

calls were summed together for data analysis and to reduce the risk of misidentifying the sound of

drying and cracking reed as bill snaps, or some "churr" warning calls as rasp calls, we conservatively

set the threshold for mobbing (propensity and intensity) at 20 combined rasps and bill snaps as per

previous studies (Welbergen and Davies, 2009; Thorogood and Davies, 2012, 2013). Three observers

carried out the experiments in Finland, each with a different set of models. We filmed each trial with

a small action camera on a pole 1 m from the nest and after the field season each observer

independently counted mobbing signals from 10 videos, blind to the field-collected data. There was

high consistency among observers' counts of bill snaps [ICC: 0.92 (95%CI 0.80–0.98), $p < 0.001$]

and rasps ICC: 0.98 (95%CI 0.95–1.0), $p < 0.001$], as assessed using the Intraclass Correlation

Coefficient (ICC) in the *irr* package in R (see Statistical analyses below). One of these same observers

also did trials in Sicily, plus two others. All field mobbing data in Sicily were confirmed by video

analysis.

To provide social information, we placed either a model cuckoo or model magpie/teal (Finland/Sicily;

treatments randomly assigned) in the reeds adjacent to a disused warbler nest, collected in previous

years, 10-15 m from the focal nest. We then played recordings of reed warblers mobbing a model

cuckoo for 10 min to simulate a close neighbour mobbing a cuckoo or teal at their nest, providing

social information for the focal pair of cuckoo activity (or magpie/teal as a nest predator/harmless

control) in the area. We used 5 mobbing audio tracks, recorded at different nests in Wicken Fen,

England, by Welbergen and Davies (2008) (also used by (Thorogood and Davies, 2016). We only

did trials at nests at least 50m from other social presentations, to avoid contamination of information.

We additionally presented social information from 4 active neighbouring nests (3 cuckoo and 1 teal;

< 20m from the focal nest) instead of disused nests in Sicily, for which none of the focal nests mobbed

a cuckoo and hence were combined with the rest of the data.

Upon visualising the mobbing data from Finland, there seemed to be greater mobbing propensities

and intensities by reed warblers towards cuckoos in bays where we had conducted this experiment

the previous year. We therefore conservatively excluded the data from these 18 bays in 2020.

Excluding these data did not qualitatively change our results and only weakened the effects we report

in this study. We also excluded 4 focal nests (1 in Finland and 3 in Sicily) where birds failed to appear

at one or both cuckoo presentations at their nest (field observations confirmed by video). In total, 81

Finnish nests (Cuckoo treatment $N = 41$; Magpie treatment $N = 40$) and 39 Sicilian nests (Cuckoo

treatment $N = 22$; Teal treatment $N = 17$) were analysed from the social information experiment.

To measure rejection rates of mimetic eggs, after the final model presentation, we chose a warbler

egg at random from the focal nest and coloured it with forty equally spaced brown dots using a

Staedtler permanent felt tip pen (Lumocolor® 314-7 (Thorogood and Davies, 2016)). In Sicily we got

rejection data from 40 nests (Cuckoo treatment $N = 21$; Teal treatment $N = 19$) and in 2019 in Finland

from 49 nests (Cuckoo treatment $N = 24$, Magpie treatment $N = 25$). In 2020 in Finland, we instead

used 3D-printed eggs with the same dimensions as redstart-gens cuckoo eggs and painted blue

(Citadel© acrylic paints, custom mixed) to simulate a non-mimetic foreign egg in 46 nests (Cuckoo

treatment $N = 29$, Magpie treatment $N = 27$). Nests were checked for rejection at least 3 days after

colouring or placement as reed warblers most often reject within 3 days (Davies and Brooke, 1988).

However, video analysis later showed that, despite vigorous pecking (4/25 nests in 2021, not included

in this experiment), reed warblers were not able to pierce and remove the 3-D printed eggs from the

nest. Therefore, we considered model eggs rejected when clear attempts had been made to reject them

or at least one own egg broken (i.e. not considered a rejection cost in this study). Finally, in Sicily we

also assessed non-mimetic egg rejection without any information of cuckoo presence ($N = 11$) and

after a cuckoo model was presented at the nest (personal information; $N = 12$) by colouring one egg

in the nest with a blue Copic felt tip pen (Copic Ciao B).

Statistical analyses

All analyses were conducted in R (v. 4.3.2; R Core Team, 2023). We used the package *glmmTMB*

(Brooks et al., 2017) for generalised linear mixed models (GLMMs) and generalised linear models

(GLMs). We present the estimated effect sizes, with their standard error, and *p*-values. We checked

for the most appropriate model error structure for each model of mobbing intensity by using QQ-

normality plots, residuals vs. predicted plots, and dispersion and outlier tests with the *DHARMa*

package (Hartig, 2022). All the models of mobbing intensity were best suited to a quasi-poisson

distribution (*nbinom1* in *glmmTMB*). Analyses of change in mobbing behaviour contained a random

effect term of nest identity to account for each nest being measured twice (before and after social

information), as did models comparing mobbing to different stimuli in Sicily to account for intruders

being presented at each nest twice. A binomial error structure and logit-link function were used for

most models of mobbing propensity, but due to the small number of nests at which mobbing occurred

in Sicily, for these data we compared mobbing propensity between treatments using Fisher's exact

tests. Reed warblers are known to assess parasitism risk using indirect cues, including distance from

nearest potential cuckoo perch (Oien et al., 1996; Welbergen and Davies, 2009). We confirmed an

even distribution of distances to possible cuckoo perches among treatments before further analysis

(gamma GLM: estimate = -0.001 ± 0.003, $z = -0.335$, $p = 0.74$). Despite data being collected across

30 different bays in Finland, including bay as a random effect did not improve model fit (Akaike

Information Criterion (AIC) > 2) so it was not part of our final models.

RESULTS

Before our social presentations, there was no difference among our treatment groups in the likelihood

that focal pairs would mob a cuckoo presented at the nest (Finland, binomial GLM: estimate = -

0.0792 ± 0.481, z = -0.165, $p = 0.87$; Sicily, Fisher's exact test: $p = 1.0$). There was also no difference

among the two years in which the experiment was conducted in Finland (binomial GLM: estimate =

0.113 ± 0.532, z = 0.213, $p = 0.83$).

We found no significant difference in reed warblers' propensity to mob cuckoos before and after

social information in Finland (binomial GLM: estimate = -1.31 ± 1.24, $z = -1.06$, $p = 0.29$), nor

between treatments (estimate = -0.993 ± 1.793, $z = -0.554$, $p = 0.58$). There was also no significant

interaction between presentation order and treatment for propensity to mob a cuckoo at the nest

(Finland: estimate = 2.93 ± 1.765, $z = 1.659$, $p = 0.097$) and therefore neither treatment independently

affected the likelihood of a cuckoo being mobbed. In Sicily, a smaller sample size combined with a

low mobbing propensity necessitated the use of a Fisher's Exact test, which suggested no association

between social treatment and the likelihood of mobbing a cuckoo ($p = 1.0$).

Of those Finnish pairs that did mob cuckoo models ($N = 32$), their rate of mobbing calls (intensity)

increased when presented with social information about cuckoos (Fig. 1A), but not magpies (Fig.

1B), i.e. there was a significant interaction between presentation (before/after social information) and

treatment (GLMM: estimate = 0.944 ± 0.458, $z = 2.05$, $p = 0.039$) but no significant differences

among presentation orders (estimate = -0.104 ± 0.340, $z = -0.301$, $p = 0.76$) or treatments (estimate =

-0.717 ± 0.482, $z = -1.49$, $p = 0.14$) separately. An increase in mobbing intensity but not propensity

in Finland suggests only reed warblers that retained recognition of cuckoos were responsive to social

information. In Sicily, there were too few nests at which birds mobbed for us to compare just

"mobbers", but there was no difference in mobbing intensity when all nests were compared between

cuckoo and teal social information (GLMM Presentation order x Treatment interaction: estimate =

0.761 ± 0.951, $z = 0.800$, $p = 0.423$; Fig. 1C & 1D).

Although only 23% (5/22) of pairs tested in Sicily mobbed model cuckoos above a threshold of 20

rasp calls and bill snaps, compared to 39% (16/41) in Finland, we did not detect a statistically

significant difference in propensity between the two countries (binomal GLM Presentation order x

Country: estimate = 5.91 ± 3.42, z = 1.726, p = 0.084). However, mobbing intensity toward cuckoos

was significantly greater in Finland than Sicily after social information (GLMM: estimate = 1.90 ±

0.67, z = -3.13, p = 0.0017). This difference between the northern and southern range edges was

specific to cuckoos, and not a general difference in nest defence, because teal, a harmless intruder

that does not require special recognition (unlike cuckoos, which mimic hawks), were mobbed at a

similar intensity between the two countries (estimate = -0.364 ± 0.400, z = -0.915, p = 0.36).

Mimetic eggs were not rejected at all in either Finland (Cuckoo treatment = 0/24; Magpie treatment

= 0/25) or Sicily (Cuckoo treatment = 0/21; Teal treatment = 0/19). Non-mimetic eggs were rejected

at low rates in Finland, but these were not influenced by social information (binomial GLM: estimate

= 0.108 ± 0.635, z = 0.170, p = 0.86): 24% (7/29) of pairs rejected a non-mimetic egg after being

presented with cuckoo social information, and 22% (6/27) with magpie social information. Although

we did not test the effect of social information on the rejection of non-mimetic eggs in Sicily, there

was no rejection of blue eggs here with either no manipulation of information (0/11) or after

presenting a cuckoo model at their nest (0/12).

289
Fig. 1. At Finnish sites, (A) social information about cuckoo activity increases mobbing intensity
towards cuckoos ($N = 41$ nests), but (B) social information about a nest predator (magpie) does not
($N = 40$). In Sicily, neither social information about (C) cuckoos ($N = 22$) nor (D) benign teal ($N = $
17) altered mobbing responses. Grey and coloured dots represent mobbing intensity prior to (grey)
and after (coloured) social information (yellow = cuckoo, blue = magpie control, green = teal control).
Any mobbing response < 20 (vertical dashed line) was considered as no response.
296
297
298

Fig. 2. Blue non-mimetic eggs were rejected at the same rate in Finland regardless of whether social
information was provided about cuckoos or magpies. Mimetic eggs were not rejected in either
country, regardless of social information.

DISCUSSION
Our results show that for some host populations at their range edges, who have escaped cuckoo
parasitism, eavesdropping on neighbours has some potential to rescue waning behavioural defences
should cuckoos invade. However, social responsiveness was contingent on birds already recognising
cuckoos: reed warbler pairs that initially mobbed cuckoos subsequently increased their mobbing
intensity after receiving social information, whereas quiet birds largely remained quiet. At the
southern range edge, reed warblers hardly mobbed cuckoos, even after witnessing the mobbing of a
cuckoo at a "neighbour's" nest. We also found that social information of a benign intruder or a nest
predator did not affect mobbing of cuckoos, suggesting that social responsiveness is a specific
strategy against cuckoos even in a population without cuckoo parasitism. Importantly, mimetic
experimental eggs all remained in nests, whereas in previous experiments elsewhere, the same social
information increased mimetic egg rejection ten-fold (Thorogood and Davies, 2016). Social

information also had no influence on the rate at which non-mimetic eggs were rejected, though 23%

were rejected in Finland after seeing a cuckoo at their nest.

Why do reed warblers hardly respond to cuckoos in Sicily, even after social information?

Small-scale geographical variation in host defences can be explained by plasticity (Lindholm, 2000;

Langmore et al., 2012), with rates of defence expression being close to optimal according to local

parasitism risk (Davies et al., 1997; Welbergen and Davies, 2009). However, by comparing

populations at a much larger geographical scale (and far from their parasitised conspecifics), we saw

that not only do defences vary, but so does their plasticity. The lack of social responsiveness we

measured in Sicily may be due to a longer (or total) absence of cuckoo parasitism, suggesting the

effects of both plasticity and adaptation. The Finnish population not only has a more recent ancestral

history of parasitism, having expanded recently from parasitised populations, but likely receives

considerable gene flow from nearby parasitised populations in the range core (Sætre et al., 2022).

This explanation – that relaxed selection has gradually lessened the expression of defences – is

supported by recent experimental evolution work on burying beetles showing that social behaviours

are less likely to be induced when the opportunity to express them is removed for multiple generations

(Bladon et al., 2023).

Evolutionary changes can also be rapid in the sudden absence of brood parasites. In as little as a

century, the eggs of African village weavers introduced to Mauritius had less between-individual

variation and within-clutch consistency of appearance, as well as less spotting, all of which are

adaptations to egg-mimicking brood parasitism (Lahti, 2005). A population of the same species

introduced a century earlier to Hispaniola, also free from cuckoos, had even more pronounced

reductions in these same egg traits (Lahti, 2005). This mirrors our results in that social responsiveness

is more reduced in the population less recently in contact with cuckoos. However, in contrast to our

results, alarm calls produced by yellow warblers that specifically warn of brood parasitic brown-

headed cowbirds seem to have persisted similarly among sites in long-term allopatry and those close

to parasitised ones (Kuehn et al., 2016), although differences in plasticity were not tested for.

Similarly, some frontline defences are retained by Australian reed warblers, which are not hosts of

its sympatric cuckoos (Chaumont et al., 2021).

Changes in introduced weaverbird egg appearance were subsequently found to prevent egg

recognition, despite recognition and rejection themselves being maintained (Lahti, 2006). Like Lahti

(2006), we saw an increase in egg rejection probability of non-mimetic eggs relative to mimetic eggs.

This implies potential changes in egg patterning rather than total loss of egg recognition, but we did

not assess egg appearance. Many species that are likely to be ex-hosts also maintain egg recognition

in the absence of parasitism (Medina and Langmore, 2015). Unlike Lahti's results, however, egg

rejection by Finnish reed warblers was still considerably less prevalent than in parasitised populations

(Stokke et al., 2008), and non-existent in Sicily, whereas the transplanted weaverbirds could reject at

the same rates as source populations when controlling for egg appearance. The costs of maintaining

phenotypic plasticity of defences may therefore be greater for some species than others (Murren et

al., 2015). Indeed, there is evidence that hosts maintaining egg rejection without selection rarely make

recognition errors when rejecting foreign eggs (Stokke et al., 2002).

Why do some pairs respond while many do not?

Reed warblers responded remarkably more to social information in previous studies in England and

mainland Italy (Campobello and Sealy, 2011), where almost identical methods were followed. At

Wicken Fen in S England, propensity to mob cuckoos increased from 46% to 75% (Davies and

Welbergen, 2009) upon experimentally learning of local cuckoo activity. A similar proportion of reed

warblers mob magpies, a nest predator, in Modena as in Finland, but in Finland cuckoos are not

treated as a greater threat to the nest than magpies like they are in Modena (Tolman et al., 2021). Why

in Finland do so few birds respond to both personal and social information of cuckoo presence?

Perhaps the Finnish population is in the process of losing defences, locally adapting to a lack of

selection by cuckoos. If we were to revisit this population many years later, would its response

diminish further and look more like that in Sicily? Genomic analyses of mobbing and non-mobbing

birds could yield some insights - perhaps mobbers are more recent immigrants from parasitised

locales. There is already some genomic evidence that reed warblers are locally adapting to the

climatic conditions at this range front (Sætre et al., 2022). However, even in parasitised populations

not all individuals exhibit defences, although it is not clear why. An alternative explanation for why

some pairs respond to social information while others do not is that the likelihood and/or costs of

recognition errors are greater for some than others (and varied between northern and southern range

edges) due to environmental and ecological differences. For example, mobbing can attract predators

(Krams et al., 2007; Marton et al., 2019) and birds with more vulnerable nests may be more reluctant

to make noise. We favour the explanation that there is local adaptation to the lack of brood parasitism,

as egg rejection behaviour also appears to be greatly diminished in both of our study locations, yet

mobbing and rejection behaviours are inherently different in their costs.

Conclusions

Overall, our results provide evidence for the degradation of socially mediated phenotypic plasticity under relaxed selection in the wild. A few experienced individuals are not sufficient to revive "ghosts of adaptation past" in the rest of the population if the tendency to use social information is reduced alongside the lack of expression of that trait. Whether we are witnessing microevolution in action (Lotem and Rothstein, 1995), or whether this reduction in responsiveness to social information is due to varied costs of plasticity, remains to be proven and warrants further research. Nevertheless, since the phenotypic traits we found to be geographically varied are defences in a coevolutionary arms race, our results also demonstrate how contemporary range expansions may influence the evolutionary outcomes of future species interactions.

ACKNOWLEDGEMENTS

We thank field assistants Pietro Di Bari, Julius Mäkinen, Juho Jolkkonen, Anna Tuominen, Anna Välkki, Purabi Deshpande, and Nora Bergman for their essential help with nest searching and monitoring, and Steve Collett for painting the model cuckoos. We would also like to thank private landowners, Uusimaa municipalities, and Metsähallitus (permission no. MH1100/2018/06.06.02) for allowing access to reed beds in Finland, and the Province of Siracusa for facilitating our work in the Fiume Ciane nature reserve in Sicily.

REFERENCES

Arnott, W. G. (2007). *Birds in the Ancient World from A to Z*. 1st ed. Routledge doi: 10.4324/9780203946626.

Bergman, T. J., and Beehner, J. C. (2023). Information Ecology: an integrative framework for studying animal behavior. *Trends in Ecology & Evolution* 38, 1041–1050. doi: 10.1016/j.tree.2023.05.017.

Bladon, E. K., Pascoal, S., Bird, N., Mashoodh, R., and Kilner, R. M. (2023). The evolutionary demise of a social interaction: experimentally induced loss of traits involved in the supply and demand of care. *Evolution Letters* 7, 168–175. doi: 10.1093/evlett/qrad016.

Brooke, M. de L., Davies, N. B., and Noble, D. G. (1998). Rapid decline of host defences in response to reduced cuckoo parasitism: behavioural flexibility of reed warblers in a changing world. *Proceedings of the Royal Society of London. Series B: Biological Sciences* 265, 1277–1282. doi: 10.1098/rspb.1998.0430.

Brooks, M. E., Kristensen, K., Benthem, K. J. van, Magnusson, A., Berg, C. W., Nielsen, A., et al.
(2017). glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated
Generalized Linear Mixed Modeling. *The R Journal* 9, 378–400. doi: 10.32614/RJ-2017-066.

Campobello, D., and Sealy, S. G. (2010). Enemy Recognition of Reed Warblers (Acrocephalus
scirpaceus): Threats and Reproductive Value Act Independently in Nest Defence
Modulation. *Ethology* 116, 498–508. doi: https://doi.org/10.1111/j.1439-
0310.2010.01764.x.

Campobello, D., and Sealy, S. G. (2011). Use of social over personal information enhances nest
defense against avian brood parasitism. *Behavioral Ecology* 22, 422–428. doi:
10.1093/beheco/arq225.

Ceresa, F., Belda, E. J., and Monrós, J. S. (2016). Similar dispersal patterns between two closely
related birds with contrasting migration strategies. *Popul Ecol* 58, 421–427. doi:
10.1007/s10144-016-0547-0.

Chaumont, M. H. J., Langmore, N. E., and Welbergen, J. A. (2021). The ghosts of parasitism past:
lingering frontline anti-brood parasite defenses in a former host. *Current Zoology* 67, 573–
583. doi: 10.1093/cz/zoab014.

Cramp, S., and Brooks, D. (1992). *Handbook of the birds of Europe, the Middle East and North
Africa. The birds of the western Palearctic, vol. VI. Warblers*. Oxford: Oxford university
press.

Dall, S. R. X., Giraldeau, L.-A., Olsson, O., McNamara, J. M., and Stephens, D. W. (2005).
Information and its use by animals in evolutionary ecology. *Trends in Ecology & Evolution*
20, 187–193. doi: 10.1016/j.tree.2005.01.010.

Davies, N. B., Brooke, M. D. L., and Kacelnik, A. (1997). Recognition errors and probability of
parasitism determine whether reed warblers should accept or reject mimetic cuckoo eggs.
Proceedings of the Royal Society of London. Series B: Biological Sciences 263, 925–931. doi:
10.1098/rspb.1996.0137.

Davies, N. B., and Brooke, M. de L. (1988). Cuckoos versus reed warblers: Adaptations and
counteradaptations. *Animal Behaviour* 36, 262–284. doi: 10.1016/S0003-3472(88)80269-0.

Davies, N. B., and Welbergen, J. A. (2009). Social Transmission of a Host Defense Against Cuckoo
Parasitism. *Science*. doi: 10.1126/science.1172227.

Duckworth, J. W. (1991). Responses of breeding Reed Warblers Acrocephalus scirpaceus to
mounts of Sparrowhawk Accipiter nisus, Cuckoo Cuculus canorus and Jay Garrulus
glandarius. *Ibis* 133, 68–74. doi: 10.1111/j.1474-919X.1991.tb04812.x.

Feeney, W. E., and Langmore, N. E. (2013). Social learning of a brood parasite by its host. *Biology
Letters* 9, 20130443. doi: 10.1098/rsbl.2013.0443.

Fong, D. W., Kane, T. C., and Culver, D. C. (1995). Vestigialization and Loss of Nonfunctional
Characters. *Annual Review of Ecology and Systematics* 26, 249–268. doi:
10.1146/annurev.es.26.110195.001341.

Foster, S. A. (2013). Evolution of behavioural phenotypes: influences of ancestry and expression.
Animal Behaviour 85, 1061–1075. doi: 10.1016/j.anbehav.2013.02.008.

Gil, M. A., Baskett, M. L., and Schreiber, S. J. (2019). Social information drives ecological outcomes
among competing species. *Ecology* 100, e02835. doi: 10.1002/ecy.2835.

Gil, M. A., Hein, A. M., Spiegel, O., Baskett, M. L., and Sih, A. (2018). Social Information Links
Individual Behavior to Population and Community Dynamics. *Trends in Ecology & Evolution*
33, 535–548. doi: 10.1016/j.tree.2018.04.010.

Hale, K., and V. Briskie, J. (2007). Response of introduced European birds in New Zealand to
experimental brood parasitism. *Journal of Avian Biology* 38, 198–204. doi:
10.1111/j.2007.0908-8857.03734.x.

Halupka, L., Wierucka, K., Sztwiertnia, H., and Klimczuk, E. (2017). Conditions at autumn stopover
sites affect survival of a migratory passerine. *J Ornithol* 158, 979–988. doi: 10.1007/s10336-
017-1472-5.

Hämäläinen, L., M. Rowland, H., Mappes, J., and Thorogood, R. (2022). Social information use by
predators: expanding the information ecology of prey defences. *Oikos* 2022, e08743. doi:
10.1111/oik.08743.

Hartig, F. (2022). *DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression
Models*. Available at: https://CRAN.R-project.org/package=DHARMa.

Ientile, R., Tagliavia, M., Cuti, N., Termine, R., Giannella, C., Nissardi, S., et al. (2023).
Morphometric versus genetic variation in the Eurasian Reed Warbler Acrocephalus
scirpaceus in Italy. *Bird Study* 0, 1–11. doi: 10.1080/00063657.2023.2228537.

Järvinen, O., and Ulfstrand, S. (1980). Species turnover of a continental bird fauna: Northern
Europe, 1850–1970. *Oecologia* 46, 186–195. doi: 10.1007/BF00540125.

Krams, I., Krama, T., Igaune, K., and Mänd, R. (2007). Long-lasting mobbing of the pied flycatcher
increases the risk of nest predation. *Behavioral Ecology* 18, 1082–1084. doi:
10.1093/beheco/arm079.

Kuehn, M. J., Peer, B. D., McCleery, R. A., and Rothstein, S. I. (2016). Yellow warbler defenses are
retained in the absence of brood parasitism but enhanced by experience with cowbirds.
Behavioral Ecology 27, 279–286. doi: 10.1093/beheco/arv145.

Lahti, D. C. (2005). Evolution of bird eggs in the absence of cuckoo parasitism. *Proc. Natl. Acad. Sci.
U.S.A.* 102, 18057–18062. doi: 10.1073/pnas.0508930102.

Lahti, D. C. (2006). PERSISTENCE OF EGG RECOGNITION IN THE ABSENCE OF CUCKOO BROOD
PARASITISM: PATTERN AND MECHANISM. *Evolution* 60, 157–168. doi: 10.1111/j.0014-
3820.2006.tb01090.x.

Lahti, D. C., Johnson, N. A., Ajie, B. C., Otto, S. P., Hendry, A. P., Blumstein, D. T., et al. (2009).
Relaxed selection in the wild. *Trends in Ecology & Evolution* 24, 487–496. doi:
10.1016/j.tree.2009.03.010.

Langmore, N. E., Feeney, W. E., Crowe-Riddell, J., Luan, H., Louwrens, K. M., and Cockburn, A.
(2012). Learned recognition of brood parasitic cuckoos in the superb fairy-wren Malurus
cyaneus. *Behavioral Ecology* 23, 798–805. doi: 10.1093/beheco/ars033.

Lawson, S. L., Leuschner, N., Enos, J. K., and Hauber, M. E. (2020). Loss of graded enemy
recognition in a Whitehead population allopatric with brood-parasitic Long-tailed Cuckoos.
Avocetta 44, 4–10. doi: 10.30456/AVO.2020104.

Lindholm, A. (2000). TESTS OF PHENOTYPIC PLASTICITY IN REED WARBLER DEFENCES AGAINST
CUCKOO PARASITISM. *Behaviour* 137, 43–60. doi: 10.1163/156853900501863.

Lindholm, A. K., and Thomas, R. J. (2000). Differences between Populations of Reed Warblers in
Defences against Brood Parasitism. *Behaviour* 137, 25–42.

Lotem, A., and Rothstein, S. I. (1995). Cuckoo-host coevolution: from snapshots of an arms race to
the documentation of microevolution. *Trends in Ecology & Evolution* 10, 436–437. doi:
10.1016/S0169-5347(00)89175-2.

Marton, A., Fülöp, A., Ozogány, K., Moskát, C., and Bán, M. (2019). Host alarm calls attract the
unwanted attention of the brood parasitic common cuckoo. *Scientific Reports* 9, 18563.
doi: 10.1038/s41598-019-54909-1.

Medina, I., and Langmore, N. E. (2015). The costs of avian brood parasitism explain variation in egg
rejection behaviour in hosts. *Biology Letters* 11, 20150296. doi: 10.1098/rsbl.2015.0296.

Murren, C. J., Auld, J. R., Callahan, H., Ghalambor, C. K., Handelsman, C. A., Heskel, M. A., et al.
(2015). Constraints on the evolution of phenotypic plasticity: limits and costs of phenotype
and plasticity. *Heredity* 115, 293–301. doi: 10.1038/hdy.2015.8.

Nakamura, H. (1991). Brood Parasitism by the Cuckoo Cuculus canorus in Japan and the Start of
New Parasitism on the Azure-winged Magpie Cyanopica cyana. *Japanese Journal of*
Ornithology 39, 1–18.

Nuismer, S. L., Thompson, J. N., and Gomulkiewicz, R. (2003). Coevolution between hosts and
parasites with partially overlapping geographic ranges. *Journal of Evolutionary Biology* 16,
1337–1345. doi: 10.1046/j.1420-9101.2003.00609.x.

Oien, I. J., Honza, M., Moksnes, A., and Roskaft, E. (1996). The Risk of Parasitism in Relation to the
Distance from Reed Warbler Nests to Cuckoo Perches. *Journal of Animal Ecology* 65, 147–
153. doi: 10.2307/5717.

Porter, M. L., and Crandall, K. A. (2003). Lost along the way: the significance of evolution in
reverse. *Trends in Ecology & Evolution* 18, 541–547. doi: 10.1016/S0169-5347(03)00244-1.

Quaglierini, A. (2004). Biologia riproduttiva della cannaiola Acrocephalus scirpaceus in una zona
umida dell'Italia Centrale. *Avocetta* 28, 15–23.

R Core Team (2023). *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R
Foundation for Statistical Computing Available at: https://www.R-project.org/.

Rayner, J. G., Sturiale, S. L., and Bailey, N. W. (2022). The persistence and evolutionary
consequences of vestigial behaviours. *Biological Reviews* 97, 1389–1407. doi:
10.1111/brv.12847.

Sætre, C. L. C., Rönkä, K., Ravinet, M., Vitulano, S., Caldarella, M., Procházka, P., et al. (2022).
Genomic consequences of range expansion and colonisation in the reed warbler
(Acrocephalus scirpaceus). 2022.11.28.518135. doi: 10.1101/2022.11.28.518135.

Stokke, B. G., Hafstad, I., Rudolfsen, G., Moksnes, A., Møller, A. P., Røskaft, E., et al. (2008).
Predictors of resistance to brood parasitism within and among reed warbler populations.
Behavioral Ecology 19, 612–620. doi: 10.1093/beheco/arn007.

Stokke, B., Honza, Moksnes, Røskaft, and Rudolfsen (2002). Costs associated with recognition and
rejection of parasitic eggs in two European passerines. *Behaviour* 139, 629–644. doi:
10.1163/15685390260136744.

Thompson, J. N. (2005). Coevolution: The Geographic Mosaic of Coevolutionary Arms Races.
Current Biology 15, R992–R994. doi: 10.1016/j.cub.2005.11.046.

Thorogood, R., and Davies, N. B. (2012). Cuckoos Combat Socially Transmitted Defenses of Reed
Warbler Hosts with a Plumage Polymorphism. *Science*. doi: 10.1126/science.1220759.

Thorogood, R., and Davies, N. B. (2013). Reed Warbler Hosts Fine-Tune Their Defenses to Track
Three Decades of Cuckoo Decline. *Evolution* 67, 3545–3555. doi: 10.1111/evo.12213.

Thorogood, R., and Davies, N. B. (2016). Combining personal with social information facilitates
host defences and explains why cuckoos should be secretive. *Sci Rep* 6, 19872. doi:
10.1038/srep19872.

Thorogood, R., Kokko, H., and Mappes, J. (2018). Social transmission of avoidance among
predators facilitates the spread of novel prey. *Nat Ecol Evol* 2, 254–261. doi:
10.1038/s41559-017-0418-x.

Tolman, D., Campobello, D., Rönkä, K., Kluen, E., and Thorogood, R. (2021). Reed Warbler Hosts Do
Not Fine-Tune Mobbing Defenses During the Breeding Season, Even When Cuckoos Are
Rare. *Frontiers in Ecology and Evolution* 9, 547. doi: 10.3389/fevo.2021.725467.

Welbergen, J. A., and Davies, N. B. (2008). Reed warblers discriminate cuckoos from sparrowhawks
with graded alarm signals that attract mates and neighbours. *Animal Behaviour* 76, 811–
822. doi: 10.1016/j.anbehav.2008.03.020.

Welbergen, J. A., and Davies, N. B. (2009). Strategic Variation in Mobbing as a Front Line of
Defense against Brood Parasitism. *Current Biology* 19, 235–240. doi:
10.1016/j.cub.2008.12.041.

Welbergen, J. A., and Davies, N. B. (2011). A parasite in wolf's clothing: hawk mimicry reduces
mobbing of cuckoos by hosts. *Behavioral Ecology* 22, 574–579. doi:
10.1093/beheco/arr008.

564