Relaxed selection diminishes social memory and expression of host defences against cuckoos

Deryk Tolman[1,2,*], Katja Rönkä[1,2], Edward Kluen[1,2], Juho Jolkkonen[3], Pietro Di Bari[1], Renzo Ientile[4], Daniela Campobello[5], Rose Thorogood[1,2]

[1]HiLIFE Helsinki Institute of Life Science, University of Helsinki, Helsinki, Finland

[2]Research Programme in Organismal and Evolutionary Biology, Faculty of Biological and Environmental Sciences, University of Helsinki, Helsinki, Finland

[3]Department of Biological and Environmental Science, University of Jyväskylä, Jyväskylä, Finland

[4]Department of Biological, Geological and Environmental Sciences, Section of Animal Biology "M. La Greca", University of Catania, Catania, Italy

[5]Dipartimento STEBICEF, Università degli Studi di Palermo, Palermo, Italy

*Corresponding author. **Email:** deryk.tolman@helsinki.fi

Keywords: relaxed selection, geographic mosaic of coevolution, host-parasite coevolution, behavioural plasticity, social information

Abstract

Behavioural defences are a key adaptation across animals, but confirming decay under relaxed selection is challenging. If expression involves decision-making, then the defence may only be expressed when the right environmental cues are available (cryptic plasticity). Even if defences wane, they could be quickly restored once selection resumes as animals can learn by observing others (social memory). Here we tested cryptic plasticity and social memory under relaxed selection in a cuckoo host, the Common Reed Warbler, by manipulating perceived parasitism risk with social information in geographically-distinct areas differing in allopatry (100 to 1000 years). Despite predicting social information of a cuckoo vs. a control would elicit otherwise cryptic behavioural defences, only birds that mobbed model cuckoos prior to manipulation upregulated this defence, and only in the more recently allopatric population. Social information also had no effect on low-to-absent rates of foreign egg rejection, a second line of defence common to cuckoo host species. These results contrast with parasitised locations where similar methods provoke mobbing and increase egg rejection ten-fold. Our study therefore demonstrates that, even when manipulating information to account for reaction norms, behavioural defences may degrade rapidly under relaxed selection, and thus facilitate a geographic mosaic of coevolution.

Main Text

Introduction

As species' ranges expand or contract, their exposure to enemies and selection pressures can also

change (Gilman *et al.* 2010). Where selection relaxes (i.e. with allopatry), traits that were previously

adaptive are expected to reduce (e.g. due to drift), particularly if maintaining the trait is costly (Lahti

et al. 2009). However, while behavioural defences are common across the animal kingdom

(Candolin & Wong 2015), whether these respond to relaxed selection is rarely tested (Foster 2013;

Lahti *et al.* 2009). Behaviours are assumed unlikely to be lost due to complex genetic architectures

(pleiotropy) and underlying trait variation that can be co-opted in response to other selective

pressures (Lahti *et al.* 2009). However, another key issue is that loss of behaviour can be

challenging to measure. Behavioural responses depend on prior experience and availability of

information about the relative costs or benefits of taking action [i.e. information ecology (Dall *et al.*

2005)]. Therefore, the absence of a response does not necessarily indicate a lack of enemy

recognition but rather cryptic plasticity (Foster 2013). Taking an informational approach (Gil *et al.*

2018; O'Connor *et al.* 2019) and manipulating cues in allopatry, however, makes it possible to

approximate the retention of behavioural reaction norms in the wild.

Social 'memory' could play a key role in determining whether behaviours are lost or retained. For

example, prey and hosts observe the behaviour of others to learn about novel threats (Feeney &

Langmore 2013; Langmore *et al.* 2012; Rowell *et al.* 2020) and assess risk of predation or

parasitism (Gil *et al.* 2018), while predators use social information to find suitable prey (Hämäläinen

et al. 2022). Therefore, how individuals adjust their behaviour in response to social information [i.e.

social responsiveness; (de Groot *et al.* 2023)] is likely to affect both their own fitness and the

strength of selection on interacting parties (Cantor *et al.* 2021; Thorogood & Davies 2012). Social

responsiveness may also determine whether populations are resilient to environmental change (Gil

et al. 2019; Thorogood *et al.* 2018), especially as the utility of social information depends on the

availability of more-informed 'demonstrators'. For example, gene flow among populations

experiencing variable selection pressures (i.e. trait re-mixing) is thought to determine how quickly

a host population responds to shifts in the strength of selection by parasites [Geographic Mosaic

Theory of Coevolution; (Thompson 2005)]. If experienced individuals from 'hotspots' of current

selection also introduce social information to naïve 'cold spots' of relaxed selection, social learning

could speed up the recovery of defences in host populations that are later reinvaded by a previous

enemy. However, while recent experimental evolution studies show that social behaviours can

diminish very rapidly after selection is removed (Bladon *et al.* 2023), the role of social cues in

maintaining behaviours in 'cold spots' is still largely unknown (Lahti *et al.* 2009; Langmore *et al.*
2012).
73
Avian brood parasitism provides an ideal system to test if behavioural defences and social
responsiveness decline in allopatry, altering the potential for hosts to respond to future (re)invasion.
Virulent brood parasites lay eggs in the nests of host species and can exert strong selection for
defensive behaviours such as mobbing or rejecting suspicious eggs. However, many brood
parasites are aggressive or Batesian mimics (Feeney *et al.* 2015; Welbergen & Davies 2011),
selecting for hosts to adjust defences according to local risk and minimise costly mistakes
(Lindholm & Thomas 2000; Thorogood & Davies 2013). Parasitism can also vary in time and space
with some host populations or species escaping parasitism for many generations (Grim & Stokke
2016). Over time in allopatry, plasticity is expected to be lost with hosts either becoming resistant
to reinvasion (i.e. fixation of defences) or susceptible to extinction if defences wane (Soler 2014)
but it is unclear which trajectory will occur. Some species long-separated from brood parasites (i.e.
hundreds of thousands to millions of years) show persistent (experimentally-induced) egg rejection
(Bolen *et al.* 2000; Briskie *et al.* 1992; Chaumont *et al.* 2021; Moskát, C. *et al.* 2002; Rothstein
2001), but this common defence appears to reduce in others (Briskie *et al.* 1992; Davies & Brooke
1989; Soler & Møller 1990; Yang *et al.* 2015, 2021). Until recently, mobbing of adult brood parasites
has received less attention (Feeney *et al.* 2012), but this frontline defence also appears to degrade
(Briskie *et al.* 1992; Lawson *et al.* 2023) or be preserved (Briskie *et al.* 1992; Chaumont *et al.* 2021;
Kuehn *et al.* 2016) under recent [e.g. lost in < 100 years; (Hale & V. Briskie 2007; Lawson *et al.*
2020)] or longer allopatry. Conversely, apparently naïve species can also obtain host defences
very rapidly upon invasion (Nakamura 1991) but whether these changes represent the cryptic
persistence of defences in the population, gene flow from parasitised areas, or novel learning is
unknown.
96
Here we use allopatric populations of Common reed warblers (*Acrocephalus scirpaceus*) at the
northern- and southernmost extents of their European breeding range to test for the retention of
behavioural defences and social responsiveness to brood-parasitic Common cuckoos (*Cuculus*
canorus). Reed warblers colonised Finland ca.100 years ago (Järvinen & Ulfstrand 1980), most
likely expanding from the southwest (Bergman *et al.* 2025) and away from regularly parasitised
areas in the range core (Avilés *et al.* 2006). Cuckoos in Finland use different host species, away
from reedbeds. In Sicily, reed warblers are likely to have escaped parasitism by cuckoos for much
longer – cuckoos only occur on mountainsides while reed warblers are largely restricted to coastal
areas, and mentions of both species in historical records are not in association with each other
(Arnott 2007). At parasitised sites in their range-core, reed warblers upregulate mobbing

(Campobello & Sealy 2011; Davies & Welbergen 2009) and egg rejection (Thorogood & Davies
2016) defences after observing their neighbours attack cuckoos. Among populations not currently
exposed to parasitism, however, egg rejection rates vary widely [e.g. from 0–40%, (Thorogood &
Davies 2013)] and mobbing responses are reduced [e.g. 25% of pairs; (47)]. However, as in other
studies of brood-parasite host defences in allopatry (see also Rothstein 2001), perceived parasitism
risk has not yet been manipulated (e.g. via social information) so the extent to which host defences
reduce under relaxed selection remains unclear.
114
Therefore, we experimentally manipulated social information and simulated parasitism in field
experiments to test if defences and/or social responsiveness decay in allopatry. We predicted that
if defences are retained, then providing social information should induce plasticity (i.e. social
responsiveness) and lead to an increase in mobbing (propensity and intensity) and egg rejection,
as previously shown in range-core populations. However, if defences are not upregulated in
response to our manipulated cues, then this would provide evidence that these behaviours are
being lost under relaxed selection. Additionally, we expected any loss of defences or weakening of
social responsiveness to be more pronounced in Sicily as reed warblers there are likely to have
been in allopatry for longer.
124
125

Materials and Methods
Study areas and host populations
We studied unparasitised populations of Common reed warblers breeding in May-July around
Helsinki, southern Finland (30 discrete reed-lined bays along 100km of coastline, 60.19848N
24.07305E to 60.34039N 25.71162E, in 2019 and 2020) and in the Fiume Ciane nature reserve
near Siracusa, Sicily, Italy (37.04285N 15.24273E, in 2022), avoiding re-use of areas across years
in Finland to avoid contamination of experience or pseudoreplication. Reed warblers nest at
relatively low densities in Helsinki (Tolman *et al.* 2021) but are more numerous in Fiume Ciane
(median distance to closest nest = 36 m, inter-quartile range = 24-53 m). We found nests by cold-
searching or monitoring singing males and recorded location and distance to the nearest potential
cuckoo vantage point [tree >3 m in height, a possible indirect cue of parasitism risk; (Oien *et al.*
1996; Welbergen & Davies 2009)] using handheld GPS (accuracy ± 1 m).
138

Experimental protocols
To compare results to previously-studied parasitised populations in the range-core, we adopted
published protocols (Davies & Welbergen 2009; Thorogood & Davies 2016) as follows.
Experiments started on the day the fourth egg was laid [clutch sizes = 3-5 eggs, incubation begins

with the penultimate egg; (Cramp & Brooks 1992)]. At each nest we measured the change in
mobbing response to a cuckoo after presentation of social information of either a cuckoo or control.
We used the same 3D-printed model cuckoos (painted by an artist, see 50 for details) in Finland
and Sicily but controls differed. In Finland we used a commercially-available plastic magpie (Live
Decoys) to assess whether changes in mobbing may be due to nest predators or specific to
cuckoos, and in Sicily we used plastic female teal hunting decoys (Flambeau Outdoors), similar to
previous range-core experiments using innocuous controls (Davies & Welbergen 2009). Although
this difference precludes comparisons of responses to controls, our focus was on probing the
generality of previous findings in the context of relaxed selection, and at parasitised sites there has
been similar variation in controls and presentation orders (Campobello & Sealy 2011; Davies &
Welbergen 2009; Thorogood & Davies 2012, 2016). To also compare how reed warblers at each
site responded to a benign control, we presented teal models at 48 additional nests in Finland in
2018-2020. Model cuckoos elicit similar responses as taxidermy specimens (Chen *et al.* 2022;
Welbergen & Davies 2008) and ours were printed using the same 3D template as independent
work (e.g. Marton *et al.* 2019).

To assay mobbing behaviour [bill snaps and rasping calls (Welbergen & Davies 2008)], we placed
a model adjacent to the nest rim and recorded latency of the first bird to arrive within 3 m (using a
stopwatch, seconds). We filmed each trial with a small action camera on a pole 1 m from the nest
while observing from 3-5 m away, depending on visibility, and counted mobbing vocalisations for 5
min (mobbing intensity). Three observers carried out experiments in Finland, each with a different
set of models. One observer also performed trials in Sicily. There was high consistency among
observers' counts of bill snaps [ICC: 0.92 (95%CI 0.80–0.98), $p < 0.001$] and rasps [ICC: 0.98
(95%CI 0.95–1.0), $p < 0.001$], as assessed using the Intraclass Correlation Coefficient (ICC) in the
irr package in R (see Statistical analyses below) from data collected later using 10 videos and blind
to field-collected data.

We used simulated neighbour presentations to provide social information 10 m from the focal nest
(cf. active neighbours < 40 m in Davies & Welbergen 2009). A previously-collected reed warbler
nest was attached to reeds at a similar height to the focal nest, with either a model cuckoo or control
adjacent to it (treatments assigned alternately). We then broadcast one of 5 playback tracks of reed
warblers mobbing a model cuckoo [recorded for range-core experiments; (Davies & Welbergen
2009; Thorogood & Davies 2016)] for 10 min from a speaker placed under the dummy nest. Focal
nests were at least 40 m from other nests or experimental neighbour playbacks to avoid
contamination of information. Live mobbing by neighbours does not influence social
responsiveness to playbacks (Davies & Welbergen 2009; Thorogood & Davies 2012) and simulated

vs. active nests elicited similar responses to social information treatments in previous experiments
(Campobello & Sealy 2011; Thorogood & Davies 2016). In Sicily, where densities were higher, we
used active neighbouring nests for social information presentations in 4 trials (3 cuckoo and 1 teal;
< 20m from the focal nest) but none of these focal pairs mobbed a cuckoo.

In Finland, we always presented the cuckoo model at the start of the experiment (P1) and after
(P3) presentation of social information (SI) of either a magpie control or a cuckoo (Fig. 1A). To
avoid pairs in the cuckoo SI treatment being exposed to cuckoos more (i.e. P1, SI, P3) than in the
magpie SI treatment (P1 & P3 only), we included a second presentation (P2) of either a cuckoo
(magpie SI treatment) or magpie (cuckoo SI treatment) before SI (Fig. 1A). This design also allowed
us to compare whether any changes in behaviour were due to repeated exposure rather than the
SI treatment. To rule out order effects, in Sicily (Fig. 1D) we followed the same design as Davies &
Welbergen (Davies & Welbergen 2009): here each nest was presented with a cuckoo and control
(teal) in random order before (P1 & P2) and after (P3 & P4) social information (SI). We waited at
least 1.5 h between presentations at both sites and conducted trials between 10:00 and 18:00 local
time as cuckoos tend to lay in the afternoon. As such, presentations were split over 2 days for each
nest.

In trials where mobbing responses were very weak (i.e. fewer than 20 vocalisations), we later
confirmed sounds from video recordings of the trial and excluded five cases where three observers
could not discern or agree on sound categorisation. We also excluded 4 nests (1 in Finland and 3
in Sicily) where birds failed to appear at one or more cuckoo presentations at their nest (field
observations confirmed by video). In total, we analysed mobbing from 83 Finnish nests (Cuckoo
treatment n = 42; Magpie treatment n = 41) and 40 Sicilian nests (Cuckoo treatment n = 22; Teal
treatment n = 18).

Measuring egg rejection defences
After the final model presentation, we coloured one egg at random from the focal nest with 40
equally-spaced brown dots using a Staedtler permanent felt tip pen (Lumocolor® 314-7) to measure
rejection (as in Thorogood & Davies 2016) of a pseudo reed warbler-gens 'cuckoo egg' (Finland:
Cuckoo treatment n = 24, Magpie treatment n = 25; Sicily: Cuckoo treatment n = 22, Teal treatment
n = 18). Due to a low response (see Results), for trials conducted in Finland in 2020 we then used
custom 3D-printed eggs with the same dimensions and colour (Citadel© blue acrylic paints, custom
mixed) as a highly non-mimetic redstart-gens cuckoo egg (Cuckoo treatment n = 29, Magpie
treatment n = 27). In all trials, nests were checked 3-6 days later for removal of the 'cuckoo egg'
(reed warblers most often reject within 3 days; Davies & Brooke 1988) or evidence of damage (reed

warblers were not able to pierce and remove 3D-printed models despite vigorous pecking: 4/25

rejection trials filmed in a subsequent year, unpublished data). Finally, in Sicily we also assessed

non-mimetic egg rejection without any information of cuckoo presence (n = 11) and with personal

information of parasitism risk (i.e. by presenting a cuckoo model at the nest; n = 12) by colouring

one egg in the nest entirely with a blue felt-tip pen (Copic Ciao B).

Statistical analyses

All analyses were conducted in R v.4.3.2 (R Core Team 2023). First, we used GLMs to check for

bias of potential indirect cues (i.e. distance to potential cuckoo vantage points, see Supporting

Information S1) and to confirm that initial responses to cuckoos (i.e. P1 in Finland, or P1|P2 in

Sicily) were similar in the treatment groups and among years of the study (in Finland). We then

used GLMMs with a random effect for nest identity when modelling change in response across

multiple presentations (i.e. P1 vs. P3, or P1 vs. P2) as implemented in the package *glmmTMB*

(Brooks *et al.* 2017). To investigate whether social information led to a change in mobbing

propensity or intensity, we only included responses to the first and last presentation of a cuckoo at

the nest (i.e. P1 and P3 in Finland). However, due to a large proportion of zeroes and lack of

suitable fit for a zero-inflated model structure, we adopted a hurdle approach to model mobbing

propensity (i.e. binary variable, binomial error structure and logit-link function) and intensity (i.e.

number of calls in at least one presentation > 0, quasi-poisson (*nbinom1*) distribution) separately

in Finland. For mobbing data from Sicily, we used Fisher's Exact tests to compare treatment effects

on mobbing propensity and quasi-binomial GLMMs to investigate changes in mobbing intensity

across all pairs, due to the smaller sample size. To investigate whether social information affected

egg rejection decisions, we used a GLM (binomial error structure, logit-link function) which included

clutch size as a covariate predictor. Despite data being collected across 30 discrete sites in Finland,

including location as a random effect did not improve model fit (Akaike Information Criterion > 2)

so it was not part of our final models.

Finally, we investigated latency to approach the nest during experiments to assess more subtle

differences in reaction to cuckoos (i.e. cryptic recognition without a mobbing response). We used

separate GLMMs with Tweedie distributions to test if change in latency to approach across

presentations differed (i) between treatments and (ii) between mobbing versus non-mobbing pairs

as we lacked power to test a Presentation order*SI treatment*Mobbing status 3-way interaction.

We used QQ-normality plots, residuals vs. predicted plots, and dispersion and outlier tests using

package *DHARMa* (Hartig 2022) to assess assumptions and fits of all models, and we present

estimated mean effect sizes (± standard error), *z*-statistics, and *p*-values. Summary tables of all

models are available in Supporting Information S2.

Results

Mobbing defences

When first presented with a cuckoo at the nest, 36% (30/83) of pairs in Finland and 13% (5/40) of pairs in Sicily showed a mobbing response (Fig. 1). This propensity to mob did not differ between social information (SI) treatment groups (Finland, GLM: estimate = 0.16 ± 0.46, z = -0.35, p = 0.73; Sicily, Fisher's Exact test: p = 1.0; Table S2.1 in Supporting Information) or among years of the study in Finland (GLM: estimate = 0.25 ± 0.48, z = 0.52, p = 0.60; Table S2.1).

In Finland, providing social information about the risk of parasitism in the local area did not induce significantly more reed warblers to mob cuckoos than pairs exposed to information about magpies (GLMM Presentation order * SI treatment: estimate = 3.99 ± 2.39, z = 1.67, p = 0.099, Fig. 1B, Fig. 2A; Table S2.2) and this lack of interaction was not because social information about magpies also induced increased mobbing propensity (Fig. 1C, Fig. 2A).

Social information about cuckoos did, however, increase mobbing intensity (i.e. number of signals) of pairs who responded (across treatments n = 36 pairs with calls >0, Fig. 1B,C; GLMM Presentation order * SI treatment: estimate = 0.97 ± 0.45, z = 2.14, p = 0.032, Fig. 2B; Table S2.2) compared to those who received social information about magpies where there was no significant change among presentations (GLMM estimate = -0.28 ± 0.33, z = -0.84, p = 0.40, Fig. 2B; Table S2.2). Instead, these pairs were just as likely to increase or reduce mobbing intensity by the third presentation (P3; Fig. 1C).

In Sicily, social information about cuckoos had no effect on either mobbing propensity (across treatments n = 10 pairs with calls >0, Fisher's exact test: p = 1.0, Fig. 2A) or intensity (GLMM Presentation order * SI treatment: estimate = -1.20 ± 0.88, z = -1.36, p = 0.17, Fig. 2B; Fig. 1E, F; Table S2.3). This meant that, overall, reed warblers in Finland were more likely to mob cuckoos presented during our experiments (19/42, 45%) than pairs in Sicily (5/22, 23%; GLM: estimate = -1.03 ± 0.42, z = -2.45, p = 0.014; Table S2.4), and at a greater intensity (n = 24, GLM: estimate = 1.44 ± 0.72, z = -2.01, p = 0.044; Table S2.4). These differences between the northern and southern

areas are likely to be specific to cuckoos, as teal were mobbed at similar intensities between both

populations (n = 96, GLM: estimate = -0.39 ± 0.31, z = -1.27, p = 0.21; Table S2.5).

Egg rejection defences

Reed warblers did not reject 'mimetic' brown-spotted eggs in either Finland (Cuckoo treatment =

0/24; Magpie treatment = 0/25; Fig. 3) or Sicily (Cuckoo treatment = 0/21; Teal treatment = 0/19),

and there was no rejection of non-mimetic blue eggs in Sicily (no cuckoo presentation = 0/11, after

cuckoo presentation at nest = 0/12). Blue eggs were however rejected at low rates in Finland,

although this did not differ between social information treatments (GLM: estimate = 0.11 ± 0.64, z

= 0.17, p = 0.97; Fig. 3; Table S2.6).

Recognition of cuckoos vs. response

In Finland, reed warblers became faster to approach their nests across subsequent presentations

of cuckoos (i.e. P1 vs P3; GLMM estimate = -0.45 ± 0.19, z = -2.37, p = 0.018; Fig. 4A; Table S2.7),

regardless of social information (i.e. difference in SI treatment at P3; GLMM estimate = 0.30 ± 0.27,

z = 1.12, p = 0.26; Fig. 4A; Table S2.7). Birds were, however, slower to approach a magpie than a

cuckoo at P2 (i.e. difference in SI treatment at P2; GLMM estimate = 0.72 ± 0.26, z = 2.81, p =

0.005; Fig. 4A; Table S2.8) even if they never mobbed (P2 data only, GLM mobbing status * SI

treatment: estimate = 0.11 ± 0.38, z = 0.29, p = 0.78; Fig. 4A; Table S2.8) suggesting that latency

was specific to the threat at the nest. Mobbing birds were quicker to approach cuckoos at first (P1

data only, GLM estimate = -0.42 ± 0.19, z = -2.29, p = 0.022; Fig. 4B; Table S2.9), but non-mobbing

birds reduced their latencies more across presentations (GLMM Presentation order * mobbing

status interaction: estimate = 0.54 ± 0.28, z = 1.96, p = 0.049; Fig. 4B; Table S2.10) so that there

was no difference by the end of the experiment (GLM estimate = -0.12 ± 0.21, z = 0.58, p = 0.56;

Fig. 4B; Table S2.9). Reed warblers in Sicily approached cuckoos at the nest more quickly than in

Finland (GLM estimate = -0.47 ± 0.20, z = -2.37, p = 0.018; Table S2.11), but latencies did not

change over the experiment, either overall (GLMM Presentation order: estimate = 0.20 ± 0.21, z =

0.95, p = 0.34; Table S2.11) or differently according to social information treatments (GLMM

Presentation order * SI treatment: estimate = 0.67 ± 0.42, z = 1.58, p = 0.12; Table S2.12). They

also approached teal as quickly as they did cuckoos (estimate = 0.065 ± 0.168, z = 0.39, p = 0.67;

Table S2.13).

Discussion

A wealth of prior research in parasitised areas has revealed that reed warblers' defences

predictably match local brood parasitism risk, including responding to social information of cuckoos

by consistently increasing mobbing and egg rejection rates (Campobello & Sealy 2011; Davies &
Welbergen 2009; Thorogood & Davies 2016). But, theory also predicts that the loss of defences in
non-parasitised populations is a key part of determining coevolutionary outcomes (Soler 2014;
Thompson 2005). Although behaviours are assumed hard to lose under relaxed selection, by using
social information to simulate increased brood parasitism risk, we found that defences and social
responsiveness of a common host of cuckoos are degrading in areas of allopatry. At the recent
range front in Finland, only pairs that mobbed a cuckoo increased their mobbing intensity after
receiving social information. The other two-thirds of birds largely remained quiet, although they
differentiated cuckoos from magpies by approaching the nest more quickly, suggesting some
recognition was retained. Our findings in Finland are enhanced by our results from the Sicilian
population, where reed warblers have been in allopatry for longer. There, they showed an even
greater reduction in defences and no evidence of social information use. Importantly, mimetic
'cuckoo' eggs remained in nests, and although non-mimetic eggs were rejected at low rates in
Finland, social information had no influence on this defence either. This suggests that both
allopatric populations would be unlikely to express plasticity or mount significant resistance should
cuckoos invade.

Our results are likely evidence for the decay of ancestral defences rather than a methodological
failure to detect responses, for three reasons. First, providing social information experimentally led
to an increase in mobbing intensity in Finland, even if not in Sicily. Second, we followed very similar
methods to previous studies conducted with reed warblers at parasitised sites, where social
information heightened near-absent egg rejection of similarly manipulated eggs ten-fold
(Thorogood & Davies 2016) and induced mobbing of cuckoos (Campobello & Sealy 2011; Davies
& Welbergen 2009; Thorogood & Davies 2012), including for pairs that showed no initial response
[e.g. 54% of quiet pairs mobbed cuckoos after social information: (Davies & Welbergen 2009)].
Third, the differences we detected were specific to cuckoos as responses to a control teal and nest
predator (magpie) were similar at our sites and to parasitised areas (Welbergen & Davies 2008;
Campobello & Sealy 2010; Tolman *et al.* 2021; Welbergen & Davies 2012) despite large ecological
differences (i.e. climate, nesting density). By using consistent experimental designs to previous
studies, testing a large number of reed warblers nesting across a broad range of environmental
variation, and considering social responsiveness explicitly, we can therefore be confident that the
weak defences in our allopatric study populations are not the result of cryptic plasticity (Scheiner &
Levis 2021). Instead, they provide compelling evidence that behavioural defences may sometimes
be lost in allopatry from cuckoos.

Given that studies of animal culture across a range of taxa demonstrate that the spread of even
arbitrary behaviours is possible (Whiten 2021), it is surprising that we did not detect evidence for
social learning among 'naïve' reed warblers (i.e. those that did not mob cuckoos). For example,
juvenile superb fairy wrens begin to mob cuckoos after observing adults do so (Feeney & Langmore
2013) and naïve individuals of several species can be trained to recognize novel predators [e.g.
introduced species (Rowell *et al.* 2020)] when paired with mobbing calls. Often social learning of
threats requires only one exposure (i.e. similar to our experimental design; reviewed by Crane &
Ferrari 2013) and can even be effective when using arbitrary cues (Hämäläinen *et al.* 2022). Why
then did so many birds not mob cuckoos or reject eggs? One possible explanation is that
ontogenetic experience or repeated exposure may be necessary before behaviours are expressed,
as in the antipredator responses of several fish and mammals (Kelley & Magurran 2003; Mineka *et*
al. 1984) and personal encounters with avian or piscine brood parasites (Blažek *et al.* 2018; Hauber
et al. 2006; Stokke *et al.* 2007). While we cannot exclude this possibility it seems unlikely to explain
the lack of response we detected here: we provided three encounters with cuckoos at or near their
nests, yet most birds still did not respond. Perhaps with relaxed selection, there is an initial loss of
recognition that then limits hosts' potential for observational learning.
369
Temporal and geographical variation in host defences has typically been explained by plasticity
(Langmore *et al.* 2012; Lindholm 2000; Welbergen & Davies 2012), particularly in reed warblers
where individuals track local parasitism risk during egg-laying and adjust defences accordingly
(Thorogood & Davies 2013). However, here we found only limited evidence for plasticity in our
allopatric populations in response to either indirect environmental or experimentally-manipulated
cues of heightened parasitism risk. This is surprising given that in a previous study at an
unparasitised site closer to the range core (Llangorse lake, Wales) and presumed to have been
allopatric for at least 130 years [i.e. similar to Finland; (Lindholm 1999)], reed warblers responded
more aggressively to cuckoo models if nesting closer to vantage points for cuckoos, if they were
present (Welbergen & Davies 2012). We found no evidence for this indirect cue use: instead, birds
nesting further from vantage points in Finland were *more* likely to mob, while mobbing in Sicily was
unaffected (Supporting Information S1). Nevertheless, of the birds that mobbed, their initial
intensities were similar across all three allopatric sites [Finland: 225 ± 43 calls per 5-min, Sicily:
280 ± 120, Llangorse: 281 ± 64; (Welbergen & Davies 2012)], suggesting that while the frequency
of birds retaining recognition may vary across allopatric populations, they can respond similarly
(see Welbergen & Davies 2012). In Sicily, a site we assume to have been in allopatry for much
longer, there was no social responsiveness, suggesting that individual reaction norms may weaken
in allopatry, and potentially lead to the eventual loss of defences in the absence of cuckoo
parasitism (Soler 2014). If we were to revisit the Llangorse or Finnish populations in future, would

their responses have diminished further and look more like those in Sicily? Quolls' antipredator
behaviours have been found to decay on a predator-free island over even fewer generations than
have existed of reed warblers in Finland (Jolly *et al.* 2018), suggesting gene flow mediates the
speed at which behaviours can be lost. There is already some genomic evidence from reed
warblers to suggest local adaptation to the strength of parasitism (Smith *et al.* 2025), but whether
the decay in defences we detected is accompanied by genetic change is yet to be determined.
395
Overall, our results suggest that it may be possible for host populations freed from parasitism to
lose expression of defences, rendering them vulnerable to future reinvasion. The recovery of
resistance is unlikely to be enhanced through social transmission, because even if there are
individuals who retain recognition of cuckoos, our results show that social responsiveness may also
be lost. Such a reduction of defences under relaxed selection is one of the key components of
geographic mosaic theory explaining the maintenance of coevolution across time and space: rather
than the arms race continually escalating, parasites are hypothesized to recurrently invade host
populations, or host species, whose defences have waned in their absence. These so-called
coevolutionary cycles (Rothstein 2001) are thought to only exist during 'intermediate phases' of
host-parasite interactions, before defences reach fixation (Soler 2014), but the rapidity with which
a brood parasite host species can acquire (or lose) successful resistance has received little
attention. Our results suggest that range expansions and a geographic mosaic of coevolution may
moderate the arms race, with defences decaying within even relatively short time frames [e.g.
approximately 35 generations in Finland (Ceresa *et al.* 2015)], allowing the longer-term persistence
of parasites until host switching becomes more favourable (Nuismer *et al.* 2003). As both social
information use and behavioural defences against enemies are common across the animal
kingdom, and species' ranges are changing rapidly, it is now crucial to determine how socially-
mediated plasticity degrades under relaxed selection in the wild.
414
415

416   **References**

417   Arnott, W.G. (2007). *Birds in the Ancient World from A to Z*. 1st edn. Routledge.
418   Avilés, J.M., Stokke, B.G., Moksnes, A., Røskaft, E., Åsmul, M. & Møller, A.P. (2006). Rapid
419   increase in cuckoo egg matching in a recently parasitized reed warbler population.
420   *Journal of Evolutionary Biology*, 19, 1901–1910.
421   Bergman, N.M., Lehikoinen, P., Kluen, E., Bensch, S., Lo Cascio Sætre, C., Eroukhmanoff, F., *et al.*
422   (2025). Combined Evidence Reveals the Origin of a Rapid Range Expansion Despite
423   Retained Genetic Diversity and a Weak Founder Effect. *Molecular Ecology*, n/a, e70077.
424   Bladon, E.K., Pascoal, S., Bird, N., Mashoodh, R. & Kilner, R.M. (2023). The evolutionary demise
425   of a social interaction: experimentally induced loss of traits involved in the supply and
426   demand of care. *Evolution Letters*, 7, 168–175.
427   Blažek, R., Polačik, M., Smith, C., Honza, M., Meyer, A. & Reichard, M. (2018). Success of cuckoo
428   catfish brood parasitism reflects coevolutionary history and individual experience of
429   their cichlid hosts. *Science Advances*, 4, eaar4380.
430   Bolen, G.M., Rothstein, S.I. & Trost, C.H. (2000). Egg Recognition in Yellow-Billed and Black-Billed
431   Magpies in the Absence of Interspecific Parasitism: Implications for Parasite-Host
432   Coevolution. *The Condor*, 102, 432–438.
433   Briskie, J.V., Sealy, S.G. & Hobson, K.A. (1992). Behavioral defenses against avian brood
434   parasitism in sympatric and allopatric host populations. *Evolution*, 46, 334–340.
435   Brooks, M.E., Kristensen, K., Benthem, K.J. van, Magnusson, A., Berg, C.W., Nielsen, A., *et al.*
436   (2017). glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated
437   Generalized Linear Mixed Modeling. *The R Journal*, 9, 378–400.
438   Campobello, D. & Sealy, S.G. (2010). Enemy Recognition of Reed Warblers (*Acrocephalus
439   scirpaceus*): Threats and Reproductive Value Act Independently in Nest Defence
440   Modulation. *Ethology*, 116, 498–508.
441   Campobello, D. & Sealy, S.G. (2011). Use of social over personal information enhances nest
442   defense against avian brood parasitism. *Behavioral Ecology*, 22, 422–428.
443   Candolin, U. & Wong, B. (2015). *Behavioural Responses to a Changing World: Mechanisms &
444   Consequences*. Oxford University Press.
445   Cantor, M., Maldonado-Chaparro, A.A., Beck, K.B., Brandl, H.B., Carter, G.G., He, P., *et al.* (2021).
446   The importance of individual-to-society feedbacks in animal ecology and evolution.
447   *Journal of Animal Ecology*, 90, 27–44.
448   Ceresa, F., Belda, E.J., Kvist, L., Rguibi-Idrissi, H. & Monrós, J.S. (2015). Does fragmentation of
449   wetlands affect gene flow in sympatric *Acrocephalus* warblers with different migration
450   strategies? *Journal of Avian Biology*, 46, 577–588.
451   Chaumont, M.H.J., Langmore, N.E. & Welbergen, J.A. (2021). The ghosts of parasitism past:
452   lingering frontline anti-brood parasite defenses in a former host. *Current Zoology*, 67,
453   573–583.
454   Chen, X., Cai, Y., Wang, J. & Yang, C. (2022). Use of 3D-printed animal models as a standard
455   method to test avian behavioral responses toward nest intruders in the studies of avian
456   brood parasitism. *Avian Research*, 13, 100061.
457   Cramp, S. & Brooks, D. (1992). *Handbook of the birds of Europe, the Middle East and North
458   Africa. The birds of the western Palearctic, vol. VI. Warblers*. Oxford: Oxford university
459   press.

Crane, A.L. & Ferrari, M.C. (2013). Social learning of predation risk: a review and prospectus.
Social learning theory: phylogenetic considerations across animal, plant, and microbial
taxa, 53–82.
Dall, S.R.X., Giraldeau, L.-A., Olsson, O., McNamara, J.M. & Stephens, D.W. (2005). Information
and its use by animals in evolutionary ecology. *Trends in Ecology & Evolution*, 20, 187–
193.
Davies, N.B. & Brooke, M. de L. (1988). Cuckoos versus reed warblers: Adaptations and
counteradaptations. *Animal Behaviour*, 36, 262–284.
Davies, N.B. & Brooke, M.D.L. (1989). An Experimental Study of Co-Evolution Between the
Cuckoo, *Cuculus canorus*, and its Hosts. I. Host Egg Discrimination. *Journal of Animal*
Ecology, 58, 207–224.
Davies, N.B. & Welbergen, J.A. (2009). Social Transmission of a Host Defense Against Cuckoo
Parasitism. *Science*.
Feeney, W.E. & Langmore, N.E. (2013). Social learning of a brood parasite by its host. *Biology*
Letters, 9, 20130443.
Feeney, W.E., Troscianko, J., Langmore, N.E. & Spottiswoode, C.N. (2015). Evidence for
aggressive mimicry in an adult brood parasitic bird, and generalized defences in its host.
Proceedings of the Royal Society B: Biological Sciences, 282, 20150795.
Feeney, W.E., Welbergen, J.A. & Langmore, N.E. (2012). The frontline of avian brood parasite–
host coevolution. *Animal Behaviour*, 84, 3–12.
Foster, S.A. (2013). Evolution of behavioural phenotypes: influences of ancestry and expression.
Animal Behaviour, Special Issue: Behavioural Plasticity and Evolution, 85, 1061–1075.
Gil, M.A., Baskett, M.L. & Schreiber, S.J. (2019). Social information drives ecological outcomes
among competing species. *Ecology*, 100, e02835.
Gil, M.A., Hein, A.M., Spiegel, O., Baskett, M.L. & Sih, A. (2018). Social Information Links
Individual Behavior to Population and Community Dynamics. *Trends in Ecology &*
Evolution, 33, 535–548.
Gilman, S.E., Urban, M.C., Tewksbury, J., Gilchrist, G.W. & Holt, R.D. (2010). A framework for
community interactions under climate change. *Trends in Ecology & Evolution*, 25, 325–
331.
Grim, T. & Stokke, B.G. (2016). In the Light of Introduction: Importance of Introduced
Populations for the Study of Brood Parasite–Host Coevolution. In: *Biological invasions*
and animal behaviour. Cambridge University Press, pp. 133–157.
de Groot, C., Wijnhorst, R.E., Ratz, T., Murray, M., Araya-Ajoy, Y.G., Wright, J., *et al.* (2023). The
importance of distinguishing individual differences in 'social impact' versus 'social
responsiveness' when quantifying indirect genetic effects on the evolution of social
plasticity. *Neuroscience & Biobehavioral Reviews*, 144, 104996.
Hale, K. & V. Briskie, J. (2007). Response of introduced European birds in New Zealand to
experimental brood parasitism. *Journal of Avian Biology*, 38, 198–204.
Hämäläinen, L., M. Rowland, H., Mappes, J. & Thorogood, R. (2022). Social information use by
predators: expanding the information ecology of prey defences. *Oikos*, 2022, e08743.
Hartig, F. (2022). *DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)*
Regression Models.
Hauber, M.E., Moskát, C. & Bán, M. (2006). Experimental shift in hosts' acceptance threshold of
inaccurate-mimic brood parasite eggs. *Biology Letters*, 2, 177–180.

Järvinen, O. & Ulfstrand, S. (1980). Species turnover of a continental bird fauna: Northern
Europe, 1850–1970. *Oecologia*, 46, 186–195.
Jolly, C.J., Webb, J.K. & Phillips, B.L. (2018). The perils of paradise: an endangered species
conserved on an island loses antipredator behaviours within 13 generations. *Biology*
Letters, 14, 20180222.
Kelley, J.L. & Magurran, A.E. (2003). Effects of relaxed predation pressure on visual predator
recognition in the guppy. *Behav Ecol Sociobiol*, 54, 225–232.
Kuehn, M.J., Peer, B.D., McCleery, R.A. & Rothstein, S.I. (2016). Yellow warbler defenses are
retained in the absence of brood parasitism but enhanced by experience with cowbirds.
Behavioral Ecology, 27, 279–286.
Lahti, D.C., Johnson, N.A., Ajie, B.C., Otto, S.P., Hendry, A.P., Blumstein, D.T., *et al.* (2009).
Relaxed selection in the wild. *Trends in Ecology & Evolution*, 24, 487–496.
Langmore, N.E., Feeney, W.E., Crowe-Riddell, J., Luan, H., Louwrens, K.M. & Cockburn, A. (2012).
Learned recognition of brood parasitic cuckoos in the superb fairy-wren Malurus
cyaneus. *Behavioral Ecology*, 23, 798–805.
Lawson, S.L., Enos, J.K., Fernandez-Duque, F., Kleindorfer, S., Ward, M.P., Gill, S.A., *et al.* (2023).
Absence of Referential Alarm Calls in Long-term Allopatry from the Referent: A Case
Study with Galapagos Yellow Warblers. *Behav Ecol Sociobiol*, 77, 99.
Lawson, S.L., Leuschner, N., Enos, J.K. & Hauber, M.E. (2020). Loss of graded enemy recognition
in a Whitehead population allopatric with brood-parasitic Long-tailed Cuckoos.
Avocetta, 44, 4–10.
Lindholm, A. (2000). Tests of phenotypic plasticity in reed warbler defences against cuckoo
parasitism. *Behaviour*, 137, 43–60.
Lindholm, A.K. (1999). Brood parasitism by the cuckoo on patchy reed warbler populations in
Britain. *Journal of Animal Ecology*, 68, 293–309.
Lindholm, A.K. & Thomas, R.J. (2000). Differences between Populations of Reed Warblers in
Defences against Brood Parasitism. *Behaviour*, 137, 25–42.
Marton, A., Fülöp, A., Ozogány, K., Moskát, C. & Bán, M. (2019). Host alarm calls attract the
unwanted attention of the brood parasitic common cuckoo. *Scientific Reports*, 9, 18563.
Mineka, S., Davidson, M., Cook, M. & Keir, R. (1984). Observational conditioning of snake fear in
rhesus monkeys. *Journal of Abnormal Psychology*, 93, 355–372.
Moskát, C., Szentpéteri, J & Barta, Z. (2002). Adaptations by Great Reed Warblers to Brood
Parasitism: A Comparison of Populations in Sympatry and Allopatry with the Common
Cuckoo. *Behaviour*, 139, 1313–1329.
Nakamura, H. (1991). Brood Parasitism by the Cuckoo *Cuculus canorus* in Japan and the Start of
New Parasitism on the Azure-winged Magpie *Cyanopica cyana*. *Japanese Journal of*
Ornithology, 39, 1–18.
Nuismer, S.L., Thompson, J.N. & Gomulkiewicz, R. (2003). Coevolution between hosts and
parasites with partially overlapping geographic ranges. *Journal of Evolutionary Biology*,
16, 1337–1345.
O'Connor, M.I., Pennell, M.W., Altermatt, F., Matthews, B., Melián, C.J. & Gonzalez, A. (2019).
Principles of Ecology Revisited: Integrating Information and Ecological Theories for a
More Unified Science. *Frontiers in Ecology and Evolution*, 7.
Oien, I.J., Honza, M., Moksnes, A. & Roskaft, E. (1996). The Risk of Parasitism in Relation to the
Distance from Reed Warbler Nests to Cuckoo Perches. *Journal of Animal Ecology*, 65,
147–153.

R Core Team. (2023). *R: A Language and Environment for Statistical Computing*. R Foundation
for Statistical Computing, Vienna, Austria.
Rothstein, S. (2001). Relic behaviours, coevolution and the retention versus loss of host
defences after episodes of avian brood parasitism. *Animal behaviour*, 61, 95–107.
Rowell, T.A.A.D., Magrath, M.J.L. & Magrath, R.D. (2020). Predator-awareness training in
terrestrial vertebrates: Progress, problems and possibilities. *Biological Conservation*,
252, 108740.
Scheiner, S.M. & Levis, N.A. (2021). The Loss of Phenotypic Plasticity Via Natural Selection:
Genetic Assimilation. In: *Phenotypic Plasticity & Evolution*. CRC Press.
Smith, W.J., Rönkä, K., Bensch, S., Bergman, N.M., Dudaniec, R.Y., Eroukhmanoff, F., *et al.*
(2025). Putative signatures of genomic adaptation to Cuckoo brood parasitism in Reed
Warbler hosts across Europe.
Soler, M. (2014). Long-term coevolution between avian brood parasites and their hosts. *Biol Rev
Camb Philos Soc*, 89, 688–704.
Soler, M. & Møller, A.P. (1990). Duration of sympatry and coevolution between the great
spotted cuckoo and its magpie host. *Nature*, 343, 748–750.
Stokke, B.G., Takasu, F., Moksnes, A. & Røskaft, E. (2007). The importance of clutch
characteristics and learning for antiparasite adaptations in hosts of avian brood
parasites. *Evolution*, 61, 2212–2228.
Thompson, J.N. (2005). Coevolution: The Geographic Mosaic of Coevolutionary Arms Races.
Current Biology, 15, R992–R994.
Thorogood, R. & Davies, N.B. (2012). Cuckoos Combat Socially Transmitted Defenses of Reed
Warbler Hosts with a Plumage Polymorphism. *Science*, 337, 578–580.
Thorogood, R. & Davies, N.B. (2013). Reed Warbler Hosts Fine-Tune Their Defenses to Track
Three Decades of Cuckoo Decline. *Evolution*, 67, 3545–3555.
Thorogood, R. & Davies, N.B. (2016). Combining personal with social information facilitates host
defences and explains why cuckoos should be secretive. *Sci Rep*, 6, 19872.
Thorogood, R., Kokko, H. & Mappes, J. (2018). Social transmission of avoidance among predators
facilitates the spread of novel prey. *Nat Ecol Evol*, 2, 254–261.
Tolman, D., Campobello, D., Rönkä, K., Kluen, E. & Thorogood, R. (2021). Reed Warbler Hosts Do
Not Fine-Tune Mobbing Defenses During the Breeding Season, Even When Cuckoos Are
Rare. *Frontiers in Ecology and Evolution*, 9, 547.
Welbergen, J.A. & Davies, N.B. (2008). Reed warblers discriminate cuckoos from sparrowhawks
with graded alarm signals that attract mates and neighbours. *Animal Behaviour*, 76,
811–822.
Welbergen, J.A. & Davies, N.B. (2009). Strategic Variation in Mobbing as a Front Line of Defense
against Brood Parasitism. *Current Biology*, 19, 235–240.
Welbergen, J.A. & Davies, N.B. (2011). A parasite in wolf's clothing: hawk mimicry reduces
mobbing of cuckoos by hosts. *Behavioral Ecology*, 22, 574–579.
Welbergen, J.A. & Davies, N.B. (2012). Direct and indirect assessment of parasitism risk by a
cuckoo host. *Behavioral Ecology*, 23, 783–789.
Whiten, A. (2021). The burgeoning reach of animal culture. *Science*, 372, eabe6514.
Yang, C., Huang, J., Liang, W. & Møller, A.P. (2021). Absence of anti-parasitic defenses in an
Asian population of the magpie, a regular host of the great spotted cuckoo in Europe.
Current Zoology, 67, 345–347.

Yang, C., Wang, L., Cheng, S.-J., Hsu, Y.-C., Stokke, B.G., Røskaft, E., *et al.* (2015). Deficiency in
egg rejection in a host species as a response to the absence of brood parasitism.
Behavioral Ecology, 26, 406–415.

Figures and Tables

Figure 1. Reed warbler mobbing defences (calls per 5 minutes) in allopatry (Finland: A-C, Sicily: D-F) before (grey dots) and after (coloured dots) experimental manipulation of social information

605   ('SI') about cuckoos (B, E) versus a control (C – magpie, F – teal). Details of experimental design

606   are shown in A, presentation order ('P') of models was alternated in D.

608
609 **Figure 2.** (A) Propensity to mob cuckoos did not change after social information about cuckoos or
610 a control in Finland or Sicily. (B) Mobbing pairs showed social responsiveness (i.e. increased
611 intensity) to cuckoos, but only in Finland (distributions of changes in mobbing intensity, vertical
612 lines = medians).
613

Figure 3. Rejection rates of experimental eggs in Finland (inset photos). Blue non-mimetic eggs (left) were rejected at the same rate regardless of whether social information was provided about cuckoos (24%) or magpies (22%). Mimetic brown-spotted eggs (right) were not rejected.

Figure 4. (A) Latency of birds to approach presentations of models at their nest (indicated by icons) before and after social information (dashed line) about cuckoos (yellow) or a magpie control (blue) in Finland. Raw data are plotted as diamonds, with darker symbols indicating pairs that mobbed cuckoos. Predicted mean values (± standard error) from a GLMM are plotted as circles and bars, with connecting lines indicating change across model presentation order (P1-P3). (B) Latency of birds to approach presentations of models at their nest according to whether they mobbed a cuckoo. Predicted mean values (± standard error) from a GLMM are plotted as circles and bars, with connecting lines indicating change across model presentation order (P1-P3).