Temporal and spatial dynamics within the fungal microbiome of grape fermentation

Cristobal A. Onetto[1,2], Chris M. Ward[1], Steven Van Den Heuvel[1], Laura Hale[1], Kathleen Cuijvers[1] and Anthony R. Borneman[1,2*]

[1] The Australian Wine Research Institute, Glen Osmond, South Australia 5064, Australia.

[2] School of Wine, Food and Agriculture, The University of Adelaide, South Australia, 5005. Australia

*Corresponding author: anthony.borneman@awri.com.au

Abstract

Wine fermentation is a highly complex and competitive environment, imposing harsh selective pressures on fungal community ecology and diversity. The composition of fungal communities inhabiting the surface of grapes will directly impact fermentation progression, wine quality, and contribute to the distinctiveness between wines from different geographical regions. Despite this, the extent of microbial community diversity between geographies, termed 'microbial terroir,' remains highly debated. We amassed a large survey of grape spontaneous ferments over six years, encompassing 3105 fungal microbiomes across 14 geographically separated grape-growing regions, and nine grape cultivars. Investigation into the biodiversity of these ferments identified that few high abundance genera form the core of the initial grape microbiome. In line with previous studies, various consistent taxa were linked to specific geographical locations and grape varieties. However, these taxa accounted for a small portion of the overall diversity in the dataset. Through unsupervised clustering, we identified three distinct community types in the grape fungal microbiome, each exhibiting variations in the abundance of key genera. Analysing ferments across temporal and spatial scales revealed significant differences in species richness and compositional heterogeneity between wineries and grape growing regions. However, microbial communities were transient between years in the same winery, regularly transitioning between the three broad community types. We then investigated microbial community composition throughout the fermentative process and observed that initial microbial community composition is predictive of the diversity during the early stages of fermentation, with *Hanseniaspora uvarum* detected as the main non-*Saccharomyces* species within this large cohort of samples. Our results help to formulate a clear understanding of the spatial and temporal characteristics of the grape juice fungal microbiome and suggest that these communities are mainly defined by the grape niche and in a minor way shaped by local environmental conditions.

Keywords: grape fungal microbiome; spontaneous fermentation; non-Saccharomyces; metabarcoding

Introduction

The surfaces of vines and grapes harbor a naturally occurring, intricate consortium of fungal and bacterial species, which exert a direct influence on vine growth and overall health [1-3]. These communities can vary significantly in their composition and may be affected by numerous factors, including grape variety, geographical location, climate and viticultural practices [4-6].

These microorganisms are transferred into the grape must during harvesting and processing, where they can play a pivotal role in shaping fermentation dynamics and contributing to the sensory

attributes of the resulting wine. This influence is particularly significant during traditional uninoculated ("wild") fermentation, where the absence of commercial yeast strains allows specific members of the microbial community to directly impact the fermentation process. Under this production framework, a complex microbial succession of yeasts and yeast-like fungi is generally observed as the grape must ferments. Initially, a diverse mixture of aerobic and apiculate yeasts, which normally reside on the surface of intact grape berries, dominate the fresh grape must. However, most of these species succumb to the decreasing oxygen levels and increasing ethanol concentrations, allowing mildly fermentative species to transiently increase in abundance [7-11]. Ultimately however, due to a higher fermentative ability, growth rate and tolerance to ethanol, most wild wine ferments converge to near monocultures of the major wine yeast, *Saccharomyces cerevisiae*. This is despite almost undetectable levels of this species being present on intact grape berries [12, 13].

While traditional microbiological techniques have provided important insights into the microbial succession that occurs in spontaneous ferments, both the breadth of ferments investigated and the depth at which individual species contributions could be resolved has been limited. Recent advances in culture-independent methods for species analysis, such as shotgun metagenomics and amplicon-based ITS phylotyping provides a high-throughput means to analyse large numbers of microbiological samples in great detail [14, 15].

Phylotyping techniques have been employed to investigate grape and fermentation microbiomes, aiming to uncover whether regional variations in wines can be partially attributed to the distinctiveness of microbial communities, supporting the notion of a "microbiome terroir". While studies have revealed geographic patterns and considerable variation in microbial community composition among specific locations [4, 5, 16, 17], the extent to which these biogeographic patterns persist on a broader ecological scale remains uncertain. To address this, an extensive survey of grape juice samples was carried out to assess the microbial diversity inherent across a large cohort of uninoculated fermentations within Australia, that extended across nine grape varieties and 14 wine regions over the six successive vintages from 2016 to 2021. This comprehensive dataset enabled a thorough exploration of the core microbial communities in grape must and fermentation at a large ecological scale, thereby enhancing our understanding of the fungal species associated with grapes and the fermentation process.

Materials and Methods
Sample collection

A total of 36 wineries were sampled during six years across 14 different wine growing regions covering six states of Australia (Fig. 1a, Table S1). Wineries were supplied with pre-labelled, 50 mL falcon tubes. Approximately 40 mL of grape must was collected at four timepoints per fermentation tank, based upon refractometry levels (T1, freshly pressed grape juice; T2, after a 1 Bé drop in sugar; T3, 50% sugar remaining; T4, 10% sugar remaining) (Fig. 1b). Samples were stored frozen until collection and processing. All samples were obtained from strictly un-inoculated spontaneous ferments.

Sample processing and ITS sequencing

Samples were thawed, homogenized, and 1.9 mL of sample was centrifuged and washed twice with 1 mL of 1X PBS. DNA was extracted from pellets using the DNeasy PowerFood Microbial DNA Isolation Kit (Qiagen) following the manufacturer's instructions. Bead-beating was carried out using a

combination of 0.1 mm and 0.5 mm zirconia/silica beads (BioSpec Products) in a Precellys Evolution
homogenizer (Bertin Instruments) at 8,000 RPM for 4 x 60 sec.

A two-step PCR was performed using 1 ng of DNA with primer sequences designed to amplify the
fungal ITS region, while adding experiment-specific inline barcodes and appropriate adaptors for the
Illumina sequencing platform as previously described by Sternes, Lee [14]. The first-round of
amplification (20–30 cycles, 55℃ annealing, 30-second extension, KAPA 2G Robust polymerase)
targeting the ITS region was performed using primers BITS (ACCTGCGGARGGATCA) and B58S3
(GAGATCCRTTGYTRAAAGTT) [18]. 2 uL of first-round PCR product was used in the second
amplification (15 cycles, 55°C annealing, 30-second extension, KAPA 2G Robust polymerase).
Sequencing was performed using 2 x 300 bp paired-end chemistry in an Illumina instrument at the
Ramaciotti Centre for Genomics (UNSW Sydney, Australia).

Sequencing data processing

Raw sequencing reads were quality trimmed using Trimmomatic v0.38 [19], adaptor trimmed using
Cutadapt v.1.16 [20] and merged using FLASH v2.2.00 [21]. Merged reads were hashed, de-replicated
and counted as previously described by Sternes, Lee [14]. Operational taxonomic unit (OTU)
clustering was performed using Swarm v2.2.2 [22]. Taxonomic classification was performed with the
assign_taxonomy.py module of QIIME v1.9.1 [23] with a 98 % similarity cut off against the UNITE
database (qiime_ver8_dynamic_10.05.2021). The taxonomic classification of all OTUs was also
manually checked and curated against the BLAST database.

Diversity analysis

All diversity analyses were performed in R v4.1.1 [24] and plots produced using a combination of
ggplot2 [25] and the Ampvis2 package v2.8.3 [26]. Samples containing less than 2000 total read
counts were removed from the dataset prior to analysis. Alpha (Shannon index) and beta diversity
(Bray-Curtis dissimilarities) were calculated on rarefied OTU counts using Vegan v2.6 [27]. Differences
of alpha diversity between categorical variables were investigated using Dunn's multiple comparison
test and Holm adjusted P-values. Investigation of differences in species diversity between categorical
variables was tested using a permutational multivariate analysis of variance (MANOVA) [28]
implemented in the adonis function with 999 permutations. Multivariate homogeneity of groups
dispersions between categories was investigated using the betadisper function of Vegan and
statistical differences between groups within categories were tested using a permutational ANOVA
with 999 permutations. Pair-wise comparison between samples was performed with the permutest
function implemented in Vegan. PERMANOVA is known to behave unreliably in unbalanced designs
with heterogenous dispersion [29]. To address this issue the F_2 statistic was [30] also used to test
categories that showed heterogeneity in group dispersion
(https://github.com/AWRI/permanovaModF). Investigation of specific taxonomic differences
between categories was tested using the LDA effect size [31] on a normalised abundance table at
genus level. Random forest supervised classification models [32] were used to evaluate the
discriminative power of fungal with an LDA effect size >2. The strength of each predictor was
evaluated using the mean decrease in Gini coefficient (MDG) as implemented in the R package
Microeco [33], using 10000 trees per category with the randomForest package v.4.7 [34]. Community
partitioning of T1 samples was performed using Dirichlet multinomial mixture (DMM) modelling [35]
on a rarefied OTU abundance level matrix. The number of groups was obtained using the lowest
Laplace estimation score. Microbiome branch structure was performed on the genus clustered
relative abundance table using the PHATE algorithm v. 1.0.7 [36] with default parameters.

Data availability

The sequencing data included in this study is available in NCBI under BioProject PRJNA1053885.

Results

The core fungal community of grapes is represented by few highly abundant species

To determine the core community of fungi associated with grapes and fermentation a large-scale
survey of grape juice was conducted that covered 14 wine grape-growing regions of Australia (Fig.
1a). Samples were obtained from uninoculated grape juice samples across four timepoints during
fermentation (Fig. 1b) from a total of 36 wineries throughout a period of six years (Fig. 1c),
representing the largest microbial survey of grape juice to date. The fungal microbial community
composition was characterized using amplicon sequencing targeting the ITS region. After quality
filtering and OTU clustering, a total of 3105 samples were obtained containing 4309 OTUs.

T1 samples, which were taken as soon as fermentation tanks were filled with freshly crushed juice,
largely reflected the composition of the fungal communities that inhabited the surface and flesh of
grapes and were selected to estimate the grape-derived core fungal microbiome. While the T1
timepoint was shown to contain the highest diversity across the 772 samples, a small number of very
abundant OTUs dominated the overall read abundance within most samples with the top 10 and 20
most abundant OTUs represented 70% (± 3.4%) and 80% (± 2.7%) of the total reads on average,
respectively (Fig. 2a). OTUs that made up at least 80% of the reads in at least one sample were
defined as abundant (Fig. 2b). To be considered a core OTU, we defined an abundance-occurrence
threshold of 50% of the samples. While no single OTU was observed as abundant in all the samples
investigated (Fig. 2b), two OTUs, one belonging to the genus *Aureobasidium* (observed in all samples,
abundant in 659) and the other *Cladosporium* (observed in 770 samples, abundant in 424), passed
the applied threshold for comprising core members of grape fungal microbiome (Fig. 2b). Species
within the genus *Aureobasidium* are recognized for their ubiquitous presence in the phyllosphere of
various plants, including grapevines [37] and these results confirm *Aureobasidium* as the
predominant fungal genus across this large cohort of grape must samples (Fig. 2b and c). OTUs
taxonomically assigned to *Cladosporium ramotenellum* (757 samples, abundant in 281),
Saccharomyces cerevisiae (690 samples, abundant in 204) and *Hanseniaspora uvarum* (689 samples,
abundant in 190) were also abundant across a large number of samples (Fig. 2b). When considering
only the occurrence of individual OTUs, 54 OTUs were observed in at least 50% of the samples, 25
OTUs in 80% (Fig. 2c) and only a single OTU (genus *Aureobasidium*) in 100% of the samples analysed.

A global overview of all samples clustered by genus showed similar profiles to the OTU analysis and is
consistent with previous studies [4, 6, 14, 38, 39]. The most abundant genera corresponded to
Aureobasidium, *Cladosporium*, *Saccharomyces*, *Hanseniaspora* and *Metschnikowia* (Fig. 2c). As
opposed to *Aureobasidium*, which comprised only four OTUs and with a single OTU accounting for
most of the abundance (Fig. 2b), the genera *Cladosporium* (13 OTUs), *Saccharomyces* (67 OTUs),
Hanseniaspora (46 OTUs) and *Metschnikowia* (53 OTUs) were comprised of high numbers of
individual OTUs and suggests a higher species diversity within these genera.

Temporal stability and regional variations in fungal communities

To investigate changes in fungal diversity over the course of the fermentation process, three
sequential samples were collected after the T1 sampling point, as delineated by sugar consumption

(Fig. 1b). The fermenting grape juice ecosystem exhibits a pronounced selective bias, favouring
species with fermentative capabilities and increased tolerance to ethanol [40, 41]. Alpha and beta-
diversity assessments across fermentation time points revealed clear reduction in both diversity and
beta dispersion (Fig. 3a and Fig. S1a), accompanied with rapid shifts in microbial community
composition (Fig. 3b) towards Saccharomycetes yeast species (Fig. S1b). In alignment with findings
from many previous studies, 94% of T4 samples were predominantly dominated (> 50% relative
abundance) by OTUs classified as *Saccharomyces cerevisiae*. Additionally, *Hanseniaspora uvarum*,
Schizozaccharomyces japonicus and *Metschnikowia* were observed as the most abundant non-
Saccharomyces at T4.

Aside from *S. cerevisiae*, *H. uvarum* was observed as the most abundant yeast during the initiation of
fermentation (1 Be drop), with 333 of 781 T2 samples containing greater than 10% relative
abundance of *H. uvarum*. Additional genera of non-*Saccharomyces* species were also observed in
high relative abundance (>10%) at the initiation of fermentation, albeit in a lower number of samples
(*Aureobasidium*: 159, *Metschnikowia*: 77, *Cladosporium*: 50, *Pichia*: 29, *Issatchenkia*: 24, *Candida*:
18, *Starmerella*: 16, *Penicillium*: 15, *Torulaspora*: 14, *Schizosaccharomyces*: 7, *Botrytis*: 4 and
Lachancea: 3 of 783 samples).

Significant differences in species richness (Dunn's, Holm-adjusted $p<0.05$) and compositional
heterogeneity (Permutest of beta dispersion $p<0.05$) were observed within T1 samples across various
wineries and grape-growing regions (Fig. S2-S3, Tables S2-S5). Additionally, species richness and
diversity exhibited substantial dispersion, indicating instability of the microbial communities (Fig. S2).
To assess the temporal microbial stability, a comparative analysis of beta diversity was conducted
across a panel of 16 wineries that were repeatedly sampled over three successive vintages (Fig. 3e).
While a small number of wineries displayed a consistent microbial composition over time, the
majority exhibited compositional alterations either after one or two years (Fig. 3e).

Previous studies have suggested that viticultural regions harbour regionally distinct microbial
communities [4, 5, 16, 17]. To investigate this effect, a Permutational MANOVA of the Bray-Curtis
dissimilarities between categories was utilized to assess fungal community structure between
wineries, grape-growing regions, states, grape varieties, and years across the dataset (Fig. 3c). All
tests were significant (Table S6), however a meaningful effect-size was only observed when
comparing between wineries (Adonis R^2 of wineries = 0.29) (Fig. 3c). To account for the observed
heterogenous dispersion between categories, a modified F_2 statistic was also calculated [30],
producing similar results (Table S6). Principal Coordinates Analysis (PCoA) revealed no discernible
clustering across the various categories examined (Fig. S4). Subsequent pairwise comparisons across
individual wineries identified that the most pronounced dissimilarities were primarily associated
with three specific wineries, namely K, E, and L (Table S7), situated within the Margaret River,
Langhorne Creek, and Canberra District wine regions. Principal Component Analysis (PCA) analysis
between these wineries indicated that they largely differ in the abundance of three OTUs
corresponding to the genera *Aureobasidium*, *Cladosporium* and *Saccharomyces* (Fig. 3d).

To investigate significant associations between specific fungal genera and categories, a differential
abundance testing method (LEfSE) combined with random forest models was implemented to
evaluate the discriminative power of the differentially abundant genera. In agreement with previous
studies [4, 5, 16, 17], significant associations (Log LDA score >2) were observed across all categories
(Table S8 – S17). When focusing on broad geographical patterns, several genera were associated with
specific states of Australia (Fig. 4a) with the largest MDG values observed for the genera
Cystofilobasidium (SA), *Botrytis* (TAS) and *Sporobolomyces* (SA). Grape varieties also showed an

influence on fungal microbiota (Fig. 4b) with *Dipodia* (Shiraz), *Seimatosporium* (Grenache) and
Cladosporium (Chardonnay) amongst the top explaining taxa.

220 **Three community types partition the grape fungal microbiome**

The large dataset provided the opportunity to investigate population stratification within the grape
fungal microbiome and uncover microbial compositional patterns that would not be easily observed
when focusing on individual taxa. To achieve this, a DMM-based approach [35], which has previously
shown recurrent compositional patterns in multiple studies of the gut microbiome, was applied to
this dataset [42].

Analysis of the 772 T1 samples indicated that the grape juice microbiomes could be partitioned into
three distinct community types (defined as DMM1 - 3) (*n* DMM1 = 430, *n* DMM2 = 250 and *n* DMM3
= 92) that displayed clear ordination (Fig. 5a and b). Inspection of the 30 most abundant genera, in
addition to the LefSE and random forest models, revealed clear differences between communities
(Figure 5a, Table S18 – S19). DMM1 was defined by a high abundance of *Aureobasidium*,
Cladosporium, *Metschnikowia* and several lower abundance genera (Fig. 5a). DMM2 displayed the
highest abundance of *Aureobasidium* and the highest species richness (Fig. 5c), including the
presence of several rare genera (Fig. 5a). DMM3 had the lowest species richness (Fig. 5c) and was
mainly characterized by a high abundance of *Saccharomyces* and *Hanseniaspora* (Fig. 5a).
Assignment of winery samples and wine growing regions to community types indicated that many
were composed of mixed fungal community types (Fig. 5d).

It is possible that these three community types represent stages of ecological progression that are
affected by external factors (e.g. grape maturity at sample time). To investigate this, the Potential of
Heat-diffusion for Affinity-based Trajectory Embedding (PHATE) algorithm [36], a method developed
to discover transitions in high dimensional data through the visualisation of branches and clusters
was applied to the dataset. PHATE analysis showed three main branches (i.e. successions of
microbiome arrangements bridging localized ecological conditions) that were associated with the
abundance of the main genera discriminating between community types (Fig. 5e), suggesting
transitions of microbial composition between these communities.

To evaluate the impact of initial community type on the microbial dynamics of fermentation, results
from T2 through T4 were partitioned based on the community type assignment of their
corresponding T1 sample. DMM1 and DMM2 retained an overall higher species richness at T2 (Fig.
S1c) compared to DMM3. However, in all community types, fermentative species such as
Saccharomyces and *Hanseniaspora* were already abundant at T2 (Fig. 5f). Non-fermentative taxa,
namely *Aureobasidium* and *Cladosporium*, appeared to be more persistent in community types that
initially contained a higher abundance of these genera in T1 (Fig. 5A). While this suggests that the
initial community type directly influences the microbial dynamics of fermentation, it is evident that
other external factors, such as grape juice composition and fermentation conditions, likely exert a
substantial influence on microbial community dynamics.

255 Discussion

The fungal communities of grapes have a direct effect on both the fermentation dynamics and the
sensory attributes of spontaneously fermented wines [16, 43, 44]. To elucidate if the regional
variation of wines could be partially explained by the distinctiveness of microbial communities,
several studies have investigated the grape microbiota across different geographical scales using

high-throughput amplicon sequencing, supporting the notion of a "microbiome terroir" [4, 5, 16, 17].
While it is clear from these studies that there can be extensive variation in microbial communities
across geographical locations, they lack the sampling depth to obtain a global overview of the fungal
communities of grape must. To address this, we carried out an extensive survey of 3105 grape juice
samples across 14 grape-growing regions throughout a period of six years (Fig. 1), representing the
most comprehensive microbial survey of grapes to date. Rather than concentrating on small scale
regional or year-specific differences we analysed the dataset as a whole to elucidate general trends
in community composition.

The core fungal community of grapes was surprisingly small, comprising only two OTUs and in
addition, fewer than 20 OTUs accounted for the majority of the sequencing reads across the 772 T1
samples analysed. Relative to other open ecosystems, where the assembly of the communities is
initiated by a multitude of species, grapes appear to display a very low number of core OTUs [45-49]
and resemble the fungal microbiome of other cultivated fruit species [50]. Grapes therefore
represent a highly selective ecosystem that differs significantly from other regions of the grapevine,
such as the rhizosphere or bark, where a higher fungal taxonomic diversity have been observed [39,
51]. *Aureobasidium* appears to be particularly well adapted to the grape environment, with a single
OTU dominating the abundance associated to this genus. While a single species could not be
confidently assigned to this OTU, it is highly probable that it corresponds to *Aureobasidium pullulans*,
given its more frequent isolation from grapes [52-55]. It is also interesting to note that community
types featuring a higher abundance of this genus exhibit an overall greater species richness,
suggesting that *Aureobasidium* species may not engage in broad antagonism but rather exhibit
targeted antagonism against specific species, as reported in several studies [56, 57]. *Cladosporium*,
known for causing post-harvest rot diseases in various fruits [58-60], is also part of the core
microbiome of grapes, implying that its presence in grapes is not necessarily linked to disease
development as previously suggested [39].

The fungal dynamics of fermentation progressed as expected [7, 8] with *S. cerevisiae* dominating by
T2 in nearly all the samples analysed, yet many samples exhibited a prolonged period of high
abundance of *Hanseniaspora uvarum*. Although prior research noted the prevalence of this species
in early fermentation stages [61], these results highlight its role as the primary non-*Saccharomyces*
yeast in grape juice fermentation. While other *Hanseniaspora* and non-*Saccharomyces* yeast were
also detected at T1, their frequency was marginal by T2. From an adaptive viewpoint, *H. uvarum*
seems less genetically equipped among *Hanseniaspora* species for fermentation due to a notable
loss of genes related to cell-cycle and DNA repair [62]. Yet recent studies suggest that losing specific
genes, including key cell-cycle regulators, might be an adaptive strategy, enabling a quicker cell cycle
[63]. This discovery prompts the need for more research on *H. uvarum* to fully uncover its
competitive advantage during early fermentation, especially compared to the various non-
Saccharomyces yeast species found in grape must.

Wineries displayed distinct and temporally unstable fungal communities, and this is consistent with
the observations reported in similar studies [4, 17, 39, 64, 65]. Despite this, the effect size of this
associations was considerably lower than previously reported, likely due to differences in sampling
size, geographic coverage, number of grape varieties and years of sampling. Large-scale geographic
denominations exhibited significant associations with several fungal genera, however, the strongest
variations between states and wineries were explained by species not considered to participate in
grape fermentation and might reflect adaptations of the fungal microbiome to local environmental
conditions, as previously suggested [4]. An example of this is the genus *Botrytis*, which was the
second most important feature distinguishing the samples obtained from Tasmania, a geographical

region known for having a colder climate and higher average rainfall, conditions that stimulate the
growth of *Botrytis* species [66]. Several taxa distinguishing broader geographical regions coincide
with microbiome reports from grape-growing areas in North America [4] and Europe [6]. Some of
these genera include *Botrytis*, *Rhodotorula*, *Cladosporium*, *Aureobasidium*, and *Penicillium*. The
congruence across studies provides support to the proposition by Bokulich [4] that biogeographical
trends in different wine regions reflect responses to local conditions rather than patterns of microbial
dispersion. Our results also suggests that grape variety plays a role in defining the abundance of
select genera. Surprisingly, *Cladosporium* was observed as the top genus distinguishing the variety
Chardonnay, which has been previously reported for grapes in North America [4], suggesting that
either a genetic component or a standardised vineyard management approach associated with this
variety drives the observed global pattern of microbial abundance. Variations in susceptibility to
grapevine pathogens between varieties may also play a major role in explaining these differences.
For example, *Diplodia* showed the highest MDG value for the variety Shiraz, which is known to be
highly susceptible to the colonisation of the grapevine trunk pathogen *Diplodia serata* [67]. While
our results support the proposition of a "microbial terroir", the overlapping results with previous
studies performed in different geographical regions suggest that global viticultural zones share a
similar grape fungal microbiome that is mainly defined by the grape niche [1] and in a minor way
shaped by local environmental conditions and grape variety.

Through an unsupervised partitioning of these large set of samples we were able to capture most of
the microbial variability in three partitions that might reflect possible ecological states of the grape
fungal microbiome. Rather than containing distinctive microbial communities, the data suggests that
the disparities among geographical regions can be largely attributed to the prevalence of a few select
genera, including *Aureobasidium*, *Cladosporium* and *Saccharomyces* and suggests that samples
transition between three broad community types characterised by differences in diversity. While we
do not know the factors driving the transitions between community types, previous reports have
seen similar dynamics in the abundance of main genera (e.g. *Aureobasidium* and *Cladosporium*)
throughout the developmental stages of grapes [39], suggesting that maturity of grapes at sampling
time might be a driver between these transitions. These community types exhibited differences in
several important genera known to impact wine composition (e.g., *Saccharomyces*, *Hanseniaspora*,
Metschnikowia, *Pichia*, *Torulaspora*, *Starmerella* etc.) [68] and likely show functional heterogeneity
with regards to fermentation. Indeed, more diverse communities retained a higher fungal diversity
during the initial stages of fermentation, likely influencing fermentation performance and the
chemical composition of wine. Therefore, future research should focus on understanding the
environmental drivers behind shifts in community types, which could serve as a valuable tool to
modulate grape health and fermentation.

Conclusion
Taken together our results help to formulate a broader understanding of the community dynamics
within grape fermentations across varietal, spatial and temporal scales utilizing the largest survey of
grape ferment microbiomes to date. A holistic approach was taken with respect to the data analysis,
investigating the role that geography, grape variety and year has on microbiome composition and
diversity. When considering all scales, little structure was identified with wineries displaying highly
unstable fungal communities that transitioned within three broad community types. In contrast to
previous work suggesting that geography might play a large role in microbiome composition, it was
observed that both regional and state level geographies showed a small effect size on the overall

diversity and composition of ferments. In general, it was observed that initial community diversity
was transient between fermentation timepoints while playing an important, but imperfect, role in
predicting the diversity of the fungal microbiome throughout fermentation. This suggests that initial
microbial community is not deterministic and highlights the role abiotic factors have in determining
microbiome composition throughout the fermentation process.

Acknowledgements

357 This work was supported by Australian grapegrowers and winemakers through their investment body
358 Wine Australia, with matching funds from the Australian Government. The AWRI is a member of the
359 Wine Innovation Cluster (WIC) in Adelaide.

Data availability

362 The sequencing data included in this study is available in NCBI under BioProject PRJNA1053885.

Conflicts of interest

365 None declared.

Figure Legends

Fig. 1: Sampling design of the study. **a** Geographical location of the Australian wineries sampled
371 throughout a six-year period. Each point represents a winery and labels show the closest town/city. **b**
372 Sampling point for each grape juice sample: T1= freshly pressed juice, T2= 1 Bé drop in sugar, T3=
373 50% sugar remaining; T4= 10% sugar remaining. **c** Number of samples collected each year for each
374 timepoint.

Fig. 2: The core fungal community of grape juice. **a** Cumulative read abundance for 774 samples and
3594 OTUs. Red line represents the mean read abundance ± SD in red shading. OTUs were ranked in
order of abundance for all samples. The 10 most abundant OTUs represent 70 % (± 3.4 %) of the total
reads on average. **b** Frequency and observation as high abundant (OTUs making up the top 80% of
the reads as in plot **a**) of OTUs across all samples. Points are coloured based on number of OTUs and
Genus classification is shown for the top 5 OTUs observed as abundant. **c** Box plot showing the
relative abundance of the top 50 genera sorted by mean read abundance. Genera in red represent
taxa observed in at least 80% of all samples. Line inside the boxplot depicts the median and upper
and lower ends of each box represent the 25th and 75th percentiles. Outer lines show the minimum
and maximum value and dots represent potential outliers.

Fig. 3: The fungal diversity of grape juice. **a** Violin-plot showing the alpha-diversity (Shannon index) of
3113 samples separated by sampling timepoint. Multiple-comparison test was performed between

samples. **$P < 0.005$ (Dunn's, Holm-adjusted). **b** Non-metric multidimensional scaling (NMDS) of all
samples based on Bray-Curtis (BC) dissimilarities. **c** Bar-plot showing the effect size (R^2) of each
metadata category calculated by PERMANOVA (Adonis). **d** Principal component analysis (PCA) of the
Hellinger transformed OTU abundances of the three wineries displaying the highest F2 values after
pairwise comparison (Table Sx). Individual OTUs are shown in grey with the three most extreme
OTUs labelled to the genus level. **e** Temporal stability of the fungal communities. Baseline represents
the average within BC distances for each winery in the year 2018. 1 and 2 years represent the
average BC distances between the baseline and 2019 and 2021, respectively.

Fig. 4: Taxa can distinguish between geographical regions and grape varieties. Relative abundance for
the top 20 genera associated with **a** States of Australia and **b** grape varieties. Only taxa with a LDA
effect size > 2 are shown. Taxa are sorted based on their Mean Decrease Gini of random forest
models.

Fig. 5: Dirichlet Multinomial Mixtures (DMM) partitioning of grape juice fungal communities. **a**
Relative abundance for the top 30 Genera observed in each DMM partition. **b** Principal coordinate
analysis (PCoA) of T1 samples based on Bray-Curtis (BC) dissimilarities. Samples are coloured by
DMM as in panel (c) **c** Violin-plot showing the alpha-diversity (Shannon index) of samples belonging
to each DMM partition. Multiple-comparison test was performed between samples. *$P < 0.05$,
$P < 0.005$ (Dunn's, Holm-adjusted). **d Total number of samples allocated to each DMM partition
for each winery with more than 10 samples. Samples are coloured by DMM as in panel (c). **e** PHATE
scatter plots where samples have been coloured based on the root squared transformed relative
abundance of *Aureobasidium*, *Cladosporium* and *Saccharomyces*. **f** Relative abundance for the top 5
genera presented in panel (a) for timepoints T1 to T4 separated by DMM partition. Only DMM
partition samples containing timepoints T2 to T4 were used to produce the plot.

References

1.   Morrison-Whittle P, Goddard MR. Quantifying the relative roles of selective and neutral processes in defining eukaryotic microbial communities. *ISME J* 2015;**9**:2003-2011. doi: 10.1038/ismej.2015.18.

2.   Taylor MW, Tsai P, Anfang N, et al. Pyrosequencing reveals regional differences in fruit-associated fungal communities. *Environ Microbiol* 2014;**16**:2848-2858. doi: https://doi.org/10.1111/1462-2920.12456.

3.   Müller DB, Vogel C, Bai Y, et al. The plant microbiota: systems-level insights and perspectives. *Ann Rev Genet* 2016;**50**:211-234. doi: 10.1146/annurev-genet-120215-034952.

4.   Bokulich NA, Thorngate JH, Richardson PM, et al. Microbial biogeography of wine grapes is conditioned by cultivar, vintage, and climate. *Proc Natl Acad Sci USA* 2014;**111**:E139-E148. doi: doi:10.1073/pnas.1317377110.

5.   Gayevskiy V, Goddard MR. Geographic delineations of yeast communities and populations associated with vines and wines in New Zealand. *ISME J* 2012;**6**:1281-1290. doi: 10.1038/ismej.2011.195.

6.   Pinto C, Pinho D, Cardoso R, et al. Wine fermentation microbiome: a landscape from different Portuguese wine appellations. *Front Microbiol* 2015;**6**. doi: 10.3389/fmicb.2015.00905

7.   Beltran G, Torija MJ, Novo M, et al. Analysis of yeast populations during alcoholic fermentation: A six year follow-up study. *Syst Appl Microbiol* 2002;**25**:287-293. doi: https://doi.org/10.1078/0723-2020-00097.

8.   Combina M, Elía A, Mercado L, et al. Dynamics of indigenous yeast populations during spontaneous fermentation of wines from Mendoza, Argentina. *Int J Food Microbiol* 2005;**99**:237-243. doi: https://doi.org/10.1016/j.ijfoodmicro.2004.08.017.

9.   Fleet GH. Growth of yeasts during wine fermentations. *J Wine Res* 1990;**1**:211-223. doi: 10.1080/09571269008717877.

10.   Fleet GH. Wine yeasts for the future. *FEMS Yeast Res* 2008;**8**:979-995. doi: 10.1111/j.1567-1364.2008.00427.x.

11.   Fleet GH, Lafon-Lafourcade S, Ribereau-Gayon P. Evolution of yeasts and lactic acid bacteria during fermentation and storage of bordeaux wines. *Appl Env Microbiol* 1984;**48**:1034-1038. doi: doi:10.1128/aem.48.5.1034-1038.1984.

12.   Martini A, Ciani M, Scorzetti G. Direct enumeration and isolation of wine yeasts from grape surfaces. *Am J Enol Viticult* 1996;**47**:435-440. doi: 10.5344/ajev.1996.47.4.435.

13.   Mortimer R, Polsinelli M. On the origins of wine yeast. *Res Microbiol* 1999;**150**:199-204. doi: https://doi.org/10.1016/S0923-2508(99)80036-9.

14.   Sternes PR, Lee D, Kutyna DR, et al. A combined meta-barcoding and shotgun metagenomic analysis of spontaneous wine fermentation. *GigaScience* 2017;**6**:1-10. doi: 10.1093/gigascience/gix040.

15.   Caporaso JG, Lauber CL, Walters WA, et al. Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *Proc Natl Acad Sci USA* 2011;**108**:4516-4522. doi: 10.1073/pnas.1000080107.

16.   Bokulich NA, Collins TS, Masarweh C, et al. Associations among wine grape microbiome, metabolome, and fermentation behavior suggest microbial contribution to regional wine characteristics. *mBio* 2016;**7**. doi:10.1128/mbio.00631-16.

17.   Knight S, Klaere S, Fedrizzi B, et al. Regional microbial signatures positively correlate with differential wine phenotypes: evidence for a microbial aspect to terroir. *Sci Rep* 2015;**5**:14233. doi: 10.1038/srep14233.

18.   Bokulich NA, Mills DA. Improved selection of internal transcribed spacer-specific primers
enables quantitative, ultra-high-throughput profiling of fungal communities. *Appl Environ Microbiol*
2013;**79**:2519-2526. doi:10.1128/AEM.03870-12.
19.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics 2014:btu170.
20.   Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet. Journal 2011;**17**:3. doi: 10.14806/ej.17.1.200.
21.   Magoč T, Salzberg SL. FLASH: Fast length adjustment of short reads to improve genome
assemblies. *Bioinformatics* 2011. doi: 10.1093/bioinformatics/btr507.
22.   Mahé F, Rognes T, Quince C, et al. Swarm: robust and fast clustering method for amplicon-
based studies. *PeerJ* 2014;**2**:e593. doi: 10.7717/peerj.593.
23.   Caporaso JG, Kuczynski J, Stombaugh J, et al. QIIME allows analysis of high-throughput
community sequencing data. *Nature methods* 2010;**7**:335-336.
24.   Team RC. R. A language and environment for statistical computing [Internet]. Vienna, Austria:
R Foundation for Statistical Computing; 2013. *Document freely available on the internet at:*
http://www. r-project. org 2015.
25.   Wickham H. *ggplot2: elegant graphics for data analysis*: Springer, 2016.
26.   Skytte Andersen KS, Kirkegaard RH, Karst SM, et al. ampvis2: an R package to analyse and
visualise 16S rRNA amplicon data. *bioRxiv* 2018:299537. doi: 10.1101/299537.
27.   Dixon P. VEGAN, a package of R functions for community ecology. *J Veg Sci* 2003;**14**:927-930.
doi: 10.1111/j.1654-1103.2003.tb02228.x.
28.   Anderson MJ. A new method for non-parametric multivariate analysis of variance. *Aust Ecol*
2001;**26**:32-46. doi: https://doi.org/10.1111/j.1442-9993.2001.01070.pp.x.
29.   Anderson MJ, Walsh DCI. PERMANOVA, ANOSIM, and the Mantel test in the face of
heterogeneous dispersions: What null hypothesis are you testing? *Ecol Monogr* 2013;**83**:557-574.
doi: https://doi.org/10.1890/12-2010.1.
30.   Anderson MJ, Walsh DCI, Robert Clarke K, et al. Some solutions to the multivariate Behrens–
Fisher problem for dissimilarity-based analyses. *Aust NZ J Stat* 2017;**59**:57-79. doi:
https://doi.org/10.1111/anzs.12176.
31.   Segata N, Izard J, Waldron L, et al. Metagenomic biomarker discovery and explanation.
Genome Biol 2011;**12**:R60. doi: 10.1186/gb-2011-12-6-r60.
32.   Breiman L; Random Forests. *Machine Learning* 2001;**45**(1):5-32. doi:
10.1023/A:1010933404324.
33.   Liu C, Cui Y, Li X, et al. microeco: an R package for data mining in microbial community
ecology. *FEMS Microbiol Ecol* 2020;**97**(2). doi: 10.1093/femsec/fiaa255.
34.   Liaw A, Wiener M. Classification and regression by randomForest. *R news* 2002;**2**(3):18-22.
35.   Holmes I, Harris K, Quince C. Dirichlet Multinomial Mixtures: generative models for microbial
metagenomics. *Plos ONE* 2012;**7**:e30126. doi: 10.1371/journal.pone.0030126.
36.   Moon KR, van Dijk D, Wang Z, et al. Visualizing structure and transitions in high-dimensional
biological data. *Nat Biotechnol* 2019;**37**:1482-1492. doi: 10.1038/s41587-019-0336-3.
37.   Zalar P, Gostinčar C, de Hoog GS, et al. Redefinition of *Aureobasidium pullulans* and its
varieties. *Stud Mycol* 2008;**61**:21-38. doi: https://doi.org/10.3114/sim.2008.61.02.
38.   Liu D, Chen Q, Zhang P, et al. The fungal microbiome is an important component of vineyard
ecosystems and correlates with regional distinctiveness of wine. *mSphere* 2020;**5**. doi:
doi:10.1128/msphere.00534-20.
39.   Liu D, Howell K. Community succession of the grapevine fungal microbiome in the annual
growth cycle. *Environ Microbiol* 2021;**23**:1842-1857. doi: https://doi.org/10.1111/1462-2920.15172.
40.   Bauer F, Pretorius IS. Yeast stress response and fermentation efficiency: how to survive the
making of wine. 2000.
41.   Ciani M, Capece A, Comitini F, et al. Yeast interactions in inoculated wine fermentation. *Front*
Microbiol 2016;**7**. doi: 10.3389/fmicb.2016.00555.

42. Costea PI, Hildebrand F, Arumugam M, et al. Enterotypes in the landscape of gut microbial
community composition. *Nat Microbiol* 2018;**3**:8-16. doi: 10.1038/s41564-017-0072-8.
43. Hawkins DL, Ryder J, Lee SA, et al. Mixed yeast communities contribute to regionally distinct
wine attributes. *FEMS Yeast Res* 2023;**23**. doi: 10.1093/femsyr/foad005.
44. Tempère S, Marchal A, Barbe J-C, et al. The complexity of wine: clarifying the role of
microorganisms. *Appl Microbiol Biotechnol* 2018;**102**:3995-4007. doi: 10.1007/s00253-018-8914-8.
45. de Souza RSC, Okura VK, Armanhi JSL, et al. Unlocking the bacterial and fungal communities
assemblages of sugarcane microbiome. *Sci Rep* 2016;**6**:28774. doi: 10.1038/srep28774.
46. Saunders AM, Albertsen M, Vollertsen J, et al. The activated sludge ecosystem contains a
core community of abundant organisms. *ISME J* 2016;**10**:11-20. doi: 10.1038/ismej.2015.117.
47. Edwards J, Johnson C, Santos-Medellín C, et al. Structure, variation, and assembly of the
root-associated microbiomes of rice. *Proc Natl Acad Sci USA* 2015;**112**:E911-E920. doi:
doi:10.1073/pnas.1414592112.
48. Fitzpatrick CR, Copeland J, Wang PW, et al. Assembly and ecological function of the root
microbiome across angiosperm plant species. *Proc Natl Acad Sci USA* 2018;**115**:E1157-E1165. doi:
doi:10.1073/pnas.1717617115.
49. Kembel SW, O'Connor TK, Arnold HK, et al. Relationships between phyllosphere bacterial
communities and plant functional traits in a neotropical forest. *Proc Natl Acad Sci USA*
2014;**111**:13715-13720. doi: doi:10.1073/pnas.1216057111.
50. Abdelfattah A, Freilich S, Bartuv R, et al.; Global analysis of the apple fruit microbiome: are
all apples the same? *Environmental Microbiology* 2021;**23**(10):6038-6055. doi:
https://doi.org/10.1111/1462-2920.15469.
51. Morrison-Whittle P, Lee SA, Goddard MR. Fungal communities are differentially affected by
conventional and biodynamic agricultural management approaches in vineyard ecosystems. *Agr*
Ecosyst Environ 2017;**246**:306-313. doi: https://doi.org/10.1016/j.agee.2017.05.022.
52. Onetto CA, Schmidt SA, Roach MJ, et al. Comparative genome analysis proposes three new
Aureobasidium species isolated from grape juice. *FEMS Yeast Res* 2020;**20**. doi:
10.1093/femsyr/foaa052.
53. Rathnayake RMSP, Savocchia S, Schmidtke LM, et al. Characterisation of *Aureobasidium*
pullulans isolates from *Vitis vinifera* and potential biocontrol activity for the management of bitter
rot of grapes. *Eur J Plant Pathol* 2018;**151**:593-611. doi: 10.1007/s10658-017-1397-0.
54. Grube M, Schmid F, Berg G. Black fungi and associated bacterial communities in the
phyllosphere of grapevine. *Fung Biol* 2011;**115**:978-986. doi:
https://doi.org/10.1016/j.funbio.2011.04.004.
55. Prakitchaiwattana CJ, Fleet GH, Heard GM. Application and evaluation of denaturing gradient
gel electrophoresis to analyse the yeast ecology of wine grapes. *FEMS Yeast Res* 2004;**4**:865-877. doi:
10.1016/j.femsyr.2004.05.004.
56. Castoria R, De Curtis F, Lima G, et al. *Aureobasidium pullulans* (LS-30) an antagonist of
postharvest pathogens of fruits: study on its modes of action. *Postharvest Biol Tech* 2001;**22**:7-17.
doi: https://doi.org/10.1016/S0925-5214(00)00186-1.
57. Sharma RR, Singh D, Singh R. Biological control of postharvest diseases of fruits and
vegetables by microbial antagonists: A review. *Biol Control* 2009;**50**:205-221. doi:
https://doi.org/10.1016/j.biocontrol.2009.05.001.
58. Briceño EX, Latorre BA. Characterization of Cladosporium Rot in grapevines, a problem of
growing importance in Chile. *Plant Dis* 2008;**92**:1635-1642. doi: 10.1094/pdis-92-12-1635.
59. Nam MH, Park MS, Kim HS, et al. *Cladosporium cladosporioides* and *C. tenuissimum* cause
blossom blight in strawberry in Korea. *Mycobiology* 2015;**43**:354-359. doi:
10.5941/MYCO.2015.43.3.354.
60. Lutz MC, Sosa MC, Colodner AD. Effect of pre and postharvest application of fungicides on
postharvest decay of Bosc pear caused by Alternaria—Cladosporium complex in North Patagonia,
Argentina. *Sci Hortic* 2017;**225**:810-817. doi: https://doi.org/10.1016/j.scienta.2017.05.007.

61.   Albertin W, Setati ME, Miot-Sertier C, et al. *Hanseniaspora uvarum* from winemaking
environments show spatial and temporal genetic clustering. *Front Microbiol* 2016;**6**. doi:
10.3389/fmicb.2015.01569.
62.   Steenwyk JL, Opulente DA, Kominek J, et al. Extensive loss of cell-cycle and DNA repair genes
in an ancient lineage of bipolar budding yeasts. *PLOS Biol* 2019;**17**:e3000255. doi:
10.1371/journal.pbio.3000255.
63.   Haase MAB, Steenwyk JL, Boeke JD. Gene loss and cis-regulatory novelty shaped core histone
gene evolution in the apiculate yeast *Hanseniaspora uvarum*. *bioRxiv* 2023:2023.08.28.551515. doi:
10.1101/2023.08.28.551515.
64.   Torija MJ, Rozès N, Poblet M, et al. Yeast population dynamics in spontaneous fermentations:
Comparison between two different wine-producing areas over a period of three years. *A Van Leeuw J*
Microb 2001;**79**:345-352. doi: 10.1023/A:1012027718701.
65.   Vigentini I, De Lorenzis G, Fabrizio V, et al. The vintage effect overcomes the terroir effect: a
three year survey on the wine yeast biodiversity in Franciacorta and Oltrepò Pavese, two northern
Italian vine-growing areas. *Microbiol* 2015;**161**:362-373. doi: https://doi.org/10.1099/mic.0.000004.
66.   Evans KJ, Pirie AJG. Weather variables for within-vineyard awareness of *Botrytis* risk. *Aust J*
Grape Wine R 2024;**2024**:6630039. doi: 10.1155/2024/6630039.
67.   Sosnowski MR, Ayres MR, McCarthy MG, et al. Winegrape cultivars (*Vitis vinifera*) vary in
susceptibility to the grapevine trunk pathogens *Eutypa lata* and *Diplodia seriata*. *Aust J Grape Wine R*
2021;**28**:166-174. doi: https://doi.org/10.1111/ajgw.12531.
68.   Varela C. The impact of non-Saccharomyces yeasts in the production of alcoholic beverages.
Appl Microbiol Biotechnol 2016;**100**:9861-9874. doi: 10.1007/s00253-016-7941-6.