Leveraging Multi-echo EPI to Enhance BOLD Sensitivity in Task-based Olfactory fMRI

Ludwig Sichen Zhao[*1,2], Clara U. Raithel[3,2], M. Dylan Tisdall[4], John A. Detre[2,4], and Jay A.
Gottfried[*2,3]

[1]*Department of Bioengineering, School of Engineering and Applied Science, University of Pennsylvania, Philadelphia,*
PA, United States
[2]*Department of Neurology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, United States*
[3]*Department of Psychology, School of Arts and Sciences, University of Pennsylvania, Philadelphia, PA, United States*
[4]*Department of Radiology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, United States*

* Corresponding Author: Ludwig Sichen Zhao (ludwigz@seas.upenn.edu), Jay A. Gottfried
(jaygottf@pennmedicine.upenn.edu)

Abstract

Functional magnetic resonance imaging (fMRI) using blood-oxygenation-level-

dependent (BOLD) contrast relies on gradient echo echo-planar imaging (GE-EPI) to

quantify dynamic susceptibility changes associated with the hemodynamic response to

neural activity. However, acquiring BOLD fMRI in human olfactory regions is particularly

challenging due to their proximity to the sinuses where large susceptibility gradients

induce magnetic field distortions. BOLD fMRI of the human olfactory system is further

complicated by respiratory artifacts that are highly correlated with event onsets in

olfactory tasks.

Multi-Echo EPI (ME-EPI) acquires gradient echo data at multiple echo times (TEs)

during a single acquisition and can leverage signal evolution over the multiple echo

times to enhance BOLD sensitivity and reduce artifactual signal contributions. In the

current study, we developed a ME-EPI acquisition protocol for olfactory task-based fMRI

and demonstrated significant improvement in BOLD signal sensitivity over conventional

single-echo EPI (1E-EPI). The observed improvement arose from both an increase in

BOLD signal changes through a T_2^*-weighted echo combination and a reduction in non-

BOLD artifacts through the application of the Multi-Echo Independent Components

Analysis (ME-ICA) denoising method. This study represents one of the first direct

comparisons between 1E-EPI and ME-EPI in high-susceptibility regions and provides

compelling evidence in favor of using ME-EPI for future task-based fMRI studies.

Keywords: multi-echo EPI, olfaction, ME-ICA, susceptibility artifacts, task-based fMRI

1 Introduction

Functional magnetic resonance imaging (fMRI) is one of the most common imaging techniques used to non-invasively study both localized neural activation and whole-brain functional connectivity in humans. Blood-oxygenation-level-dependent (BOLD) fMRI measures brain activity by detecting local magnetic susceptibility fluctuations caused by changes in deoxyhemoglobin concentration. Gradient echo echo-planar imaging (GE-EPI) is the dominant BOLD imaging strategy due to its high temporal signal-to-noise ratio and rapid acquisition capabilities (Budde et al., 2014). However, artifacts caused by the local inhomogeneous magnetic field surrounding high-susceptibility regions present a major technical challenge in utilizing GE-EPI to measure BOLD signals. Olfactory regions, including the orbitofrontal cortex (OFC), amygdala, piriform (olfactory) cortex, and entorhinal cortex are particularly affected by susceptibility-induced artifacts and signal drop-out due their location near the ethmoid sinuses (air-filled cavities in the bone surrounding the nose). These artifacts limit the applicability of GE-EPI-based BOLD fMRI in pursuit of understanding the human olfactory system (Catani et al., 2013). Moreover, respiratory artifacts, primarily caused by local field shifts during inhalation and exhalation that are highly correlated with event onsets in tasked-based olfactory-related fMRI, further compromise the sensitivity of olfactory fMRI (Glover & Law, 2001).

Several techniques have been proposed to overcome these challenges. Acquiring scans at a tilted angle allows for partial compensation of local susceptibility gradients (Deichmann et al., 2003; Weiskopf et al., 2006). Alternatively, employing pulse sequences other than GE-EPI, such as T_2-Prepared BOLD fMRI and balanced steady state free precession (bSSFP) (Parrish et al., 2008; Miao et al., 2021).

Recent work on multi-echo EPI (ME-EPI), which acquires multiple gradient echoes at each imaging timepoint, has shown potential to improve BOLD sensitivity and efficiency in resting-state fMRI (Gonzalez-Castillo et al., 2016; Lynch et al., 2020). In conventional single-echo EPI (1E-EPI), a single volume is acquired after each RF excitation at a specific echo time (TE), which determines both image signal intensity as well as BOLD contrast (Donahue et al., 2011; Gati et al., 1997; van der Zwaag et al., 2009). At a

shorter TE, the image is more proton-density weighted, has less BOLD contrast, and

less severe signal dropout in regions with high local susceptibility gradients. Conversely,

with a longer TE, the image becomes more T_2^*-weighted, has greater BOLD contrast,

and more severe signal dropout. The BOLD contrast-to-noise ratio (CNR) is expected to

vary with TE, peaking at the T_2^* of each voxel (Donahue et al., 2011; Fera et al., 2004;

Gati et al., 1997; Graham et al., 1996; Triantafyllou et al., 2011; van der Zwaag et al.,

2009). However, various brain regions exhibit distinct T_2^* values, resulting in varying

optimal TEs, making it impossible to optimize TE across the whole brain in a single-

echo acquisition.

Unlike 1E-EPI, ME-EPI acquires multiple images across a range of TEs after each RF

excitation, resulting in enhanced functional contrast by combining the echo images with

different weights (Poser et al., 2006; Posse et al., 1999). The commonly used T_2^*-

weighted echo combination method assigns the heaviest weight to the echo that is

presumed to have the greatest BOLD CNR, with weighting performed separately for

each voxel. By applying a T_2^*-weighted echo combination with ME-EPI, BOLD signal

can be maximized at the voxel level while maintaining the temporal resolution needed

for fMRI. Moreover, by leveraging the expected exponential signal decay across TEs for

true BOLD contrast, noise reduction techniques can be employed to reject artifactual

signals and thereby improve the BOLD contrast-to-noise ratio (DuPre et al., 2021;

Gonzalez-Castillo et al., 2016; Kundu et al., 2012a, 2013, 2017; Van et al., 2023).

In this study, we implemented and tested a ME-EPI acquisition protocol and analysis

pipeline specifically designed for task-based olfactory-related paradigms by

incorporating shorter TE values in an effort to improve sensitivity in ventral brain regions

with high static susceptibility. Using this approach in comparison to a standard 1E-EPI

acquisition, we demonstrate superior sensitivity to olfactory task activation resulting in a

significant sample size reduction for detecting group task activation during a

representative olfactory three-alternatives forced choice task (3AFC). Our findings

suggest that ME-EPI holds great promise for human olfactory fMRI research.

2 Method

2.1 Subjects

29 subjects (13 women; aged 22 – 30 years, mean: 26, SD: 3.15) provided informed consent as approved by the University of Pennsylvania Institutional Review Board (#827217). One subject was excluded from the study due to excessive motion during fMRI scanning session, leaving us with a final sample of n = 28. All subjects reported being right-handed nonsmokers with no history of significant neurological/psychiatric disorder or olfactory dysfunction.

2.2 Odor Stimuli and Delivery

Stimuli consisted of two odorants: lemon oil extract (12.3% v/v, Sigma-Aldrich, MO, USA) and benzaldehyde (0.42% v/v, Sigma-Aldrich, MO, USA). Both odorants were diluted in mineral oil (Sigma Aldrich, MO, USA). Mineral oil alone was used as a control stimulus. The odor stimuli were delivered through a custom-built computer-controlled air-diluted 12-channel olfactometer with a flow rate of 4 L/min controlled using the PsychToolbox package (Brainard, 1997) and MATLAB (Version 2020b, The Mathworks Inc., Natick, MA, USA.)

2.3 Behavioral Testing Session

On Day 0, subjects participated in a behavioral testing session during which odor stimuli were presented outside the MRI environment. Participants were introduced to the odors and asked to label them as well as to rate the perceived odor intensity, via the general Labeled Magnitude Scale (gLMS) (Green et al., 1993, 1996). Valence (likeability) was also assessed via the Labeled Hedonic Scale (LHS) (L. V. Jones et al., 1955; Lim et al., 2009; Schutz & Cardello, 2001; Stone et al., 2020)], as well as familiarity [Visual Analog Scale (VAS) (Dourado et al., 2021; Flint et al., 2000)]. Ratings were analyzed in MATLAB (Version 2020b, The Mathworks Inc., Natick, MA, USA), using three one-way ANOVAs to evaluate whether there were any differences among stimuli.

 2.4   fMRI Scanning Session

2-5 days after the behavioral testing session, the same participants took part in an fMRI

scanning session, completing six runs, each of which contained 24 trials. On each trial,

participants sniffed one of the three possible odorants (lemon oil extract, benzaldehyde,

or mineral oil) and were asked to identify the odor using a three-alternative forced

choice task with the labels they provided earlier during the behavioral testing session.

Following the 3AFC task, they were also asked to rate the intensity of the odor. Each

odor was presented eight times per run in a random order. The experimental paradigm

as well as the sequence of events within a given trial is shown in Figure 1b.

Figure 1. fMRI Experimental Paradigm. a) The experiment was conducted over the course of two

sessions. On Day 0, the subjects participated in a series of behavioral tests where they were required to

identify odors and rate their intensity, valence, and familiarity. Subsequently, an fMRI scanning session

was scheduled for the subjects, taking place two to five days after their initial visit. This scanning session

consisted of three runs of ME-EPI and three runs of 1E-EPI acquisition. Each run comprised 24 trials,

with an equal distribution of three odor stimuli: lemon oil extract, benzaldehyde, and mineral oil. b)

Timeline for a single trial. A trial started with a fixation cross presented for 0.5 – 4.5 s, followed by a 1.5 s

countdown. A cue to sniff was then displayed and the odor was delivered for 2 s. Participants were then

asked to identify the odor and rate its intensity.

2.5   Structural and Functional MRI Data Acquisition

All magnetic resonance images were acquired using a 3 T scanner (MAGNETOM

Prisma, Siemens Healthineers, Germany) using the vendor's 64-channel head and neck

coil. T_1-weighted structural images were acquired using ME-MPRAGE (TE =

1.69/3.55/5.41/7.27 ms, TR = 2530 ms, TI = 1100 ms, FA = 8°, 1 mm isotropic).

1E-EPI and ME-EPI fMRI data were acquired using the sequence parameters outlined

in Table 1. For both protocols, repetition time (TR), field of view (FOV), number of

slices, resolution, flip angle (FA), echo spacing, and multiband factors are identical. To

optimize the BOLD signals in OFC regions, the field of view was tilted approximately 25°

to the anterior commissure-posterior commissure line (rostral > caudal) (Deichmann et

al., 2003). The two sequence types were administered in an interleaved fashion in each

participant, with the order being counterbalanced across participants. Of note, for 1E-

EPI the echo time was set at 22 ms, which is approximately the same as the second

echo time out of five echoes in the ME-EPI sequence (see Table 1). This TE aims to

balance signal dropout and BOLD sensitivity for olfactory-related regions and is

consistent with other olfactory-related fMRI studies (Bao et al., 2019; de Groot et al.,

2021; Raithel et al., 2023; Shanahan et al., 2021; Zhou et al., 2019).

	1E- EPI	ME-EPI
TR (ms)	2041	2041
TE (ms)	22	10.60/22.92/35.24/ 47.56/59.88
FOV (mm x mm)	216 x 216	216 x 216
Slices	48	48
Resolution (mm x mm x mm)	2.4 x 2.4 x 2.4	2.4 x 2.4 x 2.4
FA (degree)	78	78
Echo Spacing (ms)	0.49	0.49
Acceleration	Multiband 2x	Multiband 2x GRAPPA 3D, 3x

Table 1. fMRI Sequence Parameters. TR: Repetition time. TE: Echo time. FOV: Field of view. FA: Flip

angle. 1E-EPI: single-echo EPI. ME-EPI: multi-echo EPI.

2.6   fMRI Data Preprocessing

fMRI and structural MRI data were preprocessed in each subject's native space using

fMRIPrep 21.0.0 based on Nipype 1.6.1 (Esteban et al., 2019; Gorgolewski et al., 2011).

All images were then registered to ICBM 152 Nonlinear Asymmetrical template version

2009c (MNI-ICBM2009c) before further processing (Fonov et al., 2009, 2011).

2.7   ME-EPI T_2^* Estimation, Echo Combination, and Multi-Echo Independent

Components Analysis (ME-ICA)

Multi-echo EPI produces separate images for each echo time at each timepoint in the

functional imaging run. At each timepoint, these images for the multiple echo times were

combined using the T_2^*-weighted combination method (ME-WC) (Posse et al., 1999).

For each voxel, T_2^* was estimated by fitting the Eq. (1) using signals from all five

echoes.

$$S = S_0 e^{-TE/T_2^*} \tag{1}$$

166 The images from the five echoes are then combined voxel-wise using a weighted

167 average and the weights for each echo (W_{TE}) are calculated using Eq. (2) based on

168 estimated T_2^*.

$$W_{TE} = \frac{TE \cdot exp(-TE/T_2^*)}{\sum_{n=1}^{5} TE_n \cdot exp(-TE_n/T_2^*)} \tag{2}$$

Independent components analysis (ICA) is a common strategy to decompose fMRI

signals into distinct independent components, allowing the isolation of BOLD-related

neuronal components from sources of noise (Beckmann, 2012; Caballero-Gaudes &

Reynolds, 2017; Mckeown et al., 1998; McKeown et al., 2003). However, a persistent

challenge lies in the manual classification of these independent components, which is

labor-intensive and can yield inconsistent results (Kelly et al., 2010). In an attempt to

tackle this problem, ME-ICA classifies each component as either a non-BOLD artifact,

such as respiratory artifact, or a BOLD-like signal based on their signal characteristics

across different echo times (Kundu et al., 2012b, 2013). Here, we specifically employed

the TE-dependency analysis pipeline (Tedana 0.0.12) (DuPre et al., 2021) to

preprocess multi-echo EPI data before registering them to the MNI template.

2.8   Physiological Data Analysis

The respiratory trace was recorded with a respiratory transducer using a pressure

sensor (BIOPAC System, CA, USA) affixed around the subject's chest, and then

preprocessed before being added as a nuisance regressor: the trace was smoothed

using a 250 ms moving window, high-pass filtered at 0.05 Hz, detrended, and

standardized. The preprocessed trace was also downsampled to 1/2.4 Hz to be

included as a nuisance regressor. Two additional nuisance regressors were introduced:

the trial-by-trial sniff volume was defined as the amplitude difference between inhalation

and exhalation, and the sniff duration was defined as the time difference between

inhalation and exhalation.

2.9   Statistical Analysis

All fMRI data were analyzed in SPM12 using a general linear model to estimate the

main effects of the odor stimulation. Each experimental condition (control, lemon oil,

and benzaldehyde) was modeled using an independent regressor for each run. This

results in a 2 (multi-echo and single-echo) by 3 (control, lemon oil, and benzaldehyde)

factorial design. The model included the following nuisance regressors: trial-by-trial sniff

volume and sniff duration, both convolved with the hemodynamic response function

(HRF), as well as the down-sampled breathing trace and 24 movement parameters (six

movement parameters derived from spatial realignment, their squares, derivatives, and

squares of derivatives). In all GLMs, data were high-pass filtered at 1/128 Hz; the

temporal autocorrelation was estimated and removed as an autoregressive AR(1)

process globally (Friston et al., 2002).

Contrasts of interest were defined as [Odor > Control], [Lemon > Control] and

[Benzaldehyde > Control]. Whole brain analysis was performed to validate that odor-

evoked activations were present in the data acquired through both protocols (1E-EPI

and ME-EPI). Small volume corrections (SVC) and ROI analysis were separately

performed on OFC, amygdala, piriform cortex and entorhinal cortex, as suggested by

previous studies highlighting the involvement of these regions in human olfactory

processing (Anderson et al., 2003). Amygdala and piriform cortex were defined

anatomically using a human brain atlas (Mai et al., 2016). Delineation of OFC and

entorhinal cortex were based on the MNI-ICBM2009c atlas (Manera et al., 2020).

2.10   Post-hoc Power Analysis

We used our ROI analysis results to estimate the sample sizes required to detect

significant olfactory-related group fMRI activation in subsequent studies. A priori power

analyses were performed with α = 0.05 and β = 0.80 using G*Power (Faul et al., 2009)

for both ME-EPI and 1E-EPI data in the same ROIs described in the previous section.

3   Results

3.1   Echo-combined Images Reduce Signal Dropout

We first examined the estimated T_2^* values for each olfactory-related region. As

expected, signal dropouts are reduced in shorter echo times (Figure 2a, TE_1 and TE_2),

while T_2^*-weighted contrast is developed in later echo times (Figure 2a, $TE_3 - TE_5$). In

particular, the medial and lateral OFC along with the entorhinal cortex, exhibited notably

shorter T_2^* values due to susceptibility artifacts (Figure 2a & 2b), and in turn the echo-

combined image more heavily weights the shorter echoes (Figure 2c). Using these

weights, the combined image (Figure 2a, WC) effectively mitigated signal dropout while

retaining notable T_2^*-weighted contrast.

Figure 2. Multi-echo fMRI Reduces Signal Dropout and Improves BOLD Contrast through T_2^*-Weighted

Combination in a Single Subject. A) Two representative coronal slices containing the ROIs of interest are

shown with a variety of contrasts: ROIs overlayed on the T_1-weighted image (ROI); estimated T_2^* maps;

all five individual echo images ($TE_1 – TE_5$) from the ME-EPI acquisition; and the T_2^*-weighted combined

images (WC). b) Plot of signal vs echo time for each anatomical ROI. c) Within-ROI mean of the voxel-

wise weights used to compute the T_2^*-weighted combined images.

3.2   BOLD Activation in Diverse Olfactory Brain Regions During Odor Stimulation

We next investigated the regions that were significantly activated when the odors were

delivered under ME-EPI acquisition with the ME-ICA denoising method [Odor > Control]

(Figure 3a) (DuPre et al., 2021). Consistent with previous literature, we observed

significant activation in the piriform cortex, amygdala, entorhinal cortex, insula, and OFC

bilaterally ($p < 0.001$, uncorrected) (Anderson et al., 2003; Dikecligil & Gottfried, 2024;

Gottfried et al., 2003; Howard & Gottfried, 2014; Patin & Pause, 2015; Plailly et al.,

2008). We next examined the BOLD effects of individual odor stimulus, namely

[Benzaldehyde > Control] and [Lemon > Control]. Despite statistical testing indicating no

significant differences in perceived intensity (p = 0.38), valence (p = 0.45), and

familiarity (p = 0.11) between benzaldehyde and lemon oil, our analysis revealed that

the main effect [Odor > Control] was driven mainly by the contrast [Lemon > Control]. In

comparison, the contrast [Benzaldehyde > Control] produced significant levels of

activation only in the bilateral insular region (Figure 3a). The difference in BOLD

response to these stimuli was not anticipated though not wholly unprecedented, as

previous research has demonstrated that different odors can elicit distinct responses

within olfactory brain regions even in the absence of differences in low-level stimulus

features (Fournel et al., 2016; Howard & Gottfried, 2014; Savic, 2002; Savic et al.,

2002). In the single-echo fMRI, a similar trend is observed, with more active regions

being detected with lemon oil (p < 0.05, uncorrected, Figure S1). Accordingly, to

compare and observe the differences between acquisition methods within olfactory

regions, we will focus on the activation under lemon oil condition [Lemon > Control] for

our further analyses.

Figure 3. Activation Maps from Olfactory-related Task for Both 1E-EPI and ME-EPI. a) Two representative

coronal slices of activation maps for all three contrasts ([Odor > Control], [Benzaldehyde > Control],

[Lemon > Control]) using ME-EPI acquisition with ME-ICA denoising methods ($p < 0.05$, FDR corrected).

b) Activation maps for [Lemon > Control] contrast for ME-EPI acquisition with ME-ICA denoising methods

(ME-ICA), and without ME-ICA denoising methods (ME-WC) as well as single-echo fMRI acquisition (1E-

EPI). The left two columns show the maps without multiple comparison correction ($p < 0.001$) while the

263 right two columns show the maps corrected for multiple comparison ($p < 0.05$, FDR corrected). No

264 activation survived correction for multiple comparisons for 1E-EPI (not shown). c) Z-score statistical maps

265 yielded from Levene's test to examine whether BOLD signal variance was lower in ME-WC [$\sigma_{ME\text{-}WC} < \sigma_{1E\text{-}}$

266 $_{EPI}$] and ME-EPI [$\sigma_{ME\text{-}ICA} < \sigma_{1E\text{-}EPI}$] compared to 1E-EPI ($p < 0.05$, FDR corrected).

3.3   ME-EPI Improves fMRI Signal Sensitivity by Both Increasing BOLD Signal Changes and Reducing non-BOLD Artifacts

3.3   ME-EPI Improves fMRI Signal Sensitivity by Both Increasing BOLD Signal

Changes and Reducing non-BOLD Artifacts

Having validated that our ME-EPI acquisition protocol is suitable to capture odor-evoked

BOLD responses, we next compared BOLD activation among different acquisition

methods (1E-EPI vs. ME-EPI) and analysis pipelines (ME-WC vs. ME-ICA). A whole-

brain analysis using either whole-brain multiple comparisons correction (FDR)

(Benjamini & Hochberg, 1995; Benjamini & Yekutieli, 2001; Yekutieli & Benjamini, 1999)

or small volume correction (SVC) did not yield any supra-threshold voxels for 1E-EPI.

However, the same whole-brain analysis using the T_2^*-weighted combined ME-EPI (ME-

WC) highlighted activation in all ROIs (piriform cortex, amygdala, OFC, and entorhinal

cortex) (Figure 3b). Moreover, by using ME-ICA denoising methods (DuPre et al.,

2021), we observed a further increase in both the number of activation voxels and peak

Z values in all ROIs (Table 2).

	Piriform Cortex		Amygdala		Entorhinal Cortex		OFC	
	# Voxel	Peak Z	# Voxel	Peak Z	# Voxel	Peak Z	# Voxel	Peak Z
ME-ICA	167	5.44	9	4.31	12	4.14	32	5.44
ME-WC	5	4.04	2	3.98	1	3.74	10	4.44
1E-EPI	0	N/A	0	N/A	0	N/A	0	N/A

Table 2. Number of Voxels and Peak Z-score Activated by Lemon Oil for each ROI (p < 0.05, SVC).

We next examined potential reasons for the observed differences. Specifically, we

hypothesized the improvement of the BOLD sensitivity in the case of the ME-EPI data

was due to more pronounced BOLD signal changes, as we performed a locally

weighted combination of all echoes for each individual voxel instead of using one single

TE across the brain. BOLD signal changes could be calculated by contrast estimates

from the general linear model (GLM). Therefore, we performed a paired t-test on

contrast estimates [Lemon > Control] between 1E- and ME-EPI with ME-ICA [ME-ICA >

1E-EPI] among all ROIs. Only piriform cortex ($p = 0.023$) and lateral OFC ($p = 0.006$)

showed greater estimates for the contrast [Lemon > Control] for ME-ICA compared to

1E-EPI.

An alternative explanation for the observed differences between acquisition protocols,

that is, the increased BOLD sensitivity for ME-EPI versus 1E-EPI, may lie in the

reduction of the background noise due to non-BOLD signals. To test the hypothesis that

ME-EPI signal changes have a lower variance, we performed a Levene's test (Levene,

1960). This test is commonly employed to assess whether two or more groups have

equal variances in a dataset (Chiarello et al., 2009; Farinha et al., 2022; Im et al., 2010;

T. B. Jones et al., 2010; Swarup et al., 2013). Using a whole-brain analysis we first

generated a mask using the voxels activated by the [Lemon > Control] contrast and $p <$

0.001 (uncorrected) through the ME-ICA acquisition and process. Within this mask, we

find that 2.98% of these masked voxels reached the significance threshold ($p < 0.05$,

FDR corrected) for Levene's test comparing ME-WC and 1E-EPI. This result aligns with

prior research indicating that ME-EPI, even without the application of ME-ICA, has the

potential to decrease non-BOLD noise in resting-state analysis (Kundu et al., 2017;

Lynch et al., 2020; Posse et al., 1999). Subsequently, we aimed to assess whether

using ME-ICA can reduce noise above and beyond ME-WC. Indeed, the number of

significant voxels further increased to 23.1% of masked voxels when comparing ME-ICA

and 1E-EPI (Figure 3c). Taken together, our data suggest that our observed increase in

BOLD detection is not only due to increased sensitivity BOLD signals, but also the

reduction of non-BOLD artifacts.

3.4   ME-EPI Improves Statistical Power and Reduces Sample Sizes

So far, we have established that ME-EPI improved BOLD signal sensitivity by both

enhancing BOLD signal and reducing non-BOLD artifacts. An outstanding question is to

what degree BOLD sensitivity is improved when using our suggested method in the

context of task-based olfactory fMRI. To quantify the change in BOLD sensitivity, we

examined the contrast [Lemon > Control] in the brain regions central to olfaction,

namely the piriform cortex, amygdala, lateral and medial OFC, and entorhinal cortex for

both acquisition methods separately and then compared the statistical outcomes.

Whereas activation in medial OFC did not reach statistical significance using data

resulting from either acquisition method, all remaining regions show much lower *p-*

values for ME-ICA compared to 1E-EPI (Table 3). Furthermore, to determine the

minimum sample size required to observe significance in each region, post-hoc power

analyses were carried out ($\alpha = 0.05$ and $\beta = 0.80$) to estimate the effect of each

acquisition method on the sample sizes needed to detect effects in similar olfactory task

fMRI studies. ME-ICA led to a substantial reduction in the sample size, ranging from by

half to one-fourth, compared to 1E-EPI, depending on the specific regions (Figure 4).

Figure 4. Estimated Sample Size for ROI Analyses Based on Post-hoc Power Analyses ($\alpha = 0.05$ and $\beta =$

0.80) in all Olfactory-related ROIs under Contrast [Lemon > Control] for Single-echo fMRI (1E-EPI), ME-

EPI BOLD Acquisition with ME-ICA Denoising Methods (ME-ICA), and without ME-ICA Denoising

Methods (ME-WC).

	Piriform Cortex	Amygdala	Entorhinal Cortex	Lateral OFC	Medial OFC
ME-ICA	< 0.0001***	0.007***	0.005**	0.006**	0.884
ME-WC	0.0006***	0.044*	0.065	0.0138*	0.612
1E-EPI	0.007**	0.024*	0.0157*	0.549	0.669

332   *p < 0.001 ***   p < 0.01 **   p < 0.05 ***

333   *Table 3. Statistical Significance of ROI Analyses for Each Olfactory-related ROI under the Contrast*

334   *[Lemon > Control] for 1E-EPI, ME-WC and ME-ICA.*

4   Discussion

Susceptibility artifacts pose a significant challenge when using GE-EPI for fMRI studies.

The primary olfactory cortex not only suffers from some of the most severe static

susceptibility artifacts within the human brain but is also affected by respiration-related

artifacts. Here, we have developed and validated a ME-EPI protocol and analysis

pipeline specifically tailored for olfactory task-based fMRI. In the context of a basic

olfactory 3AFC task, we found that ME-EPI, alongside ME-ICA denoising, systemically

improves BOLD sensitivity and reduces the required sample sizes needed to infer a

significant effect at a group level compared to conventional 1E-EPI. This is achieved by

both increasing BOLD signal sensitivity and reducing the contributions of non-BOLD

signal changes using the additional denoising features ME-EPI provided.

The echo times used for ME-BOLD provided a range of contrasts across different TEs

for each ROI of interest (Figure 2a). When evaluating the T_2^*-derived weights calculated

for the echo-combined image, we confirmed that high-susceptibility regions with short

T_2^* values, namely, lateral and medial OFC, and entorhinal cortex, had higher weights

assigned to the first two echoes. In contrast, the remaining ROIs had higher weights

assigned to the remaining (and later) three echoes. As a result, the T_2^*-weighted

combined (WC) images were able to produce more uniform T_2^*-weighted contrast

across the whole brain (Figure 2a, WC).

When investigating the region activated by odor onset using ME-ICA, we detected

activation in all major primary olfactory cortex regions (piriform cortex, amygdala,

entorhinal cortex, and OFC bilaterally) in addition to the insular region (Figure 3a).

These findings are consistent with previous literature (Anderson et al., 2003; Dikecligil &

Gottfried, 2024; Gottfried et al., 2003; Howard & Gottfried, 2014; Patin & Pause, 2015;

Plailly et al., 2008). Further analysis revealed that the activation was primarily driven by

the lemon extract stimulus in comparison to benzaldehyde. Previous studies have

demonstrated that different odor stimuli can activate distinct primary olfactory regions to

different degrees (Fournel et al., 2016; Howard & Gottfried, 2014; Savic, 2002; Savic et

al., 2002). In this study we found that that benzaldehyde activates the olfactory system

to a lesser extent than lemon oil, such that the observed activation did not reach

significance after correcting for multiple comparisons in the whole brain analysis.

When comparing the results of 1E-EPI and ME-EPI, we observed significantly increased

sensitivity in both whole brain analysis and ROI analysis, which are two commonly used

univariate fMRI analysis methods. In the case of whole brain analysis (Figure 3b), we

found that ME-EPI exhibited higher peak Z values and a greater number of activated

voxels compared to 1E-EPI, as shown in Table 2. It is worth noting that even without

using the ME-ICA denoising method, ME-EPI still outperformed 1E-EPI by performing a

T_2^*-weighted combination of data from all echoes. This improvement was expected, as

the weighting of data from each echo time was designed to enhance the sensitivity to

BOLD contrast (Posse et al., 1999).

For ROI analysis and sequential power analysis, which were used to estimate minimal

sample sizes, ME-EPI with the ICA denoising method demonstrated superiority over 1E-

EPI, as shown in Table 3 and Figure 4. We attribute these improvements to two factors:

the increase of BOLD signal intensity through T_2^*-weighted echo combination and the

reduction of noise by leveraging signal evaluation across the echo times. To support

these hypotheses, we conducted a series of statistical tests (Figure 3c) to show the

mean signals are significantly higher and the variance is significantly lower in ME-ICA

acquisition compared to 1E-EPI.

The current analysis pipeline employs a specific method of combining multiple echo

image series into a single image series: T_2^*-weighted combination, also known as

"optimal combination", is used in most ME-EPI studies as it is designed to maximize

BOLD-contrast sensitivity (Baracchini et al., 2021; Gonzalez-Castillo et al., 2016; Kundu

et al., 2017; Lynch et al., 2020; Roth et al., 2020). However, an interesting finding

emerged when we processed the ME-EPI data separately and treated each individual

echo series as a 1E-EPI dataset. Surprisingly, no supra-threshold activation was

detected at the first echo ($p < 0.05$, uncorrected), while the majority of activations were

observed at the second and third echoes ($p < 0.05$, uncorrected). When examining the

Z values for the ROI analyses across all 5 individual echoes, we also observe a similar

trend (Table S1). However, regions with short T_2^* (such as the lateral and medial OFC

and entorhinal cortex in this study) predominantly assigned higher weights to the first
two echoes.

In addition to the incongruence between apparent BOLD signal in the earliest echo and
the weight assigned to it, T_2^* values near the ethmoid sinuses fluctuate during inhalation
and exhalation due to local B_0 field shifts (Barry & Menon, 2005; Raj et al., 2000, 2001;
Van de Moortele et al., 2002). In olfactory studies, this causes significant task-related
artifacts. The T_2^*-weighted combination method used in our current analyses employed
fixed weighting throughout one scan and does not account for this dynamic T_2^*
fluctuation; the use of dynamically adjusted weights might also improve BOLD
sensitivity. Further work is required to determine if more information could be used to
choose optimal echo weightings that might further increase BOLD-sensitivity (*e.g.*, by
reducing the weights on the first echo, even in the short-T_2^* voxels). These findings
may, in part, elucidate the observed effects in Table 2 and Figure 4, wherein the
estimated sample sizes are higher, and the statistical significances are lower for ROI
analyses in certain regions for ME-WC compared to 1E-EPI.

There have also been recent advances in denoising methods that leverage phase
information, such as Noise Reduction with Distribution Corrected (NORDIC) PCA
(Dowdle et al., 2023; Moeller et al., 2021; Vizioli et al., 2021). These methods have
demonstrated the ability to further enhance BOLD sensitivity in resting-state fMRI.
However, our study did not acquire phase data, which prevented us from testing these
advanced denoising methods in our data.

In this study, we utilized a limited set of two odor stimuli, along with a control stimulus.
Although this approach was adequate to compare the ME-EPI and 1E-EPI acquisition
methods, future studies will aim to incorporate a wider range of odor stimuli. By
including multiple odor stimuli, we can expand our analysis to encompass multivariate
analysis, such as Multivariate Pattern Analysis (MVPA), which has been commonly
utilized in other olfaction task-based fMRI studies (Donoshita et al., 2021; Howard et al.,
2009; Howard & Gottfried, 2014; Perszyk et al., 2023), though we expect also to see an
improvement in multivariate analysis using ME-ICA.

In conclusion, our study demonstrated the utility of a ME-EPI protocol combined with a
pre-configured analysis pipeline for fMRI in olfactory cortex. This approach enhanced
sensitivity to task-dependent BOLD contrast changes in olfactory regions, and post-hoc
power analyses indicate that group effects can be detected with a reduced sample size
compared to standard 1E-EPI. Although our protocol and pipeline were specifically
designed for task-based fMRI in the presence of large static susceptibility, a
longstanding challenge in the field of fMRI, they showcase the superior denoising
capabilities of ME-EPI for task-based fMRI more generally. These findings strongly
support the adoption of ME-EPI over 1E-EPI for future task-based fMRI studies.

Data and Code Availability

433 The unthresholded Z-maps showing the whole-brain results for univariate analyses
434 were made available on Neurovault (https://neurovault.org/collections/KGZMOMWR/).

Author Contributions

Ludwig Sichen Zhao: Conceptualization, Methodology, Software, Formal Analysis,
Investigation, Data Curation, Visualization, Writing – Original Draft, Project
administration. **Clara U. Raithel**: Conceptualization, Methodology, Investigation, Writing
– Review & Editing. **M. Dylan Tisdall**: Conceptualization, Methodology, Writing –
Review & Editing, Supervision, Resources. **John A. Detre**: Conceptualization,
Methodology, Writing – Review & Editing, Supervision, Resources. **Jay A. Gottfried**:
Conceptualization, Methodology, Writing – Review & Editing, Supervision, Resources,
Funding acquisition

Funding

445 This work was supported by the National Institutes of Health award R01DC019405
446 [awarded to JAG].

Declaration of Competing Interests

The authors have declared that no competing interests exist.

Supplementary Materials

Figure S1. Comparison of Activation Maps Among Different Contrasts for Olfactory-Related Tasks in Both
1E-EPI and ME-EPI. Two representative coronal slices of activation maps for all three contrasts ([Odor >
Control], [Benzaldehyde > Control], [Lemon > Control]) using ME-EPI acquisition with ME-ICA denoising
methods (ME-ICA) and using single-echo fMRI (1E-EPI) acquisition ($p < 0.001$, uncorrected).

	Piriform Cortex	Amygdala	Entorhinal Cortex	Orbitofrontal Cortex
TE_1 (10.60 ms)	1.945	1.010	0.756	1.205
TE_2 (22.92 ms)	2.383	1.957	2.183	1.605
TE_3 (35.24 ms)	3.457	1.583	1.789	1.724
TE_4 (47.56 ms)	3.819	1.355	0.973	1.835
TE_5 (59.98 ms)	3.693	2.604	1.265	1.705

457 *Table S1. Z Scores of ROI Analyses for Each Olfactory-related ROI under the Contrast [Lemon > Control]*
458 *When Analyzing 5 Individual Echoes from ME-ICA Separately.*

References

Anderson, A. K., Christoff, K., Stappen, I., Panitz, D., Ghahremani, D. G., Glover, G., Gabrieli, J. D. E., & Sobel, N. (2003). Dissociated neural representations of intensity and valence in human olfaction. *Nature Neuroscience*, *6* (2), Article 2. https://doi.org/10.1038/nn1001

Bao, X., Gjorgieva, E., Shanahan, L. K., Howard, J. D., Kahnt, T., & Gottfried, J. A. (2019). Grid-like Neural Representations Support Olfactory Navigation of a Two-Dimensional Odor Space. *Neuron*, *102* (5), 1066-1075.e5. https://doi.org/10.1016/j.neuron.2019.03.034

Baracchini, G., Mišić, B., Setton, R., Mwilambwe-Tshilobo, L., Girn, M., Nomi, J. S., Uddin, L. Q., Turner, G. R., & Spreng, R. N. (2021). Inter-regional BOLD signal variability is an organizational feature of functional brain networks. *NeuroImage*, *237*, 118149. https://doi.org/10.1016/j.neuroimage.2021.118149

Barry, R. L., & Menon, R. S. (2005). Modeling and suppression of respiration-related physiological noise in echo-planar functional magnetic resonance imaging using global and one-dimensional navigator echo correction. *Magnetic Resonance in Medicine*, *54* (2), 411–418. https://doi.org/10.1002/mrm.20591

Beckmann, C. F. (2012). Modelling with independent components. *NeuroImage*, *62* (2), 891–901. https://doi.org/10.1016/j.neuroimage.2012.02.020

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)*, *57* (1), 289–300. https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Benjamini, Y., & Yekutieli, D. (2001). The control of the false discovery rate in multiple

testing under dependency. *The Annals of Statistics*, *29* (4), 1165–1188.

https://doi.org/10.1214/aos/1013699998

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10* (4), 433–436.

https://doi.org/10.1163/156856897X00357

Budde, J., Shajan, G., Zaitsev, M., Scheffler, K., & Pohmann, R. (2014). Functional MRI

in human subjects with gradient-echo and spin-echo EPI at 9.4 T. *Magnetic*

Resonance in Medicine, *71* (1), 209–218. https://doi.org/10.1002/mrm.24656

Caballero-Gaudes, C., & Reynolds, R. C. (2017). Methods for cleaning the BOLD fMRI

signal. *NeuroImage*, *154*, 128–149.

https://doi.org/10.1016/j.neuroimage.2016.12.018

Catani, M., Dell'Acqua, F., & Thiebaut de Schotten, M. (2013). A revised limbic system

model for memory, emotion and behaviour. *Neuroscience & Biobehavioral*

Reviews, *37* (8), 1724–1737. https://doi.org/10.1016/j.neubiorev.2013.07.001

Chiarello, C., Welcome, S. E., Halderman, L. K., Towler, S., Julagay, J., Otto, R., &

Leonard, C. M. (2009). A large-scale investigation of lateralization in cortical

anatomy and word reading: Are there sex differences? *Neuropsychology*, *23* (2),

210–222. https://doi.org/10.1037/a0014265

de Groot, J. H. B., Kirk, P. A., & Gottfried, J. A. (2021). Titrating the Smell of Fear: Initial

Evidence for Dose-Invariant Behavioral, Physiological, and Neural Responses.

Psychological Science, *32* (4), 558–572.

https://doi.org/10.1177/0956797620970548

Deichmann, R., Gottfried, J. A., Hutton, C., & Turner, R. (2003). Optimized EPI for fMRI

studies of the orbitofrontal cortex. *NeuroImage*, *19* (2), Article 2.

https://doi.org/10.1016/S1053-8119(03)00073-9

Dikecligil, G. N., & Gottfried, J. A. (2024). What Does the Human Olfactory System Do,

and How Does It Do It? *Annual Review of Psychology*, *75* (1), null.

https://doi.org/10.1146/annurev-psych-042023-101155

Donahue, M. J., Hoogduin, H., van Zijl, P. C. M., Jezzard, P., Luijten, P. R., &

Hendrikse, J. (2011). Blood oxygenation level-dependent (BOLD) total and

extravascular signal changes and $\Delta R2^*$ in human visual cortex at 1.5, 3.0 and

7.0 T. *NMR in Biomedicine*, *24* (1), 25–34. https://doi.org/10.1002/nbm.1552

Donoshita, Y., Choi, U.-S., Ban, H., & Kida, I. (2021). Assessment of olfactory

information in the human brain using 7-Tesla functional magnetic resonance

imaging. *NeuroImage*, *236*, 118212.

https://doi.org/10.1016/j.neuroimage.2021.118212

Dourado, G. B., Volpato, G. H., de Almeida-Pedrin, R. R., Pedron Oltramari, P. V.,

Freire Fernandes, T. M., & de Castro Ferreira Conti, A. C. (2021). Likert scale vs

visual analog scale for assessing facial pleasantness. *American Journal of*

Orthodontics and Dentofacial Orthopedics, *160* (6), 844–852.

https://doi.org/10.1016/j.ajodo.2020.05.024

Dowdle, L. T., Vizioli, L., Moeller, S., Akçakaya, M., Olman, C., Ghose, G., Yacoub, E.,

& Uğurbil, K. (2023). Evaluating increases in sensitivity from NORDIC for diverse

fMRI acquisition strategies. *NeuroImage*, *270*, 119949.

https://doi.org/10.1016/j.neuroimage.2023.119949

DuPre, E., Salo, T., Ahmed, Z., Bandettini, P. A., Bottenhorn, K. L., Caballero-Gaudes,

C., Dowdle, L. T., Gonzalez-Castillo, J., Heunis, S., Kundu, P., Laird, A. R.,

Markello, R., Markiewicz, C. J., Moia, S., Staden, I., Teves, J. B., Uruñuela, E.,

Vaziri-Pashkam, M., Whitaker, K., & Handwerker, D. A. (2021). TE-dependent

analysis of multi-echo fMRI with *tedana*. *Journal of Open Source Software*, *6*

(66), 3669. https://doi.org/10.21105/joss.03669

Esteban, O., Markiewicz, C. J., Blair, R. W., Moodie, C. A., Isik, A. I., Erramuzpe, A.,

Kent, J. D., Goncalves, M., DuPre, E., Snyder, M., Oya, H., Ghosh, S. S., Wright,

J., Durnez, J., Poldrack, R. A., & Gorgolewski, K. J. (2019). fMRIPrep: A robust

preprocessing pipeline for functional MRI. *Nature Methods*, *16* (1), Article 1.

https://doi.org/10.1038/s41592-018-0235-4

Farinha, M., Amado, C., Morgado, P., & Cabral, J. (2022). Increased Excursions to

Functional Networks in Schizophrenia in the Absence of Task. *Frontiers in*

Neuroscience, *16*. https://www.frontiersin.org/articles/10.3389/fnins.2022.821179

Faul, F., Erdfelder, E., Buchner, A., & Lang, A.-G. (2009). Statistical power analyses

using G*Power 3.1: Tests for correlation and regression analyses. *Behavior*

Research Methods, *41* (4), 1149–1160. https://doi.org/10.3758/BRM.41.4.1149

Fera, F., Yongbi, M. N., van Gelderen, P., Frank, J. A., Mattay, V. S., & Duyn, J. H.

(2004). EPI-BOLD fMRI of human motor cortex at 1.5 T and 3.0 T: Sensitivity

dependence on echo time and acquisition bandwidth. *Journal of Magnetic*

Resonance Imaging, *19* (1), 19–26. https://doi.org/10.1002/jmri.10440

Flint, A., Raben, A., Blundell, J. E., & Astrup, A. (2000). Reproducibility, power and

validity of visual analogue scales in assessment of appetite sensations in single

test meal studies. *International Journal of Obesity*, *24* (1), Article 1.

https://doi.org/10.1038/sj.ijo.0801083

Fonov, V., Evans, A. C., Botteron, K., Almli, C. R., McKinstry, R. C., & Collins, D. L.

(2011). Unbiased average age-appropriate atlases for pediatric studies.

NeuroImage, *54* (1), 313–327. https://doi.org/10.1016/j.neuroimage.2010.07.033

Fonov, V., Evans, A., McKinstry, R., Almli, C., & Collins, D. (2009). Unbiased nonlinear

average age-appropriate brain templates from birth to adulthood. *NeuroImage*,

47, S102. https://doi.org/10.1016/S1053-8119(09)70884-5

Fournel, A., Ferdenzi, C., Sezille, C., Rouby, C., & Bensafi, M. (2016). Multidimensional

representation of odors in the human olfactory cortex. *Human Brain Mapping*, *37*

(6), 2161–2172. https://doi.org/10.1002/hbm.23164

Friston, K. J., Glaser, D. E., Henson, R. N. A., Kiebel, S., Phillips, C., & Ashburner, J.

(2002). Classical and Bayesian Inference in Neuroimaging: Applications.

NeuroImage, *16* (2), 484–512. https://doi.org/10.1006/nimg.2002.1091

Gati, J. S., Menon, R. S., Uğurbil, K., & Rutt, B. K. (1997). Experimental determination

of the BOLD field strength dependence in vessels and tissue. *Magnetic*

Resonance in Medicine, *38* (2), 296–302.

https://doi.org/10.1002/mrm.1910380220

Glover, G. H., & Law, C. S. (2001). Spiral-in/out BOLD fMRI for increased SNR and

reduced susceptibility artifacts. *Magnetic Resonance in Medicine*, *46* (3), Article

3. https://doi.org/10.1002/mrm.1222

Gonzalez-Castillo, J., Panwar, P., Buchanan, L. C., Caballero-Gaudes, C., Handwerker,

D. A., Jangraw, D. C., Zachariou, V., Inati, S., Roopchansingh, V., Derbyshire, J.

A., & Bandettini, P. A. (2016). Evaluation of multi-echo ICA denoising for task

based fMRI studies: Block designs, rapid event-related designs, and cardiac-

gated fMRI. *NeuroImage*, *141*, 452–468.

https://doi.org/10.1016/j.neuroimage.2016.07.049

Gorgolewski, K., Burns, C., Madison, C., Clark, D., Halchenko, Y., Waskom, M., &

Ghosh, S. (2011). Nipype: A Flexible, Lightweight and Extensible Neuroimaging

Data Processing Framework in Python. *Frontiers in Neuroinformatics*, *5*.

https://www.frontiersin.org/articles/10.3389/fninf.2011.00013

Gottfried, J. A., O'Doherty, J., & Dolan, R. J. (2003). Encoding Predictive Reward Value

in Human Amygdala and Orbitofrontal Cortex. *Science*, *301* (5636), 1104–1107.

https://doi.org/10.1126/science.1087919

Graham, S. J., Stanchev, P. L., & Bronskill, M. J. (1996). Criteria for analysis of

multicomponent tissue T2 relaxation data. *Magnetic Resonance in Medicine*, *35*

(3), 370–378. https://doi.org/10.1002/mrm.1910350315

Green, B. G., Dalton, P., Cowart, B., Shaffer, G., Rankin, K., & Higgins, J. (1996).

Evaluating the 'Labeled Magnitude Scale' for Measuring Sensations of Taste and

Smell. *Chemical Senses*, *21* (3), 323–334.

https://doi.org/10.1093/chemse/21.3.323

Green, B. G., Shaffer, G. S., & Gilmore, M. M. (1993). Derivation and evaluation of a

semantic scale of oral sensation magnitude with apparent ratio properties.

Chemical Senses, *18* (6), 683–702. https://doi.org/10.1093/chemse/18.6.683

Howard, J. D., & Gottfried, J. A. (2014). Configural and Elemental Coding of Natural

Odor Mixture Components in the Human Brain. *Neuron*, *84* (4), Article 4.

https://doi.org/10.1016/j.neuron.2014.10.012

Howard, J. D., Plailly, J., Grueschow, M., Haynes, J.-D., & Gottfried, J. A. (2009). Odor

quality coding and categorization in human posterior piriform cortex. *Nature*

Neuroscience, *12* (7), Article 7. https://doi.org/10.1038/nn.2324

Im, K., Jo, H. J., Mangin, J.-F., Evans, A. C., Kim, S. I., & Lee, J.-M. (2010). Spatial

Distribution of Deep Sulcal Landmarks and Hemispherical Asymmetry on the

Cortical Surface. *Cerebral Cortex*, *20* (3), 602–611.

https://doi.org/10.1093/cercor/bhp127

Jones, L. V., Peryam, D. R., & Thurstone, L. L. (1955). Development of a scale for

measuring soldiers' food preferences. *Journal of Food Science*, *20* (5), 512–520.

https://doi.org/10.1111/j.1365-2621.1955.tb16862.x

Jones, T. B., Bandettini, P. A., Kenworthy, L., Case, L. K., Milleville, S. C., Martin, A., &

Birn, R. M. (2010). Sources of group differences in functional connectivity: An

investigation applied to autism spectrum disorder. *NeuroImage*, *49* (1), 401–414.

https://doi.org/10.1016/j.neuroimage.2009.07.051

Kelly, R. E., Alexopoulos, G. S., Wang, Z., Gunning, F. M., Murphy, C. F., Morimoto, S.

S., Kanellopoulos, D., Jia, Z., Lim, K. O., & Hoptman, M. J. (2010). Visual

inspection of independent components: Defining a procedure for artifact removal

from fMRI data. *Journal of Neuroscience Methods*, *189* (2), 233–245.

https://doi.org/10.1016/j.jneumeth.2010.03.028

Kundu, P., Brenowitz, N. D., Voon, V., Worbe, Y., Vértes, P. E., Inati, S. J., Saad, Z. S.,

Bandettini, P. A., & Bullmore, E. T. (2013). Integrated strategy for improving

functional connectivity mapping using multiecho fMRI. *Proceedings of the*

National Academy of Sciences, *110* (40), 16187–16192.

https://doi.org/10.1073/pnas.1301725110

Kundu, P., Inati, S. J., Evans, J. W., Luh, W.-M., & Bandettini, P. A. (2012a).

Differentiating BOLD and non-BOLD signals in fMRI time series using multi-echo

EPI. *NeuroImage*, *60* (3), 1759–1770.

https://doi.org/10.1016/j.neuroimage.2011.12.028

Kundu, P., Inati, S. J., Evans, J. W., Luh, W.-M., & Bandettini, P. A. (2012b).

Differentiating BOLD and non-BOLD signals in fMRI time series using multi-echo

EPI. *NeuroImage*, *60* (3), 1759–1770.

https://doi.org/10.1016/j.neuroimage.2011.12.028

Kundu, P., Voon, V., Balchandani, P., Lombardo, M. V., Poser, B. A., & Bandettini, P. A.

(2017). Multi-echo fMRI: A review of applications in fMRI denoising and analysis

of BOLD signals. *NeuroImage*, *154*, 59–80.

https://doi.org/10.1016/j.neuroimage.2017.03.033

Levene, H. (1960). Robust tests for equality of variances. *Contributions to Probability*

and Statistics, 278–292.

Lim, J., Wood, A., & Green, B. G. (2009). Derivation and Evaluation of a Labeled

Hedonic Scale. *Chemical Senses*, *34* (9), 739–751.

https://doi.org/10.1093/chemse/bjp054

Lynch, C. J., Power, J. D., Scult, M. A., Dubin, M., Gunning, F. M., & Liston, C. (2020).

Rapid Precision Functional Mapping of Individuals Using Multi-Echo fMRI. *Cell*

Reports, *33* (12), 108540. https://doi.org/10.1016/j.celrep.2020.108540

Mai, J. K., Majtanik, M., & Paxinos, G. (2016). *Atlas of the human brain* (4th edition).

Academic Press.

Manera, A. L., Dadar, M., Fonov, V., & Collins, D. L. (2020). CerebrA, registration and

manual label correction of Mindboggle-101 atlas for MNI-ICBM152 template.

Scientific Data, *7* (1), Article 1. https://doi.org/10.1038/s41597-020-0557-9

McKeown, M. J., Hansen, L. K., & Sejnowsk, T. J. (2003). Independent component

analysis of functional MRI: What is signal and what is noise? *Current Opinion in*

Neurobiology, *13* (5), 620–629. https://doi.org/10.1016/j.conb.2003.09.012

Mckeown, M. J., Makeig, S., Brown, G. G., Jung, T.-P., Kindermann, S. S., Bell, A. J., &

Sejnowski, T. J. (1998). Analysis of fMRI data by blind separation into

independent spatial components. *Human Brain Mapping*, *6* (3), 160–188.

https://doi.org/10.1002/(SICI)1097-0193(1998)6:3<160::AID-HBM5>3.0.CO;2-1

Miao, X., Paez, A. G., Rajan, S., Cao, D., Liu, D., Pantelyat, A. Y., Rosenthal, L. I., van

Zijl, P. C. M., Bassett, S. S., Yousem, D. M., Kamath, V., & Hua, J. (2021).

Functional Activities Detected in the Olfactory Bulb and Associated Olfactory

Regions in the Human Brain Using T2-Prepared BOLD Functional MRI at 7T.

Frontiers in Neuroscience, *15*.

https://www.frontiersin.org/articles/10.3389/fnins.2021.723441

Moeller, S., Pisharady, P. K., Ramanna, S., Lenglet, C., Wu, X., Dowdle, L., Yacoub, E.,

Uğurbil, K., & Akçakaya, M. (2021). NOise reduction with DIstribution Corrected

(NORDIC) PCA in dMRI with complex-valued parameter-free locally low-rank

processing. *NeuroImage*, *226*, 117539.

https://doi.org/10.1016/j.neuroimage.2020.117539

Parrish, T., Chen, Y., Li, W., Howard, J., & Gottfried, J. (2008). High resolution 3D

functional images of the human olfactory bulb using passband SSFP at 3T.

Proceedings of International Society of Magnetic Resonance in Medicine, 3567.

Patin, A., & Pause, B. M. (2015). Human amygdala activations during nasal

chemoreception. *Neuropsychologia*, *78*, 171–194.

https://doi.org/10.1016/j.neuropsychologia.2015.10.009

Perszyk, E. E., Davis, X. S., & Small, D. M. (2023). Olfactory decoding is positively

associated with ad libitum food intake in sated humans. *Appetite*, *180*, 106351.

https://doi.org/10.1016/j.appet.2022.106351

Plailly, J., Howard, J. D., Gitelman, D. R., & Gottfried, J. A. (2008). Attention to Odor

Modulates Thalamocortical Connectivity in the Human Brain. *Journal of*

Neuroscience, *28* (20), 5257–5267. https://doi.org/10.1523/JNEUROSCI.5607-

07.2008

Poser, B. A., Versluis, M. J., Hoogduin, J. M., & Norris, D. G. (2006). BOLD contrast

sensitivity enhancement and artifact reduction with multiecho EPI: Parallel-

acquired inhomogeneity-desensitized fMRI. *Magnetic Resonance in Medicine*, *55*

(6), 1227–1235. https://doi.org/10.1002/mrm.20900

Posse, S., Wiese, S., Gembris, D., Mathiak, K., Kessler, C., Grosse-Ruyken, M.-L.,

Elghahwagi, B., Richards, T., Dager, S. R., & Kiselev, V. G. (1999).

Enhancement of BOLD-contrast sensitivity by single-shot multi-echo functional

MR imaging. *Magnetic Resonance in Medicine*, *42* (1), 87–97.

https://doi.org/10.1002/(SICI)1522-2594(199907)42:1<87::AID-

MRM13>3.0.CO;2-O

Raithel, C. U., Miller, A. J., Epstein, R. A., Kahnt, T., & Gottfried, J. A. (2023).

Recruitment of grid-like responses in human entorhinal and piriform cortices by

odor landmark-based navigation. *Current Biology*, *33* (17), 3561-3570.e4.

https://doi.org/10.1016/j.cub.2023.06.087

Raj, D., Anderson, A. W., & Gore, J. C. (2001). Respiratory effects in human functional

magnetic resonance imaging due to bulk susceptibility changes. *Physics in*

Medicine & Biology, *46* (12), 3331. https://doi.org/10.1088/0031-9155/46/12/318

Raj, D., Paley, D. P., Anderson, A. W., Kennan, R. P., & Gore, J. C. (2000). A model for

susceptibility artefacts from respiration in functional echo-planar magnetic

resonance imaging. *Physics in Medicine & Biology*, *45* (12), 3809.

https://doi.org/10.1088/0031-9155/45/12/321

Roth, Z. N., Ryoo, M., & Merriam, E. P. (2020). Task-related activity in human visual

cortex. *PLOS Biology*, *18* (11), e3000921.

https://doi.org/10.1371/journal.pbio.3000921

Savic, I. (2002). Imaging of brain activation by odorants in humans. *Current Opinion in*

Neurobiology, *12* (4), 455–461. https://doi.org/10.1016/S0959-4388(02)00346-X

Savic, I., Gulyás, B., & Berglund, H. (2002). Odorant differentiated pattern of cerebral

activation: Comparison of acetone and vanillin. *Human Brain Mapping*, *17* (1),

17–27. https://doi.org/10.1002/hbm.10045

Schutz, H. G., & Cardello, A. V. (2001). A Labeled Affective Magnitude (lam) Scale for

Assessing Food Liking/Disliking1. *Journal of Sensory Studies*, *16* (2), 117–159.

https://doi.org/10.1111/j.1745-459X.2001.tb00293.x

Shanahan, L. K., Bhutani, S., & Kahnt, T. (2021). Olfactory perceptual decision-making

is biased by motivational state. *PLOS Biology*, *19* (8), e3001374.

https://doi.org/10.1371/journal.pbio.3001374

Stone, H., Bleibaum, R. N., & Thomas, H. A. (2020). *Sensory evaluation practices*.

Academic press.

Swarup, S., Huang, W., Mackay, T. F. C., & Anholt, R. R. H. (2013). Analysis of natural

variation reveals neurogenetic networks for Drosophila olfactory behavior.

Proceedings of the National Academy of Sciences, *110* (3), 1017–1022.

https://doi.org/10.1073/pnas.1220168110

Triantafyllou, C., Wald, L. L., & Hoge, R. D. (2011). Echo-Time and Field Strength

Dependence of BOLD Reactivity in Veins and Parenchyma Using Flow-

Normalized Hypercapnic Manipulation. *PLOS ONE*, *6* (9), e24519.

https://doi.org/10.1371/journal.pone.0024519

Van, A. N., Tisdall, M. D., Laumann, T. O., Montez, D. F., Fair, D. A., & Dosenbach, N.

U. F. (2023). Framewise distortion correction in Multi-echo Echo Planar Imaging.

Proceedings of International Society of Magnetic Resonance in Medicine.

Van de Moortele, P.-F., Pfeuffer, J., Glover, G. H., Ugurbil, K., & Hu, X. (2002).

Respiration-induced B0 fluctuations and their spatial distribution in the human

brain at 7 Tesla. *Magnetic Resonance in Medicine*, *47* (5), 888–895.

https://doi.org/10.1002/mrm.10145

van der Zwaag, W., Francis, S., Head, K., Peters, A., Gowland, P., Morris, P., & Bowtell,

R. (2009). fMRI at 1.5, 3 and 7 T: Characterising BOLD signal changes.

NeuroImage, *47* (4), 1425–1434.

https://doi.org/10.1016/j.neuroimage.2009.05.015

Vizioli, L., Moeller, S., Dowdle, L., Akçakaya, M., De Martino, F., Yacoub, E., & Uğurbil,

K. (2021). Lowering the thermal noise barrier in functional brain mapping with

magnetic resonance imaging. *Nature Communications*, *12* (1), Article 1.

https://doi.org/10.1038/s41467-021-25431-8

Weiskopf, N., Hutton, C., Josephs, O., & Deichmann, R. (2006). Optimal EPI

parameters for reduction of susceptibility-induced BOLD sensitivity losses: A

whole-brain analysis at 3 T and 1.5 T. *NeuroImage*, *33* (2), 493–504.

https://doi.org/10.1016/j.neuroimage.2006.07.029

Yekutieli, D., & Benjamini, Y. (1999). Resampling-based false discovery rate controlling

multiple test procedures for correlated test statistics. *Journal of Statistical

Planning and Inference*, *82* (1), 171–196. https://doi.org/10.1016/S0378-

3758(99)00041-5

Zhou, G., Lane, G., Cooper, S. L., Kahnt, T., & Zelano, C. (2019). Characterizing

functional pathways of the human olfactory system. *eLife*, *8*, e47177.

https://doi.org/10.7554/eLife.47177