Genomic evidence for supergene control of Darwin's "complex marriage arrangement" – the tristylous floral polymorphism

Haoran Xue[1,2], Yunchen Gong[1,3], Stephen I. Wright[1,3*], and Spencer C.H. Barrett[1*]

* These authors contributed equally to the work

1. Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, Ontario M5S 3B2, Canada

2. Institute for Biochemistry and Biology, University of Potsdam, 14476 Potsdam-Golm, Germany

3. Centre for Analysis of Genome Evolution and Function, University of Toronto, Toronto, Ontario M5S 3B2, Canada

Corresponding author: Haoran Xue, haoran.xue@mail.utoronto.ca

Keywords: tristyly, plants, genomics, supergene

Running head: Supergene control of tristyly

Abstract

Tristyly is a sexual polymorphism characterized by three flower morphs with reciprocal stigma

and anther heights controlled by two epistatically interacting diallelic loci (*S* and *M*),

hypothesized to be supergenes. Chromosome-level genome assemblies of *Eichhornia paniculata*

identified the *S*- and *M*-loci. The *S*-locus is a supergene consisting of two divergent alleles: The

S-allele (2.51 Mb), with three *S*-allele specific genes hemizygous in most S-morph plants and the

s-allele (596 kb) with five *s*-allele specific genes. Two of the *S*-allele specific genes, LAZY1-S

and HRGP-S, were specifically expressed in styles and stamens, respectively, making them

tristyly candidate genes. The *M*-locus contains one gene (LAZY1-M) present in the *M*-allele but

absent from the *m*-allele and which is homologous to LAZY1-S. Estimates of allele age are

consistent with theory predicting that the *S*-locus evolved before the *M*-locus. Re-use of the same

gene family highlights the potential role of gene duplication in the evolution of epistatic

multilocus polymorphisms.

Introduction

Flowering plants possess a remarkable diversity of floral strategies promoting cross-pollination

and outcrossed mating. This variation attracted the attention of Charles Darwin who devoted

most of his classic book "*The Different Forms of Flowers on Plants of the Same Species*"[1] to the

floral polymorphism heterostyly in which populations are characterized by two (distyly) or three

(tristyly) mating groups (floral morphs). The defining feature of heterostyly is that the floral

morphs possess a reciprocal arrangement of stigma and anther heights[2], a condition referred to as

reciprocal herkogamy[3]. However, the floral morphs usually also differ in a range of other traits

involving pollen and stigmas and these collectively make up the heterostylous syndrome[2].

Darwin described heterostyly as a 'complex marriage arrangement' promoting cross-pollination

between the floral morphs mediated by animal pollinators owing to segregated pollen placement

on their bodies (see Fig. 1 in Ref. 4). Contemporary experimental evidence on the pollination

biology of diverse heterostylous species generally supports this cross-promotion hypothesis, but

with the additional benefit that reciprocal herkogamy also limits sexual interference between

female and male function thus reducing pollen wastage[4-5]. Heterostyly is reported from 28

animal-pollinated angiosperm families[3, 6] and represents a remarkable example of adaptive

convergence in form and function maintained in populations by negative frequency-dependent

mating.

Studies on the genetics of heterostyly began with the classic early works of Hildebrand,

Darwin, Bateson and Gregory, and Barlow. Later, controlled crosses on over a dozen unrelated

species firmly established that distyly was governed by a single diallelic Mendelian locus (S)

with dominance, whereas tristyly was controlled by two diallelic epistatically interacting loci (S,

M), both with dominance[7] (Fig. 1). To resolve the apparent discrepancy between the complexity

of the heterostylous syndrome and its relatively simple inheritance it was proposed that the

heterostyly loci are in fact supergenes comprised of clusters of functionally related genes[8].

Evidence indicates that these genes are tightly linked to each other due to suppression of

recombination and passed from generation to generation as a single unit[7].

Supergenes are also known to control several other complex polymorphic traits, including

Batesian mimicry in *Heliconius* butterflies, social organizations of fire ants and plumage

variation and behaviour in white-throated sparrows and ruff[9-11]. Of particular importance for

understanding the evolution of supergenes is to determine their genetic architecture and the

mechanisms (e.g., inversions, hemizygosity, epigenetic modifications) that cause suppression of

recombination and thus the maintenance of these linkage groups. Identification of the causal loci

can also provide important insights into the age of the alleles, to assess the evidence for long-

term balanced polymorphisms maintaining variation.

Recent efforts have been made to elucidate the genetic architecture of heterostylous

supergenes in a handful of distylous systems (reviewed in[12-13]). Significantly, genomic studies on

Gelsemium, *Linum*, *Nymphoides*, *Primula,* and *Turnera*, each belonging to a different

angiosperm lineage, have demonstrated that distyly is controlled by several hemizygous genes

present in the short-styled morph (S-morph) but absent from the long-styled morph (L-morph)[14-

19]. For example, in *Primula*, the *S*-locus is a hemizygous region containing five genes controlling

the expression of different components of the distylous polymorphism[14, 15, 20]. A feature shared

by the five unrelated distylous genera is that the style in the S-morph is short in length because

the S-morph carries a hemizygous gene that degrades brassinosteroid (BR). This finding thus

serves as an excellent example of convergent evolution at both the molecular and phenotypic

levels.

All molecular investigations of heterostyly have focused on distylous species and the

supergene hypothesis has yet to be investigated in any of the six tristylous angiosperm families

in which this more complex floral polymorphism occurs[21-22]. Indeed, Charlesworth[23] suggested

that alternatives to the supergene hypothesis should be considered for tristyly because of the

absence of direct empirical evidence for supergenes in trimorphic species. Thus, the molecular

genetic basis and evolutionary assembly of this two-locus epistatically interacting genetic system

remains unclear. The primary motivation for our study was therefore to investigate the genomic

architecture of a tristylous species to evaluate the supergene hypothesis, determine whether

hemizygosity is also a feature of the loci governing sex-organ polymorphism, and to clarify the

genetic basis of the epistatic interaction between the *S*- and *M*-locus.

Here, we generate chromosome-level *de novo* genome assemblies with long-read

sequences and integrate them with data from population genomic and transcriptome analysis to

investigate the genetic architecture of tristyly in *Eichhornia paniculata*, an annual aquatic native

to the Neotropics. Earlier molecular studies using linkage mapping and a short-read genome

assembly demonstrated that the *S*- and *M*-loci in this species are closely linked (separated by a

map distance of 2.7 cM), and that the *M*-locus is located on linkage group 5 in a large region

containing 204 genes[24]. Bulk segregation analysis (BSA) identified 334 genes containing SNPs

potentially linked to the *M*-locus, 24 of which were in this interval. In this earlier work the

genomic location of the *S*-locus was not investigated. These early findings therefore do not

resolve whether the *M*-locus is a supergene or merely contains one pleiotropic gene[12].

To investigate this issue and provide novel insight into the molecular basis of tristyly here

we addressed the following questions: 1) Where are the *S*- and *M*-loci located in the genome? 2)

What is their genomic architecture and are they indeed supergenes? 3) What are the likely

candidate genes at the two loci controlling different components of the tristylous floral

polymorphism? 4) What is evolutionary timing of the origins of the *S*- and *M*-loci during the

assembly of the tristylous polymorphism? We note that in the only theoretical analysis of the

evolution of tristyly the origin of the *S*-locus was inferred to have preceded the origin of the *M*-

locus[23]. As we show below, our results are consistent with this stepwise origin of the two loci,

with recurrent gene duplication from the same ancestral gene playing a key role in the formation

of the epistatic interaction between the two loci.

Results

Comparisons of chromosome-level genome assemblies

We generated chromosome-level genome assemblies from a heterozygous S-morph individual

(*SsMm*) and an L morph individual (*ssmm*) combining long-read PacBio sequencing with

proximity ligation (Omni-C and Hi-C) scaffolding (see methods and Fig. S1). In both assemblies,

most of the two genomes were assembled into eight major chromosomal scaffolds (Table 1).

Comparisons of the two assemblies demonstrated large-scale concordance, with small-scale

differences likely reflecting inversions and/or small assembly errors (Fig. S2). There was clear

evidence for a recent whole genome duplication event in which each chromosome showed strong

synteny with one or more additional chromosomes. Based on the distribution of synonymous

substitution rate (*Ks*) values between paralogs, the most recent whole genome duplication event

has a mode of *Ks*=0.38 (Fig. S3). Assuming a substitution rate of 7×10^{-9} per year[25] and a whole

genome duplication event, this would correspond to two sets of homeologous chromosomes that

diverged roughly 30 million years ago. Because the base chromosome number in Pontederiaceae

has been inferred to be $n=8$[26], the duplication event was likely shared across the family, with

additional lineage-specific duplications due to additional rounds of polyploidy occurring in other

lineages of the family[27]. Unless otherwise noted, our analysis below uses our *SsMm* assembly as

the reference genome.

The Marey maps (plots of the linkage map against the physical map) indicated that there

was a large (likely pericentromeric) region of low recombination in the middle of each

chromosome (Fig. 2). In chromosomes 3 and 7, the region extended to one end of the

chromosome, likely corresponding to submetacentric or acrocentric chromosomes, whereas the

remaining chromosomes appeared to be more metacentric. As expected, the regions of reduced

recombination were associated with a high repeat density (particularly LTR retrotransposons)

and low gene density, whereas the more highly recombining chromosome tips were gene-dense

and comprised of fewer repeats.

Both tristyly loci occur in a large low-recombining region

Using a genome-wide association study (GWAS) we identified 4,640 SNPs that were

significantly associated with the S-morph (*S*---) versus non-S-morphs (*ss*--, i.e. L- and M-

morphs) ($p < 2.52397779$ x 10^{-8}, i.e. Bonferroni correction with $\alpha = 0.05$, Fig. 3a). 96.8% of

these SNPs occurred in a very large 63.1 Mb region on chromosome 2, embedded in an extensive

region of reduced recombination on this chromosome (Fig. 3a). As expected from earlier genetic

work[31], the *M*-locus region is partially linked to the *S*-locus. We found 4,541 SNPs significantly

associated with the M- (*ssM*-) versus L- (*ssmm*) morphs ($p < 2.23826411$ x 10^{-8}, i.e. Bonferroni

correction with $\alpha = 0.05$, Fig. 3b). 98.0% of these SNPs were in a 35.1 Mb region on

chromosome 2, at the other end of the region of reduced recombination from the *S*-locus region.

Thus, both the *S*- and *M*-loci governing tristyly occur in a large, putatively pericentromeric

region of low recombination on chromosome 2 causing extensive and significant SNP

associations across many Mb of sequence.

Elevated nucleotide diversity (π) and Tajima's *D* are expected in regions under balancing

selection[28]. However, contrary to this prediction, neither π nor Tajima's *D* were elevated in the *S*-

or *M*-locus associated regions of chromosome 2 (Fig. S2). One possible explanation is that

whereas recombination rates are low across the regions with the *S*- and *M*- loci creating

contemporary SNP associations over a large region, they may be high enough to have uncoupled

diversity across most of these regions from the causal tristyly loci over long evolutionary

timescales (see below).

The *S*-locus is a supergene consisting of divergent alleles with hemizygous genes

The *S*- and *M*-locus associated regions we identified were both extremely large with very low

rates of recombination. To further narrow down possible candidate genes and to investigate the

potential hemizygous regions associated with tristyly, we identified regions with significant

differences in read coverage between the floral morphs.

167   We identified 50 consecutive 50-kb windows that were specific to the *S*-allele (total

length: 2.5 Mb; genome coordinates: Chr2:114,300,000-116,800,000), with read coverage

significantly higher in 20 S-morph (*S*---) plants than 40 non-S-morph (*ss*--) plants (criteria: the

ratio between the average read coverage of non-S-morph plants and S-morph plants was lower

than 0.4, with the *p* value of the *t*-test lower than $9.73217066 \times 10^{-7}$, i.e. Bonferroni correction

with $\alpha = 0.05$, Fig. 4a). Downstream adjacent to this 2.5 Mb region were 13 consecutive 50-kb

windows (total length: 650 kb; genome coordinates: Chr2:116,800,000-117,450,000) that were

specific to the *s*-allele and had read coverage significantly lower in 20 S-morph (*S*---) plants than

40 non-S-morph (*ss*--) plants (criteria: the ratio between the average read coverage of non-S-

morph plants and S-morph plants was higher than 1.6, with the *p* value of the *t*-test lower than *p*

$< 9.73217066 \times 10^{-7}$, i.e. Bonferroni correction with $\alpha = 0.05$, Fig. 4a).

178   We hypothesized that given the high divergence between alleles, our collapsed haploid

179   genome assembly may have created a chimeric assembly in this region, leaving the *s*-allele

180   sequence adjacent to the *S*-allele region in our assembled scaffold and contributing to the lack of

181 a signal of high diversity when mapping to this reference genome. To investigate this hypothesis

182 further, we compared our L-morph assembly (*ss*) to this assembly in this region using whole

183 genome alignments. As expected, Anchorwave[29] alignments revealed the presence of a large

184 2.57 Mb deletion in the *ssmm* assembly relative to the *SsMm* assembly (Fig. S3), whereas the

185 adjacent non-S morph specific region showed homology. Thus, we conclude that the 2.5 Mb

186 region roughly corresponds to the *S*-allele and the 650 kb region corresponds to the *s*-allele. We

187 refined the boundaries of the two alleles using the coordinates of the closest genes in their

188 flanking regions (Fig. 4a).

189   Two genes (TIE/SPL and MPC2) were shared between the *S*- and *s*-alleles. The order of

190 the two genes are opposite in the two alleles. In the *S*-allele, both of the genes are on the negative

191 strand, whereas in the *s*–allele both are on the positive strand. This finding suggests the existence

192 of an inversion containing the two genes (Fig. 4). The rest of the genes were allele specific: three

193 genes (CYCD6;1-S, HRGP-S, and LAZY1-S) were specific to the *S*-allele, while five genes

194 (Epg15546, Epg15547, CSU1-s, PHOS32-s, and CDC20-1-s) were specific to the *s*-allele.

195 LAZY1 is a gene family that regulates the distribution of auxin[30], HRGP is a superfamily of

196 genes that forms covalently cross-linked networks in primary cell walls[31], and CYCD6;1 is a

197 gene that regulates the formative division of cells[32].

199 **The *M*-locus contains one hemizygous gene**

200 We identified five consecutive 50-kb windows (total length: 250 kb; genome coordinates:

201 Chr2:13,000,000-13,250,000) that had read coverage significantly higher in the 20 M-morph

202 (*ssM-*) plants than in the 20 L-morph (*ssmm*) plants (criteria: the ratio between the average read

203 coverage of non-S-morph plants and S-morph plants was higher than 1.6, with the *p* value of the

t-test lower than $p < 9.73217066 \times 10^{-7}$, i.e. Bonferroni correction with $\alpha = 0.05$, Fig. 4b). There

were nine coding genes in these five windows. However, among them only one gene was

missing from the *ssmm* genome assembly suggesting that it is absent from the *m*-allele (Fig. 4b).

Strikingly, this absent gene, LAZY1-M, also shows strong similarity to genes in the LAZY1

family, raising the possibility that this locus acts competitively in the same pathway as LAZY1-

S. Consistent with the SNP association mapping, the hemizygous *M*-locus region was located on

the other side of the low-recombing region on chromosome 2 (Fig. 3).

Genetic degeneration at the *S*-allele

In species with supergenes, signatures of genetic degeneration, including gene loss and

accumulation of TEs, are expected for the dominant haplotypes of the supergenes, because

dominant haplotypes are only present in heterozygotes and thus do not exhibit recombination[33].

The *S*-allele specific gene loss (the five *s*-allele specific genes) was in accord with this

prediction. In addition, we found that the density of repetitive elements in the *S*-allele, which is

only present in *Ss--* heterozygotes, was significantly higher ($p = 0.000076$) than the flanking

regions (Fig. 4f), whereas the differences between the *s*-allele or the *M*-allele and the flanking

regions were not significantly different ($p < 0.911797$ and $p < 0.075267$, respectively) (Fig. 4g

and h).

The expression pattern of hemizygous genes

Among the three *S*-allele specific genes, LAZY1-S and HRGP-S were expressed in floral

organs of S-morph plants but not non-S-morph plants. LAZY1-S was expressed in styles but not

stamens, whereas in contrast HRGP-S was expressed in the stamens but not styles (Figure 5a).

CYCD6;1-S has low expression levels in both style and stamen of S-morph plants. Therefore, it

is not considered as a candidate gene in the following discussion. The *M*-allele specific gene

LAZY1-M was expressed in both styles and stamens (Figure 5b).

Genes that were differentially expressed in different morphs

For genes that were differentially expressed genome-wide between floral morphs (see Methods

for details of comparisons and Table S1 for the number of genes that were differentially

expressed in each comparison), three sets of genes showed functional enrichment: (1) genes that

were upregulated in S-morph styles (compared with non-S-morph styles) in early buds (enriched

in three functions), (2) genes that were upregulated in S-morph styles (compared with non-S-

morph styles) in late buds (enriched in one function), and (3) genes that were upregulated in M-

morph styles (compared with L-morph styles) in late buds (enriched in one function) (Table S2;

Note, functions that were subsets of one another were only counted once). However, none of the

five functions inferred had a direct connection to growth or flower organs, suggesting that only a

small number of genes downstream to the *S*- and *M*-locus candidate genes are involved in the

phenotypic differences between the floral morphs. Other differentially expressed genes may

simply be incidental associations.

The origins of the *S* and *M* loci

To further understand the origins of the *S*- and *M*-loci, we searched for additional members of

the gene families for loci that were *S*- and *M*-specific. We identified one additional gene that was

paralogous to LAZY1-S and LAZY1-M. Hereafter, we refer to it as "LAZY1-P1". LAZY1-P1

was located 223 kb downstream to LAZY1-M (Fig. 4b). The closest paralog of HRGP-S was

found on the same chromosome but not in the same region, and the closest paralog of CYCD6;1-

S was found on another chromosome (Fig. 4c). The distance between the three ancestral genes

therefore suggests that the *S*-locus was formed via stepwise duplication rather than segmental

duplication.

Based on our gene trees and assuming that the hemizygous loci represent the derived

gene copies, LAZY1-S duplicated from LAZY1-P1 first, following by subsequent gene

duplication from LAZY1-P1 to form LAZY1-M (Fig. 6). Phylogenetic dating with BEAST

v2.7.6[34] suggested that the divergence between LAZY1-S and LAZY-P1 occurred approximately

52.4 million years ago (95% confidence interval: 44.9-59.7) and that the subsequent divergence

between LAZY1-M and LAZY-P1 was around 43.0 million years ago (95% confidence interval:

36.4-50.0).

Discussion

We show that in tristylous *Eichhornia paniculata,* both the *S-* and *M*-loci are located in a low

recombining region of Chromosome 2. The *S*-locus is comprised of two divergent alleles: the

2.51 Mb *S*-allele with three *S*-allele specific genes that are hemizygous in most S-morph plants,

and the 596 kb *s*-allele with five *s*-allele specific genes. In the *S*-allele we identified two strong

tristyly candidate genes: LAZY1-S and HRGP-S. LAZY1-S was expressed in styles but not

stamens suggesting that it may inhibit style elongation in the S-morph by decreasing auxin

concentration in styles. HRGP-S was expressed in stamens but not styles suggesting that it may

be involved in filament elongation or possibly pollen competition. Our evidence for the *S*-locus

therefore strongly supports its status as a supergene.

We found that the *M*-locus contained only one hemizygous gene – LAZY1-M – and this

gene was hemizygous in --*Mm* plants. Strikingly, this gene is in the same gene family as

LAZY1-S, and suggests the possible role of gene duplication and neofunctionalization in the

origin of the epistatic interaction between the two loci. Unlike LAZY1-S, which was only

expressed in styles, LAZY1-M was expressed in both styles and stamens and with a higher

expression level in stamens. This suggests that while LAZY1-S may only be involved in the

shortening of styles in the S-morph, LAZY1-M may also regulate auxin distribution and organ

growth in both the styles and stamens of the M-morph. Given that LAZY-M is the only *M*-allele-

specific hemizygous gene, the M-locus may not represent a supergene at this stage in its

evolutionary history but that it simply contains LAZY1-M, a pleiotropic modifier.

LAZY1-M was also expressed in the stamens but not the styles of S-morph plants (Fig.

5b), suggesting that LAZY1-S and LAZY1-M may compete for the same transcriptional

activator and that the style expression of LAZY1-M is shut down if LAZY1-S is expressed. The

expression level of LAZY1-S in S-morph styles is also much higher than LAZY1-M in M-morph

styles. For example, at early development stages the average normalized count of LAZY1-M in

samples of M-morph styles is 0.02% of that of LAZY1-S in S-morph styles. At late development

stages, the ratio is 1.9%, indicating that the promoter region of LAZY1-S may have stronger

affinity to the transcriptional activator that they compete for, potentially driving differences in

style length in the S- and M-morphs through different degrees of auxin suppression.

In a theoretical treatment of the evolutionary assembly of tristyly[23], Charlesworth

predicted that the M-morph originated last spreading in dimorphic populations of the L- and S-

morph and eventually giving rise to the tristylous polymorphism. Our results on the relative ages

of LAZY1-S and LAZY1-M are consistent with the hypothesis that the *M*-locus evolved after the

S-locus providing empirical support for this prediction. Our results also provide evidence

suggesting a major role for 're-use' through recurrent duplication of the same gene (LAZY1).

Recruitment of this gene explains the origin of the *M*-locus and birth of the third floral morph in

the tristylous system (Fig. 6) resulting in the morphological and mating complexity that

characterizes Darwin's 'complex marriage arrangement'.

Methods

Study system. Plants used in our study originated from seed collected in wild populations of *E.*

paniculata in N.E. Brazil, Cuba and Mexico. The localities of plant material and details of the

five populations used in our study are presented in Table S4. Information on the reproductive

ecology, mating systems and population genetics of *E. paniculata* are available in Ref. 35 and

references cited therein. We germinated seeds using PRO-MIX Premium Potting Mix (Premier

Tech, Rivière-du-Loup, QC, Canada) and seedlings were then transferred to individual 7 cm

plastic pots filled with the same soil mix and placed in flooded trays in a glasshouse at the

University of Toronto (Toronto, Canada) with temperatures of 25-28°C (day) and 18-21°C

(night) and a 16-hour day length. Plants were then grown and flowered regularly for several

years facilitating crosses and collection of leaves and floral buds for molecular analyses.

Genome assemblies. We conducted two chromosome-level genome assemblies of *E. paniculata*.

The first assembly was obtained from an individual of genotype *SsMm* (inferred from selfed

progeny, Table S3) obtained from an inter-population cross between an S-morph plant (*SsM-*)

from N.E. Brazil and an M-morph plant (*ssMm*) from Cuba (see Table S4 for information on the

two parents). From this individual, we obtained 79 Gb (105X coverage) PacBio CLR sequences

318 (sequenced by Arizona Genomics Institute, Tucson, AZ, USA), 55 Gb (45X coverage) Hi-C

319 (sequenced by Phase Genomics, Seattle, WA, USA) sequences, and 117 Gb (156X coverage)

320 10X Chromium sequences (sequenced by The Hospital for Sick Children, Toronto, ON, Canada).

321 The PacBio sequences were assembled with Canu[36] by the Canadian Center for Computational

322 Genomics (C3G) (Montreal, QC, Canada). The genome was phased with Falcon-Unzip[37], then

323 scaffolded into chromosome-level scaffolds with the Hi-C data by Phase Genomics (Seattle,

324 WA, USA).

325   To identify misjoins and characterize the relationship between the physical distance and

326 the genetic distance on each chromosome, we constructed a linkage map with genotyping-by-

327 sequencing data obtained from a backcross pedigree from an L-morph plant and an M-morph

328 plant, detailed in[24], accessible under Accession no. SRP069136 at the Sequence Read Archive

329 (SRA), (http://www.ncbi.nlm.nih.gov/sra, accessed 17 January 2023), and the associated

330 BioProject alias PRJNA310331). The reads were trimmed with Scythe[38] and Seqtk

331 (https://github.com/lh3/seqtk). We aligned the trimmed reads to the *SsMm* genome assembly

332 with NextGenMap V0.5.5[39], and the BAM files were processed with Picard

333 (http://broadinstitute.github.io/picard) using the four functions FixMateInformation, CleanSam,

334 SortSam, and AddOrReplaceReadGroups. SAMtools[40] was used to exclude the secondary

335 alignments. We constructed the linkage map with the alignment files and the software program

336 Lep-Map3[41]. Markers were split into eight large linkage groups using a logarithm of the odds

337 (LOD) score limit of 70 and a Theta value of 0.25 for the initial split (SeparateChromosomes2).

338 Most of the remaining single markers were put into these groups using a LOD score limit of 30

339 and a LOD difference of 5 (JoinSingles2All). We calculated marker order for each linkage group

340 with OrderMarkers2 (with default settings). The first ten linkage groups contained 2,124

markers. We then used Juicebox[42] to correct the misjoins with the linkage map. The genome was

polished with the 10X Chromium sequences, using the software Pilon[43]. The repetitive elements

were annotated with RepeatModeller V1.0.11[44] and RepeatMasker V4.0.7[45].

The second assembly of *E. paniculata* used an L-morph (*ssmm*) plant from Mexico (see

Table S4) using sequencing and assembly services from Dovetail. 59.6 Gb (~50X coverage)

PacBio HiFi reads were generated and the reads were assembled with hifiasm[46]. The assembly

was then scaffolded with Omni-C data using the HiRise software[47]. We used the linkage map

previously generated to concatenate scaffolds that belong to the same linkage group. We also

annotated TEs with RepeatModeller V1.0.11[44] and RepeatMasker V4.0.7[45]. For both of our

genome assemblies, the distribution of scaffold length was assessed with QUAST[48], and the

completeness of assemblies was assessed with BUSCO (Benchmarking Universal Single-Copy

Orthologs)[49] (Table S5).

Genome-wide associations of single nucleotide polymorphisms (SNPs) and read coverage

(RCs). We extracted genomic DNA from 60 *E. paniculata* plants (20 of each floral morph) from

a population in N.E. Brazil (hereafter "Pop 1", see Table S4), using the QIAGEN DNeasy Plant

Mini Kit (QIAGEN, Germantown, MD, USA). Library preparation and sequencing were

conducted by the McGill University and Génome Québec Innovation Centre (Montréal, QC,

Canada) using the 150-bp paired-end protocol on Illumina HiSeq 4000. The samples had an

average sequencing depth of ~8X.

The Illumina sequencing reads were trimmed with Scythe[38] and Seqtk

(https://github.com/lh3/seqtk). We aligned the trimmed reads to the *SsMm* genome assembly

with NextGenMap V0.5.5[50] and the BAM files were processed with Picard

(http://broadinstitute.github.io/picard/) using the four functions FixMateInformation, CleanSam,

SortSam, and AddOrReplaceReadGroups. We used SAMtools[40] to exclude the secondary

alignments. We used freebayes-parallel[51,52] to call SNPs jointly for all samples. We then filtered

the VCF file with VCFtools[53] and vcflib2[54] following a published guideline[55]. We removed sites

if more than 50% of the individuals had missing data, had a minor allele count of <6, or had a

quality score lower than 30. Genotypes with fewer than 4 reads were treated as "missing" and

individuals with more than 50% missing data were not considered further. Then, sites with strong

within-individual allelic bias at heterozygous sites ($0.01 < AB < 0.25$ or $AB > 0.75$), a

discrepancy in the properly paired status for reads supporting reference or alternate alleles, a low

quality relative to depth, a mean depth that was > 13, a quality score < 30, or that occurred

within 5 bp of an indel, were excluded from our analysis. The resulting VCF file had 7,358,165

filtered variant sites.

We conducted a Genome-Wide Association Study (GWAS) with the variant sites using

GEMMA[56], where a relatedness matrix and a linear mixed model were used and *p*-values of

variant sites were calculated with the Wald test.

To detect hemizygous regions associated with the *S*- and *M*-loci, we conducted another

GWAS with read-coverage, following the pipeline in Ref. 57. We split the reference genome into

25,688 windows, each of which was 50-kb, and calculated the average depth of each window in

individuals using the software mosdepth[58]. For each individual, we normalized the coverage by

dividing the coverage of each window with the average coverage across all windows, then

multiplied it by 2. Therefore, the expected normalized coverage is 1 for hemizygous regions and

2 for the rest of the genome. We conducted two *t*-tests for normalized coverage for each window:

one for the comparison between S-morph versus non-S-morph plants, and the other for the

comparison between the M-morph versus L-morph plants. We identified tristyly allele-specific

windows with the following criteria:

1) p value of the t-test was smaller than $9.73217066 \times 10^{-7}$, i.e. $p < 0.05$ after the

Bonferroni correction.

2) We set thresholds on the ratio between the average read coverage of plants of different

morphs:

For S-allele specific windows: The read coverage of a S-allele is at least 4 times of a s-

allele, i.e. the ratio between the average read coverage of non-S-morph plants and S-morph

plants was lower than 0.4: $\frac{2RC(s)}{RC(S)+RC(s)} = \frac{2 \times 1}{4+1} = 0.4$.

For s-allele specific windows: The read coverage of a s-allele is at least 4 times of a S-

allele, i.e. the ratio between the average read coverage of non-S-morph plants and S-morph

plants was higher than 1.6: $\frac{2RC(s)}{RC(S)+RC(s)} = \frac{2 \times 4}{4+1} = 1.6$

For M-allele specific windows: The read coverage of an M-allele is at least 4 times of an

m-allele, i.e. the ratio between the average read coverage of non-M-morph plants and M-morph

plants was lower than 0.25, assuming all M-morph plants were homozygous: $\frac{2RC(m)}{RC(M)+RC(m)} =$

$\frac{2 \times 1}{4+1} = 0.4$.

For m-allele specific windows: The read coverage of an s-allele is at least 4 times of a M-

allele, i.e. the ratio between the average read coverage of non-M-morph plants and M-morph

plants was higher than 1.6: $\frac{2RC(m)}{RC(M)+RC(m)} = \frac{2 \times 4}{4+1} = 1.6$

Only regions with at least two consecutive windows that were allele-specific were

considered. As a result, a 2.5 Mb S-allele, a 650 kb s-allele, and a 250 kb M-allele were

identified. We split each allele into 10 equal-sized windows (window sizes: S-allele: 250 kb; s-

allele: 65 kb; *M*-allele: 25 kb), calculated the repeat content of each window, then compared it

with the 10 windows of the same size flanking each allele (5 windows upstream and 5 windows

downstream of the locus) with Wilcoxon rank-sum tests.

Differential gene expression analysis and genome annotation with transcriptome data. To

infer genes controlling different components of the tristyly syndrome, we conducted a floral

organ-specific transcriptomic analysis. We dissected floral buds of the three morphs at two

developmental stages: "early", with buds between 3-5 mm in length, and "late", with buds

between 5-8 mm. We used sterilized forceps under a stereo dissecting microscope to open the

floral buds and placed the styles and stamens into 2 mL centrifuge tubes which were then frozen

by putting them into liquid nitrogen. We extracted RNA from the frozen tissues with Spectrum™

Plant Total RNA-Kit (Sigma-Aldrich, St. Louis, MO, USA) following the manufacturer's

instructions. For each type of sample, we collected four biological replicates, each containing

styles or stamens from 5-27 buds sampled from 5-17 individuals from each of two populations

(Pop. 1, 2) originating from N.E. Brazil (Table S4, S6).

Sequencing libraries were prepared and sequenced on the Illumina HiSeq4000 platform

(100-bp paired-end reads) by the McGill University and Génome Québec Innovation Centre

(Montréal, QC, Canada). Then we aligned the Illumina sequences to the *SsMm* genome assembly

with HISAT2[59] and assembled the transcripts with StringTie[60]. We conducted functional

annotations of the floral organ transcriptomes with Mercator4[61]. We used StringTie[60] to estimate

the abundance of each transcript in each sample. Then we used the R package DESeq2[62] to

conduct differential gene expression analysis and plot the normalized count in each sample for

the genes at the *S*- and *M*-loci. We conducted eight differential gene expression analyses for each

type of morph difference (*S* vs. non-S and M vs- L), each tissue (style vs. stamen), and each

development stage (early bud vs. late bud). In each analysis, we identified the genes that are

upregulated in each morph with false discovery rate < 0.05 and LFC (log fold change) > 2 or < -

2. We then did a Mercator BIN enrichment analysis[61] with the list of upregulated genes for each

comparison.

We also annotated the repeat-masked versions of the two genome assemblies with the

floral organ transcriptomes and also leaf and floral bud transcriptomes from Ref. 63, available

under accession number SRP049636 on the Sequence Read Archive (SRA)

(http://www.ncbi.nlm.nih.gov/sra, accessed 20 January 2023) and the associated BioProject alias

is PRJNA266681 (http://www.ncbi.nlm.nih.gov/bioproject/, accessed 20 January 2023), and also

Ref. 64, available under accession number SRP060405 at the Sequence Read Archive and

associated BioProject alias (PRJNA288861) using MAKER v3.01.03[65] and InterProScan 5[66].

This resulted in a total of 35,224 annotated genes for the *SsMm* genome and 29,592 genes for the

ssmm genome.

Origin of the *S* and *M* loci. We aligned the coding sequences of LAZY1-S, HRGP-S, and

LAZY1-M against the *SsMm* and *ssmm* genome assemblies with BLAST[67], to identify the closest

homologs of the three genes. We identified a LAZY1 homolog (LAZY1-P1) that exists in both

SsMm and *ssmm* genome assemblies.

We then downloaded the coding sequences of LAZY1 in *Oryza sativa* and *Zea mays*

from NCBI (https://www.ncbi.nlm.nih.gov/, accession numbers: NM_001138862.1 and

DQ855268.1), and two paralogs of LAZY1 in *Musa acuminata* 'DH-Pahang' from the website

Banana Genome Hub (https://banana-genome-hub.southgreen.fr/). We conducted a multiple

sequence alignment of these sequences and the coding sequences of three LAZY1 genes in *E.*

paniculata, using MAFFT[68].

We then estimated the time of divergence between LAZY1-S and LAZY1-P1 and

between LAZY1-M and LAZY1-P1 with BEAST v2.7.6[34]. First, we did model selection of

jModelTest[69] using Akaike information criterion (AIC) as the criteria. The GTR + I + Gamma

model with nCat = 4 was selected. Then we set the time to the most recent ancestors of *Oryza*

sativa and *Zea mays* (81.43 ± 0.3 MYA)[70], Zingiberales and Commelinales (79.8 ± 9.87

MYA)[71], and Poales and Commelinales (106.7 ± 8.3 MYA)[71] as calibration points. Ten

independent Markov Chain Monte Carlo runs, each with 1×10^8 generations (first 10% was

burn-in) were conducted. We combined the output of the eight runs using LogCombiner v2.7.6[34]

with a sample frequency of 5,000. Then we used TreeAnnotator v2.7.6[34] to generate the

maximum clade credibility tree, which was visualized in FigTree v1.4.4

(http://tree.bio.ed.ac.uk/software/figtree/) (Fig. 6).

Synteny analyses, whole genome alignments and structural variants. We used COGE's

SynMap2 algorithm[72] with the MAKER and InterProScan annotations to examine the synteny

between the two genome assemblies and identify structural variation. To obtain finer-scale whole

genome alignment of syntenic regions to localize putative insertion/deletion differences between

the *S* and *s* alleles and between the *M* and *m* alleles, we used Anchorwave[29] to align the two

genome assemblies, with the "minimap2" option and the "proali" function.

Population genomic statistics. We calculated four population genomic statistics with 1-Mb

sliding windows for the 60 individuals (20 of each morph) used in the GWAS analyses: (1)

Nucleotide diversity (π), (2) Tajima's D, (3) F_{ST} between individuals of the S-morph and the non-

S-morph, and (4) F_{ST} between individuals of the L- and M-morphs. We conducted these

calculations with the filtered VCF file used in the SNP-based GWAS and the functions

"window-pi", "TajimaD", and "weir-fst-pop" in the software program VCFtools[53].

Figure 1 | Tristyly in *Eichhornia paniculata*. a, Flower of the S-morph illustrating exerted

long- and mid-level stamens. The short style is concealed in the perianth tube. **b,** The

arrangement of styles and stamens in the three floral morphs of a tristylous species. L, M and S

refer to the long-, mid- and short-styled morphs, respectively. The arrows indicate "legitimate"

cross-pollinations between anthers and stigmas of equivalent height. In most tristylous species

these are the only compatible matings because of trimorphic incompatibility. However, in *E.*

paniculata "illegitimate" pollinations are permitted as the species is self- and intramorph

compatible[35]. Genotypes for the floral morphs of *E. paniculata* are indicated.

Figure 2 | The recombination landscape and genome content across the chromosomes of

Eichhornia paniculata. **a**. Marey maps plotting the position on the linkage map (cM) against the

position on the physical map (Mb). **b-h,** Densities of genes (**b**), all repetitive elements (**c**), simple

repeats (**d**), Mutator-Like Element (MuLE) (**e**), Long Interspersed Nuclear Element (LINE) (**f**),

Copia elements (**g**), and Ty3 elements (**h**), all measured as proportions of 1-Mb windows.

Figure 3 | SNPs associated with tristyly in *Eichhornia paniculata* and morph-specific read

coverage, recombination landscape, and repeat content of the *S-* and *M*-loci. a, SNPs

associated with S- vs. non-S-morphs. The blue lines indicate $p = 2.52397779$ x 10^{-8}. Inset panels

zoom in on chromosome 2. **b,** SNPs associated with M- vs. L-morphs. The blue lines indicate p

$= 2.23826411$ x 10^{-8}. Inset panels zoom in on chromosome 2. **c,** Average normalized coverages

of the L- (red), M- (blue), and S- (cyan) morph plants at the *S*-locus (*S-* and *s*-alleles assembled

next to each other). The data were based on 20 plants per morph. **d,** The average normalized

coverages of the L- (red), M- (blue), and S- (cyan) morph plants at the *M*-locus. The data were

based on 20 plants per morph. **e.** The recombination landscape of chromosome 2. **f-h,**

Comparisons of proportion of repetitive elements between the *S*-allele (ten 250-kb windows), *s*-

allele (ten 65-kb windows), and *M*-allele (ten 25-kb windows), and their flanking regions (five

windows upstream and five windows downstream to the locus, with window size equal to the

allele being compared with).

Figure 4 | The composition and proposed model of the evolution of the *S*- and *M*-loci in

Eichhornia paniculata. a, Genomic organization of the *S*-locus genes. Black dashed lines

represent the boundaries of the tristyly alleles inferred from the read coverage analysis. Blue

dashed lines represent inferred boundaries of an inversion. **b,** Genomic organization of the genes

around LAZY1-M. Green dashed lines represent the boundaries of LAZY1-M and its paralog

LAZY1-P1. **c,** Hypothesized evolutionary pathway for the organization of the *S*- and *M*-loci.

Figure 5 | Expression patterns of the three genes that are specific to the *S*- or *M*-allele and expressed in the styles or stamens of *Eichhornia paniculata*. a, Genes specific to the *S* allele. b, LAZY1-M, which is specific to the *M* allele. Samples of different morphs are separated by solid lines. For each morph, samples of different floral organs (styles or stamens) are separated by dashed lines. For each combination of morph and floral organ, the samples at the early and late development stages are on the left and right, respectively. Each type of sample had four biological replicates.

Figure 6 | Phylogenetic inference of the dates of divergence between LAZY1-P1 and

LAZY1-S and between LAZY1 and LAZY1-M in *Eichhornia paniculata* and its ancestors.

Bayesian tree of LAZY1 genes from *Oryza sativa*, *Zea mays*, *Musa acuminata*, and *E.*

paniculata. Numbers outside of parentheses represent the estimated time of divergence. Blue

rectangles and numbers in parentheses are the 95% confidence intervals of the estimated time of

divergence.

References

1. Darwin, C. R. *The different forms of flowers on plants of the same species.* (John Murray, 1877).

2. Barrett, S. C. H. Heterostylous genetic polymorphisms: model systems for evolutionary analysis. In *Evolution and function of heterostyly* (ed. Barrett, S. C. H.) 1–29 (Springer, 1992).

3. Lloyd, D. G. & Webb, C. J. The evolution of heterostyly. In *Evolution and function of heterostyly* (ed. Barrett, S. C. H.) 151–178 (Springer, 1992).

4. Barrett, S. C. H. The evolution of plant sexual diversity. *Nat. Rev. Genet.* 3, 274-284 (2002).

5. Lloyd, D. G. & Webb, C. The selection of heterostyly. In *Evolution and function of heterostyly* (ed. Barrett, S. C. H.) 179–207 (Springer, 1992).

6. Barrett, S. C. H., Jesson, L. K. & Baker, A. M. The evolution and function of stylar polymorphisms in flowering plants. *Ann. Bot.* 85, 253-265 (2000).

7. Lewis, D. & Jones, D. The genetics of heterostyly. In *Evolution and function of heterostyly* (Ed. Barrett, S. C. H.) 129–150 (Springer, 1992).

8. Mather, K. The genetical architecture of heterostyly in *Primula sinensis*. *Evolution* 340–352 (1950).

9. Schwander, T., Libbrecht, R. & Keller, L. Supergenes and complex phenotypes. *Current Biology* **24**, 288–294 (2014).

10. Thompson, M. J. & Jiggins, C. Supergenes and their role in evolution. *Heredity* **113**, 1-8 (2014).

11. Maney, D. L. & Küpper, C. Supergenes on steroids. *Philos. Trans. R. Soc. B* 377, 20200507 (2022).

12. Kappel, C., Huu, C. N. & Lenhard, M. A short story gets longer: recent insights into the

molecular basis of heterostyly. *J. Exp. Bot.* 68, 5719-5730 (2017).

13. Barrett, S. C. H. 'A most complex marriage arrangement': recent advances on heterostyly

and unresolved questions. *New Phytol.* 224, 1051-1067 (2019).

14. Huu, C. N. *et al.* Presence versus absence of CYP734A50 underlies the style-length

dimorphism in primroses. *eLife* **5**, e17956 (2016).

15. Li, J. *et al.* Genetic architecture and evolution of the *S* locus supergene in *Primula vulgaris*.

Nat. Plants **2**, 1-7 (2016).

16. Shore, J. S. *et al.* The long and short of the *S*-locus in *Turnera* (Passifloraceae). *New Phytol.*

224, 1316–1329 (2019).

17. Gutiérrez-Valencia, J. *et al.* Genomic analyses of the *Linum* distyly supergene reveal

convergent evolution at the molecular level. *Curr. Biol.* **32**, 4360–4371 (2022).

18. Zhao, Z. *et al.* Genomic evidence supports the genetic convergence of a supergene

controlling the distylous floral syndrome. *New Phytol.* **237**, 601–614 (2023).

19. Yang, J. *et al.* Haplotype-resolved genome assembly provides insights into the evolution of

S-locus supergene in distylous *Nymphoides indica*. *New Phytol.* **240**, 2058-2071(2023).

20. Potente, G. *et al.* Comparative genomics elucidates the origin of a supergene controlling

floral heteromorphism. *Mol. Biol. Evol.* **39**, msac035 (2022).

21. Barrett, S. C. H. The evolutionary biology of tristyly. In *Oxford Surveys in Evolutionary*

Biology (Eds. D. Futuyma & J. Antonovics) vol. 9, 283–326 (Oxford University Press, 1993).

22. Thompson, J. D., Pailler, T., Strasberg, D. & Manicacci, D. Tristyly in the endangered

Mascarene Island endemic *Hugonia serrata* (Linaceae). *Am. J. Bot.* **83**, 1160–1167 (1996).

23. Charlesworth, D. The evolution and breakdown of tristyly. *Evolution* **33**, 489-498. (1979)

24. Arunkumar, R., Wang, W., Wright, S. I. & Barrett, S. C. H. The genetic architecture of

tristyly and its breakdown to self-fertilization. *Mol. Ecol.* **26**, 752–765 (2017).

25. Ossowski, S. *et al.* The rate and molecular spectrum of spontaneous mutations in *Arabidopsis*

thaliana. *Science* **327**, 92–94 (2010).

26. Eckenwalder, J. E. & Barrett, S. C. H. Phylogenetic systematics of Pontederiaceae. *Syst. Bot.*

11, 373–391 (1986).

27. Ness, R. W., Siol, M. & Barrett, S. C. H. *De novo* sequence assembly and characterization of

the floral transcriptome in cross-and self-fertilizing plants. *BMC Genomics* **12**, 298 (2011).

28. Tajima, F. Statistical method for testing the neutral mutation hypothesis by DNA

polymorphism. *Genetics* **123**, 585-595 (1989).

29. Song, B. *et al.* AnchorWave: Sensitive alignment of genomes with high sequence diversity,

extensive structural polymorphism, and whole-genome duplication. *Proc. Natl. Acad. Sci.* **119**,

e2113075119 (2022).

30. Jiao, Z., Du, H., Chen, S., Huang, W. & Ge, L. *LAZY* gene family in plant gravitropism.

Front. Plant Sci. **11**, 606241 (2021).

31. Chen, Y. *et al.* Identification of the abundant hydroxyproline-rich glycoproteins in the root

walls of wild-type *Arabidopsis*, an ext3 mutant line, and its phenotypic revertant. *Plants* **4**, 85–

111 (2015).

32. Sozzani, R., Cui, H., Moreno-Risueno, M. A., Busch, W., Van Norman, J. M., Vernoux, T.,

Brady, S. M., Dewitte, W., Murray, J. A. H. and Benfey, P.N., 2010. Spatiotemporal regulation

of cell-cycle genes by SHORTROOT links patterning and growth. *Nature*, 466, 128-132.

33. Gutiérrez-Valencia, J. *et al.* The genomic architecture and evolutionary fates of supergenes.

Genome Biology and Evolution **13**, evab057 (2021).

34. Bouckaert, R. *et al.* BEAST 2.5: An advanced software platform for Bayesian evolutionary

analysis. *PLoS Comput. Biol*. **15**, e1006650 (2019).

35. Ness, R. W., Wright, S. I. & Barrett, S. C. H. Mating-system variation, demographic history

and patterns of nucleotide diversity in the tristylous plant *Eichhornia paniculata*. *Genetics* **184**,

381–392 (2010).

36. Koren, S. *et al.* Canu: scalable and accurate long-read assembly via adaptive k-mer weighting

and repeat separation. *Genome Res*. **27**, 722–736 (2017).

37. Chin, C. S. *et al.* Phased diploid genome assembly with single-molecule real-time

sequencing. *Nature Methods* **13**, 1050-1054 (2016).

38. Thress, K., Henzel, W., Shillinglaw, W. & Kornbluth, S. Scythe: a novel reaper-binding

apoptotic regulator. *EMBO J.* **17**, 6135–6143 (1998).

39. Sedlazeck, F. J., Rescheneder, P. & Von Haeseler, A. NextGenMap: fast and accurate read

mapping in highly polymorphic genomes. *Bioinformatics* **29**, 2790–2791 (2013).

40. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *GigaScience* **10**, (2021).

41. Rastas, P. Lep-MAP3: robust linkage mapping even for low-coverage whole genome

sequencing data. *Bioinformatics* **33**, 3726–3732 (2017).

42. Durand, N. C. *et al.* Juicebox provides a visualization system for Hi-C contact maps with

unlimited zoom. *Cell Syst.* **3**, 99–101 (2016).

43. Walker, B. J. *et al.* Pilon: an integrated tool for comprehensive microbial variant detection

and genome assembly improvement. *PLoS One* **9**, e112963 (2014).

44. Smit, A. F. A. & Hubley, R. RepeatModeler Open-1.0. 2008–2015. (2008).

45. Smit, A. F. A., Hubley, R. & Green, P. RepeatMasker Open-4.0. (2013).

46. Cheng, H., Concepcion, G. T., Feng, X., Zhang, H. & Li, H. Haplotype-resolved de novo

assembly using phased assembly graphs with hifiasm. *Nat. Methods* **18**, 170–175 (2021).

47. Putnam, N. H. *et al.* Chromosome-scale shotgun assembly using an *in vitro* method for long-

range linkage. *Genome Res.* **26**, 342–350 (2016).

48. Gurevich, A., Saveliev, V., Vyahhi, N. & Tesler, G. QUAST: quality assessment tool for

genome assemblies. *Bioinformatics* **29**, 1072–1075 (2013).

49. Manni, M. et al. BUSCO update: novel and streamlined workflows along with broader and

deeper phylogenetic coverage for scoring of eukaryotic, prokaryotic, and viral genomes. *Mol.*

Biol. Evol. **38**, 4647-4654 (2021).

50. Sedlazeck, F. J., Rescheneder, P. & Von Haeseler, A. NextGenMap: fast and accurate read

mapping in highly polymorphic genomes. *Bioinformatics* **29**, 2790-2791 (2013).

51. Garrison, E. & Marth, G. Haplotype-based variant detection from short-read sequencing.

ArXiv Prepr. ArXiv12073907 (2012).

52. Tange, O. Gnu parallel: the command-line power tool. *Usenix Mag.* **36**, 42 (2011).

53. Danecek, P. et al. The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158

(2011).

54. Garrison, E., Kronenberg, Z. N., Dawson, E. T., Pedersen, B. S. & Prins, P. A spectrum of

free software tools for processing the VCF variant call format: vcflib, bio-vcf, cyvcf2, hts-nim

and slivar. *PLoS Comput. Biol.* **18**, e1009123 (2022).

55. Puritz, J. B., Hollenbeck, C. M. & Gold, J. R. dDocent: a RADseq, variant-calling pipeline

designed for population genomics of non-model organisms. *PeerJ* **2**, e431 (2014).

56. Zhou, X. & Stephens, M. Genome-wide efficient mixed-model analysis for association

studies. *Nat. Genet.* **44**, 821–824 (2012).

57. Xue, L. *et al.* Evidences for a role of two Y-specific genes in sex determination in *Populus*

deltoides. *Nature Communications* **11**, 5893 (2020).

58. Pedersen, B. S. & Quinlan, A. R. Mosdepth: quick coverage calculation for genomes and

exomes. *Bioinformatics* **34**, 867–868 (2018).

59. Kim, D., Langmead, B. & Salzberg, S. L. HISAT: a fast spliced aligner with low memory

requirements. *Nat. Methods* **12**, 357–360 (2015).

60. Pertea, M. *et al.* StringTie enables improved reconstruction of a transcriptome from RNA-seq

reads. *Nat. Biotechnol.* **33**, 290–295 (2015).

61. Lohse, M. *et al.* Mercator: a fast and simple web server for genome scale functional

annotation of plant sequence data. *Plant, Cell & Environment* **37**, 1250-1258 (2014).

62. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biol.* **15**, 1–21 (2014).

63. Arunkumar, R., Ness, R. W., Wright, S. I. & Barrett, S. C. H. The evolution of selfing is

accompanied by reduced efficacy of selection and purging of deleterious mutations. *Genetics*

199, 817–829 (2015).

64. Arunkumar, R., Maddison, T. I., Barrett, S. C. H. & Wright, S. I. Recent mating-system

evolution in *Eichhornia* is accompanied by cis-regulatory divergence. *New Phytol.* **211**, 697–707

(2016).

65. Cantarel, B. L. *et al.* MAKER: an easy-to-use annotation pipeline designed for emerging

model organism genomes. *Genome Res.* **18**, 188–196 (2008).

66. Jones, P. *et al.* InterProScan 5: genome-scale protein function classification. *Bioinformatics*

30, 1236–1240 (2014).

67. Camacho, C. *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421

(2009).

68. Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. *Mol. Biol. Evol.* **30**, 772-780 (2013).

69. Posada, D. jModelTest: Phylogenetic Model Averaging. *Mol. Biol. Evol.* **25**, 1253-1256

(2008).

70. Huang, W. *et al.* A well-supported nuclear phylogeny of Poaceae and implications for the

evolution of C4 photosynthesis. *Mol. Plant* **15**, 755-777 (2022).

71. Magallón, S., Gómez-Acevedo, S., Sánchez-Reyes, L. L. & Hernández-Hernández, T. A

metacalibrated time-tree documents the early rise of flowering plant phylogenetic diversity. *New*

Phytol. **207**, 437-453 (2015).

72. Haug-Baltzell, A., Stephens, S. A., Davey, S., Scheidegger, C. E. & Lyons, E. SynMap2 and

SynMap3D: web-based whole-genome synteny browsers. *Bioinformatics* **33**, 2197–2198 (2017).

Acknowledgements

This work was supported by Discovery Grants from the Natural Sciences and Engineering

Research Council of Canada (NSERC) to S.C.H.B. and S.I.W. and student fellowships from the

University of Toronto to H.X. The Canu assembly of the *SsMm* genome was conducted by the

Canadian Center for Computational Genomics (C3G), which is a Genomics Technology

Platform (GTP) supported by the Canadian Government through Genome Canada. We thank

William Cole and Tom Gludovacz for glasshouse maintenance, Joanna Tran, SungA Choe, Ryan

Ruan, and Jianuo Feng for help with growing plants, Claire Ellis and Po-Yuan Huang for help

with tissue collection, Baharul Choudhury, Iuliia Pelikh, and Yihe Zheng for help with DNA

Author contributions

700 H.X., S.I.W., and S.C.H.B. conceived and designed the study and wrote the manuscript.

701 S.C.H.B. obtained the seeds used for the study. H.X. grew the plants, collected the tissues, and

702 extracted DNA and RNA. Y.G. performed the genome annotation. S.I.W. performed the

703 Anchorwave alignment. H.X. performed all remaining analyses. All authors read and approved

704 the final manuscript.

Competing interests

707 None declared.