Genomic evidence for supergene control of Darwin's "complex marriage arrangement" – the tristylous floral polymorphism

Haoran Xue[1,2] (https://orcid.org/0000-0003-1429-0996), Yunchen Gong[1,3], Stephen I. Wright[1,3*] (https://orcid.org/0000-0001-9973-9697), and Spencer C.H. Barrett[1*] (https://orcid.org/0000-0002-7762-3455)

* These authors contributed equally to the work

1. Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, Ontario M5S 3B2, Canada

2. Institute for Biochemistry and Biology, University of Potsdam, 14476 Potsdam-Golm, Germany

3. Centre for Analysis of Genome Evolution and Function, University of Toronto, Toronto, Ontario M5S 3B2, Canada

Corresponding author: Haoran Xue, haoran.xue@mail.utoronto.ca

Keywords: tristyly, plants, genomics, supergene

Running head: Supergene control of tristyly

Abstract

Tristyly is a polymorphism characterized by three flower morphs with reciprocal stigma and

anther heights controlled by two epistatically interacting diallelic loci (*S* and *M*), hypothesized to

be supergenes. Chromosome-level genome assemblies of *Eichhornia paniculata* identified the *S*-

and *M*-loci. The *S*-locus is a supergene consisting of two divergent alleles: The *S*-allele (2.51Mb)

with three *S*-allele specific genes hemizygous in most S-morph plants and the *s*-allele (596kb)

with five *s*-allele specific genes. Two of the *S*-allele specific genes, LAZY1-S and HRGP-S,

were specifically expressed in styles and stamens, respectively, making them tristyly candidate

genes. The *M*-locus contained one gene (LAZY1-M), homologous to LAZY1-S, that was present

in the *M*-allele but absent from the *m*-allele. Estimates of allele ages are consistent with the

prediction that the *S*-locus evolved before the *M*-locus. Re-use of the same gene family

highlights the potential role of gene duplication in the evolution of epistatic multilocus

polymorphisms.

Flowering plants possess a remarkable diversity of floral strategies promoting cross-pollination

and outcrossed mating. This variation attracted the attention of Charles Darwin who devoted

most of his classic book "*The Different Forms of Flowers on Plants of the Same Species*"[1] to the

floral polymorphism heterostyly in which populations are characterized by two (distyly) or three

(tristyly) floral morphs, which possess a reciprocal arrangement of stigma and anther heights[2].

Darwin described heterostyly as a 'complex marriage arrangement' promoting cross-pollination

between the floral morphs by animal pollinators owing to segregated pollen placement on their

bodies (see Fig. 1 in Ref. 3). Contemporary experimental evidence generally supports this

hypothesis also finding that heterostyly limits sexual interference between female and male

function thus reducing pollen wastage[3-4]. Heterostyly is reported from 28 animal-pollinated

angiosperm families[5-6] and represents a remarkable example of adaptive convergence in form

and function. Controlled crosses on over a dozen unrelated species have established that distyly

is governed by a single diallelic Mendelian locus (S) with dominance, whereas tristyly was

controlled by two diallelic epistatically interacting loci (S, M), both with dominance[7] (Fig. 1). To

resolve the discrepancy between the complexity of the heterostylous syndrome and its relatively

simple inheritance, it has been proposed that heterostyly loci are in fact supergenes comprised of

clusters of functionally related genes[8]. Evidence indicates that these genes are tightly linked to

each other due to suppression of recombination and are inherited as a single unit[7].

Supergenes are also known to control several other complex polymorphic traits, including

Batesian mimicry in *Heliconius* butterflies, social organizations of fire ants, and plumage

variation and behaviour in ruff[9-11]. Of particular importance for understanding the evolution of

supergenes is to determine their genetic architecture and the mechanisms causing suppression of

recombination. Identification of the causal loci can also provide important insights into the age

of the alleles, to assess the evidence for long-term balanced polymorphisms maintaining

variation. Recent efforts have been made to elucidate the genetic architecture of heterostylous

supergenes in a handful of distylous systems (reviewed in[12-13]). Significantly, genomic studies on

Fagopyrum, Gelsemium, *Linum*, *Nymphoides*, *Primula,* and *Turnera*, each belonging to a

different angiosperm lineage, have demonstrated that distyly is controlled by several hemizygous

genes present in the short-styled morph (S-morph) but absent from the long-styled morph (L-

morph)[14-20]. For example, in *Primula*, the S-locus is a hemizygous region containing five genes

controlling the expression of different components of the distylous polymorphism[14-15, 21].

All molecular investigations of heterostyly have focused on distylous species and the

supergene hypothesis has yet to be investigated in any of the six angiosperm families in which

the more complex tristyly occurs[22-23]. Charlesworth[24] suggested that alternatives to the supergene

hypothesis should be considered for tristyly because of the absence of empirical evidence for

supergenes. Thus, the genetic basis and evolutionary assembly of this two-locus epistatically

interacting genetic system remains unclear.

Here, we generate chromosome-level *de novo* genome assemblies with long-read

sequences and integrate them with data from population genomic and transcriptomic analysis to

investigate the genetic architecture of tristyly in *Eichhornia paniculata*, an annual aquatic native

to the Neotropics. Earlier molecular studies demonstrated that the *S*- and *M*-loci in this species

are closely linked (separated by a map distance of 2.7 cM), and that the *M*-locus is located on

linkage group 5 in a region containing 204 genes[25], but the genomic location of the *S*-locus was

not investigated, and they did not resolve whether the *M*-locus is a supergene or merely contains

one pleiotropic gene[12].

To investigate this issue and provide novel insight into the molecular basis of tristyly we

addressed the following questions: 1) What is the genomic architecture of the *S*- and *M*-loci and

are they indeed supergenes? 2) What are the likely candidate genes at the two loci controlling

different components of tristyly? 3) What is the evolutionary timing of the origins of the *S*- and

M-loci? We note that in the only theoretical analysis of the evolution of tristyly, the origin of the

S-locus was inferred to have preceded the origin of the *M*-locus[24]. Our results are consistent with

this stepwise origin of the two loci, with recurrent gene duplication from the same ancestral gene

playing a key role in the formation of the epistatic interaction between the two loci.

Results

Comparisons of chromosome-level genome assemblies

We generated chromosome-level genome assemblies from a heterozygous S-morph individual

(*SsMm*) and an L-morph individual (*ssmm*) combining long-read PacBio sequencing with

proximity ligation (Omni-C and Hi-C) scaffolding (see methods and Fig. S1). In both assemblies,

most of the two genomes were assembled into eight major chromosomal scaffolds (Table 1).

Comparisons of the two assemblies demonstrated large-scale concordance, with small-scale

differences likely reflecting inversions and/or small assembly errors (Fig. S2a). There was clear

evidence for a recent whole genome duplication event in which each chromosome showed strong

synteny with one or more additional chromosomes. Based on the distribution of synonymous

substitution rate (*Ks*) values between paralogs, the most recent whole genome duplication event

has a mode of $Ks=0.38$ (Fig. S2b). Assuming a substitution rate of 7×10^{-9} per year[26] and a

whole genome duplication event, this would correspond to two sets of homeologous

chromosomes that diverged less than roughly 30 million years ago. Since the base chromosome

number in Pontederiaceae was inferred to be $n=8$[27], the duplication event was likely shared

across the family, with additional lineage-specific duplications due to additional rounds of

polyploidy occurring in other lineages of the family[28]. Unless otherwise noted, our analysis

below uses our *SsMm* assembly as the reference genome.

The Marey maps (plots of the linkage map against the physical map) indicated that there

was a large (likely pericentromeric) region of low recombination in the middle of each

chromosome (Fig. 2a). In chromosomes 3 and 7, the region extended to one end of the

chromosome, likely corresponding to submetacentric or acrocentric chromosomes, whereas the

remaining chromosomes appeared to be more metacentric. As expected, the regions of reduced

recombination were associated with a high repeat density (particularly LTR retrotransposons),

low gene density, and low nucleotide diversity, whereas the more highly recombining

chromosome tips were gene-dense, comprised of fewer repeats, and has higher nucleotide

diversity (Fig. 2b-f, Fig. S3).

Both tristyly loci occur in a large low-recombining region

Using a genome-wide association study (GWAS) we identified 4,640 SNPs that were

significantly associated with the S-morph (*S*---) versus non-S-morphs (*ss*--, i.e. L- and M-

morphs) ($p < 2.52397779 \times 10^{-8}$, Fig. 3a). 96.8% of these SNPs occurred in a very large 63.1 Mb

region on chromosome 2, embedded in an extensive region of reduced recombination on this

chromosome (Fig. 3a). As expected from earlier genetic work[25], the *M*-locus region is partially

linked to the *S*-locus. We found 4,541 SNPs significantly associated with the M- (*ssM*-) versus

L- (*ssmm*) morphs ($p < 2.23826411 \times 10^{-8}$, Fig. 3b). 98.0% of these SNPs were in a 35.1 Mb

region on chromosome 2, at the other end of the region of reduced recombination from the *S*-

locus region. Thus, both the *S*- and *M*-loci governing tristyly occur in a large, putatively

pericentromeric region of low recombination on chromosome 2 causing extensive and significant

SNP associations across many Mb of sequence.

The *S*-locus is a supergene consisting of divergent alleles with hemizygous genes

The *S*- and *M*-locus associated regions we identified were both extremely large with very low

rates of recombination. To further narrow down possible candidate genes and to investigate the

potential hemizygous regions associated with tristyly, we identified regions with significant

differences in read coverage between the floral morphs.

We identified 50 consecutive 50-kb windows that were specific to the *S*-allele (total

length: 2.5 Mb; genome coordinates: Chr2:114,300,000-116,800,000), with read coverage

significantly higher in 20 S-morph (*S*---) plants than 40 non-S-morph (*ss*--) plants (criteria: the

ratio between the average read coverage of non-S-morph and S-morph plants was lower than 0.4,

with *p*-value of the *t*-test $< 9.73217066 \times 10^{-7}$, Fig. 4a). Downstream adjacent to this 2.5 Mb

region were 13 consecutive 50-kb windows (total length: 650 kb; genome coordinates:

Chr2:116,800,000-117,450,000) that were specific to the *s*-allele and had read coverage

significantly lower in 20 S-morph (*S*---) plants than 40 non-S-morph (*ss*--) plants (criteria: the

ratio between the average read coverage of non-S-morph and S-morph plants was higher than

1.6, with the *p*-value of the *t*-test $< 9.73217066 \times 10^{-7}$, Fig. 4a).

We hypothesized that given the high divergence between alleles, our collapsed genome

assembly may have created a chimeric assembly in this region, leaving the *s*-allele sequence

adjacent to the *S*-allele region in our assembled scaffold and contributing to the lack of a signal

of high diversity when mapping to this reference genome. To investigate this hypothesis further,

we compared our L-morph assembly (*ssmm*) to this assembly in this region using whole genome

alignments. As expected, Anchorwave[29] alignments revealed the presence of a large 2.57 Mb

deletion in the *ssmm* assembly relative to the *SsMm* assembly (Fig. S4), whereas the adjacent

region showed homology. Thus, we conclude that the 2.5 Mb region roughly corresponds to the

S-allele and the 650 kb region roughly corresponds to the *s*-allele. We refined the boundaries of

the two alleles using the coordinates of the closest genes in their flanking regions (Fig. 4a).

Two genes (TIE/SPL and MPC2) were shared between the *S*- and *s*-alleles. The order of

the two genes are opposite in the two alleles. In the *S*-allele, both genes are on the negative

strand, whereas in the *s*–allele they are on the positive strand. This finding suggests the existence

of an inversion containing the two genes (Fig. 4a). The rest of the genes were allele specific:

three genes (CYCD6;1-S, HRGP-S, and LAZY1-S) were specific to the *S*-allele, while five

genes were specific to the *s*-allele (Fig. 4a). LAZY1 is a gene family that regulates the

distribution of auxin[30], HRGP is a superfamily of genes that forms covalently cross-linked

networks in primary cell walls[31], and CYCD6;1 is a gene that regulates the formative division of

cells[32].

The *M*-locus contains one hemizygous gene

We identified five consecutive 50-kb windows (total length: 250 kb; genome coordinates:

Chr2:13,000,000-13,250,000) that had read coverage significantly higher in the 20 M-morph

(*ssM-*) plants than in the 20 L-morph (*ssmm*) plants (criteria: the ratio between the average read

coverage of M-morph and L-morph plants was higher than 1.6, with the *p*-value of the *t*-test <

$9.73217066 \times 10^{-7}$, Fig. 4b). There were nine coding genes in these five windows. However,

among them only one gene was missing from the *ssmm* genome assembly suggesting that it is

absent from the *m*-allele (Fig. 4b). Strikingly, this absent gene, LAZY1-M, also shows strong

similarity to genes in the LAZY1 family, raising the possibility that this locus acts competitively

in the same pathway as LAZY1-S. Consistent with the SNP association mapping, the

hemizygous *M*-locus region was located on the other side of the low-recombing region on

chromosome 2 (Fig. 3e).

Genetic degeneration at the *S*-allele

In species with supergenes, signatures of genetic degeneration, including gene loss and

accumulation of TEs, are expected for the dominant haplotypes of the supergenes, because

dominant haplotypes are only present in heterozygotes and thus do not exhibit recombination[33].

The *S*-allele specific gene loss (the five *s*-allele specific genes) was in accord with this

prediction. In addition, we found that the density of repetitive elements in the *S*- and *M*-alleles

was significantly higher than the flanking regions ($p = 0.00001985$ and $p = 0.04731$,

respectively, Wilcoxon signed-rank test) (Fig. 4f and h), whereas the differences between the *s*-

allele and the flanking regions were not significantly different ($p = 0.847$, Wilcoxon signed-rank

test) (Fig. 4g).

The expression pattern of hemizygous genes

Among the three *S*-allele specific genes, LAZY1-S and HRGP-S were expressed in floral organs

of the S-morph but not non-S-morph plants. LAZY1-S was expressed in styles but not stamens,

whereas in contrast HRGP-S was expressed in stamens but not styles (Fig. 5a). CYCD6;1-S has

low expression levels in both the styles and stamens of the S-morph (Fig. 5a). The *M*-allele

specific gene LAZY1-M was expressed in both styles and stamens (Fig. 5b).

Genes that were differentially expressed in different morphs

For genes that were differentially expressed genome-wide between floral morphs (see Methods

and Table S1 for details of comparisons), three sets of genes showed functional enrichment: **(**1)

genes that were upregulated in S-morph styles (compared with non-S-morph styles) in early buds

(enriched in three non-overlapping functions), (2) genes that were upregulated in S-morph styles

(compared with non-S-morph styles) in late buds (enriched in one function), and (3) genes that

were upregulated in M-morph styles (compared with L-morph styles) in late buds (enriched in

one function) (Table S2). However, none of the five functions inferred had a direct connection to

growth or flower organs, suggesting that only a small number of genes downstream to the *S*- and

M-locus candidate genes are involved in the phenotypic differences between the floral morphs.

Other differentially expressed genes may simply be incidental associations.

The origins of the *S* and *M* loci

To further understand the origins of the *S*- and *M*-loci, we identified the closest paralogs of the *S*-

and *M*-specific genes. We identified one gene that was paralogous to LAZY1-S and LAZY1-M.

Hereafter, we refer to it as "LAZY1-P1". LAZY1-P1 was located 223 kb downstream to

LAZY1-M (Fig. 4b). The closest paralog of HRGP-S was found on the same chromosome but

not in the same region (Fig. 4c). Two paralogs of CYCD6;1-S were annotated in both genome

assemblies, and they are on chromosomes 3 and 5, respectively (hereafter CYCD6;1-Chr3 and

CYCD6;1-Chr5). CYCD6;1-Chr3 shared higher sequence similarity with CYCD6;1-S (Fig. S5)

and was located in a homologous region between chromosomes 2 and 3 (Fig. S2a), which was

probably a result of the whole genome duplication estimated to have happened < 30 million

years ago, and CYCD6;1-Chr5 is likely the ancestral copy of both CYCD6;1-S and CYCD6;1-

Chr3. The distance between the three ancestral genes therefore suggests that the *S*-locus was

formed via stepwise duplication rather than segmental duplication (Fig. 4c).

223   Based on our gene trees and assuming that the hemizygous loci represent the derived

gene copies, LAZY1-S duplicated from LAZY1-P1 first, followed by subsequent gene

duplication from LAZY1-P1 to form LAZY1-M. Phylogenetic dating with BEAST v2.7.6[34]

suggested that the divergence between LAZY1-S and LAZY-P1 occurred approximately 52.4

million years ago (95% confidence interval: 44.9-59.7) and that the subsequent divergence

between LAZY1-M and LAZY-P1 was around 43.0 million years ago (95% confidence interval:

36.4-50.0) (Fig. 6a).

Discussion

We show that in tristylous *Eichhornia paniculata,* both the *S-* and *M*-loci are located in a low

recombining region of Chromosome 2. The *S*-locus is comprised of two divergent alleles: the

2.51 Mb *S*-allele with three *S*-allele specific genes that are hemizygous in most S-morph plants,

and the 596 kb *s*-allele with five *s*-allele specific genes. In the *S*-allele we identified two strong

tristyly candidate genes: LAZY1-S and HRGP-S. LAZY1-S was expressed in styles but not

stamens suggesting that it may inhibit style elongation in the S-morph by decreasing auxin

concentration in styles. HRGP-S was expressed in stamens but not styles suggesting that it may

be involved in filament elongation or possibly pollen function. Our evidence for the *S*-locus

therefore strongly supports its status as a supergene, and its origin in a large region of low

recombination may have been important during the initial origins of these genes via gene

duplication from more physically distant regions (Figure 4).

We found that the *M*-locus contained only one hemizygous gene – LAZY1-M – and this

gene was hemizygous in --*Mm* plants. Strikingly, this gene is in the same gene family as

LAZY1-S suggesting the possible role of gene duplication and neofunctionalization in the origin

of the epistatic interaction between the two loci. Unlike LAZY1-S, which was only expressed in

styles, LAZY1-M was expressed in both styles and stamens and with a higher expression level in

stamens. This suggests that while LAZY1-S may only be involved in the shortening of styles in

the S-morph, LAZY1-M may also regulate auxin distribution and organ growth in both the styles

and stamens of the M-morph. Given that LAZY-M is the only *M*-allele-specific hemizygous

gene, the M-locus may not represent a supergene at this stage in its evolutionary history but

simply contains LAZY1-M, a pleiotropic modifier.

LAZY1-M was also expressed in the stamens but not the styles of the S-morph (Fig. 5b),

suggesting that LAZY1-S and LAZY1-M may compete for the same transcriptional activator and

that the style expression of LAZY1-M is shut down if LAZY1-S is expressed. The expression

level of LAZY1-S in S-morph styles is also much higher than LAZY1-M in M-morph styles. For

example, at early development stages the average normalized count of LAZY1-M in samples of

M-morph styles is 0.02% of that of LAZY1-S in S-morph styles. At late development stages, the

ratio is 1.9%, indicating that the promoter region of LAZY1-S may have stronger affinity to the

transcriptional activator that they compete for, potentially driving differences in style length in

the S- and M-morphs through different degrees of auxin suppression.

In a theoretical treatment of the evolutionary assembly of tristyly[24], Charlesworth

predicted that the M-morph originated last spreading in dimorphic populations of the L- and S-

morph and eventually giving rise to the tristylous polymorphism. Our results on the relative ages

of LAZY1-S and LAZY1-M are consistent with the hypothesis that the *M*-locus evolved after the

S-locus providing empirical support for this prediction. Our results also provide evidence

suggesting a major role for 're-use' through recurrent duplication of the same gene (LAZY1).

Recruitment of this gene explains the origin of the *M*-locus and birth of the third floral morph in

the tristylous system (Fig. 6b) resulting in the morphological and mating complexity that

characterizes Darwin's 'complex marriage arrangement'.

Online Methods

Study system. Plants used in our study originated from seed collected in wild populations of *E.*

paniculata in N.E. Brazil, Cuba and Mexico. The localities of plant material and details of the

five populations used in our study are presented in Table S4. Information on the reproductive

ecology, mating systems and population genetics of *E. paniculata* are available in Ref. 35 and

references cited therein. We germinated seeds using PRO-MIX Premium Potting Mix (Premier

Tech, Rivière-du-Loup, QC, Canada) and seedlings were transferred to individual 7 cm plastic

pots filled with the same soil mix and placed in flooded trays in a glasshouse at the University of

Toronto (Toronto, Canada) with temperatures of 25-28°C (day) and 18-21°C (night) and a 16-

hour day length. Plants were then grown and flowered regularly for several years facilitating

crosses and collection of leaves and floral buds for molecular analyses.

Genome assemblies. We conducted two chromosome-level genome assemblies of *E. paniculata.*

The first assembly was obtained from an individual of genotype *SsMm* (inferred from selfed

progeny, Table S3) obtained from an inter-population cross between an S-morph plant (*SsM-*)

from N.E. Brazil and an M-morph plant (*ssMm*) from Cuba (see Table S4 for information on the

two parents). From this individual, we obtained 79 Gb (105X coverage) PacBio CLR sequences

(sequenced by Arizona Genomics Institute, Tucson, AZ, USA), 55 Gb (45X coverage) Hi-C

(sequenced by Phase Genomics, Seattle, WA, USA) sequences, and 117 Gb (156X coverage)

10X Chromium sequences (sequenced by The Hospital for Sick Children, Toronto, ON, Canada).

The PacBio sequences were assembled with Canu[36] and the assembly was curated with Purge

Haplotigs[37], both by the Canadian Center for Computational Genomics (C3G) (Montreal, QC,

Canada). Then the genome was phased with Falcon-Phase[38], then scaffolded into chromosome-

level scaffolds with the Hi-C data, both by Phase Genomics (Seattle, WA, USA).

To identify misjoins and characterize the relationship between the physical distance and

the genetic distance on each chromosome, we constructed a linkage map with genotyping-by-

sequencing data obtained from a backcross pedigree from an L-morph plant and an M-morph

plant, detailed in[25], accessible under Accession no. SRP069136 at the Sequence Read Archive

(SRA), (http://www.ncbi.nlm.nih.gov/sra, accessed 17 January 2023), and the associated

BioProject alias PRJNA310331). The reads were trimmed with Scythe[39] and Seqtk

(https://github.com/lh3/seqtk). We aligned the trimmed reads to the *SsMm* genome assembly

with NextGenMap V0.5.5[40], and the BAM files were processed with Picard

(http://broadinstitute.github.io/picard) using the four functions FixMateInformation, CleanSam,

SortSam, and AddOrReplaceReadGroups. SAMtools[41] was used to exclude the secondary

alignments. We constructed the linkage map with the alignment files and the software program

Lep-Map3[42]. Markers were split into eight large linkage groups using a logarithm of the odds

(LOD) score limit of 70 and a Theta value of 0.25 for the initial split (SeparateChromosomes2).

Most of the remaining single markers were put into these groups using a LOD score limit of 30

and a LOD difference of 5 (JoinSingles2All). We calculated marker order for each linkage group

with OrderMarkers2 (with default settings). The first ten linkage groups contained 2,124

markers. We then used Juicebox[43] to correct the misjoins with the linkage map. The genome was

polished with the 10X Chromium sequences, using the software Pilon[44]. The repetitive elements

were annotated with RepeatModeller V1.0.11[45] and RepeatMasker V4.0.7[46].

The second assembly of *E. paniculata* used an L-morph (*ssmm*) plant from Mexico (see

Table S4) using sequencing and assembly services from Dovetail Genomics (Scotts Valley, CA,

USA). 59.6 Gb (~50X coverage) PacBio HiFi reads were generated and the reads were

assembled with hifiasm[47]. The assembly was then scaffolded with Omni-C data using the HiRise

software[48]. We used the linkage map previously generated to concatenate scaffolds that belong to

the same linkage group. We also annotated TEs with RepeatModeller V1.0.11[45] and

RepeatMasker V4.0.7[46]. For both of our genome assemblies, the distribution of scaffold length

was assessed with QUAST V5.2.0[49], and the completeness of assemblies was assessed with

BUSCO (Benchmarking Universal Single-Copy Orthologs) V5.4.4[50] with MetaEuk[51] and the

embryophyta_odb10 (2024-01-08) dataset (Table S5).

Genome-wide associations of single nucleotide polymorphisms (SNPs) and read coverage

(RCs). We extracted genomic DNA from 60 *E. paniculata* plants (20 of each floral morph) from

a population in N.E. Brazil (hereafter "Pop 1", see Table S4), using the QIAGEN DNeasy Plant

Mini Kit (QIAGEN, Germantown, MD, USA). Library preparation and sequencing were

conducted by the McGill University and Génome Québec Innovation Centre (Montréal, QC,

Canada) using the 150-bp paired-end protocol on Illumina HiSeq 4000. The samples had an

average sequencing depth of ~8X.

The Illumina sequencing reads were trimmed with Scythe[39] and Seqtk

(https://github.com/lh3/seqtk). We aligned the trimmed reads to the *SsMm* genome assembly

with NextGenMap V0.5.5[52] and the BAM files were processed with Picard

(http://broadinstitute.github.io/picard/) using the four functions FixMateInformation, CleanSam,

SortSam, and AddOrReplaceReadGroups. We used SAMtools[41] to exclude the secondary

alignments. We used freebayes-parallel[53,54] to call SNPs jointly for all samples. We then filtered

the VCF file with VCFtools[55] and vcflib2[56] following a published guideline[57]. We removed sites

if more than 50% of the individuals had missing data, had a minor allele count of <6, or had a

quality score lower than 30. Genotypes with fewer than 4 reads were treated as "missing" and

individuals with more than 50% missing data were not considered further. Then, sites with strong

within-individual allelic bias at heterozygous sites ($0.01 < AB < 0.25$ or $AB > 0.75$), a

discrepancy in the properly paired status for reads supporting reference or alternate alleles, a low

quality relative to depth, a mean depth that was > 13, a quality score < 30, or that occurred

within 5 bp of an indel, were excluded from our analysis. The resulting VCF file had 7,358,165

filtered variant sites.

We conducted a Genome-Wide Association Study (GWAS) with the variant sites using

GEMMA[58], where a relatedness matrix and a linear mixed model were used and p-values of

variant sites were calculated with the Wald test.

To detect hemizygous regions associated with the *S*- and *M*-loci, we conducted another

GWAS with read-coverage, following the pipeline in Ref. 59. We split the reference genome into

25,688 windows, each of which was 50-kb, and calculated the average depth of each window in

individuals using the software mosdepth[60]. For each individual, we normalized the coverage by

dividing the coverage of each window with the average coverage across all windows, then

multiplied it by 2. Therefore, the expected normalized coverage is 1 for hemizygous regions and

2 for the rest of the genome. We conducted two *t*-tests for normalized coverage for each window:

one for the comparison between S-morph versus non-S-morph plants, and the other for the

comparison between the M-morph versus L-morph plants. We identified tristyly allele-specific

windows with the following criteria:

1) *p* value of the *t*-test was smaller than $9.73217066 \times 10^{-7}$, i.e. Bonferroni correction

with $\alpha = 0.05$.

2) We set thresholds on the ratio between the average read coverage of plants of different

morphs:

For *S*-allele specific windows: The read coverage of a *S*-allele is at least 4 times of a *s*-

allele, i.e. the ratio between the average read coverage of non-S-morph plants and S-morph

plants was lower than 0.4: $\frac{2RC(s)}{RC(S)+RC(s)} = \frac{2\times1}{4+1} = 0.4$.

For *s*-allele specific windows: The read coverage of a *s*-allele is at least 4 times of a *S*-

allele, i.e. the ratio between the average read coverage of non-S-morph plants and S-morph

plants was higher than 1.6: $\frac{2RC(s)}{RC(S)+RC(s)} = \frac{2\times4}{4+1} = 1.6$

For *M*-allele specific windows: The read coverage of an *M*-allele is at least 4 times of an

m-allele, i.e. the ratio between the average read coverage of non-M-morph plants and M-morph

plants was lower than 0.25, assuming all M-morph plants were homozygous: $\frac{2RC(m)}{RC(M)+RC(m)} =$

$\frac{2\times1}{4+1} = 0.4$.

For *m*-allele specific windows: The read coverage of an *s*-allele is at least 4 times of a *M*-

allele, i.e. the ratio between the average read coverage of non-M-morph plants and M-morph

plants was higher than 1.6: $\frac{2RC(m)}{RC(M)+RC(m)} = \frac{2\times4}{4+1} = 1.6$

Only regions with at least two consecutive windows that were allele-specific were

considered. As a result, a 2.5 Mb *S*-allele, a 650 kb *s*-allele, and a 250 kb *M*-allele were

identified. We split each allele into 10 equal-sized windows (window sizes: *S*-allele: 250 kb; *s*-

allele: 65 kb; *M*-allele: 25 kb), calculated the repeat content of each window, then compared it

with the 10 windows of the same size flanking each allele (5 windows upstream and 5 windows

downstream of the locus) with Wilcoxon rank-sum tests.

Differential gene expression analysis and genome annotation with transcriptome data. To

infer genes controlling different components of the tristyly syndrome, we conducted a floral

organ-specific transcriptomic analysis. We dissected floral buds of the three morphs at two

developmental stages: "early", with buds between 3-5 mm in length, and "late", with buds

between 5-8 mm. We used sterilized forceps under a stereo dissecting microscope to open the

floral buds and placed the styles and stamens into 2 mL centrifuge tubes which were then frozen

by putting them into liquid nitrogen. We extracted RNA from the frozen tissues with Spectrum™

Plant Total RNA-Kit (Sigma-Aldrich, St. Louis, MO, USA) following the manufacturer's

instructions. For each type of sample, we collected four biological replicates, each containing

styles or stamens from 5-27 buds sampled from 5-17 individuals from each of two populations

(Pop. 1, 2) originating from N.E. Brazil (Table S4, S6).

We prepared sequencing libraries and sequenced on the Illumina HiSeq4000 platform

(100-bp paired-end reads) by the McGill University and Génome Québec Innovation Centre

(Montréal, QC, Canada). Then we aligned the Illumina sequences to the *SsMm* genome assembly

with HISAT2[61] and assembled the transcripts with StringTie[62]. We conducted functional

annotations of the floral organ transcriptome with Mercator4[63]. We used StringTie[62] to estimate

the abundance of each transcript in each sample. Then we used the R package DESeq2[64] to

conduct differential gene expression analysis and plot the normalized count in each sample for

the genes at the *S*- and *M*-loci. We conducted eight differential gene expression analyses for each

type of morph difference (*S* vs. non-S and M vs. L), each tissue (style vs. stamen), and each

development stage (early bud vs. late bud). Only genes with FPKM (Fragments Per Kilobase of

transcript per Million mapped reads) > 1 and TPM (Transcript Per Million) > 1 in at least one

sample were included in the differential gene expression analyses. In each analysis, we identified

the genes that are upregulated in each morph with false discovery rate < 0.05 and LFC (log fold

change) > 2 or < -2. We then did a Mercator BIN enrichment analysis[61] with the list of

upregulated genes for each comparison.

We also annotated the repeat-masked versions of the two genome assemblies with the

floral organ transcriptomes and also leaf and floral bud transcriptomes from Ref. 65, available

under accession number SRP049636 on the Sequence Read Archive (SRA)

(http://www.ncbi.nlm.nih.gov/sra, accessed 20 January 2023) and the associated BioProject alias

is PRJNA266681 (http://www.ncbi.nlm.nih.gov/bioproject/, accessed 20 January 2023), and also

Ref. 66, available under accession number SRP060405 at the Sequence Read Archive and

associated BioProject alias (PRJNA288861) using MAKER v3.01.03[67] and InterProScan 5[68].

This resulted in a total of 35,224 annotated genes for the *SsMm* genome and 29,592 genes for the

ssmm genome.

Origin of the *S* and *M* loci. According to the annotation of the *SsMm* and *ssmm* genome

assemblies, there is only one LAZY1 homolog (LAZY1-P1) that exists in both genome

assemblies. We then downloaded the coding sequences of LAZY1 in *Oryza sativa* and *Zea mays*

from NCBI (https://www.ncbi.nlm.nih.gov/, accession numbers: NM_001138862.1 and

DQ855268.1), and two paralogs of LAZY1 in *Musa acuminata* 'DH-Pahang' from the website

Banana Genome Hub (https://banana-genome-hub.southgreen.fr/). We conducted a multiple

sequence alignment of these sequences and the coding sequences of three LAZY1 genes in *E.*

paniculata, using MAFFT v7[69] with L-INS-i method.

We then estimated the time of divergence between LAZY1-S and LAZY1-P1 and

between LAZY1-M and LAZY1-P1 with BEAST v2.7.6[34]. First, we did model selection of

jModelTest[70] using Akaike information criterion (AIC) as the criteria. The GTR + I + Gamma

model with nCat = 4 was selected. Then we set the time to the most recent ancestors of *Oryza*

sativa and *Zea mays* (81.43 ± 0.3 MYA)[71], Zingiberales and Commelinales (79.8 ± 9.87

MYA)[72], and Poales and Commelinales (106.7 ± 8.3 MYA)[72] as calibration points. Ten

independent Markov Chain Monte Carlo runs, each with 1 x 10^8 generations (first 10% was

burn-in) were conducted. We combined the output of the eight runs using LogCombiner v2.7.6[34]

with a sample frequency of 5,000. Then we used TreeAnnotator v2.7.6[34] to generate the

maximum clade credibility tree, which was visualized in FigTree v1.4.4

(http://tree.bio.ed.ac.uk/software/figtree/) (Fig. 6).

There are two CYCD6;1 paralogs that exists in both genome assemblies, and they are on

chromosomes 3 and 5, respectively (hereafter CYCD6;1-Chr3 and CYCD6;1-Chr5). We then

downloaded the coding sequence of CYCD6;1 in *O. sativa* from NCBI

(https://www.ncbi.nlm.nih.gov/, accession numbers: XM_015789871.2). We conducted a

multiple sequence alignment of the coding sequences of the three CYCD6;1 genes using MAFFT

v7[69] with the L-INS-i method. Then we constructed a Neighbor Joining tree with all the

conserved sites, Jukes-Cantor substitution model, and bootstrap resampling with 100 samples.

(Fig. S).

HRGP-S was not annotated by MAKER or InterProScan. We aligned the transcript of

HRGP-S against the *SsMm* and *ssmm* genome assemblies with the software program blastn[73], to

identify its closest paralog.

Synteny analyses, whole genome alignments and structural variants. We used COGE's

SynMap2 algorithm[74] with the MAKER and InterProScan annotations to examine the synteny

between the two genome assemblies and identify structural variation. To obtain finer-scale whole

genome alignment of syntenic regions to localize putative insertion/deletion differences between

the *S*- and *s*-alleles and between the *M*- and *m*-alleles, we used Anchorwave[29] to align the two

genome assemblies, with the "minimap2" option and the "proali" function.

Population genomic statistics. We calculated the nucleotide diversity (π) of 1-Mb sliding

windows for the 60 individuals (20 of each morph) used in the GWAS analyses. We used

SAMtools[41] to remove the reads with mapping quality < 10 in the filtered BAM files generated

in the section "Genome-wide associations of single nucleotide polymorphisms (SNPs) and read

coverage (RCs)". Then we used ANGSD[75] to calculate the nucleotide diversity (π) of the sliding

windows.

Figure 1 | Tristyly in *Eichhornia paniculata*. a, Flower of the S-morph illustrating exerted

long- and mid-level stamens. The short style is concealed in the perianth tube. **b,** The

arrangement of styles and stamens in the three floral morphs of a tristylous species. L, M and S

refer to the long-, mid- and short-styled morphs, respectively. The arrows indicate "legitimate"

cross-pollinations between anthers and stigmas of equivalent height. In most tristylous species

these are the only compatible matings because of trimorphic incompatibility. However, in *E.*

paniculata "illegitimate" pollinations are permitted as the species is self- and intramorph

compatible[35]. Genotypes for the floral morphs of *E. paniculata* are indicated.

Figure 2 | The recombination landscape, genome content, and nucleotide diversity (π)

across the chromosomes of *Eichhornia paniculata*. a, Marey maps plotting the position on the

linkage map (cM) against the position on the physical map (Mb). **b-e,** Densities of genes (**b**), all

repetitive elements (**c**), LTR retrotransposons (**d**), and non-LTR repeats (**e**), all measured as

proportions of 1-Mb windows. **h,** Nucleotide diversity (π) of 1-Mb windows.

Figure 3 | SNPs associated with tristyly in *Eichhornia paniculata* and morph-specific read

coverage, recombination landscape, and repeat content of the *S*- and *M*-loci. a, SNPs

associated with S- vs. non-S-morphs. The blue lines indicate $p = 2.52397779 \times 10^{-8}$. Inset panels

zoom in on chromosome 2. **b,** SNPs associated with M- vs. L-morphs. The blue lines indicate p

$= 2.23826411 \times 10^{-8}$. Inset panels zoom in on chromosome 2. **c**, Average normalized coverages

of the L- (red), M- (blue), and S- (cyan) morph plants at the *S*-locus (*S*- and *s*-alleles assembled

next to each other). The data were based on 20 plants per morph. **d,** The average normalized

coverages of the L- (red), M- (blue), and S- (cyan) morph plants at the *M*-locus. The data were

based on 20 plants per morph. **e,** The recombination landscape of chromosome 2. **f-h,**

Comparisons of proportion of repetitive elements between the *S*-allele (ten 250-kb windows), *s*-

allele (ten 65-kb windows), and *M*-allele (ten 25-kb windows), and their flanking regions (five

windows upstream and five windows downstream to the locus, with window size equal to the

allele being compared with).

Figure 4 | The composition and proposed model of the evolution of the *S*- and *M*-loci in

***Eichhornia paniculata*. a,** Genomic organization of the *S*-locus genes. Black dashed lines

represent the boundaries of the tristyly alleles inferred from the read coverage analysis. Blue

dashed lines represent inferred boundaries of an inversion. **b,** Genomic organization of the genes

around LAZY1-M. Green dashed lines represent the boundaries of LAZY1-M and its paralog

LAZY1-P1. **c,** Hypothesized evolutionary pathway for the organization of the *S*- and *M*-loci.

Figure 5 | Expression patterns of the three genes that are specific to the *S*- or *M*-allele and expressed in the styles or stamens of *Eichhornia paniculata*. a, Genes specific to the *S*-allele. b, LAZY1-M, which is specific to the *M*-allele. Samples of different morphs are separated by solid lines. For each morph, samples of different floral organs (styles or stamens) are separated by dashed lines. For each combination of morph and floral organ, the samples at the early and

511   late development stages are on the left and right, respectively. Each type of sample had four

512   biological replicates.

Figure 6 | a, Phylogenetic inference of the dates of divergence between LAZY1-P1 and

LAZY1-S and between LAZY1 and LAZY1-M in the ancestors of *Eichhornia paniculata*.

Bayesian tree of LAZY1 genes in *Oryza sativa*, *Zea mays*, *Musa acuminata*, and *E. paniculata*.

Numbers outside of parentheses represent the estimated time of divergence. Blue rectangles and

numbers in parentheses are the 95% confidence intervals of the estimated time of divergence. **b,**

The postulated sequence of the evolution to trimorphism from monomorphism.

References

1. Darwin, C. R. *The different forms of flowers on plants of the same species.* (John Murray,

1877).

2. Barrett, S. C. H. Heterostylous genetic polymorphisms: model systems for evolutionary

analysis. In *Evolution and function of heterostyly* (ed. Barrett, S. C. H.) 1–29 (Springer, 1992).

3. Barrett, S. C. H. The evolution of plant sexual diversity. *Nat. Rev. Genet.* **3**, 274-284 (2002).

4. Lloyd, D. G. & Webb, C. The selection of heterostyly. In *Evolution and function of heterostyly*

(ed. Barrett, S. C. H.) 179–207 (Springer, 1992).

5. Lloyd, D. G. & Webb, C. J. The evolution of heterostyly. In *Evolution and function of*

heterostyly (ed. Barrett, S. C. H.) 151–178 (Springer, 1992).

6. Barrett, S. C. H., Jesson, L. K. & Baker, A. M. The evolution and function of stylar

polymorphisms in flowering plants. *Ann. Bot.* **85**, 253-265 (2000).

7. Lewis, D. & Jones, D. The genetics of heterostyly. In *Evolution and function of heterostyly*

(Ed. Barrett, S. C. H.) 129–150 (Springer, 1992).

8. Mather, K. The genetical architecture of heterostyly in *Primula sinensis*. *Evolution* **4**, 340–352

(1950).

9. Schwander, T., Libbrecht, R. & Keller, L. Supergenes and complex phenotypes. *Current*

Biology **24**, 288–294 (2014).

10. Thompson, M. J. & Jiggins, C. Supergenes and their role in evolution. *Heredity* **113**, 1-8

(2014).

11. Maney, D. L. & Küpper, C. Supergenes on steroids. *Philos. Trans. R. Soc. B* **377**, 20200507

(2022).

12. Kappel, C., Huu, C. N. & Lenhard, M. A short story gets longer: recent insights into the

molecular basis of heterostyly. *J. Exp. Bot.* **68**, 5719-5730 (2017).

13. Barrett, S. C. H. 'A most complex marriage arrangement': recent advances on heterostyly

and unresolved questions. *New Phytol.* **224**, 1051-1067 (2019).

14. Huu, C. N. *et al.* Presence versus absence of CYP734A50 underlies the style-length

dimorphism in primroses. *eLife* **5**, e17956 (2016).

15. Li, J. *et al.* Genetic architecture and evolution of the *S* locus supergene in *Primula vulgaris*.

Nat. Plants **2**, 1-7 (2016).

16. Shore, J. S. *et al.* The long and short of the *S*-locus in *Turnera* (Passifloraceae). *New Phytol.*

224, 1316–1329 (2019).

17. Gutiérrez-Valencia, J. *et al.* Genomic analyses of the *Linum* distyly supergene reveal

convergent evolution at the molecular level. *Curr. Biol.* **32**, 4360–4371 (2022).

18. Zhao, Z. *et al.* Genomic evidence supports the genetic convergence of a supergene

controlling the distylous floral syndrome. *New Phytol.* **237**, 601–614 (2023).

19. Yang, J. *et al.* Haplotype-resolved genome assembly provides insights into the evolution of

S-locus supergene in distylous *Nymphoides indica*. *New Phytol.* **240**, 2058-2071 (2023).

20. Fawcett, J. A. *et al.* Genome sequencing reveals the genetic architecture of heterostyly and

domestication history of common buckwheat. *Nature Plants* **9**, 1236-1251 (2023).

21. Potente, G. *et al.* Comparative genomics elucidates the origin of a supergene controlling

floral heteromorphism. *Mol. Biol. Evol.* **39**, msac035 (2022).

22. Barrett, S. C. H. The evolutionary biology of tristyly. In *Oxford Surveys in Evolutionary*

Biology (Eds. D. Futuyma & J. Antonovics) vol. **9**, 283–326 (Oxford University Press, 1993).

23. Thompson, J. D., Pailler, T., Strasberg, D. & Manicacci, D. Tristyly in the endangered

Mascarene Island endemic *Hugonia serrata* (Linaceae). *Am. J. Bot.* **83**, 1160–1167 (1996).

24. Charlesworth, D. The evolution and breakdown of tristyly. *Evolution* **33**, 489-498. (1979)

25. Arunkumar, R., Wang, W., Wright, S. I. & Barrett, S. C. H. The genetic architecture of

tristyly and its breakdown to self-fertilization. *Mol. Ecol.* **26**, 752–765 (2017).

26. Ossowski, S. *et al.* The rate and molecular spectrum of spontaneous mutations in *Arabidopsis*

thaliana. *Science* **327**, 92–94 (2010).

27. Eckenwalder, J. E. & Barrett, S. C. H. Phylogenetic systematics of Pontederiaceae. *Syst. Bot.*

11, 373–391 (1986).

28. Ness, R. W., Siol, M. & Barrett, S. C. H. *De novo* sequence assembly and characterization of

the floral transcriptome in cross-and self-fertilizing plants. *BMC Genomics* **12**, 298 (2011).

29. Song, B. *et al.* AnchorWave: Sensitive alignment of genomes with high sequence diversity,

extensive structural polymorphism, and whole-genome duplication. *Proc. Natl. Acad. Sci.* **119**,

e2113075119 (2022).

30. Jiao, Z., Du, H., Chen, S., Huang, W. & Ge, L. *LAZY* gene family in plant gravitropism.

Front. Plant Sci. **11**, 606241 (2021).

31. Chen, Y. *et al.* Identification of the abundant hydroxyproline-rich glycoproteins in the root

walls of wild-type *Arabidopsis*, an ext3 mutant line, and its phenotypic revertant. *Plants* **4**, 85–

111 (2015).

32. Sozzani, R. *et al.* Spatiotemporal regulation of cell-cycle genes by SHORTROOT links

patterning and growth. *Nature*, **466**, 128-132 (2021).

33. Gutiérrez-Valencia, J. *et al.* The genomic architecture and evolutionary fates of supergenes.

Genome Biology and Evolution **13**, evab057 (2021).

34. Bouckaert, R. *et al.* BEAST 2.5: An advanced software platform for Bayesian evolutionary

analysis. *PLoS Comput. Biol***. 15**, e1006650 (2019).

35. Ness, R. W., Wright, S. I. & Barrett, S. C. H. Mating-system variation, demographic history

and patterns of nucleotide diversity in the tristylous plant *Eichhornia paniculata*. *Genetics* **184**,

381–392 (2010).

36. Koren, S. *et al.* Canu: scalable and accurate long-read assembly via adaptive k-mer weighting

and repeat separation. *Genome Res*. **27**, 722–736 (2017).

37. Roach, M. J., Schmidt, S. A., & Borneman, A. R. Purge Haplotigs: allelic contig

reassignment for third-gen diploid genome assemblies. *BMC bioinformatics* **19**, 460 (2018).

38. Kronenberg, Z. N. *et al.* Extended haplotype-phasing of long-read de novo genome

assemblies using Hi-C. *Nature Communications* **12**, 1935 (2021).

39. Thress, K., Henzel, W., Shillinglaw, W. & Kornbluth, S. Scythe: a novel reaper-binding

apoptotic regulator. *EMBO J.* **17**, 6135–6143 (1998).

40. Sedlazeck, F. J., Rescheneder, P. & Von Haeseler, A. NextGenMap: fast and accurate read

mapping in highly polymorphic genomes. *Bioinformatics* **29**, 2790–2791 (2013).

41. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *GigaScience* **10**, (2021).

42. Rastas, P. Lep-MAP3: robust linkage mapping even for low-coverage whole genome

sequencing data. *Bioinformatics* **33**, 3726–3732 (2017).

43. Durand, N. C. *et al.* Juicebox provides a visualization system for Hi-C contact maps with

unlimited zoom. *Cell Syst.* **3**, 99–101 (2016).

44. Walker, B. J. *et al.* Pilon: an integrated tool for comprehensive microbial variant detection

and genome assembly improvement. *PLoS One* **9**, e112963 (2014).

45. Smit, A. F. A. & Hubley, R. RepeatModeler Open-1.0. 2008–2015. (2008).

46. Smit, A. F. A., Hubley, R. & Green, P. RepeatMasker Open-4.0. (2013).

47. Cheng, H., Concepcion, G. T., Feng, X., Zhang, H. & Li, H. Haplotype-resolved de novo

assembly using phased assembly graphs with hifiasm. *Nat. Methods* **18**, 170–175 (2021).

48. Putnam, N. H. *et al.* Chromosome-scale shotgun assembly using an *in vitro* method for long-

range linkage. *Genome Res.* **26**, 342–350 (2016).

49. Gurevich, A., Saveliev, V., Vyahhi, N. & Tesler, G. QUAST: quality assessment tool for

genome assemblies. *Bioinformatics* **29**, 1072–1075 (2013).

50. Manni, M. *et al.* BUSCO update: novel and streamlined workflows along with broader and

deeper phylogenetic coverage for scoring of eukaryotic, prokaryotic, and viral genomes. *Mol.*

Biol. Evol. **38**, 4647-4654 (2021).

51. Levy Karin, E., Mirdita, M., & Söding, J. MetaEuk—sensitive, high-throughput gene

discovery, and annotation for large-scale eukaryotic metagenomics. *Microbiome* **8**, 48 (2020).

52. Sedlazeck, F. J., Rescheneder, P. & Von Haeseler, A. NextGenMap: fast and accurate read

mapping in highly polymorphic genomes. *Bioinformatics* **29**, 2790-2791 (2013).

53. Garrison, E. & Marth, G. Haplotype-based variant detection from short-read sequencing.

ArXiv Prepr., ArXiv12073907 (2012).

54. Tange, O. Gnu parallel: the command-line power tool. *Usenix Mag.* **36**, 42 (2011).

55. Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158

(2011).

56. Garrison, E., Kronenberg, Z. N., Dawson, E. T., Pedersen, B. S. & Prins, P. A spectrum of

free software tools for processing the VCF variant call format: vcflib, bio-vcf, cyvcf2, hts-nim

and slivar. *PLoS Comput. Biol.* **18**, e1009123 (2022).

57. Puritz, J. B., Hollenbeck, C. M. & Gold, J. R. dDocent: a RADseq, variant-calling pipeline

designed for population genomics of non-model organisms. *PeerJ* **2**, e431 (2014).

58. Zhou, X. & Stephens, M. Genome-wide efficient mixed-model analysis for association

studies. *Nat. Genet.* **44**, 821–824 (2012).

59. Xue, L. *et al.* Evidences for a role of two Y-specific genes in sex determination in *Populus*

deltoides. *Nature Communications* **11**, 5893 (2020).

60. Pedersen, B. S. & Quinlan, A. R. Mosdepth: quick coverage calculation for genomes and

exomes. *Bioinformatics* **34**, 867–868 (2018).

61. Kim, D., Langmead, B. & Salzberg, S. L. HISAT: a fast spliced aligner with low memory

requirements. *Nat. Methods* **12**, 357–360 (2015).

62. Pertea, M. *et al.* StringTie enables improved reconstruction of a transcriptome from RNA-seq

reads. *Nat. Biotechnol.* **33**, 290–295 (2015).

63. Lohse, M. *et al.* Mercator: a fast and simple web server for genome scale functional

annotation of plant sequence data. *Plant, Cell & Environment* **37**, 1250-1258 (2014).

64. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biol.* **15**, 1–21 (2014).

65. Arunkumar, R., Ness, R. W., Wright, S. I. & Barrett, S. C. H. The evolution of selfing is

accompanied by reduced efficacy of selection and purging of deleterious mutations. *Genetics*

199, 817–829 (2015).

66. Arunkumar, R., Maddison, T. I., Barrett, S. C. H. & Wright, S. I. Recent mating-system

evolution in *Eichhornia* is accompanied by cis-regulatory divergence. *New Phytol.* **211**, 697–707

(2016).

67. Cantarel, B. L. *et al.* MAKER: an easy-to-use annotation pipeline designed for emerging

model organism genomes. *Genome Res.* **18**, 188–196 (2008).

68. Jones, P. *et al.* InterProScan 5: genome-scale protein function classification. *Bioinformatics*

30, 1236–1240 (2014).

69. Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. *Mol. Biol. Evol.* **30**, 772-780 (2013).

70. Posada, D. jModelTest: Phylogenetic Model Averaging. *Mol. Biol. Evol.* **25**, 1253-1256

(2008).

71. Huang, W. *et al.* A well-supported nuclear phylogeny of Poaceae and implications for the

evolution of C4 photosynthesis. *Mol. Plant* **15**, 755-777 (2022).

72. Magallón, S., Gómez-Acevedo, S., Sánchez-Reyes, L. L. & Hernández-Hernández, T. A

metacalibrated time-tree documents the early rise of flowering plant phylogenetic diversity. *New*

Phytol. **207**, 437-453 (2015).

73. Camacho, C. *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421

(2009).

74. Haug-Baltzell, A., Stephens, S. A., Davey, S., Scheidegger, C. E. & Lyons, E. SynMap2 and

SynMap3D: web-based whole-genome synteny browsers. *Bioinformatics* **33**, 2197–2198 (2017).

75. Korneliussen, T. S., Albrechtsen, A., & Nielsen, R. ANGSD: analysis of next generation

sequencing data. *BMC bioinformatics* **15**, 356 (2014).

Acknowledgements

This work was supported by Discovery Grants from the Natural Sciences and Engineering

Research Council of Canada (NSERC) to S.C.H.B. and S.I.W. and student fellowships from the

University of Toronto to H.X. The Canu assembly and Purge Haplotigs pipeline of the *SsMm*

genome was conducted by the Canadian Center for Computational Genomics (C3G), which is a

Genomics Technology Platform (GTP) supported by the Canadian Government through Genome

Canada. We thank William Cole and Tom Gludovacz for glasshouse maintenance, Joanna Tran,

SungA Choe, Ryan Ruan, and Jianuo Feng for help with growing plants, Claire Ellis and Po-

Yuan Huang for help with tissue collection, Baharul Choudhury, Iuliia Pelikh, and Yihe Zheng

for help with DNA extraction, and Joanna Rifkin, Christian Kappel, Mathias Scharmann, Julia

Kreiner, Jingshan Yang, Meng Yuan, Liangjiao Xue, Zoë Humphries, Tyler Kent, and Ting Liu

for advice on data analysis.

Author contributions

H.X., S.I.W., and S.C.H.B. conceived and designed the study and wrote the manuscript.

S.C.H.B. obtained the seeds used for the study. H.X. grew the plants, collected the tissues, and

extracted DNA and RNA. Y.G. performed the genome annotation. S.I.W. performed the

Anchorwave alignment. H.X. performed all remaining analyses. H.X., S.I.W., and S.C.H.B.

wrote the manuscript. All authors read and approved the final manuscript.

Competing interests

None declared.

Data Availability Statement

The PacBio CLR sequences used to assemble the *Eichhornia paniculata SsMm* genome are

available under Accession no. SRR21038607 at the Sequence Read Archive (SRA)

(http://www.ncbi.nlm.nih.gov/sra, accessed 09 February 2024), and the associated BioProject

alias is PRJNA863567. The Hi-C data used to assemble the *E. paniculata SsMm* genome are

available under Accession no. SRR21038606 at the Sequence Read Archive (SRA)

(http://www.ncbi.nlm.nih.gov/sra, accessed 09 February 2024), and the associated BioProject

alias is PRJNA863567. The 10X Chromium data used to polish the *E. paniculata SsMm* genome

are available under Accession no. SRR21038605 at the Sequence Read Archive (SRA)

(http://www. ncbi.nlm.nih.gov/sra, accessed 09 February 2024), and the associated BioProject

alias is PRJNA863567. The PacBio CCS sequences used to assemble the *E. paniculata ssmm*

genome are available under Accession no. SRRXXXXXXXX at the Sequence Read Archive

(SRA) (http://www.ncbi.nlm.nih.gov/sra, accessed XX XX XXXX), and the associated

BioProject alias is PRJNAXXXXXX. The Omni-C data used to assemble the *E. paniculata*

ssmm genome are available under Accession no. SRRXXXXXX at the Sequence Read Archive

(SRA) (http://www.ncbi.nlm.nih.gov/sra, accessed XX XX XXXX), and the associated

BioProject alias is PRJNAXXXXXX. The whole genome Illumina sequences of the 60 *E.*

paniculata individuals in this study are available under Accession no. SRRXXXXXXXX -

SRRXXXXXXXX at the Sequence Read Archive (SRA) (http://www.ncbi.nlm.nih.gov/sra,

accessed XX XX XXXX), and the associated BioProject alias is PRJNA862356. The RNA-Seq

data of styles and stamens are available under Accession no. SRRXXXXXXXX -

SRRXXXXXXXX at the Sequence Read Archive (SRA) (http://www.ncbi.nlm.nih.gov/sra,

accessed XX XX XXXX), and the associated BioProject alias is PRJNAXXXXXX. The *E.*

paniculata SsMm genome and its MAKER and InterProScan annotation are available under

Genome ID XXXXX on CoGe (https://genomevolution.org/coge/, accessed XX XX XXXX).

The *E. paniculata ssmm* genome and its MAKER and InterProScan annotation are available

under Genome ID XXXXX on CoGe (https://genomevolution.org/coge/, accessed XX XX

XXXX).