***Staphylococcus aureus* counters organic acid anion-mediated inhibition of**

peptidoglycan cross-linking through robust alanine racemase activity

Sasmita Panda[1], Yahani P. Jayasinghe[2], Dhananjay D. Shinde[1], Emilio Bueno[3], Amanda

Stastny[1], Blake P. Bertrand[1], Sujata S. Chaudhari[1], Tammy Kielian[1], Felipe Cava[3], Donald R.

Ronning[2] and Vinai C. Thomas[1]*

[1]Center for Staphylococcal Research, Department of Pathology and Microbiology, University of

Nebraska Medical Center, Omaha, Nebraska 68198-5900, USA.

[2]Department of Pharmaceutical Sciences, University of Nebraska Medical Center, Omaha, NE,

68198, USA

[3]Laboratory for Molecular Infection Medicine Sweden (MIMS), Umeå Center for Microbial

Research (UCMR), Department of Molecular Biology, Umeå University, Umea SE-90187,

Sweden

*Corresponding author.

Mailing address: Department of Pathology and Microbiology

17   University of Nebraska Medical Center

18   Omaha, NE 68198-6495

19   Tel: 402-559-3640, Fax: 402-559-4077

20   E-mail: vinai.thomas@unmc.edu

21 **Keywords**: *Staphylococcus aureus*, weak acids, acetate, D-alanyl-D-alanine ligase, Alanine
22 racemase

23
24 **Running Title**: Organic acid anions inhibit Ddl
25

Abstract

Weak organic acids are commonly found in host niches colonized by bacteria, and they can inhibit bacterial growth as the environment becomes acidic. This inhibition is often attributed to the toxicity resulting from the accumulation of high concentrations of organic anions in the cytosol, which disrupts cellular homeostasis. However, the precise cellular targets that organic anions poison and the mechanisms used to counter organic anion intoxication in bacteria have not been elucidated. Here, we utilize acetic acid, a weak organic acid abundantly found in the gut to investigate its impact on the growth of *Staphylococcus aureus*. We demonstrate that acetate anions bind to and inhibit D-alanyl-D-alanine ligase (Ddl) activity in *S. aureus.* Ddl inhibition reduces intracellular D-alanyl-D-alanine (D-Ala-D-Ala) levels, compromising staphylococcal peptidoglycan cross-linking and cell wall integrity. To overcome the effects of acetate-mediated Ddl inhibition, *S. aureus* maintains a high intracellular D-Ala pool through alanine racemase (Alr1) activity and additionally limits the flux of D-Ala to D-glutamate by controlling D-alanine aminotransferase (Dat) activity. Surprisingly, the *modus operandi* of acetate intoxication in *S. aureus* is common to multiple biologically relevant weak organic acids indicating that Ddl is a conserved target of small organic anions. These findings suggest that *S. aureus* may have evolved to maintain high intracellular D-Ala concentrations, partly to counter organic anion intoxication.

Significance

Under mildly acidic conditions, weak organic acids like acetic acid accumulate to high concentrations within the cytosol as organic anions. However, the physiological consequence of organic anion accumulation is poorly defined. Here we investigate how the acetate anion impacts *S. aureus*. We show that acetate anions directly bind Ddl and inhibit its activity. The resulting decrease in intracellular D-Ala-D-Ala pools impacts peptidoglycan integrity. Since acetate is a weak inhibitor of Ddl, mechanisms that maintain a high intracellular D-Ala pools are sufficient to counter the effect of acetate-mediated Ddl inhibition in *S. aureus*.

Organic acids produced by host and bacterial metabolism are critical determinants of

infection outcomes (1, 2). During infection, the host macrophages produce millimolar amounts

of itaconate, a dicarboxylic acid known to inhibit bacterial growth (3). Conversely, many

bacterial pathogens and the gut microflora excrete short-chain organic fatty acids, which exhibit

immunomodulatory functions and can skew the host response during infection (4, 5). Upon

entry into the bacterial cell, organic acids can become toxic to bacteria when they disassociate

in the cytosol as protons and organic anions. The influx of protons can result in cytoplasmic

acidification and prove lethal for some pathogens if not adequately controlled (6). Similarly,

organic anions have been shown to accumulate to toxic levels in the bacterial cytoplasm (7).

However, the precise consequences of organic anion toxicity and the mechanisms pathogens

employ to withstand the effects of anion perturbations within cells are not clearly understood.

Here we focus on the response of *Staphylococcus aureus* to acetic acid, which is the

primary end-product of glucose catabolism under aerobic conditions. *S. aureus* also likely

encounters high concentrations (up to 100 mM) of acetic acid and other short-chain fatty acids

produced by human gut microbiota during intestinal colonization (8). On average, 20% of adults

carry *S. aureus* in their intestines (9), and the burden there often surpasses that found in nasal

passages by more than three orders of magnitude establishing the gut as a primary site for *S.*

aureus colonization (10). We have previously shown that excreted acetic acid can promote

cytoplasmic acidification in cultures of *S. aureus,* especially when the external environment

becomes sufficiently acidic (pH< 5) (11). Cytoplasmic acidification promotes protein oxidation

and triggers a staphylococcal ClpP-dependent damage response that eliminates unfit cells from

the population (12). In contrast, in mildly acidic environments (pH 5.5-6.5), although *S. aureus*

actively buffers its intracellular environment against acidification, the transmembrane pH

gradient (ΔpH) of *S. aureus* will drive the accumulation of millimolar quantities of acetate anions

into the cytoplasm. Previous studies in *Escherichia coli* have shown that acetate intoxication

causes an osmotic imbalance that can transiently be accommodated by the efflux of

physiological anions like glutamate (7). In addition, acetate anions have also been reported to

impact enzymes in the methionine biosynthetic pathway, resulting in a toxic accumulation of

homocysteine and a reduction in intracellular methionine leading to growth inhibition of *E. coli*

(13, 14). However, it remains unclear if these effects are common to other bacteria.

Here, we demonstrate that the primary target of acetate intoxication in *S. aureus* is Ddl.

This crucial enzyme produces the D-Ala-D-Ala dipeptide, that is incorporated into peptidoglycan

precursors and is necessary for cell wall cross-linking. We also demonstrate that carbon flux

through alanine racemase and a tight control of Dat activity increases the cytosolic D-Ala pools

to counter acetate-mediated inhibition of Ddl. Importantly, these phenotypic effects are not

unique to acetate but are conserved across multiple biologically important organic acid anions.

Therefore, we propose that *S. aureus* may have evolved to maintain a high intracellular D-ala

pool partly to offset the inhibition of Ddl by organic anions typically encountered during human

colonization.

Results

Alanine racemase counters acetate intoxication

To identify genetic determinants that counter the effects of acetate intoxication, we

screened the Nebraska Transposon Mutant Library (NTML) for mutants sensitive to 20 mM

acetic acid in Tryptic Soy Broth (TSB) media, pH 6.0. Under these conditions, *S. aureus*

maintains its intracellular pH approx. 1.5 units above the external pH (15) and is estimated to

accumulate over 600 mM acetate in the cytosol (16). The NTML strains were grown under static

conditions at 37°C, and the extent of growth was determined at 24 h by measuring the optical

density at 600 nm (OD_{600}). As a control, we performed an identical screen without acetic acid

supplementation. We normalized the growth of each mutant in both screens (± acetic acid) to

their isogenic wild-type (WT) strain. A comparison of growth indices ($OD_{600\ \text{Tn-mut/WT}}$) for each

mutant in the presence and absence of 20 mM acetic acid revealed that most mutants clustered

close to an index of 1 in the plot (Figure 1A), which suggested that most mutants tolerated

acetate intoxication reasonably well. A few mutants that grew poorly following acetate

intoxication due to inherent growth defects were observed close to the plot diagonal, whereas

those mutants that did not have intrinsic growth deficiencies were located further away from the

diagonal. Among the latter class of mutants, we observed that the *alr1* mutant had the most

substantial reduction in growth when subjected to acetate stress (Figure 1A, B). To confirm that

the acetate-dependent growth defect of the *alr1* mutant was not due to polar effects, we

complemented the mutant by inserting a functional copy of *alr1* under the control of its native

promoter into the *S. aureus* pathogenicity island (SaPI) attachment site. Genetic

complementation completely restored the *alr1* mutant phenotype to WT levels (Figure 1B).

These results suggest that acetate intoxication impairs the growth of *S. aureus* in the absence

of a functional alanine racemase. Further supporting this conclusion, we could reduce acetate

toxicity in the *alr1* mutant by culturing this strain in glucose-free TSB media, which alleviates

carbon catabolite repression and activates TCA cycle-dependent acetate catabolism (Figure

1C) (17). Conversely, the inactivation of citrate synthase (*citZ*), the first enzyme of the TCA-

cycle, re-imposed acetate toxicity in the *alr1* mutant when cultured in glucose-free TSB media

(Figure 1C).

Acetate intoxication alters the intracellular D-Ala-D-Ala pools

Alr1 catalyzes the conversion of L-Ala to D-Ala during staphylococcal growth (Figure 1-figure

supplement 1A). The D-Ala is further converted to D-Ala-D-Ala dipeptide by the ATP-dependent

Ddl (Figure 1-figure supplement 1A) and incorporated into peptidoglycan (PG) muropeptide,

thus playing a crucial role in PG biosynthesis, cross-linking, and integrity (18, 19). Therefore,

we hypothesized that under acetate stress, low concentrations of D-Ala in the *alr1* mutant might

concomitantly reduce D-Ala-D-Ala concentrations in the cell resulting in a growth defect. To test

this hypothesis, we determined the intracellular pool of D-Ala-D-Ala using liquid

chromatography-tandem mass spectrometry (LC-MS/MS). In regular growth media (TSB), we

observed that the inactivation of *alr1* decreased the D-Ala-D-Ala pool by approximately 80%

compared to the WT strain (Figure 1D). However, following acetate intoxication the level of D-

Ala-D-Ala was depleted by more than 99% (Figure 1D). The external supplementation of D-Ala

(5 mM) in the media fully restored the growth of the *alr1* mutant to WT levels under acetic acid

stress (Figure 1E), which suggests that increased intracellular D-Ala pools can overcome the

detrimental impact of acetate intoxication.

The depletion of D-Ala-D-Ala following acetate intoxication is surprising since *S. aureus* is

predicted to have two additional pathways that can synthesize D-Ala and channel it to the

production of this dipeptide. For instance, *S. aureus* harbors a second predicted alanine

racemase (Alr2) that could compensate for the lack of Alr1 activity (Figure 1-figure supplement

1A). Alternatively, Dat, which catalyzes the formation of D-Ala from pyruvate and D-glutamate

(D-Glu), may functionally complement the *alr1* mutant under acetate stress (Figure 1-figure

supplement 1A). However, the lack of functional complementation from these alternate

pathways of D-Ala biosynthesis following acetate intoxication suggests that not all metabolic

routes to D-Ala are operational or that regulatory bottlenecks limit pathway activity. To test

these possibilities, we constructed a series of mutants in which all three predicted routes of D-

Ala biosynthesis (*alr1*, *alr2* and *dat*) were disrupted either individually or in various

combinations and performed growth assays (Figure 1-figure supplement 1B). Surprisingly, we

observed that the inactivation of *alr1* and *dat* simultaneously (*alr1dat* mutant) was synthetic

lethal in *S. aureus*, suggesting that *alr1* and *dat* were the sole contributors of D-Ala in *S. aureus.*

Indeed, the supplementation of D-Ala fully restored the growth of the *alr1dat* mutant (Figure 1-

figure supplement 1C).

The inactivation of *alr2,* either alone or in combination with other D-alanine-generating

enzymes, did not affect growth (Figure 1-figure supplement 1B). This observation suggests that

alr2 is unlikely to be a functional alanine racemase under the growth conditions tested.

Collectively, these results indicate that Dat activity accounts for D-Ala production in the absence

of *alr1*, but its contribution is insufficient to counter acetate intoxication.

Insufficient translation of *dat* impacts the *alr1* mutant following acetate intoxication

Since Dat activity contributes to D-Ala production in the *alr1* mutant, we questioned why Dat

is insufficient to sustain D-Ala-D-Ala pools under conditions of acetate intoxication. One possible

explanation may relate to the maintenance of osmotic balance by *S. aureus*. It has been

proposed that the intracellular accumulation of acetate anions may bring about an efflux of L/D-

Glu from cells to adjust for osmolarity, thus exhausting one of the key substrates for Dat activity

and limiting D-Ala production (7). However, this hypothesis is improbable since the expression

of *dat* from a multicopy vector rescued the *alr1* mutant from the effects of acetate intoxication

(Figure 2-figure supplement 1A), suggesting that the intracellular D-Glu pools are sufficient to

support D-Ala production through Dat activity. Alternatively, we hypothesized that the *alr1*

mutant's heightened sensitivity to acetate toxicity could be due to a decrease in *dat*

transcription which would effectively reduce intracellular D-Ala. However, we found no

detrimental effect of acetate intoxication on *dat* transcription in the *alr1* mutant (Figure 2-figure

supplement 1B). Together, these observations raise the possibility that the depletion of D-Ala-D-

Ala in the *alr1* mutant following acetate intoxication may arise from a post-transcriptional

regulatory bottleneck that limits *dat* from meeting the demand for intracellular D-Ala.

In *S. aureus*, *dat* is part of a bicistronic operon (Figure 2A). The first gene, *pepV*, encodes

an extracellular dipeptidase (20, 21). Transcriptional start site (TSS) mapping of the *pepV-dat*

operon by the adaptor and radioactivity-free (ARF-TSS) method revealed a 30-nucleotide

untranslated region (5'-UTR) extending upstream from the *pepV* initiation codon. The 5'-UTR

includes a Shine-Dalgarno motif (ribosome binding site, SD1) upstream of the *pepV* start codon

(Figure 2A). In addition, a second SD motif (SD2) associated with *dat* was identified within the

pepV coding region (Figure 2A) and did not overlap with the *pepV* termination codon. The

location of SD2 within *pepV* suggests that the insufficient production of D-Ala by *dat* following

acetate intoxication could be attributed to suboptimal translation of *dat*. This could occur as

ribosomes (70S) that are moving from SD1 may interfere with the translation of *dat* from SD2.

To test this hypothesis, we engineered a nonsense mutation in *pepV* (*alr1pepV^{Q12STOP}* mutant)

that would prevent the ribosomes originating from SD1 from moving forward (Figure 2A).

However, the *alr1pepV^{Q12STOP}* mutant grew poorly compared to the *alr1* mutant following

acetate intoxication (Figure 2B). This suggested that the translation of *dat* is coupled to that of

pepV presumably through stable mRNA secondary structures that form within *pepV*. These

structures may not be effectively resolved in the *alr1pepV^{Q12STOP}* mutant due to the absence of

ribosome traffic on *pepV* mRNA.

As an alternative approach to determine if SD2 positioning within *pepV* impeded *dat*

translation, we deleted *pepV* along with SD1 in the *alr1* mutant (*alr1pepV$^{\Delta SD1\text{-}467}$*, Figure 2A). In

the resulting strain, *dat* translation was under the sole control of its native SD2. Remarkably,

the *alr1pepV$^{\Delta SD1\text{-}467}$* mutant did not display a heightened sensitivity to acetate stress and grew

identical to the WT strain following acetate intoxication (Figure 2C). Similarly, an *alr1* mutant in

which *dat* was linked to SD1 (*alr1pepV* mutant, Figure 2A) also phenocopied the WT strain

following acetate intoxication (Figure 2D). Notably, the observed growth differences in

alr1pepV$^{\Delta SD1\text{-}467}$, *alr1pepV* and *alr1pepV^{Q12STOP}* mutants following acetate intoxication did not

result from any changes in *dat* transcription (Figure 2E). These findings collectively suggest

that the native promoter elements, as well as the SD sites of *pepV* and *dat* can independently

support the robust expression and translation of *dat* to levels required for countering acetate

intoxication. However, the genetic arrangement of the *dat* translation initiation region (TIR)

within *pepV* offered tight control of *dat* translation and prevented cells from producing sufficient

enzyme following acetate intoxication.

Why is the Dat tightly controlled?

The need to tightly control Dat activity suggests that flux between D-Ala and D-Glu pools

must be carefully balanced during staphylococcal growth. To gain insight into this process, we

profiled the mass isotopologue distribution (MID) of D-Ala-D-Ala in the WT, *alr1,* and *dat*

mutants under isotopic steady-state conditions using $^{13}C_3{}^{15}N_1$-L-Ala as the tracer during growth

experiments in chemically defined medium (CDM). The flux of $^{13}C_3{}^{15}N_1$-L-Ala through Alr1

should result in $^{13}C_3{}^{15}N_1$-D-Ala production (Figure 3A, D-Ala retains labeled nitrogen). On the

other hand, staphylococcal alanine dehydrogenases (Ald1 and Ald2) catalyze the conversion of

$^{13}C_3{}^{15}N_1$-L-Ala to $^{13}C_3$-pyruvate and finally $^{13}C_3$-D-Ala through Dat activity (Figure 3A). Thus, the

labeled nitrogen in $^{13}C_3{}^{15}N_1$-L-Ala is lost as $^{15}N_1$-NH_4 when fluxed through the Ald/Dat pathway

(Figure 3A). Since the intracellular pools of D-Ala are converted to D-Ala-D-Ala, the MID of the

latter metabolite should mirror the isotopologue ratios of D-Ala produced from either Alr1 or Dat

activities.

LC-MS/MS analysis revealed that ~80% of the intracellular D-Ala-D-Ala pool had

incorporated the labeled L-Ala supplemented in media (fractional contribution, 0.80). As

expected, the majority (~55%) of the D-Ala-D-Ala in the WT was composed of the C_6N_2

isotopologue (in which both units of D-Ala contain labeled carbon and nitrogen), which

suggested that *alr1* was the major contributor of D-Ala in *S. aureus* (Figure 3B). Surprisingly,

the sole contribution of *dat* activity (C_6N_0, C_3N_0, C_0N_0) to D-Ala-D-Ala was less than 1% in the

WT strain and D-Ala-D-Ala isotopologues with at least one D-Ala originating from *dat* activity

(C_6N_1, C_3N_1, C_3N_2) although readily observed, were still in the minority. However, the D-Ala-D-

Ala originating from Dat activity expanded substantially upon *alr1* mutation (Figure 3B).

Inactivation of *dat* itself displayed few differences in the MID of D-Ala-D-Ala, compared to the

WT strain (Figure 3B). These results suggest that flux through Dat is most likely driven towards

D-Glu in the WT strain rather than D-Ala. Only upon inactivation of *alr1* does the Dat activity

reverse towards the production of D-Ala.

To confirm these predictions, we measured the levels of $^{15}N_1$-D-Glu in the WT, *alr1*, and *dat*

mutants following growth with the $^{13}C_3^{15}N_1$-L-Ala tracer. Consistent with Dat activity funneling D-

Ala to D-Glu in the WT, approximately 78% of the D-Glu pool in the WT strain was ^{15}N labeled.

Furthermore, we observed that inactivation of *dat* resulted in the complete depletion of

intracellular levels of $^{15}N_1$-D-Glu (Figure 3C). Inactivation of *alr1* also had a similar outcome with

loss of $^{15}N_1$-D-Glu pools due to the lack of $^{13}C_3^{15}N_1$-D-Ala in this mutant (Figure 3C). Together,

these results strongly suggest that in the WT strain, Dat activity diverts D-Ala towards D-Glu

production.

Given the critical need to produce D-Ala-D-Ala during acetate intoxication, any diversion of

its precursor pool (D-Ala) to D-Glu through Dat activity is bound to decrease cell fitness and thus

may justify its tight translational control. To test this hypothesis, we determined the mean

competitive fitness (*w*) of cells that overexpressed *dat* compared to those that had native levels

of expression. Accordingly, we performed coculture competition assays of the WT strain with

an isogenic mutant strain that either harbored an empty vector (pAQ59) integrated into the

SaPI chromosomal site or a vector containing *dat* under control of its native promoter (pAS8),

following acetate intoxication. Consistent with increased Dat activity in the WT strain being

detrimental to the cell, the mean competitive fitness of the *dat* overexpressing strain was

significantly lower (w_{4h}= 0.91) in the exponential growth phase than its isogenic WT strain that

harbored the empty vector (w_{4h}= 1.26) (Figure 3D). Collectively, these results suggest that Dat

catalyzes the production of D-Glu in the WT strain, and its tight regulation prevents excessive

flux of D-Ala to D-Glu which is necessary to maintain cell fitness following acetate intoxication.

Acetate intoxication impacts PG biosynthesis

Since acetate intoxication ultimately affects D-Ala-D-Ala pools (Figure 1D), we predicted

potential alterations to PG biosynthesis and cell wall integrity. To test this hypothesis, we

quantified various cytosolic PG intermediates in the WT strain by LC-MS/MS analysis. Acetate

intoxication caused a significant increase in the intracellular pools of multiple PG biosynthetic

intermediates, including Uridine diphosphate N-acetylglucosamine (UDP-NAG), UDP-N-

acetylmuramic acid (UDP-NAM), UDP-NAM-L-Ala, UDP-NAM-L-Ala-D-Glu-L-Lys and UDP-

NAM-L-Ala-D-Glu-L-Lys-D-Ala-D-Ala (UDP-NAM-AEKAA) in the WT strain when compared to

the unchallenged control (Figure 4A). However, the growth of the WT strain was slightly

inhibited by acetic acid (Figure 1B), which suggests that the observed accumulation of PG

intermediates may have been due to an imbalance between the rates of PG biosynthesis and

growth. Notably, the *alr1* mutant showed higher levels of UDP-NAM-AEK compared to the WT

and the *dat* mutant following acetate intoxication (Figure 4A), indicating a metabolic block in the

production of UDP-NAM-AEKAA due to insufficient D-Ala-D-Ala. The effect of this metabolic

block is also evident from the increased transcription of *ddl* and *murF* (Figure 4B) which encode

enzymes that incorporate D-Ala-D-Ala into PG precursors, suggesting a greater need to

maintain peptidoglycan cross-linking following acetate intoxication.

Unsurprisingly, the dysregulation of D-Ala-D-Ala homeostasis following acetate intoxication

was also reflected in the extent of cell wall cross-linking in the WT, *alr1* and *dat* mutants.

Muropeptide analysis revealed that acetate intoxication in the WT strain increased levels of

monomeric muropeptides (Figure 4C, Figure 4-figure supplement 1). Conversely, the

percentage of di- and trimeric muropeptides decreased relative to the WT control, as did the

percent cross-linking (Figure 4C). These observations suggest that acetate intoxication

constrains the D-Ala-D-Ala pool in the WT strain and alters PG cross-linking despite Alr1

activity. The extent of PG cross-linking in the *dat* mutant was similar to WT in the presence or

absence of acetate, consistent with our finding that the Dat activity plays a limited role in

maintaining the D-Ala-D-Ala pool in the WT strain (Figure 4C). In contrast, PG cross-linking in

the *alr1* mutant was lower than the WT strain by ~10% (Figure 4C). Acetate intoxication further

decreased the cross-linking approximately 20% relative to WT as well as the ratio of dimeric to

monomeric muropeptides in the *alr1* mutant, which inevitably reduced the growth of this strain

(Figure 4C).

Muropeptide analysis also revealed the accumulation of a disaccharide tripeptide (NAG-

NAM-AEK (M3); m/z. Da, 826.4080) in the peptidoglycan (PG) extracted from the *alr1* mutant

(Figure 4-figure supplement 1B). This finding suggests that the significantly elevated levels of

UDP-NAM-AEK in the *alr1* mutant could efficiently outcompete the substrate specificity of

phospho-N-acetylmuramyl pentapeptide translocase (MraY) for UDP-NAM-AEKAA, ultimately

becoming integrated into the PG structure itself. Interestingly the incorporation of UDP-NAG-

NAM-AEK into the *alr1* mutant's PG only marginally increased following acetate treatment

(Figure 4-figure supplement 1B, see inset). The increase of UDP-NAG-NAM-AEK is most likely

an underestimate since cells with higher levels of incorporation are more likely to lyse due to a

reduction in PG cross-linking. Overall, these observations support a model wherein the

immediate consequences of acetate intoxication are defects in PG crosslinking and

biosynthesis.

Acetate intoxication inhibits Ddl activity

While the above observations point to the consequences of acetate intoxication of *S.*

aureus, its molecular target was not initially identified. Since acetate intoxication dramatically

reduces D-Ala-D-Ala levels in the *alr1* mutant (Figure 1D), we reasoned that acetate might

inhibit either Dat or Ddl activity. To distinguish between these two targets, we measured the

levels of D-Ala in the *alr1* mutant following acetate intoxication. Surprisingly, we observed that

the D-Ala pools in the *alr1* mutant did not significantly change in response to acetate

intoxication compared to the untreated control (Figure 5A). This suggested that Dat activity was

preserved in the *alr1* mutant to the same extent as its untreated control and was not affected by

acetate.

Conversely, these findings also indicate that the acetate-dependent decrease of the D-Ala-

D-Ala pool in the *alr1* mutant was most likely due to the inhibition of Ddl. To test this hypothesis,

we cloned *S. aureus ddl* under the control of a cadmium inducible promoter and induced its

expression in the *alr1* mutant following acetate intoxication (Figure 5B). Indeed, the growth of

the *alr1* mutant was restored to WT levels when *ddl* was overexpressed, strongly suggesting

that Ddl was the target of acetate anion (Figure 5B).

To confirm that acetate inhibits Ddl through direct interactions, we undertook two separate

approaches. As the first approach, 6xHis-tagged *S. aureus* Ddl was purified, and in-vitro

enzyme kinetic assays were performed to determine the possible inhibitory mechanism of Ddl

by acetate. Considering the high concentration of acetate estimated to accumulate in the

cytoplasm, a concentration of 300 mM sodium acetate was used in the initial reactions to test

inhibition (Figure 5C). Interestingly, variation of acetate concentration showed that Ddl was

inhibited *in vitro*, and these conditions suggest an IC_{50} of 400.3 ± 8 mM (Figure 5D). This

indicates significant inhibition of Ddl by acetate when the cellular concentration is near the

hypothesized 600 mM, further confirming that Ddl is a direct target of inhibition by acetate

anion. Furthermore, based on kinetic experiments performed under varying concentrations of

either ATP or D-Ala, the *k*cat values are shown to be distinctly different for each acetate

concentration, which strongly suggests a mixed inhibition mechanism for acetate (Figure 5E

and F, Table S1).

Differential Scanning Fluorometry (DSF) was used as another approach to assess the

direct binding of acetate to Ddl (Table S2). The Ddl protein without any ligand bound shows a

melting temperature (Tm) of 45 °C. After adding 300 mM sodium acetate, Ddl exhibited a 3.7

°C Tm shift indicating a slight thermal stabilization upon binding acetate. This is higher than the

shift in the Tm exhibited by a Ddl/ATP complex. The addition of ADP to Ddl results in a

decrease of 3.2 °C, indicating a decrease in thermal stability compared to Ddl alone.

Intriguingly, when adding acetate to Ddl complexes with ATP or ADP, the Tm increased to 48.9

°C and 49.9 °C, respectively (Table S2). This represents a Tm increase of 2.3 °C when acetate

is added to a Ddl/ATP complex but a Tm increase of 8.1 °C when acetate is added to a

Ddl/ADP complex. The addition of D-Ala to the reaction mixture increases the Tm of Ddl by 4.2

°C and adding acetate to the Ddl/D-Ala mixture shows only a 0.3 °C Tm shift (Table S2). The

widely varying changes in Tm for the tested complexes, particularly when comparing the Tm

values for ligand-free, ADP-bound, and the ADP/Acetate complex, further support a mixed

inhibition mechanism as these data suggest acetate may bind to multiple sites on Ddl or the

location of these binding sites may change depending on the ligand-bound state of the enzyme

due to Ddl conformational changes as observed in Ddl orthologs (22).

Ddl/Acetate complex structure shows binding of acetate at both substrate binding sites

To gain further insight into the mechanism of acetate inhibition, the X-ray crystal structure of

a Ddl/acetate complex was obtained using co-crystals of Ddl and acetate. The crystal diffracted

to 1.9 Å and data were consistent with a P 2 2_1 2_1 space group possessing one molecule of Ddl

in the asymmetric unit (Table S3). The crystal structure of the Ddl/acetate complex (PDB:8FFF)

shows difference density corresponding to acetate at two different sites of the protein. One

acetate is positioned within the adenine binding subsite of the ATP binding site and the other

acetate ion is positioned in the second D-Ala binding site (Figure 5G). The acetate ion in the

ATP binding site interacts with the side chain of Lys177 and the backbone nitrogen of Val216.

Also, the methyl group of acetate forms van der Waals interactions with the side chain of

Leu145 (Figure 5H). The acetate ion that binds to the D-Ala binding site forms a bidentate polar

interaction with the side chain of Arg291 and a hydrogen-bonded interaction with the backbone

nitrogen of Gly312 (Figure 5I). These two residues are conserved in Ddl homologs and

previous structural data clearly illustrate the crucial role these residues play in D-Ala binding

(22).

The acetate-bound structure shows conformational differences compared to the previously

published ligand-free and ADP-bound structures (22). The ω loop, which is associated with

substrate binding, is disordered in both the *S. aureus* Ddl ligand-free (PDB:2I87) and the Ddl-

ADP complex structures (PDB:2I8C) as well as other available Ddl crystal structures that lack

bound substrates or ligands (PDB:3K3P, 5DMX and 6U1C) (23-25). Interestingly, this loop is

well ordered in the acetate-bound structure described here (Figure 5J), which gives the first

view of the *S. aureus* Ddl **ω** loop and the interactions it may form with substrates or inhibitors.

The structural stabilization of the **ω** loop is consistent with the DSF results exhibiting an

increase in the melting temperature upon binding acetate. The **ω** loop is shifted towards the

ATP binding site and repositions the conserved Tyr246 side chain within the ATP binding site,

which likely hinders the binding of ATP (Figure 5J). This positioning is comparable with the

Mycobacterium tuberculosis Ddl (PDB:3LWB) ligand-free structure, which also takes a closed

conformation showing the **ω** loop positioned within the ATP binding site and obstructing ATP

binding (26). Taken together, the kinetic, DSF, and structural data suggest that while acetate

can directly bind within both substrate binding pockets of Ddl, it also stimulates conformational

changes in the dynamic **ω** loop to afford more allosteric-like effects on enzyme activity. Each of

these observations support a mixed inhibition modality.

Multiple organic acids inhibit the *alr1* mutant in a D-Ala-dependent manner

Finally, we determined whether the growth inhibition of the *alr1* mutant is unique to acetate

anion or is a more general phenomenon mirrored by addition of other small organic acids.

Accordingly, we initially performed molecular docking studies of three biologically relevant

organic anions: lactate, propionate and itaconate, in both the ATP and D-Ala binding pockets of

Ddl (Figure 6A-D). The acetate anion-bound structure of Ddl was used as a reference for

analysis. The docking results suggest reasonable poses for lactate, propionate and itaconate

within the ATP binding site forming polar interactions with Ddl residues conserved for binding

ATP. Upon docking, the carboxylate moieties of both lactate and propionate form ionic

interactions with the Lys177 side chain similar to those observed in the Ddl/acetate crystal

structure (Figure 6A and B). Also, the side chain of Glu213 in Ddl forms a hydrogen bonded

interaction with the hydroxyl of lactate (Figure 6A) and van der Waals interactions between

propionate and nearby side chains of Phe175 and Phe295 (Figure 6B) were indicated. The two

carboxylate groups of itaconate form hydrogen bonded interactions with backbone amide

nitrogen atoms of Ala218 and Tyr246 as well as a van der Waals interaction with the nearby

side chain of Phe175 (Figure 6C).

The molecular docking results for lactate, propionate, and itaconate in the D-Ala binding site

of Ddl also show similar types of interactions but with variable poses and slight orientation

differences compared to that observed for acetate in the crystal structure (Figure 6D). The D-

Ala binding site, consisting of primarily charged and polar atoms, allows for a range of binding

modes for these small anions, where the ligand size is a stronger factor in determining the

binding location. Acetate and propionate, being smaller and less sterically hindered, bind

preferentially near the Arg 291 side chain that coordinates the acid moiety of D-ala during the

enzymatic reaction (Figure 6D). Meanwhile, itaconate and L-lactate bind in the more spacious

region between Lys251 and Ser317 (Figure 6D). The Glide scores from the docking results,

which provide a rough estimate of the ΔG of binding for each ligand suggest modest affinity to

the identified binding sites (Table S4).

To determine if these organic acids could impact Ddl function, the WT and the *alr1* mutant

were challenged with lactic, propionic and itaconic acids (Figure 6E-G). All three organic acids

inhibited the growth of the *alr1* mutant. The addition of D-Ala to the culture media rescued the

growth of the *alr1* mutant to WT levels (Figure 6E-G) consistent with Ddl being the target of

lactate, propionate and itaconate. Moreover, overexpression of *ddl* in the *alr1* mutant also

restored growth of the *alr1* mutant following the organic acid challenge (Figure 6-figure

supplement 1A-C). These findings collectively suggest that various organic acid anions can

inhibit Ddl activity in *S. aureus*.

Discussion

Intracellular anion accumulation has long been hypothesized to drive weak organic acid

toxicity in bacteria (16, 27, 28). However, few studies have investigated the mechanism by

which weak acid anions inhibit bacterial growth. Acetic acid is particularly interesting among

weak acids, given that it is a common byproduct of glucose catabolism in bacteria and is

excreted in high concentrations (29). *S. aureus* does not catabolize acetate as a carbon source

unless glucose is first exhausted from the environment (30). This results in the intracellular

accumulation of acetate in *S. aureus* as a function of the bacterial transmembrane pH gradient,

especially when acetic acid concentrations are high in the immediate vicinity of cells. Here we

determine that at high intracellular concentrations, acetate anions directly bind Ddl and inhibit

D-Ala-D-Ala production to adversely impact peptidoglycan cross-linking (Figure 7). However, *S.*

aureus exhibits a remarkable tolerance to acetate intoxication due to the robust production of D-

Ala by Alr1, which ultimately increases D-Ala-D-Ala pools (Figure 7).

Multiple lines of evidence demonstrate Ddl to be the target of acetate anions. First, LC-

MS/MS analysis revealed that acetate intoxication decreased D-Ala-D-Ala pools but not D-Ala in

S. aureus, pointing to Ddl as the target of acetate. Second, DSF and in-vitro enzyme kinetic

studies showed that acetate could bind and inhibit purified rDdl through a mixed inhibition

mechanism. Third, structural analysis of the Ddl-inhibitor complex confirmed that acetate binds

to both the ATP-binding and D-Ala binding sites within Ddl and further induced conformational

changes to the dynamic ω loop, which weakens the binding of ATP to the Ddl active site.

Finally, overexpression of *ddl* alone was sufficient to overcome acetate-mediated inhibition of

the *alr1* mutant and restore growth to WT levels.

Inhibitors that bind an enzyme's catalytic substrate binding sites are usually competed out

by high concentrations of substrates. However, acetate inhibits Ddl through a mixed inhibition

mechanism despite binding to the substrate binding pockets of Ddl. We suspect this is due to

additional conformational changes observed in the dynamic ω loop that affords more allosteric-

like effects on enzyme activity. However, we cannot rule out that acetate might bind to

additional sites in the Ddl-ATP complex, Ddl-ADP complex, or a Ddl-ADP-phospho-D-Ala

complex with varying affinities. The differences in the temperature shifts observed in DSF with

various substrate complexes support this possibility. The crystal structures of Ddl/acetate

complexes with different substrates could provide a more precise conclusion about the

inhibitory modality of Ddl by acetate. In line with acetate's inhibitory effect on Ddl, we observed

that acetate intoxication in the *alr1* mutant led to a disproportionate increase in the cytosolic

pool of PG tripeptide intermediate (UDP-NAM-AEK) compared to the pentapeptide form (UDP-

NAM-AEKAA). Previous reports have suggested that MraY might facilitate the integration of

UDP-NAM-tripeptide into *S. aureus* PG, especially when its concentration within cells exceeds

that of UDP-NAM-pentapeptide (31, 32). Our findings strongly support this hypothesis, as the

analysis of the *alr1* mutant's cell wall muropeptides revealed a clear elevation in the level of the

disaccharide-tripeptide NAG-NAM-AEK. The inhibition of Ddl by acetate would further reduce

the presence of terminal D-Ala-D-Ala moieties within *alr1* muropeptides which likely leaves

these cells incapable of withstanding the outward-directed cell turgor pressure, ultimately

leading to cell death (32).

Despite acetate inhibiting Ddl through a mixed inhibition mechanism, it should be noted that

a functional Alr1 or even the supplementation of D-Ala in culture media can provide significant

tolerance against acetate intoxication in *S. aureus*. These observations suggest that Ddl is only

weakly inhibited by acetate, which is also evident from the relatively high IC_{50} of approximately

400 mM observed in our kinetic experiments with *S. aureus* Ddl. The weak inhibition of Ddl

would suggest that inflating the cytosolic D-Ala pools could promote sufficient generation of D-

Ala-D-Ala to counter acetate intoxication. Indeed, it has been estimated that *S. aureus*

maintains a high concentration of roughly 30 mM intracellular D-Ala (33), which we now

demonstrate to be critical in countering acetate intoxication.

The existence of *pepV* and *dat* within the same operon suggests that these genes may

have evolved related functions. In *Lactococcus lactis* the PepV dipeptidase activity was shown

to be important for supplying cells with L-Ala which was eventually incorporated into PG (34). In

this context, *pepV* and *dat* may have a similar role in modulating the intracellular alanine pool.

A surprising finding of our study was that *dat* expression is relatively stable and tightly

controlled in *S. aureus* due to its SD motif being located within the coding region of *pepV*.

Furthermore, such a genetic arrangement has been linked to translational coupling (35),

wherein active translation from the first gene promotes the translation of the following gene in

the operon, which in the case of *dat* was not sufficient to overcome acetate toxicity in the *alr1*

mutant. Two central mechanisms of translational coupling have been proposed. The first

involves secondary and tertiary mRNA structures that either occlude or encompass the SD

motif of downstream genes and shield it from ribosomes, thus preventing its translation (36).

These mRNA structures can be relieved when a ribosome initiates translation from the first

gene of the operon and exposes the downstream intragenic SD sequences to new 30S

ribosomal subunits (34). In the second mechanism, continued translation of the first gene of the

operon is necessary to increase the abundance of ribosomes in the TIR of the second gene

resulting in its enhanced translation (35). Irrespective of the mechanism of translational

coupling, our results suggest that genetic arrangements that promote translational coupling

might also limit the overall production of *dat* and thus prevent it from functionally

complementing the *alr1* mutant following acetate intoxication. Since our data suggest that Dat

primarily promotes flux from D-Ala to D-Glu when Alr1 is active, the tight control of *dat* through

translational coupling could prevent the depletion of the intracellular reserves of D-Ala

necessary to overcome Ddl inhibition during acetate intoxication. Thus. the elevated D-Ala pool

maintained within the cell could represent a strategic adaptation by *S. aureus* to combat Ddl

inhibition caused by organic acids typically present in the niches colonized by this bacterium.

In conclusion, our findings demonstrate that Ddl is the primary target of acetate anion

intoxication in *S. aureus*. However, other biologically relevant organic anions like lactate,

propionate and itaconate could also inhibit the *alr1* mutant similar to acetate. Furthermore, the

growth inhibition of the *alr1* mutant by these organic acids could be rescued following D-Ala

supplementation, which suggests that Ddl is a *bona fide* and conserved target of various

organic acid anions. Indeed, it is tempting to speculate that the robust Alr1 activity leading to

the accumulation of millimolar levels of D-Ala may have evolved in part to offset the inhibition of

Ddl from the toxic effects of organic anions.

Acknowledgments

This work was funded by NIH/NIAID R01AI125588 and 2P01A1083211 Metabolomics Core to

VCT, 2P01AI083211 Project 4 to TK, respectively. This work was also supported in part by

NIH/NIAID R21AI151924 to DRR. X-ray diffraction data were collected at the Life Sciences

Collaborative Access Team beamline 21-ID-F at the Advanced Photon Source, Argonne

National Laboratory, which is a U.S. Department of Energy (DOE) Office of Science User

Facility operated for the DOE Office of Science by Argonne National Laboratory under Contract

No. DE-AC02-06CH11357. Use of the LS-CAT Sector 21 was supported by the Michigan

Economic Development Corporation and the Michigan Technology Tri-Corridor (Grant

085P1000817). The University of Nebraska Medical Center Mass Spectrometry and

Proteomics Core Facility is administrated through the Office of the Vice Chancellor for

Research and supported by state funds from the Nebraska Research Initiative (NRI). Research

in the Cava lab is supported by the Swedish Research Council, the Laboratory for Molecular

Infection Medicine Sweden (MIMS), Umeå University, the Knut and Alice Wallenberg

Foundation (KAW) and the Kempe Foundation. The funders had no role in the study design,

data collection, interpretation, and decision to submit this work for publication. The authors

have no conflict of interest to declare.

Data Availability

The atomic coordinates and structure factors have been deposited in the Protein Data Bank,

accessible at www.pdb.org, with the PDB ID code 8FFF.

Materials and Methods

Bacterial strains and growth conditions

The *S. aureus* WT and mutant strains described in this study were cultured in TSB

containing 14 mM glucose. *S. aureus* JE2 mutants were mainly obtained from the Nebraska

Transposon Mutant Library (37). These mutants were re-transduced into the WT strain using

Φ11- bacteriophage to eliminate any off-target effects. To generate double or triple mutants,

the Erm^R antibiotic cassette in the transposon mutants was exchanged with Kan^R or Tet^R

cassettes by allelic exchange before introducing an additional mutation. The allelic exchange

was performed as described previously (38). In-frame gene deletion mutants were created

using a temperature-sensitive vector, pJB38, as described previously (38). *S. aureus* mutants

were complemented by inserting the WT allele of mutated genes under the control of their

native promoter into the SaPI1 chromosomal site using the pJC1111 suicide vector (39). For

experiments involving the over-expression of *ddl* in *S. aureus*, *ddl* was cloned into a $CdCl_2$

inducible multicopy vector, pJB68 (38). The concentration of $CdCl_2$ was titrated to achieve full

growth complementation. All bacterial isolates, plasmids, and primers used in this study are

listed in Table S5, S6, and S7, respectively.

Nebraska Transposon Mutant Library (NTML) screen

The NTML mutants were grown in 96-well plates in the presence and absence of 20 mM

acetic acid (pH~6.1) in TSB for 24 hours at 37 °C. The growth of bacteria was determined by

measuring the optical density at 600 nm (OD_{600}) after 24 hours using a TECAN Infinite 200

spectrophotometer. To account for well-to-well variances that accompany 96-well cultures, the

WT strain was independently grown in all the wells of a 96-well plate, both in the presence and

absence of acetic acid. Area under the curve (AUC) values for each mutant under a particular

condition were obtained by normalizing the values to WT AUC. The graph was generated by

plotting the normalized AUC of a mutant under acetate stress versus the control (growth without

acetate).

Competitive fitness assay

The cultures of WT (*S. aureus* JE2) and isogenic mutant strain with either pAQ59 (empty

vector) or pAS8 (containing *dat* gene under control of its native promoter) inserted at the SaPI1

chromosomal site were used to assess competitive fitness. Following the growth of these

cultures for 24 h, 10^7 colony forming units (cfu) per milliliter of each strain were used to

measure the competitive fitness in presence of 20 mM acetate. The bacterial cfu were

enumerated on TSA plates with or without 0.1 mM cadmium chloride immediately after initiation

of competition and at 4 h between tested strains allowing the bacteria to undergo approximately

seven replications to reach 10^9 cfu/ ml. The competitive fitness was calculated using the

Malthusian parameter for competitors using the following formula: $w = \ln (M_f/M_i)/\ln (W_f/W_i)$,

where *f* and *i* represent cfu counts at final (4 h) and initial (time 0) of competition assay,

respectively (11). M and W refer to mutant and WT, respectively.

Sample collection for mass-spectrometry analysis

Overnight cultures of WT, *alr1* and *dat* mutants were inoculated to an OD_{600} of 0.06 units

into 250 ml flasks containing 25ml of TSB 14 mM glucose. Acetic acid (20 mM) was added to

the flasks whenever necessary. The flasks were incubated in a shaker incubator at 37 ⁰C and

250 rpm. A total of 10 OD_{600} units of cells were collected following 3 hours of incubation by

centrifuging the cultures at 10,000 rpm at 4 ⁰C. The pellet was then washed once with ice-cold

saline (0.85% NaCl) and centrifuged again at 10,000 rpm at 4 ⁰C. The bacterial cells were then

resuspended in ice cold quenching solution consisting of 60% ethanol, 2 µM Br-ATP and 2 µM

ribitol. The cytosolic metabolites were obtained by bead beating the cells, followed by

centrifugation. The supernatant was collected and stored at -80 ⁰C until further use. For stable

isotope experiments, overnight cultures were inoculated into a chemically defined medium

(CDM, (40)) containing $^{13}C_3^{15}N_1$-L-Ala (100 mg/ L) in place of L-Ala and the samples were

collected in the exponential phase following 4 hours of incubation at 37 ⁰C.

Chromatography for mass-spectrometry analysis

The chromatographic separation of PG intermediates was performed by liquid

chromatography using XBridge Amide (150 × 2.1 mm ID; 1.7 µm particle size, Waters, USA)

analytical column; whereas D-Ala-D-Ala was analysed using XBridge Amide (100 × 2.1 mm ID;

1.7 µm particle size, Waters, USA). A guard XBridge Amide column (20 × 2.1 mm ID; 1.7µm

particle size, Waters, USA) was connected in front of the analytical column. Mobile phase A

was composed of 10 mM ammonium acetate, 10 mM ammonium hydroxide containing 5 %

acetonitrile in LC-MS grade water; mobile phase B was 100% LC-MS grade acetonitrile. The

column was maintained at 35 °C and the autosampler temperature was maintained at 5 °C. The

gradient was started with the A/B solvent ratio at 15/85 for over 1 minute, followed by a gradual

increase of A. A was reduced to 15% after separation and elution of all the interested

compounds and equilibrated for 6.0 minutes before the next run. The needle was washed with 1

mL of strong wash solvent containing 100% acetonitrile followed by 1 mL of weak wash solvent

comprised of 10% aqueous methanol after each injection. The sample injection volume was 5µl.

Chiral separation of D- and L-isomers of alanine and glutamate was achieved on Astec

CHIROBIOTIC® T column (150 x 2.1 mm, 5 µm particle size) from Supelco. Mobile phase A

was 20 mM ammonium acetate and mobile phase B was 100% ethanol. The mobile phase

composition was 40:60 v/v of A:B in isocratic elution mode pumped at 100 µL/min flow rate. The

injection volume was 5 µL and the column was maintained at room temperature. Multiple

reaction monitoring (MRM) for D- and L- isomers of alanine are listed in Table S8. All other MS

parameters are discussed in the LC-MS/MS analysis section. The L-enantiomer of alanine and

glutamate elutes faster than their D-counterparts. The total run time was 15 minutes.

Targeted LC-MS/MS analysis

Triple-quadrupole-ion trap hybrid mass spectrometer viz., QTRAP 6500+ (Sciex, USA)

connected with Waters UPLC was used for targeted analysis. The QTRAP 6500+ was operated

in polarity switching mode for targeted quantitation of amino acids through the Multiple Reaction

Monitoring (MRM) process. LC-MS MRM data for each metabolite was acquired in centroid

mode as a default setting. MRM details for each analyte are listed in Table S8. The optimized

electrospray ionization (ESI) parameters were as follows: electrospray ion voltage of -4200 V

and 5500 V in negative and positive mode, respectively, source temperature of 500 °C, curtain

gas of 40, and gas 1 and 2 of 40 and 40 psi, respectively. Compound-specific parameters were

optimized for each compound using manual tuning. These parameters include a declustering

potential (DP) of 65 V and -60 V in positive and negative mode, respectively, entrance potential

(EP) of 10 V and -10 V in positive and negative mode, respectively, and collision cell exit

potential (CXP) maintained at 10 V and -10 V in positive and negative mode respectively. Other

compound-specific parameters, such as Q1, Q3, and collision energies, are listed in Table S8.

MRM conditions for PG intermediates were adopted from Vemula *et al* (41)**.**

High Resolution Mass Spectrometry

HRMS Orbitrap (Exploris 480) operated in polarity switching mode was used for the untargeted

analysis of isotopologues of D-Ala-D-Ala and D-Glu in data-dependent MS/MS acquisition mode

(DDA). Electrospray ionization (ESI) parameters were optimized are as follows: electrospray

ion voltage of -2700V and 3500V in negative and positive mode respectively, Ion transfer tube

temperature was maintained at 350°C, m/z scan range was 140-180 Da for non-chiral LC-

method using Amide column whereas, it was 80-160 Da for chiral column method. Sheath gas,

auxiliary gas and sweep gas were optimized according to the UHPLC flow rate. Orbitrap

resolution for precursor ion as well as for fragment ion scan was maintained at 240000 and

60000 respectively. Normalized collision energies at 30, 50 and 150% were used for the

fragmentation. Data was acquired in profile mode. Xcaliber software from Thermo was used for

instrument control and data acquisition. This software was equipped with Qual-, Quant- and

FreeStyle browsers which were used for profiling metabolites and their isotopologues in all

samples. Selected precursor ion for each isotopologue is listed in Table S9. Identification and

detection of all metabolites was aided by the Compound Discoverer (CD) software procured

from Thermo USA. The KEGG and HMDB databases plugged-in with CD software were used

for metabolite identifications and annotations. Mass accuracy for all the ions was maintained at

or below 5 ppm. To correct for natural abundance, we utilized FluxFix, an open-source online

software (42), and independently verified these calculations using the ChemCalc software (43).

Fractional contribution of D-Ala-D-Ala from imported $^{13}C_3{}^{15}N_1$-L-Ala

An estimate of the fractional contribution (FC) of labeled carbon from $^{13}C_3{}^{15}N_1$-L-Ala tracer

incorporated into the intracellular D-Ala-D-Ala pool was calculated using equation 1, as

previously described (44).

$$FC = \frac{\sum_{i=0}^{n} i.m_i}{n.\sum_{i=0}^{n} m_i} \qquad \text{eq. 1}$$

where, *n* is the number of carbon atoms in D-Ala-D-Ala, *i* represents the various carbon

isotopologues of D-Ala-D-Ala and *m* the abundance of the D-Ala-D-Ala isotopologues.

Transcription site identification of the *dat* operon

The adaptor- and radiation-free transcription start site (ARF-TSS) identification method was

employed to identify the 5′-UTR region of the *dat* operon (45). In brief, 1 ug of RNA isolated

from JE2 WT was subjected to reverse transcription by using 5′-phosphorylated primer

pepV_TSS_R1 and the first strand cDNA synthesis kit (Invitrogen, Superscript III First-Strand

Synthesis System). RNA was degraded by using 1M NaOH at 65 ºC for 30 min and then

neutralized with 1M HCl. The resultant cDNA was ligated by using T4 RNA Ligase I (Thermo

Scientific) to generate a circular cDNA. Two inverse primers: pepV_TSS_R2 and

pepV_TSS_F3 were used to amplify the circular cDNA. The amplified product was cloned into a

TOPO Cloning vector and then sequenced using M13F(-20) and M13R primers. All the primers

used in this procedure are mentioned in Table S7.

Quantitative real-time PCR

Quantitative real-time PCR was performed to estimate the transcript levels of *dat*, *ddl* and

murI in the presence and absence of acetate. The samples were collected during the

exponential growth phase and RNA was isolated using a Qiagen RNA isolation kit following the

manufacturer's protocol. A total of 500 ng of RNA was used to synthesize cDNA using the

QuantiTech reverse transcription kit (Qiagen). The cDNA samples were then diluted 1:10 and

used as a template to perform RT-qPCR. The RT-qPCR was carried out using SYBR green

master mix (Roche Applied Science) in a QuantiFast light cycler (Applied Biosystems). The

relative transcript levels were estimated by using the comparative threshold cycle method

($\Delta\Delta$CT) and *sigA* was used as the internal control for normalization. Primers used to perform

RT-qPCR are listed in Table S7.

Muropeptide analysis

The WT and isogenic mutants were inoculated to an OD_{600} of 0.06 into 1-liter flasks

containing 100 mL of TSB 14 mM glucose. Acetic acid (20 mM) was added to the media when

appropriate. A total of 95 OD_{600} units of cells were collected following 6 hours of growth at 37

ºC, 250 rpm. The pelleted cells were then resuspended in 50 % SDS and boiled for 3 hours.

Once boiled, cell wall material was pelleted by ultracentrifugation and washed with water.

Clean sacculi was digested with muramidase (100 µg/ml) and soluble muropeptides reduced

using 0.5 M sodium borate pH 9.5 and 10 mg/mL sodium borohydride. The pH of the samples

was then adjusted to 3.5 with phosphoric acid. UPLC analyses was performed on a Waters-

UPLC system equipped with an ACQUITY UPLC BEH C18 Column, 130 Å, 1.7 µm, 2.1 mm ×

150 mm (Waters Corporation, USA) and identified at Abs. 204 nm. Muropeptides were

separated using a linear gradient from buffer A (0.1 % formic acid in water) to buffer B (0.1 %

formic acid in acetonitrile). Identification of individual peaks was assigned by comparison of the

retention times and profiles to validated chromatograms (46-48). The identity of peak belonging

to disaccharide tripeptide, NAG-NAM-AEK (M3) was assigned by mass spectrometry using

UPLC system coupled to a Xevo G2/XS Q-TOF mass spectrometer (Waters Corp.). Data

acquisition and processing were performed using UNIFI software package (Waters Corp.). The

relative amount of each muropeptide was calculated relative to the total area of the

chromatogram. Representative chromatograms for each sample type are depicted in (Figure 4-

figure supplement 1). The abundance of PG (total PG) was assessed by normalizing the total

area of the chromatogram to the OD_{600}. The degree of cross-linking refers to the number of

peptide bridges and was calculated as % of dimers + % of trimers x 2 + % of tetramers x 3 (49).

Protein purification

The coding region of *ddl* was cloned into pET28a vector to generate a C-terminal 6×His tag

fusion protein before being transferred into *E. coli* BL21(DE3). The cells were grown in Luria

Broth Media (Research Product Internationals) containing 50 µg/mL Kanamycin (Gold

Biotechnology) at 37 °C. When OD_{600} reached 0.6, 1 mM IPTG (Gold Biotechnology) was

added to induce the protein expression. The cells were harvested by centrifugation (3724 *g*)

after inducing them at 16 °C for 20 h. The harvested cells were resuspended in lysis buffer

comprising 25 mM Tris pH 7.5, 150 mM NaCl, and 5 mM 2-Mercaptoethanol. The cells were

lysed by adding Lysozyme (MP-Biomedicals) and DNase I (Roche Applied Sciences) and

incubating them on ice for 30 minutes. Then cells were subjected to sonication (Sonicator

3000, Misonix) to further lyse the cells. The crude cell lysate was refined by centrifuging at

18514 *g* for 40 min (Fixed angle rotor, 5810-R Centrifuge, Eppendorf). The clarified lysate was

applied to a 5 mL HisTrap™TALON™ crude cobalt column (Cytiva) after equilibrating the

column with lysis buffer. The column was washed using the same buffer and the protein was

eluted isocratically using 150 mM imidazole-containing buffer. The purified protein was dialyzed

in 20 mM Tris pH 8.0 buffer and 0.5 mM Tris (2-carboxyethyl) phosphine to use in

crystallization experiments and biochemical assays.

Crystallization of Ddl and data collection

The crystals of Ddl in complex with acetate were obtained by co-crystallization experiments

using the hanging drop vapor diffusion method. The 10 mg/mL of protein was incubated with 30

mM potassium acetate, 5 mM magnesium chloride hexahydrate, and 1 mM ADP for 20 min

before the crystallization experiments. The co-crystals were achieved in crystallization drop

against a well solution consisting of 0.2 M sodium thiocyanate and 20 % polyethylene glycol

monomethyl ether 2000. The crystals were flash cooled in liquid nitrogen immediately after

adding 40% polyethylene glycol 3350 to the crystallization drop for cryoprotection. The data

were collected at the Advance Photon Source Argonne National Laboratory (APS-ANL, IL), LS-

CAT ID-F beamline.

Ddl enzyme kinetic assays

The Invitrogen™ EnzChek™ Phosphate Assay Kit was used to detect the release of

inorganic phosphate by continuously monitoring the absorbance at 360 nm. The reaction

components were added as specified by the kit with 200 nm Ddl (containing 1mM $MgCl_2$), 100

mM Potassium chloride, and ATP. The reaction mixture was incubated for 10 min and D-Ala

substrate was added to initiate the reaction. The inhibition of Ddl by acetate was determined

using various concentrations of sodium acetate, D-Ala, and ATP to determine kinetic

parameters.

Data processing and refinement

The data was processed by CCP4 software (50) and *S. aureus* D-alanyl D-alanine ligase

apoprotein (PDB:2I87) was used for the molecular replacement followed by a rigid body

refinement using PHENIX (51). Manual model refinement was performed using Coot (52). The

XYZ coordinate, B-factor, occupancy, and real space refinements were executed using

PHENIX between manual model refinements. The acetate was modeled using eLBOW and

positioned at the corresponding difference density. The structure was refined using PHENIX

and validated using Molprobity (53).

Molecular Docking Experiments

The docking experiments of small organic acids were performed with the acetate-bound Ddl

structure (PDB:8FFF) with acetate removed. The protein structure was first prepared with the

protein preparation wizard. The lactate, propionate and itaconate ligands were prepared by

LigPrep. The docking experiments were performed using Schrödinger Glide (New York, NY).

Differential Scanning Fluorometry

The reaction mixture was prepared using 22 µM Ddl, 5 mM magnesium chloride, 100 mM

potassium chloride, 1 mM ADP, 300 mM potassium acetate, and 20 mM Tris pH 7.5 buffer as

required. The SyPro orange dye was added to a final concentration of 1 X Protein Thermal

Shift™ Dye (Thermofisher) in the reaction mixture. The reactions were performed in triplicate.

The samples were centrifuged in MicroAmp™ Optical 96-Well Reaction Plate (Applied

Biosystems) at 2325 g for 10 minutes. The protein denaturation was monitored by obtaining the

fluorescence signal by increasing the temperature from 22 °C - 95 °C at 0.5 °C/minute rate

using QuantStudio 3 real-time PCR (ThermoFisher). The melting temperature (Tm) was

determined by calculating the derivative of the fluorescent signal and identifying the centroid of

the observed melting peak.

Figures

Figure 1. Alanine racemase activity counters acetate intoxication (A) The Nebraska Transposon Mutant library was screen against 20 mM acetic acid, pH 6.0 to identify mutants with altered growth phenotypes. The WT strain and transposon mutants were grown for 24 h in TSB ± 20 mM acetic acid. The bacterial growth at 24 h was measured spectrophotometrically (OD_{600}) and normalized to WT growth. The X and Y-axis on the plot represent normalized growth values for each mutant in the presence or absence of acetate. **(B)** The growth of the WT, *alr1* mutant, and *alr1* complemented strain in TSB supplemented with 20 mM acetic acid. **(C)** Aerobic growth of WT, *alr1*, *citZ*, *citZalr1* mutants in TSB media lacking glucose but supplemented with 20 mM acetic acid. **(D)** LC-MS/MS analysis was performed to quantify the intracellular D-Ala-D-Ala pool in strains cultured for 3 h (exponential phase) in TSB ± 20 mM acetic acid. **(E)** The growth of strains was monitored following D-Ala supplementation (5 mM) in TSB + 20 mM acetate, (n=3, mean ± SD). Ac, acetate.

795
796
797 **Figure 1-figure supplement 1. Alr1 and Dat are the primary routes of** D-**Ala production in**
798 *S. aureus.* **(A)** Schematic of the predicted D-Ala-D-Ala generating pathways in *S. aureus*.
799 Growth curves of various *S. aureus* strains grown in the **(B)** absence or **(C)** presence of 5 mM
800 D-Ala (n=3, mean ± SD).

822
823
Figure 2. Translational coupling of *dat* to *pepV* limits the *alr1* mutant from countering
acetate intoxication (A) Schematic representation of various engineered mutations in the
pepV-dat locus.SD, Shine-Dalgarno motif; TSS, transcriptional start site. **(B)-(D)** Growth of
engineered mutants was monitored spectrophotometrically (OD_{600}) in TSB supplemented with
20 mM acetate (n=3, mean ± SD). **(E)** RT-qPCR to determine *dat* transcription in various
mutants relative to the WT strain.

830
831
832 **Figure 2-figure supplement 1. Overexpression of *dat* rescues the growth defect of the**
833 ***alr1* mutant (A)** *dat* was cloned in a multicopy vector (pSP4) controlled by its native promoter.
834 The *S. aureus* strains containing pSP4 and pLI50 (empty vector) were grown in TSB
835 supplemented with 20 mM acetate **(B)** RT-qPCR analysis of *dat* expression in the *alr1* mutant in
836 the presence or absence of 20 mM acetic acid (n=3, mean ± SD).

865
866
Figure 3. Reaction orientation and fluxes through Alr1 and Dat (A) Schematic
representation of various isotopologues of D-Ala-D-Ala and D-Glu generated from $^{13}C_3^{15}N_1$
labeled L-Ala. Metabolites in blue mainly arise from Alr1, red, through the Ald1/2-Dat pathway
and yellow are unlabeled intermediates within cells. The mass isotopologue distribution of **(B)** D-
Ala-D-Ala and **(C)** D-Glu were determined by LC-MS/MS following the growth of *S. aureus* in
chemically defined media supplemented with $^{13}C_3^{15}N_1$ L-Ala (n=3, mean ± SD). Isotopologues of
D-Ala-D-Ala shown in grey color are minor species and are noted in Table S9. **(D)** The mean
competitive fitness (*w*) was determined by co-culturing the WT strain with an isogenic mutant
that contained either the empty pAQ59 vector integrated into the SaPI chromosomal site (WT^{EV})
or the pAS8 vector containing *dat* under the control of its native promoter (WT^{dat}) (n=18, the
dotted lines indicate the median and quartiles).

PG precursors

Figure 4. Acetate intoxication impacts soluble PG precursor pools and cell wall cross-
linking. **(A)** The intracellular pool of PG intermediates in exponential phase cultures of *S.*
aureus was estimated using LC-MS/MS analysis. cps, counts per second **(B)** *ddl* and *murF*
transcription in the exponential growth phase was determined by RT-qPCR analysis (n=3, mean
± SD). **(C)** Cell wall muropeptide analysis of the WT, *alr1* and *dat* mutants was determined
following growth in TSB ± 20 mM acetate for 3 h. Cell wall cross-linking was estimated as
previously described (49). Ac, acetate.

Figure 4-figure supplement 1. Muropeptide analysis. Representative **c**hromatograms of muropeptide extracts from **(A-C)** WT, *alr1* and *dat* mutants, and following **(D-F)** acetate intoxication. A unique peak corresponding to NAG-NAM-AEK (M3, m/z, Da: 826.4080) was identified in the *alr1* mutant. The peak area of M3 was normalized to the total area of peaks observed in the chromatogram and expressed as percent (see inset figure in **B**, n=3, mean ± SD). M, monomer; D, dimer; T, trimer, Tt, tetramer.

940
Figure 5. Acetate anion inhibits Ddl activity. (A) The intracellular D-Ala was determined by
LC-MS/MS analysis. **(B)** The *ddl* gene was overexpressed in *S. aureus* using a cadmium
inducible expression system (pSP36). $CdCl_2$, 0.312 μM. **(C)** Inhibition of recombinant His-
tagged Ddl activity in the presence of 300 mM sodium acetate **(D)** IC_{50} curve of the inhibition of
rDdl by acetate. Michaelis-Menten kinetics of rDdl in varying concentrations of **(E)** D-Ala, and **(F)**
ATP in the presence of acetate to assess the inhibition mechanism. **(G)** Structure of the acetate
bound Ddl (PDB:8FFF). **(H)** Acetate bound to the ATP binding site of Ddl **(I)** Acetate bound to
the second D-Ala binding site of Ddl. The calculated Fo-Fc omit maps are contoured to 3σ and
the mesh is shown in blue. **(J)** Superimposed structure of acetate bound Ddl (slate blue) with

StaDdl apo structure (PDB:2I87, beige) and StaDdl-ADP complex structure (PDB:2I8C, grey) showing a shift of ω loop (red) to ATP binding site. The D-Ala-D-Ala was modeled at the D-Ala binding site using *Thermos thermophius* HB8 Ddl structure (PDB:2ZDQ). The bound ADP (grey) of PDB:2I87 and modeled D-Ala-D-Ala (light blue) indicates the positioning of Ac at ATP and second D-Ala binding sites respectively. Ac, acetate; V, velocity.

Figure 6. Biologically relevant weak acids inhibit growth of the *alr1* mutant. Molecular docking of **(A)** lactate **(B**) propionate and **(C)** itaconate to the ATP binding site of Ddl. **(D)** The relative positions and poise of different organic anions in relation to acetate in the D-Ala binding site of Ddl was determined using Schrödinger Glide. The growth (OD_{600}) of the WT and *alr1* mutant in TSB containing **(E)** lactic acid (40 mM) **(F)** propionic acid (20 mM) and **(G)** itaconic acid (20 mM) in the presence or absence of 5 mM D-Ala.

1032
1033
1034 **Figure 6-figure supplement 1. Overexpression of *ddl* rescues the growth defect of the**
1035 ***alr1* mutant** The growth (OD_{600}) of the WT and *alr1* mutants overexpressing Ddl (pSP36;
1036 cadmium inducible expression of *ddl*) in TSB supplemented with **(A)** lactic acid (40 mM), **(B)**
1037 propionic acid (20 mM) and **(C)** Itaconic acid (20 mM) in the presence or absence of 5 mM D-Ala
1038 (n=3, mean ± SD).

Figure 7. Model depicting the role of Alr1 in countering organic acid anion-mediated inhibition of Ddl. During its growth, *S. aureus* (WT) maintains a substantial intracellular pool of D-Ala through the activity of Alr1. Any excess D-Ala is subsequently converted into D-Glu by the action of the Dat enzyme. The high concentration of D-Ala is crucial for the optimal functioning of Ddl and serves to prevent the inhibition of Ddl by acetate (Ac⁻) and other organic acid anions. This process generates sufficient D-Ala-D-Ala, which is rapidly incorporated into the PG tripeptide precursor UDP-NAM-AEKAA to form UDP-NAM-AEKAA, which ultimately contributes to a robust cross-linked PG (murein) sacculus. In the *alr1* mutant, the Dat reaction orientation is switched to preserve intracellular D-Ala. Nevertheless, this change is inadequate to maintain sufficient D-Ala pool to shield Ddl from inhibition by Ac⁻, due to tight control of *dat* translation. This results in an excess of UDP-NAM-AEK, which competes effectively with UDP-NAM-AEKAA for PG incorporation. The absence of a terminal D-Ala-D-Ala in the PG hinders crosslinking and leads to impaired growth following acetate intoxication.

Table S1: Effect of acetate on Ddl activity

Substrate	Condition	K_m (mM)	V_{max} (µM min^{-1})	k_{cat} (min^{-1})
D-**Ala**	0 mM Acetate	7.0 ± 0.6	16.0 ± 0.4	80.1 ± 2.1
	100 mM Acetate	6.8 ± 0.6	10.9 ± 0.3	54.4 ± 1.6
	300 mM Acetate	8.4 ± 0.6	6.3 ± 0.1	31.5 ± 0.7
ATP	0 mM Acetate	0.6 ± 0.1	12.3 ± 1.1	61.7 ± 5.7
	200 mM Acetate	0.8 ± 0.2	6.9 ± 0.8	34.4 ± 3.8
	300 mM Acetate	0.8 ± 0.3	4.0 ± 0.7	19.9 ± 3.6

Table S2: Impact of acetate on Ddl stability assessed by DSF

Sample	Tm D (ºC)	ΔTm D (ºC)*
Ddl	45.0 ± 0.0	-
Ddl + Sodium acetate (Ac)	48.7 ± 0.0	3.7
Ddl + ATP	46.4 ± 0.0	1.4
Ddl + ATP + Ac	48.9 ± 0.1	3.9
Ddl + ADP	41.9 ± 0.1	-3.2
Ddl + ADP + Ac	49.7 ± 0.1	4.0
Ddl + D-Ala	49.2 ± 0.1	4.2
Ddl + D-Ala + Ac	49.6 ± 0.0	4.5
Ddl + D-Ala + ATP + Ac	47.1 ± 0.1	2.0
Ddl + D-Ala + ADP + Ac	49.8 ± 0.0	4.8

*The ΔTm D values are calculated as the difference in melting temperature of the Ddl apo protein to Ddl with added substrates or acetate inhibitor.

1167

Table S3: Refinement statistics of Ddl/Acetate structure

Resolution Range	48.21 – 1.92 (1.989 – 1.92)
Space group	P 2 21 21
Unit cell	55.123 65.817 99.423 90 90 90
Total reflections	173526 (12202)
Unique reflections	20366 (2806)
Multiplicity	8.5 (9.3)
Completeness (%)	94.80 (50.46)
Mean I/sigma(I)	10.99 (1.72)
Wilson B-factor	37.29
R-merge	0.097 (0.33)
R-meas	0.104 (0.35)
R-pim	0.036 (0.12)
CC1/2	0.998 (0.95)
CC*	0.999 (0.99)
Reflections used in refinement	26870 (1416)
Reflections used for R-free	1601 (92)
R-work	0.21 (0.42)
R-free	0.27 (0.47)
CC (work)	0.223 (0.07)
CC (free)	0.217 (-0.16)
Number of non-hydrogen atoms	2963
macromolecules	2777
ligands	10
solvent	176
Protein residues	355
RMS(bonds)	0.009
RMS(angles)	1.10
Ramachandran favored (%)	93.70
Ramachandran allowed (%)	6.02
Ramachandran outliers (%)	0.29
Rotamer outliers (%)	0.00
Clashscore	6.88
Average B-factor	43.60

macromolecules	43.60
ligands	45.09
solvent	43.64

1214
1215

Table S4: Glide scores from molecular docking studies of organic anions

Organic Anion	ATP binding site	D-Ala binding site
L-lactate	-4.851	-4.431
Propionate	-2.317	-1.883
Itaconate	-3.572	-3.575

Table S5: Strains used in this study

Strains	Description	Source
E. coli Electro-Ten-Blue	General plasmid maintenance strain	Stratagene
S. aureus RN4220	Restriction-deficient strain is routinely used as a transformation intermediate	(54)
S. aureus RN4220: pRN7023	Restriction deficient strain carrying pRN7023 plasmid containing integrase gene routinely used as a transformation intermediate	(39)
E. coli DH5α	General plasmid maintenance strain	Thermo Fisher
E. coli BL21(DE3)	Protein over-expression and purification strain	Novagen
S. aureus JE2	*S. aureus* USA300 LAC cured of all 3 native plasmids	(37)
JE2 *alr1*	*bursa aurealis* transposon mutant, Erm^R	NTML
JE2 *alr1::alr1*	WT copy of *alr1* complemented at the SaPI1 site of *bursa aurealis* transposon mutant, Erm^R	This study
JE2 *citZ*	*bursa aurealis* transposon mutant, Erm^R	NTML
JE2 *citZalr1*	*bursa aurealis* transposon mutant, Erm^R, Kan^R	This study
JE2 *dat*	*bursa aurealis* transposon mutant, Erm^R	NTML
JE2 Δ*alr2*	Inframe isogenic deletion mutant of JE2	This study
JE2 *alr1*Δ*alr2*	*bursa aurealis* transposon mutant, Erm^R, transduced into inframe isogenic deletion mutant JE2 Δ*alr2*	This study
JE2 *alr1dat*	*bursa aurealis* transposon mutant, Erm^R, Tet^R	This study
JE2 alr1: *dat*	pLI50 *dat* plasmid (pSP4) transduced into *bursa aurealis* transposon mutant, Erm^R	
JE2 *pepV*$^{ΔSD1-467}$	Isogenic deletion mutant of SD1 and *pepV*	This study
JE2 *alr1pepV*$^{ΔSD1-467}$	*bursa aurealis* transposon mutant, Erm^R transduced into isogenic deletion mutant of SD1 and *pepV*	This study
JE2 Δ*pepV*	Inframe isogenic deletion mutant of JE2	This study
JE2 *alr1*Δ*pepV*	*bursa aurealis* transposon mutant, Erm^R transduced into inframe isogenic deletion mutant JE2 Δ*pepV*	This study
JE2 *pepV*^{Q12STOP}	Glutamine to STOP codon substitution at 12th amino acid position in PepV	This study
JE2 *alr1pepV*^{Q12STOP}	*Bursa aurealis* transposon mutant, Erm^R transduced into glutamine to STOP codon substitution at 12th amino acid position in PepV	This study
JE2 WT: pJB68	pJB68 transduced into WT JE2	This study
JE2 WT: pSP36	pSP36 transduced into WT JE2	This study
JE2 *alr1*: pJB68	pJB68 transduced into JE2 *alr1*	This study
JE2 *alr1*: pSP36	pSP36 transduced into WT *alr1*	This study
JE2 WT::pAQ59	pAQ59 empty vector inserted at the SaPI1 site of WT JE2	This study
JE2 WT::pAS8	*dat* (under its native promoter) inserted at the SaPI1 site of WT JE2	This study

1259
1260

Table S6: Plasmids used in this study

Plasmids	Description	Source
pLI50	*E. coli- S. aureus* shuttle vector	(55)
pJB38	*E. coli- S. aureus* allelic exchange vector	(38)
pJC1111	*E. coli- S. aureus* SaPI1 site integration vector	(39)
pJB68	*E. coli- S. aureus* cadmium inducible shuttle vector	(38)
pET28a	Expression vector for purification of protein in *E. coli* BL21 (DE3)	Novagen
pAS3	pJC1111 based vector for integration of WT copy of *alr1* at the SaPI1 site	This study
pAS2	pJB38 based vector for *alr2* chromosomal deletion	This study
pSP4	pLI50 *dat* (under control of its native promoter)	This study
pSP19	pJB38 based vector for *pepV* SD1 chromosomal deletion	This study
pSP20	pJB38 based vector for *pepV* SD1-467 chromosomal deletion	This study
pSP16	pJB38 based vector for *pepV* chromosomal deletion	This study
pSP15	pJB38 based vector for substitution of chromosomal *pepV* with *pepV*^{Q12STOP}	This study
pSP36	pJB68 based vector for overexpression of *ddl*	This study
pSP32	pET28a based vector for purification of full length Ddl (C-terminal his tag)	This study
pAQ59	*E. coli- S. aureus* SaPI1 site integration vector with pSC101 ori region	(56)
pAS8	pAQ59 based vector for integration of *dat* gene under its native promoter at the SaPI1 site	This study

Table S7: Primers used in this study

Gene/Modification	Primer name	Primer sequence (5' – 3')
alr1	alr1_F	TGCTGACGAACCAGGAGATA
	alr1_R	TGTAGTTGGGTCAGTAGCTG
alr1 complementation	alr1_comp_F	CGGCCGCTGCATGCCTGCAGACATGAGCAACGTAAAATTG
	alr1_comp_R	AGCTCGGTACCCGGGGATCCAATGACCTTTAATTACTCTAATGATAAC
citZ	1641_F	CAGCGGAGACTAAAATAAGTTC
	1641_R	CCCAATCTCAGATAACATCGTC
dat	dat_F	ACTATAGGTGGCGGTACTTA
	dat_R	ACCATCGGATATCTTCAACG
alr2 deletion	alr2_UP_F	CGAGGCCCTTTCGTCTTCAATACTTAGAAGGTAATGGCTC
	alr2_UP2_R	TCATAGCACTTGCTGTCAATGTATTACAC
	alr2_DN2_F	ATTGACAGCAAGTGCTATGAATCATGATTC
	alr2_DN_R	TTGCATGCCTGCAGGTCGACGCTTCTTCATTTCTATTAACAAG
dat complementation	dat promoter_F	CCTTTCGTCTTCAAGAATTCGATGTGAGTAGGACAGAAATG
	dat promoter_R	TTTTTTCCATTCGAAATCGACTTCCTTTTTTC
	dat_pLI50_F	TCGATTTCGAATGGAAAAAATTTTTTTAAATGGTG
	dat_pLI50_R	TTGCATGCCTGCAGGTCGACCGAAAGTTGATAAATTTAAGTAATTTAATC

TSS identification of *dat* operon	pepV_TSS_R1	P-CCATCTCTATGTGCAATTTC	
	pepV_TSS_R2	GCGTCTTCTGATGCTTTTGC	
	pepV_TSS_F3	GTCCTCGTAAGGCATTAGAC	
	M13F (-20)	GTAAAACGACGGCCAG	
	M13R	CAGGAAACAGCTATGAC	
*pepV*ΔSD1-467	RBS1pepV_UP_F	CCTTTCGTCTTCAAGAATTCAGCGACGCAATTAGGAAC	
	RBS1pepV_UP_R	TTATTCCTCCTTTTTCTATAAGTTAAATTCTATTTTACATGAAAAG	
	RBS1pepV_DN_F	TATAGAAAAAGGAGGAATAATATATGGAAAAAATTTTTTTAAATG	
	RBS1pepV_DN_R	TATAGAAAAAGGAGGAATAATATATGGAAAAAATTTTTTTAAATG	
pepV deletion	pepV_UP2_F	CCTTTCGTCTTCAAGAATTCAACAATTAAAGAAGTAAAAACAAATC	
	pepV_UP2_R	TTTTTTCCATTCGAAATCGACTTCCTTTTTTC	
	pepV_DN2_F	TCGATTTCGAATGGAAAAAATTTTTTTAAATGGTG	
	pepV_DN2_R	TTGCATGCCTGCAGGTCGACTTTCAACTGAAAATGAGAAAC	
*pepV*Q12STOP	pepV_STOP_UP_F	CCTTTCGTCTTCAAGAATTCCAAATCCGAAAGAATATGC	
	pepV_STOP_UP_R	TAATGATTTAATCTTCGTATTGTTGAACTTTTTC	
	pepV_STOP_DN_F	ATACGAAGATTAAATCATTAATGACTTAAAAGGATTATTAG	
	pepV_STOP_DN_R	TTGCATGCCTGCAGGTCGACAAAGACCTGCGTTTTCATTATC	
ddl overexpression plasmid	ddl_pJB68_F	TTTATAAGGAGGAAAAACATATGACAAAAGAAAATATTTGTATCG	

1282

	ddl_pJB68_R	GAATAGGCGCGCCTGAATTCATCCATGATTGAATTTG CTTTAATG
Ddl purification plasmid	ddl_C_Histag_F	CTTTAAGAAGGAGATATACCATGACAAAAGAAAATATT TGTATCG
	ddl_C_Histag_R	CAGTGGTGGTGGTGGTGGTGGTCAATTTTGTATTTAT TTTTCTGTTTATC
ddl RT-qPCR	ddl_RT_F	GGGCTTTTTGAAGTTTTGGA
	ddl_RT_R	TGGTAACCCTCGATGTTCAA
murF RT-qPCR	murF_RT_F	TCACAATTGATTCACGAGCA
	murF_RT_R	CCCAGCACCATCTTGTAATG
dat RT-qPCR	dat RT_F	GATGGTTACGTTGCGACATT
	dat RT_R	CACCTCGATGTTGAATTGCT
sigA RT-qPCR	JE2_RT_sigA_F	AACTGAATCCAAGTGATCTTAGTG
	JE2_RT_sigA_R	TCATCACCTTGTTCAATACGTTTG
dat insertion at the SaPI1 site (pAS8)	dat_UP2_F	GAGCCGCTGCATGCCTGCAGGATGTGAGTAGGACAG AAATG
	dat_UP_R	TTTTTTCCATTCGAAATCGACTTCCTTTTTTC
	dat_DN_F	TCGATTTCGAATGGAAAAAATTTTTTTAAATGGTG
	dat_DN_R	AGCTCGGTACCCGGGGATCCCGAAAGTTGATAAATTT AAGTAATTTAATC

1283

1284

1285

1286

1287

1288

1289

1290

1291

1292

1293

1294

1295

1296

1297

Table S8: Table of Multiple Reaction Monitoring (MRM) transitions

Metabolite	Polarity	MRM (Q1/Q3)	CE (V)	DP (V)	RT	Column
L-Ala	(+)	90.1 / 44.0	17	65	6.4	C
D-Ala	(+)	90.1 / 44.0	17	65	9.5	C
D-Ala-D-Ala	(+)	161.0 / 44.2	30.5	65	3.4	XB
UDP-NAG	(-)	606.0 / 79.0	-149	-80	12.7	XB
UDP-NAM	(-)	678.1 / 79.0	-120	-80	13.2	XB
UDP-NAM-A	(-)	749.1 / 403.0	-42	-90	13.4	XB
UDP-NAM-AE	(-)	878.2 / 403.0	-48	-105	14.3	XB
UDP-NAM-AEK	(-)	1006.2 / 403.0	-50	-130	14.9	XB
UDP-NAM-AEKAA	(-)	1148.5 / 403.0	-55	-140	14.6	XB
NAM	(-)	292.0 / 89.0	-16	-30	6.2	XB
NAG	(+)	204.0 / 138.1	18.9	30	5.6	XB
Br-ATP (IS)	(-)	588.0 / 159.0	-38.9	-60	6.0	XB
Ribitol (IS)	(-)	151.1 / 89.0	-14.8	-60	5.3	XB

1298

1299 CE: Collision energy ;DP: Declustering potential; RT: Retention Time; C: CHIROBIOTIC® T column; XB:
1300 XBridge Amide column

1301

1302

1303

1304

1305

1306

1307

1308

1309

1310

1311

1312

1313

1314

1315

1316

1317

1318

1319

1320

1321

1322

1323

1324

Table S9: HRMS base peak identification of isotopologues

	Metabolite	Isotopologues($^{13}C^{15}N$)	Base peak (m/z)
D-Ala-D-Ala (Positive mode)			
1	C6H12N2O3	C_0N_0	161.0921
2	[13]C1C5H12N2O3	C_1N_0	162.0954
3	[13]C2C4H12N2O3	C_2N_0	163.0988
4	[13]C3C3H12N2O3	C_3N_0	164.1021
5	[13]C3C3H12[15]N1N1O3	C_3N_1	165.09917
6	[13]C3C3H12[15]N2O3	C_3N_2	166.0962
7	[13]C4C2H12N2O3	C_4N_0	165.10549
8	[13]C5C1H12N2O3	C_5N_0	166.10884
9	[13]C6H12N2O3	C_6N_0	167.1122
10	[13]C6H12[15]N1N1O3	C_6N_1	168.1092
11	[13]C6H12[15]N2O3	C_6N_2	169.1063
12	[13]C1C5H12[15]N1N1O3	C_1N_1	163.09246
13	[13]C1C5H12[15]N2O3	C_1N_2	164.0895
14	[13]C2C4H12[15]N1N1O3	C_2N_1	164.0958
15	[13]C2C4H12[15]N2O3	C_2N_2	165.09285
16	[13]C4C2H12[15]N1N1O3	C_4N_1	166.1025
17	[13]C4C2H12[15]N2O3	C_4N_2	167.09956
18	[13]C5C1H12[15]N1N1O3	C_5N_1	167.10588
19	[13]C5C1H12[15]N2O3	C_5N_2	168.1029
20	C6H12[15]N1N1O3	C_0N_1	162.0891
21	C6H12[15]N2O3	C_0N_2	163.0861
D-Glu (Negative mode)			
1	C5H9NO4	C_0N_0	146.04588
2	C5H9[15]NO4	C_0N_1	147.04292
3	[13]C1C4H9NO4	C_1N_0	147.04924
4	[13]C2C3H9NO4	C_2N_0	148.05259
5	[13]C2C3H9[15]NO4	C_2N_1	149.04963
6	[13]C1C4H9[15]NO4	C_1N_1	148.04627
7	[13]C3C2H9[15]NO4	C_3N_1	150.05298
8	[13]C4C1H9[15]NO4	C_4N_1	151.05634
9	[13]C5H9[15]NO4	C_5N_1	152.05969
10	[13]C3C2H9NO4	C_3N_0	149.05595
11	[13]C4C1H9NO4	C_4N_0	150.0593
12	[13]C5H9NO4	C_5N_0	151.06266

1331

1332   RT: D-Ala-D-Ala, 3.4 mins on 10 cm XBridge amide column; D-Glu, 6.4 mins on CHIROBIOTIC® T column

1333
1334
1335
1336

1337   **References**

1338

1.   Passalacqua KD, Charbonneau ME, & O'Riordan MXD (2016) Bacterial metabolism
shapes the host-pathogen interface. *Microbiol Spectr* 4(3).
2.   Brestoff JR & Artis D (2013) Commensal bacteria at the interface of host metabolism
and the immune system. *Nat Immunol* 14(7):676-684.
3.   Tomlinson KL*, et al.* (2021) *Staphylococcus aureus* induces an itaconate-dominated
immunometabolic response that drives biofilm formation. *Nat Commun* 12(1):1399.
4.   Heim CE*, et al.* (2020) Lactate production by *Staphylococcus aureus* biofilm inhibits
HDAC11 to reprogramme the host immune response during persistent infection. *Nat*
Microbiol 5(10):1271-1284.
5.   Schlatterer K, Peschel A, & Kretschmer D (2021) Short-chain fatty acid and FFAR2
Activation - A new option for treating infections? *Front Cell Infect Microbiol* 11:785833.
6.   Salmond CV, Kroll RG, & Booth IR (1984) The effect of food preservatives on pH
homeostasis in *Escherichia coli*. *J Gen Microbiol* 130(11):2845-2850.
7.   Roe AJ, McLaggan D, Davidson I, O'Byrne C, & Booth IR (1998) Perturbation of anion
balance during inhibition of growth of *Escherichia coli* by weak acids. *J Bacteriol*
180(4):767-772.
8.   Cummings JH, Pomare EW, Branch WJ, Naylor CP, & Macfarlane GT (1987) Short
chain fatty acids in human large intestine, portal, hepatic and venous blood. *Gut*
28(10):1221-1227.
9.   Acton DS, Plat-Sinnige MJ, van Wamel W, de Groot N, & van Belkum A (2009) Intestinal
carriage of *Staphylococcus aureus*: how does its frequency compare with that of nasal
carriage and what is its clinical impact? *Eur J Clin Microbiol Infect Dis* 28(2):115-127.
10. Piewngam P*, et al.* (2023) Probiotic for pathogen-specific *Staphylococcus aureus*
decolonisation in Thailand: a phase 2, double-blind, randomised, placebo-controlled trial.
Lancet Microbe 4(2):e75-e83.
11. Thomas VC*, et al.* (2014) A central role for carbon-overflow pathways in the modulation
of bacterial cell death. *PLoS Pathog* 10(6):e1004205.
12. Alqarzaee AA*, et al.* (2021) Staphylococcal ClpXP protease targets the cellular
antioxidant system to eliminate fitness-compromised cells in stationary phase. *Proc Natl*
Acad Sci U S A 118(47).
13. Mordukhova EA & Pan JG (2013) Evolved cobalamin-independent methionine synthase
(MetE) improves the acetate and thermal tolerance of *Escherichia coli*. *Appl Environ*
Microbiol 79(24):7905-7915.
14. Roe AJ, O'Byrne C, McLaggan D, & Booth IR (2002) Inhibition of *Escherichia coli* growth
by acetic acid: a problem with methionine biosynthesis and homocysteine toxicity.
Microbiology (Reading) 148(Pt 7):2215-2222.
15. Gries CM*, et al.* (2016) Potassium uptake modulates *Staphylococcus aureus*
metabolism. *mSphere* 1(3).
16. Carpenter CE & Broadbent JR (2009) External concentration of organic acid anions and
pH: key independent variables for studying how organic acids inhibit growth of bacteria
in mildly acidic foods. *J Food Sci* 74(1):R12-15.
17. Halsey CR*, et al.* (2017) Amino acid catabolism in *Staphylococcus aureus* and the
function of carbon catabolite repression. *mBio* 8(1).

18.   Moscoso M, Garcia P, Cabral MP, Rumbo C, & Bou G (2018) A D-Alanine auxotrophic
live vaccine is effective against lethal infection caused by *Staphylococcus aureus*.
Virulence 9(1):604-620.
19.   Cava F, Lam H, de Pedro MA, & Waldor MK (2011) Emerging knowledge of regulatory
roles of D-amino acids in bacteria. *Cell Mol Life Sci* 68(5):817-831.
20.   Burlak C*, et al.* (2007) Global analysis of community-associated methicillin-resistant
Staphylococcus aureus exoproteins reveals molecules produced in vitro and during
infection. *Cell Microbiol* 9(5):1172-1190.
21.   Goldstein JM, Kordula T, Moon JL, Mayo JA, & Travis J (2005) Characterization of an
extracellular dipeptidase from *Streptococcus gordonii* FSS2. *Infect Immun* 73(2):1256-
1259.
22.   Liu S*, et al.* (2006) Allosteric inhibition of *Staphylococcus aureus* D-Alanine:D-Alanine
ligase revealed by crystallographic studies. *Proc Natl Acad Sci U S A* 103(41):15178-
15183.
23.   Lu Y, Xu H, & Zhao X (2010) Crystal structure of the apo form of D-Alanine:D-Alanine
ligase (Ddl) from *Streptococcus mutans*. *Protein Pept Lett* 17(8):1053-1057.
24.   Pederick JL, Thompson AP, Bell SG, & Bruning JB (2020) D-Alanine-D-alanine ligase as
a model for the activation of ATP-grasp enzymes by monovalent cations. *J Biol Chem*
295(23):7894-7904.
25.   Huynh KH*, et al.* (2015) The crystal structure of the D-alanine-D-alanine ligase from
Acinetobacter baumannii suggests a flexible conformational change in the central
domain before nucleotide binding. *J Microbiol* 53(11):776-782.
26.   Bruning JB, Murillo AC, Chacon O, Barletta RG, & Sacchettini JC (2011) Structure of the
Mycobacterium tuberculosis D-Alanine:D-Alanine ligase, a target of the antituberculosis
drug D-Cycloserine. *Antimicrob Agents Chemother* 55(1):291-301.
27.   Russell JB & Diez-Gonzalez F (1998) The effects of fermentation acids on bacterial
growth. *Adv Microb Physiol* 39:205-234.
28.   Russell JB (1992) Another explanation for the toxicity of fermentation acids at low pH:
anion accumulation versus uncoupling. *Journal of Applied Bacteriology* 73(5):363-370.
29.   Wolfe AJ (2005) The acetate switch. *Microbiol Mol Biol Rev* 69(1):12-50.
30.   Somerville GA & Proctor RA (2009) At the crossroads of bacterial metabolism and
virulence factor synthesis in staphylococci. *Microbiol Mol Biol Rev* 73(2):233-248.
31.   Hammes WP & Neuhaus FC (1974) On the specificity of phospho-N-acetylmuramyl-
pentapeptide translocase. The peptide subunit of uridine diphosphate-N-actylmuramyl-
pentapeptide. *J Biol Chem* 249(10):3140-3150.
32.   Sobral RG, Ludovice AM, de Lencastre H, & Tomasz A (2006) Role of *murF* in cell wall
biosynthesis: isolation and characterization of a murF conditional mutant of
Staphylococcus aureus. *J Bacteriol* 188(7):2543-2553.
33.   Vemula H, Ayon NJ, & Gutheil WG (2016) Cytoplasmic peptidoglycan intermediate
levels in *Staphylococcus aureus*. *Biochimie* 121:72-78.
34.   Huang C, Hernandez-Valdes JA, Kuipers OP, & Kok J (2020) Lysis of a *Lactococcus*
lactis dipeptidase mutant and rescue by mutation in the pleiotropic regulator CodY. *Appl*
Environ Microbiol 86(8).
35.   Huber M*, et al.* (2019) Translational coupling via termination-reinitiation in archaea and
bacteria. *Nat Commun* 10(1):4006.
36.   Rex G, Surin B, Besse G, Schneppe B, & McCarthy JE (1994) The mechanism of
translational coupling in *Escherichia coli*. Higher order structure in the *atpHA* mRNA acts
as a conformational switch regulating the access of de novo initiating ribosomes. *J Biol*
Chem 269(27):18118-18127.
37.   Fey PD*, et al.* (2013) A genetic resource for rapid and comprehensive phenotype
screening of nonessential *Staphylococcus aureus* genes. *mBio* 4(1):e00537-00512.

38. Bose JL, Fey PD, & Bayles KW (2013) Genetic tools to enhance the study of gene
function and regulation in *Staphylococcus aureus*. *Appl Environ Microbiol* 79(7):2218-
2224.
39. Chen J, Yoong P, Ram G, Torres VJ, & Novick RP (2014) Single-copy vectors for
integration at the SaPI1 attachment site for *Staphylococcus aureus*. *Plasmid* 76:1-7.
40. Hussain M, Hastings JG, & White PJ (1991) A chemically defined medium for slime
production by coagulase-negative staphylococci. *J Med Microbiol* 34(3):143-147.
41. Vemula H, Bobba S, Putty S, Barbara JE, & Gutheil WG (2014) Ion-pairing liquid
chromatography-tandem mass spectrometry-based quantification of uridine
diphosphate-linked intermediates in the *Staphylococcus aureus* cell wall biosynthesis
pathway. *Anal Biochem* 465:12-19.
42. Trefely S, Ashwell P, & Snyder NW (2016) FluxFix: automatic isotopologue
normalization for metabolic tracer analysis. *BMC Bioinformatics* 17(1):485.
43. Patiny L & Borel A (2013) ChemCalc: a building block for tomorrow's chemical
infrastructure. *J Chem Inf Model* 53(5):1223-1228.
44. Fendt SM*, et al.* (2013) Metformin decreases glucose oxidation and increases the
dependency of prostate cancer cells on reductive glutamine metabolism. *Cancer Res*
73(14):4429-4438.
45. Wang C, Lee J, Deng Y, Tao F, & Zhang LH (2012) ARF-TSS: an alternative method for
identification of transcription start site in bacteria. *Biotechniques* 52(4).
46. de Jonge BL, Chang YS, Gage D, & Tomasz A (1992) Peptidoglycan composition of a
highly methicillin-resistant *Staphylococcus aureus* strain. The role of penicillin binding
protein 2A. *J Biol Chem* 267(16):11248-11254.
47. De Jonge BL, Gage D, & Xu N (2002) The carboxyl terminus of peptidoglycan stem
peptides is a determinant for methicillin resistance in *Staphylococcus aureus*. *Antimicrob
Agents Chemother* 46(10):3151-3155.
48. Kuhner D, Stahl M, Demircioglu DD, & Bertsche U (2014) From cells to muropeptide
structures in 24 h: peptidoglycan mapping by UPLC-MS. *Sci Rep* 4:7494.
49. Alvarez L, Hernandez SB, de Pedro MA, & Cava F (2016) Ultra-sensitive, high-resolution
liquid chromatography methods for the high-throughput quantitative analysis of bacterial
cell wall chemistry and structure. *Methods Mol Biol* 1440:11-27.
50. Winn MD*, et al.* (2011) Overview of the CCP4 suite and current developments. *Acta
Crystallogr D Biol Crystallogr* 67(Pt 4):235-242.
51. Afonine PV*, et al.* (2012) Towards automated crystallographic structure refinement with
phenix.refine. *Acta Crystallogr D Biol Crystallogr* 68(Pt 4):352-367.
52. Emsley P, Lohkamp B, Scott WG, & Cowtan K (2010) Features and development of
Coot. *Acta Crystallogr D Biol Crystallogr* 66(Pt 4):486-501.
53. Chen VB*, et al.* (2010) MolProbity: all-atom structure validation for macromolecular
crystallography. *Acta Crystallogr D Biol Crystallogr* 66(Pt 1):12-21.
54. Kreiswirth BN*, et al.* (1983) The toxic shock syndrome exotoxin structural gene is not
detectably transmitted by a prophage. *Nature* 305(5936):709-712.
55. Lee CY, Buranen SL, & Ye ZH (1991) Construction of single-copy integration vectors for
Staphylococcus aureus. *Gene* 103(1):101-105.
56. Jacquet R*, et al.* (2019) Dual gene expression analysis identifies factors associated with
Staphylococcus aureus virulence in diabetic mice. *Infect Immun* 87(5):e00163-00119.

1478