Applying gradient tree boosting to QTL mapping with Shapley additive

explanations

Tomohiro Ishibashi, Akio Onogi*

Department of Life Sciences, Faculty of Agriculture, Ryukoku University, Otsu, Shiga, 520-2194,

Japan

Corresponding author:

Akio Onogi

Postal address: 1-5, Yokotani, Oe-cho, Seta, Otsu, Shiga, 520-2194, Japan

Phone: +81-(0)77-599-5719

E-mail: onogiakio@gmail.com

ORCID: 0000-0003-1707-9539

Abstract

Mapping quantitative trait loci (QTLs) is one of the major goals of quantitative genetics; however, identifying the interactions between QTLs remains challenging. Recently developed machine learning methods, such as deep learning and gradient boosting, are transforming the real world. These methods could advance QTL mapping methodologies because of their high capability for capturing complex relationships among features. One problem with applying such complex models to QTL mapping is evaluation of feature importance. In this study, XGBoost, a popular gradient tree boosting algorithm, was applied for QTL mapping in biparental populations with Shapley additive explanations (SHAPs). SHAP is a local (i.e., instance-wise) importance index with the desired properties as feature importance indices. The SHAP-assisted XGBoost (SHAP-XGB) was compared with conventional methods, including likelihood ratio tests (LRT), composite interval mapping (CIM), multiple interval mapping (MIM), and BayesC, using simulations and rice heading date data. SHAP-XGB performed comparable to CIM, MIM, and BayesC in mapping main QTL effects and was superior to conventional methods in mapping QTL interaction effects. As SHAP can evaluate local importance, interactions between markers can be visualized by plotting SHAP interaction values for each instance (plant/line). These results illustrated the strength of SHAP-XGB in detecting and interpreting epistatic QTLs and suggest the possibility that SHAP-XGB complements conventional methods.

Keywords

Association mapping, gene interaction, variable selection, SHAP, explainability

Introduction

Detecting quantitative trait loci (QTLs) underpinning variations in quantitative traits is one of the major goals in analyzing quantitative traits. Detecting QTLs can lead to useful applications such as efficient selection based on QTL genotypes, genomic engineering to modify phenotypes, development of medical technologies by understanding molecular mechanisms, and advancing quantitative genetics

theories by opening "black boxes" (Falconer and Mackay 1996). Interval mapping (IM) is a typical method to detect QTLs in inbred line crosses (Lander and Botstein 1989). IM tests associations of each marker and marker interval with the phenotypes independently. In practice, composite IM (CIM), an extension of IM, is the most popular method (Jansen and Stam 1994; Zeng 1994). CIM considers markers linked with major QTLs as cofactors to enhance detection power and precision when multiple QTLs exist.

Interactions between QTLs, known as epistasis, have also been of interest in quantitative genetics because genes (proteins) function in corporations. Detecting epistatic QTLs will broaden the applications of the QTLs described above and will be helpful to depict gene networks underlying complex quantitative traits. A successful example of QTL and epistatic QTL mapping in plant breeding is the rice heading date. To date, a number of QTL/genes and their networks under short- and long-day conditions have been elucidated (Hori et al. 2016; Wei et al. 2020). Methods for detecting epistatic QTLs include multiple interval mapping (MIM, Laurie et al. 2014) and inclusive CIM (Li et al. 2008). Bayesian regression, where whole genome-wide markers are used as explanatory variables simultaneously, were also applied in detecting epistatic QTLs (Zhang and Xu 2005; Li and Sillanpää 2012). Detection of QTL/gene interactions has been of interest in genome-wide association studies (GWASs), although detection is more difficult than in inbred line crosses because of the generally low allele frequencies. These methods include hierarchical Bayesian modeling for high-order gene interactions (Bayesian high-order interaction toolkit (BHIT) (Wang et al. 2015) and a marginal epistasis test (MAPIT) (Crawford et al. 2017).

Recent machine learning models, such as gradient boosting and deep neural networks, have shown high predictive abilities in various applications. Because these models are inherently capable of capturing complex interactions among features, they are expected to be useful for QTL mapping, particularly for complex QTL interactions. However, unlike linear regressions, the contributions of each feature and feature interaction are not explicitly outputted. Thus, such complex models should be used with efficient and reliable methods to extract the importance of features for selection. For example,

Johnsen et al. (2021) applied Shapley additive explanations (SHAP) to estimate the feature importance of extreme gradient boosting (XGBoost) (Chen and Guestrin 2016) in a GWAS using UK Biobank data. Cui et al. (2022) applied a Shapley-based interaction score (Sundararajan et al. 2020) to deep neural networks to identify gene-gene interactions in a GWAS. The Shapley value was originally proposed to evaluate players' contributions to rewards in cooperative games (Shapley 1953). By considering players and rewards as features and prediction outcomes, respectively, the Shapley value can be used to evaluate the feature and feature combination importance of the prediction models. SHAP is based on the Shapley value and is characterized by the local importance of the feature; that is, the importance of the feature for each instance is estimated (Lundberg and Lee 2017). Because the summation of the SHAP values of the features for an instance is equal to the predicted values of the instance (a property called local accuracy), it is feasible to determine which features have a large impact on the prediction of the instance (Lundberg and Lee 2017; Lundberg et al. 2018). SHAP also has two additional properties–missingness and consistency–that are desirable as feature importance indices.

It may be possible to map QTLs and QTL interactions using SHAP or Shapley value-based scores and recently advanced machine learning more efficiently and effectively than conventional methods. The GWASs mentioned above partially prove this possibility; however, evaluation studies are limited. In particular, the performance and properties in mapping populations, such as bi-parental populations, have not been elucidated, and comparisons with conventional methods, such as CIM, MIM, and Bayesian regressions, have not been conducted. In this study, we evaluated the potential of XGBoost for QTL and epistatic QTL mapping using SHAP. XGBoost was selected because it is one of the most prevalent gradient boosting algorithms. We characterized this SHAP-assisted XGBoost (SHAP-XGB) by applying the method to simulated and real biparental populations and comparing it with conventional methods. The results illustrated the strength of SHAP-XGB in detecting and interpreting epistatic QTLs, as expected, and suggested the possibility that SHAP-XGB complements conventional methods.

Materials and Methods

Extreme gradient boosting

Extreme gradient boosting (XGBoost) is an algorithm for gradient boosting, a class of ensemble learning (Chen and Guestrin 2016). Gradient tree boosting is based on decision trees and represents the response of instance i, y_i, with the linear combination of K decision trees as

$$\hat{y}_i = \sum_{k=1}^{K} f_k(x_i),$$

where f is a tree with structure q and leaf weight w and x_i is the vector of feature. The objective function of XGBoost at iteration t, that is, when adding tth tree, can be written as

$$L^t = \sum_{i=1}^{n} l\left(y_i, \ \hat{y}_i^{t-1} + f_t(x_i)\right) + \Omega(f_t),$$

where l is a loss function and Ω is a penalty function on the tree structure (i.e., the number of leaves and weights). By approximating f with Taylor expansions and dropping the constant terms, the objective function becomes

$$\tilde{L}^t = \sum_{j=1}^{T} \left[\left(\sum_{i\in I_j} g_i\right) w_j + \frac{1}{2}\left(\sum_{i\in I_j} h_i + \lambda\right) w_j^2\right] + \gamma T,$$

where g_i and h_i are the first and second order gradients of the loss function, I_j is the set of instances assigned to leaf j, T is the number of leaves, and λ and γ are the regularization parameters. The optimal weight of leaf j (w_j^*) and the corresponding optimal value of $\tilde{L}^{t*}$ are

$$w_j^* = -\frac{\sum_{i\in I_j} g_i}{\sum_{i\in I_j} h_i + \lambda} \ \text{and} \ \tilde{L}^{t*} = -\frac{1}{2}\sum_{j=1}^{T} \frac{\left(\sum_{i\in I_j} g_i\right)^2}{\sum_{i\in I_j} h_i + \lambda} + \gamma T,$$

respectively. Using these results, whether a leaf I should be split into I_L and I_R can be evaluated with the following equation:

$$\frac{1}{2}\left[\frac{\left(\sum_{i\in I_L} g_i\right)^2}{\sum_{i\in I_L} h_i + \lambda} + \frac{\left(\sum_{i\in I_R} g_i\right)^2}{\sum_{i\in I_R} h_i + \lambda} - \frac{\left(\sum_{i\in I} g_i\right)^2}{\sum_{i\in I} h_i + \lambda}\right] - \gamma.$$

XGBoost uses this formula to search for the tree structures to be added, making the method fast and scalable. The authors proposed exact greedy and approximate algorithms for this type of search.

Shapley values

In cooperative game theory, the Shapley value measures how much each player contributes to the game's reward. Assessing the contributions of individual players independently is insufficient in cooperative games because there are usually interactions between players. The core concept of Shapley's value is the 'marginal contribution'. Marginal contribution represents the extent to which a player's participation in a cooperative game increases the game's rewards. This varies according to the combination of players and order of participation. Therefore, the Shapley value of a player is defined by calculating the marginal contribution of a player to all possible player combinations and participation orders and taking their average. In other words, the Shapley value is the average marginal contribution of the player. The Shapley value for player i is defined as follows:

$$\varphi_i = \sum_{S \subseteq N \setminus \{i\}} \frac{|S|!(M-|S|-1)!}{M!} [f(S \cup \{i\}) - f(S)] \ \ (1),$$

where N is a set of players, S is a subset of players, M is the number of players, and f is the function that outputs rewards for the given players.

SHAP

Players of Shapley values are replaced with features, and rewards are replaced with the model output (e.g., fitter/predicted phenotypes) in SHAP. The model function to output the rewards given to players takes the form of the expected values, that is, $f(S)$ in Eq. (1) is expressed as $E[f(x)|x_S]$ in SHAP, where x and x_S indicate all features and the features in subset S, respectively. Thus, the SHAP value of feature i of an instance is

$$\varphi_i = \sum_{S \subseteq N \setminus \{i\}} \frac{|S|!(M-|S|-1)!}{M!} \left[E[f(x)|x_{S \cup \{i\}}] - E[f(x)|x_S] \right]. \ (2)$$

SHAP is an additive feature-contribution method with three properties: local accuracy, missingness, and consistency. Local accuracy indicates that the summation of the feature contributions defined by Eq. (2) and is equal to the output of the model. Missingness means that importance is not assigned to a missing feature. Consistency implies that if the model is changed such that a feature has

a greater impact on the model, the contribution assigned to that feature does not decrease.

The interaction index was used to assess the impact of one feature of a model when combined

with another. The Shapley interaction index between features i and j is calculated as (Fujimoto et al.

2006),

$$\varphi_{ij} = \sum_{S \subseteq N \setminus \{i,j\}} \frac{|S|!(M-|S|-2)!}{2(M-1)!} \delta_{ij}(S), (3)$$

where $\delta_{ij}(S) = f(S \cup \{i,j\}) - f(S \cup \{i\}) - f(S \cup \{j\}) + f(S)$. The SHAP interaction value was

calculated by replacing the function $f(S)$ with expectations, as shown in Eq. (2). The SHAP

interaction value is split equally between each feature combination, that is, $\varphi_{ij} = \varphi_{ji}$. Thus, total

interaction effects are calculated as $\varphi_{ij} + \varphi_{ij} = 2\varphi_{ij}$. The SHAP value ($\varphi_i$) of feature i for an instance

is decomposed to the main effect of the feature (φ_{ii}) and the interaction effects between the feature

and the others as

$$\varphi_i = \varphi_{ii} + \sum_{j \neq i} \varphi_{ij}.$$

That is, the main effect of feature i is obtained by subtracting the SHAP interaction value between

feature i and j ($j \neq i$) from the SHAP value of feature i.

Lundberg et al. (2018) proposed an efficient algorithm for calculating SHAP values from tree-

based machine learning models (referred to as Tree SHAP). The algorithm exploits the differences in

outputs between models with and without a particular feature using a tree structure to evaluate

$E[f(x)|x_S]$. The algorithm was implemented in R package xgboost (ver. 1.7.5.1).

Algorithm of SHAP-XGB

The algorithm of SHAP-XGB is as follows.

Step 1: Create a bootstrap dataset.

Step 2: Fit an XGBoost model to the dataset, with early stopping determined from the remaining

samples. The xgboost R package was used in this study.

Step 3: Calculate the SHAP values (φ_i) from the fitted samples.

Step 4: Decompose the SHAP values into the SHAP interaction values (φ_{ij}) and the remaining values

(φ_{ii}).

Step 5: Repeat steps 1–4 100 times and calculate the average of φ_i, φ_{ij}, and φ_{ii}. Note that these

values are measures of local importance (i.e., for each line).

Step 6: Average the absolute values of the values obtained in Step 5 across lines to obtain global

importance. The averages of the absolute values of φ_i or φ_{ii} are referred to as the global SHAP

scores and the averages of the absolute values of $2\varphi_{ij}$ are referred to as the global SHAP interaction

scores.

Four notifications were added to these algorithms. First, although the absolute SHAP values are

averaged across lines to obtain the global importance in Step 6, the consistency property still holds for

the global importance scores (Lundberg et al. 2018). Second, bootstrapping was conducted to separate

the data into training and validation sets, which were used for early stopping to prevent overfitting.

Therefore, other resampling methods could be applied. The number of bootstrap replicates (100) was

determined arbitrarily. Thus, fewer numbers—even one—may be acceptable, albeit with a loss of

reliability of the SHAP values. Third, default values were used for the hyperparameters of xgboost

except for the "early_stopping_rounds" argument, which was set to three. This argument specifies the

number of rounds in which the boosting iterations stop if the prediction performance does not improve.

These settings resulted in a moderate fit of the models to data, as shown later. We also attempted

settings that lead to underfitting and overfitting and compared them (see the "Algorithm assessment"

section). Lastly, there would be two choices for detecting the main effects of QTLs: using SHAP values

that contain interaction scores (i.e., φ_i) or using SHAP values from which interaction scores are

removed (i.e., φ_{ii}). In this study, the former score is referred to as "compound" scores, and the latter

is referred to as the "pure" scores for detecting QTL main effects. These two scores were compared as

described in the "Algorithm assessment" section.

Comparing methods

The SHAP-XGB method was compared with other methods using simulated and real data. These

methods were selected based on their prevalence in the QTL analyses of inbred line crosses and/or

ease of execution in the R environment. All methods, including the SHAP-XGB method, used only

marker genotypes as covariates and did not use the genotype probabilities of loci between markers.

Likelihood ratio test (LTR)

LRT was conducted using the interactionPval function of the R package SNPassoc (ver. 2.1.0,

González et al. 2007). The function simultaneously carried out several types of tests, including LRTs

for the main QTL and interaction effects. The likelihood ratios of these tests can be written as $\frac{L_m(i)}{L_0}$,

and $\frac{L_f(i+j+i*j)}{L_m(i+j)}$, respectively, where L_m indicates the likelihood of the models that include the main

effects of markers i and j, L_0 indicates the likelihood of the null model (i.e., model without marker

covariates), and L_f indicates the likelihood of the full model that include the interaction effects

between markers i and j.

Composite interval mapping (CIM)

CIM was performed using the cim function of the R package qtl (ver. 1.60, Broman et al. 2003). The

default parameter values were used. Briefly, the expectation-maximization algorithm was used, three

covariates were used, and the window size was 10 cM. CIM was used to detect the main QTL effects.

Multiple interval mapping

MIM was conducted using QTL cartographer (ver. 2.5_011, Wang et al. 2012). We followed the

algorithm of Laurie et al. (2014) which was illustrated in the section "Instruction for using the

procedures in Windows QTL Cartographer" of the article. The algorithm consists of multiple steps,

including a forward search of the main and interaction effects, optimization of QTL positions, and

testing of QTL effects. These steps require manual program operation, which hampers massive

simulation analysis. Out of 40 replications of the simulations (10 replications each for simulation

Scenarios 1–4, described later), the program abnormally ended in nine replications. These replications

were excluded when summarizing the performance of the MIM.

BayesC

BayesC (Habier et al. 2011) was performed using the vigor function of R package VIGoR (ver. 1.1.0, Onogi and Arakawa 2022). This package adopts fast variational Bayesian algorithms to estimate the posterior distributions of Bayesian alphabets. For genotype i, the model can be written as:

$$y_i = \mu + \sum_p \beta_p x_{p,i} + \sum_p \sum_{k \neq p} \beta_{pk} x_{p,i} x_{k,i} + e_i$$

where y_i is the phenotypic values, μ is the intercept, β_p is the effect of marker p, $x_{p,i}$ is the genotype of marker p, β_{pk} is the interaction effect between marker p and k, and e_i is the residual. The prior distributions of β_p and β_{pk} were

$$\beta_p \sim N(0, \ \sigma_M^2) \ \text{if} \ \gamma_p = 1, \text{and} \ \beta_p = 0 \ \text{if} \ \gamma_p = 0, \text{and}$$

$$\beta_{pk} \sim N(0, \ \sigma_I^2) \ \text{if} \ \gamma_{pk} = 1, \text{and} \ \beta_{pk} = 0 \ \text{if} \ \gamma_{pk} = 0,$$

where γ_p and γ_{pk} are the marker and marker combination inclusion probabilities, respectively. The prior distributions of γ_p and γ_{pk} were the Bernoulli distributions with hyperparameters $\kappa_M = 0.05$ and $\kappa_I = 0.001$, respectively. Default values were used for the other hyperparameters. Bootstrapping was applied to make the inferences on the inclusion probability more robust. That is, the model was fitted to bootstrap samples, and the posterior means of the inclusion probabilities were averaged across the bootstrap samples. In this study, 1000 bootstrap samples were used.

Simulations

The usefulness of SHAP-XGB was first investigated using simulations that mimicked biparental F_2 and RIL populations. The conditions of the simulations were as follows: The number of chromosomes was 8, the length of each chromosome was 120 cM, the number of markers (SNPs) per chromosome was 20, the number of QTLs was five, and the number of individuals was 200. The QTLs were assigned so as to be one QTL per chromosome, and chromosomes harboring the QTLs were randomly selected. Among the 10 combinations of QTLs, half were assumed to have interaction effects, and the combinations with interaction effects were randomly selected. The main and interaction effects of the

QTLs were generated using the standard normal distribution. The broad-sense heritability was set at

0.8.

We simulated six scenarios using these conditions: (1) F_2 population with only additive effects

(QTL genotypes were coded as −1, 0, and 1). (2) RIL population with additive effects only. (3) F_2

population with only complete dominance effects (QTL genotypes were coded as 0, 1, and 1). (4) F_2

population with only overdominance effects (QTL genotypes were coded as 0, 1, and 0). (5) F_2

population with only additive effects and with epistatic variances increased by setting the variance of

interaction effects to 16. (6) F_2 population had only additive effects, but the number of QTLs with

nonzero additive effects was limited to two (i.e., three QTLs only had interaction effects without main

effects). When only additive (dominance) effects were simulated, interaction effects were simulated as

additive-by-additive (dominance-by-dominance) effects by multiplying the corresponding QTL

genotypes. A summary of the simulation scenarios and realized proportions of variances due to the

additive and interaction effects are presented in Table 1. The QTLs and marker genotypes were

generated using the R package qtl (ver. 1.60). QTL effects and phenotypes were generated using house-

developed R scripts. The number of replications per scenario was 10, which was determined by

considering the MIM manual operations.

The accuracy of QTL detection was evaluated using receiver operating characteristic (ROC)

curves and the area under the curve (AUC). When drawing ROC curves, markers located within 20

cM of the QTLs were regarded as positive markers. For interaction effects, combinations of markers

located within 20 cM of QTLs relevant to the interactions were regarded as positive combinations.

ROC curves and AUC were calculated using the R package pROC ver. 1.18.0 (Robin et al. 2011). The

global SHAP and global SHAP interaction scores were regarded as QTL signals for SHAP-XGB. The

p values were regarded as the signals for LRT and CIM, and the bootstrap means of the inclusion

probabilities (γ_p and γ_{pk}) were regarded as the signals for BayesC. The output results were lists of

positive markers (for the main effects) and combinations of markers (for the interaction effects) for

MIM, and ROC curves could not be drawn. Instead, the true and false positive rates were calculated.

Mapping dominance effects

Whereas the cim function of the qtl package and MIM by the QTL cartographer test the additive and dominance effects automatically, LRT by the interactionPval function, BayesC by vigor, and SHAP-XGB need a specification which effect is targeted. For LRT, the argument "model" of the function was set as "co" (codominant) to test the additive effects, "do" to test the dominance effects, and "ov" to test the overdominance effects. For BayesC and SHAP-XGB, genotype codes were modified depending on the target effects. The genotypes were coded as $[-1, 0, 1]$, $[0, 1, 1]$, and $[0, 1, 0]$ to test the additive, complete dominance, and over dominance effects, respectively.

MIM tested multiple combinations of interaction effects, that is, additive-by-additive, additive-by-dominance, and dominance-by-dominance. Thus, even when only dominance-by-dominance interactions were simulated, the results occasionally included additive-by-additive or additive-by-dominance interactions for the correct marker combinations. Because it was too strict to regard these results as false positives, they were considered true positives.

Algorithm assessment

The performance of the compound and pure global SHAP scores in detecting QTL main effects was compared using simulation scenarios (1) (F_2 population with only additive effects), (5) (F_2 population with only additive effects and with increased epistatic variances), and (6) (F_2 population with only additive effects, but the number of QTLs that have non-zero additive effects was limited to two).

Two additional settings of XGBoost were compared to investigate the effects of the degree of fit of the XGBoost models to the data on the QTL detection power. In the underfitting setting, the learning rate "eta" was set to 0.1 (default is 0.3) and the "max_depth," which specifies the maximum depth of trees, was set to 3 (the default is 6). The "early_stopping_rounds" was set to 3. In the overfitting setting, the hyperparameters were the default values and the number of rounds (boosting iterations) was set to 50 without early stopping. Simulation Scenarios 1 and 2 were used for comparison.

Real data

The real data were obtained from Onogi et al. (2016). Briefly, days to heading of rice were evaluated

for 174 Koshihikari/Kasalath//Koshihikari backcross inbred lines (BILs) and their parental lines. The

experiments were conducted in six experimental fields in Japan and Vietnam over three years, resulting

in nine environments. The nine environments are summarized in Table 2, and details were provided in

Table S1 in Onogi et al. (2016). The lines were genotyped for 162 biallelic markers.

Dependency plots

SHAP interaction values were calculated for each instance (line) because SHAP was originally

proposed as a local importance measure. Thus, the interactions between QTLs (markers) can be

visualized by plotting the SHAP interaction values of the marker combinations of interest for all lines.

These plots are referred to as dependency plots. Dependency plots were constructed using the R

package SHAPforxgBoost (ver. 0.1.1) with some modifications.

312 **Table 1** Simulation scenarios[1].

Scenario	Population	Main and interaction effects	Prop. of additive variance	Prop. of epistatic variance
1	F$_2$ lines	Additive and additive-by-additive	0.476 ± 0.104	0.328 ± 0.112
2	Recombinant inbred lines	Additive and additive-by-additive	0.408 ± 0.175	0.385 ± 0.176
3	F$_2$ lines	Complete dominance and dominance-by-dominance	0.417 ± 0.222	0.407 ± 0.212
4	F$_2$ lines	Overdominance and dominance-by-dominance	0.462 ± 0.134	0.322 ± 0.130
5	F$_2$ lines	Additive and additive-by-additive	0.115 ± 0.037	0.679 ± 0.032
6	F$_2$ lines	Additive and additive-by-additive (three QTLs had only interaction effects)	0.313 ± 0.178	0.476 ± 0.176

[1]All populations were assumed to be derived from biparental crosses. The other conditions common

to the scenarios were as follows: the number of chromosomes was 8, the length of each chromosome

was 120 cM; the number of SNPs per chromosome was 20, the number of QTLs was 5, the number of

individuals was 200, and the number of QTL interactions was 5. The main and interaction effects of

QTLs were generated from the standard normal distribution, except for the interaction effect of

Scenario 5, where the variance increased to 16.

Table 2 Nine environments in the rice heading date data[1].

Environment name	Location	Year	Transplanting date
Tsukuba2007	Tsukuba (N36°01'/E140°06')	2007	8 Jun.
Fukuoka2008	Fukuoka (N33°37' /E130°28')	2008	24 Jun.
HaNoi2008	HaNoi (N21°01' /E105°80')	2008	3 Jul.
Ishigaki2008	Ishigaki (N24°23' /E124°11')	2008	17 Jul.
Ishikawa2008	Ishikawa (N36°30' /E136°36')	2008	16 May
ThaiNguyn2008	ThaiNguyn (N21°35' /E105°49')	2008	15 Jul.
Tsukuba2008E	Tsukuba (N36°01' /E140°06')	2008 (Early sowing)	25 Apr.
Tsukuba2008L	Tsukuba (N36°01' /E140°06')	2008 (late sowing)	9 Jul.
Tsukuba2009	Tsukuba (N36°01' /E140°06')	2009	21 May

321   [1]Details of these environments are provided in Table S1 of Onogi et al. (2016)

Results

There are two choices when applying SHAP values for mapping QTL main effects: using compound global SHAP scores that contain interaction scores or using pure global SHAP scores from which interaction scores are removed. We first compared these two types of values for mapping the QTL main effects in simulation Scenarios 1, 5, and 6. Scenarios 1 and 5 differed in the proportion of phenotypic variance explained by epistatic variance (0.328 vs. 0.679 on average; Table 1), and Scenarios 1 and 6 differed in the number of QTLs that had main effects (5 vs. 2; Table 1). The AUC values and ROC curves were comparable between the compound and pure scores (Fig. S1). A slight increase in the detection power was observed for the compound scores in Scenario 5. In contrast, the false positive rates increased slightly for the compound scores in Scenario 6. This is reasonable because the three QTLs lacked main effects and only had interaction effects in Scenario 6. Thus, we concluded that the compound SHAP scores were comparable with or even better than the pure SHAP scores in QTL mapping as long as the QTLs that have non-zero main effects are involved in the interactions. Hereafter, we used compound scores to map the main QTL effects.

Next, we investigated the effect of the fitting degree of the XGBoost models on QTL mapping accuracy. We compared the three settings that led to underfitting, moderate fitting, and overfitting in simulation Scenarios 1 and 2. The SHAP values were either underestimated or overestimated when underfitting or overfitting was induced (Fig. S3). Nevertheless, as the relative values between the markers did not vary significantly, the AUC values and ROC curves were hardly changed by the fitting status (Fig. S3). We used a setting that induced a moderate fit.

In the comparison analyses to map the main QTL effects, SHAP-XGB generally showed higher or comparable AUC values than the other methods (Fig. 1). However, in the F_2 populations (Scenarios 1, 3, and 4), SHAP-XGB showed lower true positive rates than CIM or BayesC when the false positive rates were less than 0.01. In mapping dominance effects (Scenarios 3 and 4), SHAP-XGB showed comparable or only slightly decreased AUC values when additive effects were targeted compared to when dominance effects were targeted. These results were in contrast to those of BayesC, where the

misspecification of target effects resulted in a more prominent decrease in accuracy. These results

suggested that SHAP-XGB is insensitive to the effect type.

Fig. 1 Area under the curve (AUC) values and receiver operation characteristics (ROC) curves in

mapping main QTL effects. The top and bottom panels are the AUC values and ROC curves,

respectively. The ROC curves are the mean curves of 10 replications. Scenarios 1 and 2 simulated F_2

and recombinant inbred line populations, respectively, and Scenarios 3 and 4 simulated F_2 populations

that comprised only complete and over dominance effects, respectively. Small letters a and d in

Scenarios 3 and 4 indicate methods that targeted the additive and dominance effects, respectively. LRT:

likelihood ratio test, CIM: composite interval mapping, SHAP-XGB: Shapley additive explanations-

assisted XGBoost, MIM: multiple interval mapping.

In the comparison analyses to map QTL interaction effects, SHAP-XGB showed higher AUC

values than the other methods (Fig. 2), and the true positive rates of SHAP-XGB were superior to those

of the other methods in any range of false positive rates (Fig. 2). As observed in the main effect

detection, SHAP-XGB was found to be insensitive to misspecification of target effects, that is,

dominance-by-dominance effects can be detected even when additive-by-additive effects were targeted.

Interactions between markers (QTLs) can be interpreted using dependency plots. The SHAP

interaction values (local importance metrics) for a replication of Scenario 2 are shown in Fig. 3. The

effects of two QTLs, *QTL2* and *QTL3* tagged with *D3M19* and *D4M11* markers, respectively, were

inverted depending on the genotype of *QTL4* tagged with *D6M1*. These figures illustrate that

dependency plots using local importance scores can facilitate the interpretation of interactions.

Fig. 2 Area under the curve (AUC) values and receiver operation characteristics (ROC) curves in

mapping QTL interaction effects. The top and bottom panels are the AUC values and ROC curves,

respectively. The ROC curves are the mean curves of 10 replications. Scenarios 1 and 2 simulated F_2

and recombinant inbred line populations, respectively, and Scenarios 3 and 4 simulated F_2 populations

that comprised only complete and over dominance effects, respectively. Small letters a and d in

Scenarios 3 and 4 indicate methods that targeted the additive and dominance effects, respectively. LRT:

likelihood ratio test, SHAP-XGB: Shapley additive explanations-assisted XGBoost, MIM: multiple

interval mapping.

Fig. 3 Dependencies between QTLs in a simulated dataset. The Shapley additive explanations (SHAP) interaction values of each line are shown as dots. The left panel shows the interaction between the 2nd and 4th QTLs, and the right panel shows that between the 3rd and 4th QTLs. *D3M19*, *D4M11*, and *D6M1* are the markers linked to the 2nd, 3rd, and 4th QTLs, respectively. Genotypes of *D6M1* are shown using different colors. Dots are scattered towards the x-axis. The regression lines were added for each genotype of *D6M1*. The shades denote the standard errors. The left panel shows that, when the genotype of *D6M1* is −1, SHAP interaction values (i.e., phenotypes) tend to decrease as the genotype of *D3M19* changes from −1 to 1, and when the genotype of *D6M1* is 1, SHAP interaction values tend to increase. The right panel shows inverted tendencies between *D4M11* and *D6M1*.

The XGBoost models explained 94.3% (± 1.9%) of the phenotypic variance with the setting of moderate fitting in the real rice data analyses. SHAP-XGB and the compared methods showed similar results in mapping the main QTL effects (Fig. S4 and Fig. 4), and markers linked to *Hd6* (chr. 3, Takahashi et al. 2001), *Hd1* (chr. 6, Yano et al. 1997), and *DTH8* (chr. 8, Wei et al. 2010) showed higher signals for each method. The BayesC results were generally stringent (or conservative) when mapping epistatic QTLs (Fig. S5 and Fig. 5). LRT showed wide positive regions for each interacting combination, probably because of linkage disequilibrium (LD) between the markers (Fig. S5 and Fig. 5). In contrast, SHAP-XGB could pinpoint interacting combinations within reasonably narrow margins. The global SHAP interaction scores in each environment showed interesting trends across latitudes (Fig. 6). Interactions between *Hd1* and *Hd6*, *Hd1* and *DTH8*, and *Hd1* and *Hd2*/*OsPRR37* (Yano et al.

1997; Koo et al. 2013) showed higher scores at higher latitudes (environments Tsukuba2007, Tsukuba2008E, Tsukuba2009, Ishikawa2008, and Fukuoka2008), but lower scores at lower latitudes (environments Ishigaki2008, HaNoi2008, and ThaiNguyn2008). In contrast, the interaction between *DTH8* and *Hd2* showed larger scores at lower latitudes but lower scores at higher latitudes.

Fig. 4 Comparison of QTL mapping methods to detect main effects in the rice heading date data. The y-axis denotes the QTL signals (LRT, −log10p; CIM, LOD score; BayesC, posterior inclusion probability; SHAP-XGB, global SHAP score). The x-axis denotes the marker positions. The neighboring chromosomes are distinguished using different colors. The vertical bars indicate the four major QTLs (left to right, *Hd6*, *Hd1*, *Hd2*, and *DTH8*). The horizontal bars in LRT and CIM are the 5% thresholds after the Bonferroni correction and the 95% thresholds obtained by permutation tests, respectively. Three representative environments (Tsukuba2007, Fukuoka2008, and HaNoi2008) are presented. The full results are shown in Fig. S4. LRT: likelihood ratio test, CIM: composite interval mapping, SHAP-XGB: Shapley additive explanations-assisted XGBoost.

Fig. 5 Comparison of QTL mapping methods to detect interaction effects in the rice heading date data. The y- and x-axes denote the relative positions of markers. The signal levels obtained with each QTL mapping method (LRT, −log10p; BayesC, posterior inclusion probability; SHAP-XGB, global SHAP interaction score) are shown using warm colors. The gray broken lines indicate the four major QTLs (left to right or bottom to top, *Hd6*, *Hd1*, *Hd2*, and *DTH8*). Three representative environments (Tsukuba2007, Fukuoka2008, and HaNoi2008) are presented. The full results are shown in Fig. S5. LRT: likelihood ratio test, SHAP-XGB: Shapley additive explanations-assisted XGBoost.

Fig. 6 Summary of the global Shapley additive explanations (SHAP) interaction scores obtained in the

rice heading date data. The y-axis denotes the global SHAP interaction scores, which are the averages

of the absolute SHAP interaction values across all lines. The global SHAP interaction scores were

horizontally jittered according to the densities of scores. The interaction values for the combinations

of markers that were located within 20 cM from the known QTLs (*Hd6*, *Hd1*, *Hd2*, and *DTH8*) were

regarded as the signals for the corresponding QTL interactions and highlighted using different colors

and point shapes. The interactions between neighboring markers (< 20 cM apart) were removed for

ease of visualization.

The interaction between *Hd1* and three QTLs (*DTH8*, *Hd6*, and *Hd2*) in Tsukuba2009 and the

interaction between *DTH8* and *Hd2* in Ishigaki2008 were visualized using dependence plots (Fig. 7).

The Koshihikari allele (1) of *DTH8* and *Hd6* tended to shorten days to heading at Tsukuba2009 when

the Koshihikari alleles were harbored at *Hd1*, whereas the Kasalath allele (−1) tended to prolong days

to heading when the Kasalath alleles were harbored at *Hd1*. Contrasting tendencies were observed for

the interactions between *Hd1* and *Hd2* in Tsukuba2009 and *Hd2* and *DTH8* in Ishigaki2008.

Fig. 7 Dependencies between major QTLs shown with the Shapley additive explanations (SHAP) interaction values. The left top, right top, and left bottom panels show the dependency between *Hd1* and the three QTLs (*DTH8*, *Hd6*, and *Hd2*) at Tsukuba2009, respectively. The right bottom panel shows the dependency between *DTH8* and *Hd2* at Ishigaki2008. Each dot represents the SHAP interaction values for each line. Marker genotypes were regarded as the QTL genotypes. These markers were selected when the corresponding marker combinations showed the highest global SHAP interaction scores for the QTL interactions. For each panel and each QTL (marker), 1 and −1 indicate the Koshihikari and Kasalath alleles, respectively.

Discussion

455
We applied SHAP-XGB and SHAP-assisted XGBoost to QTL mapping in biparental populations and

compared it with conventional mapping methods in this study. SHAP-XGB generally showed

comparable accuracy in finding QTL main effects with conventional methods and higher accuracy in

finding epistatic QTL effects. The dependency plots enabled easy interpretation of the interactions.

These results suggested that SHAP-XGB complements existing methods, particularly for identifying

novel interactions between QTLs.

The simulation results revealed several properties of the SHAP-XGB. First, compound and pure

global SHAP scores worked similarly to map the main effects of QTLs. While the compound scores

showed slightly increased detection power when epistasis variances were larger, they also showed a

slight increase in false positives when QTLs with interaction effects lacked the main effects. Thus,

using compound scores may be advantageous as long as QTLs with non-zero main effects are involved

in the interactions. Using compound scores may circumvent the reduction in pure scores due to LD, as

described later. Second, the degree of fit of the XGBoost models did not affect the QTL detection

power. This is because the ROC curves and AUC values were calculated based on the relative rankings

of markers that did not differ significantly depending on the fitting status. However, the result of over-

or underfitting is the inflation or deflation of SHAP values, which in turn may make interpretations of

SHAP values difficult. Thus, careful tuning of the fitting degree is recommended to improve

interpretability. Finally, SHAP-XGB is robust to the misspecification of target effects, i.e., additive or

dominance. That is, dominance main effects or dominance-by-dominance interaction effects can be

mapped as accurately as in the existing methods with correct target specifications even when additive

effects are targeted. This property may be advantageous; however, it may be disadvantageous in some

cases because the gene actions of the detected QTLs cannot be identified.

There are ways to measure feature importance for gradient boosting other than SHAP, including

"gain" (the average increase of the objective function when the feature is added to trees) and "split

count" (the number of counts that trees split using the feature). Although these indices are provided by

the R package of XGBoost, we did not adopt them because they do not hold the consistency property (Lundberg et al. 2018) and the package did not support them as importance of feature interactions. Nevertheless, considering that there are various indices to measure feature importance, such as local interpretable model-agnostic explanations (LIME) (Ribeiro et al. 2016) and the Shapley–Taylor interaction index (Sundararajan et al. 2020), it would still be worth investigating which indices are suitable for QTL mapping and whether the properties revealed here are applicable when SHAP is replaced with other indices. Such investigations are required to mitigate the negative effects of LD, as will be discussed later.

Although the simulated and real data analyses consistently suggest the usefulness of SHAP-XGB in QTL mapping, the marker genotypes have an undesirable property, i.e., LD between markers, for evaluating feature importance. Because XGBoost randomly samples markers to build a tree f, QTL effects are diluted with the markers linked to the QTLs, consequently decreasing the Shapley values of each linking marker. In fact, the global SHAP scores and SHAP interaction values in Fig. 4 and 6 shrank to 0 (2.0 at most) considering that the units of the scores/values are the same as the phenotypic values (i.e., days). These decreased values were probably due to the correlated marker genotypes. The LD between the markers affected another aspect. In rice heading date analyses, interactions between neighboring markers were occasionally assigned large SHAP interaction values, resulting in a reduction in the main pure scores (these interactions were removed from Fig. 6 for ease of visualization). However, such interactions are caused by LD between neighboring markers and thus should be interpreted as part of the main QTL effects. Although this decrease in pure main effects can be circumvented using compound scores for mapping, caution is required when interpreting pure scores. It has also been pointed out that approximations of Shapley values with SHAP become inaccurate when features are correlated (Aas et al. 2021) because the evaluations of the expectation in Eq. (2) assume independence between the features. Thus, taken together, LD would affect the detection power of SHAP-XGB. One solution is to replace SHAP with other indices that consider the correlations among features (Aas et al. 2021).

Interactions between the major heading date QTLs weakened at lower latitudes (Fig. 6 and Table 1). These results were reasonable because *Hd6*, *Hd2*, and *DTH8* are involved in heading date variation under long-day conditions (Hori et al. 2016; Wei et al. 2020). Interactions between *Hd1* and three QTLs (*Hd6*, *Hd2*, and *DTH8*) are plausible under natural long-day conditions. For example, interactions between functional *Hd6* alleles (e.g., Kasalath allele) and *Hd1* alleles (e.g., Nipponbare allele) were reported under long-day conditions (Ogiso et al. 2010). Interactions between *Hd1* and *Hd2* under long-day conditions (Lin et al. 2000), and interactions among *Hd1*, *Hd2*, *DTH8*, and *Ghd7* under natural long-day conditions were also reported (Zhang et al. 2019). The interactions between *Hd2* and *DTH8*, which is prominent in lower latitudes, are interesting but also plausible because interactions between the functional/non-functional alleles of *Hd2* and *DTH8* alleles under natural short-day conditions have been reported (Zhang et al. 2019). Thus, these results indicated the usefulness of epistatic QTL mapping using SHAP-XGB for identifying novel interactions.

SHAP-XGB may be applicable to GWAS; however, the computational time and required memory size will increase. The increase in computational time depends on the XGBoost fitting and SHAP value calculations. Fitting XGBoost to the rice real data ($n = 176$ and $p = 162$) took 0.111 ± 0.055 seconds using a machine with Intel Corei7-13700k (3.40 GHz), 24 logical cores, and 128GB RAM, and calculation of SHAP and SHAP interaction scores took 0.007 ± 0.014 and 0.316 ± 0.144 seconds, respectively. We generated a feature matrix with $n = 1,760$ and $p = 162,000$ from the standard normal distribution and fitted the SHAP-XGB to investigate the scalability of SHAP-XGB. XGBoost took 51.34 seconds, and the calculation of SHAP took 430.89 seconds using the same machine, which is sufficiently practical. However, the calculation of the SHAP interaction scores failed because of a memory allocation error, and this error could not be fixed until p decreased to 2,000. SHAP interaction scores are calculated for all marker combinations in all samples (i.e., an $n \times p \times p$ array is created), and the memory requirement may be a major bottleneck. Thus, it may be necessary to prune markers (SNPs) using, for example, t-tests or sure independence screening (Fan and Lv 2008) in practical GWAS applications. Population stratification is another issue in GWAS. One solution would be to

adjust the phenotypic values with population stratification before application.

The lack of thresholds for determining significant markers is a practical drawback for each

application of SHAP-XGB (QTL mapping and GWAS). Permutation tests are one possible solution.

Permutation tests are generally used to determine the thresholds for IM and CIM (e.g., Broman et al.

2003) and have been applied to detect gene-gene interactions based on the Shapley-based interaction

score and neural networks (Cui et al. 2022). However, further investigation is required to determine

whether these tests work as expected.

Acknowledgements

542   The authors would like to thank Ryukoku University for providing financial support.

Declarations

Funding

546   This study was supported by Ryukoku University.

Competing interests

548   The authors declare no competing interests.

Data availability

550   Real data are shared at https://github.com/Onogi/HeadingDatePrediction. Simulation data are

551   available upon request.

Code availability

553   R scripts are available upon requests.

Author contributions

555   TI and AO collaborated in analyzing the data and drafting the manuscript. AO conceived the study.

Ethics approval

557   Not applicable

Consent to participate

Not applicable

Consent to publication

Not applicable

References

Aas K, Jullum M, Løland A (2021) Explaining individual predictions when features are dependent:
More accurate approximations to Shapley values. Artif Intell 298:103502.
https://doi.org/10.1016/j.artint.2021.103502

Broman KW, Wu H, Sen S, Churchill GA (2003) R/qtl: QTL mapping in experimental crosses.
Bioinformatics 19:889–890. https://doi.org/10.1093/bioinformatics/btg112

Chen T, Guestrin C (2016) Xgboost: A scalable tree boosting system. Proceedings of the 22nd ACM
SIGKDD international conference on knowledge discovery and data mining 785–794.
https://doi.org/10.1145/2939672.2939785

Crawford L, Zeng P, Mukherjee S, Zhou X (2017) Detecting epistasis with the marginal epistasis test
in genetic mapping studies of quantitative traits. PLOS Genet 13:e1006869.
https://doi.org/10.1371/journal.pgen.1006869

Cui T, El Mekkaoui K, Reinvall J, Havulinna AS, Marttinen P, Kaski S (2022) Gene-gene interaction
detection with deep learning. Commun Biol 5:1238. https://doi.org/10.1038/s42003-022-
04186-y

Falconer DS, Mackay TFC (1996). Introduction to quantitative genetics (fourth edition). Pearson
Education Ltd, London.

Fan J, Lv J (2008) Sure independence screening for ultrahigh dimensional feature space. J R Stat Soc
B 70:849–911. https://doi.org/10.1111/j.1467-9868.2008.00674.x

Fujimoto K, Kojadinovic I, Marichal J (2006) Axiomatic characterizations of probabilistic and
cardinal-probabilistic interaction indices. Games Econ Behav 55:72–99.
https://doi.org/10.1016/j.geb.2005.03.002

González JR, Armengol L, Solé X, Guinó E, Mercader JM, Estivill X, Moreno V (2007) SNPassoc:

An R package to perform whole genome association studies. Bioinformatics 23:644–645.
https://doi.org/10.1093/bioinformatics/btm025

Habier D, Fernando RL, Kizilkaya K, Garrick DJ (2011) Extension of the Bayesian alphabet for
genomic selection. BMC Bioinform 12:186. https://doi.org/10.1186/1471-2105-12-186

Hori K, Matsubara K, Yano M (2016) Genetic control of flowering time in rice: Integration of
Mendelian genetics and genomics. Theor Appl Genet 129:2241–2252.
https://doi.org/10.1007/s00122-016-2773-4

Jansen RC, Stam P (1994) High resolution of quantitative traits into multiple loci via interval
mapping. Genetics 136:1447–1455. https://doi.org/10.1093/genetics/136.4.1447

Johnsen PV, Riemer-Sørensen S, DeWan AT, Cahill ME, Langaas M (2021) A new method for
exploring gene-gene and gene-environment interactions in GWAS with tree ensemble methods
and SHAP values. BMC Bioinform 22:230. https://doi.org/10.1186/s12859-021-04041-7

Koo BH, Yoo SC, Park JW, Kwon CT, Lee BD, An G, Zhang Z, Li J, Li Z, Paek NC (2013) Natural
variation in *OsPRR37* regulates heading date and contributes to rice cultivation at a wide range
of latitudes. Mol Plant 6:1877–1888. https://doi.org/10.1093/mp/sst088

Lander ES, Botstein D (1989) Mapping Mendelian factors underlying quantitative traits using RFLP
linkage maps. Genetics 121:185–199. https://doi.org/10.1093/genetics/121.1.185

Laurie C, Wang S, Carlini-Garcia LA, Zeng ZB (2014) Mapping epistatic quantitative trait loci.
BMC Genet 15:112. https://doi.org/10.1186/s12863-014-0112-9

Li H, Ribaut JM, Li Z, Wang J (2008) Inclusive composite interval mapping (ICIM) for digenic
epistasis of quantitative traits in biparental populations. Theor Appl Genet 116:243–260.
https://doi.org/10.1007/s00122-007-0663-5

Li Z, Sillanpää MJ (2012) Estimation of quantitative trait locus effects with epistasis by variational
Bayes algorithms. Genetics 190:231–249. https://doi.org/10.1534/genetics.111.134866

Lin HX, Yamamoto T, Sasaki T, Yano M (2000) Characterization and detection of epistatic
interactions of 3 QTLs, *Hd1*, *Hd2*, and *Hd3*, controlling heading date in rice using nearly
isogenic lines. Theor Appl Genet 101:1021–1028. https://doi.org/10.1007/s001220051576

Shapley LS (1953) A value for n-person games. Contributions to the Theory of Games 2.28, pp.
307–317

Lundberg SM, Lee S (2017) A unified approach to interpreting model predictions. Nips'17.
Proceedings of the 31st international conference on neural information processing systems
4768–4777. Available from https://dl.acm.org/doi/10.5555/3295222.3295230

Lundberg SM, Erion GG, Lee S (2018) Consistent individualized feature attribution for tree
ensembles. arXiv preprint arXiv:1802.03888. https://doi.org/10.48550/arXiv.1802.03888

Ogiso E, Takahashi Y, Sasaki T, Yano M, Izawa T (2010) The role of casein kinase II in flowering
time regulation has diversified during evolution. Plant Physiol 152:808–820.
https://doi.org/10.1104/pp.109.148908

Onogi A, Watanabe M, Mochizuki T, Hayashi T, Nakagawa H, Hasegawa T, Iwata H (2016) Toward
integration of genomic selection with crop modelling: The development of an integrated
approach to predicting rice heading dates. Theor Appl Genet 129:805–817.
https://doi.org/10.1007/s00122-016-2667-5

Onogi A, Arakawa A (2022) An R package VIGoR for joint estimation of multiple linear learners
with variational Bayesian inference. Bioinformatics 38:3306–3309.
https://doi.org/10.1093/bioinformatics/btac328

Ribeiro MT, Singh S, Guestrin C (2016) 'Why should i trust you?' Explaining the predictions of any
classifier. Proceedings of the 22nd ACM SIGKDD international conference on knowledge
discovery and data mining 1135–1144. https://doi.org/10.1145/2939672.2939778

Robin X, Turck N, Hainard A, Tiberti N, Lisacek F, Sanchez JC, Müller M (2011) pROC: an open-
source package for R and S+ to analyze and compare ROC curves. BMC Bioinform 12:77.
https://doi.org/10.1186/1471-2105-12-77

Sundararajan M, Dhamdhere K, Agarwal A (2020) The Shapley Taylor interaction index.
Proceedings of the machine learning research 119:9259–9268. Available from
https://proceedings.mlr.press/v119/sundararajan20a.html

Takahashi Y, Shomura A, Sasaki T, Yano M (2001) *Hd6*, a rice quantitative trait locus involved in
photoperiod sensitivity, encodes the alpha subunit of protein kinase CK2. Proc Natl Acad Sci U
S A 98:7922–7927. https://doi.org/10.1073/pnas.111136798

Wang J, Joshi T, Valliyodan B, Shi H, Liang Y, Nguyen HT, Zhang J, Xu D (2015) A Bayesian
model for detection of high-order interactions among genetic variants in genome-wide
association studies. BMC Genom 16:1011. https://doi.org/10.1186/s12864-015-2217-6

Wang S, Basten CJ, Zeng ZB (2012) Windows QTL Cartographer 2.5. Department of Statistics,
North Carolina State University, Raleigh, NC. (http://statgen.ncsu.edu/qtlcart/WQTLCart.htm)

Wei H, Wang X, Xu H, Wang L (2020) Molecular basis of heading date control in rice. aBIOTECH
1:219–232. https://doi.org/10.1007/s42994-020-00019-w

Wei X, Xu J, Guo H, Jiang L, Chen S, Yu C, Zhou Z, Hu P, Zhai H, Wan J (2010) *DTH8* suppresses
flowering in rice, influencing plant height and yield potential simultaneously. Plant Physiol
153:1747–1758. https://doi.org/10.1104/pp.110.156943

Yano M, Harushima Y, Nagamura Y, Kurata N, Minobe Y, Sasaki T (1997) Identification of
quantitative trait loci controlling heading date in rice using a high-density linkage map. Theor
Appl Genet 95:1025–1032. https://doi.org/10.1007/s001220050658

Zeng ZB (1994) Precision mapping of quantitative trait loci. Genetics 136:1457–1468.
https://doi.org/10.1093/genetics/136.4.1457

Zhang B, Liu H, Qi F, Zhang Z, Li Q, Han Z, Xing Y (2019) Genetic interactions among *Ghd7*,
Ghd8, *OsPRR37* and *Hd1* contribute to large variation in heading date in rice. Rice (N Y)
12:48. https://doi.org/10.1186/s12284-019-0314-x

Zhang YM, Xu S (2005) A penalized maximum likelihood method for estimating epistatic effects of
QTL. Heredity 95:96–104. https://doi.org/10.1038/sj.hdy.6800702

Short Legends of Supporting Information

Fig. S1: Comparison of area under the curve (AUC) values and receiver operating characteristics

(ROC) curves between compound and pure global SHAP scores.

Fig. S2: Comparison of the XGBoost settings in the simulated data sets.

Fig. S3: Comparison of the compound global SHAP scores obtained using the different XGBoost

settings.

Fig. S4: Comparison of QTL mapping methods to detect main effects in the rice heading date data.

Fig. S5: Comparison of QTL mapping methods to detect interaction effects in the rice heading date

data.