A mechanosensitive channel Pkd2 modulates the assembly of myosin II and actin filaments

in the cytokinetic contractile ring

Pritha Chowdhury, Debatrayee Sinha, Abhishek Poddar, Madhurya Chetluru and Qian Chen[*]

Department of Biological Sciences

The University of Toledo

2801 Bancroft St, Toledo, OH 43606

* Correspondence: qian.chen3@utoledo.edu

Keywords: Cytokinesis, Fission yeast, Myosin, Actin, Polycystin, Pkd2

Running title: Pkd2 regulates contractile ring assembly

Abstract

Cytokinesis, the last step in cell division, leads to the separation of daughter cells. This

evolutionarily conserved process requires the force generated by an actomyosin contractile ring

assembled at the equatorial plane. A newly discovered ion channel Pkd2 localizes to the

equatorial plane and its activation by membrane-stretching promotes the cytokinetic calcium

spikes in the fission yeast *Schizosacchromyces pombe* cells. Here, we investigated whether Pkd2

plays a direct role in the assembly and constriction of the contractile ring through quantitative

fluorescence microscopy and genetic analyses. The localization of type II myosin Myo2, its

regulatory light chain Rlc1 and actin filaments all increased in the contractile ring of a

hypomorphic mutant *pkd2-81KD*, compared to the wild type. This was accompanied by a

significant increase of the intracellular concentration of Rlc1 and actin filaments. A strong

negative interaction existed between *pkd2-81KD* and the temperature-sensitive mutant *myo2-E1*.

The *pkd2 myo2* double mutant cells failed to assemble a complete contractile ring and separate

efficiently, even at the permissive temperature. We conclude that Pkd2 modulates the assembly

of myosin and actin in the contractile ring, suggesting that this ion channel mediates a novel

pathway to regulate the actin cytoskeletal dynamics during cytokinesis. (197 words)

Introduction

Cytokinesis is the last stage of cell division when the daughter cells separate physically through

mechanical force. In most eukaryotes, this process requires an actomyosin contractile ring

specifically assembled at the equatorial division plane. Decades of work have discovered several

key signaling pathways that regulate the assembly and constriction of the contractile ring (for a

review see (Pollard and Wu, 2010)). They include those mediated by Polo kinase, the Central

Spindlin complex, and RhoA. These pathways in turn reorganize the actin cytoskeletal structures

to assemble the cytokinetic contractile ring which consists of dozens actin-binding proteins

including formin, cofilin and myosins (Balasubramanian et al., 1998; Chen and Pollard, 2011;

Kovar et al., 2011; Wu and Pollard, 2005). Among them, type II non-muscle myosins (Myosin

II) play a critical role by driving both the assembly of actin filaments in the contractile ring and

the ensuing ring constriction and disassembly of actin filaments from the ring (Balasubramanian

et al., 1998; Laplante et al., 2015; Malla et al., 2022). Myosin II is a motor protein whose

activities depend on its regulatory light chain (Rlc). However, it remains unknown how myosin

II toggles switch its initial function of assembling the contractile ring to the later role of

disassembly as cytokinesis progresses.

We recently made a surprising finding that an essential ion channel Pkd2 is required for

cytokinesis of the model organism fission yeast *Schizosacchromyces pombe* (Morris et al., 2019;

Palmer et al., 2005). As a transmembrane protein, Pkd2 is trafficked to the plasma membrane

where its distribution is cell-cycle dependent (Malla et al., 2023). During interphase, this channel

mostly localizes at the cell tips. Upon entering cytokinesis, Pkd2 is recruited to the cleavage

furrow where it stays until the end of cell separation. Pkd2 is an essential gene. A hypomorphic

mutant *pkd2-81KD*, which replaces the endogenous promoter with an inducible 81nmt1

promoter, reduces the expression of *pkd2* gene by more than 70%. Although this mutation

accelerates the cleavage furrow, the mutant cells often fail to separate at the end of cytokinesis

(Morris et al., 2019).

Pkd2 is a putative Ca^{2+}-permissive channel, activated by the membrane stretching. In

vitro reconstituted Pkd2 in giant unilamellar vesicles (GUV) allows the passage of Ca^{2+} when the

membrane is stretched by hydraulic force (Poddar et al., 2022). In vivo, Pkd2 promotes the

separation calcium spikes, the transient influx of Ca^{2+} following the daughter cell separation at

the end of cytokinesis (Poddar et al., 2021). Besides cytokinesis, Pkd2 also promotes the tip

extension of fission yeast cells during interphase (Sinha et al., 2022). To our knowledge, Pkd2 is

the only essential mechanosensitive ion channels identified so far in yeast.

Evolutionally, Pkd2 belongs to a conserved family of cation channels. Compared to

fission yeast, most vertebrates possess two polycystins, PC-1 and PC-2, encoded by Pkd1 and

Pkd2 genes respectively (for a review, see (Hu and Harris, 2020)). Both are essential but they

play distinct developmental roles (for reviews, see (Hardy and Tsiokas, 2020; Ta et al., 2020)).

PC-1 is a G-protein Coupled Receptor (GPCR) like protein. It possesses a large extracellular

domain (>100 KD) and a 11-helix transmembrane domain. In comparison, PC-2 is a classical

Transient Receptor Potential (TRP) channel with six transmembrane helices. Although PC2 can

oligomerizes by itself into a tetrameric cation channel (Shen et al., 2016), PC-1 and PC-2 are

also capable of assembling into a heterologous tetrameric channel in 3:1 ratio (Su et al., 2018).

Mutations of human polycystins lead to a common genetic disorder, Autosomal Dominant

Polycystic Kidney Disorder (ADPKD) (1994; Mochizuki et al., 1996). However, the cellular

function of polycystins, particularly in the context of cell division (AbouAlaiwi et al., 2011), has

remained largely unknown.

In this study, we focused on answering one question, does Pkd2 have a role in the

assembly of contractile ring during cytokinesis. Using quantitative fluorescence microscopy, we

measured the number of type II myosin Myo2, myosin regulatory light chain Rlc1 and actin in

the contractile ring of a hypomorphic mutant *pkd2-81KD* (Morris et al., 2019), in comparison to

the wild-type. Surprisingly, the number of all three proteins increased in the contractile ring of

the *pkd2* mutant cells. Genetic analysis revealed a negative interaction between two *pkd2*

mutants and the temperature-sensitive *myo2-E1* mutant. The *pkd2-B42 myo2-E1* mutant cells

grew very slowly even at the permissive temperature. These *pkd2 myo2* mutant cells assembled

their contractile ring very slowly. They failed to septate and lysed frequently. Overall, our results

would demonstrate that Pkd2 does modulate the assembly of myosin II and actin in the

contractile ring, suggesting the existence of a novel calcium signaling pathway mediated by this

mechanosensitive channel for reorganizing the actin cytoskeletal structures during cell division.

Materials and methods

Cell culture and Yeast genetics

Yeast cell culture and genetics were carried out according to the standard protocols (Moreno et

al., 1991). Unless specified, YE5S was used in all experiments. For genetic crosses, tetrads were

dissected using Sporeplay+ dissection microscope (Singer, England). Yeast strains used in this

study are listed in Table 1.

92   For ten-fold dilution series of yeast, overnight cultures were diluted and inoculated for

additional 5 hours before being spotting onto YE5s agar plates. The plates were incubated for 2-3

days before being scanned using a photo scanner (Epson).

Microscopy

For live microscopy, fission yeast cells were first inoculated in YE5s liquid media for two days

before being harvested through centrifugation at 1500g for 1 min during exponential phase at a

density between 5×10^6 /ml and 1.0×10^7/ml. The cells were resuspended in 50µl of YE5s and 6 µl

of the resuspended cells were spotted onto a gelatin pad (25% gelatin in YE5s) on a glass slide.

The cells were sealed under a coverslip (#1.5) with VALAP (1:1:1 mixture of Vaseline, lanolin

and paraffin) before imaging.

102   Confocal microscopy was carried out on a spinning disk confocal microscope equipped

with an EM-CCD camera. The Olympus IX71 microscope was equipped with three objective

lenses of 40x (NA=1.30, oil), 60x (NA=1.40, oil) and 100x (NA=1.40, oil), a motorized Piezo Z

Top plane (ASI, USA) and a confocal spinning disk unit (CSU-X1, Yokogawa, Japan). The

images were captured on an Ixon-897 EMCCD camera (Andor). Solid state lasers of 488 and

561nm were used to excite green and red fluorescence proteins respectively. Unless specified,

the cells were imaged with Z-series of 15 slices at a step size of 0.5um.

109   To stain the cell wall with calcofluor, exponentially growing cell culture was fixed with

4% paraformaldehyde, prepared from 16% fresh stock solution (Electron Microscopy Science,

USA). The fixed cells were washed with TEMK buffer (50mM Tris-HCL pH 7.4, 1mM $MgCl_2$,

50mM KCl,1mm EGTA, pH8.0) and stored in 1ml of TEMK overnight at 4ºC. The cells were

stained with 1µg/ml calcofluor, prepared with the stock solution (1mg/ml, Sigma Aldrich) on a

rocker for 10 minutes at the room temperature in the dark. The stained cells were pelleted and

resuspended in 50ul TEMK, 6ul of which was spotted on a glass slide directly and sealed under a

coverslip (#1.5) with VALAP. The images were acquired in an Olympus IX81 microscope

equipped with 60x objective lens (NA=1.42), a LED lamp for bright-field microscopy, and a

mercury lamp for fluorescence microscope. The images were captured on an ORCA C-11440

digital camera (Hamamatsu, Japan), operated with CellSens (Olympus).

Image analysis

We used ImageJ (NIH), its plug-ins and macros to process all the microscopy images. For

quantitative analysis, the time-lapse series were first corrected for X-Y drifting using the plugin

StackReg (Thevenaz et al., 1998). Average intensity projections of Z-series were used for

quantification. The localization of GFP-Myo2 and Rlc1-tdTomato at the contractile ring was

quantified by measuring the fluorescence intensities in a 1µm by 4µm rectangle centered on the

equatorial division plane. The measurements were corrected for the background fluorescence,

measured in a 0.2µm by 4µm rectangle adjoined to the equatorial division plane. To measure

fluorescence intensities of GFP-Lifeact in the contractile ring, we used the fluorescence of Rlc1-

tdTomato to manually segment the contractile ring. To determine the start of the contractile ring

constriction, we constructed kymographs based on the time-lapse series of the contractile ring.

All figures were prepared with Canvas X (ACD Systems) and the plots were prepared

using Origin 2021 (OriginLab, USA).

Results

Pkd2 modulates the assembly of Myo2, Rlc1, and actin in the contractile ring

We first determined whether Pkd2 promotes the localization of Myo2 to the contractile ring. To

test this hypothesis, we compared the localization of the GFP tagged Myo2, expressed from its

endogenous locus, at the equatorial division plane of the hypomorphic *pkd2-81KD* mutant cells

(Morris et al., 2019) to that of the wild type cells using quantitative fluorescence microscopy

(Malla et al., 2022) (Fig. 1A). As in the wild type, GFP-Myo2 localized to a broad band of

cytokinesis nodes in the *pkd2* mutant cells before the nodes condensed into a complete

contractile ring (Fig. 1A and B). However, the contractile ring of the *pkd2* mutant cells matured

and constricted more quickly than that of the wild type cells (Fig. 1B), as reported previously

(Morris et al., 2019). GFP-Myo2 in the contractile ring of *pkd2* mutant cells increased by more

than 20% compared to the wild type cells (Fig. 1C and D). In contrast, the intracellular

concentration of GFP-Myo2, measured by its average fluorescence in a cell, was unchanged in

the mutant cells (Fig. 1E). Thus, Pkd2 modulates the assembly of Myo2 in the contractile ring

during cytokinesis.

Next, we considered whether Pkd2 helps recruit Rlc1, the regulatory light chain of Myo2

(Naqvi et al., 2000), to the contractile ring. This hypothesis is particularly attractive because Rlc1

possesses four EF-hands that bind the Pkd2-permeated Ca^{2+}. As we did above, we compared the

fluorescence of Rlc1 tagged endogenously with the fluorescence protein tdTomato (Rlc1-

tdTomato) in the contractile ring of *pkd2-81KD* cells to that of the wild type cells (Fig. 2A).

Similar to Myo2, the number of Rlc1-tdTomato molecules in the ring of the *pkd2* mutant cells

was ~20% higher than that of the wild type (Fig. 2B). Surprisingly, there was also a significant

increase (26%) of the intracellular concentration of Rlc1-tdTomato in the *pkd2* mutant cells,

compared to the wild type (Fig. 2E). Therefore, Pkd2 also modulates the recruitment of Rlc1 to

the contractile ring during cytokinesis.

158   Lastly, we determined whether Pkd2 has a role in the assembly of actin filaments in the

cytokinetic contractile ring. We employed GFP-Lifeact as the fluorescence reporter (Malla et al.,

2022) to quantify the number of actin molecules in the ring of *pkd2-81KD* and the wild type cells

(Fig. 3A). Overall, the actin patches of mutant cells were far more numerous than those of the

wild type cells and these patches distributed in a less polarized fashion (Fig. 3A). Nevertheless,

the assembly and disassembly of actin filaments in the contractile ring of the mutant cells

proceeded normally (Fig. 3B). In fact, the number of actin molecules in the contractile ring of

pkd2-81KD cells increased by 12% from that of the wild type cells (Fig. 3C). Interestingly, the

intracellular concentration of GFP-Lifeact in the mutant cells increased dramatically by almost

90% (Fig. 3D). Again, we concluded that Pkd2 modulates the assembly of actin filaments in the

cytokinetic contractile ring.

***pkd2* and *myo2* play complementary roles in cytokinesis**

In an alternative approach, we examined the relationship between *pkd2* and *myo2,* both of which

are essential for cytokinesis through genetic studies. First, we confirmed the previously

discovered negative genetic interaction between *pkd2-81KD* and a temperature-sensitive mutant

myo2-E1 (Morris et al., 2019). This finding is consistent with the negative genetic interaction

between the temperature-sensitive mutant *pkd2-B42* (Sinha et al., 2022) and *myo2-E1* (Fig. 4A).

As a result of this genetic interaction, the double mutant *pkd2-B42 myo2-E1* grew much more

slowly than either *pkd2-B42* or *myo2-E1* (Fig. 4A). Next, we compared the morphogenesis of

pkd2-81KD, myo2-E1, and *pkd2-81KD myo2-E1* cells. Morphologically, the double *pkd2 myo2*

mutant cells were similar to the *pkd2-81KD* cells. Both were rounder than the wild type.

However, the *pkd2 myo2* double mutant stood out for its large number of lysed cells (11±1.7%)

even at the permissive temperature of 25°C, far more than either the *pkd2* (5±0.5%) or *myo2*

(0.6±0.8%) mutant cells (Fig. 4B). The septation index of the double mutant (74%) was also

much higher than those of either single mutant (Fig. 4C). The septum of *pkd2-81KD myo2-E1*

cells were abnormal in their appearance. They were often wavy, a rare phenotype among either

of the single mutants. Lastly, a much higher fraction of the double mutant cells was multi-

septated (21±3%, averaged ± standard deviation, n>500) than either *pkd2-81KD* (10±1%) or

myo2-E1 (1±0.1%). Based on these genetic analyses of the double *pkd2 myo2* mutant, we

concluded that *pkd2* and *myo2* play complementary roles in cytokinesis.

Pkd2 synergizes with Myo2 to regulate the assembly of the contractile ring

To determine how Pkd2 and Myo2 work synergistically in cytokinesis, we measured the

contractile ring assembly and constriction of the *pkd2-81KD myo2-E1* cells at the permissive

temperature through live fluorescence microscopy (Fig. 5A). These double mutant cells

assembled their contractile ring, marked by Rlc1-tdTomato, much more slowly than either *pkd2-*

81KD or *myo2-E1* cells (Fig. 5B). Strikingly, the contractile rings of most double mutant cells

(n=26) never appeared to be fully assembled. There were often many gaps in these contractile

rings (Fig. 5B). These rings nevertheless constricted at a rate comparable to those of either wild

type or *myo2-E1* cells at 25°C, but much more slowly than *pkd2-81KD* cells (Fig. 5C). We

concluded that both Pkd2 and Myo2 regulate the contractile ring assembly likely through

complementary pathways.

Lastly, we determined whether Pkd2 and Myo2 synergistically recruit the regulatory light

chain Rlc1 to the contractile ring. We measured the number of Rlc1-tdTomato molecules in the

contractile ring of the *pkd2-81KD myo2-E1* mutant cells through quantitative fluorescence

microscopy. Although Rlc1-Tdtomato continued to localize to the contractile ring of the *pkd2*

myo2 double mutant cells (Fig. 5D), its number decreased dramatically (-60%) from that of the

myo2-E1 cells (Fig. 5E). The intracellular concentration of Rlc1-tdTomato in the *pkd2 myo2*

double mutant cells decreased by a similar extent (-70%) (Fig. 5F). We concluded that both

Myo2 and Pkd2, likely through two complementary pathways, regulate the assembly of Rlc1 in

the contractile ring.

Discussion

In this study, we investigated whether the mechanosensitive calcium channel Pkd2 plays a role in

the assembly and constriction of the actomyosin contractile ring during cytokinesis. Through

quantitative fluorescence microscopy, we found surprisingly that Pkd2 mostly modulates the

assembly of the essential type II myosin Myo2, its regulatory light chain Rlc1 and actin filaments

in the contractile ring. Through analysis of the negative genetic interactions between *pkd2* and

myo2 mutants, we found that Pkd2 and the type II myosin Myo2 work synergistically to promote

the assembly of the cytokinetic contractile ring.

The modulating role of Pkd2 in the contractile ring assembly is consistent with the

cytokinesis defects of the *pkd2-81KD* mutant cells. Our earlier works has found that the cleavage

furrow ingressed almost 50% more quickly in the *pkd2* mutant cells compared to the wild type

(Morris et al., 2019). This phenotype is in line with our new finding that the contractile ring of

the mutant cells contains 20% more Myo2, Rlc1 and actin than the wild type ones. The increased

densities of Myosin II and actin filaments shall in theory enhance the mechanical force generated

by the contractile ring. Combining this increased compressing force by the contractile ring of the

pkd2 mutant with the reduced resistance from the outward turgor pressure (Sinha et al., 2022), it

is not surprising that the cleavage furrow ingresses so quickly in *pkd2-81KD* mutant cells.

225   The modulating effect of Pkd2 in the assembly of contractile ring most likely is rooted its

activity as a Ca^{2+}-permeable ion channel. In animal cells, Ca^{2+}-dependent myosin light chain

kinase (MLCK) phosphorylate Rlc in order to trigger the contractile ring constriction (Craig et

al., 1983). However, fission yeast does not possess a MLCK. Instead, its regulatory light chain

Rlc1 is phosphorylated by the kinase Pak1 during cytokinesis (Loo and Balasubramanian, 2008).

Although such phosphorylation of Rlc1 is not essential the temporal regulation of cytokinesis

(McCollum et al., 1999), it does promotes the assembly of actin filaments in the contractile ring

under stress condition (Prieto-Ruiz et al., 2023). Although Pak1 is not known as a

Ca^{2+}/calmodulin-dependent kinase, the contractile ring-localized phosphotase calcineurin is

(Martin-Garcia et al., 2018). We propose that calcineurin, a potential downstream target of Pkd2

channel, could dephosphorylate Rlc1 to modulate the assembly of the contractile ring.

236   It is surprising that mutations of *pkd2* significantly increase the intracellular

concentration of Rlc1 and actin. This could be due to the regulation of transcription by Pkd2. The

human polycystin PC1 does regulate gene transcription through its cleaved C-terminal

cytoplasmic tail (Chauvet et al., 2004). However, we have not observed nuclear localization of

Pkd2 so far. The alternative explanation could lie in the channel activity of Pkd2. Mutation of

this calcium permissive ion channel changes intracellular Ca^{2+} homeostasis (Poddar et al., 2022).

Ca^{2+}-sensitive phosphotase calcineurin. Calcineurin promotes the nuclear translocation and

activation of the transcription activator Prz1 (Hirayama et al., 2003). Therefore, Pkd2 may also

regulate the global gene transcription indirectly through the activation of calcineurin.

Considering the modulating effects of Pkd2 in the assembly of Myo2 at the contractile

ring, the negative genetic interaction between the *pkd2* and *myo2* mutants is surprising. Even

more surprising is our discovery that the abundance of Rlc1 in the contractile ring decreased

dramatically in the contractile ring of the *myo2 pkd2* double mutant cells. This is despite the fact

that *pkd2-81KD* mutation increases the number of Rlc1 in the contractile ring. A likely

explanation is that both Pkd2 and Myo2 promote the stability of the contractile ring. The Pkd2

channel may stabilize the contractile ring through modulating the assembly of actin and Myo2.

In contrast, Myo2 promotes the contractile ring stability through its motor activity, which is

comprised in the *myo2-E1* mutant (Balasubramanian et al., 1998). Our discovery suggests that

Pkd2 and Myo2 may play complementary roles in stabilizing the contractile ring during

cytokinesis.

Overall, our study demonstrated that the force-sensitive calcium channel Pkd2, although

not essential for the assembly of the contractile ring, plays a regulatory role in the assembly of

contractile ring through modulating the recruitment of actomyosin filaments to the ring. Future

work would determine whether Pkd2 exerts its regulatory function through its calcium

permeability and through it activation of the calcium-binding proteins such calmodulin and

calcineurin in cytokinesis.

Author contributions

Conceptualization, QC.; Methodology, QC, PC and DS; Software, QC; Validation, QC, PC, DS,

AP and MD; Formal Analysis, QC and PC; Investigation, QC, PC, DS, AP and MD; Resources,

QC, PC and DS; Data Curation, QC, PC and DS; Writing – Original Draft Preparation, QC and

PC; Writing – Review & Editing, QC and PC; Visualization, QC and PC; Supervision, QC;

Project Administration, QC; Funding Acquisition, QC.

Competing interests

The authors declare no competing interests.

Acknowledgements

We would like to thank the Chen lab members at The University of Toledo for technical

assistance. This work has been supported by the National Institutes of Health grant

R01GM144652 to QC and the National Science Foundation grant 2144701 to QC. The content is

solely the responsibility of the authors and does not necessarily represent the official views of the

National Institutes of Health. The authors declare no conflicts of interests.

Table 1. List of yeast strains used in this study.

Name	Genotype	Source
FY527	*h- leu1-32 ura4-D18 his3-D1 ade6-M216*	Lab stock
JW766	*h+ kanMX6-Pmyo2-GFP-myo2 ade6-M210 leu1-32 ura4-D18*	Wu Lab
JW1341	*h- rlc1-tdTomato-natMX6 ade6-M210 leu1-32 ura4-D18*	Wu Lab
QC-Y601	*h+ leu2::kanMX-Pcof1-mEGFP-Lifeact rlc1-tdTomato-Nat*	Lab stock
QC-Y799	*h+ myo2-E1 rlc1-tdTomato-NatMX6 sad1-mGFP-KanMX6*	This study
QC-Y813	*h- kanMX6-81xnmt1-pkd2 rlc1-tdTomato-NatMX6*	Lab stock
QC-Y817	*h+ kanMX6-81xnmt1-pkd2 leu1-32 ura4-D18 his3-D1 ade6-M210*	Lab stock
QC-Y839	*h- myo2-E1*	
QC-Y840	*h+ myo2-E1 kanMX6-81xnmt1-Pkd2*	This study
QC-Y1033	*h? myo2-E1 kanMX6-81xnmt1-pkd2 rlc1-tdTomato-natMX6*	This study
QC-Y1112	*h? kanMX6-Pmyo2-GFP-myo2 ade6-M210 leu1-32 ura4-D18 kanMX6-81xnmt1-pkd2*	This study
QC-Y1156	*h? leu2::kanMX-Pcof1-mEGFP-Lifeact rlc1-tdTomato-Nat KanMX6-P81nmt1-pkd2 leu1-32 ura4-D18 his3-D1*	Lab stock
QC-Y1752	*h? pkd2::pkd2-B42-ura4+-hist5+ leu1-32 myo2-E1*	This study

Figure and Figure Legends

Figure 1: Localization of Myo2 in the contractile ring increased in *pkd2-81KD* cells. (A)
Micrographs of wild type and *pkd2-81KD* cells expressing GFP-Myo2. Arrow: broad band of
nodes. Arrowhead: the contractile ring. **(B)** Time lapse series of the equatorial plane of a wild
type and a *pkd2-81KD* cells expressing GFP-Myo2. Top: a diagram of the contractile ring
assembly, maturation and constriction in fission yeast cells. Asterisk: the start of contractile ring
constriction. Number indicates time in minutes. Scale bar=5μm. **(C)** Time course of the
fluorescence intensity of GFP-Myo2 at the equatorial plane after the start of contractile ring
assembly (time zero). Cloud: standard deviations. **(D-E)** The dot plots of the average
fluorescence intensity of GFP-Myo2 in either the contractile ring (D) or the whole cell (E). *:
P<0.05. **: P<0.01. ***: P<0.001. N.S.: not significant. Statistics were calculated using two-
tailed Student's t tests. All the data are pooled from at least two independent biological repeats.
Scale bar=5μm.

Figure 2

Figure 2: Localization of myosin light chain Rlc1 increased in *pkd2-81KD* mutant cells. (A)
Micrographs of *wild type* and *pkd2-81KD* expressing Rlc1-tdTomato. Arrowhead: the contractile
ring. **(B)** Time-lapse series of the equatorial plane of a *wild type* and a *pkd2-81KD* cell
expressing Rlc1-tdTomato. Dashed box: the start of contractile ring constriction. Number
indicates time in minutes. Scale bar=5µm. **(C)** Time course of the fluorescence intensity of Rlc1-
tdTomato in the equatorial plane. Cloud: standard deviations. **(D-E)** Dot plots of the average
fluorescence intensity of Rlc1-tdTomato in the mature contractile ring (D) or the whole cell (E).
*: P<0.05. **: P<0.01. ***: P<0.001. N.S.: not significant. Statistics were calculated using two-
tailed Student's t tests. All the data are pooled from at least two independent biological repeats.

Figure 3

Figure 3: The number of actin molecules decreased in the contractile ring of *pkd2-B42*
mutant cells. (A) Micrographs of the *wild type* and *pkd2-81KD* co-expressing GFP-Lifeact
(green) and Rlc1-tdTomato (magenta). Arrow: GFP-Lifeact in division plane.Arrow-head: Rlc1-
tdTomato in contractile ring. **(B)** Time series micrographs of a *wild type* and a *pkd2-81KD* cell.
Number: time in minutes. Interval=2 mins. Scale bar=5µm. **(C-D)** Dot plots of the average
fluorescence intensity of GFP-Lifeact in either the contractile ring (C) or the whole cell (D). *:
P<0.05. **: P<0.01. ***: P<0.001. N.S.: not significant. Statistics were calculated using two-
tailed Student's t tests. All the data are pooled from at least two independent biological repeats.

Figure 4

314
Figure 4: Genetic interactions among *pkd2* and *myo2* mutants. (A) Ten-fold dilution series of
yeast cells at the indicated temperature after two days. **(B)** Bright-field micrographs of *pkd2-*
81KD, myo2-E1, and pkd2-81KD myo2-E1 cells at 25°C. Arrow: lysed cells. Number: average
percentage of lysed cells ± standard deviations (n>200). **(C)** Micrographs of calcofluor stained
cells at 25°C. Arrow: abnormal septum. Arrowhead: curved septum (n>500). All the data are
pooled from at least two independent biological repeats. Scale bar=5µm.
321

Figure 5

Figure 5: Localization of myosin regulatory light chain Rlc1 in the contractile ring
decreased in the *pkd2-81KD myo2-E1* mutant cells. (A) Micrographs of *myo2-E1* and *pkd2-*
81KD myo2-E1 expressing Rlc1-tdTomato at 25°C. Arrow-heads: Rlc1-tdTomato in contractile
ring. **(B)** Time-lapse series of the equatorial plane of a *myo2-E1* and a *pkd2-81KD myo2-E1* cell
expressing Rlc1-tdTomato. Number indicates time in minutes. Scale bar=5μm. **(C)** Dot plots of
the constriction rate of contractile ring at 25°C. **(D)** Time course of the fluorescence intensity of
Rlc1-tdTomato at the equatorial plane. **(E-F)** Dot plots of average fluorescence intensity of Rlc1-
tdTomato in either the contractile ring (E) or the whole cell (F). *: P<0.05. **: P<0.01. ***:
P<0.001. N.S.: not significant. Statistics were calculated using two-tailed Student's t tests. All
the data are pooled from at least two independent biological repeats.

334   **Reference**

335   Morris, Z., D. Sinha, A. Poddar, B. Morris, and Q. Chen. 2019. Fission yeast TRP channel Pkd2p
336   localizes to the cleavage furrow and regulates cell separation during cytokinesis.
337   *Molecular biology of the cell*. 30:1791-1804.

338   Courtemanche, N., T.D. Pollard, and Q. Chen. 2016. Avoiding artefacts when counting
339   polymerized actin in live cells with LifeAct fused to fluorescent proteins. *Nature cell*
340   *biology*. 18:676-683.
341   Malla, M., T.D. Pollard, and Q. Chen. 2022. Counting actin in contractile rings reveals novel
342   contributions of cofilin and type II myosins to fission yeast cytokinesis. *Molecular*
343   *biology of the cell*. 33:ar51.

344   1994. The polycystic kidney disease 1 gene encodes a 14 kb transcript and lies within a
345   duplicated region on chromosome 16. The European Polycystic Kidney Disease
346   Consortium. *Cell*. 77:881-894.
347   AbouAlaiwi, W.A., S. Ratnam, R.L. Booth, J.V. Shah, and S.M. Nauli. 2011. Endothelial cells
348   from humans and mice with polycystic kidney disease are characterized by polyploidy
349   and chromosome segregation defects through survivin down-regulation. *Human*
350   *molecular genetics*. 20:354-367.
351   Balasubramanian, M.K., D. McCollum, L. Chang, K.C. Wong, N.I. Naqvi, X. He, S. Sazer, and
352   K.L. Gould. 1998. Isolation and characterization of new fission yeast cytokinesis
353   mutants. *Genetics*. 149:1265-1275.
354   Chauvet, V., X. Tian, H. Husson, D.H. Grimm, T. Wang, T. Hiesberger, P. Igarashi, A.M.
355   Bennett, O. Ibraghimov-Beskrovnaya, S. Somlo, and M.J. Caplan. 2004. Mechanical
356   stimuli induce cleavage and nuclear translocation of the polycystin-1 C terminus. *The*
357   *Journal of clinical investigation*. 114:1433-1443.
358   Chen, Q., and T.D. Pollard. 2011. Actin filament severing by cofilin is more important for
359   assembly than constriction of the cytokinetic contractile ring. *The Journal of cell biology*.
360   195:485-498.
361   Craig, R., R. Smith, and J. Kendrick-Jones. 1983. Light-chain phosphorylation controls the
362   conformation of vertebrate non-muscle and smooth muscle myosin molecules. *Nature*.
363   302:436-439.
364   Hardy, E., and L. Tsiokas. 2020. Polycystins as components of large multiprotein complexes of
365   polycystin interactors. *Cellular signalling*. 72:109640.
366   Hirayama, S., R. Sugiura, Y. Lu, T. Maeda, K. Kawagishi, M. Yokoyama, H. Tohda, Y. Giga-
367   Hama, H. Shuntoh, and T. Kuno. 2003. Zinc finger protein Prz1 regulates Ca2+ but not
368   Cl- homeostasis in fission yeast. Identification of distinct branches of calcineurin
369   signaling pathway in fission yeast. *The Journal of biological chemistry*. 278:18078-
370   18084.
371   Hu, J., and P.C. Harris. 2020. Regulation of polycystin expression, maturation and trafficking.
372   *Cellular signalling*. 72:109630.
373   Kovar, D.R., V. Sirotkin, and M. Lord. 2011. Three's company: the fission yeast actin
374   cytoskeleton. *Trends in cell biology*. 21:177-187.
375   Laplante, C., J. Berro, E. Karatekin, A. Hernandez-Leyva, R. Lee, and T.D. Pollard. 2015. Three
376   myosins contribute uniquely to the assembly and constriction of the fission yeast
377   cytokinetic contractile ring. *Curr Biol*. 25:1955-1965.

Loo, T.H., and M. Balasubramanian. 2008. Schizosaccharomyces pombe Pak-related protein,
Pak1p/Orb2p, phosphorylates myosin regulatory light chain to inhibit cytokinesis. *The*
Journal of cell biology. 183:785-793.
Malla, M., T.D. Pollard, and Q. Chen. 2022. Counting actin in contractile rings reveals novel
contributions of cofilin and type II myosins to fission yeast cytokinesis. *Molecular*
biology of the cell. 33:ar51.
Malla, M., D. Sinha, P. Chowdhury, B.T. Bisesi, and Q. Chen. 2023. The cytoplasmic tail of the
mechanosensitive channel Pkd2 regulates its internalization and clustering in eisosomes.
Journal of cell science. 136.
Martin-Garcia, R., V. Arribas, P.M. Coll, M. Pinar, R.A. Viana, S.A. Rincon, J. Correa-Bordes,
J.C. Ribas, and P. Perez. 2018. Paxillin-Mediated Recruitment of Calcineurin to the
Contractile Ring Is Required for the Correct Progression of Cytokinesis in Fission Yeast.
Cell reports. 25:772-783 e774.
McCollum, D., A. Feoktistova, and K.L. Gould. 1999. Phosphorylation of the myosin-II light
chain does not regulate the timing of cytokinesis in fission yeast. *The Journal of*
biological chemistry. 274:17691-17695.
Mochizuki, T., G. Wu, T. Hayashi, S.L. Xenophontos, B. Veldhuisen, J.J. Saris, D.M. Reynolds,
Y. Cai, P.A. Gabow, A. Pierides, W.J. Kimberling, M.H. Breuning, C.C. Deltas, D.J.
Peters, and S. Somlo. 1996. PKD2, a gene for polycystic kidney disease that encodes an
integral membrane protein. *Science (New York, N.Y)*. 272:1339-1342.
Moreno, S., A. Klar, and P. Nurse. 1991. Molecular genetic analysis of fission yeast
Schizosaccharomyces pombe. *Methods in enzymology*. 194:795-823.
Morris, Z., D. Sinha, A. Poddar, B. Morris, and Q. Chen. 2019. Fission yeast TRP channel Pkd2p
localizes to the cleavage furrow and regulates cell separation during cytokinesis.
Molecular biology of the cell. 30:1791-1804.
Naqvi, N.I., K.C. Wong, X. Tang, and M.K. Balasubramanian. 2000. Type II myosin regulatory
light chain relieves auto-inhibition of myosin-heavy-chain function. *Nature cell biology*.
2:855-858.
Palmer, C.P., E. Aydar, and M.B. Djamgoz. 2005. A microbial TRP-like polycystic-kidney-
disease-related ion channel gene. *The Biochemical journal*. 387:211-219.
Poddar, A., Y.Y. Hsu, F. Zhang, A. Shamma, Z. Kreais, C. Muller, M. Malla, A. Ray, A. Liu,
and Q. Chen. 2022. Membrane stretching activates calcium-permeability of a putative
channel Pkd2 during fission yeast cytokinesis. *Molecular biology of the*
cell:mbcE22070248.
Poddar, A., O. Sidibe, A. Ray, and Q. Chen. 2021. Calcium spikes accompany cleavage furrow
ingression and cell separation during fission yeast cytokinesis. *Molecular biology of the*
cell. 32:15-27.
Pollard, T.D., and J.Q. Wu. 2010. Understanding cytokinesis: lessons from fission yeast. *Nature*
reviews. 11:149-155.
Prieto-Ruiz, F., E. Gomez-Gil, R. Martin-Garcia, A.J. Perez-Diaz, J. Vicente-Soler, A. Franco,
T. Soto, P. Perez, M. Madrid, and J. Cansado. 2023. Myosin II regulatory light chain
phosphorylation and formin availability modulate cytokinesis upon changes in
carbohydrate metabolism. *eLife*. 12.
Shen, P.S., X. Yang, P.G. DeCaen, X. Liu, D. Bulkley, D.E. Clapham, and E. Cao. 2016. The
Structure of the Polycystic Kidney Disease Channel PKD2 in Lipid Nanodiscs. *Cell*.
167:763-773 e711.

Sinha, D., D. Ivan, E. Gibbs, M. Chetluru, J. Goss, and Q. Chen. 2022. Fission yeast polycystin
Pkd2p promotes cell size expansion and antagonizes the Hippo-related SIN pathway.
Journal of cell science. 135.
Su, Q., F. Hu, X. Ge, J. Lei, S. Yu, T. Wang, Q. Zhou, C. Mei, and Y. Shi. 2018. Structure of the
human PKD1-PKD2 complex. *Science (New York, N.Y)*. 361.
Ta, C.M., T.N. Vien, L.C.T. Ng, and P.G. DeCaen. 2020. Structure and function of polycystin
channels in primary cilia. *Cellular signalling*. 72:109626.
Thevenaz, P., U.E. Ruttimann, and M. Unser. 1998. A pyramid approach to subpixel registration
based on intensity. *IEEE Trans Image Process*. 7:27-41.
Wu, J.Q., and T.D. Pollard. 2005. Counting cytokinesis proteins globally and locally in fission
yeast. *Science (New York, N.Y)*. 310:310-314.

Figure 5

Figure 4

A

	25°C	30°C	36°C
WT			
myo2-E1			
pkd2-81KD			
pkd2-81KD myo2-E1			
pkd2-B42			
pkd2-B42 myo2-E1			

B

WT	pkd2-81KD	myo2-E1	pkd2-81KD myo2-E1
25°C			
0.4±0.2%	5.3±0.5%	0.6±0.8%	11.4±1.7%(***)

C

Calcoflour

| 13±0.2% | 49±0.1% | 17±0.9% | 74±1.0%(***) |

Figure 3

Figure 2

Figure 1

A

WT   *pkd2-81KD*

GFP-Myo2

B

Assembly → Maturation → Constriction

WT

pkd2-81KD

GFP-Myo2

Interval=2'

C

GFP-Myo2 at the division plane

pkd2-81KD (n=20)

WT (n=30)

Fluorescence intensity (A.U.)

Time (min)

D

GFP-Myo2 in the contractile ring

Mean ± 1 SD

**

n=30   n=20

WT   *pkd2-81KD*

E

GFP-Myo2 in whole cell

NS

n=200   n=190

WT   *pkd2-81KD*