Adaptive thermogenesis is mediated by GDF15 via the GFRAL neuronal axis in mice

Ji Eun Kim[#1], Sang-Hyeon Ju[#1,2], Minhee Lee[3], Hyun Jung Hong[1,4], Uzma Yaseen[1], Benyuan Zhang[1], Seong Eun Lee[1], Yea Eun Kang[1,2], Jung Tae Kim[1,4], Hyon-Seung Yi[1,2], Yoon-Sun Yi[5], Sangmi Jun[5], Minsung Park[4], Jinkuk Kim[4], Johan Auwerx[6], Jong-Woo Sohn[4], Ju Hee Lee*[1,2], Minho Shong*[1,4]

[1]Research Center for Endocrine and Metabolic Diseases, School of Medicine, Chungnam National University, Daejeon 35015, Republic of Korea

[2]Department of Internal Medicine, Chungnam National University Hospital, Daejeon, 35015, Republic of Korea

[3]THOR Therapeutics, Inc., 99 Chungnam National University, Daehak-ro, Yuseong-gu, Daejeon 34134, Republic of Korea

[4]Graduate School of Medical Science and Education, Korea Advanced Institute of Science and Technology, 291 Daehak-ro, Yuseong-gu, Daejeon 34141, Republic of Korea

[5]Center for Research Equipment, Korea Basic Science Institute, Cheongju 28119, Republic of Korea

[6]Laboratory for Integrative Systems Physiology, Institute of Bioengineering, École Polytechnique Fédérale de Lausanne, Lausanne 1015, Switzerland

[#]These authors contributed equally to this study.

*Corresponding Authors:
Ju Hee Lee
Chungnam National University,
Munwharo 266, Jung-gu, Daejeon 35015, Republic of Korea
82-42-280-6801
serenaj@cnu.ac.kr

Minho Shong
Korea Advanced Institute of Science and Technology,
291 Daehak-ro, Yuseong-gu, Daejeon 34141, Republic of Korea
minhos@kaist.ac.kr

Author Contributions: Conceptualization, J.H.L., J.W.S., and M.S.; Methodology, J.H.L., J.E.K., S.H.J., M.H.L., H.J.H., Z.B., Y.E.K., J.T.K., H.S.Y., S.J., and J.K.; Formal Analysis, J.H.L., J.E.K., S.H.J., M.H.L., and S.E.L.; Investigation, J.E.K., S.H.J., M.H.L., U.Y., Y.S.Y. and M.P.; Writing – Original Draft, J.H.L., J.E.K., and S.H.J.; Writing – Review & Editing, J.H.L., J.A., J.W.S., and M.S.; Supervision, M.S.; Funding Acquisition, J.H.L. and M.S. All the authors read and edited the manuscript, and approved the final version.

Competing Interest Statement: The authors declare no competing financial or other interests in this study.

43 **Keywords:** Brown adipose tissue, mitochondrial unfolded protein response, growth differentiation
44 factor 15, glial cell-derived neurotrophic factor receptor alpha-like neuron, sympathetic nervous
45 system
46
47 **This file includes:**
48   Main Text
49   Figures 1 to 7
50
Abstract
Adaptive thermogenesis is a key homeostatic mechanism that primarily occurs in brown adipocytes
and enables the maintenance of body temperature. Although this process involves coordinated
responses in multiple tissues, including the browning of white adipocytes, the precise inter-organ
crosstalk underlying adaptive thermogenesis is unclear. Here, we investigate the pivotal role of the
GDNF family receptor alpha-like (GFRAL) neuronal axis in modulating compensatory thermogenic
responses in brown and white adipose depots under stress conditions, specifically the mitochondrial
unfolded protein response resulting from genetic modification and cold exposure. We employed a
mouse model with targeted deletion of *Crif1* in the mitoribosomes of brown adipocytes, and cold-
exposed mice and immortalized adipocytes, to uncover the mechanism by which mitochondrial stress-
induced growth differentiation factor 15 (GDF15) expression affects metabolism and facilitates
adaptive thermogenesis. We found that *Crif1* deletion resulted in browning of inguinal white adipose
depots, increased energy expenditure, reduced food intake, and resistance to weight gain.
Retrograde neuronal tracing established that GFRAL-positive neurons in the hindbrain and
sympathetic preganglionic neurons in the spinal cord mediated the GDF15-associated browning of
inguinal white adipose tissue. Intervention studies using antisense oligonucleotides to inhibit *Gfral*
expression blunted the effect of *Crif1* deletion on energy expenditure and food intake, further
confirming the essential role the GFRAL axis plays in GDF15-driven thermogenic adaptation in white
adipose tissue. Our findings suggest that the GFRAL neuronal axis is key in coordinating the adaptive
thermogenic response across multiple tissues and adipose depots, thereby ensuring metabolic
homeostasis during mitochondrial stress.
72
73

Introduction

Adaptive thermogenesis in adipose tissue is a crucial homeostatic mechanism that helps protect animals against cold environments. While brown adipose tissue (BAT) is the primary site of thermogenesis, beige adipocytes within white adipose tissue (WAT) also display a thermogenic response to cold-induced hypothermia. The coordinated actions of brown and beige adipocytes, facilitated by activation of the sympathetic nervous system (SNS), are essential for the regulation of body temperature in mammals[1]. The SNS not only triggers lipolysis in white adipocytes, supplying substrates for brown adipocyte thermogenesis, but also appears to orchestrate the temporary emergence of beige adipocytes, indicating a complex regulatory hierarchy underlying responses to cold in adipose tissue[2].

In circumstances where BAT is absent or functionally compromised, WAT has the capacity to undergo compensatory browning, a process driven by sympathetic nerve activity and proteins secreted from BAT, to preserve energy homeostasis[3-6]. One way BAT function may be impaired is through disruption to mitochondrial function, such as that seen in brown adipocyte-specific Lkb1 or mitochondrial transcription factor A knockout rodent models[7]. In these examples, there is a paradoxical increase in metabolic efficiency with mitochondrial dysfunction, even under high-fat diet conditions and irrespective of environmental temperature. This phenomenon implies a decoupling of the thermogenic activity of BAT from energy expenditure, which may be related to targeted suppression of mitochondrial DNA (mtDNA) gene expression and resulting electron transport chain (ETC) imbalances[7]. Such imbalances may elicit a mitochondrial unfolded protein response (UPR[mt]) in BAT, which subsequently influences systemic metabolism independent of thermogenesis by promoting the appearance of beige adipocytes to compensate for BAT dysfunction and maintain metabolic stability. However, the precise interplay between BAT and WAT depots in response to the UPR[mt] in BAT is unclear.

The mammalian UPR[mt] involves a cascade of transcription factors, such as activating transcription factor 4 (ATF4), which are involved in optimizing thermoregulatory functions in BAT[8]. In particular, cold exposure upregulates ATF protein levels, emphasizing their integral role in maintaining thermal homeostasis. These transcription factors, including ATF4, ATF5, and C/EBP homologous protein (CHOP), promote mitochondrial biogenesis and expression of genes involved in protein folding quality control, and also mediate the release of mitokines like growth differentiation factor 15 (GDF15) and fibroblast growth factor 21 (FGF21), in response to the UPR[mt9-11]. However, the role of these mitokines in BAT where the UPR[mt] is activated is not fully understood. The glial-derived neurotrophic factor (GDNF)-family receptor alpha-like (GFRAL) has been identified by four research groups as the receptor for GDF15, which signals through the coreceptor RET, primarily located in the hindbrain[12-15].

Our study reveals the pivotal role of GFRAL neurons in this context: we demonstrate that GDF15, secreted by brown adipocytes undergoing UPR[mt], initiates beige adipogenesis through the activation of GFRAL neurons. This discovery underscores previously unrecognized BAT–brain crosstalk via the GDF15–GFRAL axis, which is activated by either physiological stress from cold

113 exposure or pathological stress from mitochondrial dysfunction, and, in turn, orchestrates
114 compensatory adaptive thermogenesis in beige adipocytes. Thus, GFRAL neurons emerge as master
115 regulators in the thermogenic network, underpinning the relationship between BAT function and
116 whole-body energy homeostasis.
117
118 **Results**
119
Induction of the UPR^{mt} and the integrated stress response in the BAT of Crif1 (CR6-interacting
factor 1)-deficient mice
To investigate the effects of abnormalities in brown adipocyte-specific mitochondrial
proteostasis, we generated a mouse model of mitochondrial dysfunction induced by the genetic
inactivation of the *Crif1* gene in brown adipocytes. CRIF1 is critical to the function of the mitoribosome
and the mitochondrial translation of mtDNA-encoded oxidative phosphorylation (OXPHOS)
polypeptides. Inactivation of the *Crif1* gene results in significant reductions in the production of
OXPHOS components and ETC activity[16]. BAT-specific *Crif1*-knockout mice (Crif1[BKO], generated
using *Ucp1-Cre* and *Crif1[f/f]* mice) had significantly lower *Crif1* mRNA and protein expression in BAT
than control mice (Fig. 1a and Extended Data Fig. 1a–c). In addition, Crif1[BKO] mice had significantly
lower expression of the OXPHOS complexes I (NDUFB8, nuclear DNA-encoded), III (UQCRC2,
nuclear [n]DNA-encoded), and IV (MTCO1, mtDNA-encoded) than control mice (Fig. 1a and
Extended Data Fig. 1C). Electron microscopy of brown adipocytes from Crif1[BKO] mice revealed
swollen mitochondria with abnormal cristae (Fig. 1b). Thus, the ablation of the *Crif1* gene in BAT
results in substantial suppression of OXPHOS complex expression and mitochondrial structural
abnormalities. The BAT of Crif1[BKO] mice had larger lipid droplets and reduced multilocularity,
indicative of potential defects in fatty acid oxidation and/or triglyceride storage (Fig. 1c and Extended
Data Fig. 1d). UCP1 protein levels were either unchanged (males) or significantly increased (females)
(Fig. 1c and Extended Data Fig. 1c).
Abnormal expression of OXPHOS components in *Crif1*-deficient cells induces transcriptomic
changes, including those relevant to the UPR^{mt}, which consists of both cell-autonomous and non-cell-
autonomous responses[11]. The ATF family, including ATF4 and ATF5, and CHOP are implicated in the
mammalian UPR^{mt}[17,18]. In the present study, we evaluated the induction of the integrated stress
response (ISR) and the UPR^{mt} in the BAT of Crif1[BKO] mice and found that the expression of ATF5, but
not ATF4 or CHOP, was significantly higher in the BAT of *Crif1*[BKO] mice than control mice, suggesting
a role for brown adipocytes in the maintenance of UPR^{mt} activity during mitochondrial stress (Fig. 1d
and Extended Data Fig. 1e). In addition, mitochondrial chaperones and intrinsic proteases, such as
heat shock protein family D member 1 (HSPD1), DnaJ heat shock protein family member A3
(DNAJA3), and lon peptidase 1 (LONP1), were expressed at significantly higher levels in the BAT of
Crif1[BKO] mice (Fig. 1e and Extended Data Fig. 1f). However, the expression of another mitochondrial
intrinsic protease, caseinolytic mitochondrial matrix peptidase proteolytic subunit (CLPP), was

significantly lower than in control mice in both male and female Crif1[BKO] mice (Fig. 1e and Extended
Data Fig. 1f).

These results demonstrate the consequences of mitoribosomal dysfunction in brown
adipocytes: less biogenesis of OXPHOS components, structural abnormalities of the mitochondria,
and an induction of both the ISR and UPR[mt], as evidenced by elevated expression of ATF5,
mitochondrial chaperones such as DNAJA3, and intrinsic proteases including LONP1.

158 Metabolic characteristics of thermogenic compensation by Crif1[BKO] mice subjected to cold
159 exposure or a high-fat diet (HFD)

Crif1[BKO] mice housed at room temperature (RT) underwent assessments for body weight, daily
food intake, and the weight of various organs including adipose tissue, liver, and skeletal muscle.
Both male and female Crif1[BKO] mice exhibited significantly lower body weight and daily food intake
compared with control mice. Both of the epididimal and inguinal WAT mass in Crif1[BKO] mice were
lower than those in control mice (Figs. 2a–c and Extended Data Figs. 2a–c). An analysis of
covariance showed that Crif1[BKO] mice exhibited significantly higher energy expenditure (kcal/day)
independent of body weight during paired feeding (Figs. 2d,e).

To evaluate the ability of Crif1[BKO] mice to maintain their body temperature in a cold
environment, we measured body temperature after housing them at 5°C for 1 week. The body
temperatures of Crif1[BKO] mice were significantly lower than those of the control mice during the first 4
h of cold exposure ($P<0.05$), but they subsequently maintained body temperatures comparable with
the control mice and showed no mortality (Fig. 2f). There were no differences in the glucose or insulin
tolerance test results of control and Crif1[BKO] mice (Extended Data Figs. 2d–g). These observations
indicate that despite impaired BAT OXPHOS, Crif1[BKO] mice have higher energy expenditure than
control mice, although glucose metabolism was not affected.

Next, to evaluate the contribution of inguinal WAT to thermogenic compensation and systemic
metabolic homeostasis, we measured the expression of UCP1 in this thermogenic fat depot of
Crif1[BKO] mice and found significantly higher expression of UCP1 and peroxisome proliferator-activated
receptor gamma co-activator 1-alpha (PPARGC1A) than in control mice (Fig. 2g and Extended Data
Fig. 2h). Histological examination revealed multilocular adipocytes and more UCP1 immunostaining in
Crif1[BKO] mice than in control mice (Fig. 2h and Extended Data Fig. 2i). Transmission electron
microscopy revealed more abundant and larger mitochondria in the inguinal WAT of Crif1[BKO] than in
control mice (Fig. 2i). These findings suggest that WAT browning may contribute to the maintenance
of systemic energy homeostasis despite BAT OXPHOS defects in Crif1[BKO] mice.

To further characterize the differences in the systemic metabolism of Crif1[BKO] mice, we
evaluated metabolic parameters after HFD feeding for 12 weeks. Crif1[BKO] mice fed an HFD (Crif1[BKO]-
HFD) had significantly lower expression of CRIF1; the OXPHOS complexes I, III, and IV; and UCP1
(Extended Data Figs. 3a–c). Crif1[BKO] mice fed an HFD also had significantly higher expression of
DNAJA3 and LONP1 in BAT, although CLPP was expressed at a significantly lower level (Extended
Data Fig. 3a). The body weight of Crif1[BKO]-HFD mice was significantly lower than that of control mice

(Fig. 2j and Extended Data Fig. 3c), and the WAT mass and fat percentages, determined using dual-
energy X-ray absorptiometry, were significantly lower in Crif1[BKO] mice, at only one-third or one-half of
those of the control group (Figs. 2k,l and Extended Data Figs. 3d,e). The daily food consumption of
male Crif1[BKO]-HFD mice was lower than that of the controls (Fig. 2m and Extended Data Fig. 3f).
Glucose and insulin tolerance testing showed no differences between the control and Crif1[BKO]-HFD
mice (Extended Data Figs. 3g–j). The inguinal WAT of Crif1[BKO]-HFD mice had significantly higher
expression of UCP1 and PPARGC1A, a greater number of multilocular adipocytes, and greater UCP1
immunostaining (Figs. 2n,o and Extended Data Figs. 3k,l), consistent with the results obtained in
normal chow-fed Crif1[BKO] mice (Figs. 2g–h and Extended Data Figs. 2h,i). In summary, Crif1[BKO] mice
exhibited resistance to weight gain when fed an HFD. In addition, Crif1[BKO] mice consumed less food
and had more beige adipocytes in the inguinal WAT depot than control mice.
Taken together, these findings suggest that Crif1[BKO] mice maintain a lower body weight and
higher energy expenditure than control mice, despite impaired BAT OXPHOS. Moreover, when fed an
HFD, Crif1[BKO] mice resist weight gain, effects associated with increased browning of the inguinal WAT
depot.
205

206   Identification and mechanism of GDF15 expression by the UPR[mt] in the BAT of Crif1[BKO] mice

To confirm the non-cell-autonomous effect of UPR[mt] activation in BAT, we aimed to identify
proteins secreted from BAT using both the UniProt database (www.uniprot.org) and RNA-seq data.
Bulk RNA-seq of Crif1[BKO] BAT showed 54 genes encoding secretory proteins that were significantly
differentially expressed, with log$_2$-fold differences >2 and adjusted P-values <0.01 (Fig. 3a). Of these,
the expression of *Gdf15* had one of the highest fold changes and lowest P-values. This is consistent
with previous findings that GDF15 is a mitokine, which in mice and humans is secreted during
mitochondrial dysfunction[9-11,19,20]. The difference in BAT gene expression was accompanied by
significantly higher plasma concentrations of GDF15 in Crif1[BKO] mice fed either normal chow or an
HFD than control mice (Figs. 3b–e and Extended Data Figs. 4a–d). By contrast, the expression of
Fgf21 and serum levels of FGF21 in Crif1[BKO] mice were significantly higher than control mice only with
a normal chow diet, with no significant changes observed under HFD feeding (Extended Data Figs.
4e–l). After both a normal chow diet and HFD feeding, *Gdf15* and *Fgf21* mRNA levels in the liver were
not significantly different between control and Crif1[BKO] mice, except for *Fgf21* expression under
normal chow feeding in females (Extended Data Figs. 4m–p).
To further characterize the relationship between mitochondrial OXPHOS dysfunction, UPR[mt]
activation, and GDF15 secretion, we treated differentiated immortalized brown adipocytes with
OXPHOS complex inhibitors and measured the expression of genes encoding proteins involved in the
ISR and UPR[mt] pathways. The expression of the mitochondrial chaperone *Dnaja3* and the intrinsic
protease *Lonp1* was significantly higher after treatment with 10 μg/mL oligomycin (a complex V
inhibitor) and 2 μg/mL carbonyl cyanide m-chlorphenyl hydrazine (CCCP, an uncoupler), and
expression of the transcription factors *Atf4*, *Atf5*, and *Chop* was significantly higher after treatment
with oligomycin, CCCP, and 1 μM rotenone (a complex I inhibitor) treatment than after control

treatment (DMSO) (Figs. 3f,g). In addition, brown adipocytes treated with OXPHOS complex inhibitors
had significantly higher *Gdf15* mRNA expression and a 2–3-fold higher concentration of GDF15 in the
medium, with similar responses seen in the mitokine FGF21 (Figs. 3h,i and Extended Data Figs. 4q,r).
To investigate the connection between UPR[mt] and the induction of mitokine expression in brown
adipocytes, we induced cellular proteostasis stress in the mitochondrial matrix using gamitrinib-
triphenylphosphonium (GTPP), an Hsp90 inhibitor[21]. As expected, GTPP treatment resulted in
significantly higher mRNA expression of *Hspd1*, *Clpp*, and *Lonp1* than after the control treatment (Fig.
3j). The expression of *Chop* and *Gdf15* and the concentration of GDF15 in the medium was also
significantly higher after 2 µM GTPP treatment than after control treatment (Figs. 3k–m). In addition,
the concentration of FGF21 in the medium was significantly higher after 2 µM GTPP treatment than
after control treatment, in line with the tendency toward higher expression observed at the mRNA
level (Extended Data Figs. 4s,t).

Overall, these results indicate that GDF15 production increases in BAT in response to
OXPHOS dysfunction and abnormalities in proteostasis, induced by both pharmacological
intervention and genetic manipulation, *in vitro* and *in vivo*. The higher resulting concentrations of
GDF15 in Crif1[BKO] mice likely affect systemic metabolism and adaptive thermogenesis.

Induction of the UPR[mt] and GDF15 expression in the BAT of mice during cold exposure

The above results demonstrate that in Crif1[BKO] mice the UPR[mt] is induced, proteostasis is
dysfunctional, and GDF15 expression is increased, which is compensated for by adaptive
thermogenesis in inguinal WAT. To determine the effects of environmental, rather than genetic, BAT
manipulation, we assessed UPR[mt] activation and GDF15 expression in mice exposed to cold. Cold
exposure increases mitochondrial demand in thermogenic adipose tissue and triggers induction of the
mitochondrial stress response[22-24]. We assessed the induction of the UPR[mt] in the BAT of C57BL/6
wild-type (WT) mice in response to cold exposure (7°C) for 24 h. After 6 h of exposure, the expression
of *Ucp1* mRNA in BAT was 26-fold higher than at baseline (Fig. 4a). We measured the expression of
genes involved in the UPR[mt] and found that expression of the mitochondrial chaperones *Hspd1* and
Dnaja3, the mitochondrial matrix proteases *Clpp* and *Lonp1*, and the inner-membrane protease
Yme1l was 2–5-fold higher after 6 and 24 h of cold exposure than at baseline (Extended Data Figs.
4b–e), but expression of the inner-membrane protease *Imm2l* and the outer-membrane protease
Usp30 was not affected. The mRNA expression of *Atf4* and *Atf5*, transcription factors known to
regulate the UPR[mt], was approximately 20-fold higher after 6 and 24 h of cold exposure than at
baseline (Fig. 4f), and the protein expression of ATF4 and CHOP, but not that of ATF5, was also
significantly higher after cold exposure (Fig. 4g). Thus, the expression of mitochondrial chaperones
and proteases involved in the UPR[mt] was increased in the BAT of mice during cold-induced
thermogenesis. In addition, the expression of the mitokines *Gdf15* and *Fgf21* was significantly higher
in the BAT of cold-exposed mice after 6 and 24 h of cold exposure than at baseline, as were the
serum concentrations of GDF15 and FGF21 after 24 h of cold exposure (Figs. 4h–i and Extended
Data Figs. 5a–b).

To further explore the induction of UPR^{mt} *in vitro*, we treated differentiated immortalized brown
adipocytes with 1 μM CL 316,243, a β3-agonist. *Ucp1* expression in these cells was ~20-fold higher
after 3 h of treatment than at baseline (Fig. 4j), and the mRNA expression of mitochondrial
chaperones (*Hspd1* and *Dnaja3*) and proteases (*Clpp*, *Lonp1*, *Yme1l1*, *Htra2*, and *Usp30*) was 4–6-
fold higher after 6 h of treatment, while *Imm2l* expression was significantly lower after 3 and 6 h of
treatment than at baseline (Figs. 4k–n). The protein expression of HSPD1, DNAJA3, CLPP, and
LONP1 was also higher after CL 316,243 treatment than at baseline (Fig. 4o). The expression of the
related transcription factors *Atf5* and *Chop* was significantly higher, while expression of *Atf4* was
significantly lower, from 1.5 to 3 h of treatment compared with baseline (Fig. 4p). The mRNA
expression of *Gdf15* and *Fgf21*, as well as the concentration of GDF15 and FGF21 in the media, was
also significantly higher in differentiated immortalized brown adipocytes after CL 316,243 treatment
than at baseline (Figs. 4q,r and Extended Data Figs. 5c,d). Additionally, the expression of *Ucp1* and
mitochondrial chaperones and proteases in undifferentiated brown adipocytes were only slightly
higher (2–4-fold higher than controls) after 1.5 h of treatment with 1 μM CL 316,243 (Extended Data
Figs. 5e–i). The mRNA expression of *Gdf15*, *Fgf21*, *Atf4*, *Atf5*, and *Chop* was significantly lower in
undifferentiated brown adipocytes, unlike the differentiated cells, which have the mitochondrial
capacity for thermogenesis (Extended Data Figs. 5j,k). These results demonstrate a specific induction
of the UPR^{mt} and mitokine secretion in thermogenic brown adipocytes, as evidenced by greater
induction of the UPR^{mt} in differentiated cells than in undifferentiated cells.

Previous studies show that an imbalance between nDNA and mtDNA-encoded OXPHOS
proteins activates the UPR^{mt} [25]. To determine whether this mitochondrial/nuclear protein imbalance
was present in cold-exposed mice, we measured the mRNA expression of OXPHOS components in
previously published RNA sequencing data from the BAT of C57BL/6 mice after 24 h of cold exposure
(4°C) [26]. Of the 13 mtDNA-encoded OXPHOS complex genes, eight were present at significantly
lower levels after cold exposure, and of the nDNA-encoded oxidative phosphorylation proteins, 21
were present at significantly higher levels (Fig. 4s). We therefore concluded that these changes in
transcript levels would induce a mitochondrial/nuclear protein imbalance that may have been
responsible for the induction of the UPR^{mt} in cold-exposed BAT.

In summary, cold exposure increases mitochondrial demand in BAT, resulting in UPR^{mt}
induction, both in mice subjected to cold exposure and in differentiated immortalized brown
adipocytes treated with a β3-agonist. *Ucp1*, mitochondrial chaperones and proteases, and mitokines
increase in cold-exposed BAT and *in vitro* in β3-agonist-exposed differentiated brown adipocytes, but
less so in undifferentiated brown adipocytes. Activation of the UPR^{mt} in cold-exposed mice may result
from an imbalance between nDNA and mtDNA-encoded OXPHOS proteins.

The UPR^{mt} and Gdf15 expression in the BAT of mice is low under thermoneutral conditions

As shown in Fig. 4, the UPR^{mt} was induced by cold exposure; however, RT (22–24°C) housing
of mice also causes thermal stress[27]. To avoid thermal stress in Crif1^{BKO} mice, animals were housed
under thermoneutral conditions (30°C) for 8 weeks, which resulted in lower UPR^{mt}-related protein and

mRNA expression, as well as significantly lower UCP1 expression, in the BAT of 30°C-housed
Crif1[BKO] mice than RT-housed Crif1[BKO] mice (Figs. 5a,b). At both RT and 30°C, Crif1[BKO] mice had
lower protein expression of OXPHOS complexes I, III, and IV, as well as lower *Crif1* mRNA
expression, than control mice housed in the same conditions (Extended Data Figs. 6a,b). On
histological examination, both Crif1[BKO] and control mice were found to have unilocular adipocytes and
low UCP1 expression under thermoneutral conditions (Extended Data Fig. 6c). mRNA expression of
Gdf15 in BAT and the concentration of serum GDF15 in the 30°C-housed Crif1[BKO] mice was also
significantly lower than RT-housed Crif1[BKO] mice, implying an attenuated UPR[mt] response (Figs. 5c–d).
mRNA expression of *Fgf21* in the BAT of 30°C-housed Crif1[BKO] mice was significantly lower than RT-
housed Crif1[BKO] mice, while serum FGF21 concentration was not different between Ctrl and Crif1[BKO]
mice in 30°C housing (Extended Data Figs. 6d,e).

In the inguinal WAT of Crif1[BKO] mice, the expression of UCP1 and tyrosine hydroxylase (TH),
the rate-limiting enzyme in catecholamine synthesis, was significantly higher in RT-housed mice than
30°C-housed mice (Figs. 5e,f). *Gdf15* expression in the livers of Crif1[BKO] mice did not differ from that
of control mice (Extended Data Fig. 6f). The body weights of Crif1[BKO] mice were significantly lower
than control mice (Fig. 5g and Extended Data Figs. 6g,h), which is consistent with significantly higher
serum GDF15 concentrations (Fig. 5d) and significantly lower food intakes of 30°C-housed Crif1[BKO]
mice than control mice housed at the same temperature (Fig. 5h).

These results suggest that under thermoneutral conditions, an absence of thermal stress
blunts the UPR[mt] activation, as well as elevated *Gdf15* mRNA expression and serum GDF15
concentrations observed in Crif1[BKO] mice. Under these conditions, adaptive thermogenesis in the BAT
of Crif1[BKO] mice is not required, meaning mitochondrial proteostasis continues to function, even with
OXPHOS deficits. Therefore, sub-thermoneutral stress amplifies UPR[mt] activation in BAT, triggering a
response in inguinal WAT via GDF15 expression.

332 High circulating concentrations of GDF15 in Crif1[BKO] mice activate GFRAL-positive neurons in
333 the hindbrain

Neurons located in the area postrema (AP) and the nucleus of the solitary tract (NTS)
regions of the hindbrain are well positioned to detect blood-borne cues and project neurites into the
perivascular spaces of local fenestrated capillaries. Therefore, the AP is involved in interoception and
is a key detector of several peptides, including GDF15 and GLP-1, which play important roles in the
regulation of feeding behavior and metabolism[28,29]. Key studies demonstrate that GFRAL is the
central mediator of the metabolic effects of GDF15[12-15]. These studies converge on the AP and NTS in
the brainstem as the primary sites of GFRAL expression, linking GDF15 signaling to reduced food
intake and body weight regulation in rodent and nonhuman primate models[12-15].

To determine whether high circulating concentrations of GDF15 in Crif1[BKO] mice activate
subsets of glutamatergic neurons, including GFRAL-positive neurons, we assessed c-Fos expression
in GFRAL-positive neurons in the hindbrain of control and Crif1[BKO] mice. We first examined the
distribution of GFRAL-positive neurons in the AP and NTS of WT mice using immunohistochemical

staining. GFRAL-positive neurons were concentrated in the AP and NTS (Fig. 6a). By contrast, the
paraventricular nucleus of the hypothalamus (PVH) and arcuate nucleus, which regulate metabolism,
did not have GFRAL immunostaining (Figs. 6b,c). Next, we sectioned the brains of Crif1[BKO] and
control mice at the level of the medulla oblongata, focusing on the AP and NTS, and performed
GFRAL and c-Fos immunostaining. We found that the c-Fos expression of GFRAL-stained cells was
higher in Crif1[BKO] mice than in controls, especially in the AP (Figs. 6d,e). This implies that GFRAL
activity is higher in the AP of Crif1[BKO] mice than controls, which may account for the lower food
consumption of Crif1[BKO]-HFD mice (Fig. 2m).
Taken together, these results suggest that high circulating concentrations of GDF15 in
Crif1[BKO] mice are sufficient to activate GFRAL-positive neurons in the AP and NTS of the hindbrain,
which may explain the metabolic phenotype and resistance to obesity of Crif1[BKO] mice.
357

Retrograde neuronal tracing reveals the role of the sympathetic nervous system and GFRAL-
358
positive neurons in the GDF15-mediated browning of inguinal WAT in Crif1[BKO] mice
359
We next investigated the involvement of sympathetic neuronal activation in the browning of
inguinal WAT in Crif1[BKO] mice[30]. c-Fos expression was significantly higher in the sympathetic
preganglionic neurons of the intermediolateral nucleus region of the T7–T10 segment of the spinal
cord, which innervates the inguinal WAT, of Crif1[BKO] mice than control mice (Figs. 6f,g). To confirm
communication between GFRAL-stained neurons in the brain and sympathetic preganglionic neurons
of the spinal cord, we performed retrograde neuronal tracing using a pseudorabies virus. We injected
an enhanced green fluorescent protein (EGFP)-expressing pseudorabies virus unilaterally into the
inguinal WAT of ChAT-IRES-Cre:tdTomato mice, which express tdTomato in their sympathetic
preganglionic neurons, and WT mice (Fig 6h). Ninety-six (96) h post-injection, EGFP-stained neurons
were present in the intermediolateral nucleus region of the T6–L1 segment alone of the T1–L2 spinal
cord and in multiple regions of the brain, including the paraventricular nucleus of the hypothalamus,
locus coeruleus, and nucleus raphe pallidus (Fig. 6i and Extended Data Fig. 7a). At 120 h post-
injection, EGFP-stained neurons were present in a wider range of brain regions, including the AP and
NTS, primary motor cortex, hypothalamus, and reticular nuclei (Extended Data Fig. 7b). In the AP and
NTS of the mice (n = 2), 11.8% and 27.8% of GFRAL-stained neurons colocalized with EGFP,
respectively (Fig. 6j and Extended Data Fig. 7c). Moreover, compared with baseline levels, 24 h of
cold exposure in control mice resulted in a pronounced increase in c-Fos expression in GFRAL-
positive neurons (Extended Data Fig. 7d,e). These results suggest that GDF15-induced activation of
GFRAL in the AP and NTS could activate sympathetic nerves innervating inguinal WAT and induce
browning, as in Crif1[BKO] and/or cold-induced mice (Figs. 2g,h and Figs. 5e,f).
We next investigated whether GDF15 activates the preganglionic sympathetic neurons of the
spinal cord directly through cerebrospinal fluid (CSF). Significantly higher concentrations of GDF15
were found in the CSF of Crif1[BKO] mice than control mice, and the CSF concentration was significantly
positively correlated with the serum concentration (Figs. 6k,l). However, GFRAL was not detected by
RNA fluorescence *in situ* hybridization in the spinal cords of WT C57BL/6 mice (data not shown). This

suggests that the greater GFRAL activity in the AP of Crif1[BKO] mice was caused by higher serum
and/or CSF GDF15 concentrations activating the preganglionic sympathetic nerves of the spinal cord.
Despite this activation, the blood pressure of Crif1[BKO] mice was significantly lower, and their pulse
rates did not differ from those of control mice (Extended Data Figs. 7f,g).

Taken together, these findings indicate that activation of GFRAL in the AP and NTS of
Crif1[BKO] mice by high serum and/or CSF GDF15 concentrations activates the preganglionic SNS in
the spinal cord, leading to the browning of inguinal WAT. They suggest a direct link between systemic
metabolism and defective adaptive thermogenesis mediated by GDF15 in mice with BAT
mitochondrial dysfunction.

***Gfral* antisense oligonucleotides prevent GDF15-mediated phenotypic changes in Crif1[BKO]**
mice

To assess the role of GFRAL in adaptive thermogenesis within Crif1[BAT], we synthesized
antisense oligonucleotides (ASOs) targeting the *Gfral* gene. *Gfral* ASOs were designed to anneal to
specific complementary regions of *Gfral* mRNA, triggering RNase H-mediated degradation (Fig. 7a).
The phosphorothioate-modified backbone and bases enhance *Gfral* ASO stability and reduce
immunogenicity, ensuring sustained knockdown (Fig. 7b). We explored *Gfral* suppression in the AP
and NTS of the hindbrain using mouse models, with *Gfral* depletion achieved via
intracerebroventricular (ICV) injections of *Gfral* ASOs. Crif1[BKO] mice (n = 5 per group) fed *ad libitum*
over 12 weeks received either *Gfral* ASOs or control ASOs in the final 2 weeks. RNA scope analysis
of the hindbrain showed a significant 90% reduction in *Gfral* mRNA in the AP after 2 weeks of
treatment with *Gfral* ASOs (Figs. 7c–e).

We assessed energy expenditure and UCP1 expression in the inguinal WAT of Crif1[BKO] mice
treated with *Gfral* ASOs. These mice exhibited significantly higher food intake than control ASO-
treated Crif1[BKO] mice, although the body weight did not show significant change (Figs. 7f,g). Notably,
Gfral ASO diminished differences in energy expenditure caused by *Crif1* deletion (Figs. 7h,i).
Furthermore, we observed lower UCP1 expression in the inguinal WAT of *Gfral* ASO-treated Crif1[BKO]
mice compared with control ASO-treated mice (Fig. 7j). Histological assessment showed that *Gfral*
ASO-injected mice had fewer multilocular adipocytes than control ASO-injected mice (Fig. 7k). To
identify the role of GFRAL neurons in beige fat formation in mice exposed to cold, we housed GDF15
knockout (KO) mice in a cold environment for 1 days at 7°C and measured UCP1 expression. When
we cold-exposed GDF15 KO mice, serum concentrations of GDF15 and GFRAL-expressing neuronal
activation in the AP and NTS were significantly lower than cold-exposed WT mice (Extended Data
Figs. 7h,i). After 24 h of cold exposure, *Ucp1* mRNA expression was significantly lower in the inguinal
WAT of GDF15 KO mice than WT mice, while UCP1 protein expression in inguinal WAT tended to be
lower in GDF15 KO mice than WT mice (Extended Data Figs. 7j,l).

In summary, these results demonstrate that knockdown of *Gfral* in the hindbrain via ASOs
prevents the suppression of food intake, along with the elevation in energy expenditure, associated
with GDF15 expression. *Gfral* knockdown also results in lower UCP1 expression in inguinal WAT,

indicating a diminished thermogenic response. Moreover, the suppression of WAT browning in in
cold-exposed GDF15 KO mice confirms the critical role of GFRAL in regulating beige fat formation.
These findings underscore the significant impact of hindbrain GFRAL pathways on whole-body
energy balance in response to GDF15, which increases with mitochondrial stress in BAT (Fig. 7l).
428

Discussion
Adaptive thermogenesis in adipose tissue is a key homeostatic mechanism by which animals
maintain body temperature in cold environments. Brown adipocytes have developed a range of
compensatory mechanisms to offset these deficiencies and sustain thermogenesis; as such, this
process persists even when there are defects in the mitochondrial respiratory chain[5-7]. For example, in
response to mitochondrial dysfunction, brown adipocytes increase UCP1 expression, which elevates
metabolic rate and heat production[7]. Additionally, activation of the SNS increases catecholamine
release, which stimulates lipid and glucose metabolism for heat production. Alternative pathways,
such as ketogenesis, also support thermogenesis when mitochondrial function is compromised[31].
These compensatory mechanisms enable brown adipocytes to maintain thermogenic activity despite
mitochondrial dysfunction; however, they may be less efficient and therefore insufficient for
maintaining metabolic rate. As a result, communication between brown and white adipose depots is
required to offset OXPHOS defects in brown adipocytes. In line with this, previous findings indicate an
association between BAT mtDNA-ETC subunit mRNA expression, proteome imbalance, impaired
thermogenesis, and reduced adiposity. Likewise, mitochondrial transcription factor A deficiency in
BAT is associated with ETC proteome imbalance, defective respiration, and reduced adiposity[7].
However, the precise mechanisms linking mitochondrial stress in BAT to a systemic adaptive
metabolic response remains unclear.
Previous studies indicate that GDF15 expression increases in response to cellular stress, such
as oxidative stress and mitochondrial respiratory chain dysfunction[10]. Mitochondrial dysfunction
typically leads to an increase in oxidative stress, which can trigger the ISR[32] and lead to subsequent
upregulation of GDF15. In addition, ablation of CRIF1 function, specifically in liver and adipose tissue,
results in elevated circulating levels of GDF15[9,11] and GFRAL-positive neuron activation in the
hindbrain, which coincides with increased UCP1 expression in inguinal WAT. UCP1 transgenic mice
that overexpress UCP1 with skeletal muscle specificity also exhibit elevated plasma GDF15 levels
and WAT browning[33]. Conversely, GDF15 deficiency leads to diminished inguinal WAT UCP1
expression in these models[9,30]. Therefore, GDF15, upregulated by cellular stress, appears to be
instrumental in triggering adaptive thermogenesis in WAT. Indeed, several studies show that GDF15
regulates metabolism and increases thermogenesis in WAT when mitochondrial defects in peripheral
organs, such as skeletal muscle, exist[9,30].
In line with previous findings, we observed a relationship between cellular stress and
metabolic regulation mediated by GDF15. In our study, UPR^{mt} and the ISR were activated in Crif1^{BKO}
mice: this was likely responsible for higher BAT GDF15 expression and serum GDF15 concentrations
in Crif1^{BKO} mice. The deletion or dysfunction of CRIF1 in tissues typically results in impaired OXPHOS

function. This often triggers the UPR[mt] due to mitochondrial stress from accumulated misfolded
proteins[34,35]. While the impairment of BAT OXPHOS is directly linked to CRIF1 deletion, downstream
effects, such as protection from obesity and insulin resistance, are consistently observed across
different tissues with *Crif1* gene deletion[9-11]. Furthermore, plasma levels of GDF15 in mice with *Crif1*
deletion—whether in skeletal muscle, WAT, or liver—are comparable to those we observed in Crif1[BKO]
mice, levels that stimulate activation of GFRAL neurons in the hindbrain. Therefore, the metabolic
benefits of UPR[mt] activation, combined with the increase in GDF15, appear to result from activation of
the GDF15–GFRAL axis in the hindbrain.

The involvement of the GDF15–GFRAL axis in SNS activation has been previously
investigated. GDF15-expressing xenografts induce thermogenic gene expression in BAT and lipolytic
gene expression in WAT[36]. GDF15 also stimulates hepatic triglyceride export via β-adrenergic
signaling, while chemical sympathectomy completely reverses GDF15-induced body mass loss[37,38].
The mechanism behind SNS activation by GDF15 remains elusive, but it may involve hypothalamic
neurons in the arcuate nucleus and PVH, which regulate energy homeostasis[39]. GDF15 might also
influence the hypothalamic-pituitary-adrenal axis[40]: this plays a crucial role in the stress response,
although the impact of GDF15 on the hindbrain receptor GFRAL within the context of hypothalamic-
pituitary-adrenal axis and proopiomelanocortin neuron activation is yet to be clarified in Crif1[BKO] mice.
Spinal preganglionic neurons, the final controllers of the SNS crucial for environmental stress
adaptation, are not believed to express GFRAL[41]; therefore, activation of these neurons in Crif1[BKO]
mice may be mediated by neuronal or humoral pathways. We confirmed the activation of T7–T10
spinal preganglionic neurons in Crif1[BKO] mice and identified the responsible neural pathway using
retrograde tracing with a pseudorabies virus. This virus traced axons from inguinal WAT to cell bodies
in the spinal cord preganglion and hindbrain. Fluorescence microscopy enabled visualization of
neurons connected to inguinal WAT. Our results suggest that the GDF15–GFRAL axis forms a unique
neural network involving GFRAL-positive neurons, spinal preganglionic neurons, and the end organ,
inguinal WAT, which results in beige adipocyte formation in mice with high GDF15 concentrations.

Recent studies have demonstrated the therapeutic potential of activating the GFRAL–SNS
through GDF15 administration[42,43]. Administration of GDF15 not only suppresses appetite but also
counteracts reductions in energy expenditure, which leads to more effective weight loss and a greater
reduction in non-alcoholic fatty liver disease than caloric restriction alone[32]. The influence of GDF15
on energy expenditure is mediated by a GFRAL–β-adrenergic signaling mechanism that enhances
fatty acid oxidation in mouse skeletal muscle. In particular, the GFRAL–β-adrenergic signaling axis
increases fatty acid oxidation and stimulates futile calcium cycling in mouse skeletal muscle[42].
Additionally, GDF15 enhances insulin action in obese rodents without the need for weight loss,
increasing insulin sensitivity due to decreased glucose production and increased glucose uptake. This
insulin-enhancing effect operates through the GFRAL receptor and β-adrenergic signaling, impacting
both the liver and adipose tissue[43]. Nonetheless, tissue responses and therapeutic effects of a
pharmacological dose of GDF15 vary considerably across species and even among individuals within
a species because of genetic, metabolic, and environmental factors[44]. In addition, there are significant

differences in GDF15 plasma levels resulting from pharmacological GDF15 administration and those
resulting from gene editing, such as in Crif1[BKO] mice. Although we observed higher GDF15 plasma
concentrations in Crif1[BKO] (about 500-1000 pg/mL) than in control mice, these levels are of a lower
magnitude than those resulting from pharmacological GDF15 administration (about 50,000-110,000
pg/mL)[42], meaning that the energy expenditure seen in the skeletal muscle of mice treated with
GDF15 might be less pronounced in Crif1[BKO] mice. However, even in Crif1[BKO] mice, we observed
lower food intake and body weights and higher energy expenditure than in control mice. These results
suggest that even at a subtherapeutic concentration, GDF15 can have a significant impact on appetite
and metabolic health.

Cold exposure stimulates mitochondrial biogenesis in brown adipocytes through a complex
process involving several signaling pathways. Specifically, it stimulates expression of the
transcriptional co-activator PPARGC1A, which regulates mitochondrial biogenesis[45]. PPARGC1A
activates several transcription factors, including nuclear respiratory factor 1 and mitochondrial
transcription factor A. These transcription factors control the expression of genes involved in
mitochondrial biogenesis and function in brown adipocyte[46], which contribute to greater thermogenic
capacity of BAT in response to cold exposure. We found that GDF15 expression was induced in the
BAT of mice exposed to cold temperatures, and that this was accompanied by increases in the
expression of genes involved in mitochondrial biogenesis and the UPR[mt], as well as mitochondrial
chaperones and proteases. The induction of the UPR[mt] was more marked in differentiated brown
adipocytes, which are capable of thermogenesis, than in undifferentiated cells; however, the induction
of the UPR[mt] was less marked under thermoneutral conditions in mice with defective OXPHOS. These
results suggest that the induction of the mitochondrial stress response is dependent on the metabolic
workload of the cells. This may be analogous to the induction of the ISR by mitochondrial defects
dependent on the metabolic state of the cell[32], and the induction of endoplasmic reticulum stress in
WAT resulting from the elevated metabolic workload present in obesity[47,48].

In conclusion, GDF15 expression induced by the UPR[mt] in BAT causes browning of inguinal
WAT through the activation of GFRAL-expressing spinal preganglionic sympathetic neurons. These
findings outline a compensatory mechanism involving both BAT and WAT that may provide novel
targets for therapeutic interventions aimed at increasing energy expenditure in patients with obesity
and the metabolic syndrome.

532
533 **Materials and Methods**

534
Animals
To generate brown adipocyte-specific Crif1-deficient mice, Crif1 floxed mice[49] were cross-mated with
UCP1-Cre mice (Jackson Laboratory, Cat. 024670). C57BL/6J male mice were purchased from Orient
Bio (Seongnam, Korea). GDF15 KO mice derived from the inbred C57BL/6 strain were provided by Dr.
S. Lee (Johns Hopkins University School of Medicine, Baltimore, MD). All mice had free access to a
standard chow diet (ENVIGO, Teklad global 18% protein, Cat. 2918C) and water unless indicated. To

induce obesity, mice were fed a high-fat diet (HFD; 60% of total energy intake as fat, ENVIGO, Cat.

TD.06414). Animals were housed under controlled temperature conditions (22 ± 1°C) unless indicated

and subjected to a 12 h light–dark cycle, with light from 08:00 to 20:00 h. During the cold challenge

(5~7°C), body temperatures of Ctrl or Crif1^{BKO} were monitored with an implantable programmable

temperature transponder (Bio Medic Data Systems, Seaford, USA). All animal procedures were

approved by the Institutional Animal Care and Use Committee of the Chungnam National University

School of Medicine (Daejeon, Korea).

Metabolic phenotyping

Food intake levels and body weights were monitored weekly after weaning until sacrifice. Body

composition (lean mass and fat mass) was measured using dual X-ray absorptiometry (InAnlyzer,

MEDIKORS, Seongnam, Korea) during the late light phase at the indicated ages. Energy expenditure

were determined using an OxyletProTM system (Panlab, Barcelona, Spain). Mice were placed in the

OxyletProTM for 48 h to acclimatize to these conditions before measurement. During this period, day–

night cycles were the same as those during the initial housing of the animals and food was provided

as pellets on the floor of the OxyletProTM. Data were analyzed using METABOLISM software,

version 2.2 (Panlab). For the glucose tolerance test, mice were fasted for 16 h, and then 2 g glucose

per kilogram of body weight was injected into the intraperitoneal cavity. The insulin tolerance test was

performed by measuring blood glucose after 6 h of fasting followed by intraperitoneal injection of 0.75

U/kg insulin (Humalog Lilly, Indianapolis, IN, USA). Blood samples were obtained from the tail vein for

glucose measurement immediately before, and at the indicated times after, injections. Glucose levels

were measured using a glucometer (ACCU-CHEK, Roche Diabetes Care, Inc., IN, USA).

Immunostaining of brain and spinal cord sections

Mice were anesthetized with isoflurane and perfused with 20 mL normal saline followed by 20 mL 4%

paraformaldehyde via the left ventricle of the heart. Whole brain was collected and fixed with 4%

paraformaldehyde for 24 h at 4°C, before being dehydrated in a PBS-based 30% sucrose solution

until the tissues sank to the bottom of the container, usually after 48 h at 4°C. Coronal brain sections

and transverse spinal cord sections were collected at a 30 µm thickness using a cryostat (Leica,

Wetzlar, Germany). Brain slices were blocked with 5% donkey serum in 0.3% PBS-T for 1 h at RT

and then incubated with a sheep anti-GFRAL antibody (1:500, Thermofisher, Cat. PA5-47769) and/or

a rabbit anti-c-Fos antibody (1:500, Abcam, Cat. ab190289) at 4°C overnight. In the case of spinal

cord staining, heat-induced epitope retrieval was required before the blocking step. The heat-induced

antigen retrieval method was modified from a reference method[50] to use 10 mM sodium citrate buffer

and double boil the slides within the buffer using a 60°C water bath. After antigen retrieval, spinal cord

slices were blocked in the same way as brain slices and incubated with a goat anti-ChAT antibody

(1:100, Chemicon, Cat. AB144P) and a rabbit anti-c-Fos antibody (1:500, Abcam, Cat. ab190289) at

4°C overnight. Slices were then incubated with combinations of secondary antibodies from Invitrogen,

including donkey anti-rabbit 647 (1:500, Cat. A31573), donkey anti-goat plus 488 (1:200, Cat.

A32814), and donkey anti-sheep 488 (1:500, Cat. A11015), for 1.5 h at RT. For nuclear staining,
slides were treated with DAPI (1:10000) for 10 min before mounting. *Z*-stack images were acquired
with a confocal microscope (Zeiss, LSM 780), and maximum-intensity *z*-stacks were produced with
Zen Black software (Zeiss). Cross-sectional images of the intermediolateral nucleus were taken at
40× magnification, and brain images and regional images of spinal cords were taken at 20×
magnification.
586

CSF collection for GDF15 quantification

Mice were anesthetized with 2% isoflurane and mounted on stereotaxic frames using ear bars. The
mouse snout was tilted down to form an angle of 30–45° between the frame base and imaginary line
connecting bregma to lambda following the reference method[51]. Next, the scalp and muscle covering
the dura mater over the fourth ventricle was gently removed, a 36-gauge beveled needle (WPI, Cat.
NF36BV) connected to 1 mL syringe with tubing was slowly inserted into the dura mater, and the CSF
was gently aspirated; approximately 8–10 µL of CSF was collected and used for GDF15 quantification.
594

Retrograde tracing the neurons innervating inguinal WAT

Mice were anesthetized with 2% isoflurane and the flank was exposed using the lateral decubitus
position. PRV-CAG-EGFP (BrainVTA, Cat. P03001) was injected into unilateral inguinal WAT (4 ×
200 nL) using a 36-gauge beveled needle (WPI, Cat. NF36BV). After the injection, mice recovered on
a heating pad until they were awake and mobile. Mice were sacrificed at 96 and 120 h post-injection,
and the brain and spinal cord were isolated to identify sites of EGFP expression and evaluate co-
localization of EGFP with ChAT in the spinal cord and GFRAL in the AP/NTS.
602

ICV administration of Gfral ASOs

Mice at 8–13 weeks of age with average weights of 25–30 g were anesthetized with 3% isoflurane
and anesthesia was maintained with 2% isoflurane administered at 0.5 L/min. The head was fixed to a
stereotaxic platform (RWD Life Science, China, Cat. 68045). Next, a 36-gauge NanoFil needle was
inserted through a burr hole into the right lateral cerebral ventricle using stereotaxic coordinates –0.58
mm posterior from bregma, 1.05 mm lateral from bregma, and 2.2 mm deep from the bregma plane.
Injection of a contro or *Gfral* ASO (GGTATATTTTGATAGAAGAACC) was performed using a Pump
11 Elite Nanomite pump and Nanomite Injector Unit (Harvard Apparatus; Catalog# 70-4507, Serial#
D-301251) at a rate of 2 µL/min. Food intake and body weight measurements, as well as metabolic
chamber studies, were performed 1 week postoperatively.
613

Western blotting

Mouse tissues were homogenized in lysis buffer (50 mM Tris-HCl, pH 7.4; 150 mM NaCl; 1 mM EDTA,
pH 8.0; 0.1% Triton X-100) containing a protease inhibitor cocktail (#11836145001, Roche, Basel,
Switzerland) and phosphatase inhibitors (04906837001, Roche) on ice for 30 min using a TissueLyser
II (Qiagen, Venlo, Netherlands). After centrifugation at 16,000 × *g* for 15 min, the protein

concentrations of the supernatants were measured using a BCA protein assay (#23227, Thermo
Fisher Scientific). From each sample, 50 µg protein was loaded onto 8–12% polyacrylamide gels and
electrophoresis was performed. The separated proteins were then electrotransferred to 0.45 µm
PVDF membranes (#IPVH00010, Millipore) at 200 mA for 2 h. Membranes were blocked with 5%
skimmed milk (#T145.2, Roth) in TBS/T buffer (20 mM Tris, 150 mM NaCl, 0.1% Tween 20, pH 7.6)
for 1 h and then incubated with primary antibodies overnight at 4°C. After washing three times with
TBS/T, the membranes were incubated with secondary antibodies for 1 h at RT and visualized using
ECL solution (#34580, Thermo Fisher Scientific). Target protein levels were normalized to those of β-
actin, α-tubulin, or glyceraldehyde 3-phosphate. The antibodies used are listed in Table S1.
628

Cell cultures
Immortalized brown adipocytes were provided by Dr. Shingo Kajimura (University of California San
Francisco, CA). Immortalized brown adipocytes were maintained in Dulbecco's Modified Eagle's
Medium (DMEM; Welgene, Daegu, South Korea, Cat. LM001-05) supplemented with 10% fetal
bovine serum (CYTIVA, Utah, US, Cat. SH30919.03) and 1% penicillin and streptomycin (Welgene,
Daegu, South Korea, Cat. LS202-02). Forty-eight hours post-confluence, the cells were differentiated
with 0.5 mM 3-isobutyl-1-methylxanthine, 1 µM dexamethasone, 1 µg/mL insulin, 1 nM T3, and 1 µM
rosiglitazone. Two days later, the medium was replaced with DMEM containing 10% FBS, 1 nM T3
and 1 µg/mL insulin, and was changed every 2 days for a total of 7 days to achieve complete
differentiation. The cells were treated with CL 316,243 (1 µM) for 1.5, 3, or 6 h, or oligomycin (Sigma
Aldrich, Cat. O4876, 10 µg/mL), CCCP (Sigma Aldrich, Cat. C2759, 2 µg/mL), rotenone (Sigma
Aldrich, Cat. R8875, 1 µM/mL), or gamitrinib (Legochem Biosciences, 1 or 2 µM) for 24 h at 37°C.
641

RNA sequencing (RNA-seq)
RNA was isolated from the BAT of 8-week-old control and Crif1[BKO] mice. To construct cDNA libraries,
1 µg RNA and a TruSeq RNA Library Prep Kit v2 (RS-122-2001, Illumina, San Diego, CA, USA) were
used, and the results were quantified by qPCR using an 2100 Bioanalyzer (Agilent Technology Inc.,
Santa Clara, CA, USA). The libraries were used for 100 nt paired-end sequencing by an Illumina
HiSeq4000 (Illumina). After removing the low-quality and adapter sequences using Trimmomatic, the
reads were aligned with the Mus musculus genome (mm10) using HISAT (ver. 2.0.5). Two types of
indexes were used for alignment (a global, whole-genome index and tens of thousands of small local
indexes), which were downloaded from the UCSC table browser (http://genome.uscs.edu). StringTie
(ver. 1.3.3b) was used to assemble the transcript, and provided the relative abundance estimates as
fragments per kilobase of exon per million fragments mapped (FPKM) values of the transcript or gene.
After excluding the genes with one more than zero FPKM values, the signal value (FPKM+1) was
transformed to a base 2 logarithm and normalized by quantile normalization methods to reduce
systematic bias. These values were used for the analysis of differentially expressed genes in the
mouse groups. The statistical significance of differentially expressed gene values was calculated
using independent t-tests ($P<0.05$) and fold change ($|FC|≥2$). The false discovery rate, which

estimates the frequency of type I statistical errors, was determined by adjusting the *P*-value using the

Benjamini and Hochberg algorithm.

Enzyme-linked immunosorbent assay

Cell supernatant was obtained from primary hepatocytes by centrifugation at 16,000 × *g* for 5 min.

Blood was collected from a retro-orbital sinus, incubated at RT for 2 h and then centrifuged at 600 × *g*

for 5 min to obtain serum or CSF. GDF15 (R&D Systems, Minneapolis, MN, USA, Cat. MGD150) and

FGF21 (R&D Systems, Cat. MF2100) concentrations were measured using ELISAs according to the

manufacturer's instructions.

Statistical analysis

Statistical analyses were performed using SPSS software, version 26.0 (IBM, Armonk, NY, USA).

Data are reported as mean ± standard errors of the mean (SEMs). A two-tailed Student's t-test was

used to determine differences between the groups. A *P*-value <0.05 was considered statistically

significant.

Data Availability

RNA-Seq data associated with this study are available from Gene Expression Omnibus [GSE233960]

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE233960).

Acknowledgments

This study was supported by a National Research Foundation of Korea (NRF) grant funded by the

Korea government, Ministry of Science and ICT (RS-2023-00210819) and NRF-

2023R1AC3003438, South Korea, the European Research Council (ERC-AdG-787702), and Global

Research Laboratory Program through the NRF of Korea (NRF 2017K1A1A2013124).

References

1   Cannon, B. & Nedergaard, J. Brown adipose tissue: function and physiological significance.
Physiological reviews (2004).
2   Nedergaard, J. & Cannon, B. The browning of white adipose tissue: some burning issues.
Cell metabolism **20**, 396-407 (2014).
3   Schulz, T. J. *et al.* Brown-fat paucity due to impaired BMP signalling induces compensatory
browning of white fat. *Nature* **495**, 379-383 (2013).
4   Piao, Z. *et al.* Reduced adiposity by compensatory WAT browning upon iBAT removal in mice.
Biochemical and biophysical research communications **501**, 807-813 (2018).
5   Pereira, R. O. *et al.* OPA1 deletion in brown adipose tissue improves thermoregulation and
systemic metabolism via FGF21. *Elife* **10**, e66519 (2021).
6   Verdeguer, F. *et al.* Brown adipose YY1 deficiency activates expression of secreted proteins
linked to energy expenditure and prevents diet-induced obesity. *Molecular and cellular biology*
(2015).
7   Masand, R. *et al.* Proteome imbalance of mitochondrial electron transport chain in brown
adipocytes leads to metabolic benefits. *Cell metabolism* **27**, 616-629. e614 (2018).
8   Paulo, E. *et al.* Brown adipocyte ATF4 activation improves thermoregulation and systemic
metabolism. *Cell reports* **36** (2021).
9   Kang, S. G. *et al.* Differential roles of GDF15 and FGF21 in systemic metabolic adaptation to
the mitochondrial integrated stress response. *IScience* **24** (2021).
10   Choi, M. J. *et al.* An adipocyte-specific defect in oxidative phosphorylation increases systemic
energy expenditure and protects against diet-induced obesity in mouse models. *Diabetologia*
63, 837-852 (2020).
11   Chung, H. K. *et al.* Growth differentiation factor 15 is a myomitokine governing systemic
energy homeostasis. *J Cell Biol* **216**, 149-165 (2017).
12   Emmerson, P. J. *et al.* The metabolic effects of GDF15 are mediated by the orphan receptor
GFRAL. *Nature medicine* **23**, 1215-1219 (2017).
13   Mullican, S. E. *et al.* GFRAL is the receptor for GDF15 and the ligand promotes weight loss in
mice and nonhuman primates. *Nature medicine* **23**, 1150-1157 (2017).
14   Yang, L. *et al.* GFRAL is the receptor for GDF15 and is required for the anti-obesity effects of
the ligand. *Nature medicine* **23**, 1158-1166 (2017).
15   Hsu, J.-Y. *et al.* Non-homeostatic body weight regulation through a brainstem-restricted
receptor for GDF15. *Nature* **550**, 255-259 (2017).
16   Kim, S. J. *et al.* CRIF1 is essential for the synthesis and insertion of oxidative phosphorylation
polypeptides in the mammalian mitochondrial membrane. *Cell Metab* **16**, 274-283 (2012).
17   Fiorese, C. J. *et al.* The Transcription Factor ATF5 Mediates a Mammalian Mitochondrial UPR.
Curr Biol **26**, 2037-2043 (2016).
18   Quiros, P. M. *et al.* Multi-omics analysis identifies ATF4 as a key regulator of the mitochondrial
stress response in mammals. *J Cell Biol* **216**, 2027-2045 (2017).
19   Khan, N. A. *et al.* mTORC1 Regulates Mitochondrial Integrated Stress Response and
Mitochondrial Myopathy Progression. *Cell Metab* **26**, 419-428 e415 (2017).
20   Montero, R. *et al.* GDF-15 Is Elevated in Children with Mitochondrial Diseases and Is Induced
by Mitochondrial Dysfunction. *PLoS One* **11**, e0148709 (2016).
21   Münch, C. & Harper, J. W. Mitochondrial unfolded protein response controls matrix pre-RNA
processing and translation. *Nature* **534**, 710-713 (2016).
22   Chouchani, E. T. *et al.* Mitochondrial ROS regulate thermogenic energy expenditure and
sulfenylation of UCP1. *Nature* **532**, 112-116 (2016).
23   Lu, Y. *et al.* Mitophagy is required for brown adipose tissue mitochondrial homeostasis during
cold challenge. *Sci Rep* **8**, 8251 (2018).
24   Forner, F. *et al.* Proteome differences between brown and white fat mitochondria reveal
specialized metabolic functions. *Cell Metab* **10**, 324-335 (2009).
25   Houtkooper, R. H. *et al.* Mitonuclear protein imbalance as a conserved longevity mechanism.
Nature **497**, 451-457 (2013).
26   Quesada-Lopez, T. *et al.* The lipid sensor GPR120 promotes brown fat activation and FGF21
release from adipocytes. *Nat Commun* **7**, 13479 (2016).
27   Feldmann, H. M., Golozoubova, V., Cannon, B. & Nedergaard, J. UCP1 ablation induces
obesity and abolishes diet-induced thermogenesis in mice exempt from thermal stress by
living at thermoneutrality. *Cell Metab* **9**, 203-209 (2009).

28 Tsai, V. W. *et al.* GDF15 mediates adiposity resistance through actions on GFRAL neurons in
the hindbrain AP/NTS. *Int J Obes (Lond)* **43**, 2370-2380 (2019).
29 Zhang, C. *et al.* Area Postrema Cell Types that Mediate Nausea-Associated Behaviors.
Neuron **109**, 461-472 e465 (2021).
30 Huesing, C. *et al.* Sympathetic innervation of inguinal white adipose tissue in the mouse. *J
Comp Neurol* **529**, 1465-1485 (2021).
31 Panic, V. *et al.* Mitochondrial pyruvate carrier is required for optimal brown fat thermogenesis.
Elife **9**, e52558 (2020).
32 Mick, E. *et al.* Distinct mitochondrial defects trigger the integrated stress response depending
on the metabolic state of the cell. *Elife* **9**, e49178 (2020).
33 Ost, M. *et al.* Muscle-derived GDF15 drives diurnal anorexia and systemic metabolic
remodeling during mitochondrial stress. *EMBO reports* **21**, e48804 (2020).
34 Durieux, J., Wolff, S. & Dillin, A. The cell-non-autonomous nature of electron transport chain-
mediated longevity. *Cell* **144**, 79-91 (2011).
35 Nargund, A. M., Pellegrino, M. W., Fiorese, C. J., Baker, B. M. & Haynes, C. M. Mitochondrial
import efficiency of ATFS-1 regulates mitochondrial UPR activation. *Science* **337**, 587-590
(2012).
36 Chrysovergis, K. *et al.* NAG-1/GDF-15 prevents obesity by increasing thermogenesis,
lipolysis and oxidative metabolism. *International journal of obesity* **38**, 1555-1564 (2014).
37 Luan, H. H. *et al.* GDF15 is an inflammation-induced central mediator of tissue tolerance. *Cell*
178, 1231-1244. e1211 (2019).
38 Suriben, R. *et al.* Antibody-mediated inhibition of GDF15–GFRAL activity reverses cancer
cachexia in mice. *Nature medicine* **26**, 1264-1270 (2020).
39 Johnen, H. *et al.* Tumor-induced anorexia and weight loss are mediated by the TGF-β
superfamily cytokine MIC-1. *Nature medicine* **13**, 1333-1340 (2007).
40 Cimino, I. *et al.* Activation of the hypothalamic–pituitary–adrenal axis by exogenous and
endogenous GDF15. *Proceedings of the National Academy of Sciences* **118**, e2106868118
(2021).
41 McCarthy, M. Allen Brain Atlas maps 21 000 genes of the mouse brain. *The Lancet Neurology*
5, 907-908 (2006).
42 Wang, D. *et al.* GDF15 promotes weight loss by enhancing energy expenditure in muscle.
Nature **619**, 143-150 (2023).
43 Sjøberg, K. A. *et al.* GDF15 increases insulin action in the liver and adipose tissue via a β-
adrenergic receptor-mediated mechanism. *Cell Metabolism* **35**, 1327-1340. e1325 (2023).
44 Moon, J. S. *et al.* Growth differentiation factor 15 protects against the aging-mediated
systemic inflammatory response in humans and mice. *Aging cell* **19**, e13195 (2020).
45 Puigserver, P. *et al.* A cold-inducible coactivator of nuclear receptors linked to adaptive
thermogenesis. *Cell* **92**, 829-839 (1998).
46 Wu, Z. *et al.* Mechanisms controlling mitochondrial biogenesis and respiration through the
thermogenic coactivator PGC-1. *Cell* **98**, 115-124 (1999).
47 Ozcan, U. *et al.* Endoplasmic reticulum stress links obesity, insulin action, and type 2 diabetes.
Science **306**, 457-461 (2004).
48 Zhao, Q. *et al.* A mitochondrial specific stress response in mammalian cells. *The EMBO
journal* **21**, 4411-4419 (2002).
49 Kwon, M. c. *et al.* Crif1 is a novel transcriptional coactivator of STAT3. *The EMBO journal* **27**,
642-653 (2008).
50 Jiao, Y. *et al.* A simple and sensitive antigen retrieval method for free-floating and slide-
mounted tissue sections. *Journal of neuroscience methods* **93**, 149-162 (1999).
51 Šakić, B. Cerebrospinal fluid collection in laboratory mice: Literature review and modified
cisternal puncture method. *Journal of neuroscience methods* **311**, 402-407 (2019).

Figures and Tables

795
796
Figure 1. Crif1[BKO] **mice with a brown adipocyte-specific defect in oxidative phosphorylation**
show higher brown adipose tissue (BAT) expression of mitochondrial unfolded protein
response proteins. a, Immunoblots and band densities for CRIF1, oxidative phosphorylation
proteins, and UCP1 in the BAT of male control (Ctrl) and Crif1[BKO] mice. **b**, Representative electron
microscopic images of brown adipocytes. L: lipid, M: mitochondria. Bar, 0.5 μm. **c**, Hematoxylin and
eosin (H&E)-stained and UCP1-immunostained sections of BAT. Bar, 50 μm. **d**, Immunoblots and
band densities for mitochondrial unfolded protein response-associated transcription factors. **e,**
Immunoblots and band densities for mitochondrial chaperones and proteins. *, $P < 0.05$; **, $P < 0.01$;
***, $P < 0.001$ *vs.* Ctrl by Student's t-test. Data are represented as mean ± standard error of mean
(SEM). See also Extended Data Figure 1.
807

808
809
Figure 2. Crif1^{BKO} mice exhibit browning of inguinal white adipose tissue (WAT) and
demonstrate resistance to high-fat diet-induced obesity. a–c, Body weight (**a**), daily food
consumption (**b**), and organ/body weight ratios (c) for male control (Ctrl) and Crif1^{BKO} mice (n=5 per

group). **d, e,** ANCOVA of energy expenditure (EE) against body weight (**d**) and ANCOVA-adjusted EE

(**e**) in male Ctrl and Crif1^{BKO} mice (n=8 per group) during paired feeding. **f**, Body temperatures of male

Ctrl and Crif1^{BKO} mice housed at room temperature (22–24℃) following cold exposure at 5℃ for 1

week (n=4 per group). **g**, Immunoblots and band densities for UCP1 and PPARGC1A in inguinal WAT.

h, Representative images of H&E-stained and UCP1-immunostained sections of inguinal WAT in

mice fed a normal chow diet (NCD). Bar, 50 µm. **i,** Representative electron microscopic images of

inguinal white adipocytes. L: lipid, M: mitochondria. Bar, 0.5 µm. **j–l**, Body weights (**j**), organ/body

weight ratios (**k**), and fat percentages (**l**), according to densitometry. **m**, Daily food intake of male Ctrl-

HFD and Crif1^{BKO}-HFD mice fed a HFD for 8 weeks (n=3–4 per group). **n**, Immunoblots and band

densities for UCP1 and PPARGC1A in inguinal WAT. **o**, Representative images of H&E-stained and

UCP1-immunostained sections of inguinal WAT. Bar, 50 µm. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$

vs. Ctrl by Student's t-test. Data are represented as mean ± SEM. See also Extended Data Figure 2

for *ad libitum* feeding results and Extended Data Figure 3 for HFD results.

827
828
Figure 3. *Gdf15* expression increases in response to mitochondrial unfolded protein response
activation in brown adipose tissue (BAT). a, Volcano plot of differentially expressed secretory
protein-encoding genes in the BAT of Crif1[BKO] mice, identified using RNA sequencing. **b**, Relative
Gdf15 expression in the BAT of male control (Ctrl) and Crif1[BKO] mice (n=5 per group), as
characterized in Fig. 1. **c**, Serum GDF15 concentrations in male Ctrl and Crif1[BKO] mice. **d**, Relative
Gdf15 expression in the BAT of the male Ctrl-HFD and Crif1[BKO]-HFD mice, as characterized in Fig. 3.
e, Serum GDF15 concentration in male Ctrl-HFD and Crif1[BKO]-HFD mice. **f–i**, Relative mRNA
expression of mitochondrial unfolded protein response-associated genes (**f**), transcription factors (**g**),
and *Gdf15* (**h**) in differentiated brown adipocytes treated with oxidative phosphorylation inhibitors (10
μg/mL oligomycin, 2 μg/mL CCCP, or 1 μM rotenone) for 24 h. (**i**) GDF15 concentration in the
medium. **j–l,** Relative mRNA expression of mitochondrial unfolded protein response-associated genes
(**j**), transcription factors (**k**), and *Gdf15* (**l**) in differentiated brown adipocytes treated with gamitrinib

(G-TTP) for 24 h. **m**, GDF15 concentration in cell medium. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$ *vs.*

0 h, Ctrl, Ctrl-HFD, or DMSO by Student's t-test. Data are represented as mean ± SEM. See also

Extended Data Figure 4.

845
846
Figure 4. The expression of mitochondrial chaperones and proteases is high in the brown
adipose tissue (BAT) of mice with cold-induced thermogenesis. a–f, Relative mRNA expression
of *Ucp1* (**a**), and genes encoding mitochondrial chaperones (**b**), mitochondrial matrix proteins (**c**),
inner-membrane (**d**), inter- and outer-membrane proteases (**e**), and transcription factors (**f**) in BAT
from wild-type mice housed at 30°C following cold exposure at 5°C for 24 h (n=5–7 per group). **g**,
Representative immunoblots and band densities for transcription factors in the BAT of cold-exposed
mice. **h**, Relative GDF15 expression in the BAT of cold-exposed mice. (**i**) Serum GDF15
concentrations of cold-exposed mice. **j–n**, Relative mRNA expression of *Ucp1* (**j**), mitochondrial
chaperones (**k**) and proteases **l-n,** in differentiated immortalized brown adipocytes treated with 1 µM
CL 316,243. **o,** Representative immunoblots for UCP1, mitochondrial chaperones, and proteases. **p,**

Representative immunoblots for transcription factors and relative mRNA expression. **q**,**r**, Relative *Gdf15* expression and GDF15 concentration in cell medium. **s**, Heatmap of differentially expressed oxidative phosphorylation genes encoded by mitochondrial DNA (mtDNA) and nuclear DNA (nDNA) from previously published RNA-seq data[26]. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$ *vs*. 0 h by Student's t-test. Data are represented as mean ± SEM. See also Extended Data Figure 5.

863
864
Figure 5. Activation of the mitochondrial unfolded protein response and *Gdf15* expression in
brown adipose tissue (BAT) is low under thermoneutral conditions. a, Immunoblots and band
densities for CRIF1, mitochondrial unfolded protein response, and UCP1 in the BAT of 12–13-week-
old male control (Ctrl) and Crif1^{BKO} mice housed at room temperature (22–24°C) (Ctrl-RT or Crif1^{BKO}-
RT) or 30°C (Ctrl-30°C or Crif1^{BKO}-30°C) for 8 weeks. **b**, Relative mRNA expression of *Ucp1* and
mitochondrial unfolded protein response genes in BAT. **c**, Relative *Gdf15* expression in BAT. **d**,
Serum GDF15 concentration. **e**, Immunoblots and band densities for UCP1, tyrosine hydroxylase
(TH), and PPARGC1A in inguinal white adipose tissue (WAT). **f**, Representative images of UCP1-
immunostained inguinal WAT. Bar, 50 μm. **g,h**, Body weight (**g**) and daily food intake (**h**) of 15–16-
week-old male Ctrl and Crif1^{BKO} mice housed at room temperature (22–24°C) (Ctrl-RT or Crif1^{BKO}-RT)
or 30°C (Ctrl-30°C or Crif1^{BKO}-30°C) for 9 weeks (n=3–4 per group). *, $P < 0.05$; **, $P < 0.01$; ***, $P <$
0.001 *vs.* Ctrl by Student's t-test; #, $P < 0.05$; ##, $P < 0.01$; ###, $P < 0.001$ *vs.* RT by Student's t-test.
Data are represented as mean ± SEM. See also Extended Data Figure 6.
878

879
880
Figure 6. Activation of GFRAL-positive neurons in the hindbrain, combined with retrograde
neuronal tracing of both the sympathetic nervous system and GFRAL neurons in Crif1[BKO] **mice.**
a–c, Representative images of the area postrema (AP) and nucleus of the solitary tract (NTS) (**a**),
paraventricular nucleus of the hypothalamus (PVH; **b**), and arcuate nucleus (ARH; **c**) of wild-type
mice stained with DAPI and immunostained for GFRAL. **d**, Representative images of the AP and NTS
of control (Ctrl) and Crif1[BKO] mice stained using DAPI and immunostained for GFRAL and c-Fos. **e**, c-
Fos expression in GFRAL-expressing cells and non-GFRAL-expressing neurons in the AP. **f**,
Representative images of the intermediolateral nucleus region of the thoracic spinal cords of Ctrl and
Crif1[BKO] mice stained with DAPI and immunostained for ChAT and c-Fos. **g**, c-Fos expression in
ChAT-expressing cells. **h**, Schematic model of retrograde neuronal tracing study using a
pseudorabies virus that express EGFP under CMV promoter (PRV-CMV-EGFP). AP, area postrema;
NTS, nucleus tractus solitarius; IML, intermediolateral nucleus; WAT, white adipose tissue. **i**,

Representative images of the intermediolateral nucleus region of the spinal cord of ChAT-IRES-
Cre:tdTomato mice, 96 h after injection of enhanced green fluorescent protein (EGFP)-expressing
pseudorabies virus into the inguinal WAT, stained with DAPI, and immunostained for EGFP and
ChAT. **j**, Representative images of the AP and NTS of wild-type mice 120 h after injection of EGFP-
expressing pseudorabies virus into the inguinal WAT, stained with DAPI and immunostained for
EGFP and GFRAL. **k**, GDF15 concentrations in the cerebrospinal fluid of Ctrl and Crif1[BKO] mice. **i**,
Correlation between the serum and cerebrospinal fluid GDF15 concentrations in Crif1[BKO] mice. *, $P <$
0.05; ***, $P < 0.001$ *vs.* Ctrl by Student's t-test. Data are represented as mean ± SEM. See also
Extended Data Figure 7.
902

Figure 7. *Gfral* antisense oligonucleotides (ASOs) prevent GDF15-mediated phenotypic changes in Crif1[BKO] mice. **a**, Design and sequence of ASO targeting *Gfral* mRNA (*Gfral* ASO). The *Gfral* gene in mice include nine exons, and the *Gfral* ASO targets 22 nucleotides within exon 5. **b**, Representative immunoblots for GFRAL 1 week after treatment with lipid-based carriers (lipo), non-targeting control ASOs (Ctrl ASOs), and *Gfral* ASOs. **c**, Intracerebroventricular (ICV) injection of *Gfral* ASO to Crif1[BKO] mouse. **d,e**, *Gfral* mRNA expression 2 week post-injection with PBS, Ctrl ASO, and *Gfral* ASO (**d**), and relative expression of *Gfral*+ neurons in the AP and NTS (**e**). In (**e**), the number of *Gfral*+ neurons were normalized to those in PBS-injected mice. **f–i**, Body weight (**f**) and food intake (**g**) and energy expenditure (EE; **h** and **i**) in Crif1[BKO] mice, post-injection of Ctrl ASOs and *Gfral* ASOs. **j**, UCP1 expression in inguinal white adipose tissue (iWAT) in mice housed at room temperature. WT: wild type **k**, Representative images of UCP1-immunostained sections of inguinal WAT. Bar, 100 μm. **l**, Schematic model of the consequences of mitochondrial unfolded protein response (UPR[mt]) activation and *Gdf15* expression in brown adipose tissue (BAT) on systemic energy homeostasis. *, $P < 0.05$; **, $P < 0.01$; vs. PBS or Ctrl-ASO by Student's t-test. Data are represented as mean ± SEM.