Comprehensive study of subfamily evolution through sequences of conserved nuclear genes

and whole chloroplast genomes extracted from the same raw read data

Eranga Pawani Witharana[1,2,3, *], Takaya Iwasaki[4], Myat Htoo San[2,5], Nadeeka U. Jayawardana[1,6],

Nobuhiro Kotoda[3,5,7], Masashi Yamamoto[5,8], Yukio Nagano[2,3,5, *]

[1]*Faculty of Agriculture, University of Peradeniya, Peradeniya, Sri Lanka*

[2]*Analytical Research Center for Experimental Sciences, Saga University, Saga, Japan*

[3]*Graduate School of Advanced Health Science, Saga University, Saga, Japan*

[4]*Faculty of Science, Ochanomizu University, Tokyo, Japan*

[5]*The United Graduate School of Agricultural Sciences, Kagoshima University, Kagoshima, Japan*

[6]*Applied BioSciences,205B, Culloden Road, Macquarie University, Sydney, NSW, Australia*

[7]*Faculty of Agriculture, Saga University, Saga, Japan*

[8]*Faculty of Agriculture, Kagoshima University, Kagoshima, Japan*

*Corresponding author:

[✉]E-mail: erangav8@gmail.com

[✉]E-mail: nagano@cc.saga-u.ac.jp

Abstract

The chloroplast (cp) genome is a widely used tool for exploring plant evolutionary relationships, yet

its effectiveness in fully resolving these relationships remains uncertain. Integrating cp genome data

with nuclear DNA information offers a more comprehensive view but often requires separate

datasets. Nuclear DNA analyses are hindered by the need for high-quality sequences and high costs.

In response, we employed the same raw read sequencing data to construct cp genome-based trees

and nuclear DNA phylogenetic trees using Read2Tree, a cost-efficient method for extracting

conserved nuclear gene sequences from raw read data, focusing on the Aurantioideae subfamily,

which includes *Citrus* and its relatives. The resulting nuclear DNA trees were consistent with

existing nuclear evolutionary relationships derived from high-throughput sequencing, but diverged

from cp genome-based trees. To elucidate the underlying complex evolutionary processes causing

these discordances, we implemented an integrative workflow that utilized multiple alignments of

each gene generated by Read2Tree, in conjunction with other phylogenomic methods. Our analysis

revealed that incomplete lineage sorting predominantly drives these discordances, while

introgression and ancient introgression also contribute to topological discrepancies within certain

clades. This study underscores the cost-effectiveness of using the same raw sequencing data for both

cp and nuclear DNA analyses in understanding plant evolutionary relationships.

Keywords: Chloroplast genome sequences, Read2Tree, Aurantioideae subfamily, Phylogenetic

discordance, Incomplete lineage sorting, Introgression

Introduction

Phylogenetic analysis plays a crucial role in comprehending organism evolution and taxonomy. Maximizing the utilization of variable sites during the construction of a phylogenetic tree enhances accuracy[1,2]. The advent of high-throughput sequencing has vastly increased the number of available variable sites. In recent years, chloroplast (cp) genome sequences have gained popularity in plant molecular systematics[3,4,5]. Characterized by typical maternal inheritance, stable gene content, low recombination, and manageable size, the cp genome has emerged as a cost-effective tool for phylogenetic studies[6,7,8]. Assembling the complete cp genome, typically ranging from 120 to 170 kbp, from as little as 1 Gbp of raw sequencing data enables detailed analysis of evolutionary relationships among plant species[9,10]. However, sole reliance on cp genome sequences in plant phylogenetics has its limitations. These include potential instances of biparental or paternal inheritance of cpDNA in some species[6,11], and the phenomenon of 'chloroplast capture,' where cpDNA is transferred between different lineages through hybridization[12]. Additionally, because cpDNA represents a single genetic lineage inherited separately from the nuclear genome, it is subject to independent genetic drift and chloroplast capture that do not affect the nuclear genome in the same way. These factors can lead to differences in the phylogenetic signals from cpDNA and nuclear DNA, resulting in cytonuclear discordance[13,14,15]. Given these limitations, integrating cpDNA data with other molecular markers, particularly nuclear DNA, can provide a more comprehensive approach to phylogenetic reconstruction.

Nuclear DNA markers, owing to their biparental inheritance and greater genetic diversity, offer a broader and more accurate understanding of evolutionary relationships compared to cpDNA[16,17]. However, the phylogenetic analysis of many plants encounters challenges due to the limited availability of high-quality nuclear sequences[18,19]. This scarcity results from the high costs and technical difficulties associated with sequencing and annotation[20]. To address these limitations, transcriptomic data are increasingly being employed in phylogenetic reconstructions[21,22]. However, variations in gene expression among species, dependent on tissue type, can affect the accuracy of these reconstructions, potentially leading to skewed interpretations of phylogenetic signals[23,24].

The Read2Tree tool may offer a promising alternative for overcoming the limitations associated

with nuclear molecular markers in reconstructing phylogenies[25]. This method allows for the direct

extraction of conserved nuclear gene sequences from raw sequencing reads, effectively bypassing

the conventional steps such as genome assembly and annotation. Additionally, the potential to

employ the same short-read sequence data for both whole cp genome analysis and Read2Tree offers

a promising approach for phylogenetic reconstructions. Read2Tree produces two primary outputs:

concatenated alignment data for tree inference and multiple alignments for each gene across the

analyzed species. The concatenated alignment data is a primary focus of Read2Tree's publication,

which has been employed in reconstructing the phylogeny of yeast and various coronaviruses[25]. Its

effectiveness was confirmed by validating the reconstructed yeast phylogeny through comparison

with results from a previous study based on whole genome assemblies[25]. Additionally, the versatility

of Read2Tree was demonstrated by the alignment of the reconstructed coronavirus phylogeny with

lineage classifications from UniProt reference proteomes[25]. However, only one study has applied

Read2Tree to plant phylogenetics[26], and it did not address the validation of Read2Tree results as

comprehensively as the original publication[25]. The application of Read2Tree in plant phylogenetics

is particularly significant because plants are more prone to interspecies hybridization compared to

animals[27,28]. This frequent hybridization can lead to complex evolutionary histories, including

phenomena like chloroplast capture, which are challenging to resolve using chloroplast DNA alone.

Therefore, employing phylogenetic analyses based on many nuclear genes becomes crucial for

accurately reconstructing plant evolutionary relationships. Given this context, the effectiveness of

Read2Tree in plant phylogenetics remains relatively underexplored and requires further validation

through comprehensive comparisons with existing nuclear-based phylogenetic findings.

The second output type, multiple alignments for each gene, may hold significant potential for

analyzing detailed genetic differentiation processes as a downstream application of Read2Tree.

However, attempts to use Read2Tree for this purpose have not been comprehensively explored in

previous studies [25,26]. The genetic differentiation processes such as incomplete lineage sorting (ILS),

introgression, and ancient introgression often generate discordances between gene trees and species

trees, driving significant interest in understanding the underlying sources of these

discrepancies[29,30,31,32,33,34]. Consequently, constructing gene trees from multiple gene alignments can

facilitate species tree construction while also identifying conflicting phylogenetic signals. Integrating

Read2Tree with other phylogenomic methods can address these processes more effectively.

Additionally, this integration can pinpoint their contributions to cytonuclear discordance.

This study aims to elucidate and harness the potential of Read2Tree in advancing phylogenetic

reconstructions and discordance analysis, focusing on the phylogenetic relationships within

Aurantioideae, a subfamily of Rutaceae that encompasses *Citrus* and its relatives. We selected

Aurantioideae as the ideal model for this investigation for two primary reasons. First, our previous

research successfully utilized nuclear DNA sequences for this subfamily based on RAD-Seq data[35].

While RAD-Seq focuses on nuclear DNA near specific restriction sites, Read2Tree specializes in

analyzing a broader set of conserved nuclear gene sequences. This allows for a more comprehensive

evaluation of nuclear gene sequences. By comparing the nuclear gene trees generated by Read2Tree

with RAD-Seq results, we can more effectively evaluate the phylogenetic relationships and identify

potential sources of discordance within the subfamily. Second, the genus *Citrus*, along with other

members of Aurantioideae, holds global importance due to its nutritional value, high vitamin C

content, and widespread culinary use. Understanding the phylogenetic relationships within this

group has implications not only for evolutionary biology but also for agriculture and conservation

efforts. Many Aurantioideae species are valued for their edible and medicinal properties across

various regions, making accurate phylogenetic reconstructions crucial for biodiversity preservation

and resource management[36,37].

In this study, we employed the Read2Tree method to conduct nuclear-based phylogenetic

reconstructions of approximately 40 plant species within the subfamily Aurantioideae. We integrated

the nuclear phylogenies with whole cp genomes analyses to explore evolutionary relationships more

precisely and to investigate the potential causes of cytonuclear discordance. By comparing the

nuclear phylogenetic trees generated by Read2Tree with those obtained from previous RAD-Seq

analyses, we aimed to validate the robustness and utility in plant phylogenetics. Additionally, we

developed a comprehensive workflow that integrates Read2Tree with other phylogenomic methods

to analyze both species-gene tree discordance and cytonuclear discordance. This workflow enabled

us to assess whether the observed discordances were due to factors such as ILS, introgression, or

ancient introgression followed by ILS. By identifying the sources of these discordances, we aimed to

gain insights into the complex evolutionary history of Aurantioideae and the potential mechanisms

driving cytonuclear discordance in plant phylogenetics.

Results

The phylogenetic tree constructed using Read2Tree demonstrates strong concordance with the tree derived from RAD-Seq data

The samples utilized in this investigation are detailed in Supplementary Table S1. Initially, Read2Tree facilitated the retrieval of sequences from various conserved nuclear genes. To establish the root of the phylogenetic tree within the subfamily Aurantioideae, a Maximum Likelihood (ML) tree was generated for 43 species of the Rutaceae family (Supplementary Fig. S1). Subsequently, we focused on 39 Aurantioideae species to generate a ML phylogenetic tree using the conserved nuclear gene sequences (Fig.1), with rooting determined from the Rutaceae analysis. The ML phylogenetic trees largely aligned with findings from a prior RAD-Seq based study[35], with noteworthy exceptions observed for *Micromelum minutum* and *Murraya koenigii*. Their phylogenetic positions, previously uncertain in the prior study[35], remained unresolved in our current analysis, as evident from the comparison between Fig.1 and Supplementary Fig. S1. An in-depth analysis within the genus *Citrus* was not conducted using Read2Tree due to the presence of recombination events occurring during hybridization[38].

Phylogenetic analysis employed whole cp genome sequences revealing disparities when compared with nuclear gene-based trees

Initially, a cp phylogenetic tree encompassing 59 members of the Rutaceae family was constructed, to determine the root position for subsequent analyses (Supplementary Fig. S2). Subsequently, cp phylogenetic trees were developed based on DNA sequences for the same 39 members of the Aurantioideae subfamily used in Read2Tree, utilizing ML and Bayesian Inference (BI) methods (Supplementary Fig. S3 and Supplementary Fig. S4, respectively). The inclusion of identical taxa in both the whole cp genome and nuclear analyses was a deliberate approach to minimize discrepancies arising from taxon sampling, thereby reducing variations associated with

lineage-specific evolutionary histories. The topologies of these cp phylogenetic trees remained

consistent across the two analyses.

Comparison of the topologies of the two phylogenetic trees, one from whole cp genome

sequences and the other from the concatenated nuclear gene tree analysis (Fig. 2A), revealed two

primary groups within the Aurantioideae subfamily, referred to as group A and group B. However,

this comparison detected cytonuclear discordances within each group (Fig. 2A). Inconsistencies

within the *Citrus*, *Atalantia*, and *Clausena* genera were noted due to interspecific hybridization

within the genus. Moreover, marked differences were observed in the intergeneric relationships. For

instance, the cpDNA-based analysis positioned the *Limonia*/*Feroniella* subgroup as one of the

closest outgroups to *Citrus*, contrary to the nuclear DNA analysis, which identified this subgroup as

the nearest outgroup to a clade comprising *Citrus, Atalantia, Citropsis*, and *Naringi*. Furthermore,

the cpDNA analysis grouped *Swinglea* with *Citropsis* and *Naringi*, while nuclear DNA analysis

aligned *Swinglea* with the *Limonia*/*Feroniella* subgroup. Discrepancies between the two analyses

were also evident in the phylogenetic structures formed by *Paramignya*, *Triphasia*, and

Aegle/*Afraegle*. The phylogenetic positions of *Micromelum minutum* and *Murraya koenigii* remained

ambiguous.

Integrating Read2Tree for species tree estimation and revealing phylogenetic discordances

The detection of cytonuclear discordances strongly suggests the presence of broader phylogenetic

discordances. By integrating Read2Tree with other phylogenomic methods, these discordances were

determined. Orthologous groups (OGs) consisting of genes or DNA sequences in different species

that diverged from a common ancestral gene through speciation, typically maintain similar functions

across species, making them critical for studying evolutionary relationships[39]. In this analysis,

Read2Tree generated multiple sequence alignments for each OG, which were used for constructing

each gene tree. Specifically, alignments from 579 OGs with a ratio of N characters to total

nucleotides below 10%, as identified through Read2Tree analyses, were selected for gene trees

constructions. The gene trees were subsequently used to infer the ASTRAL species tree, as shown in

Fig. 2B (left panel), Fig. 3A (right panel), and detailed in Supplementary Table S3. The resulting

species tree has a normalized quartet score (QT) of 0.7, indicating that 70% of the quartets in the

gene trees are concordant with the species tree. This score reflects a high level of overall

concordance. Additionally, the grouping into groups A and B in the ASTRAL species tree aligns

with those in the cp phylogeny and concatenated nuclear phylogeny as demonstrated in the Fig.2B

and Fig.3A.

When comparing the species tree to the concatenated nuclear tree constructed using Read2Tree,

notable discrepancies are observed within the *Citrus* genus and in the intergeneric relationships

involving *Swinglea* and *Murraya koenigii* (Fig. 3A). In the species tree, *Swinglea* is placed as an

outgroup to *Citrus*, *Atalantia*, and *Citropsis/Naringi*, whereas in the concatenated tree, *Swinglea*

clusters with *Limonia* and *Feroniella*. Additionally, *Murraya koenigii* acts as a near outgroup

exclusively to *Clausena* species in the species tree, while in the concatenated tree, it serves as an

outgroup to both *Clausena* and *Glycosmis* species. This ambiguous positioning of *Murraya koenigii*

mirrors the unresolved status observed in previous RAD-Seq studies as well as in our Read2Tree and

cp genome trees (Fig.2A).

PhyParts conflict analysis was performed to investigate the complex discordances between gene

trees and the species tree, with *Boenninghausenia albiflora* used as the outgroup (Fig. 3B). The

analysis revealed that 511 out of 579 gene trees were concordant with the topology supporting the

presence of two primary groups, groups A and B, within the subfamily Aurantioideae. However,

significant conflict was observed among informative gene trees at the node encompassing the *Citrus*,

Atalantia, Citropsis/Naringi, *Swinglea*, *Paramignya/Triphasia*, *Murraya koenigii*, and *Micromelum*

minutum.

As mentioned above, cytonuclear discordance was observed between the concatenated nuclear

and cp trees (Figure 2A). Additionally, significant cytonuclear discordances were evident between

the species tree and cp tree, particularly within the *Citrus*, *Atalantia*, and *Clausena* genera, as well as

in intergeneric relationships (Fig.2B). The species tree positions the *Limonia/Feroniella* subgroup as

an outgroup to a clade containing *Citrus*, *Atalantia, Citropsis/Naringi*, and *Swinglea*, while the cp

tree places this subgroup as the closest outgroup to *Citrus* and *Atalantia*. Furthermore, in the species

tree, *Swinglea* is positioned as an outgroup to *Citrus, Atalantia, Citropsis*, and *Naringi*, but the cp

tree groups *Swinglea* with *Citropsis/Naringi*. Discrepancies between the species tree and cp tree are

also apparent in the phylogenetic structures of *Paramignya, Triphasia*, and *Aegle/Afraegle*.

Additionally, in the ASTRAL species tree, *Micromelum minutum* is positioned as an outgroup to

both *Clausena* and *Glycosmis*, whereas in the cp tree, it appears as a near outgroup exclusively to

Glycosmis.

Consistent subgrouping patterns unveiled by nuclear and cp trees

Both nuclear and cp analyses demonstrated similar grouping and subgrouping patterns, as

depicted in Figures 2 and 3. The phylogenies derived from both nuclear and cp data consistently

revealed two primary groups, designated as group A and group B, within the subfamily

Aurantioideae (Figs. 1, 2, and 3). Historically, Aurantioideae has been classified into the Citreae

and Clauseneae tribes[40]. However, our analysis led to a reassignment of species within these tribes.

Specifically, *Murraya caloxylon* and *Murraya paniculata*, historically classified within the

Clauseneae tribe, have been reassigned to the Citreae tribe, now defined as group A. The remaining

species historically classified under the Clauseneae tribe have been consolidated into group B, which

corresponds to the newly defined Clauseneae tribe in this study (Fig.2 and Fig.3).

Within group A, six distinct subgroups were identified. The "True citrus fruit trees" subgroup

comprises *Citrus* species, consistent with traditional taxonomy[40]. The "Yaki Naran" subgroup

encompasses all *Atalantia* species, inspired by Ayurvedic medicine in Sri Lanka including *Atalantia*

ceylanica. The "Thanaka" subgroup, named after the traditional cosmetic from Myanmar, comprises

Citropsis and *Naringi* species. The "Wood apple" subgroup includes *Limonia* and *Feroniella* species,

while the "Bael fruit" subgroup comprises *Aegle* and *Afraegle,* reflecting their historical names[40].

The final subgroup, the *Murraya* subgroup, consists of *Murraya caloxylon* and *Murraya paniculata*.

However, three species *Swinglea glutinosa*, *Paramignya lobata*, and *Triphasia trifolia*, did not fit

neatly into any defined subgroup.

Signatures of ILS, introgression, and ancient introgression

Discordances were detected among three main subgroups: (1) True citrus fruit trees and its near

subgroups, (2) the Bael fruit subgroup and its associated species, and (3) Group B (Fig.4). These

discordances underscore the significant influence of genetic differentiation processes, such as ILS,

introgression, and ancient introgression. To assess the impact of these processes on the discordance

patterns observed in these subgroups, one representative species from each subgroup was selected

for further analysis. To elucidate whether ILS or introgression was responsible for the observed

phylogenetic discordances, Quantifying Introgression via Branch Lengths (QuIBL) was initially

employed[41]. This method evaluates triplet topologies from gene trees by comparing branch length

distributions under two models: one accounting for ILS alone and the other incorporating both ILS

and introgression. Model selection was guided by Bayesian Information Criterion (BIC) values, with

ΔBIC used to differentiate between ILS and introgression. Specifically, a ΔBIC greater than 10

indicates ILS, a value lower than -10 suggests introgression, and values within this range imply that

ILS and introgression cannot be clearly distinguished. To further assess introgression, Patterson's D

statistic[42] was applied (P < 0.05), while D_{FOIL} was employed to detect ancient introgression events (P

< 0.05)[43].

In the True citrus fruit trees and its near subgroups, we assessed 860 gene trees, focusing on 30

triplets. Among these, 63.33% (19 of 30) indicated ILS (ΔBIC > 10), 10% (3 of 30) supported

introgression (ΔBIC < -10), and 26.67% were inconclusive (ΔBIC between -10 and 10). This

suggests that ILS was the predominant cause of phylogenetic conflict (Fig.5A and Supplementary

Table S4). Specifically, the discordant topology involving wood apple, *Swinglea glutinosa*, and

Citropsis gabunensis (used as an outgroup) was primarily attributed to introgression (ΔBIC < -10).

Patterson's D statistic and D_{FOIL} analyses indicated that ancient introgression was the main factor,

followed by ILS (Supplementary Table S5). Additionally, QuIBL and Patterson's D statistic

identified introgression events between True citrus fruit trees and Yaki Naran, as well as between

Yaki Naran and wood apple (Supplementary Table S4 and Supplementary Table S6). Furthermore,

PhyloNet analyses provided evidence of introgression events within True citrus fruit trees and its

near subgroups (Fig.5B and Supplementary Fig.S5).

In the analysis of the Bael fruit subgroup and its associates, based on 1964 gene trees, ILS was

identified as the primary cause of discordance (66.67%). However, triplet analysis involving

Paramignya lobata, Triphasia trifolia, and *Aegle marmelos* (used as an outgroup) yielded ΔBIC

values between -10 and 10, suggesting the potential presence of either ILS or introgression (Fig.5A,

and Supplementary Table S7). Further clarification from Patterson's D statistic and PhyloNet

analyses indicated that the observed inconsistencies were due to introgression between *Paramignya*

lobata and *Triphasia trifolia* (Fig.4C, Supplementary Fig.S6 and Supplementary Table S8).

Within the Group B, based on 1135 gene trees, 58.33% of the triplets provided evidence for ILS

(7 of 12 triplets, ΔBIC > 10), 16.67% indicated introgression (2 of 12 triplets, ΔBIC < -10), and 25%

were ambiguous (ΔBIC between -10 and 10). This suggests that ILS was the predominant cause of

phylogenetic discordance. Discordant topologies involving *Glycosmis citrifolia, Murraya koenigii*,

and *Clausena anisata* were mainly attributed to ILS (Supplementary Table S11, Supplementary

Table S12 and Supplementary Table S13). Additionally, D_{FOIL} analysis suggested ancient

introgression involving *Glycosmis citrifolia* and *Micromelum minutum* with other taxa (Fig.4A).

Consequently, the inconsistencies within this group were primarily attributed to ancient introgression,

followed by ILS (Supplementary Table S11 and Supplementary Table S13).

The source of cytonuclear discordance was inferred using QuIBL and D-statistics results. The

QuIBL analysis suggested that ILS is the primary driver of the observed cytonuclear discordance

within True citrus fruit trees and its near subgroups, as well as the Bael fruit subgroup and its

associates (Supplementary Table S4, Supplementary Table S7, and Supplementary Table S9).

Additionally, both QuIBL and D_{FOIL} analyses revealed that the discordant topologies involving

Micromelum minutum and *Murraya koenigii* are predominantly due to ancient introgression,

followed by ILS (Supplementary Table S11 and Supplementary Table S13).

Introgression events between Asian and African Plants

The generated phylogenetic trees revealed two instances of close relationships between Asian and African plants in the Thanaka and Bael fruit subgroups (Fig.2 and Fig.3). Within the Thanaka subgroup, *Citropsis* is native to Africa, while *Naringi* originates from Asia. Similarly, within the Bael fruit subgroup, *Aegle* is indigenous to Asia, and *Afraegle* is native to Africa. To investigate the discordance in topologies involving *Citropsis* and *Naringi*, we analyzed a total of 1,084 gene trees. Similarly, we evaluated 1,080 gene trees to explore the discordance in topologies involving *Afraegle* and *Aegle*. Our QuIBL and Patterson's D-statistic analyses revealed compelling evidence of introgression events between these Asian and African plant lineages (Fig.5A, Supplementary Table S12 and Supplementary Table S13). Specifically, the introgression event within the Bael fruit subgroup occurred over 1 to 2 coalescent time scales, while the introgression event in the Thanaka subgroup occurred over 0.5 to 2 coalescent time scales (Supplementary Table S12).

Variability in structural variations and pseudogenization across phylogenetic tree branches using publicly available whole cp genome sequences

Due to the lack of publicly available raw sequencing data, an in-depth phylogenetic analysis using Read2Tree was not feasible. Instead, we focused on analyzing publicly accessible whole cp genome sequences to assess phylogenetic relationships within the Aurantioideae subfamily. We constructed a phylogenetic tree for 45 Aurantioideae members using both ML and BI methods (Supplementary Figs. S7 and S8), with congruent topologies across analyses. These cp phylogenetic trees reinforced the close relationships between Asian and African plants within the Thanaka and Bael subgroups and allowed us to estimate divergence times, suggesting that these dispersal events occurred around five million years ago (Supplementary Fig. S9). However, our study did not resolve the unique biogeographical distribution of *Clausena anisata*, the sole *Clausena* species found in both Asia and Africa, due to inconsistencies in its interrelationships across different phylogenetic trees.

Given the little or absence of recombination in hybridization processes affecting cpDNA[11], the

construction of phylogenetic trees for the *Citrus* genus using whole cp genome sequences was

considered valuable. Consequently, phylogenetic trees for 56 *Citrus* species were generated

(Supplementary Fig. S10 and Supplementary Fig. S11). We identified eight clusters within the

Citrus genus based on whole cp genome sequences. The cpDNAs of the two wild species, *Citrus*

latipes, and *Citrus hongheensis*, were closely related to those of the pummelo cluster.

We identified a ~20,000 bp inversion in the *psbM* to *trnG-UCC* region of *Atalantia ceylanica*,

Atalantia roxburghiana, and *Naringi crenulata*, consistently occurring at the same nucleotide

position (Supplementary Fig. S12 and Supplementary Fig. S13). This suggests that similar cpDNA

structural variations can arise at different phylogenetic nodes. Notably, 53 bp inverted repeats at both

ends of this region suggest their involvement in these inversions. These repeats were found in most

Aurantioideae, except in *Citrus*, *Feroniella, Triphasia*, and *Aegle.* Additionally, pseudogenes were

identified through frameshifts, premature stop codons, or missing start codons. The

pseudogenization of *infA* occurred in all species, with *rpl22* and *rps19* present in various

configurations, along with *ycf68*, *orf56*, *cemA*, and *ycf15* (Supplementary Fig. S14, and

Supplementary Fig. S15). Furthermore, in *Atalantia kwangtungenensis,* the pseudogenization of the

accD gene was confirmed[44], and a new pseudogenization event was identified in the *ycf1* gene

(Supplementary Fig. S16).

Discussion

In this study, we leveraged an accessible dataset to analyze a large number of nuclear gene sequences alongside whole cp genome sequences for phylogenetic reconstructions. Rather than relying on a single method, the integration of both nuclear and cp genome sequences offers a more comprehensive and informative perspective in plant phylogenetics[14,15,45,46,47]. Unlike previous studies that required separate datasets for nuclear and cp genome sequences[45,46,47], our method is cost-efficient as it uses the same short-read sequence data for both analyses. This not only reduces costs but also enhances research efficiency, demonstrating its validity as a robust tool for plant phylogenetic reconstructions.

Our analysis revealed discrepancies between phylogenetic trees derived from nuclear gene sequences and those based on whole cp genome sequences. Despite these differences, all trees consistently identified two primary groups with similar subgrouping patterns. These results underscore the need for reclassification within the Aurantioideae subfamily to better align with the evolutionary relationships revealed by both nuclear and cp genome data. In response, we propose a reclassification that refines the taxonomy of Aurantioideae, providing a more accurate reflection of the subfamily's genetic and evolutionary relationships.

The Read2Tree method extends its utility beyond core phylogenetic reconstruction by addressing phylogenetic discordance. It facilitates the direct acquisition of multiple alignments for each OG, crucial for constructing individual gene trees. Integrating Read2Tree with other phylogenomic tools allows for the effective identification of genetic differentiation processes such as ILS, introgression, and ancient introgression that contribute to discordances. These evolutionary processes pose challenges to phylogenomics by complicating the inference of accurate evolutionary relationships[13,31,32,33,34]. ILS, often identified as a primary cause of gene tree discordance, arises when ancestral polymorphisms persist through successive speciation events, especially during rapid radiations with short intervals between these events[31,32,33,34,48,49]. This persistence can lead to discordant gene tree topologies. Introgression, which involves the transfer of genetic material

between species through hybridization, can blur species boundaries and distort gene trees[29,32,33,50,51].

Ancient introgression, which occurs between species that have diverged significantly, can

incorporate historical gene flow into the phylogenetic signal[29,30,31,32]. Our analysis of nuclear gene-

derived trees indicates that ILS is the primary driver of discordance, rather than introgression. The

presence of ILS suggests that the Aurantioideae subfamily has experienced rapid diversification,

which has obscured the true phylogenetic relationships and complicating the reconstruction of its

evolutionary history.

Our discordance analysis concentrated on three key groups: True citrus fruit trees and its near

subgroups, the Bael fruit subgroup and its associates, and group B. We found that some of the

discordant topologies in these areas were primarily attributed to introgression and ancient

introgression, followed by ILS. These introgressive events likely facilitated gene flow between

distinct lineages, resulting in discrepancies between gene trees and the species tree. Additionally,

ILS preserves ancestral genetic polymorphisms, further complicating the reconstruction of accurate

phylogenetic histories[32,33,48,51,52]. In the context of True citrus fruit trees and its near subgroups, our

results align with previous studies that reported hybridization between *Citrus* and intergeneric

species, emphasizing the substantial impact of introgression on shaping phylogenetic relationships

within this group[53,54]. Prior research also highlights how hybridization frequently leads to

incongruence among gene trees, underscoring its importance in accurately inferring the phylogeny of

Citrus species[55].

Topological conflicts between nuclear and organellar phylogenies, known as cytonuclear

discordance, are common in phylogenetic studies[14,15,45,46,47,56,57]. In our investigation of the

Aurantioideae subfamily, we found that ILS is often a more likely cause of cytonuclear discordance

than introgression. Rapid speciation events within Aurantioideae may have occurred with

insufficient time for gene lineages to coalesce, resulting in discordant gene trees between nuclear

and organellar genomes[58]. Additionally, large effective population sizes can also extend the

persistence of multiple gene lineages, increasing the likelihood of discordant phylogenies[58,59]. While

historical hybridization and introgression introduce additional genetic material and create a mosaic

of gene trees[60,61,62], our findings suggest that ILS plays a primary role in the observed cytonuclear

discordance, underscoring the complex evolutionary history of the Aurantioideae subfamily.

Structural variations and pseudogenization events provide valuable insights into the evolutionary

history and genetic diversity within a genus but have limited utility for intergeneric classification.

Variations in genes such as *rpl22*, *ycf68*, and *orf56* have appeared independently of their

phylogenetic positions, likely due to hybridization[63] or ILS[15]. Our analysis highlighted significant

patterns of pseudogenization, including the co-pseudogenization of *accD* and *ycf1*, which supports

the co-evolution hypothesis[64,65,66]. Additionally, the pseudogenization of the *cemA* gene suggests

functional compensation by the nuclear-encoded *DLDG1* or a transfer to the nuclear genome with a

transit peptide sequence[67,68]. Independent structural changes, such as inversions in the *psbM* to *trnG-*

UCC region, may arise from similar mechanisms across different lineages, complicating

classification efforts. These findings emphasize the necessity for a multifaceted approach in plant

phylogenetics, rather than relying solely on structural variations or pseudogenization patterns.

In conclusion, our study highlights the effectiveness of the Read2Tree method in plant

phylogenetics, showcasing its utility not only in reconstructing phylogenies but also in addressing

phylogenetic discordance. By integrating Read2Tree with other phylogenomic approaches, we were

able to identify key genetic differentiation processes, including ILS, introgression, and ancient

introgression, which contribute to phylogenetic conflicts. This sheds light on the method's robustness

and versatility, advancing our understanding of complex evolutionary relationships and offering a

valuable tool for future research. Additionally, our findings emphasize the critical importance of

complete data submission for ensuring the transparency, reproducibility, and robustness of

phylogenetic analyses. While many studies have submitted only assembled whole cp genome

sequences to public databases, the inclusion of the raw sequence data used in the assembly process is

essential, especially for methods like Read2Tree that rely on these data. Ensuring the availability of

raw data allows for more comprehensive analyses, facilitating validation, replication, and deeper

insights into evolutionary relationships.

Methods

DNA sequence data collection and processing

The DNA used in this study was either sourced from the same batch used in a previous

experiment[35] or purified using the method described therein. DNA quality was assessed by

electrophoresis on a 1% agarose gel, and DNA concentration was measured using a DNA BR Assay

Kit (Thermo Fisher Scientific, USA). Sequencing libraries were prepared from total DNA samples

by Novogene (Singapore), using the NEBNext Ultra DNA Library Prep Kit for Illumina (NEB,

USA). These libraries were then sequenced to produce 150-bp paired-end reads using a NovaSeq

6000 instrument (Illumina, USA) by Novogene. Additional DNA sequence data were acquired from

public databases, with accession numbers detailed in Supplementary Table S1. For preprocessing all

DNA sequence data, FastP (version 0.23.2) was employed with its default settings[69].

Concatenated phylogenetic analysis based on nuclear gene sequences

For the concatenated phylogenetic analysis, we used Read2Tree (version 0.1.5)[25]. Marker genes

for *Citrus clementina*, *Eucalyptus grandis*, all five species of Malvaceae, and all seven species of

Brassicaceae were obtained from the OMA browser (https://omabrowser.org/oma/export_markers).

The OMA browser parameters were set with a minimum fraction of covered species at 0.8 and a

maximum number of markers at –1. In total, 16,258 OGs were identified among the selected 14

species in the OMA browser. Read2Tree was utilized to generate alignment files using these marker

genes. To ensure the specificity of the phylogenetic trees, the 14 species obtained from the OMA

browser were excluded from the resulting alignment files. Aligned data were trimmed using trimAl

(v1.4. rev15) with the – automated1 option[70]. ModelTest-NG (version 0.1.7) was used to identify the

optimal evolutionary model[71]. Subsequently, Maximum Likelihood (ML) trees were generated using

RAxML (version 8.2.12) with 100 bootstrap replicates[72]. The visualization of the resulting trees was

achieved using Dendroscope (version 3.8.8)[73].

***De novo* assembly and annotation of cp genomes**

The *de novo* assembly of cp genomes was performed using GetOrganelle (version 1.7.5)[74]. The

parameters for this process were set to – R 15, – k 21,45,65,85,105, and – F embplant_pt. To

validate the assembly, we utilized Bowtie2 (version 2.4.4) for mapping[75]. The sorting of the mapped

data and the generation of bam files were conducted using Samtools (version 1.14)[76]. Assembly

quality was confirmed, and heteroplasmic sites were identified using the Integrated Genome

Viewer[77]. Despite detecting heteroplasmic sites, the assemblies obtained from GetOrganelle were

used for subsequent analysis.

For annotation of the assembled cp genomes, we used GeSeq[78]. Within the GeSeq framework,

protein-coding sequences (CDS) were predicted using Chloë (version 0.1.0)[79], tRNAs were

identified using tRNAscan-SE (version 2.0.7)[80], and rRNAs were predicted using HMMER[81].

Thereafter, the annotation results were curated manually. Circular genome maps of the annotated cp

genomes were generated using OGDRAW (version 1.3.1)[82].

Phylogenetic analysis based on cp genome sequences

For phylogenetic analysis of cp genome sequences, DNA sequence alignments were performed

using HomBlocks (version 1.0) with its default settings[83]. The optimal evolutionary model for the

alignments was identified using ModelTest-NG (version 0.1.7)[71]. Subsequently, ML trees were

constructed using RAxML (version 8.2.12), employing 1000 bootstrap replicates[72]. The visualization

of the resultant ML trees was accomplished using Dendroscope (version 3.8.8)[73].

For the construction of Bayesian Inference (BI) trees, BEAST (version 1.10.4) was used,

implementing the Markov Chain Monte Carlo (MCMC) method over a total of 10,000,000

generations[84]. Tree construction samples were collected every 10,000 generations, with the initial

10% of generations discarded as 'burn-in' to reduce potential bias from the early stages of the

MCMC run. The completed BI trees were visualized using FigTree (version 1.4.4)

(http://tree.bio.ed.ac.uk/software/figtree/).

Species tree inference from Read2Tree data

To construct the species tree, alignments for 16,284 OGs across all species were utilized from the

Read2Tree output file "06_align_merge_dna". The OG alignments were processed using trimAl

(version 1.4, revision 15)[70] with the -automated1 option and eliminate species present in the OMA

browser. The alignments were then assessed based on the ratio of missing data (N characters) to total

nucleotide characters, with a threshold set at 10%. From this, 579 OGs were selected for further

analysis (Supplementary Dataset 16). Model testing[71] was performed for each OG, and the ML gene

tree of each OG was constructed using RAxML (version 8.2.12) with 100 bootstrap replicates[72]. The

species tree was subsequently inferred using ASTRAL-III (version 5.7.8)[85].

Phylogenetic discordance analysis

PhyParts was utilized to evaluate the concordance and discordance between gene trees and the

species tree[86]. Gene trees were processed by collapsing branches with bootstrap support below 33%

and subsequently rerooted using *Boenninghausenia albiflora* as the outgroup. Similarly, the

ASTRAL species tree was rerooted with *B. albiflora*, and this rerooted tree served as the reference

for the conflict analysis. The PhyParts output was visualized on the ASTRAL topology using the

phypartspiecharts.py script available at github.com/mossmatters/phyloscripts. The analysis provided

metrics on the number of gene trees that were concordant or discordant with the species tree

topology at each branch, as well as those that were uninformative (i.e., gene trees with support

values below 33% at the given branch).

The discordance analysis was focused on three specific issues: (1) *Citrus* and its near subgroups

using 860 gene trees (Supplementary Dataset 17), (2) Bael fruit subgroup and its associated species

using 1,964 gene trees (Supplementary Dataset 18), and (3) Group B using 1,135 gene trees

(Supplementary Dataset 19), with one representative from each subgroup included. Gene trees for

these analyses were generated using similar criteria and procedures as those applied in species tree

estimation. Additionally, 1,084 gene trees were generated to assess discordance in topologies

involving *Citropsis* and *Naringi* (Supplementary Dataset 20)*,* while 1,080 gene trees were used to

examine discordance involving *Afraegle* and *Aegle* (Supplementary Dataset 21). QuIBL was

employed to distinguish between ILS and introgression as potential causes of the observed

discordances[41]. Model selection was guided by BIC values, with ΔBIC values interpreted as follows:

ΔBIC > 10 indicates ILS only, ΔBIC < -10 suggests introgression, and -10 < ΔBIC < 10 indicates

either ILS or introgression. Patterson's D-statistic analysis was performed using D-suite (version

0.4)[42] with the Dtrios option to identify introgression events ($P < 0.05$). Additionally, D_{FOIL}

statistics[43] were employed to detect ancient introgression events, with significance determined by $P <$

0.05.

To further investigate reticulation events, Maximum Pseudolikelihood (MPL) analysis in

PhyloNet (version 3.8.2)[87] was conducted. We performed 10 runs per search, each returning five

optimal networks. These networks were then optimized for branch lengths and inheritance

probabilities under full likelihood, using default settings and the -po option. The best-fitting network

was selected based on the Akaike Information Criterion (AIC)[88,89], with the number of parameters

(k) being the sum of branch lengths and reticulations, and L representing the likelihood value.

Detection of Structural Variations in Chloroplast Genomes

503   The process of identifying structural variations within the cp genomes involved a thorough visual

504   inspection of the circular cp genome map combined with an analysis of BLAST results[90]. To

505   visualize these identified structural variations, Easyfig (version 2.2.5) was employed[91].

Divergence Time Estimation

The ML phylogenetic tree for the Rutaceae family was constructed based on the aligned

sequences of cp genomes using MEGA11 (version 11.0.13)[92]. The optimal evolutionary model was

determined using a model test conducted within MEGA11. To generate the ML tree used in

divergence time estimation, the General Time Reversible (GTR) model incorporating gamma

distribution (G), and proportion of invariable sites (I) was employed. Additionally, complete deletion

parameters were used. The RelTime-ML option was utilized to produce the time tree by employing

the GTR+G+I model and complete deletion parameters. Calibration of the time tree was established

based on the divergence time between *Murraya paniculata* and *Murraya caloxylon* retrieved from

the TimeTree database[93]. The calibration parameters were defined using a Normal Distribution, with

a mean of 10, a standard deviation of 2, and a confidence interval ranging from 6.08 to 13.92 million

years ago.

Identification of Pseudogenes

Protein sequences were extracted from each annotation file using genbank_to (version 0.42)

(https://github.com/linsalrob/genbank_to). OrthoFinder (version 2.5.4) was used to extract all

combinations of orthologous proteins[94]. For accurate sequence alignment, MAFFT (version 7.515)

was used to align the amino acid sequences[95]. The aligned results were visualized using MEGA11

(version 11.0.13) for comparative analysis[92], which facilitated the identification of pseudogenes.

Pseudogenes were characterized by the presence of frameshift mutations, premature stop codons,

and the absence of start codons. These features were systematically compared to the intact gene

sequences, with additional support provided by translation mapping using the tool available at

https://www.bioinformatics.org/sms2/trans_map.html.

Use of generative AI

ChatGPT (https://chat.openai.com/) was used to rewrite the manuscript into more appropriate

English and proofread the English text. Subsequently, the authors verified the correctness of the

revisions.

References

1. Kaya, G., Ezekannagha, C., Heider, D. & Hattab, G., Context-aware phylogenetic trees for phylogeny-based taxonomy visualization. *Front. Plant Sci.* **13**, 891240; 10.3389/fgene.2022.891240 (2022).

2. Steenwyk, J.L. & King, N., The promise and pitfalls of synteny in phylogenomics. *PLoS Biol.* **22,** e3002632; 10.1371/journal.pbio.3002632 (2024).

3. Tian, Y. et al. Research progress in plant molecular systematics of Lauraceae. *Biology (Basel)* **10**, 391; 10.3390/biology10050391 (2021).

4. Jiang, D. et al. Complete chloroplast genomes provide insights into evolution and phylogeny of *Zingiber* (Zingiberaceae). *BMC Genom.* **24**, 30; 10.1186/s12864-023-09115-9 (2023).

5. Li, C.C. et al. Insights into chloroplast genome evolution in Rutaceae through population genomics. *Horticulture Advances* **2**, 13; 10.1007/s44281-024-00032-9 (2024).

6. Park, H.S. et al. Inheritance of chloroplast and mitochondrial genomes in cucumber revealed by four reciprocal F1 hybrid combinations. *Sci. Rep.* **11**, 2506; 10.1038/s41598-021-81988-w (2021).

7. Sun, J. et al. Evolutionary and phylogenetic aspects of the chloroplast genome of *Chaenomeles* species. *Sci. Rep.* **10,** 11466; 10.1038/s41598-020-67943-1 (2020).

8. Feng, J. et al. Analysis of complete chloroplast genome: Structure, phylogenetic relationships of *Galega orientalis* and evolutionary inference of Galegeae. *Genes (Basel)* **14**, 176; 10.3390/genes14010176 (2023).

9. Kahraman, K. & Lucas, S.J., Comparison of different annotation tools for characterization of the complete chloroplast genome of *Corylus avellana* cv Tombul. *BMC Genom.* **20**, 874; 10.1186/s12864-019-6253-5 (2019).

556   10.   Wang, Y. et al. Chloroplast genome variation and phylogenetic relationships of *Atractylodes*
557   species. *BMC Genom.* **22**, 103 10.1186/s12864-021-07394-8 (2021).

558   11.   Greiner, S., Sobanski, J. & Bock, R., Why are most organelle genomes transmitted maternally?
559   *Bioessays* **37**, 80–94 (2015).

560   12.   Rieseberg, L. H. & Soltis, D. E. Phylogenetic consequences of cytoplasmic gene flow in plants.
561   *Evol. Trend. Plant.* **5**, 65–84 (1991).

562   13.   Pamilo, P. & Nei, M. Relationships between gene trees and species trees. *Mol. Biol. Evol.* **5**,
563   568–583 (1988).

564   14.   Soltis, D.E. and Kuzoff, R.K., Discordance between nuclear and chloroplast phylogenies in the
565   Heuchera group (Saxifragaceae). *Evolution* **49**, 727–742 (1995).

566   15.   Thureborn, O., Wikström, N., Razafimandimbison, S.G. & Rydin, C., Plastid phylogenomics
567   and cytonuclear discordance in Rubioideae Rubiaceae. *PLoS One* **19**, e0302365;
568   10.1371/journal.pone.0302365 (2024).

569   16.   Krawczyk, K., Nobis, M., Nowak, A., Szczecińska, M. & Sawicki, J., Phylogenetic
570   implications of nuclear rRNA IGS variation in *Stipa* L. (Poaceae). *Sci. Rep*. **7**,11506;
571   10.1038/s41598-017-11804-x (2017).

572   17.   Wang, W., Li, H. & Chen, Z., Analysis of plastid and nuclear DNA data in plant
573   phylogenetics—evaluation and improvement. *Sci. China Life Sci.* **57**, 280–286 (2014).

574   18.   Lutz, K.A., Wang, W., Zdepski, A. & Michael, T.P., Isolation and analysis of high quality
575   nuclear DNA with reduced organellar DNA for plant genome sequencing and resequencing.
576   *BMC biotechnol.* **11**, 54; 10.1186/1472-6750-11-54 (2011).

577   19.   Amarasinghe, S.L. et al. Opportunities and challenges in long-read sequencing data analysis.
578   *Genome Biol.* **21**, 30; 10.1186/s13059-020-1935-5 (2020).

20. Mani, I., Current status and challenges of DNA sequencing. *Advances in synthetic biology*, 71–
80 (2020).

21. Liu, X. et al. Phylogenetic inference from single-cell RNA-seq data. *Sci. Rep.* **13**, 12854;
10.1038/s41598-023-39995-6 (2023).

22. Washburn, J.D. et al. Genome-guided phylo-transcriptomic methods and the nuclear
phylogenetic tree of the Paniceae grasses. *Sci. Rep.* **7**, 13528; 10.1038/s41598-017-13236-z
(2017).

23. Spillane, J.L., LaPolice, T.M., MacManes, M.D. & Plachetzki, D.C., Signal, bias, and the role
of transcriptome assembly quality in phylogenomic inference. *BMC Ecol. Evol.* **21**, 43;
10.1186/s12862-021-01772-2 (2021).

24. Quek, Z.B.R. and Huang, D., Effects of missing data and data type on phylotranscriptomic
analysis of stony corals (Cnidaria: Anthozoa: Scleractinia). *Mol. Phyl. Evol.* **134**, 12–23 (2019).

25. Dylus, D., Altenhoff, A., Majidian, S., Sedlazeck, F. J. & Dessimoz, C. Inference of
phylogenetic trees directly from raw sequencing reads using Read2Tree. *Nat. Biotechnol.* **42**,
139–147 (2024).

26. Park, S., Kwak, M. & Park, S., Complete organelle genomes of Korean fir, *Abies koreana* and
phylogenomics of the gymnosperm genus *Abies* using nuclear and cytoplasmic DNA sequence
data. *Sci. Rep.* **14**, 7636; 10.1038/s41598-024-58253-x (2024).

27. Soltis, P. S., & Soltis, D. E., The Role of Hybridization in Plant Speciation, *Annu. Rev. Plant
Biol.* **60**, 561–588 (2009).

28. Rieseberg, L. H., & Willis, J. H., Plant Speciation. *Science* **317**, 910–914 (2007).

29. Nelson, T.C. et al. Ancient and recent introgression shape the evolutionary history of pollinator
adaptation and speciation in a model monkeyflower radiation (*Mimulus* section *Erythranthe*).
PLoS Genet. **17**, e1009095; 10.1371/journal.pgen.1009095 (2021).

30.   Suvorov, A. et al. Deep ancestral introgression shapes evolutionary history of dragonflies and
damselflies. *Syst. Biol.* **71**, 526–546 (2022).

31.   Maddison, W. P. Gene trees in species trees. *Syst. Biol.* **46**, 523–536 (1997).

32.   Meleshko, O. et al. Extensive genome-wide phylogenetic discordance is due to incomplete
lineage sorting and not ongoing introgression in a rapidly radiated *Bryophyte* genus. *Mol Biol*
Evol. **38**, 2750–2766 (2021).

33.   Kandziora, M., Sklenář, P., Kolář, F. & Schmickl, R., How to tackle phylogenetic discordance
in recent and rapidly radiating groups? Developing a workflow using *Loricaria* (Asteraceae) as
an example. *Front. Plant Sci.* **12**, 765719; 10.3389/fpls.2021.765719 (2022).

34.   Wanke, S. and Wicke, S., Phylogenomic discordance in plant systematics. *Front. Plant Sci.* **14**,
1308126; 10.3389/fpls.2023.1308126 (2023).

35.   Nagano, Y. et al. Phylogenetic relationships of Aurantioideae (Rutaceae) based on RAD-Seq.
Tree Genet. Genomes **14**, 6; 10.1038/nrg.2015.28 (2018).

36.   Herrero, R., Asins, M.J., Carbonell, E.A. & Navarro, L., Genetic diversity in the orange
subfamily Aurantioideae. I. Intraspecies and intragenus genetic variability. *Theor. Appl. Genet.*
92, 599–609 (1996).

37.   Penjor, T. et al. Phylogenetic relationships of *Citrus* and its relatives based on *matK* gene
sequences. *PLoS One* **8**, e62574; 10.1371/journal.pone.0062574 (2013).

38.   Iwamasa, M., Nito, N. & Ling, J. T. Intra- and intergeneric hybridization in the orange
subfamily, Aurantioideae *in Proceedings of the Sixth International Citrus Congress* (eds.
Goren, R. & Mendel, K.) 123–130 (Balaban Publishers, Philadelphia/Rehovot Margraf
Scientific Books, Weikersheim, 1988).

39.   Gabaldón, T. & Koonin, E.V., Functional and evolutionary implications of gene orthology. *Nat.*
Rev. Genet. **14**, 360–366 (2013).

40. Swingle, W. T., Reece, P. C. The botany of Citrus and its wild relatives in the orange subfamily

in *The citrus industry, vol 1* (eds. Reuther, W., Webber, H. J. & Bachelor, L. D.) 190–430

(University of California, Berkeley, 1967).

41. Edelman, N.B. et al. Genomic architecture and introgression shape a butterfly radiation.

Science **366**, 594–599 (2019).

42. Malinsky, M., Matschiner, M. & Svardal, H., Dsuite‑Fast D‑statistics and related admixture

evidence from VCF files. *Mol. Ecol. Resour.* **21**, 584–595 (2021).

43. Pease, J.B. & Hahn, M.W., Detection and polarization of introgression in a five-taxon

phylogeny. *Syst. Biol.* **64**, 651–662 (2015).

44. Shi, W., Song, W., Zhao, Y., Shi, C., & Wang, S. Complete chloroplast genomes of four

Atalantia (Rutaceae) species: insights into comparative analysis, phylogenetic relationships,

and divergence time estimation. *Plant Syst. Evol.* **309**, 31; 10.1007/s00606-023-01868-w

(2023).

45. Hodel, R. G. J., Zimmer, E. A., Liu, B. B., & Wen, J. Synthesis of nuclear and chloroplast data

combined with network analyses supports the polyploid origin of the apple tribe and the hybrid

origin of the Maleae-Gillenieae Clade. *Front. Plant Sci.* **12**, 820997; 10.3389/fpls.2021.820997

(2022).

46. Yu, W. B., Huang, P. H., Li, D. Z. & Wang, H. Incongruence between nuclear and chloroplast

DNA phylogenies in *Pedicularis* section *Cyathophora* (Orobanchaceae). *PLoS One* **8**, e74828;

10.1371/journal.pone.0074828 (2013).

47. Bruun-Lund, S., Clement, W. L., Kjellberg, F., & Rønsted, N. First plastid phylogenomic study

reveals potential cyto-nuclear discordance in the evolutionary history of *Ficus* L. (Moraceae).

Mol. Phylogenet. Evol. **109**, 93–104 (2017).

48. Morales-Briones, D.F. et al. Disentangling sources of gene tree discordance in phylogenomic

data sets: testing ancient hybridizations in Amaranthaceae s.l *Syst. Biol.* **70**, 219–235 (2021).

49. Hibbins, M.S., Breithaupt, L.C. & Hahn, M.W., Phylogenomic comparative methods: Accurate

evolutionary inferences in the presence of gene tree discordance. *PNAS* **120**, e2220389120;

10.1073/pnas.2220389120 (2023).

50. Mendes, F.K., Hahn, Y. & Hahn, M.W., Gene tree discordance can generate patterns of

diminishing convergence over time. *Mol. Biol. Evol.* **33**, 3299–3307 (2016).

51. Liu, X. et al. Genomic insights into zokors' phylogeny and speciation in China. *PNAS* **119**,

e2121819119; 10.1073/pnas.2121819119 (2022).

52. Hernández-Gutiérrez, R. et al. Localized Phylogenetic Discordance Among Nuclear Loci Due

to Incomplete Lineage Sorting and Introgression in the Family of Cotton and Cacao

(Malvaceae). *Front. Plant Sci.* **13**, 850521; 10.3389/fpls.2022.850521 (2022).

53. Yasuda, K., et al. Identification of parental chromosomes in sexual intergeneric hybrid

progenies between *Citrus* cultivar 'Nanpu' tangor and *Citropsis schweinfurthii* in the subfamily

Aurantioideae. *J. Japan. Soc. Hort. Sci.* **79**, 129–134 (2010).

54. Smith, M.W., Gultzow, D.L. & Newman, T.K., First fruiting intergeneric hybrids between

Citrus and *Citropsis*. *J. Am. Soc. Hort. Sci.* **138**, 57–63 (2013).

55. Ramadugu, C. et al. A six nuclear gene phylogeny of *Citrus* (Rutaceae) taking into account

hybridization and lineage sorting. *PLoS One* **8**, e68410; 10.1371/journal.pone.0068410 (2013).

56. Fu, X.G., et al. Phylogenomic analysis of the hemp family (Cannabaceae) reveals deep

cyto☐nuclear discordance and provides new insights into generic relationships. *J. Syst. Evol.*

61, 806-826 (2023).

57. Aardema, M.L., Schmidt, K.L. & Amato, G., Patterns of cytonuclear discordance and

divergence between subspecies of the scarlet macaw (*Ara macao*) in Central America. *Genetica*

151, 281-292 (2023).

58. Duan, L., Fu, L. & Chen, H.F., Phylogenomic cytonuclear discordance and evolutionary
histories of plants and animals. *Sci. China Life Sci.* **66**, 2946–2948 (2023).

59. Phuong, M.A., Bi, K. & Moritz, C., Range instability leads to cytonuclear discordance in a
morphologically cryptic ground squirrel species complex. *Mol. Ecol.* **26**, 4743–4755 (2017).

60. Hibbins, M.S. & Hahn, M.W., The timing and direction of introgression under the multispecies
network coalescent. *Genetics* **211**,1059–1073 (2019).

61. Seixas, F.A., Boursot, P. & Melo-Ferreira, J., The genomic impact of historical hybridization
with massive mitochondrial DNA introgression. *Genome Biol.* **19**, 91; 10.1186/s13059-018-
1471-8 (2018).

62. Nge, F.J., Biffin, E., Thiele, K.R. & Waycott, M., Reticulate evolution, ancient chloroplast
haplotypes, and rapid radiation of the Australian plant genus *Adenanthos* (Proteaceae). *Front.*
Ecol. Evol. **8**, 616741; 10.3389/fevo.2020.616741 (2021).

63. Stull, G. W., Pham, K. K., Soltis, P. S., & Soltis, D. E. Deep reticulation: the long legacy of
hybridization in vascular plant evolution. *Plant J.* **114**, 743–766 (2023).

64. Sasaki, Y. et al. Chloroplast-encoded protein as a subunit of acetyl-CoA carboxylase in pea
plant. *J. Biol. Chem.* **268**, 25118–25123 (1993).

65. Sasaki, Y. & Nagano, Y. Plant acetyl-CoA carboxylase: structure, biosynthesis, regulation, and
gene manipulation for plant breeding. *Biosci. Biotechnol. Biochem.* **68**, 1175–1184 (2004).

66. de Vries, J., Sousa, F. L., Bölter, B., Soll, J. & Gould, S. B. YCF1: a green TIC?. *Plant Cell* **27**,
1827–1833 (2015).

67. Sasaki, Y., Sekiguchi, K., Nagano, Y. & Matsuno, R. Chloroplast envelope protein encoded by
chloroplast genome. *FEBS Lett.* **316**, 93–98 (1993).

68. Trinh, M. D. L. et al. Lack of plastid-encoded Ycf10, a homolog of the nuclear-encoded
698   DLDG1 and the cyanobacterial PxcA, enhances the induction of non-photochemical quenching
699   in tobacco. *Plant Direct* **5**, e368; 10.1002/pld3.368 (2021).

700   69. Chen, S., Zhou, Y., Chen, Y., & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
701   *Bioinform.* **34**, i884–i890 (2018).

702   70. Capella-Gutiérrez, S., Silla-Martínez, J. M. & Gabaldón, T. trimAl: A tool for automated
703   alignment trimming in large-scale phylogenetic analyses. *Bioinform.* **25**, 1972–1973 (2009).

704   71. Darriba, D. et al. ModelTest-NG: A new and scalable tool for the selection of DNA and protein
705   evolutionary models. *Mol. Biol. Evol.* **37**, 291–294 (2020).

706   72. Stamatakis, A. RAxML version 8: A tool for phylogenetic analysis and post-analysis of large
707   phylogenies. *Bioinform.* **30**, 1312–1313 (2014).

708   73. Huson, D. H. & Scornavacca, C. Dendroscope 3: An interactive tool for rooted phylogenetic
709   trees and networks. *Syst. Biol.* **61**, 1061–1067 (2012).

710   74. Jin, J. J. et al. GetOrganelle: A fast and versatile toolkit for accurate de novo assembly of
711   organelle genomes. *Genome Biol.* **21**, 241; 10.1186/s13059-020-02154-5 (2020).

712   75. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**,
713   357–359 (2012).

714   76. Li, H. et al. The sequence alignment/map format and SAMtools. *Bioinform.* **25**, 2078–2079
715   (2009).

716   77. Robinson, J. T., Thorvaldsdóttir, H., Turner, D. & Mesirov, J. P. igv. js: An embeddable
717   JavaScript implementation of the Integrative Genomics Viewer (IGV). *Bioinform.* **39**, btac830;
718   10.1093/bioinformatics/btac830 (2023).

719   78. Tillich, M. et al. GeSeq: versatile and accurate annotation of organelle genomes. *Nucleic Acids*
720   *Res.* **45**, W6–W11 (2017).

79. Zhong, X. Assembly, annotation and analysis of chloroplast genomes. Doctoral thesis in *the*

University of Western Australia, *Australia* (2020).

80. Chan, P. P., Lin, B. Y., Mak, A. J. & Lowe, T. M. tRNAscan-SE 2.0: improved detection and

functional classification of transfer RNA genes. *Nucleic Acids Res.* **49**, 9077–9096 (2021).

81. Potter, S. C. et al. HMMER web server: 2018 update. *Nucleic Acids Res.* **46**, W200–W204

(2018).

82. Greiner, S., Lehwark, P. & Bock, R. *OrganellarGenomeDRAW (OGDRAW)* version 1.3.1:

Expanded toolkit for the graphical visualization of organellar genomes. *Nucleic Acids Res.* **47**,

W59–W64 (2019).

83. Bi, G., Mao, Y., Xing, Q. & Cao, M. HomBlocks: A multiple-alignment construction pipeline

for organelle phylogenomics based on locally collinear block searching. *Genomics* **110**, 18–22

(2018).

84. Drummond, A. J., Suchard, M. A., Xie, D. & Rambaut, A. Bayesian phylogenetics with

BEAUti and the BEAST 1.7. *Mol. Biol. Evol.* **29**, 1969–1973 (2012).

85. Zhang, C., Rabiee, M., Sayyari, E. & Mirarab, S., ASTRAL-III: polynomial time species tree

reconstruction from partially resolved gene trees. *BMC Bioinform.* **19,** 153; 10.1186/s12859-

018-2129-y (2018).

86. Smith, S.A., Moore, M.J., Brown, J.W. & Yang, Y., Analysis of phylogenomic datasets reveals

conflict, concordance, and gene duplications with examples from animals and plants. *BMC*

Ecol. Evol. **15**, 150; 10.1186/s12862-015-0423-0 (2015).

87. Wen, D., Yu, Y., Zhu, J. and Nakhleh, L., Inferring phylogenetic networks using PhyloNet.

Syst. Biol. **67**, 735-740 (2018).

88. Sullivan, J., & Joyce, P. Model selection in phylogenetics. *Annu. Rev. Ecol. Evol. Syst.* **36**,

445–466 (2005).

89. Akaike, H. In *S*elected Papers of Hirotugu Akaike (eds Emanuel Parzen, Kunio Tanabe, &
Genshiro Kitagawa) 199–213 (Springer New York, 1998).

90. Altschul, S. F. et al. Gapped BLAST and PSI-BLAST: a new generation of protein database
search programs. *Nucleic Acids Res.* **25**, 3389–3402 (1997).

91. Sullivan, M. J., Petty, N. K. & Beatson, S. A. Easyfig: a genome comparison visualizer.
Bioinform. **27**, 1009–1010; (2011).

92. Tamura, K., Stecher, G. & Kumar, S. MEGA11: Molecular evolutionary genetics analysis
version 11. *Mol. Biol. Evol.* **38**, 3022–3027 (2021).

93. Kumar, S. et al. TimeTree 5: An expanded resource for species divergence times. *Mol. Biol.*
Evol. **39**, msac174; 10.1093/molbev/msac174 (2022).

94. Emms, D. M. & Kelly, S. OrthoFinder: Phylogenetic orthology inference for comparative
genomics. *Genome Biol.* **20**, 238; 10.1186/s13059-019-1832-y (2019).

95. Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment software version 7:
Improvements in Performance and Usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

Acknowledgements

In the initial phase of our study's data analysis conducted in Sri Lanka; we extend our profound gratitude to Dr. Dhammika Elkaduwe from the Faculty of Engineering at the University of Peradeniya for his exceptional support in granting access to high-performance computing facilities. We would like to thank Ms. Akiko Sakai for technical assistance, and Mr. Takashi Arita for providing the experimental samples. This work was partially supported by a Grant-in-Aid for Scientific Research (18K05623) from the Japan Society for the Promotion of Science to Yukio Nagano. E.P.W. was supported by the Japanese Government (MEXT) scholarship. We would also like to thank Editage (https://www.editage.com) for the English language editing. This study was part of a dissertation submitted by the first author toward the partial fulfillment of her Ph.D. All authors provided consent for submission and publication.

Author contributions

E.P.W., Y.N., and T.I. designed the study. M.Y. and N.K. supplied the experimental samples. Y.N. performed the DNA experiments. E.P.W., T.I., M.H.S., and Y.N. conducted the bioinformatics analysis. N.U.J. established the framework for conducting the study, including an initial phase of data analysis conducted in Sri Lanka. As experts on Aurantioideae, M.Y. and N.K. reviewed the descriptions of Aurantioideae. All authors were involved in the data analysis. The manuscript was written by E.P.W. and Y.N. and reviewed by all authors.

Data Availability Statement

Raw sequencing data have been deposited in the DDBJ Sequence Read Archive

(https://www.ddbj.nig.ac.jp/dra/index-e.html; accession no. DRA017638). The cp genome sequences

assembled and annotated from these raw sequencing data have been deposited in

DDBJ/EMBL/GenBank under accession numbers LC794878 to LC794908. The cp genome

sequences, assembled and annotated from publicly available raw sequencing data, are available in

GenBank format in Supplementary Dataset 1. The alignments, tree data and Nexus files used in this

study are included in the Supplementary Datasets 2 through 21.

Competing interests

The authors declare that they have no conflicts of interest.

Figure Legends

Figure 1. ML phylogenetic tree based on conserved nuclear gene sequences of 39 species in the

Aurantioideae subfamily.

The alignment used for phylogenetic tree construction included a total of 6,674,283 nucleotide

positions with 605,597 parsimony-informative sites, as calculated using custom Python scripts

(Supplementary Dataset 15). For the analysis, the optimal evolutionary model, GTRGAMMAIX,

was employed in RAxML. The root position corresponded to that identified in the phylogenetic tree

of the Rutaceae family (Supplementary Fig. S1). Bootstrap values (percentages over 100 replicates)

are displayed at the nodes, with the scale bar indicating genetic divergence in substitutions per site.

Bootstrap values below 50% are omitted. The subgroups identified in this study are represented by

different colors. The alignment data in FASTA format and tree data in Newick format are available

in Supplementary Dataset 2.

Figure 2. Discordance between nuclear gene-derived trees and cp tree

(A) Comparison of the concatenated nuclear tree and the cp tree. Left panel: ML phylogenetic tree

generated using Read2Tree, based on conserved nuclear gene sequences from 39 species within the

Aurantioideae subfamily. Bootstrap support values (percentages from 100 replicates) are displayed

at each node, with values below 50% omitted. Subgroups identified in this study are represented by

different colors.

(B) Comparison of the ASTRAL species tree and the cp tree. Left panel: Species tree inferred using

ASTRAL-III, with local posterior probability (LPP) values displayed as percentages for each

corresponding node. Subgroups identified in this study are represented by different colors. Tree data

in Newick format are available in Supplementary Dataset 4.

Right panel in both A and B: ML phylogenetic tree constructed using the whole cp genome

sequences of 39 species in the Aurantioideae subfamily. Bootstrap support values (percentages from

1,000 replicates) are displayed at the nodes, with values below 50% omitted. Asterisks mark

sequences that were assembled *de novo* in this study. Subgroups identified are represented by

distinct colors. Dotted lines and shapes highlight conflicting relationships between the nuclear and

cp phylogenies, revealing areas of topological incongruence between the two trees. Alignment data

(FASTA format) and tree data (Newick format) are available in Supplementary Dataset 3.

Figure 3. Species tree and concatenated nuclear gene tree

(A) Comparison of the concatenated nuclear gene tree and ASTRAL species tree. Left panel: ML

phylogenetic tree generated using Read2Tree, based on conserved nuclear gene sequences from 39

species within the Aurantioideae subfamily. Bootstrap support values (percentages from 100

replicates) are displayed at each node, with values below 50% omitted. Subgroups identified in this

study are highlighted in distinct colors. Right panel: Species tree inferred using ASTRAL-III, with

local posterior probability (LPP) values displayed as percentages for each corresponding node.

Subgroups identified in this study are denoted by distinct colors. Dotted lines indicate areas of

incongruence between the two trees.

(B) Gene tree discordance displayed on the ASTRAL-inferred species tree. Pie charts at each node

illustrate the proportion of gene trees falling into one of four categories: concordant with the species

tree (blue), discordant but representing the most common alternative topology (green), discordant

with other alternative topologies (red), and uninformative due to low branch support (gray, <33%

bootstrap support at the node). The number above each branch indicates the count of gene trees

concordant with the species tree topology at that node (blue portion of the pie). The number below

each branch represents the sum of informative discordant topologies at that node (green + red

portions of the pie). Different colors denote the subgroups identified in this study.

Figure 4. Discordance analysis among three subgroups: (1) True citrus fruit trees and its near

subgroups, (2) Bael fruit subgroup and its associates, and (3) the Group B. One representative from

each subgroup/genus is used.

Figure 5. Summary of Phylogenetic Discordance Sources

(A) ASTRAL-inferred species tree depicting the relationships among members of the Aurantioideae subfamily. Arrows indicate various evolutionary events: the dashed blue arrow represents ILS; green arrows represent introgression; and pink arrows indicate ancient hybridization followed by ILS. Detailed results are provided in Supplementary Tables 4 to 13. One representative from each genus/subgroup is used.

Phylogenetic networks for True citrus fruit trees and its near subgroups (B), as well as the Bael fruit subgroup and associated species (C), were inferred using PhyloNet with Maximum Pseudolikelihood under models allowing for one, two, and three reticulations. The Akaike Information Criterion (AIC) indicates that the three-reticulation model is the best fit for both groups. In Figure (B), green branches highlight lineages involved in reticulate evolutionary histories, with numerical values representing the inheritance probabilities for each reticulation event. In Figure (C), purple branches similarly indicate lineages associated with reticulation, along with their corresponding inheritance probabilities. The Nexus files for each reticulation model can be found in Supplementary Datasets 5 and 6. Supplementary Figs. S5 and S6 present all inferred phylogenetic networks generated by PhyloNet for True citrus fruit trees and its near subgroups and the Bael fruit subgroup with associated species, respectively.

A)

Concatenated nuclear gene tree from Read2Tree

ASTRAL species tree

B)

1. Conflicts in True citrus fruit trees and its near subgroups

Concatenated gene tree

ASTRAL-III Species tree

Whole Chloroplast tree

2. Conflicts in Bael fruit subgroup and its associates

Concatenated gene tree

ASTRAL-III Species tree

Whole Chloroplast tree

3. Conflicts in Group B

Concatenated gene tree

ASTRAL-III Species tree

Whole Chloroplast tree

A)

True citrus fruit trees
Yaki Naran
Citropsis
Naringi
Swinglea
Wood apple
Afraegle
Aegle
Paramignya
Triphasia
Murraya
Clausena
Murraya koengii
Glycosmis
Micromelum minutum

Group A

Group B

B)

Under a one reticulation model
log probability: -4690.716

Under a two reticulation model
log probability: -4668.391

Under a three reticulation model
log probability: -4683.586

C)

Under a one reticulation model
log probability: -2723.046

Under a two reticulation model
log probability: -2621.664

Under a three reticulation model
log probability: -2673.374