miR-100 and *miR-125b* Contribute to Enhanced 3D Growth and Invasiveness and can be Functionally Transferred to Silence Target Genes in Recipient Cells

Hannah M. Nelson[1], Shimian Qu[1], Sydney N. Chapman[1], Nicole L. Luthcke[1], Sara A. Schuster[1], Lauren A. Turnbull[1], Muhammad Shameer[1], Lucas L. Guy[1], Xiao Liu[2], Kevin C. Corn[3], Kasey C. Vickers[4], Qi Liu[2], Jeffrey L. Franklin[5], Alissa M. Weaver[6], Marjan Rafat[3], Robert J. Coffey[7], James G. Patton[1*]

[1]Laboratory of James G. Patton, Department of Biological Sciences, Center for Extracellular Vesicle Research, Vanderbilt University, Nashville, TN, United States

[2]Laboratory of Qi Liu, Department of Biostatistics, Department of Biomedical Informatics, Center of Extracellular Vesicle Research, Vanderbilt University Medical Center, Nashville, TN, United States

[3]Laboratory of Marjan Rafat, Department of Biomedical Engineering, Department of Chemical and Biomolecular Engineering, Department of Radiation Oncology, Vanderbilt University, Nashville, TN, United States.

[4]Laboratory of Kasey C. Vickers, Department of Pathology, Microbiology and Immunology, Department of Medicine, Department of Molecular Physiology and Biophysics, Center for Extracellular Vesicle Research, Vanderbilt University Medical Center, Nashville, TN, United States.

[5]Department of Cell and Developmental Biology, Center for Extracellular Vesicle Research, Vanderbilt University, Vanderbilt University Medical Center, Nashville, TN, United States.

[6]Laboratory of Alissa M. Weaver, Department of Cell and Developmental Biology, Department of Pathology, Microbiology and Immunology, Center for Extracellular Vesicle Research, Vanderbilt University Medical Center, Nashville. TN, United States.

[7]Laboratory of Robert J. Coffey, Department of Medicine, Division of Gastroenterology, Hepatology and Nutrition, Department of Cancer Research, Department of Cell and Developmental Biology, Center for Extracellular Research, Vanderbilt University Medical Center, Nashville, TN, United States.

*Corresponding Author:
James G Patton
Department of Biological Sciences, Vanderbilt University
james.g.patton@vanderbilt.edu

Keywords: miRNA, miR-100, miR-125b, Colorectal Cancer, Cingulin, Invasiveness, Tight Junctions

Number of words in main text: 4,578
Number of figures/tables in main text: 7

Abstract

Extracellular communication via the transfer of vesicles and nanoparticles is now recognized to play an important role in tumor microenvironment interactions. Colorectal cancer (CRC) cells upregulate and secrete abundant levels of *miR-100* and *miR-125b* that can alter gene expression by cell- and non-cell-autonomous mechanisms. We previously showed that these miRNAs activate Wnt signaling through noncanonical pairing with 5 negative regulators of Wnt signaling. To identify additional targets of *miR-100* and *miR-125b*, we used bioinformatic approaches comparing multiple CRC cell lines, including knockout lines lacking one or both miRNAs. From an initial list of 96 potential mRNA targets, we tested 15 targets with 8 showing significant downregulation in the presence of *miR-100* and *miR-125b*. Among these, Cingulin (CGN) and Protein tyrosine phosphatase receptor type-R (PTPRR) are downregulated in multiple cancers, consistent with regulation by increased levels of *miR-100* and *miR-125b*. We also show that increased levels of *miR-100* and *miR-125b* enhance 3D growth and invasiveness. Lastly, we show that *miR-100* and *miR-125b* can be functionally transferred between cells to silence both reporter and endogenous mRNA targets.

1   Introduction

MicroRNAs (miRNAs) are small (22-23nt) RNAs that base pair with target mRNAs to regulate gene expression by decreasing translation and promoting RNA degradation (He and Hannon, 2004, Krol et al., 2010). In cancer, miRNA expression patterns are frequently altered, contributing to all aspects of tumor progression from proliferation to metastasis (Peng and Croce, 2016). We and others have focused on the role of *miR-100* and *miR-125b* in cancer (Cha et al., 2015, Qin et al., 2015, Peng et al., 2021, Liu et al., 2022, Lu et al., 2017, Zhang et al., 2021). These miRNAs are encoded in the third intron of *MIR100HG*, a 3kb lncRNA that is transcribed from the large 330kb MIR100HG locus (Emmrich et al., 2014). Increased expression of *miR-100, miR-125* and *MIR100HG* can promote cetuximab (anti-EGFR) resistance in colorectal cancer (CRC) cells, drive progression of pancreatic cancer, and promote the epithelial to mesenchymal transition (Lu et al., 2017, Liu et al., 2022, Ottaviani et al., 2018). These effects are driven in part by regulating the expression of target mRNAs including mTOR (Cha et al., 2015), negative regulators of Wnt signaling (Lu et al., 2017), Cystic Fibrosis Transmembrane Regulator (CFTR) (Zhang et al., 2021), Cingulin (CGN) (Zhang et al., 2021), and checkpoint kinases (CHK) (Ottaviani et al., 2018). These are just a small subset of targets for these miRNAs; TargetScan predicts a large number of potential targets for *miR-100* (59 targets) and *miR-125b* (931 targets)(https://www.targetscan.org/vert_80/). Base pairing between miRNAs and target 3'-UTRs is typically imperfect, with canonical targets displaying extensive pairing between nucleotides 2-8 of the miRNA (the "seed" sequence) and variable complementarity throughout the rest of the miRNA (Bartel, 2009). However, even within the seed sequence, the extent of base pairing between miRNAs and their mRNA targets is highly variable making prediction of mRNA targets a challenge (Li et al., 2008, Riolo et al., 2021). Indeed, all 5 of the negative regulators of Wnt (DKK1, DKK3, ZNRF3, RNF43, and APC2) that we previously found to be targeted by *miR-100* and *miR-125b* are noncanonical targets, raising the question as to whether additional targets might contribute to cetuximab resistance and altered 3D growth.

Extracellular vesicles (EVs) and nanoparticles participate in cell-cell communication through the transfer of RNA, protein and lipids (Couch et al., 2021, Dixson et al., 2023). Among RNA cargo for EVs, transfer of miRNA is the most well characterized due to their small size and

the ability to experimentally monitor silencing of target mRNAs in recipient cells (O'Brien et al.,
2020). We previously showed that *miR-100* and *miR-125b* are enriched in small EVs (sEVs)
from CRC cells with Transwell co-culture experiments supporting extracellular transfer of these
miRNAs to recipient cells (Cha et al., 2015). However, since the recipient cells in those transfer
experiments expressed endogenous *miR-100* and *miR-125b*, it remains a possibility that the
silencing of mRNA targets in recipient cells could be due to unexpected activation of the
endogenous genes (Gruner and McManus, 2021).

To identify targets of *miR-100* and *miR-125b* and to definitively demonstrate
extracellular transfer, we created cell lines with CRISPR/Cas9-mediated knockouts of *miR-100,*
miR-125b, or both. Through extensive bioinformatic analyses of RNAseq data, we identified and
tested 15 candidate targets for *miR-100* and *miR-125b* with the two most statistically significant
being Cingulin (CGN) and Protein Tyrosine Phosphatase Receptor Type-R (PTPRR). We also
show that *miR-100* and *miR-125b* contribute to enhanced 3D growth and invasiveness of CRC
cells. Lastly, we demonstrate functional transfer of *miR-100* and *miR-125b* from donor to
recipient cells leading to silencing of both reporter and endogenous mRNAs.

2   Materials and Methods

2.1 Cell Culture
CC and CC-CR CRC cells were obtained from the Coffey lab (Lu et al., 2017). All cell
lines were confirmed to be free of mycoplasma contamination (Venor^{TM}GeM Mycoplasma
Detection Kit, PCR-based, Sigma Aldrich Cat #MP0025). Cells were grown in Dulbecco's
Modified Eagle's Medium (DMEM, Corning) supplemented with 10% fetal bovine serum (FBS,
Corning), 1% L-glutamine (200 nM, Gibco), 1% MEM nonessential amino acids (Sigma
Aldrich), and 1% penicillin-streptomycin (10,000 U/mL, Gibco) in 5% CO_2 at 37°C. Cells were
passaged a maximum of 8 times before discarding.

2.2 Generation of Knockout Cell Lines
We used CRISPR/Cas9 multiplex genome engineering (Kabadi et al., 2014) to delete
miR100, miR125b, or both. Guide RNAS (gRNAs) flanking the targeted locus were designed
using the Benchling CRISPR gRNA Design Tool (https://www.benchling.com/crispr).
Oligonucleotides containing gRNA sequences were first cloned into sgRNA expression vectors
(Addgene #53186-53189) and then multiple gRNA cassettes were cloned into lentiviral
expression vectors *(*pLV hUbC-Cas9-T2A-GFP; Addgene, #53190) using Golden Gate cloning
technology. Lentiviral stocks were prepared in 293T cells and used to transduce CC-CR cells.
Single GFP$^+$ cells were sorted into 96 cell plates and expanded. PCR amplification of genomic
DNA from clonal GFP$^+$ cells was performed using primers flanking the gRNA target sequences
to identify deletion clones.

2.3 Extracellular Vesicle Isolation
Cells were plated at 9.0×10^6 cells in T-175 flasks (Corning), with each EV collection
using at least 3 flasks. Cells were grown to 80% confluency and washed 3 times with 1X PBS
before media were replaced with DMEM lacking FBS and grown for 48 hours. Culture media
were then collected and adherent cells were trypsinized and counted. Conditioned media were
then centrifuged at increasing speeds and time: 1,000xg for 10 minutes (room temperature),
2,000xg for 20 minutes (4°C), and 10,000xg for 30 minutes (4°C). Crude EV pellets were

obtained after the conditioned media was centrifuged for 17 hours at 100,000xg (4°C). EV
pellets were then washed two times by resuspension in 1mL of 1X DPBS (Corning) and
centrifugation at 100,000xg for 70 minutes. Final pellets were resuspended in 100µl 1X DPBS.
Under these conditions, the EV pellets consist of a heterogenous mixture of sEVs with some
non-vesicular material (Jeppesen et al., 2019). This protocol was used to directly compare with
earlier results examining export of *miR-100 and miR-125b* into CRC EVs (Cha et al., 2015).
sEVs were quantified by protein concentration (Pierce BCA Protein Assay) and by Nanoparticle
Tracking Analysis (NTA, ZetaView Nanoparticle Tracking Analysis Instrument).

2.4 qRT-PCR
131
Total RNA from both whole cell and sEVs was isolated using TRIzol (Life
Technologies). Taqman small RNA assays (Life Technologies) were used to quantify miRNA
levels. A total of 10ng of RNA were utilized for each RT reaction; 0.67 µl of cDNA was used in
each 10µl qPCR reactions. qPCR was completed in either 96-well or 384-well plates using
Biorad CFX96 or CFX384 instruments. Fold changes were calculated as previously described
(Cha et al., 2015).

2.5 Transfer Assays
138
139 For Transwell co-culture assays, cells were plated in Transwell dishes with 0.4mm filters
140 (Corning, 3460) with 50,000 cells in the donor well and 300,000 cells in the recipient well.
141 Transwell co-cultures were grown for five days, changing media every 72 hours. Cells were
142 washed 3 times with 2 mL of 1X PBS; RNA was then collected from either donor or recipient
143 cells.

2.6 AGO2 Immunoprecipitation
144
For immunoprecipitation, anti-AGO2 (clone 11A9 antibody; MABE253, Sigma-Aldrich)
or control IgG antibodies were used. Ten 150cm dishes of either CC or CC-CR cells were grown
to 95% confluency. Cells were then scraped in ice cold PBS and centrifuged to collect cell
pellets in ice cold lysis buffer (20mM Tris HCl pH 7.5, 150 mM KCl, 0.5% NP40, 2mM EDTA,
1mM NaF, 0.5mM DTT). After centrifugation of insoluble cell debris, cell lysates were
incubated with Dynabeads. Prior to cell lysis, Dynabeads™ Protein G (ThermoFisher, 10003D)
were conjugated with antibodies by incubating overnight with rotation at 4°C and washed two
times with ice cold lysis buffer. Cell lysates were then added to the conjugated beads and
incubated at 4°C for six hours. Beads were gently washed with fresh lysis buffer and proteins
were eluted. For RNA isolation, beads were incubated with DNase and Proteinase K for 10
minutes at room temperature. Trizol was then added and incubated at room temp for 10 minutes
and RNA was isolated.

2.7 Small RNA Sequencing
157
RNA libraries were generated using the NEXTFlex Small RNA Library Preparation Kits
v3 (Perkin) with the following modifications: (1) 3′- and 5′-adaptors were diluted 1:8, (2) 3′-
adaptor ligations were performed overnight in multiple steps –25°C for 2 h, 20°C for 4 h and
16°C overnight, (3) following cDNA synthesis and before barcoding PCR, step F protocol was
followed, and (4) PCR amplification was 20 cycles. Following PCR amplification, individual
libraries were size-selected (136–200 bp product) using Pippin Prep (Sage Sciences). Size-
selected libraries were quantified using a Qubit Fluorometer. Libraries were checked for quality
and sequenced using Illumina short-read technology. Libraries were pooled and paired-end

sequencing (PE-150) of the pool (equimolar multiplexed libraries) was performed on the
NovaSeq6000 platform by the VANTAGE core (Vanderbilt University). Demultiplexing and
bioinformatic analyses were performed using the TIGER pipeline (Allen et al., 2018). Briefly,
Cutadapt (v1.16) was used to trim 3′ adaptors and all reads with <16 nucleotides (nts) were
removed (Martin, 2011). Quality control on both raw reads and adaptor-trimmed reads was
performed using FastQC (v0.11.9)(www.bioinformatics.babraham.ac.uk/projects/fastqc). The
adaptor-trimmed reads were mapped to the hg19 genome, with additional rRNA and tRNA
reference sequences using Bowtie1 (v1.1.2), allowing only one mismatch (Langmead and
Salzberg, 2012).

2.8 Total RNA Sequencing
Bulk RNA sequencing libraries were prepared using Universal RNAseq kits (Tecan).
Libraries were cleaned (Zymo), checked for quality using the Agilent bioanalyzer, quantified
(Qubit), and pooled based on equimolar concentrations. Pooled libraries were sequenced using
Illumina short-read technology based on PE150 on the NovaSeq6000 (Illumina). After
sequencing, samples (libraries) were demultiplexed and analyzed. Briefly, adapter sequences
were removed using Cutadapt v2.10)(Martin, 2011). Quality control of both raw reads and
adaptor-trimmed reads was performed using FastQC (v0.11.9). After adaptor trimming, reads
were aligned to the Gencode GRCh38.p13 genome using STAR (v2.7.8a)(Dobin et al., 2013).
FeatureCounts (v2.0.2)(Liao et al., 2014) was used to count the number of reads mapped to each
gene. Heatmap3 (Zhao et al., 2014) was used for cluster analysis and visualization. Differential
expression was analyzed by DESeq2 (v1.30.1)(Love et al., 2014). Significant differentially
expressed genes were determined with fold change >2 or <0.5, and adjusted p value (padj) <0.05.
Genome Ontology and KEGG pathway over-representation analyses were performed using the
WebGestaltR package (NULL)(Wang et al., 2017). Gene set enrichment analysis (GSEA) was
performed using GSEA package (v4.3.2)(Subramanian et al., 2005) on database v2022.1.Hs.

2.9 Gene Expression Analysis
The Gene Expression database of Normal and Tumor Tissues (GENT2) was used to
compare levels of expression between 72 paired normal and tumor tissues utilizing the GPL570
platform. Statistical significance was determined using two-sample t-tests (GPL570 platform
(HG-U133_Plus_2)).

2.10 Luciferase Reporter Assays
3'-UTRs from mRNAs of interests were cloned using RT/PCR into the pMIR-REPORT™
miRNA Expression Reporter Vector System (Firefly Luciferase) using NEBuilder HiFi DNA
Assembly (New England BioLabs). PCR amplification primers are listed in **Table S1**. All
plasmids were verified by sequencing (Genewiz). Transfected cells were cultured alone or in co-
cultures using Transwells with 0.4mm membranes for 72 hours followed by washing with 1X
PBS. Luciferase levels were then detected using the Dual-Glo® Luciferase Assay System
(Promega). Changes in luciferase levels were normalized to Renilla luciferase levels and
compared to luminescence from the empty (no 3'UTR) vector.

2.11 Transfections
Cells were grown to approximately 50-60% confluency with transfections performed
using TransIT®-2020 (Mirus, MIR5404). The Renilla luciferase construct (pCIneo-RL, Plasmid
115366, Addgene) was transfected at 0.5ug/mL and firefly luciferase construct was transfected

209 at 1ug/mL. DNA and TransIT-2020 were incubated for 30 minutes at room temperature in
210 reduced serum media (Opti-MEM; Gibco). Transfection complexes were diluted 1:10 in DMEM
211 and added to cells.

2.12 Immunofluorescence Staining and Imaging of Cells
Cells were plated on coverslips and grown for three days after which coverslips were
washed three times in 1x PBS and fixed in 4% paraformaldehyde for 30 minutes at room
temperature. Samples were then washed and allowed to block for 1 hour at room temperature in
blocking buffer (1X PBS with 0.001% Triton-X and 0.03% donkey serum). Primary antibody
incubation (CGN, ab244406) was overnight at 4°C, followed by three 1X PBS wash steps, and
incubation with secondary antibodies (Cy3 anti-Rabbit antibody and 488 conjugated phalloidin)
for 2 hours at room temperature. Samples were washed three times with 1X PBS and mounted
on glass slides using Vectashield with DAPI and imaged using a Zeiss LSM 880 microscope
(Vanderbilt University, CISR). All images were analyzed using ImageJ (Aebi et al., 1986).

2.13 Spheroid Cultures
223   Cells (10,000) were plated in Nunclon Sphera low adherent plates (Thermo Scientific)
224 and centrifuged at 1,000xg for 10 minutes at room temperature, as previously described (Carey et
225 al., 2013). Cells were grown for three days before transfer into 2mg/ml Type I collagen
226 (Advanced Biomatrix). Colonies were imaged using Muvicyte Live Cell imager for five days.
227 Images were processed and analyzed using ImageJ(Aebi et al., 1986).

3   Results
To facilitate identification of mRNA targets for *miR-100* and *miR-125b* and to develop
cell lines to test for functional miRNA transfer, we used CRISPR/Cas9 technology to generate
deletions in the MIR100HG locus for these two miRNAs (**Fig. 1A**). A single lentiviral vector
expressing multiple gRNAs, Cas9, and GFP was created and lentiviruses were transduced into
CC-CR cells. CC-CR cells are cetuximab resistant cells that were generated from CC cells by
iterative cycles of selection under 2D (resistant) and 3D (sensitive) growth conditions (Lu et al.,
2017). The gRNAs were designed to base pair with regions flanking *miR-100, miR-125b,* or
both (**Fig. S1**). Individual GFP+ clones were sorted, targeted deletions were identified by PCR,
and verified by sequencing. While CC-CR cells express high levels of *miR-100* and *miR-125b*,
CC cells express ~50-fold less of both miRNAs (Lu et al., 2017), and our different knockout
lines showed undetectable levels of *miR-100 or miR-125b or both* **(Fig. 1B, C).**

3.1 Identification of mRNA Targets of *miR-100 and miR125b*
RNAseq was performed on CC, CC-CR, and knockout cells (Δ*miR-100*, Δ*miR-125b*, and
Δ*miR-100/miR-125b*) and bioinformatic analyses were performed to compare gene expression
patterns (**Table S2**). Potential mRNA targets for *miR-100* and *miR-125b* were identified using a
three-step strategy (**Fig. 2A).** First, we identified upregulated mRNAs in both CC and Δ*miR-*
100/miR-125b cells compared to CC-CR cells (Lu et al., 2017). Second, we compiled a list of
predicted targets for *miR-100* and *miR-125b* using three different prediction algorithms
(TargetScan, TargetMiner, and miRDB; https://www.targetscan.org/vert_80/;
https://www.isical.ac.in/~bioinfo_miu/targetminer20.htm; https://mirdb.org/). Third, we
performed RNA immunoprecipitation of AGO2 followed by bulk and short RNA sequencing
(**Tables S3,4**). AGO2 is a component of the RNA Induced Silencing Complex (RISC) in which
miRNAs pair with their target mRNAs (Müller et al., 2020, Cenik and Zamore, 2011, Farazi et

al., 2008, Liu et al., 2004). Immunoprecipitation of AGO2-containing complexes from CC-CR cells allowed for enrichment of *miR-100* and *miR-125b* targets because these miRNAs are so highly expressed in CC-CR cells (Lu et al., 2017). By comparing the three sets of data, we were able to identify overlapping targets of *miR-100* and *miR-125b* (**Fig. 2A**). Together, we identified 51 potential targets for *miR-100* and 45 potential targets for *miR-125b* (**Fig. 2B**). When we subjected all 96 potential targets to Gene Ontology analysis using ShinyGo (Ge et al., 2020), we found significant enrichment for cell migration, cell motility, actin filament-based processes, and MAPK signaling (**Fig. 2C; Table S5**).

3.2 Silencing of mRNA Targets of *miR-100* and *miR-125b*
From the targets generated above, we examined miRNA:mRNA pairing (**Fig. S2**) and selected 15 candidate mRNAs to test for the ability of *miR-100* (3 candidates), *miR-125b* (7 candidates) or both (5 candidates) to silence luciferase reporter constructs. We designed three different categories of luciferase vectors in which the firefly luciferase open reading frame was fused to (1) no 3'-UTR (empty vector), (2) 3'-UTRs containing three perfect binding sites for *miR-100* or *miR-125b* (100/125 vector), or (3) 3'-UTRs from candidate targets (experimental vectors)(**Fig. 3A**). These vectors were transfected into either CC-CR or Δ*miR-100/miR-125b* cells and luciferase levels were determined. *Bona fide* targets of *miR-100 or miR-125b* should show increased luciferase levels in Δ*miR-100/miR-125b* cells as compared to CC-CR cells. Of the 15 targets we tested, 14 showed increased luciferase levels in Δ*miR-100/miR-125b* cells with 8 being statistically significant (**Fig. 3B**). Of the 8, CGN and PTPRR showed the most significant increase of luciferase activity when comparing Δ*miR-100/miR-125b* and CC-CR cells (**Fig. 3B**). The fact that the two most significant genes are targets of *miR-125b* is consistent with more dramatic overexpression of *miR-125b* in CC-CR cells as compared to *miR-100* (**Fig. 1B**).

3.3 CGN and PTPRR are Downregulated in Colorectal Cancer
miR-100 and *miR-125b* display elevated gene expression and target multiple mRNAs in CRC and other cancer types (Li et al., 2015, Wang et al., 2020, Ottaviani et al., 2018, Cha et al., 2015, Lu et al., 2017, Zhang et al., 2021, Vallacchi and Rodolfo, 2018). Consistent with that, CC-CR cells form solid, disorganized colonies when grown in collagen, indicative of aggressive behavior, as opposed to CC cells which form hollow cysts with a central lumen lined by polarized cells (Lu et al., 2017). We thus sought to test if decreased expression of CGN and PTPRR due to increased expression of *miR-100* and *miR-125b* might be observed in CRC and other cancers. We used the GENT2 database to compare the levels of CGN and PTPRR between normal and cancer tissues (**Fig. 4A,B**). We found that CGN and PTPRR are both downregulated across a variety of cancers, including CRC. There are likely additional mechanisms that can lead to decreased levels of CGN and PTPRR because not all cancers show increased expression of *miR-100* (Li et al., 2015) or *miR-125b* (Wang et al., 2020), but the data are in agreement that decreased levels of CGN and PTPRR correlate with cancer.

3.4 Extracellular Transfer of *miR-100* and *miR-125b*
Experiments proposing extracellular miRNA transfer can generate false positives if the recipient cell lines express the same miRNAs being tested (Gruner and McManus, 2021). Our knockout cell lines provide an ideal control to test for extracellular transfer of *miR-100* and *miR-125b* because Δ*miR-100/miR-125b* cells do not express either miRNA. As donor cells, we used CC-CR cells because they not only express high levels of *miR-100* and *miR-125b*, they also secrete EVs enriched in both miRNAs (**Fig. 5A**). Thus, we set up Transwell assays with CC-CR

donor cells and Δ*miR-100/miR-125b* recipient cells and co-cultured the cells for 5 days before
isolation of RNA from the recipient cells. Compared to control transfer experiments in which
Δ*miR-100/miR-125b* cells were used as both the donor and recipient cells, we observed dramatic
increases in the levels of *miR-100* and *miR-125b* in Δ*miR-100/miR-125b* recipient cells (~150-
and 60-fold, respectively) when CC-CR cells were used as the donor (**Fig. 5B**).

To determine whether the transfer of *miR-100* and *miR-125b* from CC-CR cells to Δ*miR-100/miR-125b* recipient cells results in functional miRNA transfer, we used Transwell co-culture
assays in which the Δ*miR-100/miR-125b* recipient cells were transfected with vectors in which
the firefly luciferase open reading frame was fused to the 3'-UTRs from either CGN or PTPRR
(**Fig. 5C**). Compared to controls, we observed a significant reduction in luciferase levels when
CC-CR cells were the donor cells and Δ*miR-100/miR-125b* were the recipient cells (**Fig. 5D**).
These data demonstrate that *miR-100 and miR-125b* can be functionally transferred between cells
to silence reporter mRNAs.

As an additional test of functional extracellular transfer of *miR-100 and miR-125b,* we
next tested whether we could observe silencing of the endogenous CGN gene in recipient cells
by immunofluorescence staining. Cingulin connects tight junctions to the cytoskeleton and
localizes to apical junctions joining adjacent epithelial cells (Zhang et al., 2021, Citi et al., 1988).
In cells that express low to undetectable levels of *miR-100 or miR-125b* (CC and Δ*miR-100/miR-125b* cells), CGN was readily detected and localized to cell-cell junctions (**Fig. 6A**).
Immunofluorescent signals were more readily visualized in CC cells compared to Δ*miR-100/miR-125b* cells, potentially due to regulation of other factors by *miR-100 or miR-125b* that
also seem to affect overall morphology, as indicated by differences in F-actin staining and
changes in CGN localization. Nevertheless, in cells expressing high levels of *miR-100* and *miR-125b* (CC-CR cells), CGN detection was dramatically reduced, and at least partially reversed by
deletion of *miR-100* and *miR-125b* (**Fig. 6A**). These data support targeting of CGN by *miR-100*
and *miR-125b.*

With baseline levels of CGN immunofluorescence in hand, we next used Transwell co-culture experiments with CC-CR donor cells and Δ*miR-100/miR-125b* recipient cells and
performed immunofluorescent staining of CGN in recipient cells. With Δ*miR-100/miR-125b*
donor cells, we were able to detect CGN in recipient cells (**Fig. 6B**) at levels similar to those
observed in **Fig. 6A**. However, when CC-CR cells were used as the donor cells, CGN was
undetectable in Δ*miR-100/miR-125b* recipient cells when measured by immunofluorescence
(**Fig. 6B**). These results provide strong evidence that CGN protein levels are downregulated by
functional transfer of *miR-100* and *miR-125b*.

3.5 *miR-100* and *miR-125b* Contribute to Enhanced 3D Growth and Invasiveness
Gene Ontology analysis of targets of miR-100 and miR-125b showed enrichment in
genes associated with cell migration and motility (**Fig. 2C**). This is consistent with
downregulation of CGN which leads to increased invasiveness, metastatic potential, and EMT
(Guillemot and Citi, 2006, Mangan et al., 2016, Schossleitner et al., 2016, Zhang et al., 2021).
To test the effect of *miR-100* and *miR-125b* and targeting of CGN on 3D growth, we cultured
CC, CC-CR and Δ*miR-100/miR-125b* cells under spheroid growth conditions (Carey et al., 2013)
in low adherence plates for 3 days before transfer to growth in Type I collagen. Under these

conditions at both day 0 and day 5 in collagen, the morphology of Δ*miR-100/miR-125b* colonies
was more similar to CC colonies than to colonies derived from parental CC-CR cells (**Fig. 7A**;
Supp. Movies 1-3). At day 5, we also observed significant differences in surface dynamics
when comparing colonies derived from the three lines **(Supp. Movies 1-3)**. Increased
projections emanating from the edges of the colonies and protruding into collagen were observed
with CC-CR cells, whereas far fewer projections were observed in colonies derived from CC
cells grown under the same conditions. As with overall morphology, Δ*miR-100/miR-125b*
colony dynamics and projections were more similar to that observed in CC colonies than CC-CR
colonies **(Supp. Movies 1-3).**

Quantification of overall spheroid growth revealed that colonies derived from CC-CR
cells showed a significant increase in total spheroid area when compared to colonies derived
from CC or Δ*miR-100/miR-125b* cells (**Fig. 7B**). We also quantified the number and size of
projections into the collagen as a measure of invasiveness by determining the increase in
projection surface area after growth in collagen for five days. CC-CR colonies showed a
significantly greater increase in projection area compared to CC and Δ*miR-100/miR-125b*
colonies (**Fig. 7C**). While invasiveness is likely due to multiple changes, the decreased
projections we observe in CC and Δ*miR-100/miR-125b* colonies are consistent with increased
expression of CGN due to lower levels of *miR-100* and *miR-125b*.

4 Discussion

Several reports have supported the hypothesis that miRNAs can transfer from one cell to
another via extracellular mechanisms (Mittelbrunn et al., 2011, Valadi et al., 2007, Skog et al.,
2008, Boelens et al., 2014, Squadrito et al., 2014, Cha et al., 2015, Barman et al., 2022). While
these reports show that miRNA can be transferred between cells, the recipient cells often express
the same miRNAs, potentially complicating interpretation of target mRNA silencing in recipient
cells (Gruner and McManus, 2021). Silencing under these conditions could be due to an
unexpected increased expression of endogenous miRNAs as opposed to functional miRNA
transfer. Our knockout cell lines provide an ideal control to obviate such concerns and
demonstrate functional miRNA transfer. Using donor cells that express high levels of *miR-100*
and *miR-125b* and recipient cells with targeted deletions of these genes, we identified mRNA
targets and demonstrated functional transfer of *miR-100* and *miR-125b* between cells to silence
both reporter constructs and endogenous mRNA transcripts in recipient cells. Decreased
expression of CGN in recipient cells is especially important because it validates silencing
without the use of reporter constructs.

4.1 Tumor Microenvironment Implications

Invasiveness and metastasis are indicative of differing stages of colorectal cancer and can
also be indicative of patient prognosis (Hanahan and Weinberg, 2011, Zeineddine et al., 2023).
EMT is a pivotal stage of metastasis involving the breakdown of cell-cell adhesions (Pretzsch et
al., 2019). Located on apical regions of adjacent epithelial cells, tight junctions play a crucial
role in cell-cell adhesion and tissue integrity. Disruption of tight junctions can allow cells to
become invasive and metastatic (Martin, 2014). CGN serves as a tether between the
cytoskeleton and tight junction components, interacting with ZO-1 at apical junctions (Zihni et
al., 2016, Citi et al., 1988). Loss of CGN in epithelial cells has been shown to disrupt tight
junction formation and cell polarity (Mangan et al., 2016, Guillemot and Citi, 2006,
Schossleitner et al., 2016). Coupled to our finding that *miR-100* and *miR-125b* are enriched in

EVs from CRC cells and that *miR-100* and *miR-125b* can be transferred between cells, our data support a model whereby cell-cell communication within the tumor microenvironment can alter tight junctions leading to altered growth and increased metastatic potential in recipient cells. For PTPRR, transfer of *miR-100* and *miR-125b* leading to decreased levels of PTPRR would be expected to activate Wnt signaling since it is a negative regulator of the Wnt pathway (Wang et al., 2019). This is consistent with activation of Wnt signaling in aggressive CC-CR cells (Lu et al., 2017).

4.2 Tumor Growth and Invasiveness

Beyond demonstrating miRNA transfer between cells, our knockout lines provided a model to test the effects of 3D growth in the presence and absence of *miR-100* and *miR-125b*. In 3D growth in collagen, CC cells form hollow cysts with a central lumen lined by polarized cells whereas CC-CR cells grow into the central lumen with disorganized, solid colonies (Lu et al., 2017). A prior deletion of exon 4 within the MIR100HG locus supported the idea that MIR100HG contributes to both cetuximab resistance and metastasis (Liu et al., 2022). Our precise deletions of *miR-100* and *miR-125b* allowed us to test the effect of these miRNAs on 3D growth and showed that loss of these miRNAs results in slower overall growth with decreased edge dynamics and fewer invasive projections. Since expression of MIR100HG is unaffected in our knockout lines, our data agree with previous results that *MIR100HG* can induce EMT, but also that *miR-100* and *miR-125b* play a similar role through downregulation of CGN (Zhang et al., 2021). The data are also consistent with our GO analysis of the targets of *miR-100* and *miR-125b* being enriched in cell motility and migration (**Fig 2C**).

In summary, our data support findings that increased expression of *miR-100* and *miR-125b* in cancer promotes disruption of tight junctions, activation of EMT, and increased metastatic potential. Given that these miRNAs are abundant in EVs secreted from CRC cells, the data support both cell-autonomous and non cell-autonomous roles for *miR-100* and *miR-125b*.

5 Acknowledgements

The authors would like to thank all members of the Coffey, Weaver, and Patton labs for advice and suggestions.

6 Funding

This work was supported by PO1 CA229123 to RJC, AMW, KCV, QL, and JGP.

7 Author Contributions

HMN, SQ, SC, NL, SS, LT, MS, LG, and KCC generated all data in the manuscript. KCV, QL, JLF, AMW, MR, and RJC provided scientific advice and suggestions and edited the manuscript. HMN, SQ, JLF, MR and JGP wrote the paper.

8 References

AEBI, M., HORNIG, H., PADGETT, R. A., REISER, J. & WEISSMAN, C. 1986. Sequence requirements for splicing of higher eukaryotic nuclear pre-mRNA. *Cell,* 47**,** 555-565.

ALLEN, R. M., ZHAO, S., RAMIREZ SOLANO, M. A., ZHU, W., MICHELL, D. L., WANG, Y., SHYR, Y., SETHUPATHY, P., LINTON, M. F., GRAF, G. A., SHENG, Q. & VICKERS, K. C. 2018. Bioinformatic analysis of endogenous and exogenous small RNAs on lipoproteins. *J Extracell Vesicles,* 7**,** 1506198.

BARMAN, B., SUNG, B. H., KRYSTOFIAK, E., PING, J., RAMIREZ, M., MILLIS, B., ALLEN, R., PRASAD, N., CHETYRKIN, S., CALCUTT, M. W., VICKERS, K., PATTON, J. G., LIU, Q. & WEAVER, A. M. 2022. VAP-A and its binding partner CERT drive biogenesis of RNA-containing extracellular vesicles at ER membrane contact sites. *Dev Cell,* 57**,** 974-994 e8.

BARTEL, D. P. 2009. MicroRNAs: target recognition and regulatory functions. *Cell,* 136**,** 215-33.

BOELENS, M. C., WU, T. J., NABET, B. Y., XU, B., QIU, Y., YOON, T., AZZAM, D. J., TWYMAN-SAINT VICTOR, C., WIEMANN, B. Z., ISHWARAN, H., TER BRUGGE, P. J., JONKERS, J., SLINGERLAND, J. & MINN, A. J. 2014. Exosome transfer from stromal to breast cancer cells regulates therapy resistance pathways. *Cell,* 159**,** 499-513.

CAREY, S. P., STARCHENKO, A., MCGREGOR, A. L. & REINHART-KING, C. A. 2013. Leading malignant cells initiate collective epithelial cell invasion in a three-dimensional heterotypic tumor spheroid model. *Clin Exp Metastasis,* 30**,** 615-30.

CENIK, E. S. & ZAMORE, P. D. 2011. Argonaute proteins. *Curr Biol,* 21**,** R446-9.

CHA, D. J., FRANKLIN, J. L., DOU, Y., LIU, Q., HIGGINBOTHAM, J. N., BECKLER, M. D., WEAVER, A. M., VICKERS, K., PRASAD, N., LEVY, S., ZHANG, B., COFFEY, R. J. & PATTON, J. G. 2015. KRAS-dependent sorting of miRNA to exosomes. *eLife,* 4**,** e07197.

CITI, S., SABANAY, H., JAKES, R., GEIGER, B. & KENDRICK-JONES, J. 1988. Cingulin, a new peripheral component of tight junctions. *Nature,* 333**,** 272-276.

COUCH, Y., BUZAS, E. I., DI VIZIO, D., GHO, Y. S., HARRISON, P., HILL, A. F., LOTVALL, J., RAPOSO, G., STAHL, P. D., THERY, C., WITWER, K. W. & CARTER, D. R. F. 2021. A brief history of nearly EV-erything - The rise and rise of extracellular vesicles. *J Extracell Vesicles,* 10**,** e12144.

DIXSON, A. C., DAWSON, T. R., DI VIZIO, D. & WEAVER, A. M. 2023. Context-specific regulation of extracellular vesicle biogenesis and cargo selection. *Nat Rev Mol Cell Biol*.

DOBIN, A., DAVIS, C. A., SCHLESINGER, F., DRENKOW, J., ZALESKI, C., JHA, S., BATUT, P., CHAISSON, M. & GINGERAS, T. R. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics,* 29**,** 15-21.

EMMRICH, S., STRELTSOV, A., SCHMIDT, F., THANGAPANDI, V. R., REINHARDT, D. & KLUSMANN, J. H. 2014. LincRNAs MONC and MIR100HG act as oncogenes in acute megakaryoblastic leukemia. *Mol Cancer,* 13**,** 171.

FARAZI, T. A., JURANEK, S. A. & TUSCHL, T. 2008. The growing catalog of small RNAs and their association with distinct Argonaute/Piwi family members. *Development,* 135**,** 1201-14.

GE, S. X., JUNG, D. & YAO, R. 2020. ShinyGO: a graphical gene-set enrichment tool for animals and plants. *Bioinformatics,* 36**,** 2628-2629.

GRUNER, H. N. & MCMANUS, M. T. 2021. Examining the evidence for extracellular RNA function in mammals. *Nat Rev Genet,* 22**,** 448-458.

GUILLEMOT, L. & CITI, S. 2006. Cingulin regulates claudin-2 expression and cell proliferation
through the small GTPase RhoA. *Mol Biol Cell,* 17**,** 3569-77.
HANAHAN, D. & WEINBERG, R. A. 2011. Hallmarks of cancer: the next generation. *Cell,* 144**,**
646-74.
HE, L. & HANNON, G. J. 2004. MicroRNAs: small RNAs with a big role in gene regulation. *Nat Rev*
Genet, 5**,** 522-31.
JEPPESEN, D. K., FENIX, A. M., FRANKLIN, J. L., HIGGINBOTHAM, J. N., ZHANG, Q., ZIMMERMAN,
L. J., LIEBLER, D. C., PING, J., LIU, Q., EVANS, R., FISSELL, W. H., PATTON, J. G., ROME, L.
H., BURNETTE, D. T. & COFFEY, R. J. 2019. Reassessment of Exosome Composition. *Cell,*
177**,** 428-445 e18.
KABADI, A. M., OUSTEROUT, D. G., HILTON, I. B. & GERSBACH, C. A. 2014. Multiplex
CRISPR/Cas9-based genome engineering from a single lentiviral vector. *Nucleic Acids*
Res, 42**,** e147.
KROL, J., LOEDIGE, I. & FILIPOWICZ, W. 2010. The widespread regulation of microRNA
biogenesis, function and decay. *Nat Rev Genet,* 11**,** 597-610.
LANGMEAD, B. & SALZBERG, S. L. 2012. Fast gapped-read alignment with Bowtie 2. *Nat*
Methods, 9**,** 357-9.
LI, C., GAO, Y., ZHANG, K., CHEN, J., HAN, S., FENG, B., WANG, R. & CHEN, L. 2015. Multiple
Roles of MicroRNA-100 in Human Cancer and its Therapeutic Potential. *Cell Physiol*
Biochem, 37**,** 2143-59.
LI, N., FLYNT, A., KIM, H. R., SOLNICA-KREZEL, L. & PATTON, J. 2008. Dispatched Homolog 2 is
targeted by miR-214 through a combination of three weak microRNA recognition sites.
Nucleic Acids Res, 36**,** 4277-85.
LIAO, Y., SMYTH, G. K. & SHI, W. 2014. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics,* 30**,** 923-30.
LIU, H., LI, D., SUN, L., QIN, H., FAN, A., MENG, L., GRAVES-DEAL, R., GLASS, S. E., FRANKLIN, J. L.,
LIU, Q., WANG, J., YEATMAN, T. J., GUO, H., ZONG, H., JIN, S., CHEN, Z., DENG, T., FANG,
Y., LI, C., KARIJOLICH, J., PATTON, J. G., WANG, X., NIE, Y., FAN, D., COFFEY, R. J., ZHAO,
X. & LU, Y. 2022. Interaction of lncRNA MIR100HG with hnRNPA2B1 facilitates m(6)A-
dependent stabilization of TCF7L2 mRNA and colorectal cancer progression. *Mol Cancer,*
21**,** 74.
LIU, J., CARMELL, M. A., RIVAS, F. V., MARSDEN, C. G., THOMSON, J. M., SONG, J. J., HAMMOND,
S. M., JOSHUA-TOR, L. & HANNON, G. J. 2004. Argonaute2 is the catalytic engine of
mammalian RNAi. *Science,* 305**,** 1437-41.
LOVE, M. I., HUBER, W. & ANDERS, S. 2014. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol,* 15**,** 550.
LU, Y., ZHAO, X., LIU, Q., LI, C., GRAVES-DEAL, R., CAO, Z., SINGH, B., FRANKLIN, J. L., WANG, J.,
HU, H., YANG, M., YEATMAN, T. J., LEE, E., SAITO-DIAZ, K., HINGER, S., PATTON, J. G.,
CHUNG, C. H., EMMRICH, S., KLUSMANN, J.-H., FAN, D. & COFFEY, R. J. 2017. lncRNA
MIR100HG-derived miR-100 and miR-125b mediate Cetuximab resistance via Wnt/beta
catenin signaling. *Nature Medicine,* 23**,** 1331-1341.
MANGAN, A. J., SIETSEMA, D. V., LI, D., MOORE, J. K., CITI, S. & PREKERIS, R. 2016. Cingulin and
actin mediate midbody-dependent apical lumen formation during polarization of
epithelial cells. *Nat Commun,* 7**,** 12426.

MARTIN, M. 2011. Cutadapt removes adapator sequences from high-throughput sequencing
reads. *EMBnet.journal,* 17**,** 10-12.
MARTIN, T. A. 2014. The role of tight junctions in cancer metastasis. *Semin Cell Dev Biol,* 36**,**
224-31.
MITTELBRUNN, M., GUTIERREZ-VAZQUEZ, C., VILLARROYA-BELTRI, C., GONZALEZ, S., SANCHEZ-
CABO, F., GONZALEZ, M. A., BERNAD, A. & SANCHEZ-MADRID, F. 2011. Unidirectional
transfer of microRNA-loaded exosomes from T cells to antigen-presenting cells. *Nat*
Commun, 2**,** 282.
MÜLLER, M., FAZI, F. & CIAUDO, C. 2020. Argonaute Proteins: From Structure to Function in
Development and Pathological Cell Fate Determination. *Frontiers in Cell and*
Developmental Biology, 7.
O'BRIEN, K., BREYNE, K., UGHETTO, S., LAURENT, L. C. & BREAKEFIELD, X. O. 2020. RNA delivery
by extracellular vesicles in mammalian cells and its applications. *Nat Rev Mol Cell Biol,*
21**,** 585-606.
OTTAVIANI, S., STEBBING, J., FRAMPTON, A. E., ZAGORAC, S., KRELL, J., DE GIORGIO, A.,
TRABULO, S. M., NGUYEN, V. T. M., MAGNANI, L., FENG, H., GIOVANNETTI, E., FUNEL,
N., GRESS, T. M., JIAO, L. R., LOMBARDO, Y., LEMOINE, N. R., HEESCHEN, C. &
CASTELLANO, L. 2018. TGF-beta induces miR-100 and miR-125b but blocks let-7a
through LIN28B controlling PDAC progression. *Nat Commun,* 9**,** 1845.
PENG, B., THENG, P. Y. & LE, M. T. N. 2021. Essential functions of miR-125b in cancer. *Cell Prolif,*
54**,** e12913.
PENG, Y. & CROCE, C. M. 2016. The role of MicroRNAs in human cancer. *Signal Transduction*
and Targeted Therapy, 1**,** 15004.
PRETZSCH, E., BÖSCH, F., NEUMANN, J., GANSCHOW, P., BAZHIN, A., GUBA, M., WERNER, J. &
ANGELE, M. 2019. Mechanisms of Metastasis in Colorectal Cancer and Metastatic
Organotropism: Hematogenous versus Peritoneal Spread. *J Oncol,* 2019**,** 7407190.
QIN, C., HUANG, R. Y. & WANG, Z. X. 2015. Potential role of miR-100 in cancer diagnosis,
prognosis, and therapy. *Tumour Biol,* 36**,** 1403-9.
RIOLO, G., CANTARA, S., MARZOCCHI, C. & RICCI, C. 2021. miRNA Targets: From Prediction Tools
to Experimental Validation. *Methods and Protocols,* 4.
SCHOSSLEITNER, K., RAUSCHER, S., GROGER, M., FRIEDL, H. P., FINSTERWALDER, R.,
HABERTHEUER, A., SIBILIA, M., BROSTJAN, C., FODINGER, D., CITI, S. & PETZELBAUER, P.
2016. Evidence That Cingulin Regulates Endothelial Barrier Function In Vitro and In Vivo.
Arterioscler Thromb Vasc Biol, 36**,** 647-54.
SKOG, J., WÜRDINGER, T., RIJN, S. V., MEIJER, D. H., GAINCHE, L., JR, W. T. C., CARTER, B. S.,
KRICHEVSKY, A. M. & BREAKEFIELD, X. O. 2008. Glioblastoma microvesicles transport
RNA and proteins that promote tumour growth and provide diagnostic biomarkers.
Nature Cell Biology, 10**,** 1470.
SQUADRITO, M. L., BAER, C., BURDET, F., MADERNA, C., GILFILLAN, G. D., LYLE, R., IBBERSON,
M. & DE PALMA, M. 2014. Endogenous RNAs Modulate MicroRNA Sorting to Exosomes
and Transfer to Acceptor Cells. *Cell Rep,* 8**,** 1432-46.
SUBRAMANIAN, A., TAMAYO, P., MOOTHA, V. K., MUKHERJEE, S., EBERT, B. L., GILLETTE, M. A.,
PAULOVICH, A., POMEROY, S. L., GOLUB, T. R., LANDER, E. S. & MESIROV, J. P. 2005.

Gene set enrichment analysis: a knowledge-based approach for interpreting genome-
wide expression profiles. *Proc Natl Acad Sci U S A,* 102**,** 15545-50.
VALADI, H., EKSTROM, K., BOSSIOS, A., SJOSTRAND, M., LEE, J. J. & LOTVALL, J. O. 2007.
Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic
exchange between cells. *Nat Cell Biol,* 9**,** 654-9.
VALLACCHI, V. & RODOLFO, M. 2018. miR-100 and miR-125b regulate epithelial-mesenchymal
transition and drug resistance in tumors. *Non-coding RNA Investigation,* 2**,** 57-61.
WANG, J., VASAIKAR, S., SHI, Z., GREER, M. & ZHANG, B. 2017. WebGestalt 2017: a more
comprehensive, powerful, flexible and interactive gene set enrichment analysis toolkit.
Nucleic Acids Res, 45**,** W130-W137.
WANG, Y., CAO, J., LIU, W., ZHANG, J., WANG, Z., ZHANG, Y., HOU, L., CHEN, S., HAO, P., ZHANG,
L., ZHUANG, M., YU, Y., LI, D. & FAN, G. 2019. Protein tyrosine phosphatase receptor
type R (PTPRR) antagonizes the Wnt signaling pathway in ovarian cancer by
dephosphorylating and inactivating beta-catenin. *J Biol Chem,* 294**,** 18306-18323.
WANG, Y., ZENG, G. & JIANG, Y. 2020. The Emerging Roles of miR-125b in Cancers. *Cancer*
Manag Res, 12**,** 1079-1088.
ZEINEDDINE, F. A., ZEINEDDINE, M. A., YOUSEF, A., GU, Y., CHOWDHURY, S., DASARI, A., HUEY,
R. W., JOHNSON, B., KEE, B., LEE, M. S., MORELLI, M. P., MORRIS, V. K., OVERMAN, M. J.,
PARSEGHIAN, C., RAGHAV, K., WILLIS, J., WOLFF, R. A., KAWAGUCHI, Y., VAUTHEY, J.-N.,
SUN, R., KOPETZ, S. & SHEN, J. P. 2023. Survival improvement for patients with
metastatic colorectal cancer over twenty years. *npj Precision Oncology,* 7**,** 16.
ZHANG, X., LI, T., HAN, Y. N., GE, M., WANG, P., SUN, L., LIU, H., CAO, T., NIE, Y., FAN, D., GUO,
H., WU, K., ZHAO, X. & LU, Y. 2021. miR-125b Promotes Colorectal Cancer Migration and
Invasion by Dual-Targeting CFTR and CGN. *Cancers (Basel),* 13.
ZHAO, S., GUO, Y., SHENG, Q. & SHYR, Y. 2014. Advanced heat map and clustering analysis using
heatmap3. *Biomed Res Int,* 2014**,** 986048.
ZIHNI, C., MILLS, C., MATTER, K. & BALDA, M. S. 2016. Tight junctions: from simple barriers to
multifunctional molecular gates. *Nature Reviews Molecular Cell Biology,* 17**,** 564-580.
575

9   Figure Legends

9.1   Figure 1. Expression levels of *miR-100* and *miR-125b* in CC, CC-CR and Δ*miR-100/miR-125b* cell lines. (A) One of the transcription start sites within the 330kb MIR100HG locus encodes a 115kb transcript that is spliced (black exons) to generate a 3kb lncRNA (*MIR100HG*). *miR-100, miR-125b, let-7a* and the BLID protein are encoded in the third intron of *MIR100HG*. In CC-CR cells, *MIR100HG*, *miR-100* and *miR-125b* are highly overexpressed but *let-7a* is not (Lu et al., 2017). (B) CC-CR cells are a line of cetuximab-resistant cells, derived from cetuximab-sensitive CC cells (Lu et al., 2017). *miR-100* and *miR-125b* are dramatically overexpressed in CC-CR cells, as measured by RT-qPCR. (C) RT-qPCR analysis of *miR-100* and *miR-125b* in CC-CR, Δ*miR-100*, Δ*miR-125b*, Δ*miR-100/miR-125b* cells. Fold changes were determined using the ΔΔC(t) method. ** indicates $p<0.005$ and **** indicates $p<0.0001$.

9.2   Figure 2. Identification of targets for *miR-100* and *miR-125b* in colorectal cancer cells. (A) Differentially expressed genes were identified by mining RNAseq data comparing CC, CC-CR, Δ*miR-100*, Δ*miR-125b*, Δ*miR-100/miR-125b* cells. Venn diagram showing overlap between genes upregulated in CC and knockout cell lines, mRNA targets of *miR-100* and *miR-125b* using three different prediction algorithms, and association with AGO2 in CC-CR cells. (B) Heatmaps were generated from the differential RNAseq data in Fig. 2A to identify potential targets upregulated in CC cells (>2 fold) and knockout cells (>1.5 fold). (C) Gene Ontology analysis of the targets of *miR-100* and miR-125b show enrichment in genes associated with cell migration and motility. Graph was created using ShinyGO (Ge et al., 2020).

9.3   Figure 3. Silencing of reporter constructs containing target 3'-UTRs for *miR-100* and *miR-125b*. (A) Reporter constructs were generated in which the firefly luciferase open reading frame was fused to either plasmid 3'-UTR sequences (empty vector), 3'-UTR sequences containing three perfect binding sites for *miR-100 or miR-125b* (100/125 vector), or 3'-UTR sequences from candidate mRNA targets of *miR-100 or miR-125b* (experimental vector). (B) Quantification of relative luciferase levels after transfection of the indicated constructs into CC-CR cells or Δ*miR-100/miR-125b* cells. Relative firefly luciferase levels were calculated by normalizing to a co-transfected internal control vector expressing Renilla luciferase. * indicates $p<0.05$, ** indicates $p< 0.01$, *** indicates $p<0.001$, **** indicates $p<0.0001$.

9.4   Figure 4. Expression of CGN and PTPRR are increased in cancer. (A) Boxplot of CGN (A) and PTPRR (B) expression across 72 paired tissues. Red indicates cancer samples, blue indicates normal samples.

9.6   Figure 5. Functional transfer of *miR-100* and *miR-125b*. (A) RNA was collected from extracellular vesicles (EVs) from CC and CC-CR cells and the levels of *miR-100* and *miR-125b* were quantified by RT-qPCR. The left panel shows the fold-change enrichment of these miRNAs in EVs compared to cellular levels. The right panel shows the fold-change in EV levels of *miR-100* and *miR-125b* when comparing CC-CR and CC cells. (B) Transwell co-culture experiments were performed with Δ*miR-100/miR-125b* recipient cells and the donor cells as indicated on the X axis. RNA was collected from the recipient cells after 5 days in culture and fold-changes in *miR-100* and *miR-125b* levels were determined by RT-qPCR. (C) Schematic of Transwell co-culture experiments to test silencing of CGN and PTPRR luciferase reporter constructs in Δ*miR-100/miR125b* recipient cells. (D) Δ*miR-100/miR-125b* recipient cells were

transfected with luciferase reporters (x-axis) and co-cultured with either Δ*miR-100/miR-125b*
(grey) or CC-CR (black) donor cells. Relative luciferase values were calculated in recipient
cells, normalizing as in Fig. 3. *indicates $p<0.05$, **indicates $p< 0.01$, ***indicates $p<0.001$,
**** indicates $p<0.0001$; ns is not significant.

9.7   Figure 6. Transfer of *miR-100* and *miR-125b* decreases CGN immunofluorescence in
recipient cells. (A) The indicated cells were fixed and immunostained with DAPI to stain nuclei
(blue), antibodies against F-actin (green), or antibodies against CGN (white). Merged images are
shown at the far right. (B) Transwell co-culture experiments were performed with Δ*miR-*
100/miR-125b recipient cells and the indicated donor cells. After 5 days, recipient cells were
immunostained as in Fig. 6A. Scale bar indicates 10mm.

9.8   Figure 7. *miR-100* and *miR-125b* contribute to enhanced 3D growth and invasiveness.
(A) The indicated cell types were plated in low adherent, spheroid cultures for 3 days before
transfer to growth in Type I collagen and live imaging over 5 days. (B) The increase in colony
area was determined by measuring the surface area of spheroids from day 0 to day 5. Surface
area tracing was performed using ImageJ, shown in yellow. (C) Increases in area due to colony
projections into the collagen (invasiveness) were calculated by measuring the total surface area
including projections (yellow) and subtracting the surface area of the main spheroid (red).
**indicates $p< 0.01$, ***indicates $p<0.001$; ns is not significant. Scale bar indicates 500mm.

A

	CC	CC-CR	ΔmiR-100/miR-125b
Day 0			
Day 5			

B

Area Increase (%)

CC-CR *** CC *** ΔmiR100/Δmir125b-1 ** ΔmiR100/Δmir125b-2

C

Projection area increase (x1000 um2)

CC-CR *** CC ns ΔmiR-100/miR-125b ***