The transgenerational consequences of paternal social isolation and predation exposure in

threespined sticklebacks

Jennifer K. Hellmann* [1,2], Michaela M. Rogers [1,2]

[1]Department of Biology, University of Dayton, Dayton, Ohio, USA, 45469

[2]Department of Evolution, Ecology, and Organismal Biology, The Ohio State University,

Columbus, Ohio, USA, 43210

*Corresponding author: Hellmann.13@osu.edu
11

Abstract

1. Although parents routinely encounter stress in the ecological environment that can affect offspring development (transgenerational plasticity: TGP), nearly all organisms interact with conspecifics that can either independently induce parental stress or can alter how parents respond to ecological stressors.

2. During social buffering, the presence of conspecifics can reduce the response to or increase the speed of recovery from a stressor. This may have cascading effects on offspring if the presence of conspecifics can mitigate parental responses to ecological stress in ways that blunt transmission of stress-induced transgenerational effects.

3. Here, we simultaneously manipulated both paternal social isolation and experience with predation risk prior to fertilization in threespined stickleback (*Gasterosteus aculeatus*). If social buffering mitigates TGP induced by paternal exposure to predation risk, then we expected fewer phenotypic effects in offspring when fathers are exposed to predation risk with a conspecific with present compared to fathers exposed to predation risk while isolated.

4. Offspring of predator-exposed fathers showed reduced anxiety-like behavior. Fathers who were socially isolated had offspring that were captured faster by a live predator, suggesting that social isolation may be a stressor that has maladaptive consequences for offspring. Further, we also found evidence that the presence of a conspecific buffered fathers against the effects of predation risk: offspring of predator-exposed fathers tended to be captured faster by the predator, especially when fathers were isolated compared to when they had neighbors.

5. Our results suggest that socially-induced stress is an important, yet underappreciated,

mediator of TGP and can elicit transgenerational effects even in species that do not form

permanent social groups. Future studies should therefore consider the broader social

environment when predicting both within and trans-generational responses to ecological

change.

Key words: *Gasterosteus aculeatus*, paternal effects, phenotypic plasticity, predation risk, social

buffering

Introduction

Organisms encounter a wide variety of threats and stressors in their environment that can

push individuals beyond their homeostatic limits and induce physiological and behavioral

changes that allow them to cope with those stressors (Romero, Dickens & Cyr 2009). It is well

known that individuals differ drastically in how they respond to these environmental stressors:

the same external stressor can induce strong responses in some individuals and weak response in

others (Koolhaas *et al.* 1999; Cockrem 2013). Although some of this variation can be attributed

to traits of the organism itself (e.g., personality (Cockrem 2013), sex (DeVries, Glasper &

Detillion 2003)), some of this variation is environmentally dependent. In particular, there is

ample evidence that the social environment alters how animals cope with ecological stress,

influencing both the perception of and response to ecological cues (DeVries, Glasper & Detillion

2003; Brandl, Pruessner & Farine 2022). In many cases, the presence of conspecifics can reduce

physiological and behavioral responses to stress via social buffering, which occurs when the

presence of a conspecific reduces the response to or increases the speed of recovery from an

external stressor (Kikusui, Winslow & Mori 2006; Gilmour & Bard 2022). For example, socially

isolated mice showed greater immune impairment and a higher basal corticosterone increase in

response to a mild stress compared to group housed mice (Bartolomucci *et al.* 2003). Similarly,

in humans, anxiety and negative emotions induced by a social stress test were attenuated when

women had social support (McQuaid *et al.* 2016). This social buffering may be an adaptive

response to minimize the transmission of stress among group members (Gilmour & Bard 2022),

which can occur when a stress response system (e.g., glucocorticoids) in one individual is

activated after interacting with another stressed individual (Brandl, Pruessner & Farine 2022).

Further, social buffering can even increase the limits of the physical conditions organisms are

willing to tolerate (Gilmour & Bard 2022), suggesting that sociality can mediate response to

ecological stressors.

While social buffering has been well studied in highly social mammals, there is mounting

evidence that social buffering of stress also occurs in less social species, including solitary

species and species with aggregations or schooling behavior (Gilmour & Bard 2022). For

example, in lake sturgeon, group-housed fish had significantly reduced endocrine and cellular

stress markers after exposure to heat stress compared to isolated individuals (Yusishen *et al.*

2020). This social buffering effect can even occur in small groups of two individuals (Faustino,

Tacão-Monteiro & Oliveira 2017; Schumann *et al.* 2023) or in response to mere visual cues of a

conspecific (da Costa *et al.* 2004; Faustino, Tacão-Monteiro & Oliveira 2017), suggesting that

these social buffering effects are robust and occur in a wide range of taxonomic groups with

differing levels of sociality.

While social buffering effects can have consequences for the individuals that directly

experience the stressor, social buffering in parents may also have consequences for offspring.

Transgenerational plasticity (TGP) occurs when the experiences of a previous generation alter

the phenotypes of future generations (Bell & Hellmann 2019). TGP can occur in response to a

variety of ecological stressors, including predation risk (Tariel, Plénet & Luquet 2020; MacLeod

et al. 2022), heat exposure (Donelson *et al.* 2016; Pessato, McKechnie & Mariette 2022),

immune challenges (Beemelmanns & Roth 2016; Moore, Kaletsky & Murphy 2019), and food

limitation (Bygren *et al.* 2014; Hagmayer *et al.* 2022). These ecological stressors induce changes

in parental state, including the activation of the hypothalamic-pituitary-adrenal axis (HPA axis),

that influence offspring development, such as stress responsiveness, growth, and behavior (Bell

& Hellmann 2019; Potticary & Duckworth 2020). Given that the magnitude of the stress

response in parents can be linked to the magnitude of the phenotypic changes in offspring
(Sheriff, Krebs & Boonstra 2010), social buffering may affect both the strength and frequency
with which parental experiences affect offspring traits. This is supported by a recent study in
humans which found that the adverse health outcomes differentiating offspring of healthy and
psychologically stressed mothers could be specifically linked to a lack of social support in
psychologically stressed mothers (Walsh *et al.* 2019). Similarly, in rats, offspring of socially-
isolated mothers showed cognitive impairments and increased anxiety-like behaviors (McDonald
et al. 2023). These effects may arise because conspecifics provide a buffering effect that abates
maternal response to external stress (e.g., brings glucocorticoids back to baseline quickly
(Kikusui, Winslow & Mori 2006)), thereby mitigating adverse outcomes for offspring (Séguin *et
al.* 1995; Nylen, O'Hara & Engeldinger 2013).

Although this previous work suggests the importance of the social environment in
mediating transgenerational effects for mothers, we know nothing about the transgenerational
effects of social isolation for fathers. This is problematic, as paternal experiences can affect
offspring phenotypes in magnitudes that are similar to maternal experiences (Bonduriansky &
Head 2007; Beemelmanns & Roth 2016; Gilad & Scharf 2019; Hellmann *et al.* 2020). Further,
because males and females differ in their susceptibility to stress (DeVries, Glasper & Detillion
2003) and the effects of paternal experiences often have different effects on offspring than
maternal experiences (Bonduriansky & Head 2007; Bonduriansky, Runagall-McNaull & Crean
2016; Bell & Hellmann 2019; Gilad & Scharf 2019; Hellmann *et al.* 2020), it is likely that the
transgenerational effects of paternal isolation may be different from maternal isolation. Thus,
understanding how the social environment mediates paternal effects can help us better predict the
extent to which paternal experiences influence offspring phenotypes and understand how we

might predict the effects of paternal stress if, for instance, social buffering during or after a

stressor mitigates or reverses changes in parental state that induce transgenerational effects.

We sought to understand the ways that the social environment alters transgenerational

consequences of paternal stress in threespined stickleback *(Gasterosteus aculeatus)*. There is

abundant evidence for predator-induced paternal effects in stickleback, which alter offspring

behavior, growth, gene expression, survival, and stress responsiveness (Hellmann *et al.* 2020;

Hellmann, Carlson & Bell 2020; Chen *et al.* 2021; Hellmann, Carlson & Bell 2021; Afseth *et al.*

2022). There is also previous evidence of social buffering effects in stickleback: fish that are

exposed to simulated predation risk while isolated had higher circulating levels of cortisol

compared to fish in a group (Mommer & Bell 2013) and pairs of sticklebacks co-regulate their

cortisol responses in stressful environments (Fürtbauer & Heistermann 2016). However, we have

very little understanding as to how the social environment might modulate parental responses to

ecological stress in ways that might have cascading effects on the adaptive nature and frequency

of transgenerational plasticity.

We used a full factorial design to compare the transgenerational consequences of

exposing fathers to predation risk while fathers were isolated or with another conspecific.

Offspring were generated via in-vitro fertilization to isolate paternal effects mediated via sperm

and/or seminal fluid. We then measured offspring behavioral traits relevant to predation defense,

including anti-anxiety behavior, shoaling behavior, and survival against a live predator. If social

buffering mitigates TGP induced by paternal exposure to predation risk, then we expect to see

fewer phenotypic effects in offspring when fathers are exposed to predation risk with a

conspecific with present compared to fathers exposed to predation risk while isolated.

Methods

Housing information and F0 exposures. In June 2021, we collected adult three-spined

stickleback *(Gasterosteus aculeatus)* from Putah Creek (CA, USA), a freshwater stream in

northern California, and shipped them to University of Dayton. We maintained the F0 generation

on a summer photoperiod schedule (16 L : 8 D) at $20° ± 1°C$ and fed them ad libitum daily with a

mix of frozen bloodworms (*Chironomus* spp.), brine shrimp (Artemia spp.) Mysis shrimp, and

Cyclop-eeze. We obtained 10-15 cm long rainbow trout *(Oncorhynchus mykiss)* from Freshwater

Farms of Ohio in June 2021. Putah Creek has piscivorous predators, including the prickly sculpin

(*Cottus asper*); while rainbow trout are present in Putah Creek, they are not present in the

isolated section of Putah Creek where we collected stickleback (Beaver Pond), suggesting that

rainbow trout is an environmentally relevant, but novel predator.

In August 2021, we moved males into experimental 37.9 L (53L x 33W x 25H cm) tanks

divided into 3 equal sections using removable barriers. Males in the neighbor treatment were

separated with clear barriers with 9, 3mm holes to allow for water flow between the sections.

Males in the isolated treatment were separated with permanent opaque barriers that did not allow

for water flow between sections. In both treatments, we placed one male stickleback in each of

the outer sections and left the middle section empty. Each male section contained gravel, one

fake plant, and nesting materials: a container of sand, a terra cotta pot for cover, and algae. The

middle section contained only gravel. Next to each nesting male tank, we placed an undivided

37.9L tank containing gravel and 2 sponge filters, which either contained one live rainbow trout

or was left empty depending on the predator treatment. All experimental tanks were separated

visually from each other with a removable opaque divider.

We exposed male stickleback to predator cues once a day for 2 hours, for 11-14

consecutive days depending on the availability of gravid females. We used a short stressor to

avoid potential habituation to predation risk (Dellinger *et al.* 2018). Further, by beginning the

treatment when males were transferred to a nesting arena, we sought to mimic the change in

predator regime that males may encounter as they move into a different habitat to nest. At a

random time of day between 0800 hrs and 2000 hrs, we turned off the bubblers in the tanks and

removed the opaque divider to reveal the adjacent tank (which was empty or contained a live

trout). At the same time, we added either 100mL of control RO water or the prepared predator

cue to each end compartment containing one male stickleback. We prepared predator cues using

methods consistent with Crane and Ferrari (2015). Briefly, we placed a trout in a 37.9 L tank

with clean, conditioned tap water proportional to the size of the trout (500mL per gram of fish).

We fed the trout one juvenile stickleback per day for four days and after 96hrs, we froze the tank

water into 200 mL aliquots.

In the control treatment, males saw an empty tank and received 100 mL of RO water. In

the predator treatment, males saw a live trout and received 100 mL of the prepared predator cue.

Every other day, we fed the trout one juvenile stickleback during the two hour exposure to

reinforce the visual cue. Importantly, despite holes between the sections, we found very limited

diffusion of water between the sections (tested using dyed water) when bubblers were turned off,

suggesting that these olfactory cues did not disperse throughout the tank during the predation

exposure and therefore, the neighbor treatment did not receive weaker olfactory cues or

predation risk than the isolated group. After two hours, we replaced the dividers and completed a

50% water change on all treatments. We removed males after the last day of exposure.

Generating Offspring. The day after the last exposure, we generated offspring by manually

squeezing the eggs out of a gravid, unexposed female and fertilizing them with sperm from the

testes of the dissected experimental male. We checked clutches for successful fertilization via

microscope and then placed into a small cup with a mesh bottom above an air bubbler in an

18.9L (38L x 25W x 21H) tank containing gravel and two fake plants, with each clutch housed

individually on a recirculating flow-through system. Once hatched, we fed offspring newly

hatched brine shrimp for two months and then gradually transitioned to frozen food (as above).

We switched offspring to a winter light schedule (8 L: 16 D) prior to starting behavioral assays.

We generated a total of n = 35 clutches: n = 9 of control fathers with a neighbor, n = 9 of

control, isolated fathers, n = 8 of predator-exposed fathers with a neighbor, and n = 9 of predator-

exposed, isolated fathers. By artificially fertilizing the eggs and incubating the embryos using an

air bubbler, we controlled for the possibility that predator-exposed or isolated fathers might be

less likely to successfully mate with females or parent offspring.

Offspring assays.

Shoaling assays. We used experimental 37.9L (53 x 33 x 24 cm high) tanks lined with gravel.

On one end of the tank, we positioned three small plastic plants; on the other end of the tank, we

created a small compartment for shoaling fish by inserting a clear plastic barrier (10 cm from the

edge) perforated with small holes to allow for flow of water and olfactory cues. The evening

before a day of trials, we placed five size-matched juveniles into the shoaling side compartment

of each tank to habituate overnight. These fish were unrelated and unfamiliar to the focal fish.

At the start of the shoaling trial, we placed an experimental offspring (focal fish) into the

open area in the middle of the tank and allowed the focal fish to habituate for 10 minutes. After

the habitation, we used BORIS to record total time spent shoaling, total time spent oriented to the

shoal while shoaling (the focal fish's head had to be pointing in the direction of at least one

member of the shoal), total time spent hiding in plants, and time spent in the gravel in the center

of the tank over a 10 minute period. We defined a focal fish to be shoaling or hiding in the plant

if it was within one body length of the shoal barrier or plants, respectively. Upon completion of

the assay, we gently removed the focal fish from the tank with a cup, weighed and measured

them, and then placed them into an individual holding tanks. We assayed and measured n = 242

fish: n = 61 of control fathers with a neighbor, n = 57 of control, isolated fathers, n = 66 of

predator-exposed fathers with a neighbor, and n = 58 of predator-exposed, isolated fathers.

Measuring anxiety/cautiousness. Scototaxis (light/dark preference) protocols have been

developed to test anti-anxiety/cautious behavior in fish (Maximino *et al.* 2010). On the day after

the shoaling assay, offspring were gently caught with a cup from their holding tank and placed in

a clear cylinder (10.5cm diameter) in the center of a half-black, half-white tank (51L x 28W x

19H cm). After a 5-minute acclimation period, we lifted the cylinder, and fish explored the tank

for 15 minutes, during which we measured the latency to enter the white section, total time in the

white section, and thigmotaxis behavior (time spent within a body length of the wall) within the

white section. After the 15-minute testing period, we removed the fish from the tank and placed

them back in their holding tank. We assayed n = 234 fish: n = 58 of control fathers with a

neighbor, n = 55 of control, isolated fathers, n = 64 of predator-exposed fathers with a neighbor,

and n = 57 of predator-exposed, isolated fathers.

Survival assay. 48-72 hours after the scototaxis assay, we conducted survival assays in groups of

two siblings (same paternal treatment and genetic family). We filled circular portable pools with

conditioned tap water (1m diameter, 13 cm water depth) and evenly placed 6 plants around each

arena. We placed an individually marked rainbow trout in each pool 24-hours before the first day

of assays. We fasted the trout for 48 hours prior to being placed in the pools and we fed them one

non-experimental juvenile stickleback after being placed in the pool to keep hunger levels

standardized across days. On each assay day, we used each trout for two assays, once in the

morning and once in the afternoon. For each assay, we placed two siblings in a clear cylinder

(20.3 cm diameter) in the middle of the arena. After acclimating for 20 minutes, we lifted the

cylinder and used a stopwatch to time how long it took the trout to capture one of the fish

(survival time). We had a total of n=91 trials: n = 29 of control fathers with a neighbor, n = 15 of

control, isolated fathers, n = 25 of predator-exposed fathers with a neighbor, and n = 22 of

predator-exposed, isolated fathers. We analyzed all trials where one of the fish was captured

within one hour (n=88 trials).

Statistical methods. To get an estimate of shoaling behavior, we used principal components

analysis (R package factoextra (Kassambara & Mundt 2017)) to combine several behaviors: time

spent hidden in the plant, time spent shoaling, time spent oriented to the shoal, and time spent in

the gravel in the center of the tank. We extracted an eigenvalue of 2.74 that captured 68.5% of

the variance in these behaviors; positive values indicate more social individuals who spent more

time shoaling (contribution: 33.4%) and oriented to the shoal (31.3%), and spent less time hiding

in plants (25.1%) and gravel (10.2%). This PC was used as the dependent variable in the model

described below (Gaussian distribution). For the scototaxis assay, latency to enter the white arena

was negatively correlated with time spent in the white section (Spearman rank correlation: ρ = -

0.42, p<0.001), so we analyzed variation in time spent in the white section of the tank, as well as

time spent performing thigmotaxis in the white section of the tank (both Poisson distribution).

We used the package MCMCglmm (Hadfield 2010) to analyze offspring traits. We ran

models with a weak prior on the variance (V=1, nu=0.002). We ran models for 200,000

iterations, with a burn-in of 3000 iterations and thin = 3. Models testing variation in shoaling and

scototaxis behavior included fixed effects of paternal predation exposure, paternal isolation, and

offspring standard length and random effects of clutch identity and observer identity. For the

thigmotaxis model, we also included a fixed effect of total time in the white section of the tank.

The survival model (time to capture: Poisson distribution) included fixed effects of paternal

predation exposure and paternal isolation, and random effects of clutch identity, observer

identity, and trout identity. The length model (Gaussian distribution) included fixed effects of

paternal predation exposure, paternal isolation, and offspring age, while the mass model

(Gaussian distribution) included fixed effects of paternal predation exposure, paternal isolation,

and offspring length; both models included random effects of clutch identity and observer

identity. For all models, we tested for interactions between paternal predation exposure and

paternal isolation and removed all non-significant interactions.

Animal welfare note. All methods were approved by Institutional Animal Care and Use

Committee of University of Dayton (protocol ID 021-05), including the use of live predators.

Results

We examined how paternal exposure to predation risk and paternal social isolation altered offspring behavior, size, and survival against live predators.

Anti-anxiety behavior. We found that offspring of predator-exposed fathers spent more time in the white section of the tank compared to offspring of control fathers (MCMC GLMM, 95% CI [0.04, 0.29], p = 0.01; Figure 1). After controlling for the total time in the white area, we found that offspring of predator-exposed fathers tended to do more thigmotaxis in the white section compared to offspring of control fathers ([-0.49, 42.5], p = 0.06). We found no effect of paternal isolation on time spent in the white section ([-0.20, 0.06], p = 0.30) or on thigmotaxis behavior ([-24.5, 18.5], p = 0.77).

Shoaling behavior. We found no change in offspring shoaling behavior based on either paternal isolation (MCMC GLMM: 95% CI [-0.54, 0.40], p = 0.77) or paternal predation exposure ([-0.60, 0.33], p = 0.59), although longer offspring spent more time shoaling than smaller offspring ([0.06, 0.24], p <0.001).

Offspring size. We also found no differences in length or mass based on either paternal isolation (length: 95% CI [-2.32, 1.21], p = 0.49; mass: [-0.01, 0.01], p = 0.90) or paternal predation exposure (length: [-1.15, 2.39], p = 0.49; mass: [-0.01, 0.01], p = 0.71).

Survival. We found that offspring of fathers who were isolated were captured significantly faster by the predator compared to offspring of fathers with neighbors (95% CI [-1.55, -0.09], p = 0.03;

Figure 2). Similarly, offspring of fathers who were predator-exposed tended to be captured

significantly faster by the predator compared to offspring of control fathers (95% CI [-1.38,

0.07], p = 0.07; Figure 2). We found some evidence for social buffering: considering paternal

treatment as a 4-level factor, we found a borderline significant reduction in survival time

between offspring of predator-exposed fathers who had a neighbor and offspring of control

fathers with a neighbor (95% CI [-1.83, 0.04], p = 0.052), with a strong difference between

offspring of isolated, predator-exposed fathers relative to offspring of control fathers with a

neighbor (95% CI [-2.40, -0.43], p = 0.006).

302
303

Figure 1: Offspring of predator-exposed fathers (blue) spent more time in the white section of

the tank compared to offspring of control fathers (grey), suggesting an anti-anxiety phenotype

(median with interquartile range, dots correspond to individuals). P- indicates control, unexposed

fathers while P+ indicates predator-exposed fathers; I- indicates fathers who had a male

neighbor, while I+ indicates isolated fathers.

Figure 2. Offspring of socially isolated fathers (darker shade), regardless of paternal predation

exposure, were captured more quickly by the predator compared to offspring of a father with a

male neighbor (lighter color). Data are median with interquartile range, dots correspond to

individuals. Although raw integer data were used for the statistical analysis, we log-transformed

the data here for better visibility of results.

Discussion

Although parents routinely encounter stress in the ecological environment that can affect

offspring development, the biological reality is that nearly all organisms are embedded within a

social environment that can either independently induce parental stress or can alter how parents

respond to ecological stressors. Here, we sought to understand how paternal social buffering (via

the presence or absence of conspecific neighbors) altered the transgenerational consequences of

paternal predation exposure. Offspring of predator-exposed fathers spent more time in the white

section of the tank, consistent with an anti-anxiety phenotype. We found that fathers who were

socially isolated had offspring who were captured faster by a live predator, suggesting that social

isolation in parents may be a stressor that leads to maladaptive effects in offspring. Further, we

found some evidence for social buffering: offspring of fathers who were exposed to predation

risk were more likely to get eaten by the predator, particularly when the father was isolated

compared to when he had a neighbor. This suggests that the social environment can mediate

transgenerational effects of paternal predation risk, with the presence of a conspecific buffering

fathers from potentially stressful environmental cues.

Previous studies have found that social isolation itself can have a detrimental effect on

those directly experiencing it, including an immunosuppressive effect in greylag geese (Ludwig

et al. 2017), oxidative stress in fish (Schumann *et al.* 2023), and elevated adrenocorticotropic

hormone and corticosterone in mice (Dronjak *et al.* 2004). This likely occurs because social

contact improves homeostatic control over the HPA axis and reduces exposure to glucocorticoids

(DeVries, Glasper & Detillion 2003). Here, we demonstrate that social isolation in adult males

does not just affect individuals within a lifetime, but also has detrimental consequences for

offspring, leading to lower survival against a predator. This may arise because ancestral stress

can make offspring more vulnerable to stress by reprograming the HPA axis, hampering

offspring ability to respond appropriately to stress encountered within their own lifetime (Yehuda

et al. 2008; Rodgers *et al.* 2013; Faraji *et al.* 2017; McDonald *et al.* 2023). Although no studies

(to our knowledge) have looked at the transgenerational consequences of paternal isolation prior

to fertilization, our results are consistent with previous studies demonstrating that other forms of

paternal social stress, including maternal separation early in life (Gapp *et al.* 2014), social defeat

(Dietz *et al.* 2011), and social instability (Kong *et al.* 2021), alter offspring anxiety and

corticosterone levels. Consequently, although nearly all TGP studies focus on the effects of

parental exposure to ecological stress on offspring, here we demonstrate the importance of

considering how social stress can also have transgenerational consequences for offspring.

The mechanism by which paternal social isolation affects offspring is unknown, but is

clearly different than mechanisms of maternal social isolation, largely because those maternal

social effects seem to be passed down via glucocorticoids in eggs or during gestation (Walsh *et*

al. 2019; MacLeod *et al.* 2023; McDonald *et al.* 2023). Dietz *et al.* (2011) found that paternal

social defeat in mice increased offspring susceptibility to depression and anxiety-like behaviors,

but that this effect was largely not present when offspring were generated via IVF, suggesting

that these transgenerational effects were mediated by changes in maternal investment in her

offspring in response to perceived paternal quality (acquired during behavioral interactions

between the mother and father). Here, we did not allow for the effects of maternal compensation

on paternal phenotypes, as offspring were generated via in-vitro fertilization without behavioral

interactions between the mother and father. Therefore, our study demonstrates that these paternal

effects can be mediated by fathers alone; as our fathers also did not interact behaviorally with

their offspring (no paternal care), these effects are presumably mediated via epigenetic changes

to sperm and/or seminal fluid. Future studies linking the precise mechanism (e.g., small RNAs or

methylation in sperm (Immler 2018)) to phenotypic effects in offspring would be useful for

predicting when and how paternal social experiences affect offspring.

We found that offspring of predator-exposed fathers spent more time in the white side of

the scototaxis arena, suggesting that they were less anxious (Maximino *et al.* 2010). This is

consistent with previous studies demonstrating that paternal stress results in offspring with

reduced anxiety and HPA responsiveness (Rodgers *et al.* 2013; Gapp *et al.* 2014; He *et al.* 2016;

Brass *et al.* 2020). Further, offspring of predator-exposed fathers tended to be consumed faster by

the predator, which is consistent with previous work in sticklebacks that found that paternal

exposure to predation risk lowered offspring survival against a predator (Hellmann *et al.* 2020).

We found some evidence that the presence of a conspecific buffered fathers against the effects of

predation risk: there was a stronger reduction in survival time in offspring of isolated, predator-

exposed fathers compared to offspring of predator-exposed fathers with a neighbor. This is

consistent with previous work demonstrating that isolated individuals show high levels of stress

in response to environmental stimuli, while individuals in social groups recover more quickly

from or have a reduced response to stressful experiences (Dronjak *et al.* 2004; Kikusui, Winslow

& Mori 2006; Ozbay *et al.* 2007; Culbert, Gilmour & Balshine 2019). Interestingly, we show that

social buffering can occur even under circumstances where we wouldn't necessarily predict

strong effects of social buffering: we used an experimental design where we paired two

unfamiliar males together, both of whom were exposed to the predator. This would typically lead

to weak effects of social buffering, as previous studies have found stronger effects of social

buffering when individuals are paired with a known conspecific compared to an unknown

conspecific (Hennessy, Kaiser & Sachser 2009), when individuals are in groups compared to

pairs (Stanton, Patterson & Levine 1985; Kikusui, Winslow & Mori 2006), and when the partner

is unstressed compared to stressed (Kiyokawa *et al.* 2004; Kikusui, Winslow & Mori 2006).

Consequently, future studies manipulating the quality of the social environment could help us

better understand how transgenerational effects of social buffering may vary with the number,

familiarity, and state of the other conspecifics.

Conclusions

We have long known that the social environment is an important source of selection, and

that any given trait (e.g., body condition) is not necessarily the property of a single individual

because social interactions will influence the traits and fitness of other individuals, even if those

individuals have never physically met (Fisher & McAdam 2017; Fisher *et al.* 2019). Currently,

we view transgenerational plasticity as a linear and relatively isolated relationship between

parent and offspring. Here, we demonstrate that the broader social environment can induce

transgenerational effects even in relatively asocial species such as stickleback, which do not form

permanent social groups. Previous work has shown that the social environment can also mediate

parental response to ecological cues, providing an opportunity for parents to learn about

ecological change in their environment in ways that have cascading effects on offspring (Afseth

et al. 2022). Consequently, the social environment can provide information about ecological cues

(Silva, Garcia de Leaniz & Luchiari 2019), act as a buffer against ecological stress (Kikusui,

Winslow & Mori 2006; Gilmour & Bard 2022), and induce stress (via both stress transmission

(Sterley *et al.* 2018), as well as via aggression, social instability, social isolation (Brandl,

Pruessner & Farine 2022)). Future studies exploring how these socially-induced effects alter both

within and trans-generational responses to ecological change can help us better predict when and

to what extent individuals will be able to adaptively response to rapidly changing environments.

References

Afseth, C., Shim, A., Anderson, S., Bell, A. & Hellmann, J.K. (2022) Vertical transmission of horizontally-acquired social information in sticklebacks: Implications for transgenerational plasticity. *Proceedings of the Royal Society B,* **289,** 20220571.

Bartolomucci, A., Palanza, P., Sacerdote, P., Ceresini, G., Chirieleison, A., Panerai, A.E. & Parmigiani, S. (2003) Individual housing induces altered immuno-endocrine responses to psychological stress in male mice. *Psychoneuroendocrinology,* **28,** 540-558.

Beemelmanns, A. & Roth, O. (2016) Biparental immune priming in the pipefish *Syngnathus typhle*. *Zoology,* **119,** 262-272.

Bell, A.M. & Hellmann, J.K. (2019) An integrative framework for understanding the mechanisms and multigenerational consequences of transgenerational plasticity. *Annual Review of Ecology, Evolution and Systematics,* **50,** 97-118.

Bonduriansky, R. & Head, M. (2007) Maternal and paternal condition effects on offspring phenotype in *Telostylinus angusticollis* (Diptera: Neriidae). *Journal of Evolutionary Biology,* **20,** 2379-2388.

Bonduriansky, R., Runagall-McNaull, A. & Crean, A.J. (2016) The nutritional geometry of parental effects: maternal and paternal macronutrient consumption and offspring phenotype in a neriid fly. *Functional Ecology,* **30,** 1675-1686.

Brandl, H.B., Pruessner, J.C. & Farine, D.R. (2022) The social transmission of stress in animal collectives. *Proceedings of the Royal Society B,* **289,** 20212158.

Brass, K.E., Herndon, N., Gardner, S.A., Grindstaff, J.L. & Campbell, P. (2020) Intergenerational effects of paternal predator cue exposure on behavior, stress reactivity, and neural gene expression. *Hormones and Behavior,* **124,** 104806.

Bygren, L.O., Tinghög, P., Carstensen, J., Edvinsson, S., Kaati, G., Pembrey, M.E. & Sjöström, M. (2014) Change in paternal grandmothers´ early food supply influenced cardiovascular mortality of the female grandchildren. *BMC Genetics,* **15**.

Chen, E., Zielinski, C., Deno, J., Singh, R., Bell, A.M. & Hellmann, J.K. (2021) The specificity of sperm-mediated paternal effects in threespine sticklebacks. *Behavioral Ecology and Sociobiology,* **75,** 68.

Cockrem, J.F. (2013) Individual variation in glucocorticoid stress responses in animals. *General and Comparative Endocrinology,* **181,** 45-58.

Crane, A.L. & Ferrari, M.C.O. (2015) Minnows trust conspecifics more than themselves when faced with conflicting information about predation risk. *Animal Behaviour,* **100,** 184-190.

Culbert, B.M., Gilmour, K.M. & Balshine, S. (2019) Social buffering of stress in a group-living fish. *Proceedings of the Royal Society B,* **286,** 20191626.

da Costa, A.P., Leigh, A.E., Man, M.-S. & Kendrick, K.M. (2004) Face pictures reduce behavioural, autonomic, endocrine and neural indices of stress and fear in sheep. *Proceedings of the Royal Society B,* **271,** 2077-2084.

Dellinger, M., Zhang, W., Bell, A.M. & Hellmann, J.K. (2018) Do male sticklebacks use visual and/or olfactory cues to assess a potential mate's history with predation risk? *Animal Behaviour,* **145,** 151-159.

DeVries, A.C., Glasper, E.R. & Detillion, C.E. (2003) Social modulation of stress responses. *Physiology & Behavior,* **79,** 399-407.

Dietz, D.M., LaPlant, Q., Watts, E.L., Hodes, G.E., Russo, S.J., Feng, J., Oosting, R.S., Vialou, V. & Nestler, E.J. (2011) Paternal transmission of stress-induced pathologies. *Biological Psychiatry,* **70,** 408-414.

Donelson, J.M., Wong, M., Booth, D.J. & Munday, P.L. (2016) Transgenerational plasticity of reproduction depends on rate of warming across generations. *Evolutionary Applications,* **9,** 1072-1081.

Dronjak, S., Gavrilović, L., Filipović, D. & Radojčić, M.B. (2004) Immobilization and cold stress affect sympatho–adrenomedullary system and pituitary–adrenocortical axis of rats exposed to long-term isolation and crowding. *Physiology & Behavior,* **81,** 409-415.

Faraji, J., Soltanpour, N., Lotfi, H., Moeeini, R., Moharreri, A.-R., Roudaki, S., Hosseini, S.A., Olson, D.M., Abdollahi, A.-A., Soltanpour, N., Mohajerani, M.H. & Metz, G.A.S. (2017) Lack of social support raises stress vulnerability in rats with a history of ancestral stress. *Scientific Reports,* **7,** 5277.

Faustino, A.I., Tacão-Monteiro, A. & Oliveira, R.F. (2017) Mechanisms of social buffering of fear in zebrafish. *Scientific Reports,* **7,** 44329.

Fisher, D.N., Haines, J.A., Boutin, S., Dantzer, B., Lane, J.E., Coltman, D.W. & McAdam, A.G. (2019) Indirect effects on fitness between individuals that have never met via an extended phenotype. *Ecology Letters,* **22,** 697-706.

Fisher, D.N. & McAdam, A.G. (2017) Social traits, social networks and evolutionary biology. *Journal of Evolutionary Biology,* **30,** 2088-2103.

Fürtbauer, I. & Heistermann, M. (2016) Cortisol coregulation in fish. *Scientific Reports,* **6,** 30334.

Gapp, K., Jawaid, A., Sarkies, P., Bohacek, J., Pelczar, P., Prados, J., Farinelli, L., Miska, E. & Mansuy, I.M. (2014) Implication of sperm RNAs in transgenerational inheritance of the effects of early trauma in mice. *Nature Neuroscience,* **17,** 667-669.

Gilad, T. & Scharf, I. (2019) Separation between maternal and paternal effects on offspring following exposure of adult red flour beetles to two stressors. *Ecological Entomology,* **44,** 494-501.

Gilmour, K.M. & Bard, B. (2022) Social buffering of the stress response: insights from fishes. *Biology Letters,* **18,** 20220332.

Hagmayer, A., Lankheet, M.J., Bijsterbosch, J., van Leeuwen, J.L. & Pollux, B.J.A. (2022) Maternal food restriction during pregnancy affects offspring development and swimming performance in a placental live-bearing fish. *Journal of Experimental Biology,* **225,** jeb242850.

He, N., Kong, Q.-Q., Wang, J.-Z., Ning, S.-F., Miao, Y.-L., Yuan, H.-J., Gong, S., Cui, X.-Z., Li, C.-Y. & Tan, J.-H. (2016) Parental life events cause behavioral difference among offspring: Adult pre-gestational restraint stress reduces anxiety across generations. *Scientific Reports,* **6,** 39497.

Hellmann, J.K., Bukhari, S.A., Deno, J. & Bell, A.M. (2020) Sex-specific plasticity across generations I: Maternal and paternal effects on sons and daughters. *Journal of Animal Ecology,* **89,** 2788-2799.

Hellmann, J.K., Carlson, E.R. & Bell, A.M. (2020) Sex-specific plasticity across generations II: Grandpaternal effects are lineage- and sex-specific. *Journal of Animal Ecology,* **89**.

Hellmann, J.K., Carlson, E.R. & Bell, A.M. (2021) The interplay between sperm-mediated and care-mediated paternal effects in threespined sticklebacks. *Animal Behaviour,* **179,** 267-277.

Hennessy, M.B., Kaiser, S. & Sachser, N. (2009) Social buffering of the stress response:
Diversity, mechanisms, and functions. *Frontiers in Neuroendocrinology,* **30,** 470-482.
Immler, S. (2018) The sperm factor: paternal impact beyond genes. *Heredity,* **121,** 239-247.
Kassambara, A. & Mundt, M. (2017) factoextra: extract and visualize the results of multivariate
data analyses. R package version 1.0.5. https://CRAN.R-project.org/package=factoextra.
Kikusui, T., Winslow, J.T. & Mori, Y. (2006) Social buffering: relief from stress and anxiety.
Philosophical Transactions of the Royal Society B, **361,** 2215-2228.
Kiyokawa, Y., Kikusui, T., Takeuchi, Y. & Mori, Y. (2004) Partner's stress status influences social
buffering effects in rats. *Behavioral Neuroscience* **118**, 798-804.
Kong, Q.-Q., Tian, X.-D., Wang, J., Yuan, H.-J., Ning, S.-F., Luo, M.-J. & Tan, J.-H. (2021) A
next-generation sequencing study on mechanisms by which restraint and social instability
stresses of male mice alter offspring anxiety-like behavior. *Scientific Reports,* **11,** 7952.
Koolhaas, J.M., Korte, S.M., De Boer, S.F., Van Der Vegt, B.J., Van Reenen, C.G., Hopster, H.,
De Jong, I.C., Ruis, M.A.W. & Blokhuis, H.J. (1999) Coping styles in animals: current
status in behavior and stress-physiology. *Neuroscience & Biobehavioral Reviews,* **23,**
925-935.
Ludwig, S.C., Kapetanopoulos, K., Kotrschal, K. & Wascher, C.A.F. (2017) Effects of mate
separation in female and social isolation in male free-living Greylag geese on behavioural
and physiological measures. *Behavioural Processes,* **138,** 134-141.
MacLeod, K.J., English, S., Ruuskanen, S.K. & Taborsky, B. (2023) Stress in the social context:
a behavioural and eco-evolutionary perspective. *Journal of Experimental Biology,* **226,**
jeb245829.
MacLeod, K.J., Monestier, C., Ferrari, M.C.O., McGhee, K.E., Sheriff, M.J. & Bell, A.M. (2022)
Predator-induced transgenerational plasticity in animals: a meta-analysis. *Oecologia,* **200,**
371-383.
Maximino, C., Marques de Brito, T., Dias, C.A.G.d.M., Gouveia Jr, A. & Morato, S. (2010)
Scototaxis as anxiety-like behavior in fish. *Nature Protocols,* **5,** 209.
McDonald, R.J., Hong, N.S., Trow, J.S., Kaupp, C., Balog, R.J., Gokarn, L., Falkenberg, E.A.,
McCreary, K.J., Soltanpour, N., Witbeck, C., McKenna, A. & Metz, G.A.S. (2023)
Effects of maternal social isolation on adult rodent offspring cognition. *Scientific Reports,*
13, 7748.
McQuaid, R.J., McInnis, O.A., Paric, A., Al-Yawer, F., Matheson, K. & Anisman, H. (2016)
Relations between plasma oxytocin and cortisol: The stress buffering role of social
support. *Neurobiology of Stress,* **3,** 52-60.
Mommer, B.C. & Bell, A.M. (2013) A test of maternal programming of offspring stress response
to predation risk in threespine sticklebacks. *Physiology and Behavior,* **122,** 222-227.
Moore, R.S., Kaletsky, R. & Murphy, C.T. (2019) Piwi/PRG-1 Argonaute and TGF-α Mediate
Transgenerational Learned Pathogenic Avoidance. *Cell,* **177,** 1827-1841.e1812.
Nylen, K.J., O'Hara, M.W. & Engeldinger, J. (2013) Perceived social support interacts with
prenatal depression to predict birth outcomes. *Journal of Behavioral Medicine,* **36,** 427-
440.
Ozbay, F., Johnson, D.C., Dimoulas, E., Morgan, C.A., Charney, D. & Southwick, S. (2007)
Social support and resilience to stress: from neurobiology to clinical practice. *Psychiatry*
(Edgmont), **4,** 35-40.

Pessato, A., McKechnie, A.E. & Mariette, M.M. (2022) A prenatal acoustic signal of heat affects
544   thermoregulation capacities at adulthood in an arid-adapted bird. *Scientific Reports,* **12,**
545   5842.
546   Potticary, A.L. & Duckworth, R.A. (2020) Multiple environmental stressors induce an adaptive
547   maternal effect. *American Naturalist,* **196,** 487-500.
548   Rodgers, A.B., Morgan, C.P., Bronson, S.L., Revello, S. & Bale, T.L. (2013) Paternal stress
549   exposure alters sperm microRNA content and reprograms offspring HPA stress axis
550   regulation. *Journal of Neuroscience,* **33,** 9003-9012.
551   Romero, L.M., Dickens, M.J. & Cyr, N.E. (2009) The reactive scope model — A new model
552   integrating homeostasis, allostasis, and stress. *Hormones and Behavior,* **55,** 375-389.
553   Schumann, S., Mozzi, G., Piva, E., Devigili, A., Negrato, E., Marion, A., Bertotto, D. &
554   Santovito, G. (2023) Social buffering of oxidative stress and cortisol in an endemic
555   cyprinid fish. *Scientific Reports,* **13,** 20579.
556   Séguin, L., Potvin, L., St.-Denis, M. & Loiselle, J. (1995) Chronic stressors, social support, and
557   depression during pregnancy. *Obstetrics & Gynecology,* **85,** 583-589.
558   Sheriff, M.J., Krebs, C.J. & Boonstra, R. (2010) The ghosts of predators past: population cycles
559   and the role of maternal programming under fluctuating predation risk. *Ecology,* **91,**
560   2983-2994.
561   Silva, P.F., Garcia de Leaniz, C. & Luchiari, A.C. (2019) Fear contagion in zebrafish: a
562   behaviour affected by familiarity. *Animal Behaviour,* **153,** 95-103.
563   Stanton, M.E., Patterson, J.M. & Levine, S. (1985) Social influences on conditioned cortisol
564   secretion in the squirrel monkey. *Psychoneuroendocrinology,* **10,** 125-134.
565   Sterley, T.-L., Baimoukhametova, D., Füzesi, T., Zurek, A.A., Daviu, N., Rasiah, N.P.,
566   Rosenegger, D. & Bains, J.S. (2018) Social transmission and buffering of synaptic
567   changes after stress. *Nature Neuroscience,* **21,** 393-403.
568   Tariel, J., Plénet, S. & Luquet, É. (2020) Transgenerational plasticity in the context of predator-
569   prey interactions. *Frontiers in Ecology and Evolution,* **8,** 319.
570   Walsh, K., McCormack, C.A., Webster, R., Pinto, A., Lee, S., Feng, T., Krakovsky, H.S.,
571   O'Grady, S.M., Tycko, B., Champagne, F.A., Werner, E.A., Liu, G. & Monk, C. (2019)
572   Maternal prenatal stress phenotypes associate with fetal neurodevelopment and birth
573   outcomes. *Proceedings of the National Academy of Sciences,* **116,** 23996-24005.
574   Yehuda, R., Bell, A., Bierer, L.M. & Schmeidler, J. (2008) Maternal, not paternal, PTSD is
575   related to increased risk for PTSD in offspring of Holocaust survivors. *Journal of*
576   *Psychiatric Research,* **42,** 1104-1111.
577   Yusishen, M.E., Yoon, G.R., Bugg, W., Jeffries, K.M., Currie, S. & Anderson, W.G. (2020) Love
578   thy neighbor: Social buffering following exposure to an acute thermal stressor in a
579   gregarious fish, the lake sturgeon (*Acipenser fulvescens*). *Comparative Biochemistry and*
580   *Physiology Part A: Molecular & Integrative Physiology,* **243,** 110686.