1   **The transgenerational consequences of paternal social isolation and predation exposure in**

2   **threespined sticklebacks**

Abstract

1. Parents routinely encounter stress in the ecological environment that can affect offspring development (transgenerational plasticity: TGP); however, parents' interactions with conspecifics may alter how parents respond to ecological stressors.

2. During social buffering, the presence of conspecifics can reduce the response to or increase the speed of recovery from a stressor. This may have cascading effects on offspring if conspecifics can mitigate parental responses to ecological stress in ways that blunt the transmission of stress-induced transgenerational effects.

3. Here, we simultaneously manipulated both paternal social isolation and experience with predation risk prior to fertilization in threespined stickleback (*Gasterosteus aculeatus*). We generated offspring via in-vitro fertilization to allow us to isolate paternal effects mediated via sperm alone (i.e., in the absence of paternal care). If social buffering mitigates TGP induced by paternal exposure to predation risk, then we expect the transgenerational effects of predation exposure to be weaker when a conspecific is present compared to when the father is isolated.

4. Offspring of predator-exposed fathers showed reduced anxiety-like behavior and tended to be captured faster by the predator. Fathers who were socially isolated also had offspring that were captured faster by a live predator, suggesting that paternal social isolation may have maladaptive effects on how offspring respond to ecological stressors. Despite additive effects of paternal social isolation and paternal predation risk, we found no evidence of an interaction between these paternal treatments, suggesting that the presence of a conspecific did not buffer fathers and/or offspring from the effects of predation risk.

5. Our results suggest that socially-induced stress is an important, yet underappreciated,

mediator of TGP and can elicit transgenerational effects even in species that do not form

permanent social groups. Future studies should therefore consider how the parental social

environment can affect both within and trans-generational responses to ecological

stressors.

Key words: *Gasterosteus aculeatus*, paternal effects, phenotypic plasticity, predation risk, social

buffering, stress, intergenerational

Introduction

Organisms encounter a wide variety of threats and stressors in their environment that can push individuals beyond their homeostatic limits and induce physiological and behavioral changes that allow them to survive those stressors (Romero, Dickens & Cyr 2009). It is well known that individuals differ drastically in how they respond to these environmental stressors: the same external stressor can induce strong responses in some individuals and weak responses in others (Koolhaas *et al.* 1999; Cockrem 2013). Although some of this variation can be attributed to traits of the organism itself (e.g., personality (Cockrem 2013), sex (DeVries, Glasper & Detillion 2003)), some of this variation is environmentally dependent. In particular, there is ample evidence that the social environment alters how animals cope with ecological stress, influencing both the perception of and response to ecological cues (DeVries, Glasper & Detillion 2003; Brandl, Pruessner & Farine 2022). In many cases, the presence of conspecifics can reduce physiological and behavioral responses to stress via social buffering, which occurs when the presence of a conspecific reduces the response to or increases the speed of recovery from an external stressor (Kikusui, Winslow & Mori 2006; Gilmour & Bard 2022). For example, socially isolated mice showed greater immune impairment and a higher basal corticosterone increase in response to a mild stress compared to group housed mice (Bartolomucci *et al.* 2003). Similarly, in humans, anxiety and negative emotions induced by a social stress test were attenuated when women had social support (McQuaid *et al.* 2016). This social buffering may be an adaptive response to minimize the transmission of stress among group members (Gilmour & Bard 2022), which can occur when a stress response system (e.g., HPA axis) in one individual is activated after interacting with another stressed individual (Brandl, Pruessner & Farine 2022). Further, social buffering can even increase the limits of the physical conditions organisms are willing to

tolerate (Gilmour & Bard 2022), suggesting that sociality can mediate responses to ecological

stressors.

While social buffering has been well studied in highly social mammals, there is mounting

evidence that social buffering of stress also occurs in less social species, including solitary

species and species with aggregations or schooling behavior (Gilmour & Bard 2022). For

example, in lake sturgeon (*Acipenser fulvescens*), group-housed fish had significantly reduced

endocrine and cellular stress markers after exposure to heat stress compared to isolated

individuals (Yusishen *et al.* 2020). This social buffering effect can even occur in small groups of

two individuals (Faustino, Tacão-Monteiro & Oliveira 2017; Schumann *et al.* 2023) or in

response to mere visual cues of a conspecific (da Costa *et al.* 2004; Faustino, Tacão-Monteiro &

Oliveira 2017), suggesting that these social buffering effects are robust and occur in a wide range

of taxonomic groups with differing levels of sociality.

While social buffering effects can have consequences for the individuals that directly

experience the stressor, social buffering in parents may also have consequences for offspring.

Transgenerational plasticity (TGP) occurs when the experiences of a previous generation alter

the phenotypes of future generations (Bell & Hellmann 2019). TGP can occur in response to a

variety of ecological stressors, including predation risk (Tariel, Plénet & Luquet 2020; MacLeod

et al. 2022), heat exposure (Donelson *et al.* 2016; Pessato, McKechnie & Mariette 2022),

immune challenges (Beemelmanns & Roth 2016; Moore, Kaletsky & Murphy 2019), and food

limitation (Bygren *et al.* 2014; Hagmayer *et al.* 2022). These ecological stressors induce changes

in parental state, including the activation of the hypothalamic-pituitary-adrenal axis (HPA axis),

that influence offspring development, such as stress responsiveness, growth, and behavior (Bell

& Hellmann 2019; Potticary & Duckworth 2020). Given that the magnitude of the stress

response in parents can be linked to the magnitude of the phenotypic changes in offspring

(Sheriff, Krebs & Boonstra 2010), social buffering may affect both the strength and frequency

with which parental experiences affect offspring traits. This is supported by a recent study in

humans which found that the adverse health outcomes differentiating offspring of healthy and

psychologically stressed mothers could be specifically linked to a lack of social support in

psychologically stressed mothers (Walsh *et al.* 2019). These effects may arise because

conspecifics provide a buffering effect that abates maternal response to external stress (e.g.,

brings glucocorticoids back to baseline quickly (Kikusui, Winslow & Mori 2006)), mitigating

adverse outcomes for offspring (Nylen, O'Hara & Engeldinger 2013).

Although this previous work suggests the importance of the social environment in

mediating transgenerational effects for mothers, we know little about the transgenerational

effects of social isolation for fathers. Fathers have at least two mechanisms by which they can

influence offspring non-genetically: epigenetic changes to sperm/seminal fluid and behavioral

changes to parental care (Immler 2018). Paternal experiences can affect offspring phenotypes in

magnitudes that are similar to maternal experiences (Akkerman *et al.* 2016; Beemelmanns &

Roth 2016; Hellmann *et al.* 2020), although it is difficult to compare the strength of parental

effects in most systems because the effects of paternal experiences often have different effects on

offspring than maternal experiences (Bonduriansky & Head 2007; Bouwhuis, Vedder & Becker

2015; Bonduriansky, Runagall-McNaull & Crean 2016; Bell & Hellmann 2019; Gilad & Scharf

2019; Hellmann *et al.* 2020). This is likely because males and females differ in their

susceptibility to stress (DeVries, Glasper & Detillion 2003) and because the mechanisms of

inheritance are different (e.g., eggs/placenta versus sperm). Consequently, the transgenerational

effects of paternal isolation are likely different from maternal isolation. Thus, understanding how

the social environment mediates paternal effects can help us better predict when and how

paternal experiences influence offspring phenotypes and if, for instance, social buffering during

or after a stressor mitigates or reverses changes in parental state that induce transgenerational

effects.

We sought to understand the ways that the social environment alters the transgenerational

consequences of paternal stress in threespined stickleback *(Gasterosteus aculeatus).* There is

abundant evidence for predator-induced paternal effects mediated via sperm (i.e., in the absence

of paternal care) in stickleback, which alter offspring behavior, growth, gene expression,

survival, and stress responsiveness (Hellmann *et al.* 2020; Hellmann, Carlson & Bell 2020; Chen

et al. 2021; Hellmann, Carlson & Bell 2021; Afseth *et al.* 2022; Hellmann *et al.* 2024). There is

also previous evidence of social buffering effects in stickleback: fish that are exposed to

simulated predation risk while isolated had higher circulating levels of cortisol compared to fish

in a group (Mommer & Bell 2013) and pairs of sticklebacks co-regulate their cortisol responses

in stressful environments (Fürtbauer & Heistermann 2016). However, we have very little

understanding as to how the social environment might modulate parental responses to ecological

stress in ways that might have cascading effects on the adaptive nature and frequency of

transgenerational plasticity.

We used a full factorial design to compare the transgenerational consequences of

exposing fathers to predation risk while fathers were isolated or with another conspecific.

Offspring were generated via in-vitro fertilization; because fathers had no behavioral interactions

with offspring post-fertilization, this allowed us to isolate paternal effects mediated via sperm

and/or seminal fluid. We chose to focus on sperm-mediated paternal effects rather than paternal

care because, in care-mediated paternal effects, it can be difficult to isolate changes in offspring

phenotypes that come directly from the father (transgenerational effects) versus changes that

result from the early life environment (within-generational effects). Further, a previous study

found that the presence or absence of paternal care did not alter the transgenerational effects

mediated via sperm alone (Hellmann, Carlson & Bell 2021). We then measured offspring

behavioral traits relevant to predation defense, including anti-anxiety behavior, shoaling

behavior, and survival against a live predator. If social buffering mitigates TGP induced by

paternal exposure to predation risk, then we expect the transgenerational effects of predation

exposure to be weaker when a conspecific is present compared to when the father is isolated.

Materials and Methods

Housing information and F0 exposures. In June 2021, we collected adult three-spined

stickleback *(Gasterosteus aculeatus)* from Putah Creek (CA, USA), a freshwater stream in

northern California, and shipped them to the University of Dayton. We maintained the F0

generation on a summer photoperiod schedule (16 L : 8 D) at $20° \pm 1°C$ and fed them ad libitum

daily with a mix of frozen bloodworms (*Chironomus* spp.), brine shrimp (Artemia spp.) Mysis

shrimp, and Cyclop-eeze. We obtained 10-15 cm long rainbow trout *(Oncorhynchus mykiss)*

from Freshwater Farms of Ohio in June 2021. Putah Creek has piscivorous predators, including

the prickly sculpin (*Cottus asper*); while rainbow trout are present in Putah Creek, they are not

present in the isolated section of Putah Creek where we collected stickleback (Beaver Pond),

suggesting that rainbow trout is an environmentally relevant, but novel predator.

In August 2021, we moved males into experimental 37.9 L (53L x 33W x 25H cm) tanks

divided into three equal sections using removable barriers. Males in the neighbor treatment were

separated with clear barriers with nine 3mm holes to allow for water flow between the sections.

Males in the isolated treatment were separated with permanent opaque barriers that did not allow

for water flow between sections. In both treatments, we transferred males from their home tank

using a cup (males were never removed from water to minimize stress) and placed one male

stickleback in each of the outer sections and left the middle section empty. Each male section

contained gravel, one fake plant, and nesting materials: a container of sand, a terra cotta pot for

cover, and algae. The middle section contained only gravel. Males that shared a tank were

assigned to the same treatment group. Next to each nesting male tank, we placed an undivided

37.9L tank containing gravel and two sponge filters, which either contained one live rainbow

trout or was left empty depending on the predator treatment. All experimental tanks were

separated visually from each other with a removable opaque divider.

We exposed male stickleback to predator cues once a day for two hours, for 11-14

consecutive days depending on the availability of gravid females. We used a short stressor to

avoid potential habituation to predation risk (Dellinger *et al.* 2018). Further, by beginning the

treatment when males were transferred to a nesting arena, we sought to mimic the change in

predator regime that males may encounter as they move into a different habitat to nest. At a

random time of day between 0800 hrs and 2000 hrs, we turned off the bubblers in the tanks and

removed the opaque divider to reveal the adjacent tank (which was empty or contained a live

trout). At the same time, we added either 100mL of control RO water or the prepared predator

cue to each end compartment containing one male stickleback. We prepared predator cues using

methods consistent with Crane and Ferrari (2015). Briefly, we placed a trout in a 37.9 L tank

with clean, conditioned tap water proportional to the size of the trout (500mL per gram of fish).

We fed the trout one juvenile stickleback per day for four days and after 96hrs, we froze the tank

water into 200 mL aliquots.

In the predator-naïve treatment, males saw an empty tank and received 100 mL of RO

water. In the predator treatment, males saw a live trout and received 100 mL of the prepared

predator cue. Every other day, we fed the trout one juvenile stickleback during the two hour

exposure to reinforce the visual cue. Importantly, despite holes between the sections, we found

very limited diffusion of water between the sections (tested using dyed water) when bubblers

were turned off, suggesting that these olfactory cues did not disperse throughout the tank during

the predation exposure and therefore, the neighbor treatment did not receive weaker olfactory

cues or predation risk than the isolated group. However, water did disperse between sections

when the bubblers were on, suggesting that neighboring males exchanged olfactory cues outside

of the two-hour exposure period. After two hours, we replaced the dividers and completed a 50%

water change on all treatments. We removed males after the last day of exposure.

Generating Offspring. The day after the last exposure, we generated offspring by manually

squeezing the eggs out of a gravid female (n = 29 different females across the 35 clutches, each

sired by a unique male) and fertilizing them with sperm from the testes of the dissected

experimental male. Females were all predator naïve and housed in group tanks on recirculating

racks with biological, particulate, and UV filters. We checked clutches for successful fertilization

via microscope and then placed into a small cup with a mesh bottom above an air bubbler in an

18.9L (38L x 25W x 21H) tank containing gravel and two fake plants, with each clutch housed

individually on a recirculating system. By artificially fertilizing the eggs and incubating the

embryos using an air bubbler, we controlled for possible pre-fertilization effects mediated by

interactions between mothers and fathers (Mashoodh *et al.* 2012; McGhee *et al.* 2015), post-

fertilization effects mediated by paternal care (Stein & Bell 2014), and the possibility that

stressed parents might be less likely to successfully mate or parent offspring. Once hatched, we

fed offspring newly hatched brine shrimp for two months and then gradually transitioned to

frozen food (as above). We switched offspring to a winter light schedule (8 L: 16 D) prior to

starting behavioral assays. We generated a total of n = 35 clutches: n = 9 of predator-naïve

fathers with a neighbor, n = 9 of predator-naïve, isolated fathers, n = 8 of predator-exposed

fathers with a neighbor, and n = 9 of predator-exposed, isolated fathers.. We assayed offspring at

approximately 5 months of age; all offspring went through the shoaling assay first, followed by

the scototaxis assay 24 hours later, and then the survival assay 48-72 after that.

Offspring assays.

Shoaling assays. We used experimental 37.9L (53 x 33 x 24 cm high) tanks lined with gravel.

On one end of the tank, we positioned three small plastic plants; on the other end of the tank, we

created a small compartment for shoaling fish by inserting a clear plastic barrier (10 cm from the

edge) perforated with small holes to allow for flow of water and olfactory cues. The evening

before a day of trials, we placed five size-matched juveniles (similarly aged juveniles that were

not part of the experiment) into the shoaling side compartment of each tank to habituate

overnight. These fish were unrelated and unfamiliar to the focal fish.

At the start of the shoaling trial, we placed an experimental offspring (focal fish) into the

open area in the middle of the tank and allowed the focal fish to habituate for 10 minutes. After

the habitation, we used BORIS (Friard & Gamba 2016) to record total time spent shoaling, total

time spent oriented to the shoal while shoaling (the focal fish's head had to be pointing in the

direction of at least one member of the shoal), total time spent hiding in plants, and time spent in

the gravel in the center of the tank over a 10 minute period. We defined a focal fish to be

shoaling or hiding in the plant if it was within one body length of the shoal barrier or plants,

respectively. Upon completion of the assay, we gently removed the focal fish from the tank with

a cup, weighed and measured them, and then placed them into individual holding tanks. We

assayed and measured n = 242 fish: n = 61 of predator-naïve fathers with a neighbor, n = 57 of

predator-naïve, isolated fathers, n = 66 of predator-exposed fathers with a neighbor, and n = 58

of predator-exposed, isolated fathers.

***Measuring anxiety/cautiousness*.** Scototaxis (light/dark preference) protocols have been

developed to test anti-anxiety/cautious behavior in fish (Maximino *et al.* 2010). On the day after

the shoaling assay, offspring were gently caught with a cup from their holding tank and placed in

a clear cylinder (10.5cm diameter) in the center of a half-black, half-white tank (51L x 28W x

19H cm). After a five minute acclimation period, we lifted the cylinder, and fish explored the

tank for 15 minutes, during which we measured the latency to enter the white section, total time

in the white section, and thigmotaxis behavior (time spent within a body length of the wall)

within the white section. We only measured thigmotaxis in the white area because we felt that

thigmotaxis in a light, open area is more reflective of anxiety like behavior compared to

thigmotaxis in a darker area. After the 15-minute testing period, we removed the fish from the

tank and placed them back in their holding tank. We assayed n = 234 fish: n = 58 of predator-

naïve fathers with a neighbor, n = 55 of predator-naïve, isolated fathers, n = 64 of predator-

exposed fathers with a neighbor, and n = 57 of predator-exposed, isolated fathers.

Survival assay. 48-72 hours after the scototaxis assay, we conducted survival assays in groups of

two siblings (same paternal treatment and genetic family). We filled circular portable pools with

conditioned tap water (1m diameter, 13 cm water depth) and evenly placed six plants around

each arena. We placed an individually marked rainbow trout (n = 4) in each pool 24-hours before

the first day of assays. We fasted the trout for 48 hours prior to being placed in the pools and we

fed them one non-experimental juvenile stickleback after being placed in the pool to keep hunger

levels standardized across days. On each assay day, we used each trout for two assays, once in

the morning and once in the afternoon. For each assay, we placed two siblings in a clear cylinder

(20.3 cm diameter) in the middle of the arena. After acclimating for 20 minutes, we lifted the

cylinder and used a stopwatch to time how long it took the trout to capture one of the fish

(survival time). We had a total of n=81 trials and we analyzed all trials where one of the fish was

captured within one hour (n=79 trials: n = 26 of predator-naïve fathers with a neighbor, n = 13

of predator-naïve, isolated fathers, n = 22 of predator-exposed fathers with a neighbor, and n =

18 of predator-exposed, isolated fathers)

Statistical methods. To get an estimate of shoaling behavior, we used principal components

analysis (R package factoextra (Kassambara & Mundt 2017)) to combine several behaviors: time

spent hidden in the plant, time spent shoaling, time spent oriented to the shoal, and time spent in

the gravel in the center of the tank. We extracted an eigenvalue of 2.74 that captured 68.5% of

the variance in these behaviors; positive values indicate more social individuals who spent more

time shoaling (contribution: 33.4%) and oriented to the shoal (31.3%), and spent less time hiding

in plants (25.1%) and gravel (10.2%). This PC was used as the dependent variable in the model

described below (Gaussian distribution). For the scototaxis assay, latency to enter the white arena

was negatively correlated with time spent in the white section (Spearman rank correlation: ρ = -

0.42, p<0.001), so we analyzed variation in time spent in the white section of the tank, as well as

time spent performing thigmotaxis in the white section of the tank (both Poisson distribution).

We used the package MCMCglmm (Hadfield 2010) to analyze offspring behavior,

offspring length, and offspring mass. We ran models with a weak prior on the variance (V=1,

nu=0.002) because our data were heteroskedastic. We ran models for 200,000 iterations, with a

burn-in of 3000 iterations and thin = 3; we confirmed model fit and convergence. Models testing

variation in shoaling and scototaxis behavior included fixed effects of paternal predation

exposure, paternal isolation, and offspring standard length as well as random effects of clutch

identity and observer identity. For the thigmotaxis model, we also included a fixed effect of total

time in the white section of the tank. The survival model (time to capture, log-transformed,

Gaussian distribution) included fixed effects of paternal predation exposure and paternal

isolation, and random effects of clutch identity, observer identity (n = 4), and trout identity (n =

4). The model testing predictors of offspring length (Gaussian distribution) included fixed effects

of paternal predation exposure, paternal isolation, and offspring age, while the model testing

predictors of offspring mass (Gaussian distribution) included fixed effects of paternal predation

exposure, paternal isolation, and offspring length; both models included random effects of clutch

identity and observer identity. For all models, we included the interaction between paternal

predation exposure and paternal isolation as this was part of our *a priori* predictions. We

estimated effect sizes for all significant effects of paternal treatment using rank-epsilon-

squared tests (R package effectsize (Ben-Shachar, Lüdecke & Makowski 2020)) to account for

the non-parametric nature of our data. Effect sizes less than 0.06 indicate a small effect, while

effect sizes between 0.06 and 0.14 indicate a medium effect size (Ben-Shachar, Lüdecke &

Makowski 2020).

288  *Animal welfare note.* All methods were approved by Institutional Animal Care and Use

289 Committee of University of Dayton (protocol ID 021-05), including the use of live predators.

291 **Results**

292   We examined how paternal exposure to predation risk and paternal social isolation altered

293 offspring behavior (shoaling and anti-anxiety behavior), size, and survival against live predators.

295  *Offspring behavior and size*. We found no change in offspring shoaling behavior based on either

paternal isolation or paternal predation exposure, although longer offspring spent more time

shoaling than smaller offspring (Table 1). However, in the scototaxis assay testing anti-anxiety

behavior, we found that offspring of predator-exposed fathers spent more time in the white

section of the tank compared to offspring of predator-naïve fathers (Table 1; Figure 1; rank ε^2 =

0.03). We found no significant effect of paternal isolation on time spent in the white section

(Table 1; Figure 1). We found no significant effect of paternal isolation or paternal predation

exposure on offspring thigmotaxis behavior (Table 1). We also found no differences in offspring

length or mass based on either paternal isolation or paternal predation exposure (Table 1).

305  *Offspring survival*. We found that offspring of fathers who were isolated were captured

significantly faster by the predator compared to offspring of fathers with neighbors (mean 45 sec

$\pm$ 19 s.e. versus 153 sec $\pm$ 49 s.e.; rank ε^2 = 0.07; Table 1; Figure 2). Similarly, offspring of

fathers who were predator-exposed tended to be captured significantly faster by the predator

compared to offspring of predator-naïve fathers (mean 87 sec $\pm$ 49 s.e. versus 135 sec $\pm$ 39 s.e.;

rank $\varepsilon^2 = 0.07$; Table 1; Figure 2). While the combined influence of paternal social isolation and

predation risk seemed to have an additive effect on offspring survival (Figure 2), we did not find

a significant interaction of paternal isolation and paternal predation exposure (Table 1),

suggesting no evidence for social buffering.

Table 1: Results of mixed effects models on shoaling behavior (PCA), anti-anxiety behavior in

the scototaxis assay (time in white section, thigmotaxis), survival against a live predator, mass,

and length.

	Shoaling		
	Mean	*95% CI*	*P*
Paternal predation exposure	-0.10	-0.80, 0.56	0.77
Paternal isolation	-0.04	-0.73, 0.61	0.89
Standard length	0.15	0.06, 0.24	**0.001**
Paternal isolation * predation	-0.05	-1.00, 0.91	0.92
	Time in white section		
	Mean	*95% CI*	*P*
Paternal predation exposure	0.18	0.01, 0.36	**0.047**
Paternal isolation	-0.05	-0.24, 0.14	0.59
Standard length	-0.01	-0.03, 0.01	0.39
Paternal isolation * predation	-0.04	-0.29, 0.22	0.78
	Thigmotaxis		
	Mean	*95% CI*	*P*
Paternal predation exposure	0.05	-0.05, 0.16	0.33
Paternal isolation	-0.01	-0.13, 0.11	0.85
Time in white section	0.003	0.0028, 0.0033	**<0.001**
Length	0.02	0.005, 0.03	**0.009**
Paternal isolation * predation	-0.01	-0.17, 0.14	0.85
	Survival time		
	Mean	*95% CI*	*P*
Paternal predation exposure	-0.89	-1.80, 0.07	0.058
Paternal isolation	-1.17	-2.24, -0.13	**0.03**
Paternal isolation * predation	0.64	-0.85, 2.09	0.38
	Length		
	Mean	*95% CI*	*P*
Paternal predation exposure	1.26	-1.24, 3.74	0.31
Paternal isolation	0.04	-2.50, 2.47	0.97
Age	0.05	0.01, 0.09	**0.009**
Paternal isolation * predation	-1.31	-4.81, 2.34	0.46
	Mass		
	Mean	*95% CI*	*P*
Paternal predation exposure	-0.005	-0.02, 0.01	0.60
Paternal isolation	-0.003	-0.02, 0.01	0.72
Length	0.01	0.010, 0.0014	**<0.001**
Paternal isolation * predation	0.005	-0.02, 0.03	0.71

318
319

Figure 1: Offspring of predator-exposed fathers (blue) spent more time in the white section of

the tank compared to offspring of predator-naïve fathers (grey), suggesting an anti-anxiety

phenotype (median with interquartile range, dots correspond to individuals). P- indicates

predator-naïve, unexposed fathers while P+ indicates predator-exposed fathers; I- indicates

fathers who had a male neighbor, while I+ indicates isolated fathers.

Figure 2. Offspring of socially isolated fathers (darker shade) were captured more quickly by the

predator compared to offspring of a father with a male neighbor (lighter color). Similarly,

offspring of predator-exposed fathers (blue) tended to be captured more quickly by the predator

compared to offspring of predator-naïve fathers (grey). Data are median with interquartile range,

dots correspond to individuals. Although raw integer data were used for the statistical analysis,

we log-transformed the data here for better visibility of results.

Discussion

Parents routinely encounter stress in the ecological environment that can affect offspring development. However, the social environment may alter how parents respond to ecological stressors, and therefore, affect how those parental experiences get integrated into offspring phenotypes. Here, we sought to understand how paternal social buffering (via the presence or absence of conspecific neighbors) altered the transgenerational consequences of paternal predation exposure. Offspring of predator-exposed fathers spent more time in the white section of the tank, consistent with an anti-anxiety phenotype, and had offspring that tended to be captured faster by the predator. We found that fathers who were socially isolated had offspring who were captured faster by a live predator, suggesting that social isolation in parents may be a stressor that alters offspring ability to cope with ecological stress. Despite evidence that paternal social isolation and predation risk have additive effects on offspring survival, we did not find evidence for a social buffering effect, where the presence of a conspecific buffers fathers and/or offspring from the effects of predation risk.

Previous studies have found that social isolation itself can have a detrimental effect on those directly experiencing it, including an immunosuppressive effect in greylag geese (Ludwig *et al.* 2017), oxidative stress in fish (Schumann *et al.* 2023), and elevated adrenocorticotropic hormone and corticosterone in mice (Dronjak *et al.* 2004). This likely occurs because social isolation can elevate glucocorticoids, which can have detrimental effects if individuals are chronically stressed (DeVries, Glasper & Detillion 2003). Here, we demonstrate that social isolation in adult males does not just affect individuals within a lifetime, but also has detrimental consequences for offspring, leading to lower survival against a predator. The effect size of paternal isolation on offspring survival was moderate and biologically meaningful, with

offspring captured, on average, over three times faster when their father was isolated compared

to when he had a neighbor. These maladaptive effects are consistent with a previous study in rats,

where offspring of socially-isolated mothers showed cognitive impairments and increased

anxiety-like behaviors (McDonald *et al.* 2023). These transgenerational effects of social isolation

may arise because parental stress can hamper the ability of offspring to respond appropriately to

stress encountered within their own lifetime (Yehuda *et al.* 2008; Rodgers *et al.* 2013; Faraji *et*

al. 2017; McDonald *et al.* 2023), thus potentially leading to maladaptive behaviors in stressful

situations (e.g., predation events (Hellmann *et al.* 2020)). Although no studies (to our

knowledge) have looked at the transgenerational consequences of paternal isolation prior to

fertilization, our results are consistent with previous studies demonstrating that other forms of

paternal social stress, including maternal separation early in life (Gapp *et al.* 2014), social defeat

(Dietz *et al.* 2011), and social instability (Kong *et al.* 2021), alter offspring anxiety and

corticosterone levels. Consequently, although most TGP literature focuses on the effects of

parental exposure to ecological stress (e.g., heat, predation, toxins, disease), we suggest that

social stress can also have transgenerational consequences for offspring in ways that have not yet

been fully appreciated.

The mechanism by which paternal social isolation affects offspring is unknown, but is

clearly different than mechanisms of maternal social isolation, likely because those maternal

social effects seem to be passed down via glucocorticoids in eggs or during gestation (Walsh *et*

al. 2019; MacLeod *et al.* 2023; McDonald *et al.* 2023). Dietz *et al.* (2011) found that paternal

social defeat in mice increased offspring susceptibility to depression and anxiety-like behaviors,

but that this effect was largely not present when offspring were generated via IVF, suggesting

that these transgenerational effects were mediated by changes in maternal investment in her

offspring in response to perceived paternal quality (acquired during behavioral interactions

between the mother and father). Here, we did not allow for the effects of maternal compensation

on paternal phenotypes, as offspring were generated via in-vitro fertilization without behavioral

interactions between the mother and father. Further, as our fathers also did not interact

behaviorally with their offspring (no paternal care), our results demonstrate that these effects can

be mediated via epigenetic changes to sperm and/or seminal fluid alone. Future studies linking

the precise mechanism (e.g., small RNAs or methylation in sperm (Immler 2018)) to phenotypic

effects in offspring would be useful for predicting when and how paternal social experiences

affect offspring.

We found that offspring of predator-exposed fathers spent more time in the white side of

the scototaxis arena, suggesting that they were less anxious (Maximino *et al.* 2010). This is

consistent with previous studies demonstrating that paternal stress results in offspring with

reduced anxiety and HPA responsiveness (Rodgers *et al.* 2013; Gapp *et al.* 2014; He *et al.* 2016;

Brass *et al.* 2020), although results on this are mixed (Batchelor & Pang 2019). Further, offspring

of predator-exposed fathers tended to be consumed faster by the predator, which is consistent

with previous work in sticklebacks that found that paternal exposure to predation risk lowered

offspring survival against a predator (Hellmann *et al.* 2020). While the effects of paternal

isolation and predation exposure appeared to be additive, we found no evidence that the presence

of a conspecific buffered fathers against the effects of predation risk (i.e., no evidence of an

interaction between paternal predation exposure and paternal isolation). This was surprising,

given abundant evidence that isolated individuals show high levels of stress in response to

environmental stimuli, while individuals in social groups recover more quickly from or have a

reduced response to stressful experiences (Dronjak *et al.* 2004; Kikusui, Winslow & Mori 2006;

Ozbay *et al.* 2007; Culbert, Gilmour & Balshine 2019). Additionally, while some species do not

show signs of social buffering (Gilmour & Bard 2022), there is previous evidence of social

buffering in sticklebacks (Mommer & Bell 2013).

There may be several reasons why we did not see social buffering effects in this

experiment. First, it is possible that the presence of a conspecific did alter how fathers responded

to the presence of the predator, but not in a way that had consequences for offspring. Future

studies linking paternal physiology and behavior to phenotypic effects in offspring would be

useful for assessing this possibility. Second, it is possible that allowing for paternal care would

have enhanced the social buffering effect, as the presence of a caring parent can buffer offspring

against stressors (Sanchez, McCormack & Howell 2015). Finally, it is possible that the potential

for social buffering was limited in our experiment, given that we paired two unfamiliar males

together, both of whom were exposed to the predator. This would typically lead to weak effects

of social buffering, as previous studies have found stronger effects of social buffering when

individuals are paired with a known conspecific compared to an unknown conspecific

(Hennessy, Kaiser & Sachser 2009), when individuals are in groups compared to pairs (Stanton,

Patterson & Levine 1985; Kikusui, Winslow & Mori 2006), and when the partner is unstressed

compared to stressed (Kiyokawa *et al.* 2004; Kikusui, Winslow & Mori 2006). Consequently,

future studies manipulating the quality of the social environment could help us better understand

how transgenerational effects of social buffering may vary with the number, familiarity, and state

of the other conspecifics.

Conclusions

We have long known that the social environment is an important source of selection, and

that any given trait is not necessarily the property of a single individual because social

interactions will influence the traits and fitness of other individuals, even if those individuals

have never physically met (Fisher & McAdam 2017; Fisher *et al.* 2019). Currently, we view

transgenerational plasticity as a linear and relatively isolated relationship between parent and

offspring. Here, we demonstrate that the broader social environment can induce transgenerational

effects even in relatively asocial species such as stickleback, which do not form permanent social

groups. Although we did not find social buffering effects, previous work has shown that the

social environment can mediate parental response to ecological cues, providing an opportunity

for parents to learn about ecological change in their environment in ways that have cascading

effects on offspring (Afseth *et al.* 2022). Further, we demonstrate that social stress can combine

with ecological stress to have additive effects on offspring phenotypes. Future studies exploring

the within and transgenerational consequences of these socially-induced stress effects, either by

themselves or in conjunction with ecological stress, can help us better predict when and to what

extent individuals will be able to adaptively response to rapidly changing environments.

References

Afseth, C., Shim, A., Anderson, S., Bell, A. & Hellmann, J.K. (2022) Vertical transmission of horizontally-acquired social information in sticklebacks: Implications for transgenerational plasticity. *Proceedings of the Royal Society B,* **289,** 20220571.

Akkerman, K.C., Sattarin, A., Kelly, J.K. & Scoville, A.G. (2016) Transgenerational plasticity is sex-dependent and persistent in yellow monkeyflower (*Mimulus guttatus*). *Environmental Epigenetics,* **2**.

Bartolomucci, A., Palanza, P., Sacerdote, P., Ceresini, G., Chirieleison, A., Panerai, A.E. & Parmigiani, S. (2003) Individual housing induces altered immuno-endocrine responses to psychological stress in male mice. *Psychoneuroendocrinology,* **28,** 540-558.

Batchelor, V. & Pang, T.Y. (2019) HPA axis regulation and stress response is subject to intergenerational modification by paternal trauma and stress. *General and Comparative Endocrinology,* **280,** 47-53.

Beemelmanns, A. & Roth, O. (2016) Biparental immune priming in the pipefish *Syngnathus typhle*. *Zoology,* **119,** 262-272.

Bell, A.M. & Hellmann, J.K. (2019) An integrative framework for understanding the mechanisms and multigenerational consequences of transgenerational plasticity. *Annual Review of Ecology, Evolution and Systematics,* **50,** 97-118.

Ben-Shachar, M.S., Lüdecke, D. & Makowski, D. (2020) effectsize: Estimation of effect size indices and standardized parameters. *Journal of Open Software,* **5,** 2815.

Bonduriansky, R. & Head, M. (2007) Maternal and paternal condition effects on offspring phenotype in *Telostylinus angusticollis* (Diptera: Neriidae). *Journal of Evolutionary Biology,* **20,** 2379-2388.

Bonduriansky, R., Runagall-McNaull, A. & Crean, A.J. (2016) The nutritional geometry of parental effects: maternal and paternal macronutrient consumption and offspring phenotype in a neriid fly. *Functional Ecology,* **30,** 1675-1686.

Bouwhuis, S., Vedder, O. & Becker, P.H. (2015) Sex-specific pathways of parental age effects on offspring lifetime reproductive success in a long-lived seabird. *Evolution,* **69,** 1760-1771.

Brandl, H.B., Pruessner, J.C. & Farine, D.R. (2022) The social transmission of stress in animal collectives. *Proceedings of the Royal Society B,* **289,** 20212158.

Brass, K.E., Herndon, N., Gardner, S.A., Grindstaff, J.L. & Campbell, P. (2020) Intergenerational effects of paternal predator cue exposure on behavior, stress reactivity, and neural gene expression. *Hormones and Behavior,* **124,** 104806.

Bygren, L.O., Tinghög, P., Carstensen, J., Edvinsson, S., Kaati, G., Pembrey, M.E. & Sjöström, M. (2014) Change in paternal grandmothers´ early food supply influenced cardiovascular mortality of the female grandchildren. *BMC Genetics,* **15**.

Chen, E., Zielinski, C., Deno, J., Singh, R., Bell, A.M. & Hellmann, J.K. (2021) The specificity of sperm-mediated paternal effects in threespine sticklebacks. *Behavioral Ecology and Sociobiology,* **75,** 68.

Cockrem, J.F. (2013) Individual variation in glucocorticoid stress responses in animals. *General and Comparative Endocrinology,* **181,** 45-58.

Crane, A.L. & Ferrari, M.C.O. (2015) Minnows trust conspecifics more than themselves when faced with conflicting information about predation risk. *Animal Behaviour,* **100,** 184-190.

Culbert, B.M., Gilmour, K.M. & Balshine, S. (2019) Social buffering of stress in a group-living fish. *Proceedings of the Royal Society B,* **286,** 20191626.

da Costa, A.P., Leigh, A.E., Man, M.-S. & Kendrick, K.M. (2004) Face pictures reduce
behavioural, autonomic, endocrine and neural indices of stress and fear in sheep.
Proceedings of the Royal Society B, **271,** 2077-2084.
Dellinger, M., Zhang, W., Bell, A.M. & Hellmann, J.K. (2018) Do male sticklebacks use visual
and/or olfactory cues to assess a potential mate's history with predation risk? *Animal*
Behaviour, **145,** 151-159.
DeVries, A.C., Glasper, E.R. & Detillion, C.E. (2003) Social modulation of stress responses.
Physiology & Behavior, **79,** 399-407.
Dietz, D.M., LaPlant, Q., Watts, E.L., Hodes, G.E., Russo, S.J., Feng, J., Oosting, R.S., Vialou,
V. & Nestler, E.J. (2011) Paternal transmission of stress-induced pathologies. *Biological*
Psychiatry, **70,** 408-414.
Donelson, J.M., Wong, M., Booth, D.J. & Munday, P.L. (2016) Transgenerational plasticity of
reproduction depends on rate of warming across generations. *Evolutionary Applications,*
9, 1072-1081.
Dronjak, S., Gavrilović, L., Filipović, D. & Radojčić, M.B. (2004) Immobilization and cold
stress affect sympatho–adrenomedullary system and pituitary–adrenocortical axis of rats
exposed to long-term isolation and crowding. *Physiology & Behavior,* **81,** 409-415.
Faraji, J., Soltanpour, N., Lotfi, H., Moeeini, R., Moharreri, A.-R., Roudaki, S., Hosseini, S.A.,
Olson, D.M., Abdollahi, A.-A., Soltanpour, N., Mohajerani, M.H. & Metz, G.A.S. (2017)
Lack of social support raises stress vulnerability in rats with a history of ancestral stress.
Scientific Reports, **7,** 5277.
Faustino, A.I., Tacão-Monteiro, A. & Oliveira, R.F. (2017) Mechanisms of social buffering of
fear in zebrafish. *Scientific Reports,* **7,** 44329.
Fisher, D.N., Haines, J.A., Boutin, S., Dantzer, B., Lane, J.E., Coltman, D.W. & McAdam, A.G.
(2019) Indirect effects on fitness between individuals that have never met via an extended
phenotype. *Ecology Letters,* **22,** 697-706.
Fisher, D.N. & McAdam, A.G. (2017) Social traits, social networks and evolutionary biology.
Journal of Evolutionary Biology, **30,** 2088-2103.
Friard, O. & Gamba, M. (2016) BORIS: a free, versatile open-source event-logging software for
video/audio coding and live observations. *Methods in Ecology and Evolution,* **7,** 1325-
1330.
Fürtbauer, I. & Heistermann, M. (2016) Cortisol coregulation in fish. *Scientific Reports,* **6,**
30334.
Gapp, K., Jawaid, A., Sarkies, P., Bohacek, J., Pelczar, P., Prados, J., Farinelli, L., Miska, E. &
Mansuy, I.M. (2014) Implication of sperm RNAs in transgenerational inheritance of the
effects of early trauma in mice. *Nature Neuroscience,* **17,** 667-669.
Gilad, T. & Scharf, I. (2019) Separation between maternal and paternal effects on offspring
following exposure of adult red flour beetles to two stressors. *Ecological Entomology,* **44,**
494-501.
Gilmour, K.M. & Bard, B. (2022) Social buffering of the stress response: insights from fishes.
Biology Letters, **18,** 20220332.
Hagmayer, A., Lankheet, M.J., Bijsterbosch, J., van Leeuwen, J.L. & Pollux, B.J.A. (2022)
Maternal food restriction during pregnancy affects offspring development and swimming
performance in a placental live-bearing fish. *Journal of Experimental Biology,* **225,**
jeb242850.

He, N., Kong, Q.-Q., Wang, J.-Z., Ning, S.-F., Miao, Y.-L., Yuan, H.-J., Gong, S., Cui, X.-Z., Li, C.-Y. & Tan, J.-H. (2016) Parental life events cause behavioral difference among offspring: Adult pre-gestational restraint stress reduces anxiety across generations. *Scientific Reports,* **6,** 39497.

Hellmann, J.K., Bukhari, S.A., Deno, J. & Bell, A.M. (2020) Sex-specific plasticity across generations I: Maternal and paternal effects on sons and daughters. *Journal of Animal Ecology,* **89,** 2788-2799.

Hellmann, J.K., Carlson, E.R. & Bell, A.M. (2020) Sex-specific plasticity across generations II: Grandpaternal effects are lineage- and sex-specific. *Journal of Animal Ecology,* **89**.

Hellmann, J.K., Carlson, E.R. & Bell, A.M. (2021) The interplay between sperm-mediated and care-mediated paternal effects in threespined sticklebacks. *Animal Behaviour,* **179,** 267-277.

Hellmann, J.K., Keagy, J., Carlson, E.R., Kempfer, S. & Bell, A.M. (2024) Predator-induced transgenerational plasticity of parental care behaviour in male three-spined stickleback fish across two generations. *Proceedings of the Royal Society B,* **291,** 20232582.

Hellmann, J.K., Rogers, M.R. (2024). Data from: The transgenerational consequences of paternal social isolation and predation exposure in threespined sticklebacks.

Hennessy, M.B., Kaiser, S. & Sachser, N. (2009) Social buffering of the stress response: Diversity, mechanisms, and functions. *Frontiers in Neuroendocrinology,* **30,** 470-482.

Immler, S. (2018) The sperm factor: paternal impact beyond genes. *Heredity,* **121,** 239-247.

Kassambara, A. & Mundt, M. (2017) factoextra: extract and visualize the results of multivariate data analyses. R package version 1.0.5. https://CRAN.R-project.org/package=factoextra.

Kikusui, T., Winslow, J.T. & Mori, Y. (2006) Social buffering: relief from stress and anxiety. *Philosophical Transactions of the Royal Society B,* **361,** 2215-2228.

Kiyokawa, Y., Kikusui, T., Takeuchi, Y. & Mori, Y. (2004) Partner's stress status influences social buffering effects in rats. *Behavioral Neurosceince,* **118,** 798-804.

Kong, Q.-Q., Tian, X.-D., Wang, J., Yuan, H.-J., Ning, S.-F., Luo, M.-J. & Tan, J.-H. (2021) A next-generation sequencing study on mechanisms by which restraint and social instability stresses of male mice alter offspring anxiety-like behavior. *Scientific Reports,* **11,** 7952.

Koolhaas, J.M., Korte, S.M., De Boer, S.F., Van Der Vegt, B.J., Van Reenen, C.G., Hopster, H., De Jong, I.C., Ruis, M.A.W. & Blokhuis, H.J. (1999) Coping styles in animals: current status in behavior and stress-physiology. *Neuroscience & Biobehavioral Reviews,* **23,** 925-935.

Ludwig, S.C., Kapetanopoulos, K., Kotrschal, K. & Wascher, C.A.F. (2017) Effects of mate separation in female and social isolation in male free-living Greylag geese on behavioural and physiological measures. *Behavioural Processes,* **138,** 134-141.

MacLeod, K.J., English, S., Ruuskanen, S.K. & Taborsky, B. (2023) Stress in the social context: a behavioural and eco-evolutionary perspective. *Journal of Experimental Biology,* **226,** jeb245829.

MacLeod, K.J., Monestier, C., Ferrari, M.C.O., McGhee, K.E., Sheriff, M.J. & Bell, A.M. (2022) Predator-induced transgenerational plasticity in animals: a meta-analysis. *Oecologia,* **200,** 371-383.

Mashoodh, R., Franks, B., Curley, J.P. & Champagne, F.A. (2012) Paternal social enrichment effects on maternal behavior and offspring growth. *Proceedings of the National Academy of Sciences,* **109,** 17232-17238.

Maximino, C., Marques de Brito, T., Dias, C.A.G.d.M., Gouveia Jr, A. & Morato, S. (2010)
Scototaxis as anxiety-like behavior in fish. *Nature Protocols,* **5,** 209.
McDonald, R.J., Hong, N.S., Trow, J.S., Kaupp, C., Balog, R.J., Gokarn, L., Falkenberg, E.A.,
McCreary, K.J., Soltanpour, N., Witbeck, C., McKenna, A. & Metz, G.A.S. (2023)
Effects of maternal social isolation on adult rodent offspring cognition. *Scientific Reports,*
13, 7748.
McGhee, K.E., Feng, S., Leasure, S. & Bell, A.M. (2015) A female's past experience with
predators affects male courtship and the care her offspring will receive from their father.
Proceedings of the Royal Society B, **282**.
McQuaid, R.J., McInnis, O.A., Paric, A., Al-Yawer, F., Matheson, K. & Anisman, H. (2016)
Relations between plasma oxytocin and cortisol: The stress buffering role of social
support. *Neurobiology of Stress,* **3,** 52-60.
Mommer, B.C. & Bell, A.M. (2013) A test of maternal programming of offspring stress response
to predation risk in threespine sticklebacks. *Physiology and Behavior,* **122,** 222-227.
Moore, R.S., Kaletsky, R. & Murphy, C.T. (2019) Piwi/PRG-1 argonaute and TGF-α mediate
transgenerational learned pathogenic avoidance. *Cell,* **177,** 1827-1841.e1812.
Nylen, K.J., O'Hara, M.W. & Engeldinger, J. (2013) Perceived social support interacts with
prenatal depression to predict birth outcomes. *Journal of Behavioral Medicine,* **36,** 427-
440.
Ozbay, F., Johnson, D.C., Dimoulas, E., Morgan, C.A., Charney, D. & Southwick, S. (2007)
Social support and resilience to stress: from neurobiology to clinical practice. *Psychiatry
(Edgmont),* **4,** 35-40.
Pessato, A., McKechnie, A.E. & Mariette, M.M. (2022) A prenatal acoustic signal of heat affects
thermoregulation capacities at adulthood in an arid-adapted bird. *Scientific Reports,* **12,**
5842.
Potticary, A.L. & Duckworth, R.A. (2020) Multiple environmental stressors induce an adaptive
maternal effect. *American Naturalist,* **196,** 487-500.
Rodgers, A.B., Morgan, C.P., Bronson, S.L., Revello, S. & Bale, T.L. (2013) Paternal stress
exposure alters sperm microRNA content and reprograms offspring HPA stress axis
regulation. *Journal of Neuroscience,* **33,** 9003-9012.
Romero, L.M., Dickens, M.J. & Cyr, N.E. (2009) The reactive scope model — A new model
integrating homeostasis, allostasis, and stress. *Hormones and Behavior,* **55,** 375-389.
Sanchez, M.M., McCormack, K.M. & Howell, B.R. (2015) Social buffering of stress responses
in nonhuman primates: Maternal regulation of the development of emotional regulatory
brain circuits. *Social Neuroscience,* **10,** 512-526.
Schumann, S., Mozzi, G., Piva, E., Devigili, A., Negrato, E., Marion, A., Bertotto, D. &
Santovito, G. (2023) Social buffering of oxidative stress and cortisol in an endemic
cyprinid fish. *Scientific Reports,* **13,** 20579.
Sheriff, M.J., Krebs, C.J. & Boonstra, R. (2010) The ghosts of predators past: population cycles
and the role of maternal programming under fluctuating predation risk. *Ecology,* **91,**
2983-2994.
Stanton, M.E., Patterson, J.M. & Levine, S. (1985) Social influences on conditioned cortisol
secretion in the squirrel monkey. *Psychoneuroendocrinology,* **10,** 125-134.
Stein, L.R. & Bell, A.M. (2014) Paternal programming in sticklebacks. *Animal Behaviour,* **95,**
165-171.

Tariel, J., Plénet, S. & Luquet, É. (2020) Transgenerational plasticity in the context of predator-
prey interactions. *Frontiers in Ecology and Evolution,* **8,** 319.
Walsh, K., McCormack, C.A., Webster, R., Pinto, A., Lee, S., Feng, T., Krakovsky, H.S.,
O'Grady, S.M., Tycko, B., Champagne, F.A., Werner, E.A., Liu, G. & Monk, C. (2019)
Maternal prenatal stress phenotypes associate with fetal neurodevelopment and birth
outcomes. *Proceedings of the National Academy of Sciences,* **116,** 23996-24005.
Yehuda, R., Bell, A., Bierer, L.M. & Schmeidler, J. (2008) Maternal, not paternal, PTSD is
related to increased risk for PTSD in offspring of Holocaust survivors. *Journal of*
Psychiatric Research, **42,** 1104-1111.
Yusishen, M.E., Yoon, G.R., Bugg, W., Jeffries, K.M., Currie, S. & Anderson, W.G. (2020) Love
thy neighbor: Social buffering following exposure to an acute thermal stressor in a
gregarious fish, the lake sturgeon (*Acipenser fulvescens*). *Comparative Biochemistry and*
Physiology Part A: Molecular & Integrative Physiology, **243,** 110686.
633
634