Title: Monocytic fibrocyte-like cell enrichment and myofibroblastic adaptation causes nucleus pulposus fibrosis and associates with disc degeneration severity

Running title: Fibrocytes modify disc fibrosis and degeneration

Yi Sun[1][#], Yan Peng[2][#], Zezhuo Su[2][#], Kyle, KH SO[2], Qiu-ji Lu[2], Mao-jiang Lyu[1], Jianwei Zuo[1], Yong-can Huang[3], Zhi-ping Guan[3], Kenneth MC Cheung[2], Zhao-min Zheng[4], Xin-tao Zhang[1], Victor YL Leung[2]*

[1]Department of Sports Medicine, Peking University-Shenzhen Hospital, Shenzhen, Guangdong, China 518034

[2]Department of Orthopaedics and Traumatology, The University of Hong Kong, Hong Kong SAR, China

[3]Shenzhen Engineering Laboratory of Orthopaedic Regenerative Technologies, Department of Spine Surgery, Peking University Shenzhen Hospital, Shenzhen, Guangdong, China 518034

[4]Department of Spine Surgery, the First Affiliated Hospital, Sun Yat-sen University, Guangzhou, Guangdong, China 510062

[#] These authors contribute equally to this work.

*Corresponding Authors:

Dr. Victor YL Leung, PhD

Department of Orthopedics and Traumatology, Li Ka Shing Faculty of Medicine, The University of Hong Kong

L9-12, Lab Block, 21 Sassoon Road, Hong Kong SAR, China

Tel.: (852) 3917 6985;   Fax: (852) 2818 5210

E-mail: vicleung@hku.hk

Abstract (<250 words)

Fibrotic remodeling of nucleus pulposus (NP) leads to structural and mechanical anomalies of intervertebral discs that prone to degeneration, leading to low back pain incidence and disability. Emergence of fibroblastic cells in disc degeneration has been reported, yet their nature and origin remain elusive. In this study, we performed an integrative analysis of multiple single-cell RNA sequencing datasets to interrogate the cellular heterogeneity and fibroblast-like entities in degenerative human NP specimens. We found that disc degeneration severity is associated with an enrichment of fibrocyte-like phenotype, characterized by CD45 and collagen I dual positivity, and expression of myofibroblast marker α-smooth muscle actin. Refined clustering and classification distinguished the fibrocyte-like populations as subtypes in the NP cells - and immunocytes-clusters, expressing disc degeneration markers *HTRA1* and *ANGPTL4* and genes related to response to TGF-β. In injury-induced mouse disc degeneration model, fibrocyte-like cells were found recruited into the NP undergoing fibrosis and adopted a myofibroblast phenotype. Depleting the fibrocyte-like cells in CD11b-DTR mice in which monocytic lineages were ablated by diphtheria toxin could markedly attenuate fibrous modeling and myofibroblast formation in the NP of the degenerative discs, and prevent disc height loss and histomorphological abnormalities. Marker analysis supports that disc degeneration progression is dependent on a function of $CD45^+COL1A1^+$ and αSMA^+ cells. Our findings reveal that fibrocyte-like cells play a pivotal role in NP fibrosis and may therefore be a target for modifying disc degeneration and promoting its repair.

Introduction

The intervertebral discs (IVDs) are the largest avascular connective tissues in the body, playing a central role in spine movement. IVD degeneration (IDD) is associated with fibrosis of the nucleus pulposus (NP), the innermost colloidal core of IVDs, along with a loss of hydration and swelling pressure. The degeneration involves a degradation of hyaline extracellular matrix (ECM), in particular aggrecan and collagen II, and accumulation of fibrotic components including collagen I and small leucin-rich proteoglycans such as biglycan and fibronectin[1]. This leads to reduced tissue mechanical strength and hence disc prolapse under load, ultimately causing back pain and myelopathy[2]. Previous study has shown that transplantation of mesenchymal stromal cells (MSCs)[3] can alleviate IDD via inhibiting NP fibrosis. Understanding and controlling the profibrotic events may therefore hold the key to modifying the degeneration process. However, the mechanism of NP fibrosis and the origin of the fibroblastic population in the IDD remains elusive.

The primitive NP is initially populated by forkhead transcription factor (*FOXA2*)- or brachyury (*TBXT*)-expressing vacuolated cells derived from the notochord[4-6]. In humans, these vacuolated cells are gradually replaced by the chondrocyte-like cells after birth[7]. Tissue fibrosis relates to an excessive fibroblast and myofibroblast activity[8]. Reports have shown an increased number of $\alpha SMA^+/FSP1^+$/collagen I^+ fibroblastic cells in the degenerative IVDs[4,9]. In fact, NP cells (NPC) have a capacity to become $FSP1^+$ cells when exposed to excessive TGF-β[5,10] as well as undergo fibroblastic/myofibroblastic transition ($\alpha SMA^+/COL1A1^+/FSP1^+/FAP-\alpha^+$) in an injury-induced IDD model[4]. However, tracing study indicated that not all myofibroblastic cells are derived from local cells[4], implying alternative origins of effector cells for NP fibrosis. In line with this notion, single cell transcriptomic studies have revealed that the NP of human degenerative IVDs[7,11-14] and animal model of IDD[6] contain cell types other than IVD cells. These include endothelial cells,

myeloid granulocytic suppressor cells, CCR2$^+$ monocytes and derivatives of macrophages.

Consistent with the finding, studies showed that bone marrow cells could migrate into mechanically

overloaded IVD explants[15] and that focal enrichment of fibroblastic cells could be found near the

infiltrating blood vessels in the degenerative IVDs[16]. These evidences support that access of extrinsic

cells to the degenerated IVD may mediate NP fibrosis.

We hypothesized that cell sources other than resident disc cells can contribute to NP fibrosis. In this

study, we interrogated the cell hierarchy in the NP of degenerative IVDs through an integrative

analysis of the published single cell transcriptomes and characterization in human tissue and cells.

Our findings indicated an emergence of a small hematopoietic population that possess a fibrocyte-

like phenotype and myofibroblastic identity in the degenerative samples. Using a mouse model that

labels or depletes monocyte-derived cells, we analyzed the time course of the disc fibrocytes

appearance and identified their role in NP fibrosis and IDD progression. Our study implies a function

of circulatory fibrocytes in NP fibrosis and as a target for modifying IDD.

Results

Identification of myofibroblastic NPC in disc degeneration and NP fibrosis

We examined the cellular heterogeneity and fibroblastic cell entities by an integrative analysis of 4

published single-cell RNA sequencing (scRNA-seq) data derived from cells isolated from NP

tissues[7,12,14] or recognized as NPC [11]. The datasets consist of 6 non-degenerative human IVDs, 13

mildly (Pfirrmann II-III) and 12 severely (Pfirrmann IV-V) degenerative human IVDs (**Table 1**). A

number of existing NPC phenotypes were previously identified from these datasets, and their marker

genes were summarized in **Supplementary Table 1**. The meta-analysis identified 12 cell clusters

from a total of 192,184 cells and the cell types were annotated based on their cluster differentially

expressed genes (DEGs, Table SI). In addition to 5 NPC clusters, we also identified neutrophils

(*LYZ+*), myeloid derived suppressor cells (MDSC+; *MPO+*), macrophages (*CD68+*), T cells

(*TRBC2+*), endothelial cells (*CD34+* and *PECAM1+*), pericytes (*RGS5+* and *ACTA2+*), and

erythrocytes (*HBB+*) (**Fig. 1a** and **Supplementary Figure S1**). These clusters were mostly present

across the four datasets although their proportion and distribution varied (**Fig. 1b**). Clustered NPC

were identified by their expression of *ACAN*, *COL2A1* and *SOX9* (Table SI). The most abundant

NPC population was the chondrogenic cluster (ChonNP) (51%) which expressed *COL2A1*, *ACAN*

and *CYR61*. The regulatory cluster (RegNP) (21%) expressed *MMP3*, *GPX3* and *CHI3L2/1*. These

RegNP gene markers were also expressed in the inflammatory C1/3 chonNPC and regulatory NPC

reported by Han et al[7] and Tu et al[12], respectively (**Supplementary Table S1**). The fibroblastic

cluster (FibroNP; 8%) expressed *COL1A1*, *COL3A1*, *POSTN* and *ASPN*, which are known to

associate with tissue fibrosis. We found that the proportion of FibroNP significantly increased with

the severity of disc degeneration and became one of the major NPC populations (30%) in severely

degenerated NP (**Fig. 1c and Supplementary Figure S2**). The cycling cluster (CyclingNP) (1%)

expressed the cell proliferation genes including *TOP2A* and *STMN1*, containing two distinct subsets

expressing high level of either collagen I or II gene (*COL1A1^{high}* and *COL2A1^{high}*). The progenitor

cluster (ProgNP) (1%), distinguished by the expression of *PDGFRA* and *PLA2G2A*, was unique to

the non-degenerative samples as reported by Gan et al[14]. Endothelial cells (2%) were also found,

which highly expressed *PECAM1* and *CD34*. Erythrocytes (1%) were defined by gene expression of

HBB (**Fig. 1d**).

Contractile myofibroblasts are primary effector cells of tissue fibrosis[8]. FAPα and FSP1 mark the

active fibroblasts or myofibroblast precursors[17], and αSMA marks the mature myofibroblasts[18].

These three markers were found enriched in the FibroNP (**Fig. 1e**). More fibrillar collagens,

specifically thick collagen I and thin collagen III, were deposited in human degenerative IVDs,

which together with reduction of hyaline matrix aggrecan and collagen II, indicated NP fibrosis

(**Supplementary Figure S3**). Immunofluorescence showed that the number of cells expressing

αSMA (14.0% vs 48%), FAPα (31.4% vs 78.8%) and FSP1 (28.4% vs 65%) in the nNP was far

fewer than that in the dNP (**Fig. 1f**), indicating the augmented myofibroblast sprouting in dNP. In

line with our single-cell transcriptome analysis, cells from either dNP or nNP tissues were found to

contain a lower amount of αSMA$^+$ cells than FAPα$^+$ or FSP1$^+$ cells, which was further supported by

fluorescence-activated cell sorter (FACS) analysis of isolated NPC (**Fig. 1g**). FACS also

demonstrated an increased myofibroblast population in the dNP compared to the nNP. Notably, nNP

(88.2%) and dNP (93.1%) cells presented a comparable amount of cells expressing intracellular

collagen I (**Supplementary Figure S4**). However, under monolayer expansion, dNP cells became

more flattened and morphologically less refractive, and preferentially expressed higher levels of

myofibroblastic markers including *ACTA2*, *FAP* and *COL1A1* (**Supplementary Figure S5**). These

findings indicated that myofibroblastic cells became the dominant cell type in the degeneration and

supported their contribution to NP fibrosis.

Figure 1. Integrative single-cell transcriptome analysis and identification of fibroblastic NPC in human disc degeneration. (a) Identification of 12 distinct cell populations by unsupervised clustering. **(b)** Distribution of the cell populations in individual datasets. **(c)** Degeneration-related FibroNP abundance. **(d)** Signature gene expression for cell clusters defined in A. **(e)** Expression distribution of fibroblastic markers *ACTA2*, *FAP*, and *S100A4*. **(f)** Immunofluorescence and quantification of fibroblastic marker expression in human NP tissues (n=5). Scale bars: 50μm. Nuclei were counterstained with DAPI (Blue). **(g)** Representative flow cytometry plots to identify myofibroblastic NPC. Percentage of αSMA[+], Fap-α[+] or Fsp1[+] cells was averaged from three independent experiments. *$p < 0.05$, **$p < 0.01$, and ***$p < 0.001$ by two-tailed unpaired *t*-tests.

Disc degeneration contains fibroblastic NPC with hematopoietic features

Previous histological studies suggested that immune cells such as CD68[+] macrophages and

neutrophils may infiltrate into the NP during disc degeneration[19]. In the single cell analysis,

immunocyte-like populations could be found at all degeneration grades, expressing markers for

neutrophils (*LYZ* and *HLA-DRA*), macrophages (*CD163* and *CD68*), granulocyte-like myeloid

derived suppressor cells (G-MDSC) (*ITGAM* and *MPO*) and T cells (*TRAC* and *TRBC2*) (**Table SI**).

Interestingly, we found that *COL1A1*, *COL3A1* and *POSTN* were also expressed in the immunocytes,

predominantly the neutrophil and macrophage populations (**Fig. 2a**). These *COL1A1+* immunocytes

expressed both myeloid lineage genes (e.g. *S100A9*, *PTPRC* and *ITGB1*) and fibroblastic genes (e.g.

VIM, *TAGLN2* and *COL1A1*; (**Table SI**). Previous studies reported that monocytes contain a

subpopulation that co-expresses collagen I and III in addition to markers of hematopoietic cells

(CD11b, CD45 or CD34), being referred as fibrocytes[20-22]. The hematopoietic progenitor-associated

marker CD34 was immunolocalized with collagen I and contractile protein αSMA in the NP tissue

(**Supplementary Figure S6**). However, CD34 is also highly expressed in other cell types such as

endothelial cells and muscle satellite cells. *CD34[+]COL1A1[+]* or *CD34[+]ACTA2[+]* cells were in fact

mostly enriched in the endothelial cell and pericyte clusters. We further identified *COL1A1+*

subpopulations co-expressing *ITGAM* (coding CD11b) (1094 cells) or *PTPRC* (coding CD45) (1738

cells) (**Fig. 2b**). The abundance of these subpopulations increased with degeneration severity and

occupied up to 0.1% cell population dNP tissue (**Fig. 2c**). Immunostaining and flow cytometry

verified an increase of CD45[+]COLI[+] cells in the dNP tissue (16.4% vs 5.4%, **Fig. 2d**) or dNP-derived

cell culture (0.80% vs 0.30%, **Fig. 2e**) compared to the nNP controls. These CD45[+]COLI[+] cells

adopted an elongated morphology (insert, **Fig. 2d**). Based on the findings, we regarded the

CD45[+]COLI[+] cells as disc fibrocytes.

We identified disc fibrocytes within the NPC clusters, present mostly in the FibroNP (7.5%, as fib-fNP), ChonNP (4.2%, as fib-cNP) and RegNP (3.0%, as fib-rNP), in addition to immunocyte clusters of macrophages (35.4%, as fib-M), neutrophil (34.2%, as fib-N), T cell (11.4%, as fib-T) and G-MDSC (3.0%, fib-G) (**Supplementary Table S2**). By comparing these fibrocyte populations, we identified specific gene markers (**Table SI**) and examined their expression (**Figure 2f**). *GNB1* and gluconeogenesis-related gene *ALDOA*, together with marker genes of *COL1A1* and *ITGAM*, were enriched in both immunocytes- (fib-M/N) and NPC-clustered (fib-cNP/fNP) fibrocytes. Within immunocyte-clustered fibrocyte subpopulations (fib-M/N/T/G), a total of 47 genes (e.g. *PTPRC* and *CAPZA1*) were considerably enhanced and mainly linked to the actin organization and immune process (**Table 2 and Supplementary Figure S7**). *RORA* was dominantly expressed in fib-T. Fib-M-related genes *LGALS1* and *LGALS3BP* have been linked to monocyte-derived fibrocyte differentiation[23]. 43 genes (e.g. *COL3A1*, *ASPN* and *ITGB5*) that are commonly expressed in the three NPC-derived fibrocyte subsets are related to the response to TGF-β (**Supplementary Figure S7**). This finding is consistent with the notion that, similar to other fibrotic diseases, disc fibrocyte maturation may involve TGF-β-mediated endothelial- mesenchymal transition[24]. We also performed GO analysis for DEGs specific to the NPC-fibrocytes (**Table 3**). 35 genes distinctly enriched in fib-fNP were mostly related to cell adhesion function. 26 genes enriched in the Fib-cNP were related to matrix organization. Two degeneration-featured genes, *HTRA1*[25] and *ANGPTL4*[26] were distinctly expressed in the NPC-fibrocytes (**Fig. 2g**). Fibroinflammatory molecules of TGFBI and SPP1 from the disc fibrocytes were inferred to act on the FibroNP and stimulate the fibrotic process using CellChat analysis (**Supplementary Figure S8**). Fibrocyte-FibroNP communication may also rely on POSTN-ITGB5. Moreover, *ACTA2* was found expressed in *PTPRC*[+] FibroNP cells (**Fig. 2h**), supporting their potential in myofibroblastic differentiation[23,24]. Consistent with the scRNA-seq finding, co-immunostaining showed a markedly higher expression of CD45 and αSMA in the dNP compared to the nNP (**Fig. 2h**). We also observed αSMA[+] cells negative for CD45.

Figure 2. Identification of disc fibrocytes in the disc degeneration. (a) Featured gene plots of fibroblast markers of *COL1A1* and *COL3A1* in disc immunocytes. **(b)** Feature plots of expression distribution for monocyte lineage marker *ITGAM* and pan-hematopoietic marker *PTPRC* with *COL1A1*. **(c)** Fibrocyte abundance in disc degeneration. **(d)** Representative co-immunofluorescence micrographs of CD45 and Collagen I (COL1) in human NP tissues from scoliosis (nNP) and degenerative (dNP) IVDs, and quantification of co-expressing cells (n=5). **(e)** Flow cytometric evaluation of disc fibrocytes in isolated human NP cells. Representative flow cytometry was plotted and the percentages of CD45⁺COL1⁺ cells were labeled. **(f)** Heatmap showing and highlighting differentially expressed genes for each disc fibrocytes subtype. (**g**) Venn diagram and GO (gene ontology) functions of fibrocytes residing within three NP cells subclusters: FibroNP (fib-fNP), ChonNP (fib-cNP) and RegNP (fib-rNP). **(h)** UMAP distribution and immunodetection of *PTPRC*

(coding CD45)⁺ACTA2 (coding αSMA)⁺ cells sin disc NP tissues. Arrowhead: CD45⁺ αSMA⁺ cell;

sale bar: 50 μm. Insert: representative staining micrograph showing positive cellular signals; scale

bar: 12μm. *p < 0.05, **p < 0.01 by two-tailed unpaired *t*-tests.

Given the fibrocytes being a lineage of monocytes expressing CD11b, we studied their

temporospatial distribution in the puncture injury induced disc degeneration in CD11b-DTR mice, in

which the expression of EGFP-diphtheria toxin (DT) receptor fusion protein is controlled by the

ITGAM promoter, thereby directing transgene expression and expressing membrane-localized GFP in

monocytes/macrophages[22,27] (**Supplementary Figure S9**). Normal mouse NP, which contains

predominantly notochordal cells[4], was negative for GFP expression (**Supplementary Figure S9**).

Annulus puncture injury led to disc degeneration with NP fibrosis[3,4]. We observed an emergence of

GFP⁺ cells in the NP (~15% of NPC population) by day 3 after puncture, which continued to peak at

roughly 35% after 2 weeks and gradually subsided thereafter (**Fig. 3a**). GFP⁺ cells were also

observed in the annulus fibrosus (AF) (**Supplementary Figure S9**). We could observe GFP⁺ cells

that barely expressed CD11b in the NP at 2 weeks post-puncture (2wpp, **Fig. 3b**), possibly because

of a longer turnover of GFP compared to CD11b[27]. Up to 95% of the GFP⁺ NPC were CD45-

positive, indicating their myeloid origin (**Fig. 3c**). In line with previous report of fibrocyte

differentiation into myofibroblast, we observed αSMA expression in nearly 60% GFP⁺ cells at 4wpp

(**Fig. 3d**). The abundance of these dual-positive cells decreased significantly by 8wpp, indicating that

the emergence of fibrocytes might be transient or they underwent differentiation. Furthermore, we

detected CD45⁺COLI⁺ and CD45⁺αSMA⁺ cells in the punctured disc (**Fig. 4b&c**), and the amount of

CD45⁺αSMA⁺ NPC showed a two-fold reduction by 8wpp compared to 4wpp. These findings

collectively support the notion of fibrocyte infiltration into the degenerating discs and adopting a

myofibroblastic phenotype (CD45⁺αSMA⁺) during NP fibrosis.

Figure 3. Hematopoietic fibrocytes contribute to myofibroblastic NPCs. (a) Immunostaining and

quantification of GFP expression in the degenerative caudal discs of CD11b-DTR in 3 days (dpp) to

8 weeks post-puncture (wpp) window. **(b)** Co-immunostaining of hematopoietic marker CD45 and

(c) αSMA in CD11b⁺ cells with GFP with quantification (GFP⁺CD34⁺, **d**; GFP⁺αSMA⁺, **e**). Cellular

positivity counting was conducted in five representative sections, and expressed as the mean ± S.D.

Sale bar: 50 μm. * p<0.05, One-way ANOVA with Bonferroni post-hoc test.

Fibrocyte depletion rescues NP fibrosis

We tested whether depletion of CD11b-expressing cells by DT administration[27] could lead to a

reduction of fibrocytes in the puncture induced IDD. We first examined the frequency and dosage of

DT injection to enable extended depletion while minimizing lethality due to compromised immunity

(**Supplementary Table S3**). The number of CD11b⁺ monocytes were largely reduced in peripheral

blood (>50%) after twice administration of DT at 10μg/kg (**Supplementary Figure S10**). For the

IDD model, the first injection was performed 3 days prior to the puncture surgery (**Fig. 4a**). We

found a marked decrease of the GFP$^+$ cells in the NP at 2 and 8wpp, supporting the effect of DT on

the transgene expressing cells (**Supplementary Figure S10**). Interestingly, a lower number of

CD45$^+$COL1$^+$ cells was observed in the NP after the second dose of DT injection by 4wpp when

compared to vehicle control (**Fig. 4b**). The amount of CD45$^+$αSMA$^+$ NPC also tended to decrease

although the difference is not statistically significant (**Fig. 4c**). These findings indicate that DT

administration caused a depletion of monocytic fibrocytes in the NP which emerged during injury-

induced IDD.

We questioned if the loss of the myeloid derived fibrocytes leads to a modification of IDD and NP

fibrosis. Puncture induced IDD showed a significant disc height loss by 50% at 2wpp and 30% at

8wpp (**Fig. 4d**). DT administration could induce a gradual recovery and reduce the disc height loss

by >80% at 8wpp. Disc integrity assessed by FAST staining[3] showed a prominent improvement with

DT administration at both 4 and 8wpp, including reduced lacune formation, preserved NP-AF

boundary, and increased alcian blue intensity in the NP (**Fig. 4e**). Moreover, alcian blue staining in

the NP of the DT-injected mice was largely maintained, implying that the loss of proteoglycans was

prevented. The amount of myofibroblastic NPC positive for αSMA steadily increased after puncture

and took up to 76% of the whole population by 8wpp (**Fig. 4c**). DT administration could lower its

positivity to 41%. It is noteworthy that the majority of αSMA$^+$ cells detected in the punctured discs,

in particular in the DT-treated group, was negative for CD45 (>90%). Picrosirius red staining and

immunostaining of collagen I further consolidated a decrease in fibrillar collagens (**Fig. 4f-h**).

Pericellular expression of collagen I was noted in the NP after DT administration. Collectively, these

findings demonstrated that IDD progression is dependent on a function of myeloid derived cells

where the CD45$^+$COL1A1$^+$ fibrocytes and αSMA$^+$ myofibroblastic cells may mediate NP fibrosis.

Figure 4. Fibrocyte deficiency suppresses disc degeneration and NP fibrosis. (a) Time-line

illustration for DT administration and puncture surgery on CD11b-DTR mice. PBS serves as vehicle

control. **(b)** Co-staining of CD45 (red) with COL1 **(**green**)** or αSMA **(c,** green**)** and dual-positive cells

counts. αSMA positivity in NPC was also counted. **(d)** Time course assessment of disc height

changes (n=5 per group). **(e)** FAST staining and degeneration scoring based on compartmental

deformation in NP, annulus fibrosus (AF), endplate (EP) and interface. **(f)** Picrosirius red staining

and **(g)** their polarized microscopy. **(h)** Collagen I immunostaining. In immunofluorescence, nuclei

were counterstained by DAPI. Scale bar: 50μm, insert scale bar: 12μm. Dot line: NP region. Wpp:

weeks post-puncture; DP: disc puncture. Data are expressed as mean ± S.D. Two-way ANOVA with

Bonferroni post-hoc test, $* p<0.05$.

Discussion

NPC are conventionally defined as collagen II-, aggrecan- and Sox9-expressing cells. Recent studies

of single-cell transcriptome have indicated cell heterogeneity in the human NP, which contains cell

populations expressing notochordal, chondrogenic, and fibroblastic phenotypes[7,11-14]. Our meta-

analysis of the human disc scRNA-seq datasets has outlined the commonly featured NPC hierarchy

and stratified their fibroblastic subtypes. *COL1A1*[+] FibroNP is previously defined as

adhesion/fibroNPC[12] or C4 NPC[7]. The FibroNP population was not reported in Wang et al scRNA-

seq dataset, possibly because their cells were derived from the whole disc tissue without NP isolation

and therefore may have dominated by *COL1A1*[+] AF cells[11]. Nonetheless, they defined the FibroNP

genes (*POSTN*, *COL1A1*, *COL3A1* and *TMSB4X*) in MK167[+] progenitor and NP progenitor cells

(**Supplementary Table 1**), implying the fibroblast presence within the two NP subtypes. Han et al

found a small fraction of fibrochondrocyte progenitors in the NP, which partially constitute the

aforementioned two NP subsets. The regulatory NPC and C1/3 chonNPC subtypes reported by Tu[12]

and Han[7] are merged to the RegNP cluster in this study, sharing the expression of *MMP3*, *CHI3L2/1*

and *GPX3*. The CyclingNP cluster is a pool of *TOP*2A[+] cells fraction from the *COL2A1*[+] ChonNP or

COL1A1[+] FibroNP subsets, implying they may be derived from a common replicative progenitor

origin. Pseudotime ordering of the NP clusters and velocity analysis may help construct their

relationship.

The combination of collagen production (e.g. collagen I) and the expression of leukocytic common

antigen (CD45) or one of the hematopoietic or myeloid antigens (e.g. CD34 or CD11b) is considered

as a sufficient criterion for fibrocyte identity[21]. $CD34^+COL1A1^+$ cells are mainly enriched in the

endothelial cells and pericytes subpopulations (**Supplementary Figure S6**). *ITGAM* (coding CD11b)

or *PTPRC* (coding CD45) is therefore preferrable to CD34 as the marker of fibrocytes in human NP

tissues. Our single cell transcriptome analysis showed a wider UMAP distribution and higher number

of $PTPRC^+$ than $ITGAM^+$ cells, supporting the notion that CD11b is a more selective yet less

representative fibrocyte marker compared to the pan-hematopoietic markers Vav-1 or CD45[21].

Pronase digestion was commonly deployed for releasing single cells from the NP in scRNA-seq and

flow cytometry assays[28]. However, the digestion can also results in extensively membrane proteins

degradation, and thereby affect the antibody recognition of surface molecules[29]. This may in part

explain the discrepancy in measurements of $CD45^+COL1A1^+$ cells between in situ

immunohistochemistry and flow cytometry (**Fig. 2d&e**).

In addition to the NP clusters, $CD11b^+COL1A1^+$ or $CD45^+COL1A1^+$ cells are mostly located within

the macrophages, neutrophils and T cells clusters, implying the fibrocytes may be derived from

multiple sources or differentiation routes of the immune system. Blood immune cells are casually

linked with IDD, such as macrophages[19] and $CD39^+CD4^+$ T cells[30]. Interestingly, differentiation of

monocyte ($CD11b^+CD115^+Gr1^+$) into fibrocyte could be dependent on $CD4^+$ T cells activation[31].

Immune cells can infiltrate through focal defects in cartilaginous endplate (EP) and AF advancing the

fibroinflammatory process[32]. Monocytes are defined as $CD45^+CD11b^+$ cells and constitute the

myeloid-derived macrophages ($CD45^+CD11b^+F4/80^+$), neutrophils ($CD45^+CD11b^+HLA-DRA^+$) and

MDSCs ($CD45^+CD11b^+OLR1^+$)[12,13,21]. G-MDSCs, shown in cardiac fibrosis with myofibroblast

activation capabilities, contribute to T-cell suppression and ROS production in IDD[12]. Monocyte-

derived inflammatory macrophages could transit into myofibroblasts[33,34] and were identified in the

degenerative NP[19,35]. CD11b$^+$CD45$^+$ monocytes were previously reported to support the proliferation

and colony forming ability of in vitro-cultivated MSCs[36]. Whether the monocytic cells infiltrated into

the degenerating discs may exert a similar effect on local mesenchymal progenitors to promote

fibrogenesis or repair is not clear. Moreover, CCR2$^+$ monocytes have been identified in the

degenerative IVDs from herniation patients and constitute the majority of CD11b$^+$ and F4/80$^+$ cells[37].

However, as reported in myocardial infarction model[38], loss of CCR2 expression is required for

fibrocyte differentiation of the infiltrated monocytes. This is in line with our observation that the disc

fibrocytes are negative for *CCR2*. In a mouse model of renal fibrosis where CCR2 is depleted,

migration of circulating fibrocytes to the kidney was interfered[27]. Whether CCR2 plays a role in

monocyte infiltration and hence differentiation into disc fibrocytes await to be determined.

Fibrocytes could be derived not only from local monocyte differentiation but also migrated from

peripheral blood in the form of partially differentiated collagen-producing cells[21,27]. Recruitment of

monocytes and peripheral blood fibrocytes may rely on disc neo-vascularization. Normally, blood

vessels recede from the outer AF and the EP in adults. However, under degenerative conditions the

NP can become re-vascularized and blood vessel infiltration is frequently observed and spatially

associated with immunocytes infiltration[39]. Most of the degenerative NP tissues in this study for

scRNA-seq analysis were dissected from extruded NP of herniated discs that normally show

pronounced neo-vascularization[40]. The increased number of GFP$^+$ cells found in the AF and EP of

the punctured CD11b-DTR discs may indicate the infiltration through the AF and EP routes.

Pericytes (CD34$^+$αSMA$^+$) were immunodetected in the degenerative human NP (**Supplementary**

Figure S6), supporting the presence of neo-vascularization. In fact, pericytes can contribute to scar-

forming myofibroblasts in kidney and subretinal fibrosis[8]. Studying disc vascularization and its

relation to the emergence of disc fibrocytes may provide further insights into the pathway of NP

fibrosis.

Trafficking fibrocytes express chemokine receptors CCR5[41], CCR7[21] and CXCR4[21,42]. Studies have

reported increased CCR5 expression in AF and NP cells[43] and chemokines CCL2/7 in degenerated

NP[44]. Bone marrow cells, such as $CD146^+$ MSCs[45], may have a potential to migrate into the NP,

possibly by control of SDF1[46]. CCL5 could interact with CCR1/3/5 in NPC and was up-regulated

under inflammatory stimuli[47]. CXCR4 is a receptor of SDF1[48] and CXCL12[42], which was found

upregulated in degenerated discs and implicated in disc angiogenesis[48] and cellular apoptosis[49] via

PI3K/NF-κB pathway. Interestingly, we find that the disc fibrocytes highly express *CXCR4* and

CCR1 (**Table 2**), and may communicate with the ProgNP population via the CXCL12-CXCR4 axis.

Future study may test if CXCL12 has a function to recruit $CD45^+COL1A1^+$ fibrocytes.

Tissue fibrosis is mediated by myofibroblast from local or extrinsic sources[8] and fibrocytes may

serve as one of the progenitors of myofibroblasts[21,24]. Expression of αSMA is the hallmark of mature

myofibroblasts and essential to their cellular contractility[18]. Expression of αSMA in NPC may

therefore indicate their myofibroblastic property[24]. FAP-α is a well-known biomarker of

(myo-)fibroblasts, and together with FSP1 it can label most of the collagen I-producing cells in the

bone[50]. FSP1 marks an active fibroblast population and represents a non-overlapping fibroblast entity

to $αSMA^+$ subtype in heart, kidney and skin fibrosis disease[51]. FSP1 is also expressive in atypical

myeloid-derived F4/80-positive inflammatory macrophages in liver injury[52]. FSP1 was found

expressed in the $COL2A1^+$ ChonNP cells in addition to the $COL1A1^+$ FibroNP cells.

Our results demonstrate αSMA expression in the CD45[+] and CD11b[+] fibrocytes, supporting the

conversion of fibrocytes to myofibroblasts in NP fibrosis. Yet, a vast number of αSMA[+]

myofibroblastic NPC in both human degenerative IVDs and mouse punctured discs are not positive

for CD45 and CD11b. In the CD11b-DTR mice where the monocytes and their derived lineages

(including fibrocytes) were depleted[27], αSMA expressing cells could still be observed (>55%, 2 wpp

in **Fig. 4d**) in the degenerative NP. This may be due to the existence of non-hematopoietic origins of

the myofibroblast pool, such as those derived from resident NPC[4]. Alternatively, fibrocytes may lose

CD45 and CD11b expression during fibrocyte differentiation[33]. We also note the presence of residual

CD45[+]COL1[+] cells and GFP[+] cells after DT administration. This likely arises from the incomplete

monocyte ablation and a continuation of fibrocyte recruitment.

Fibrocyte-to-myofibroblast conversion acquires activation of TGF-β/SMAD signaling[53], and

interleukin-18 receptor 1[54] and muscarinic receptor M3[55] are shown to promote the fibrocyte

formation and contractile function. Moreover, fibrocytes can secret soluble factors such as periostin[56]

and TIMP1[57] and semaphorin-7A[58] to regulate myofibroblast activity and collagen expression in

pulmonary and intestinal fibrosis[20]. Class 3 semaphorins expression are found associated with disc

innervation and angiogenesis[59]. Interestingly, TIMP1[60] and POSTN[61] are up-regulated in the

degenerated discs and enriched in the FibroNP. Our CellChat data suggest that POSTN may interact

with ITGB5 of disc fibrocytes. Whether disc fibrocytes regulate NP fibrosis through a paracrine

manner awaits to be addressed. Furthermore, the NPC-enriched disc fibrocytes express *HTRA1* and

ANGPTL4. HTRA1 is a proteolytic enzyme for degrading the matrix either on its own[62] or by

upregulating ADAMTS-5[25] or MMPs, whereas expression of ANGPTL4 is positively associated with

IDD severity [26]. Notably, our data suggest that ANGPTL4 is a major mediator of the intercellular

communication between the disc fibrocytes and the CyclingNP, RegNP and FibroNP subpopulations.

Their regulatory role in the fibrocytes and NP fibrosis worths to be investigated.

In conclusion, this study implicates a function of monocytic fibrocytes in NP fibrosis and IDD

progression, shedding a new insight to cell fate control of disc immunocytes and their modulatory

effect on IVD degeneration. Understanding the effects and regulation of the fibrocytes may

potentiate the development of new diagnostic and therapeutic strategies for modifying IDD.

Materials and methods

scRNA-seq data acquisition and downstream analysis

Four single cell RNA sequencing data sets (GSE160756, GSE165722, CNP0002664,

PRJCA014236) were downloaded and processed for next-step analysis. 31 individual samples were

rearranged according to Pfirrmann grade, which Pfirrmann grade I samples were classified as non-

degenerative NP, Pfirrmann grade II/III as mildly degenerated NP, and Pfirrmann grade IV/V as

severely degenerated NP (**Table 1**). Seurat package (version: 3.1.4, https://satijalab.org/seurat/) was

used for cell normalization and regression according to the UMI counts of each sample and percent

of mitochondria rate to obtain the scaled data. MNN (mutual nearest neighbor) was used to remove

potential batch effect. The top 10 principles were used for unsupervised UMAP construction via

graph-based cluster method (resolution = 0.8), and marker genes were determined with p-value <

0.01 and log(fold-change) > 0.25 by the FindAllMarkers function. CellChat analysis was conducted

to identify potential intercellular interactions, and probability values for each interaction were

calculated. Significant mean and cell communication significance (p-value < 0.01) were calculated

based on the interaction and the normalized cell-matrix achieved by Seurat normalization.

Human samples

All procedures described in the present study were approved by the Institutional Ethics Review

Board of The University of Hong Kong. IVD tissues were obtained from patients under informed

consent. Non-degenerative (n) and degenerative (d) IVDs were harvested from respectively

adolescent idiopathic scoliosis (n=7) and lumbar disc degeneration patients (n=9). The age of the

patients ranged from 14 to 72 years, and tissue was evaluated with the Pfirrmann classification

system (**Supplementary Table 4**).

Cell isolation and culture

Jelly-like NP tissues were carefully dissected from the cartilaginous EP and lamellar-structured AF.

In severely degenerated IVDs (grade IV-V), NP tissues were more fibrotic and hardly distinguishable

from inner AF, so we harvested the tissue presumably at the central region of the NP. Primary NPC

were extracted by sequential enzymes digestion of pronase (0.25%, Roche Diagnostics) and

collagenase II (600U/ml, Worthington Biochemical)[28], and maintained in high-glucose Dulbecco's

modified Eagle's medium (DMEM) (Invitrogen) supplemented with 10% fetal bovine serum (FBS,

Biosera) and 1% penicillin/streptavidin (P/S, Invitrogen) at 37 °C and 5% CO_2.

Flow cytometry

Newly extracted NPC were washed by PBS and re-suspended in PBS. Cells were fixed in 4%

paraformaldehyde (PFA) and blocked by 5% BSA. Permeabilization with 0.1% triton X-100 (Sigma-

Aldrich) was applied where necessary. Cells were then treated with primary and secondary

antibodies, and signals from cells labeled with conjugated fluorophores were detected by using BD

FACS CantoII Analyzer (BD Biosciences). The antibodies used for different flow cytometry analysis

are listed in **Supplementary Table S5**. Appropriate IgG control fluorescence compensation was

applied to avoid false positive signals. Data were further analyzed by BD FACS Diva software (BD

Bioscience).

Real-time quantitative PCR

Total RNA was prepared using RNeasy mini kit (Invitrogen) and cDNA synthesis was performed

using High-capacity RNA-to-cDNA kit (Applied Biosystems). PCR was performed using PowerUp

SYBR green master mix (Applied Biosystems), and data are presented as expression levels relative

to *GAPDH* using the $2^{-\Delta CT}$ method. The primer sequences were listed in **Supplementary Table S6**.

Mouse disc puncture model

Tg (ITGAM-DTR/eGFP)34Lan/J mice express membrane localized GFP fluorescence with diphtheria toxin (DT) receptor under the control of CD11b and were used for disc puncture. The animal experiments were approved by The University of Hong Kong Committee on the Use of Live Animals in Teaching and Research (CULATR). CD11b-DTR mice (female, 12 weeks of age, *n*=5 per group and time points) received needle puncture at levels of C5/6 and C7/8 of tail IVDs as described previously[4]. At four days prior to the timepoints and every other week after the puncture surgery, these mice were subjected to intraperitoneal injection of saline or DT (25μg/kg body weight) (Sigma-Aldrich). At assigned end points, mice were euthanized by intravenous injection of sodium pentobarbital (1.2 g/kg), and intact discs with attached vertebral bones were harvested for histochemistry. Peripheral bloods were harvested at days 0, 3, and 7 after DT injection, and number of GFP$^+$ monocytes was measured by fluorescence-activated cell sorting (FACS).

Disc height measurement

Anterior-to-posterior radiographs of tail discs of CD11b-DTR mice or C57BL/6J were taken using a digital radiography machine (Siemens) with an exposure at 25 kV, for 5 seconds. Disc height index was calculated as previously described [4] and presented as percent relative to the unoperated level of tail disc C6/7. Images were analyzed biweekly in all groups up to 8 weeks.

Histology and immunostaining

Samples from human and CD11b-DTR mice tissues were fixed with 4% paraformaldehyde in phosphate buffer for 24 h. Decalcification was performed for mouse IVDs using mouse solution. Paraffin-embedded tissue blocks of human NP and mice IVDs were sectioned at 7 μm. For histological staining, sections were deparaffinized and rehydrated, and subjected to FAST staining (Alcian blue, Safranin O, Fast green and Tartrazine stain) (Sigma-Aldrich) as previously described[63], or Masson's trichrome stain and Picrosirius red stain following standard procedures. Polarized microscopy was combined with picrosirius red staining to observe the collagens network in human NP tissue and mouse

IVDs[64]. According to the histomophological changes, disc degeneration in CD11b-DTR mice was

scored with reference to Tam's scoring system[63].

For immunofluorescence, antigen retrieval was performed following rehydration by incubating with

0.8% hyaluronidase (Sigma-Aldrich) at 37°C for 30 min and antigen retrieval buffer (100 mM Tris,

5% EDTA, pH 9.5) at 95°C for 10 min. The sections were blocked in protein block solution (DAKO)

for 60 min before incubating with antibodies. Fluorophore conjugated antibodies were used to enable

fluorescence detection. Information for these primary and secondary antibodies were summarized in

Supplementary Table S5. The sections were mounted with VECTASHIELD Mounting Medium

with DAPI for nuclei staining. Isotype-matched mouse, rabbit or rat IgG was used as negative

controls. Tissue sections were imaged by fluorescence microscope (Nikon Eclipse E600), and

fluorescent positive cells in five fields of view (FOV) were manually counted in ImageJ with

reference to total number of cells.

Data analysis

Data are presented as mean $\pm$ S.D. of four to six experiments, as indicated in respective figure

legends. Unpaired t-test (two tailed) was used to determine significant differences. Comparisons

among multiple groups were assessed using one-way ANOVA with Bonferroni post-hoc test. A p

value < 0.05 was considered statistically significant.

Acknowledgements

This work was jointly supported by the General Research Fund (17121619) of the Research Grant

Council of Hong Kong and the Scientific Research Foundation of PEKING UNIVERSITY

SHENZHEN HOSPITAL (KYQD202100X).

Competing Interests statement

The authors have no competing interest to declare.

Author contributions

YS, YP and ZS contributed equally to this manuscript. YS contributed to the study design, data

collection and interpretation, producing figures and obtaining the fund. ZS, KKS and YS conducted

the bioinformatic analysis. YS, YP, QL and ML performed the experiments and helped in data

analysis. YS, YP, ZS and VYL wrote the manuscript. JZ, YH, ZG, KMC, ZZ and XZ collected

samples and clinical data, and edited the manuscript. KMC, ZZ and XZ reviewed the data and

provided critical data interpretation. VYL conceptualized and supervised the whole project, provided

supports and coordinated the collaboration. All authors have read and approved the manuscript.

References

1. Sun, Y., Lyu, M., Lu, Q., Cheung, K. & Leung, V. Current Perspectives on Nucleus Pulposus Fibrosis in Disc Degeneration and Repair. *Int J Mol Sci* **23**(2022).
2. Zhang, S., *et al.* The role of structure and function changes of sensory nervous system in intervertebral disc-related low back pain. *Osteoarthritis Cartilage* **29**, 17-27 (2021).
3. Leung, V.Y., *et al.* Mesenchymal stem cells reduce intervertebral disc fibrosis and facilitate repair. *Stem Cells* **32**, 2164-2177 (2014).
4. Au, T.Y.K., *et al.* Transformation of resident notochord-descendent nucleus pulposus cells in mouse injury-induced fibrotic intervertebral discs. *Aging Cell* **19**, e13254 (2020).
5. Zhang, Y., *et al.* Directed Differentiation of Notochord-like and Nucleus Pulposus-like Cells Using Human Pluripotent Stem Cells. *Cell Rep* **30**, 2791-2806 e2795 (2020).
6. Zhou, T., *et al.* Spatiotemporal Characterization of Human Early Intervertebral Disc Formation at Single-Cell Resolution. *Adv Sci (Weinh)* **10**, e2206296 (2023).
7. Han, S., *et al.* Single-Cell RNA Sequencing of the Nucleus Pulposus Reveals Chondrocyte Differentiation and Regulation in Intervertebral Disc Degeneration. *Front Cell Dev Biol* **10**, 824771 (2022).
8. Kuppe, C., *et al.* Decoding myofibroblast origins in human kidney fibrosis. *Nature* **589**, 281-286 (2021).
9. Lv, F.J., *et al.* Matrix metalloproteinase 12 is an indicator of intervertebral disc degeneration co-expressed with fibrotic markers. *Osteoarthritis Cartilage* **24**, 1826-1836 (2016).
10. Chen, C., *et al.* Autologous fibroblasts induce fibrosis of the nucleus pulposus to maintain the stability of degenerative intervertebral discs. *Bone Res* **8**, 7 (2020).
11. Wang, D., *et al.* Single-cell transcriptomics reveals heterogeneity and intercellular crosstalk in human intervertebral disc degeneration. *iScience* **26**, 106692 (2023).

12. Tu, J*., et al.* Single-Cell Transcriptome Profiling Reveals Multicellular Ecosystem of
Nucleus Pulposus during Degeneration Progression. *Adv Sci (Weinh)* **9**, e2103631 (2022).

13. Ling, Z*., et al.* Single-Cell RNA-Seq Analysis Reveals Macrophage Involved in the
Progression of Human Intervertebral Disc Degeneration. *Front Cell Dev Biol* **9**, 833420
(2021).

14. Gan, Y*., et al.* Spatially defined single-cell transcriptional profiling characterizes diverse
chondrocyte subtypes and nucleus pulposus progenitors in human intervertebral discs.
Bone Res **9**, 37 (2021).

15. Sakai, D*., et al.* Migration of bone marrow-derived cells for endogenous repair in a new
tail-looping disc degeneration model in the mouse: a pilot study. *Spine J* **15**, 1356-1365
(2015).

16. Melrose, J*., et al.* Spatial and temporal localization of transforming growth factor-beta,
fibroblast growth factor-2, and osteonectin, and identification of cells expressing alpha-
smooth muscle actin in the injured anulus fibrosus: implications for extracellular matrix
repair. *Spine (Phila Pa 1976)* **27**, 1756-1764 (2002).

17. Esteve, D*., et al.* Lobular architecture of human adipose tissue defines the niche and fate
of progenitor cells. *Nat Commun* **10**, 2549 (2019).

18. Hinz, B. & Lagares, D. Evasion of apoptosis by myofibroblasts: a hallmark of fibrotic
diseases. *Nat Rev Rheumatol* **16**, 11-31 (2020).

19. Li, X.C*., et al.* Investigation of macrophage polarization in herniated nucleus pulposus of
patients with lumbar intervertebral disc herniation. *J Orthop Res* **41**, 1335-1347 (2023).

20. Abe, S*., et al.* Functional analysis of human fibrocytes derived from monocytes reveals
their profibrotic phenotype through paracrine effects. *J Med Investig* **67**, 102-112 (2020).

21. Reinhardt, J.W. & Breuer, C.K. Fibrocytes: A Critical Review and Practical Guide. *Front
Immunol* **12**, 784401 (2021).

22. Ozono, Y*., et al.* Neoplastic fibrocytes play an essential role in bone marrow fibrosis in
Jak2V617F-induced primary myelofibrosis mice. *Leukemia* **35**, 454-467 (2021).

23. White, M.J., Roife, D. & Gomer, R.H. Galectin-3 Binding Protein Secreted by Breast Cancer
Cells Inhibits Monocyte-Derived Fibrocyte Differentiation. *J Immunol* **195**, 1858-1867
(2015).

24. Abu El-Asrar, A.M*., et al.* Myofibroblasts in proliferative diabetic retinopathy can originate
from infiltrating fibrocytes and through endothelial-to-mesenchymal transition
(EndoMT). *Exp Eye Res* **132**, 179-189 (2015).

25. Li, D*., et al.* HtrA1 upregulates the expression of ADAMTS-5 in HNPCs via the ERK/NF-
kappaB/JNK signaling pathway. *Am J Transl Res* **11**, 5114-5121 (2019).

26. Liu, F.J*., et al.* Expression of ANGPTL4 in Nucleus Pulposus Tissues Is Associated with
Intervertebral Disc Degeneration. *Dis Markers* **2021**, 3532716 (2021).

27. Reich, B*., et al.* Fibrocytes develop outside the kidney but contribute to renal fibrosis in a
mouse model. *Kidney Int* **84**, 78-89 (2013).

28. Basatvat, S*., et al.* Harmonization and standardization of nucleus pulposus cell extraction
and culture methods. *JOR Spine* **6**, e1238 (2023).

29. Srakocic, S*., et al.* Proposed practical protocol for flow cytometry analysis of microglia

from the healthy adult mouse brain: Systematic review and isolation methods' evaluation.
Front Cell Neurosci **16**, 1017976 (2022).

30. Qin, C.*, et al.* Causal relationship between the blood immune cells and intervertebral disc
degeneration: univariable, bidirectional and multivariable Mendelian randomization.
Front Immunol **14**, 1321295 (2023).

31. Niedermeier, M.*, et al.* CD4+ T cells control the differentiation of Gr1+ monocytes into
fibrocytes. *Proc Natl Acad Sci U S A* **106**, 17892-17897 (2009).

32. Bermudez-Lekerika, P.*, et al.* Immuno-Modulatory Effects of Intervertebral Disc Cells.
Front Cell Dev Biol **10**, 924692 (2022).

33. Sinha, M.*, et al.* Direct conversion of injury-site myeloid cells to fibroblast-like cells of
granulation tissue. *Nat Commun* **9**, 936 (2018).

34. Meng, X.M.*, et al.* Inflammatory macrophages can transdifferentiate into myofibroblasts
during renal fibrosis. *Cell Death Dis* **7**, e2495 (2016).

35. Burt, K.G., Kim, M.K.M., Viola, D.C., Abraham, A.C. & Chahine, N.O. Nuclear Factor Kappa B
Over-Activation in the Intervertebral Disc Leads to Macrophage Recruitment and Severe
Disc Degeneration. *bioRxiv* (2023).

36. Hong, H.S., Ahn, W. & Son, Y. CD45+/CD11b+monocytes are required for mesenchymal
stem cell proliferation. *Tissue Engineering and Regenerative Medicine* **11**, 224-229 (2014).

37. Jin, L.*, et al.* CCR2 monocytes as therapeutic targets for acute disc herniation and
radiculopathy in mouse models. *Osteoarthritis Cartilage* (2023).

38. Mouton, A.J.*, et al.* Mapping macrophage polarization over the myocardial infarction time
continuum. *Basic Res Cardiol* **113**, 26 (2018).

39. Zhang, S.*, et al.* Inhibiting Heat Shock Protein 90 Attenuates Nucleus Pulposus Fibrosis
and Pathologic Angiogenesis Induced by Macrophages via Down-Regulating Cell
Migration-Inducing Protein. *Am J Pathol* **193**, 960-976 (2023).

40. Yasuma, T., Arai, K. & Yamauchi, Y. The histology of lumbar intervertebral disc herniation.
The significance of small blood vessels in the extruded tissue. *Spine (Phila Pa 1976)* **18**,
1761-1765 (1993).

41. Ishida, Y.*, et al.* Essential roles of the CC chemokine ligand 3-CC chemokine receptor 5 axis
in bleomycin-induced pulmonary fibrosis through regulation of macrophage and
fibrocyte infiltration. *Am J Pathol* **170**, 843-854 (2007).

42. Phillips, R.J.*, et al.* Circulating fibrocytes traffic to the lungs in response to CXCL12 and
mediate fibrosis. *J Clin Invest* **114**, 438-446 (2004).

43. Liu, W.*, et al.* Annulus fibrosus cells express and utilize C-C chemokine receptor 5 (CCR5)
for migration. *Spine J* **17**, 720-726 (2017).

44. Phillips, K.L.*, et al.* The cytokine and chemokine expression profile of nucleus pulposus
cells: implications for degeneration and regeneration of the intervertebral disc. *Arthritis
Res Ther* **15**, R213 (2013).

45. Wangler, S.*, et al.* CD146/MCAM distinguishes stem cell subpopulations with distinct
migration and regenerative potential in degenerative intervertebral discs. *Osteoarthritis
Cartilage* **27**, 1094-1105 (2019).

46. Leite Pereira, C.*, et al.* Stromal Cell Derived Factor-1-Mediated Migration of Mesenchymal

Stem Cells Enhances Collagen Type II Expression in Intervertebral Disc. *Tissue Eng Part A* (2018).

47. Gruber, H.E.*, et al.* Production and expression of RANTES (CCL5) by human disc cells and modulation by IL-1-beta and TNF-alpha in 3D culture. *Exp Mol Pathol* **96**, 133-138 (2014).

48. Zhang, H.*, et al.* SDF1/CXCR4 axis facilitates the angiogenesis via activating the PI3K/AKT pathway in degenerated discs. *Mol Med Rep* **22**, 4163-4172 (2020).

49. Liu, Z.*, et al.* SDF-1/CXCR4 axis induces apoptosis of human degenerative nucleus pulposus cells via the NF-kappaB pathway. *Mol Med Rep* **14**, 783-789 (2016).

50. Chen, Y.*, et al.* Type-I collagen produced by distinct fibroblast lineages reveals specific function during embryogenesis and Osteogenesis Imperfecta. *Nat Commun* **12**, 7199 (2021).

51. Saraswati, S., Marrow, S.M.W., Watch, L.A. & Young, P.P. Identification of a pro-angiogenic functional role for FSP1-positive fibroblast subtype in wound healing. *Nat Commun* **10**, 3027 (2019).

52. Osterreicher, C.H.*, et al.* Fibroblast-specific protein 1 identifies an inflammatory subpopulation of macrophages in the liver. *Proc Natl Acad Sci U S A* **108**, 308-313 (2011).

53. Mitsuhashi, A.*, et al.* Identification of fibrocyte cluster in tumors reveals the role in antitumor immunity by PD-L1 blockade. *Cell Rep* **42**, 112162 (2023).

54. Niu, X.H.*, et al.* IL-18/IL-18R1 promotes circulating fibrocyte differentiation in the aging population. *Inflamm Res* **69**, 497-507 (2020).

55. Henrot, P.*, et al.* Muscarinic receptor M3 activation promotes fibrocytes contraction. *Front Pharmacol* **13**, 939780 (2022).

56. Ashley, S.L., Wilke, C.A., Kim, K.K. & Moore, B.B. Periostin regulates fibrocyte function to promote myofibroblast differentiation and lung fibrosis. *Mucosal Immunol* **10**, 341-351 (2017).

57. Kuroda, N.*, et al.* Infiltrating CCR2(+) monocytes and their progenies, fibrocytes, contribute to colon fibrosis by inhibiting collagen degradation through the production of TIMP-1. *Sci Rep* **9**, 8568 (2019).

58. Gan, Y.*, et al.* Role of semaphorin 7a signaling in transforming growth factor beta1-induced lung fibrosis and scleroderma-related interstitial lung disease. *Arthritis Rheum* **63**, 2484-2494 (2011).

59. Binch, A.L.*, et al.* Class 3 semaphorins expression and association with innervation and angiogenesis within the degenerate human intervertebral disc. *Oncotarget* **6**, 18338-18354 (2015).

60. Vo, N.V.*, et al.* Expression and regulation of metalloproteinases and their inhibitors in intervertebral disc aging and degeneration. *Spine J* **13**, 331-341 (2013).

61. Zhu, D.*, et al.* Periostin: An Emerging Molecule With a Potential Role in Spinal Degenerative Diseases. *Front Med (Lausanne)* **8**, 694800 (2021).

62. Furtwangler, T., Chan, S.C., Bahrenberg, G., Richards, P.J. & Gantenbein-Ritter, B. Assessment of the matrix degenerative effects of MMP-3, ADAMTS-4, and HTRA1, injected into a bovine intervertebral disc organ culture model. *Spine (Phila Pa 1976)* **38**, E1377-1387 (2013).

656 63. Tam, V*., et al.* Histological and reference system for the analysis of mouse intervertebral
657   disc. *J Orthop Res* **36**, 233-243 (2018).
658 64. Lattouf, R*., et al.* Picrosirius red staining: a useful tool to appraise collagen networks in
659   normal and pathological tissues. *J Histochem Cytochem* **62**, 751-758 (2014).

Table 1 Demographic data. A total of 31 subjects from 4 studies were selected with degeneration graded I-V, and their single cell RNA sequence data were extracted and integrated for further analysis. Samples defined as Pfirrmann grade I were considered as Non-degenerative; grades II/III and IV/V as respectively Mild and Severe degeneration. F: Female; M: Male.

Data availability	Sample ID	Gender	Age	Pfirrmann grade	Diagnosis	Labeling in analysis	Ref
GSE160756	GSM4878538	M	16	I	Spine fracture	Non-degenerative	14
	GSM4878539	M	31	I	Brain death	Non-degenerative	
	GSM4878540	M	31	I	Brain death	Non-degenerative	
CNP0002664	Ctrl	M	37	I	Spinal cord injury	Non-degenerative	7
	NP2	F	65	V	Herniation	Severe	
	NP4	M	59	III	Herniation	Mild	
	NP8	M	50	IV	Herniation	Severe	
	NP9	M	56	III	Herniation	Mild	
	NP10	F	48	III	Herniation/spondylolisthesis	Mild	
GSE165722	S1	M	63	II	Burst fracture	Mild	12
	S2	M	41	II	Burst fracture	Mild	
	S3	F	56	III	Herniation	Mild	
	S4	F	65	III	Herniation	Mild	
	S5	F	64	IV	Herniation	Severe	
	S6	F	53	IV	Herniation	Severe	
	S7	M	54	V	Herniation	Severe	
	S8	M	56	V	Herniation	Severe	
PRJCA014236	D1_1	M	7	I	Scoliosis	Non-degenerative	11

D1_2	F	6	I	Scoliosis	Non-degenerative
D2_1	M	19	II	Herniation/spondylolisthesis	Mild
D2_2	F	46	II	Herniation/spondylolisthesis	Mild
D2_3	M	15	II	Herniation/spondylolisthesis	Mild
D3_1	M	37	III	Herniation/spondylolisthesis	Mild
D3_2	F	31	III	Herniation/spondylolisthesis	Mild
D3_3	M	30	III	Herniation/spondylolisthesis	Mild
D4_1	F	44	IV	Herniation/spondylolisthesis	Severe
D4_2	F	43	IV	Herniation/spondylolisthesis	Severe
D4_3	M	54	IV	Herniation/spondylolisthesis	Severe
D5_1	M	68	V	Herniation/spondylolisthesis	Severe
D5_2	F	50	V	Herniation/spondylolisthesis	Severe
D5_3	F	57	V	Herniation/spondylolisthesis	Severe

Table 2. Analysis of DEGs for disc fibrocytes. *PTPRC⁺COL1A1⁺* or *ITGAM⁺COL1A1⁺* NPC were considered as disc fibrocytes, and subdivided to two major clusters based on their UMAP distribution: 1) immunocytes-clustered fibrocytes containing fib-M (Macrophage), fib-N (Neutrophil), fib-T (T cells) and fib-G (G-MDSC); 2) NP cells (NPC)-clustered fibrocytes containing fib-fNP (FibroNP), fib-cNP (ChonNP), fib-rNP (RegNP). Positively and negatively expressed DEGs of different fibrocytes subsets were demonstrated.

DEGs for immunocytes-clustered fibrocytes: fib-M; fib-N; fib-T; fib-G	
Commonly shared positive genes	*LAPTM5 PTPRC CXCR4 CFLAR HCLS1 PLEKHA2 FAM49B CELF2 SRGN ARHGDIB LCP1 EVI2B FMNL1 COL1A1 GMFG ITGB2 TMSB4X SAMSN1 MT-ND1 RAC2 B2M MT-ND3 SLA WDR26 FAM107B DEDD2 PFN1 MT-ND4 CFL1 FYB1 HLA-B LASP1 WIPF1 PPP1R18 GNB1 HLA-E HLA-A MSN MT-ATP8 MYH9 RAB8B ATP5E BAZ1A ARPC2 CAPZA1 MT-ND6 PABPC1*
Commonly shared negative genes	*GLT8D2 NCKAP1 NGEF FAP TMEM98 SOX5 DIXDC1 CD320 GSTM3 PDGFRL SERPINA1 SERTAD4-AS1 KDELR3 BARX1 TGM2 ZNF385D CFB FKBP7 ECM2 GPC6 CAV2 CHPF CDH19 AOPEP PERP LARP6 CNN3 CCDC85B TUBB2B MIR100HG MPP6 MAGI2-AS3 TIMP4 GFPT2 ARL3 MDFI SCIN SMOC1 VASN DPT LAMB2 CHST3 PPFIBP1 ACKR3 HSPA2 EVA1B GOLM1 METRN GNG11 WWTR1 SGCB ITGA10 MELTF PYURF THY1 FXYD1 LY6E EFEMP2 SNAI2 SSPN VGLL4 ADPRHL1 FHL2 DDR2 LTBP3 CAVIN3 RGS3 CILP2 ANGPTL2 PAX1 FSTL1 FGFRL1 P4HA2 FAM114A1 FAM3C MRC2 ANGPTL5 TMEM45A COX7A1 ANGPTL4 MXRA7 C1R H19 DANCR COL6A3 RHOBTB3 BAMBI TPM1 NFIA TRPS1 FKBP10 MXRA8 COL5A2 MEG3 CRIP2 TCEAL2 RCN3 COPZ2 IFITM3 CDO1 PDPN ID1 NRN1 TCF4 ID4 PDLIM4 SIVA1 ERRFI1 ADIRF ENPP1 PPIC EDIL3 FOXC1 TPM2 FGF2 NENF FERMT2 NOVA1 PCOLCE CHAD APOD ATP5IF1 RCN2 SLC39A14 EMP2 TMEM59 MAGED2 GPRC5A SBDS COL11A2 PRDX2 WDR83OS ISLR CALD1 AK1 CAVIN1 FHL1 COL6A1 BNIP3 TM4SF1 FXYD6 C1S*

	CLEC3A FBXO2 SOX9 TIMP3 RRBP1 CAV1 MYDGF SERPINA5 UAP1 PPDPF TNFRSF11B SPARC ITM2A FN1 CP EID1 LINC00632 RBP4 BEX3 MT1M KRT10 SERPING1 CPE VCAN VKORC1 PLOD2 PDIA6 CAPS COL6A2 SNHG5 KCNMA1 PMEPA1 MFGE8 COL9A3 SOD3 RABAC1 TNFRSF12A CCDC80 C12orf57 CCN1 DST HAPLN1 PRDX4 COL11A1 GPX3 PLAC9 COL2A1 PPIB FMOD HTRA1 CLU S100B NUPR1 TSC22D1 ZFAS1 ABI3BP SERPINA3 PCOLCE2 OGN S100A13 SERPINE2 PRELP IGFBP7 DSTN SSR4 AEBP1 C11orf96 MIF MIA CNMD SELENOM NNMT S100A1 BGN CCN2 SCRG1 MT1G RPS26 FGFBP2 SNORC ACAN ECRG4 LUM CRYAB DCN TIMP1 MGP NDUFA4L2 MT2A MT1X RPLP0 MT1E COMP RPL7
Fib-M specific	*CSTB LGMN ANXA2 DAB2 LGALS1 CTSL PLD3 CST3 NRP2 LMNA MGST3 PDE4DIP*
Fib-T specific	*RORA*
Fib-G specific	*NUCB2 DUT*
Fib-N specific	*N/A*
DEGs for NPC-clustered fibrocytes: fib-cNP; fib-fNP; fib-rNP	
Commonly shared positive genes	*COL1A1 COL3A1 COL1A2 ASPN TGFBI CTSB CTGF MT-ATP8 ATP5E ALDOA CDR1 HTRA1 ITGB5 MATR3 AC112491.1 ATP5G2 CYR61 ATP5J ATP5H HMOX1 NBEAL1 ATP5G3 ITGAM ATP5A1 ATP5F1 ATP5B ANKH LRRC75A ZNF90 TRAM1 ATP5L CERS2 DUSP5 PPP3CA PXDC1 CDK2AP1 OAT CKS2 ANGPTL4 SLC1A4 LDHA ZFAND5 MAN1A2*
Commonly shared negative genes	*RPL21 MALAT1*

704 **Table 3. Analysis of DEGs for NPC-clustered disc fibrocytes.** NP cells (NPC)-clustered fibrocytes

705 were subdivided based on their UMAP distribution: fib-cNP (ChonNP), fib-fNP (FibroNP) and fib-

706 rNP (RegNP). Featured genes of each subset that were negatively expressed by immunocytes-

707 clustered fibrocytes were summarized.

Featured DEGs of NPC-derived fibrocytes	
fib-cNP	*PLOD2 FGFBP2 SPARC FMOD HTRA1 TIMP3 ANGPTL2 DPT MELTF CCDC80 PCOLCE2 FHL2 OGN CILP2 GFPT2 SCIN PRELP SERPINA5 CPE HAPLN1 SCRG1 COL2A1 ANGPTL4 ABI3BP SOX9 COL5A2 FXYD6 P4HA2 SMOC1 FGF2 NRN1 MXRA7 CDO1 FSTL1 PPIC SERPINA1 ID4 TUBB2B TMEM45A LUM CDH19 FAM114A1*
fib-fNP	*COL6A3 COL5A2 COL6A1 DPT FSTL1 SPARC THY1 FAP COL6A2 ISLR CALD1 ANGPTL2 HTRA1 PCOLCE MXRA8 C1R CAV1 EDIL3 CAVIN1 CNN3 ECM2 TPM1 PPIC FKBP10 DDR2 PLOD2 GOLM1 SMOC1 SSPN FAM114A1 PDGFRL FN1 ERRFI1 TIMP3 FERMT2 PDPN GLT8D2 MXRA7 SLC39A14 BGN NCKAP1 CAVIN3 ANGPTL4 DST WWTR1 SGCB DIXDC1 P4HA2 C1S TCF4 RCN3 ACKR3 EFEMP2 IGFBP7*
fib-rNP	*DCN FAP NNMT FN1 MFGE8 GFPT2 CLU SERPINE2 SLC39A14 ANGPTL4 GPX3 ERRFI1 SOD3 C11orf96 CDO1 ACKR3 HTRA1*