Root-specific secondary metabolism at the single-cell level: a case study of theanine metabolism and regulation in the roots of tea plants (*Camellia sinensis*)

Shijia Lin[1,†], Yiwen Zhang[1,†], Shupei Zhang[1], Yijie Wei[1], Mengxue Han[1], Yamei Deng[1], Jiayi Guo[1], Biying Zhu[1], Tianyuan Yang[1], Enhua Xia[1], Xiaochun Wan[1], William J. Lucas[2], Zhaoliang Zhang[1,*]

[1]State Key Laboratory of Tea Plant Biology and Utilization, Anhui Agricultural University, Hefei, Anhui 230036, China

[2]Department of Plant Biology, College of Biological Sciences, University of California, Davis, CA 95616, USA

[†]These authors contributed equally to this work

[*]Corresponding author (Email: zhlzhang@ahau.edu.cn)

Abstract

Root-synthesized secondary metabolites are critical quality-conferring compounds of foods, plant-derived medicines, and beverages. However, information at a single-cell level on root-specific secondary metabolism remains largely unexplored. L-theanine, an important quality component of tea, is primarily synthesized in roots, from which it is then transported to new shoots of tea plant. In this study, we present a single-cell RNA sequencing (scRNA-seq)-derived map for the tea plant root, which enabled cell-type-specific analysis of glutamate and ethylamine (two precursors of theanine biosynthesis) metabolism, and theanine biosynthesis, storage, and transport. Our findings support a model in which the theanine biosynthesis pathway occurs via multicellular compartmentation and does not require high co-expression levels of transcription factors and their target genes within the same cell cluster. This study provides novel insights into theanine metabolism and regulation, at the single-cell level, and offers an example for studying root-specific secondary metabolism in other plant systems.

Introduction

Different root cell types produce diverse and complex secondary metabolites in response to environmental and developmental clues (Hong et al., 2017; Shaw et al., 2021); furthermore, such secondary metabolites can also participate in environmental adaptation and development (Garcia-Lemos et al., 2020; Hartman et al., 2017; Santelia et al., 2008; Schulz-Bohm et al., 2018; Steppuhn et al., 2004; Wan et al., 2018; Xu et al., 2017). As examples, the alkaloid nicotine is synthesized, specifically, in the root and is transported to the shoot for resistance to biotic stresses (Steppuhn et al., 2004; Xu et al., 2017). Root-synthesized flavonoids regulate root tip growth through affecting auxin transport and metabolism (Santelia et al., 2008; Wan et al., 2018). Legume roots secrete flavonoids as signaling agents to attract symbiotic bacteria, such as *Rhizobium* for nitrogen fixation (Hartman et al., 2017). In *Abies nordmanniana*, volatile organic compounds (e.g., propanal, g-nonalactone, and dimethyl disulfide) function to recruit certain bacteria or fungi, such as *Paenibacillus*. *Paenibacillus* sp. S37 produces high quantities of indole-3-acetic acid that can then promote plant root growth (Garcia-Lemos et al., 2020; Schulz-Bohm et al., 2018).

In many medicinal and beverage plants, root-synthesized secondary metabolites are critical quality-conferring compounds (Bailly, 2021; Hu et al., 2019; Schmid et al., 2018). Lobetyolin (LBT), a *Codonopsis pilosula* (known as Radix codonopsis or Dangshen) root-synthesized polyacetylene glycoside, exhibits activities against various cancers, notably, gastric cancer (Bailly, 2021). Polysaccharides and lectins isolated from *Pseudostellaria heterophylla* (known as Taizishen or Haiershen) roots, have multiple pharmacological activities (Hu et al., 2019). Root-synthesized polyphenols and triterpenoid saponins are crucial quality compounds extracted from *Glycyrrhiza glabra* (known as Licorice) and are employed in the food and beverage industry (Schmid et al., 2018). Due to the importance of secondary metabolites, the biosynthesis, transport, and regulation of plant root-synthesized secondary metabolites have been extensively studied (Fernie and Tohge, 2017; Wang et al., 2018b). However, in most cases, whole root materials were employed with very few assessing secondary metabolite biosynthesis and regulation, at the single-cell level.

Tea plant (*Camellia sinensis* L.), a crop with high economic value, is a member of the

eudicot family, *Theaceae*, has diverse and active secondary metabolic activities (Yu et al., 2020). Tea plants produce and accumulate extensive amounts of secondary metabolites, including theanine, catechins and caffeine (Zhao et al., 2020). These secondary metabolites confer tea with both unique flavor and multiple health benefits (Zhao et al., 2020). Theanine, a non-protein amino acid, endows the "umami" taste and relaxing effect of the tea infusion (Lin et al., 2022). It is the most abundant free amino acid in the tea plant, accounting for 1% - 2% dry weight of the tender shoots (Lin et al., 2022). It is primarily synthesized in tea plant roots and is then transported to the shoots, via the vasculature system (Figure 1A, B) (Lin et al., 2022).

Tea plant roots acquire inorganic nitrogen, especially ammonium, from the soil and assimilate it into amino acids, such as glutamate (Glu), glutamine (Gln), and alanine (Ala) (Lin et al., 2022). Ala can be decarboxylated, by alanine decarboxylase (CsAlaDC), to produce ethylamine (EA) (Bai et al., 2019; Zhu et al., 2021), which can then be combined with Glu, through the action of theanine synthase (CsTSI), to produce theanine (She et al., 2022; Wei et al., 2018). The high level of EA availability is likely a key factor in the elevated accumulation of theanine in tea plants (Cheng et al., 2017; Zhu et al., 2021). Importantly, in the tea plant, CsAlaDC and CsTSI are both indispensable for theanine synthesis (Zhu et al., 2021). Theanine synthesis is tightly regulated, at multiple levels, especially at the transcriptional level (Lin et al., 2022, Guo et al., 2022); CsMYB6 bound to the *CsTSI* promoter can regulate theanine synthesis (Zhang et al., 2021c). Furthermore, CsMYB40 and CsHHO3 bound to the *CsAlaDC* promoter can regulate theanine synthesis in "accelerator" or "brake" mode, respectively, in response to N levels (Guo et al., 2022). Theanine transporters, including the tonoplast-localized CsCAT2 and plasma membrane-localized CsAAPs, have been proposed to mediate in theanine storage within the root and root-to-shoot transport, respectively (Dong et al., 2020; Feng et al., 2021). In addition, in response to N status in tea plant roots, theanine may affect apoplastic H_2O_2 accumulation to regulate lateral root development (Chen et al., 2022). Taken together, these studies have partially revealed the mechanisms for theanine synthesis, transport, and regulation, but information on the specific root cell types involved in these processes is lacking.

Recently, with the in-depth study of biological structure and function, it has become

increasingly clear that differences exist in gene expression levels between cells, even if they

appear to be within the same cell population. Bulk RNA-seq can be employed to investigate

the average level of gene expression within whole tissues, whereas single-cell RNA sequencing

(scRNA-seq) can capture single cells, from whole tissues, to detect the heterogeneity of cells,

and obtain information on gene expression within single cells (Shaw et al., 2021). Thus,

scRNA-seq provides a powerful approach to elucidate the potential involvement of cell

heterogeneity in secondary metabolism in tea plant roots. Currently, scRNA-seq has been

performed in several plants, including *Arabidopsis thaliana* (Ryu et al., 2019), *Oryza sativa*

(Zhang et al., 2021b), and *Zea mays* (Nelms and Walbot, 2019). Importantly, algorithms are

available to decode the cellular regulatory codes from scRNA-seq datasets (Becht et al., 2018;

Haghverdi et al., 2015; Haghverdi et al., 2016; Trapnell et al., 2014), and these databases can

provide a critical foundation for analyzing the cellular-based information on gene expression

(Chen et al., 2021; Xu et al., 2022).

In plants, scRNA-seq has yet to be used for studying root-specific secondary metabolism.

In this study, using high-throughput scRNA-seq, we analyzed the cell heterogeneity of

theanine metabolism and regulation in the tea plant root. Our findings offer important insights

into the molecular mechanisms underlying theanine synthesis, transport, and regulation, and

provide a basis for studies on root-specific secondary metabolism in other plant systems.

Results

Tea plant root cell clusters identified by scRNA-seq

Tea plant roots synthesize and accumulate large amount of theanine (Figure 1A and 1B). The

tea plant root cell types, including cortical (Co), xylem (Xy), phloem (Ph), cambial (Ca),

pericycle (Pe), endodermal (En), epidermal (Ep) and root hair (RH), are shown in Figure 1C.

To understand the cellular characteristics of theanine biosynthesis in the tea plant root,

protoplasts were isolated from seedling roots (from the root tip back ~2.0 cm) of the tea plant

cultivar 'Shuchazao' (Figure 1A). These protoplasts were subjected to scRNA-seq, using the

commercial 10× Genomics platform (Figure 1D), and the sequencing data were pre-filtered, at

the cell and gene levels, resulting in a pool of 10,435 cells (Supplemental Figure 1A;

Supplemental Data 1). The *t*-distributed stochastic neighborhood embedding (*t*-SNE) algorithm

was employed to visualize local similarities (Haghverdi et al., 2015), resulting in these 10,435

cells being grouped into 8 cell clusters (Figure 2A; Supplemental Figure 1B).

The cell types within these 8 clusters were assessed by an analysis of homologs of cell

type marker genes according to protein sequence similarity, confirmed in other scRNA-seq

studies, or in the model plant *Arabidopsis thaliana*. In *A. thaliana,* GLABRA3 (GL3) is

involved in epidermal cell fate specification (Bernhardt et al., 2003). In our scRNA-seq,

CsGL3 transcripts were detected in Cluster 2, indicating the presence of epidermal cells

(Supplemental Figure 2A). The *SCARECROW* (*SCR*) gene is required for the asymmetric

division of the cortex/endodermis initial (CEI) cells, and its expression is confined to the

endodermis, CEI cell, and the quiescent center (QC) (Dong et al., 2021). In our tea plant root

data, *CsSCR* transcripts were detected primarily in Clusters 3 and 4 (Supplemental Figure 2A),

suggesting that cortical and endodermal cell division occurs in these two clusters. However, as

CsAAP1 transcripts were detected in Cluster 4 (Figure 2B), and the gene encodes for AMINO

ACID PERMEASE 1, which generally mediates in uptake of amino acids, into the cortex and

endodermis, this cluster likely contained a population of cortical or endodermal cells.

The stele system in the root is comprised of the pericycle and vasculature, in which

LONESOME HIGHWAY (*LHW*) has been established as a pericycle marker gene (De Rybel et

al., 2013). As the *CsLHW* transcripts were detected in Cluster 8, this cell cluster likely contains

a pericycle cell population (Figure 2B). The *WALLS ARE THIN LIKE 1* (*WAT1*) and

IRREGULAR XYLEM 9 (*IRX9*) are characterized vasculature and xylem development marker

genes, respectively (Li et al., 2021). Hence, the high level of *CsWAT1* transcripts detected in

Clusters 1 and Cluster 3 (Figure 2B), along with *CsIRX9* transcripts present in Cluster 3

(Supplemental Figure 2A) support the notion that these two clusters represent vasculature or

xylem-related cells in the tea plant root. Finally, the *PHLOEM PROTEIN 2* (*PP2*) gene

participates in phloem development and is a well-characterized phloem marker (Dinant et al.,

2003) and the presence in Cluster 7 of abundant *CsPP2* transcripts is consistent with phloem

cells being located within this Cluster 7 (Figure 2B).

To confirm the annotation of these cell clusters, we performed *in situ* RT-PCR analyses.

These assays indicated expression of the following genes in their respective cell types:

CsWAT1 in the vasculature (including pericycle, xylem parenchyma and cambium); *CsPP2* in

the protophloem; *CsAAP1* mainly in the cortex and endodermis; and *CsLHW* in the pericycle

(Figure 2C). These findings were consistent with the above annotation of these cell clusters

(Figure 2B and 2C).

Given the lack of more marker genes in tea plants, we next applied an unbiased approach,

developed by Denyer et al. (2019), in which cluster-enriched genes are used as markers to

define cell cluster identity. Cluster-enriched genes are defined as those being expressed, at

significantly higher level, in each cluster, than in all other clusters, and expressed in ≥ 10%

of cells within this specific cluster. The information on the cell-specific expression of these

cluster-enriched genes was confirmed in *A. thaliana*. Therefore, we analyzed expression

patterns of the top 10 cluster-enriched genes (Figure 2D; Supplemental Figures 2-6;

Supplemental data 3). Here, we could establish enrichment of *LACCASE 7* (*CsLAC7*) and

PECTIN METHYLESTERASE 41 (*CsPME41*) in Cluster 2, *ASPARAGINE RICH PROTEIN1*

(*CsNRP1*) in Clusters 3 and 5, and *MITOCHONDRIAL IMPORT INNER MEMBRANE*

TRANSLOCASE subunit TIM17 (*CsTIM17*) in Cluster 5 (Figure 2D). Based on these

scRNA-seq results from *A. thaliana*, and previous studies showing that *AtLAC7* and *AtPME41*

are expressed mainly in epidermal cells, *AtNRP1* functions as a root meristem maker, including

stem cell niche (SCN) and root cap (RC), and *AtTIM17* is expressed in the root cap (Passarinho

et al., 2001; Zhu et al., 2006; Supplemental Figure 2C), we assigned Cluster 3 and 5 as SCN and

RC cells, respectively.

Taken together, according to the above results, Clusters 1, 2, 4 and 7 were annotated as

xylem, epidermal, cortex/endodermal, and phloem cells, respectively (Figure 2E). In addition,

Cluster 8 was annotated as pericycle cells (Figure 2E), whereas Cluster 6 included primarily

procambium or transition cells during differentiation (Figure 2E), and Cluster 3 was defined as

the SCN, which serves as the cluster containing founder cells for differentiation (Figure 2E).

Differentiation trajectories of tea plant root cells supported cell cluster
annotations

During root cell differentiation, all root cell types originate from stem cells generated by the

root apical meristem (RAM). In the RAM, stem cells, including stelar initial (SI) cells, CEI

cells, epidermis/lateral root cap initial (ELI) cells and columella initial (CI) cells, surround the

QC (Efroni et al., 2016) (Figure 3A). The QC cells are required for specification of the SCN

and maintaining the undifferentiated state of the stem cell initials. Stem cells continuously

undergo asymmetric division to produce daughter cells, which then begin to differentiate into

outer, inner, and middle cell layers (Efroni et al., 2016) (Figure 3A). We noticed that Cluster 3

could probably be divided into 4 sub-cell states, C3.1, C3.2, C3.3 and C3.4 (Figure 3A), with

C3.1 expressing the QC marker gene, *WUS-RELATED HOMEOBOX 5* (*CsWOX5,* Kong et al.,

2015), C3.2 expressing the xylem development gene *CsIRX9*, C3.3 expressing the cortex and

endodermal cell fate decision gene, *CsSCR,* and C3.4 being close to the Cluster 2 epidermal cell

population (Figure 3A; Supplemental Figure 2A). Thus, these four Cluster 3 sub-cell states

were defined as QC, SI, CEI, and ELI cells, respectively (Figure 3A).

The differentiation trajectories of these clusters were next assessed to confirm our annotation

of the grouped cell clusters. To this end, we conducted pseudo-time analysis by ordering cells

of these clusters along a reconstructed trajectory (Figure 3B and Supplemental Figure 7). This

analysis indicated that Cluster 3 was primarily located at the start of the differentiation

trajectories, and that the trajectories of these cells bifurcated into two major and one minor

branch (Figure 3B). Here, the two major branches reflected differentiation trajectories of the

ELI and SI cells from the SCN (Cluster 3), with the ELI cells differentiating into the outer cell

layer (red branch) and Cluster 2 (epidermal cells) and Cluster 5 (root cap cells) located at the

end of this divergence (Figure 3B). The SI cells differentiated to give an inner cell layer (blue

branch): Cluster 7 (phloem cells) and Cluster 1 (xylem cells) were arranged successively

(Figure 3B). The minor branch resulted in CEI cell trajectories that differentiated to give a

middle cell layer (black branch); Cluster 4 (cortex and endodermal cells) were located at the

end of this divergence. Importantly, Cluster 4 (cortex and endodermis), Cluster 6 (cambium)

and Cluster 8 (pericycle) cells were also located within the region where cell differentiation was

initiated, thus suggesting these three cell clusters can differentiate into other cell types (Figure

3B). Indeed, the cambium differentiates into the xylem and the phloem, especially in woody

plants (Li et al., 2021); endodermal cells assume a SCI to generate a new epidermal layer

(Efroni et al., 2016); the founder cells, derived from the pericycle, can form lateral roots (Péret

et al., 2009). The results of our pseudo-time analyses matched well with these earlier

established findings.

We next constructed a differentiation heatmap of xylem and epidermal cells (Figure 3C).

Consistently, highly expressed genes in Cluster 1 (xylem cells) were most enriched in cell wall

biogenesis, lignin catabolic process and amino sugar catabolic process, etc. (Figure 3C and

Supplemental data 4). Meanwhile, genes highly expressed in Cluster 2 (epidermal cells) were

enriched in trichoblast differentiation, root hair elongation and plant epidermal cell

differentiation, etc. (Figure 3C and Supplemental data 4). These in-depth differentiation

trajectories of root cells further indicated reliability of the cell cluster annotations.

Finally, a differentiation map of tea plant root cells was constructed, based on the

pseudo-time analyses (Figure 3D). This map was generated to assist in developing an

understanding of tea plant root development, to provide a basis for studying secondary

metabolism, in tea plant roots, at the single-cell level.

Cell heterogeneity of N transport and metabolism in tea plant roots

In tea plant roots, theanine accounts for approx. 60~80% of the total free amino acids (Yang

et al., 2020). Nitrogen (N) uptake and transport, in the form of nitrate and ammonium, and N

assimilation into amino acids, are prerequisites for theanine biosynthesis (Yang et al., 2020)

(Figure 4A). We therefore investigated the cell heterogeneity of these processes by analyzing

the scRNA-seq data.

The NITRATE TRANSPORTERS (CsNRTs) and AMMONIUM TRANSPORTERS

(CsAMTs) function in N uptake in tea plants (Zhang et al., 2021a; Zhang et al., 2020) (Figure

4A). The heatmap of relative expression revealed that most *CsNRTs* and *CsAMTs*, including

CsAMT1.1, *CsAMT1.2*, *CsNRT1.1*, *CsNRT1.5*, *CsNRT1.7*, *CsNRT3.1* and *CsNRT3.2*, were

expressed primarily in Cluster 4 (cortex and endodermal cells); *CsAMT1.2* and *CsNRT1.5*

were also highly expressed in Cluster 7 (phloem cells); *CsAMT3.1* was expressed primarily in

Cluster 8 (pericycle cells) and Cluster 6 (cambium cells)*; CsAMT3.1* and *CsNRT3.2* were

relatively specifically expressed in Cluster 2 (epidermal cells); *CsNRT2.4* was highly and

specifically expressed in Cluster 3 (SCN) (Figure 4B). At the same time, our *in situ* RT-PCR

results for *CsAMT1.1, CsAMT3.1*, *CsNRT1.1* and *CsNRT3.2* confirmed these scRNA-seq

findings (Figure 4B and 4C). Furthermore, these results were similar to previous reports in *A.*

thaliana and rice (Wang et al., 2018a; Wang et al., 2018b).

Intracellular NO_3^- is assimilated in the cytoplasm into NO_2^-, by NITRATE REDUCTASE

(NR), which then enters the plastids where it is further assimilated into NH_4^+, by NITRITE

REDUCTASE (NiR). Subsequently, NH_4^+ enters the amino acid metabolism pathway (Galili et

al., 2016; Xu et al., 2012) (Figure 4A). Our scRNA-seq assays revealed that expression of

CsNR1, *CsNR2*, *CsNiR1* and *CsNiR2* was more active in Cluster 8 (pericycle cells) (Figure

4B); *CsNR1* and *CsNR2* expression was also relatively high in Cluster 1 (xylem cells). These

data supported the notion that N assimilation occurs primarily within stelar cells in tea plant

roots.

Previously, we grouped amino acid metabolism into five pathways (P1-P5), based on their

metabolic relationships (Yang et al., 2020): glutamate pathway (P1: Glu, Gln, arginine [Arg],

proline [Pro], citrulline [Cit]); asparate (Asp) pathway (P2: Asp, isoleucine [Ile], lysine [Lys],

threonine [Thr] and methionine [Met]); pyruvate pathway (P3: Ala, leucine [Leu] and valine

[Val]); aromatic amino acid pathway (P4: phenylalanine [Phe], tyrosine [Tyr] and tryptophan

[Trp]); and the 3-phosphoglycerate pathway (P5: serine [Ser], glycine [Gly] and cysteine

[Cys]) (Figure 4D and Supplemental data 5). Theanine is synthesized from Glu and

Ala-derived ethylamine (Lin et al., 2022).

Based on the scRNA-seq analysis, we observed that the expression level of genes encoding

for Glu biosynthetic pathway enzymes was the highest (especially in Clusters 1, 3, 4, 6 and 8)

within these five pathways (Figure 4D and Supplemental data 5), suggesting that biosynthetic

activity for Glu synthesis is highly active in the roots. Meanwhile, the expression level of

genes encoding alanine biosynthetic pathway enzymes was also high in all these clusters,

especially in Clusters 3 (SCN) and 8 (pericycle cells) (Figure 4D and Supplemental data 5).

These findings indicated that Glu and Ala metabolism represent the major amino acid

metabolism in tea plant roots. This is the basis for N metabolism into theanine in these roots.

Cell types participating in theanine biosynthesis, storage, and transport in tea
plant roots

The genes encoding enzymes in the theanine biosynthetic pathway, theanine transporters, and

transcription factors regulating theanine biosynthesis, were recently identified (Dong et al.,

2020; Feng et al., 2021; Guo et al., 2022; She et al., 2022; Wei et al., 2018; Zhang et al.,

2021c) (Figure 5A). We analyzed the cell-specific expression patterns of these genes using

our scRNA-seq data (Figure 5B and Supplemental data 6). Impressively, although *CsTSI* was

highly expressed in all clusters (Figure 5C), it was expressed, more specifically, in Cluster 8

(pericycle cells) (Figure 5B). Here, our *in situ* RT-PCR assays also revealed that *CsTSI* is

expressed primarily in the pericycle (Figure 5D). Consistently, Glu biosynthetic pathway

genes, including *CsGOGAT1*, *CsGDH2* and *CsGDH3*, also had a relatively high expression

level in Cluster 8 (Figure 5B), although some *CsGOGAT* and *CsGDH* members were also

relatively highly expressed in Clusters 1 (xylem cells), 2 (epidermis cells), 3 (SCN), 4 (cortex

and endodermal cells), 5 (root cap cells), 6 (cambium cells) and 7 (phloem cells) (Figure 5B).

The *CsTSI* and *CsAlaDC* are the most important genes for theanine synthesis (She et al.,

2022; Zhu et al., 2021). However, we observed that *CsTSI* and *CsAlaDC* were not enriched in

the same cluster. Different from *CsTSI*, *CsAlaDC* was most highly expressed in Cluster 1

(xylem cell) and was also relatively highly expressed in Cluster 6 (cambium cell) (Figure 5B

and 5C). The *in situ* RT-PCR assays verified the expression pattern of *CsAlaDC* in the

vasculature and cambium, and also showed that *CsAlaDC* was expressed, at a low level, in

the pericycle (Figure 5D). Therefore, generally, *CsTSI* and *CsAlaDC* are co-expressed in the

pericycle, where *CsTSI* is most highly expressed, whereas *CsAlaDC* is mostly highly

expressed in the vasculature, where *CsTSI* has a lower expression level. These results suggest

that the theanine biosynthesis pathway has a multicellular compartmentation characteristic, as

EA would be synthesized primarily in the vasculature, whereas theanine would be more

actively synthesized in the pericycle.

Generally, root-synthesized theanine is stored in the root cells, or transported from root to

shoot (Dong et al., 2020; Lin et al., 2022). In this regard, CsAAPs (CsAAP1, CsAAP2, CsAAP4,

CsAAP5, CsAAP6 and CsAAP8) and CsCAT2 were recently identified as theanine transporters

(Dong et al., 2020; Feng et al., 2021). Among them, CsCAT2 probably mediates theanine storage

in the vacuole, with CsAAP1 functioning in theanine loading into the xylem transpiration stream

for root-to-shoot transport (Dong et al., 2020; Feng et al., 2021). Interestingly, *CsCAT2* was shown

to be expressed most highly in the same cluster as *CsTSI*; i.e., in Cluster 8 (pericycle cells) (Figure

5B). *CsCAT2* was also relatively highly expressed in Cluster 4 (cortex and endodermal cells),

suggesting that CsCAT2 may function in theanine storage in the pericycle, cortex and endodermis.

In contrast, *CsAAP1* was expressed primarily in Cluster 4 (cortex and endodermal cells). This is

consistent with its proposed role in retrieving extracellular theanine for its transport through the

endodermis prior to xylem loading (Dong et al., 2020; Lin et al., 2022). The expression of other

CsAAPs was also shown to be highly cell cluster-specific; for example, *CsAAP2* in Cluster 4

(cortex and endodermal cells), *CsAAP4* in Cluster 6 (cambium cells), *CsAAP6* in Cluster 3 (SCN)

and Cluster 4 (cortex and endodermal cells), and *CsAAP8* in Cluster 5 (root cap cells) (Figure 5B),

suggesting cell type-specific functions for these CsAAPs in theanine transport.

Transcriptional regulation of theanine biosynthesis

Theanine biosynthesis is regulated at the transcriptional level, based on recent findings that

transcription factors (TFs) can control *CsTSI* or *CsAlaDC* expression to regulate its pathway

(Zhang et al., 2021; Guo et al., 2022). Here, Zhang et al. reported that CsMYB6, a root-specific

expressed TF, promoted theanine biosynthesis through activation of *CsTSI* expression. Thus, we

were surprised that *CsMYB6* was expressed in Cluster 3 (SCN), Cluster 6 (cambium cells) and

Cluster 1 (xylem cells), rather than in Cluster 8 (the high *CsTSI* expression cell cluster), as shown

in Figure 5B. A similar situation was also observed in terms of *CsAlaDC* transcriptional regulation.

We previously identified CsMYB40 and CsHHO3 to be an activator and repressor of *CsAlaDC*

expression, respectively, in response to N levels (Guo et al., 2022). Analysis of our scRNA-seq

data indicated that *CsMYB40* and *CsHHO3* were not highly expressed in Cluster 1 (the high

CsAlaDC expression cell cluster); *CsMYB40* was relatively highly expressed in Cluster 4 (cortex

and endodermal cells) and Cluster 7 (phloem cells); and *CsHHO3* was relatively highly expressed

in Clusters 3 (SCN) and 6 (cambium cells) (Figure 5B). Above regulatory relationships all have a

common feature, the TFs and their target genes were highly expressed in different cell types.

Next, to further explore whether this regulatory relationship also exists in other plants, we

analyzed expression of some TFs and their target genes in the model plant *Arabidopsis*, using

published single cell RNA-seq data (Ryu et al., 2019; Wendrich et al., 2020; Zhang et al.,

2019; Denyer et al., 2019; Jean-Baptiste et al. 2019; Shulse et al., 2019; Shahan et al., 2022)

and databases (Root Cell Atlas, https://rootcellatlas.org/; BAR,

https://bar.utoronto.ca/#GeneExpressionAndProteinTools). By integrating findings from

various studies (Fukaki et al., 2006; Okushima et al.,2007; Stracke et al., 2007; Bustos et al.,

2010; Huang et al., 2018; Liu et al., 2022; Tang et al., 2022), we observed that there is a

hierarchical regulatory network, in which IAA14 functions as the upstream regulator

(Supplemental figure 8A). Similar to the situation in tea plants, the regulators were not

present in exactly the same cell types in which their target genes were highly expressed. For example, *AtARF7* and *AtARF19* were highly expressed in the cortex and stele, respectively, whereas their target genes *AtLBD16* and *AtLBD29* were highly expressed in endodermal cells (Okushima et al.,2007; Supplemental figure 8B and 8C); *AtPHR1* was highly expressed in root epidermal and pericyte cells, but its target gene *AtF3'H* was highly expressed in the cortex and *AtRALF23* was highly expressed in xylem cells (Liu et al., 2022; Tang et al., 2022; Supplemental figure 8B and 8C). In this hierarchical regulatory network, some TFs have multiple target genes, and TFs participate in multiple biological processes (root development, metabolism regulation and plant immunity) through regulation of different target genes (Supplemental figure 8A).

Based on these findings, we hypothesized that, for the regulation of theanine biosynthesis, TFs and target genes are not necessarily always highly expressed in the same cells. Moreover, the regulation of theanine biosynthesis is probably coupled with the regulation of other biological processes. In the theanine biosynthesis pathway, CsAlaDC evolved a tea plant-specific activity in catalyzing EA biosynthesis (Cheng et al., 2017), which is essential for theanine biosynthesis (Zhu et al., 2021). *CsAlaDC* is highly and specifically express in tea plant roots (Zhu et al., 2021), and its expression levels were shown to be highly correlated with theanine content in the roots of various tea plant cultivars, and further, was under TF regulation (Zhu et al., 2021). Thus, taking the transcriptional regulation of *CsAlaDC* as an example, we next analyzed the TFs that were co-expressed with *CsAlaDC* to test this notion.

Identification of CsLBD37, a novel TF that co-regulates both theanine synthesis and lateral root development

In tea plant roots, both theanine biosynthesis and root development are regulated by N level and its form (Yang et al., 2020). Moreover, theanine is probably involved in N-regulated lateral root development (Chen et al., 2022). This implicates the presence of TFs that can co-regulate theanine biosynthesis and development in the tea plant root. To test this notion, we identified genes co-expressed with *CsAlaDC*, using our previous transcriptome data obtained for roots that were supplied with various N treatments, including without N (0N), 1.4 mM nitrate (NO_3^-), 1.4 mM EA (EA), 1.4 mM ammonium (NH_4^+), or 1.4 mM nitrate plus

ammonium (1:1) (CK) (Supplemental data 9 and 10). By filtering of these genes, using

|correlation coefficients| $\geq$ 0.85 and $p \leq$ 0.05, we constructed a *CsAlaDC* co-expression

network that consisted of 133 genes (Figure 6A and Supplemental data 10).

The expression patterns of these *CsAlaDC* co-expressing genes were assessed in eight

tissues, based on comprehensive tea plant transcriptome data (Wei et al., 2018; Xia et al.,

2019) (Figure 6B). Given that *CsAlaDC* expression is root-specific, we chose 34 root-specific

co-expressed genes (RSCGs) for further analysis (Figure 6B). Expression analyses of these

RSCGs showed that the genes which were positively correlated with *CsAlaDC* expression

were induced by N, especially by EA, and the negatively correlated genes were induced by a

0 N stress treatment (Figure 6C). Surprisingly, the results of the scRNA-seq analysis indicated

that most of these genes, which were highly correlated with *CsAlaDC*, were not highly

expressed in Cluster 1 (xylem cell) (Figure 6D).

Within these 34 root-specific genes, there were 7 genes potentially encoding for TFs

LBD37, LBD38, bZIP, WRKY, AP2 and MYB18 (Figure 6C). Among them, *CsLBD37*

(CSS0024718) had the most positive correlation with *CsAlaDC,* and was induced by N (CK,

EA and NH_4^+) and was not expressed highly in Cluster 1 (Figure 6C and 6D). Du et al. (2021)

predicted that *CsLBD38*, named *CsLBD37* in this study, is a highly credible TF that

regulates theanine synthesis and is in an evolution node of theanine-associated

regulatory module (TARM) in *Camellia* and Non-*Camellia* species. Therefore,

CsLBD37 probably regulates *CsAlaDC* expression in tea plants, but its cell type expression

pattern is different from that of *CsAlaDC*.

To further explore this situation, we first confirmed the root-specific expression for

CsLBD37 (Figure 7A and 7B). Next, we performed *in situ* RT-PCR assays, which indicated

that the *CsLBD37* expression pattern did not exactly reflect that observed for *CsAlaDC*.

CsLBD37 was expressed in the cambium; however, it was also expressed in the pericycle,

whereas *CsAlaDC* was barely expressed in these cells (Figure 5C; Figure 7C and 7D). As

expected, CsLBD37 was localized in the nucleus (Figure 7E), and furthermore, yeast

one-hybrid (Y1H) assays, combined with *CsAlaDC* promoter-*LUC* assays conducted in

tobacco leaves, were consistent with CsLBD37 binding directly to and repressing the activity

of the *CsAlaDC* promoter (Figure 7F and 7H). We further conducted an electrophoretic

mobility shift assay (EMSA) with recombinant CsLBD37 protein and showed that it bound to the proximal region (-1734 to -1695 bp) of the *CsAlaDC* promoter (Figure 7G and Supplemental Figure 9).

To test whether *CsLBD37* was able to inhibit *CsAlaDC* expression, *in vivo*, we developed transiently *CsLBD37*-silenced or over-expression tea seedlings, by antisense oligonucleotide (asODN) interference (Figure 7I) and generation of transgenic hairy roots, respectively (Figure 7K). *CsLBD37* expression was down-regulated and *CsAlaDC* was up-regulated in these asODN-treated roots (Figure 7I), whereas EA and theanine levels were increased (Figure 7J). In contrast, in *CsLBD37* transgenic hairy roots, *CsLBD37* was up-regulated and *CsAlaDC* was down-regulated (Figure 7K), whereas EA and theanine levels were decreased (Figure 7L). These findings offered support for the hypothesis that *CsLBD37* regulates *CsAlaDC* expression to modulate EA and theanine synthesis. *CsLBD37* probably acts as a "brake" to maintain the expression level of *CsAlaDC* within a certain range, in response to high levels of N. This may be why *CsLBD37* is a repressor but is positively correlated with *CsAlaDC*.

Some studies have reported that the LBD family of TFs can regulate both plant secondary metabolism and root development (Goh et al., 2012; Okushima et al., 2007; Ye et al., 2021). Lateral root founder cells originate from the pericycle, located opposite to the xylem poles (Péret et al., 2009). We noted that *CsLBD37* was also expressed in the tea plant root pericycle (Figure 7C). Therefore, we overexpressed *CsLBD37* in *A. thaliana* (Supplemental Figure 10A and 8B) and observed that the number of lateral roots, in these overexpression (OE) lines, was significantly less than in the wild-type (WT) plants (Supplemental Figure 10B and 10C). We further observed that *CsLBD37* overexpression repressed the initial steps of lateral root development (Supplemental Figure 10D-10G). These results implied CsLBD37 plays a role in regulates lateral root development in tea plant. Moreover, *CsLBD37* expression was induced by exogenous theanine treatment and this result underscores once again the importance of CsLBD37 in co-regulating theanine biosynthesis and root development (Supplemental Figure 10H).

Collectively, our findings support a model in which transcriptional regulation of theanine biosynthesis does not require high expression of both TFs and target genes in the same cell

cluster, but only requires partial co-expression. We also established that *CsLBD37* co-regulates both theanine biosynthesis and lateral root development in tea plants under N-rich conditions.

Discussion

Application of single-cell transcriptome sequencing to plant studies has afforded progress in understanding root development in model plants, such as *A. thaliana* and rice (Ryu et al., 2019; Zhang et al., 2021b). In this regard, important aspects of secondary and specialized metabolism occur in the plant root system and are important for plant growth and adaption to environmental challenges. These metabolic products are also essential for food and beverage quality, and pharmaceutical products. Bird et al. (2005) reported that biosynthesis of the narcotic analgesics, morphine and codeine, is localized to sieve elements in the opium poppy, which extends the function of sieve elements beyond transport, and into the realm of metabolism. However, significantly less information is currently available regarding both the cell heterogeneity of these secondary metabolic pathways and the underlying regulatory events that occur in the plant root system, especially in woody plants. The information regarding gene expression, in different type cells, is important with respect to establishing insights into both the mechanisms underlying these secondary metabolites and the regulatory events that control their syntheses. In this study, we performed a scRNA-seq-based study of tea plant roots, as an example, to elucidate the biosynthesis and regulatory processes for theanine, a root-specific secondary metabolite. In this process, we established the first map of tea plant roots comprised of 8 cell clusters (Figures 2A and 3D).

Cell cluster annotation of non-transgenic plants

Cell cluster annotation is a critical step for scRNA-seq. In this regard, we employed candidate marker genes to annotate cell clusters, as in previous studies (Ryu et al., 2019; Zhang et al., 2021b). In some non-model plants, including tea, transgenic technologies are not currently available and, hence, we used *in situ* RNA hybridization to establish the location(s) for gene expression. In some studies, isolation of different cell types was combined with q-RT-PCR to detect cell-type marker gene expression (Wang et al., 2022). However, this approach has two

limitations in that it cannot display the gene location directly and has only low resolution.

After numerous trials, we were able to optimize *in situ* RT-PCR assays (detailed in the

Methods), which enabled a cell-specific characterization of gene expression in tea plant root

cells, prior to establishing a genetic transformation system for tea. However, we note the

challenge associated with weak calling of homologous marker genes, which may reflect the

differences between tea and *A. thaliana*, along with differences in sampling sites, but this only

slightly impacted our findings, as in the annotation of cell clusters, *in situ* RT-PCR was

employed to identify expression patterns for weakly expressed tea plant root genes, such as

CsAAP1, CsLHW, CsAMT1.1, CsNRT1.1. Importantly, the results of *in situ* RT-PCR and

scRNA-seq analysis were generally consistent.

Nitrogen metabolism and transport of tea plant root at the single cell level

N is one of the most critical mineral nutrients essential for crop growth and yield performance

and provides precursors for the biosynthesis of secondary metabolites. Roots acquire NO_3^--N

and NH_4^+-N from the soil (Xu et al., 2012), with CsNRTs (CsNRT2.5 and CsNRT3.2) and

CsAMTs (CsAMT3.1) acting in N uptake and transport in tea plant roots (Zhang et al., 2021a;

Zhang et al., 2020). Our current findings provided further insight into the cellular locations

where these transporters function. We show that many *CsNRTs* and *CsAMTs* are in the cortex,

endodermal and stelar cells, whereas the N assimilation genes, *CsNRs* and *CsNiRs*, were

primarily located within the stelar cells (Figure 4B). Our data support a model in which these

CsNRTs and CsAMTs function in N uptake, into the cytoplasm of cortical and endodermal

cells, but also play an important role in NO_3^- and NH_4^+ loading into the cells within the stele

for N assimilation and amino acids synthesis.

The Glu pathway is the most abundant and active pathway for amino acid biosynthesis in

all root cell types. However, our findings indicated that stelar cells and the SCN have a higher

capacity for amino acid synthesis, compared with epidermal, cortical, and endodermal cells

(Figure 4C). For 20 amino acids, their rates of synthesis appeared to varying in different tea

plant root cells; a similar situation was recently reported in *A. thaliana* leaves (Kim et al.,

2021), but has not yet been reported in the root systems of other plants. These findings deepen

our understanding of N metabolism and transport in plant roots.

Numerous studies have established that the processes underlying plant N sensing, uptake,

transport, and utilization are critical for the next novel green revolution (Liu et al., 2022b). In

this regard, high-throughput scRNA-seq studies should offer greater insights into the cell-type

specificity of many N-associated genes, which could help to select cell-type-specific genes

more efficiently for future designed breeding programs.

Multicellular compartmentation of theanine metabolism and transport

Compared with N metabolism in other plant species, theanine metabolism is a special

component of N metabolism in tea plants (Xu et al., 2012; Zhao et al., 2020). Tea plants prefer

NH_4^+ and synthesize large amounts of non-toxic theanine to decrease toxicity associated with

high NH_4^+ levels (Lin et al., 2022). In the theanine biosynthesis pathway, and consistent with

the findings of She et al. (She et al., 2022), our study established that *CsTSI* is expressed

primarily in the pericycle cells (Cluster 8). We further clarified that *CsAlaDC* is generally

expressed in the vasculature (Clusters 1). This differential enrichment of *CsTSI* and *CsAlaDC*

expression suggests that EA and theanine are synthesized in different cell types (Figure 5),

presumably reflecting multicellular compartmentation (Figure 8A).

It is well known that theanine is the most abundant amino acid in the tea plant, and that

theanine synthesis does not always occur, but is regulated to maintain growth and

development in balance or adapted to environmental conditions. Indeed, the observed

multicellular compartmentation may contribute to the regulation of theanine biosynthesis

through precursor syntheses and transport within different cells. Thus, we propose a model in

which multicellular compartmentation of theanine and its precursors functions like an innate

design for regulation of secondary metabolism. This regulation model may be a way to avoid

excessive or inhibition of theanine synthesis. (Figure 8A). Compartmentalization of theanine

biosynthesis may effectively prevent interference between secondary metabolic pathways.

Similar mechanisms may also have evolved in other plants. We also cannot rule out that

synthesis of theanine or EA can occur in the same cell type; after all, *CsTSI* and *CsAlaDC* are

two of the most highly expressed genes in tea plant roots and their cell type-specific expression

partially overlapped (Yang et al., 2020) (Figure 4F). In addition, the spatial distribution of

theanine in the roots can be affected by transport processes. So, it is likely that the cell types

in which theanine is distributed do not exactly correspond to the cell types in which theanine

is synthesized.

On the other hand, multicellular compartmentation of theanine and its precursors EA implied

EA transport from its location of biosynthetic into theanine biosynthetic cells. EA is a

colorless gas, and hence it may be transported by way of simple diffusion, or via a gas channel.

Perhaps aquaporins (AQPs), a class of channel proteins that transport water, gas, and nutrients

(Hachez and Chaumont, 2010), may be involved. At present, the mechanism of EA transport

remains unknown, and the involvement of cell-type-specific AQPs would be worthy to explore

in the future.

Previously, we identified theanine transporters acting in theanine root-to-shoot transport

and theanine storage in the vacuole of tea plant root cells (Dong et al., 2020; Feng et al.,

2021). In the current study, we elucidated the cell heterogeneity of theanine transporters (Figure

5B and 8B), and among them *CsAAP1* is located primarily in cortical and endodermal cells

(Figures 2E and 5B), and *CsCAT2* is co-expressed with *CsTSI* in the same cell type. This nicely

reflects that CsAAP1 and CsCAT2 mediate in the physiological function of theanine transport

and storage (Dong et al., 2020; Feng et al., 2021). However, these theanine transporters are all

importers (Dong et al., 2020; Feng et al., 2021). The exporter(s) mediating in theanine

transport from the xylem parenchyma for loading into the xylem transpiration stream, within

the roots, along with other theanine export processes in the shoot, remain to be identified

(Figure 8B). Combining the cell type information, provided by scRNA-seq data, may assist in

the identification of these specific-cell-located theanine exporter(s), and may also provide

information for identifying amino acid exporters in other plant species. At present, all the

exporters identified are bidirectional transporters, such as UMAMITs (Besnard et al., 2016).

Therefore, specific-cell-located transporters, with bidirectional transport functions, could be

potential theanine exporters.

The regulation of theanine biosynthesis at the single cell level

Interestingly, genes highly correlated with *CsTSI* or *CsAlaDC* were identified, including some

confirmed TFs (*CsMYB6*, *CsMYB40*, *CsHHO3*) (Guo et al., 2022; Zhang et al., 2021c), which

were not always highly co-expressed in the same cell type. The cell heterogeneity of

regulatory genes for *CsTSI* or *CsAlaDC* expression may reflect the "accelerator" and "brake"

regulatory patterns that could act to fine-regulate theanine biosynthesis in specific cells (Guo

et al., 2022). Another possibility is that these theanine-associated genes are promiscuous,

having many target genes and regulate multiple biological processes in tea plants. In the model

plant *Arabidopsis*, AtARF7 and AtARF19 regulate *AtLBD16* and *AtLBD29*, thereby

participating in the regulation root development (Okushima et al.,2007; Supplemental figure

8B and 8C). Meanwhile, AtARF7 and AtARF19 acted in *AtPHR1*, AtPHR1 activated *AtF3'H*

and *AtRALF23* to co-regulate flavonoid metabolism and plant immunity (Liu et al., 2022;

Tang et al., 2022; Supplemental figure 8B and 8C). The highest expressed cell types of these

regulators and target genes were different. Thus, TFs that regulate the theanine biosynthesis

pathway may not always be co-express highly with their target genes, in the same cell type.

Indeed, this point was confirmed in a subsequent study on CsLBD37 regulation of *CsAlaDC*

expression.

Of course, TFs also will regulate their target genes and are highly co-expressed in the same

cell type. As an example, AtMYB12 activated *AtCHS* and they were both highly expressed in

cortical cells (Stracke et al., 2007; Supplemental figure 8B and 8C). However, members of the

MYB-bHLH-WD40 (MBW) complex encoded genes were expressed in different cell clusters

(Xu et al., 2014; Supplemental figure 8). Also, the processes of gene regulation may be

dynamically changing. Some TFs, such as HY5, SHR, and TMO, can alter their expression in

different cell types, due to their mobile characteristics or changes in cell differentiation and

external conditions. (Ortiz-Ramírez et al., 2021; Chen et al., 2024; Mankotia S et al., 2024).

In short, gene regulation is complicated at the single cell level. This complexity includes the

underlying nature of the biological processes involved, along with the complexity and

dynamics of their regulatory mechanisms. In this regard, ScRNA-seq better enabled us to

identify more details about gene regulation that were not previously noted.

To date, some identified TFs have been shown to play a role in the fine-regulation of

CsTSI and *CsAlaDC* expression; however, the TFs that impart this *CsTSI* and *CsAlaDC*

root-specificity and high expression remain to be identified. Recently, it was established that,

in roots, hypomethylation in the promoters of *CsTSI* and *CsAlaDC* provided an epigenetic

basis for their high expression (Kong et al., 2023). Although hypomethylation in the

promoters provides accessibility for transcription, activating TFs is still required for their high

expression. In this regard, the current cell-type-specific expression database offers a valuable

resource to identify such unknown TF activators. For example, root-specific and Cluster

1-specific CsMYB18 could well be an activator of *CsAlaDC* expression (Figure 7B-D).

Crosstalk between theanine metabolism and root development

Recently, some TFs regulating *CsTSI* or *CsAlaDC* were identified and characterized and

indicted that theanine regulates lateral root development (Chen et al., 2022; Guo et al., 2022;

Zhang et al., 2021c); however, the molecular link between these processes was not elucidated.

Previous studies have shown that several LBD family members regulate proanthocyanidin

(PA) metabolism and are involved in root development in response to N in *Arabidopsis*

thaliana (Rubin et al., 2009). In tea plant roots, theanine, PA metabolism and root

development are also regulated by N (Wang et al., 2021). In addition, *CsLBD37* was also

N-inducible and was one of the most positively correlated TF encoding genes with *CsAlaDC*

(Figure 6C). This information provided important insights into the roles for *CsLBD37* in

theanine biosynthesis and root development. In this study, we identified CsLBD37 as a

repressor of *CsAlaDC* transcription and root development (Figure 7 and 8C; Supplemental

Figure 10). CsLBD37 may act in a similar manner to CsHHO3, namely through being a

repressor of *CsAlaDC* transcription to "brake" theanine biosynthesis under high N availability

(Guo et al., 2022). Here, we also revealed that *CsLBD37* probably inhibits lateral root

development, at the root initial stage (Supplemental Figure 10D-10G), but the underlying

mechanism remains unknown. *CsLBD37* expression could be induced by exogenous theanine

treatment in tea seedlings (Supplemental Figure 10H). This implied that CsLBD37 may also

be involved in a process by which theanine regulates lateral root development. Besides, it has

been reported that LBD family TFs were regulated by, or interacted with, regulators of

hormone pathways (e.g., ARFs) to regulate the process of root morphogenesis (Goh et al.,

2012; Ye et al., 2021). Based on these findings, we speculated that CsLBD37 is likely

regulated by, or interacts with, other proteins to form a complex to regulate root development

or theanine biosynthesis. Then, is *CsLBD37* involved in regulating root development through

the hormones and ROS homeostasis pathways as well as theanine? This question is worthy of

future investigation.

In summary, we generated a transcriptional map of tea plant roots, at the single cell level,

which provided cell-type specific information on the expression of some 40,000 genes.

Moreover, these datasets provide an important resource for future studies on secondary

metabolism within tea plant roots, as well as other plant root systems. Recently, spatial

transcriptomics and metabolomics techniques were applied for studies in both animals and

plants (La Manno et al., 2021; Liu et al., 2022a). An integration of these techniques with

scRNA-seq would afford a powerful approach to further explore cell specific secondary

metabolism, at higher resolution, in various crop root systems.

Methods

Plant materials and growth conditions. Tea plants (*Camellia sinensis* var. *sinensis* cv.

Shuchazao) seeds were soaked in water for 7 days until their seed coats burst: water was

changed daily. Tea seeds were then cultured in vermiculite and watered (twice weekly) for 30

days, until radicle emergence. These tea seeds were then transplanted, singly, to new

vermiculite, and watered (three times weekly) until the two-leaf stage, when they were

transferred to nutrient solution for further growth; these hydroponic conditions and nutrient

solution were as described previously (Konishi et al., 1985).

Preparation of root samples for scRNA-seq. The root tip regions (~2 cm in length from

root tip) were cut, longitudinally, into 3 - 4 small strips and then processed for protoplast

isolation. Here, the root strips were transferred immediately and gently into the prepared

enzyme solution (0.1 mol·L^{-1} KCl, 0.08 mol·L^{-1} MES, 0.02 mol·L^{-1} CaCl$_2$, 1.5%

cellulase R10 (Yakult, Japan), 1% Pectolyase (Yakult, Japan), 0.4 mol·l-1 mannitol, and

0.1% BSA), and digested at 25°C, with shaking (40 rpm), in the dark for the indicated period.

An equal volume of washing buffer (WB) solution (0.1 mol·L^{-1} KCl, 0.08 mol·L^{-1} MES, 0.4

mol·L^{-1} mannitol and 0.1% BSA) was added in the enzyme solution and the mixture was

then filtered using nylon mesh (200 mesh/inch). The flow-through was centrifuged at 200 x g

for 2 min to pellet protoplasts. To evaluate protoplast yield, under different isolation conditions,

protoplasts were re-suspended with suitable WB solution and counted with a hemocytometer.

Protoplast viability was determined by trypan blue staining, with the ratio of viable cells being

more than 85%. The concentration of protoplasts was adjusted to 700–1200 cells/μL and cells

were then processed with the 10x Genomics Single Cell Protocol (CG00052, RevC).

Construction and sequencing of the scRNA-seq library. Approx. 12,593 counted cells

were loaded onto a Single Cell Gchip. The libraries were constructed using $10\times$ Genomics

Chromium Next GEM Single Cell 3□ Reagent Kits v3.1(1000268). In brief, cell

suspensions in a chip were loaded on a Chromium Controller (10xGenomics, Pleasanton, CA)

to generate single-cell GEMs (gel beads in emulsion). The scRNA-seq libraries were then

prepared, according to the manufacturer's protocol. Qualitative analysis of the DNA library

was performed on an Agilent 2100 Bioanalyzer. The concentration of DNA library was

measured by Qubit (Invitrogen). Libraries were sequenced by an Illumina NovaSeq sequencer

(Genergy Biotechnology Shanghai) and the raw scRNA-seq dataset comprised Read1, Read2,

and i7 index read. The 26-bp read length of Read1 contained the sequence of the 16 bp

10xBarcode and 10 bp UMI (unique molecular identifiers). The 98-bp read length of Read2

was the sequence of the cDNA fragment. In total, 332,593,050 reads were obtained.

Pre-processing of raw scRNA-seq data. The raw files were analyzed by Cell Ranger 5.0.0

(10x Genomics). The tea plant genome and GTF annotation files, which excluded the organelle

genomes, were downloaded from the Tea plants Information Archive (TPIA) website

(http://tpia.teaplants.cn) (Xia et al., 2019). Running 'cellranger mkref' with ''–genome, –fasta

and –genes'' inputs was employed to build the reference. Then 'cellranger count' with ''–id,

–transcriptome, –fastqs, –sample" was run. Some 79.8% of the reads in all of the samples were

aligned to the tea plant genome by the aligner STAR (v.2.5.1b). Mean reads per cell were

26,411. Median genes per cell was in the range of 680.

Data integration, clustering, and annotation. Downstream analyses were mainly

performed with the Seurat package (v.3.1.2), as previously described (Butler et al., 2018). To

remove low quality cells and likely multiplet captures, which is a major concern in

microdroplet-based experiments, we applied a criteria to filter out cells with UMI/gene numbers

out of the limit of the mean value +/- 2-fold of the standard deviations, assuming a Guassian

distribution of the UMI/gene number for each cell. Additionally, we applied DoubletFinder

package (version 2.0.2) potential doublet identification (McGinnis et al., 2019).

Library size normalization was performed with the "NormalizeData" function (LogNormalize

method, scaling factor of 10,000). We then detected variable genes with the

"FindVariableFeatures" function (vst method, 4000 features), scaled data with "ScaleData"

function, performed PCA analysis with "RunPCA" function (14 principal components),

determined statistical significance of PCA scores by "JackStraw" function, constructed the

SNN graph, clustered cells based on Louvain ("FindNeighbors" and"FindClusters"), and

visualized data with non-linear dimensional reduction algorithms ("RunTSNE"). The

cluster-enriched genes were computed with the FindAllMarkers function in Seurat, using the

following parameters: a bimod test; above 1 – fold difference (logfc. Threshold ≥0) between

the two groups of cells; test genes that having a minimum fraction was at least 0.25.

Pseudo-time analysis. Monocle2 (V2.9.0) package was used to infer the trajectory of cell

differentiation (Trapnell et al., 2014), and the specific steps were as follows: First, the import

CDS function of Monocle2 package was used to transform the Seurat object into the Cell Data

Set object. 0.01), next the reduce Dimension function was used for dimensionality reduction

clustering, and finally, order Cells function was used to infer the differentiation trajectory.

***In situ* RT-PCR.** *In situ* RT-PCR was improved and performed based on previous methods

(Munns et al., 2012). The experimental material was the primary root of a 2-month-old

'shuchazao' tea cultivar. The young root tissue was soaked in fresh FAA solution,

vacuum-infiltrated for 15-20 min, then put at 4□ for 12 h. The samples were then eluted, three

times, for 10 min each, in a mixture solution (63% ethanol and 5% acetic acid), and twice for 5

min each in 1× PBS. Root tissue was then embedded in 5% low-melting point agarose (agarose

dissolved in 1× PBS) and the embedded blocks were sectioned using a Leica RM2255

microtome (Leica, Nussloch, Germany). The root tissue sections were separated from agarose

by rinsing, twice, with Rnase-Free water, 3 μg/ml proteinase K was then added for 30 min at

25°C. A heat treatment, at 85°C for 2 min, was given to inactivate proteinase K, and samples

were then washed, once, separately with 1×PBS and then Rnase-Free water for 5-10 min,

followed by the addition of 1 U/μl Dnase I, at 37°C, for 20 min, or overnight. Liquid was then

removed and 15 mM EDTA (pH 8.0) was added, at 75°C for 10 min, followed by 1-2 washes

with Rnase-Free water to remove excess liquid. Reverse transcription reactions were performed

using the PrimeScript II 1[st] strand cDNA Synthesis Kit (TAKARA, Cat. No. 9750), solution

was then removed, followed by a wash, once, with Rnase-Free water. Mixed tissue sections and

reaction reagents: 2 μl 10× Tag DNA polymerase (NEB, Cat. No. M0267V), 0.4 μL Buffer Tag

DNA polymerase (5 U/μL, NEB, Cat. No. B9004S), 1.6 μL dNTP (TAKARA, Cat. No. 4030),

0.32 μL DIG-11-dUTP (25 nmol, Sigma, Cat. No. 11093088910), 1 μL Forward Primer, 1 μL

Reverse Primer, 0.6 μL MgCl$_2$ (1.5 mM, Thermo Fisher, Cat. No. R0971) and 13.08 μL ddH$_2$O.

PCR program: 95°C for 30 s, 95°C for 30 s, 55°C for 50 s, 68°C for 45 s, 30-35 cycle, 68°C for

5 min 10□. Washed twice with 1×PBS for 5 min each, and blocked with confining liquid (5%

skim milk) for 30 min. Alkaline phosphatase antibody was diluted in confining liquid at 1:500,

50 μL and was added and allowed to stand for 1 h. Wash twice for 15 min each time using 10×

Washing Buffer. Sample staining was performed using BM purple AP substrate, precipitating

for 30 min or 1 h, then washed twice with Rnase-Free water, and observed under a microscope

(Carl Zeiss, Gottingen, Germany) and photographed. All primers used for *in situ* PCR are listed

in Supplemental data 11.

Analysis of theanine and ethylamine content. Theanine and ethylamine were exacted

and detected as previously described, with slight modification (Yang et al., 2020; Cheng et al.,

2017). Theanine was analyzed by a Waters e2695 HPLC system equipped with 2489 UV/Vis

detector (Waters, U.S.A). Ethylamine was analyzed by GC-MS detecting (Agilent, Santa

Clara, U.S.A).

WGCNA analysis. The WGCNA package, in R software, was used in co-expression

analyses to detect relative relationships among genes (Langfelder and Horvath, 2008). A

weighted adjacency matrix was created, following unsupervised hierarchal clustering analysis

of genes, as follows: (1) removal of outliner genes and samples by the variation efficient and

hierarchical clustering analysis; (2) choosing of an applicable soft threshold (power = 12); (3)

identification of co-expressed gene modules using the function blockwise Modules; (4)

visualization of the co□expression network using Cytoscape3.9.0.

GO analysis. Tbtools (v1.098769) software was used to analyze the GO enrichment of

co-expression genes (Chen et al., 2020). GO annotation texts of tea plants were summarized

from TPIA database (http://tpia.teaplants.cn) (Xia et al., 2019).

Subcellular localization. Sequence information on *CsLBD37* was obtained from TPIA (http://tpia.teaplants.cn) (Xia et al., 2019). The ORF of *CsLBD37* within the entry vector pDONR207 was cloned into the destination binary vector, pK7WGF2.0, for subcellular localization studies. The plasmid, pK7WGF2.0-*CsLBD37*-GFP was transformed into *Agrobacterium tumefaciens* strain EHA105 to select for positive colonies for infiltration into *Nicotiana benthamiana*. After 48h-72h post-infiltration, the GFP fluorescence was imaged with an Olympus FV1000 confocal microscope (Olympus, Tokyo, Japan) to reveal the subcellular localization of *CsLBD37*. GFP fluorescence signals were excited with a 488-nm laser, and the emitted light was recorded from 500 to 530 nm to display the subcellular localization of *CsLBD37*.

Yeast one-hybrid assays (Y1H). We inserted *CsAlaDC* promoter fragments (26 approx. 2000 bp) into the pAbAi vector to generate the Pro*CsAlaDC*-pAbAi reporter construct, which was then introduced into the yeast one-hybrid gold strain. The transformed cells were grown on synthetic dropout medium (SD–Ura) (Coolaber, Beijing, China). The ORF of *CsLBD37* was inserted into vector pGADT7 AD to generate AD-CsLBD37, which was then introduced into the yeast strain with pro*CsAlaDC*-pAbAi. Transformants were grown on synthetic dropout medium (SD-Leu) with 200 ng ml^{-1} Aureobasidin A (AbA) (Coolaber, Beijing, China) for 3 days at 30°C. pGADT7 AD empty vectors were used as controls.

Luciferase activity assay. Approx. 2 kb *CsAlaDC* promoter fragments were inserted into the pGreenII 0800-LUC vector to generate the luciferase reporter construct, pro*CsAlaDC*-LUC. The *CsLBD37* ORF was inserted into the pGreenII 62-SK vector to generate the effector construct *35S-CsLBD37*. Empty vectors were used as negative controls. The above-described constructs were introduced into *A. tumefaciens* strain GV3101 with the pSoup plasmid, respectively. Then effector and reporter cell suspensions were mixed, in a 5:1 (v/v) ratio, and infiltrated into 4-week-old *N. benthamiana* leaves, followed by harvesting after 3 days. Sections of leaves were sprayed with 150 mg L^{-1} D-luciferin, potassium salt (Sciencelight, Shanghai, China) and LUC signals were captured using a Tanon 5200 Multi charge-coupled device camera (Tanon, Shanghai, China). LUC and REN activities of other *N benthamiana* leaves were measured using a dual-luciferase assay kit (Yeasen Biotech, Shanghai, China) and a Spectra Max M2 (Molecular Devices, America) according to the manufacturer's instructions.

Electrophoretic mobility shift assay (EMSA). The *CsLBD37* ORF was inserted into the

GST vector pGEX-4T-1 to generate GST-CsLBD37 construct, which was then transferred into

Escherichia coli BL21 cells. The positive BL21 strains were induced by 0.5 mM

isopropyl-beta-D-thiogalactopyranoside (IPTG), at 16°C for 20 h, and the GST-CsLBD37

recombinant proteins were purified by immobilized glutathione beads (TransGen Biotech, Beijing,

China)(Supplemental Figure 8). DNA probes were synthesized and labelled with biotin, at the 3'

end, by using the EMSA Probe Biotin Labeling Kit (Beyotime Biotech, Shanghai, China). The

EMSA was performed using the instructions of the Chemiluminescent EMSA Kit (Beyotime

Biotech, Shanghai, China).

Antisense oligonucleotides assay. The antisense oligonucleotides assay was performed,

as previously described (Xie et al., 2014), to silence the target gene *CsLBD37*. AsOND was

designed and selected using the SOLIGO software

(https://sfold.wadsworth.org/cgi-bin/soligo.pl). The sequence of the sODN and AsODN

are listed in Supplemental data 11. To silence the genes, the primary roots of tea plant

cultivar 'Shuchazao' were wounded and then incubated in 10 ml 50 μM asODN-*CsLBD37*

solution for 6 h and 12 h. The solution with sense oligonucleotides (sOND) of the gene

serves as the control. After treatment, the roots were collected for RNA isolation and

metabolite detection.

Overexpression of *CsLBD37* in a tea plant hairy root system. The *CsLBD37* ORF

was inserted into pK7WGF2.0 driven by the *35S* promoter, using the BP and LR reaction

(Gateway system) to generate the pK7WGF2.0-CsLBD37 overexpression construct. The

resultant vector was transformed into Ar.A4 Chemically Competent Cell (Weidi Biotech,

Shanghai, China). The selected positive transformants were collected and resuspended in

1/2 Murashige and Skoog (MS) culture medium to an $OD_{600} = 0.6$ to transform 3-month-old

tea seedlings, as described previously (Alagarsamy et al., 2018).

Arabidopsis **transformation.** The Gateway system was employed to generate the

pB2GW7-CsLBD37 construct. *A. tumefaciens* strain GV3101, carrying pB2GW7-*CsLBD37*,

was used for the genetic complementation of *Arabidopsis* wild-type (Col-0) plants.

Transformants were selected on 0.25 ‰ Basta, and further verified by genomic PCR and

quantitative RT-PCR (qRT-PCR). The T3 progeny plants were used for these analyses. All

primers are listed in Supplemental data 11.

Theanine treatment of tea seedlings. Tea seedlings with a 5-cm primary root were

cultured in 1/2 nutrient solution for 1 week, and then treated with 1/2 nutrient solution

containing theanine for 2 weeks. The theanine concentration of the control group (CK)

was 0 mM and that for the treatment group (Theanine) was 1 mM.

RNA isolation and q-RT–PCR. RNA isolation was performed according to the manual of

RNAprep Pure Plant Plus Kit (TIANGEN, Beijing, China), and cDNA synthesis was

accomplished using the TransScript II One-Step gDNA Removal and cDNA Synthesis

SuperMix kit (TransGen Biotech, Beijing, China). A QuantStudio 6 Flex Real-Time PCR

System (Thermo, Singapore) was used to measure relative expression of genes, which were

calculated based on the $2^{-\Delta\Delta Ct}$ method (Schmittgen et al., 2000), using *CsGAPDH* as an internal

control. All primers used for qRT-PCR are listed in Supplemental data 11.

Data available. The raw data and processed files have been submitted to the Gene Expression

Omnibus (https://www.ncbi.nlm.nih.gov/geo/) under accession number GSE267845.

Reference

Alagarsamy, K., Shamala, L. F. & Wei, S. (2018). Protocol: high-efficiency in-planta
   agrobacterium-mediated transgenic hairy root induction of *Camellia sinensis* var. *sinensis*.
   Plant Methods. **14**:17.

Bai, P. et al. (2019). Identification of a novel gene encoding the specialized alanine
   decarboxylase in tea (*Camellia sinensis*) plants. *Molecules* **24**:540.

Bailly, C. (2021). Anticancer properties of lobetyolin, an essential component of *Radix
   Codonopsis* (Dangshen). *Nat. Prod. Bioprospect.* **11**:143-153.

Becht, E. et al. (2018). Dimensionality reduction for visualizing single-cell data using UMAP.
   Nat. Biotechnol.

Bernhardt, C. et al. (2003). The bHLH genes *GLABRA3* (*GL3*) and *ENHANCER OF
   GLABRA3* (*EGL3*) specify epidermal cell fate in the Arabidopsis root. *Development*. **130**:

6431-6439.

Besnard, J. et al. (2016). UMAMIT14 is an amino acid exporter involved in phloem
unloading in Arabidopsis roots. *J. Exp. Bot.* **67**:6385-6397.

Bird, D. et al. (2003). A tale of three cell types: alkaloid biosynthesis is localized to sieve
elements in opium poppy. *Plant Cell*. **11**:2626-35.

Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. (2018). Integrating
single-cell transcriptomic data across different conditions, technologies, and species. *Nat.*
Biotechnol. **36**:411-420.

Chen, C. et al. (2020). TBtools: an integrative toolkit developed for interactive analyses of
big biological data. Mol. Plant **13**:1194-1202.

Chen, H. et al. (2021). PlantscRNAdb: a database for plant single-cell RNA analysis. Mol.
Plant **14**:855-857.

Chen, T. et al. (2022). Theanine, a tea plant specific non-proteinogenic amino acid, is
involved in the regulation of lateral root development in response to nitrogen status. Hortic.
Res. **10**:uhac267.

Chen, M. et al. (2024). TARGET OF MONOPTEROS: key transcription factors
orchestrating plant development and environmental response. J Exp Bot. **75**:2214-2234.

Cheng, S. et al. (2017). Studies on the biochemical formation pathway of the amino acid
l-theanine in tea (*Camellia sinensis*) and other plants. J. Agric. Food Chem. **65**:7210-7216.

De Rybel, B. et al. (2013). A bHLH complex controls embryonic vascular tissue
establishment and indeterminate growth in *Arabidopsis*. Dev. Cell **24**:426-437.

Denyer, T. et al. (2019). Spatiotemporal developmental trajectories in the *Arabidopsis* root
revealed using high-throughput single-cell RNA sequencing. Dev. Cell **48**:840-852.e845.

Dinant, S. et al. (2003). Diversity of the superfamily of phloem lectins (phloem protein 2) in
angiosperms. Plant Physiol. **131**: 114-128.

Dong, C. et al. (2020). Theanine transporters identified in tea plants (*Camellia sinensis* L.).
Plant J. **101**:57-70.

Dong, W. et al. (2021). An SHR–SCR module specifies legume cortical cell fate to enable
nodulation. Nature **589**:586-590.

Du, J. et al. (2021). Gene coexpression network reveals insights into the origin and evolution

of a theanine-associated regulatory module in Non-*Camellia* and *Camellia* Species. J. Agric.

Food Chem. **69**:615-626.

Efroni, I. et al. (2016). Root regeneration triggers an embryo-like sequence guided by

hormonal interactions. Cell **165**:1721-1733.

Feng, L. et al. (2021). Tonoplast-localized theanine transporter CsCAT2 may mediate

theanine storage in the root of tea plants (*Camellia sinensis* L.). Front. Plant Sci.

12:797854.

Fernie, A. R. & Tohge, T. (2017). The genetics of plant metabolism. Annu. Rev. Genet.

51:287-310.

Galili, G., Amir, R. & Fernie, A. R. (2016). The regulation of essential amino acid synthesis

and accumulation in plants. Annu. Rev. Plant Biol. **67**:153-178.

Garcia-Lemos, A. M. et al. (2020). Identification of root-associated bacteria that influence

plant physiology, increase seed germination, or promote growth of the christmas tree

species *Abies nordmanniana*. Front. Microbiol. **11**:566613.

Goh, T., Joi, S., Mimura, T. & Fukaki, H. (2012). The establishment of asymmetry in

Arabidopsis lateral root founder cells is regulated by LBD16/ASL18 and related LBD/ASL

proteins. Development **139**:883-893.

Guo, J. et al. (2022). Potential "accelerator" and "brake" regulation of theanine biosynthesis

in tea plant (*Camellia sinensis*). Hortic. Res. **9**:uhac169.

Hachez, C. & Chaumont, F. (2010). Aquaporins: a family of highly regulated

multifunctional channels. Adv. Exp. Med. Biol. **679**:1-17.

Haghverdi, L., Buettner, F. & Theis, F. J. (2015). Diffusion maps for high-dimensional

single-cell analysis of differentiation data. Bioinformatics **31**:2989-2998.

Haghverdi, L., Büttner, M., Wolf, F. A., Buettner, F. & Theis, F. J. (2016). Diffusion

pseudotime robustly reconstructs lineage branching. Nat. Methods **13**: 845-848.

Hartman, K., van der Heijden, M. G., Roussely-Provent, V., Walser, J. C. & Schlaeppi,

K. (2017). Deciphering composition and function of the root microbiome of a legume plant.

Microbiome **5**:2.

Hong, J. H. et al. (2017). A sacrifice-for-survival mechanism protects root stem cell niche

from chilling stress. Cell **170**:102-113.e114.

Hu, D. J., Shakerian, F., Zhao, J. & Li, S. P. (2019). Chemistry, pharmacology and analysis

of *Pseudostellaria heterophylla*: a mini-review. Chin. Med. **14**: 21.

Kim, J. Y. et al. (2021). Distinct identities of leaf phloem cells revealed by single cell

transcriptomics. Plant Cell **33**:511-530.

Kong W. L. et al. (2023). 5mC DNA methylation modification-mediated regulation in tissue

functional differentiation and important flavor substance synthesis of tea plant (*Camellia*

sinensis L.). Hortic Res. **10**:uhad126.

Kong, X. et al. (2015). WOX5 is shining in the root stem cell niche. Trends Plant Sci.

20:601-603.

Konishi, S. et al. (1985). Stimulatory effects of aluminum on tea plants grown under low and

high phosphorus supply. *Soil Sci. Plant Nutr.* **31**:361–368

La Manno, G. et al. (2021). Molecular architecture of the developing mouse brain. Nature

596:92-96.

Langfelder, P. & Horvath, S. (2008). WGCNA: an R package for weighted correlation

network analysis. BMC. Bioinformatics **9**:559.

Li, H. et al. (2021). Single-cell RNA sequencing reveals a high-resolution cell atlas of xylem

in populus. J. Integr. Plant Biol. **63**:1906-1921.

Lin, S. et al. (2022). Theanine metabolism and transport in tea plants (*Camellia sinensis* L.):

advances and perspectives. Crit. Rev. Biotechnol. 1-15.

Liu, C. et al. (2022). A spatiotemporal atlas of organogenesis in the development of orchid

flowers. Nucleic Acids Res. **50**:9724-9737.

Liu, Q. et al. (2022). Improving crop nitrogen use efficiency toward sustainable green

revolution. Annu. Rev. Plant Biol. **73**:523-551.

Liu, W., Xu, L., Lin, H. & Cao, J. (2021). Two expansin genes, *AtEXPA4* and *AtEXPB5*, are

redundantly required for pollen tube growth and *AtEXPA4* is involved in primary root

elongation in *Arabidopsis thaliana*. Genes (Basel). **12**:249.

Liu, Z. et al. (2022). PHR1 positively regulates phosphate starvation-induced anthocyanin

accumulation through direct upregulation of genes *F3'H* and *LDOX* in *Arabidopsis*. Planta.

256:42.

Mankotia, S. et al. (2024). HY5: a key regulator for light-mediated nutrient uptake and

utilization by plants. New Phytol. **241**:1929-1935.

McGinnis, C. S., Murrow, L. M. & Gartner, Z. J. (2019). DoubletFinder: doublet detection

in single-cell RNA sequencing data using artificial nearest neighbors. Cell Syst. **8**:329-337.

Munns, R. et al. (2012). Wheat grain yield on saline soils is improved by an ancestral Na□

transporter gene. Nat. Biotechnol. **30**:360-364.

Nelms, B. & Walbot, V. (2019). Defining the developmental program leading to meiosis in

maize. Science **364**:52-56.

Okushima, Y., Fukaki, H., Onoda, M., Theologis, A. & Tasaka, M. (2007). ARF7 and

ARF19 regulate lateral root formation via direct activation of LBD/ASL genes in

Arabidopsis. Plant Cell **19**:118-130.

Ortiz-Ramírez C. et al., (2021). Ground tissue circuitry regulates organ complexity in maize

and *Setaria*. Science. **374**:1247-1252.

Passarinho, P. A., Van Hengel, A. J., Fransz, P. F. & de Vries, S. C. (2001). Expression

pattern of the *Arabidopsis thaliana AtEP3/AtchitIV* endochitinase gene. Planta

212:556-567.

Péret, B. et al. (2009). *Arabidopsis* lateral root development: an emerging story. Trends Plant

Sci. **14**:399-408.

Rubin, G., Tohge, T., Matsuda, F., Saito, K. & Scheible, W. R. (2009). Members of the

LBD family of transcription factors repress anthocyanin synthesis and affect additional

nitrogen responses in *Arabidopsis*. Plant Cell **21**:3567-3584.

Ryu, K. H., Huang, L., Kang, H. M. & Schiefelbein, J. (2019). Single-cell RNA sequencing

resolves molecular relationships among individual plant cells. Plant Physiol.

179:1444-1456.

Santelia, D. et al. (2008). Flavonoids redirect PIN-mediated polar auxin fluxes during root

gravitropic responses. J. Biol. Chem. **283**:31218-31226.

Schmid, C., Dawid, C., Peters, V. & Hofmann, T. (2018). Saponins from european licorice

roots (*Glycyrrhiza glabra*). J. Nat. Prod. **81**:1734-1744.

Schmittgen, T. D. et al. (2000). Quantitative reverse transcription-polymerase chain reaction

to study mRNA decay: comparison of endpoint and real-time methods. Anal. Biochem.

285:194-204.

Schulz-Bohm, K. et al. (2018). Calling from distance: attraction of soil bacteria by plant root
volatiles. Isme J. **12**:1252-1262.

Shahan, R. et al. (2022). A single-cell *Arabidopsis* root atlas reveals developmental
trajectories in wild-type and cell identity mutants. Dev Cell. **57**:543-560.e9.

Shaw, R., Tian, X. & Xu, J. (2021). Single-cell transcriptome analysis in plants: advances
and challenges. Mol. Plant **14**:115-126.

She, G. et al. (2022). Characterization of *CsTSI* in the biosynthesis of theanine in tea plants
(*Camellia sinensis*). J. Agric. Food Chem. **70**:826-836.

Shulse, C. et al. (2019). High-throughput single-cell transcriptome profiling of plant cell
types. Cell Rep. **27**:2241-2247.e4.

Silva, L. S., Seabra, A. R., Leitão, J. N. & Carvalho, H. G. (2015). Possible role of
glutamine synthetase of the prokaryotic type (GSI-like) in nitrogen signaling in *Medicago*
truncatula. Plant Sci. **240**:98-108.

Sláviková, S. et al. (2005). The autophagy-associated Atg8 gene family operates both under
favourable growth conditions and under starvation stresses in *Arabidopsis* plants. J. Exp.
Bot. **56**:2839-2849.

Steppuhn, A., Gase, K., Krock, B., Halitschke, R. & Baldwin, I. T. (2004). Nicotine's
defensive function in nature. PLoS Biol. **2**:E217.

Tang, J. et al. (2022). Plant immunity suppression via PHR1-RALF-FERONIA shapes the
root microbiome to alleviate phosphate starvation. EMBO J. **41**:e109102.

Trapnell, C. et al. (2014). The dynamics and regulators of cell fate decisions are revealed by
pseudotemporal ordering of single cells. Nat. Biotechnol. **32**:381-386.

Wan, J. et al. (2018). UV-B radiation induces root bending through the flavonoid-mediated
auxin pathway in *Arabidopsis*. Front. Plant Sci. **9**:618.

Wang, Q. et al. (2022). Single-cell transcriptome atlas reveals developmental trajectories and
a novel metabolic pathway of catechin esters in tea leaves. Plant Biotechnol. J.
20:2089-2106.

Wang, W. et al. (2018). Expression of the nitrate transporter gene *OsNRT1.1A/OsNPF6.3*
confers high yield and early maturation in rice. Plant Cell **30**:638-651.

Wang, Y. et al. (2021). Nitrogen-regulated theanine and flavonoid biosynthesis in tea plant

940 roots: protein-level regulation revealed by multiomics analyses. J. Agric. Food Chem.
941 **69**:10002-10016.

942 **Wang, Y. Y., Cheng, Y. H., Chen, K. E. & Tsay, Y. F.** (2018). Nitrate transport, signaling,
943 and use efficiency. Annu. Rev. Plant. Biol. **69**:85-122.

944 **Wei, C. et al.** (2018). Draft genome sequence of *Camellia sinensis var. sinensis* provides
945 insights into the evolution of the tea genome and tea quality. Proc. Natl. Acad. Sci. U. S. A.
946 **115**:E4151-e4158.

947 **Wendrich, J.R., et al.** (2020). Vascular transcription factors guide plant epidermal responses
948 to limiting phosphate conditions. Science. **370**:eaay4970.

949 **Xia, E. H. et al.** (2019). Tea plant information archive: a comprehensive genomics and
950 bioinformatics platform for tea plant. Plant Biotechnol. J. **17**:1938-1953.

951 **Xie, Z. et al.** (2014). A selection strategy in plant transformation based on antisense
952 oligodeoxynucleotide inhibition. Plant J. **77**:954-961.

953 **Xu, G., Fan, X. & Miller, A. J.** (2012). Plant nitrogen assimilation and use efficiency. Annu.
954 Rev. Plant Biol. **63**:153-182.

955 **Xu, S. et al.** (2017). Wild tobacco genomes reveal the evolution of nicotine biosynthesis. Proc.
956 Natl. Acad. Sci. U. S. A. **114**:6133-6138.

957 **Xu, Z. et al.** (2022). Plant single cell transcriptome hub (PsctH): an integrated online tool to
958 explore the plant single-cell transcriptome landscape. Plant Biotechnol. J. **20**:10-12.

959 **Zhang, T. et al.** (2019). A single-cell RNA sequencing profiles the developmental landscape
960 of arabidopsis root. Mol Plant. **12**:648-660.

961 **Yang, T. et al.** (2020). Transcriptional regulation of amino acid metabolism in response to
962 nitrogen deficiency and nitrogen forms in tea plant root (*Camellia sinensis* L.). Sci. Rep.
963 **10**:6868.

964 **Ye, L. et al.** (2021). Cytokinins initiate secondary growth in the *Arabidopsis* root through a
965 set of LBD genes. Curr. Biol. **31**:3365-3373.e3367.

966 **Yu, X. et al.** (2020). Metabolite signatures of diverse *Camellia sinensis* tea populations. Nat.
967 Commun. **11**:5586.

968 **Zhang, F. et al.** (2020). Identification of regulatory networks and hub genes controlling
969 nitrogen uptake in tea plants [*Camellia sinensis* (L.) O. Kuntze]. J. Agric. Food Chem.

68:2445-2456.

Zhang, F. et al. (2021). Transcriptome analysis identifies CsNRT genes involved in nitrogen

uptake in tea plants, with a major role of CsNRT2.4. Plant Physiol. Biochem. **167**:970-979.

Zhang, T. et al. (2021). Single-cell transcriptome atlas and chromatin accessibility landscape

reveal differentiation trajectories in the rice root. Nat. Commun. **12**:2053.

Zhang, Y. et al. (2021). Molecular basis of the distinct metabolic features in shoot tips and

roots of tea plants (*Camellia sinensis*): characterization of MYB regulator for root theanine

synthesis. J. Agric. Food Chem. **69**:3415-3429.

Zhao, J., Li, P., Xia, T. & Wan, X. (2020). Exploring plant metabolic genomics: chemical

diversity, metabolic complexity in the biosynthesis and transport of specialized metabolites

with the tea plant as a model. Crit. Rev. Biotechnol. **40**:667-688.

Zhu, B. et al. (2021). CsAlaDC and CsTSI work coordinately to determine theanine

biosynthesis in tea plants (*Camellia sinensis* L.) and confer high levels of theanine

accumulation in a non-tea plant. Plant Biotechnol. J. **19**:2395-2397.

Zhu, Y. et al. (2006). *Arabidopsis* NRP1 and NRP2 encode histone chaperones and are

required for maintaining postembryonic root growth. Plant Cell **18**:2879-2892.

Acknowledgements

This work was supported by the National Key R&D Program of China (2022YFF1003103,

2021YFD1601101), and grants from the National Natural Science Foundation of China

(32072624), Anhui Provincial Major Science and Technology Project (202103b06020024)

and   Anhui   Educational   Committee   Excellent   Youth   Talent   Support   project

(gxyqZD2022018). We want to thank Oebiotech company provide 10× scRNA-seq service.

Author contributions

ZZ and WX conceived the research and finalized the manuscript. SL, YZ, SZ, YW, MH,

YD, JG and BZ collected materials and carried out the experiments. SL and YZ analyzed

the data. TY and EX offered guidance on experimental techniques and bioinformatics

analysis. SL and ZZ and wrote the manuscript. WJL offered suggestions on the study and

made revisions to the writing of this manuscript. All authors read and approved the

manuscript.

Competing interests

The authors declare no competing interests.

Figures and legends

 Figure 1 Process employed for scRNA-seq of tea plant roots. **(A)** Tea seedlings were used

 for theanine detection and scRNA-seq. Tea plant roots (~2.0 cm in length from the root tip)

 were harvested and used for scRNA-seq. **(B)** Theanine contents of leaf, stem, cotyledon, and

 root. **(C)** Cross section of tea plant roots (from the root tip back ~2.0 cm). Co, Cortex; Xy,

 Xylem; Ph, Phloem; Ca, Cambium; Pe, Pericycle; En, Endodermis; EP, Epidermis. Scale bar =

 200 μm. **(D)** Flow chart of scRNA-seq analyses.

 Figure 2 Annotation of cell clusters based on scRNA-seq analysis of tea plant roots. **(A)**

 t-SNE visualization plot of 10,435 tea plant root cells showed that they were grouped into 8 cell

 clusters. Each dot denoted a single cell. **(B)** *t*-SNE visualization of cell-type marker genes of

 root cells. Color bar indicates gene expression level. **(C)** *In situ* RT-PCR of *CsWAT1*, *CsPP2*,

CsAAP1 and *CsLHW*. The blue areas of sliced tissue represent regions where genes are

expressed. The red box represents the magnified areas of root sections. Scale bar = 200 μm. **(D)**

Violin plot showed the expression patterns for the top 5 marker genes of each cell clusters. The

height of violin represents the gene expression level, and the width represents the proportion of

cells expressing in the cluster. Asterisks represent homologous *Arabidopsis thaliana* marker

genes that can be found in PlantscRNAdb. **(E)** Schematic of the tea plant root anatomy, cell

types and their associated clusters.

Figure 3 Trajectory of root cell differentiation in the tea plant root. **(A)** Schematic

representation of root meristem organization (in median longitudinal section). *t*-SNE

visualization and cell type annotation of Cluster 3 (bottom). **(B)** Simulation of the successive

differentiation trajectory of the stem cell niche (SCN) over pseudo-time. Each dot denotes a

unique cell. Pseudo-time analysis of all 10,435 tea plant root cells (center), and the detailed

distribution of 8 cell clusters along pseudo-time trajectory (outer). Colors of the dots represent

the pseudo-time score. Red and blue line mark major differentiation trajectories, black lines

mark minor differentiation trajectories. **(C)** Heatmap illustrating the expression patterns of

differential genes, along the pseudo-time trajectory, during cell differentiation of the

vasculature and epidermis. Each row represents one gene. Color bars indicate the relative

expression levels. Biological processes are given on the right. **(D)** Proposed tea plant roots cell

differentiation trajectories for cell clusters shown on the *t*-SNE visualization plots. Dotted

arrows represent assumed cell differentiation routes.

Figure 4 Cell heterogeneity of nitrogen (N) transport and assimilation in tea plant roots.

(A) Schematic of N transport and metabolism pathway in tea plant roots. Membrane-located transporter NRTs and AMTs uptake nitrate and ammonium into root cells, respectively, and subsequently NO_3^- is assimilated into NH_4^+ and then metabolized into various amino acids. (B) Heatmap shows cell cluster expression patterns of N transport and assimilation genes in tea plant roots. Row normalization for gene expression according to "normalized" mode. (C) *In situ* RT-PCR of *CsAMT1.1*, *CsAMT3.1*, *CsNRT1.1* and *CsNRT3.2*. The blue areas of sectioned

tissues represent regions where genes are expressed. The red boxes represent the magnified

areas of root sections shown on the right. Scale bar = 200 μm. **(D)** Expression of amino acid

synthesis pathway genes in different cell clusters. Color bars indicate the expression levels of

amino acids synthesis pathway genes. Dot size indicates the percentage of amino acid synthesis

genes expressed in a cell cluster.

Figure 5 Cell heterogeneity of theanine synthesis, transport, and regulation in tea plant

roots. (A) Model of the putative theanine synthesis, transport, and regulation in tea plant root

cells. Black full lines represent unique metabolic processes in tea plants; grey full lines

represent common metabolic pathways in plants; dotted lines indicate the multi-step process.

(B) Heatmap for cell cluster expression patterns of genes encoding theanine transporters

CsAAPs, key enzymes and transcription factors in the theanine metabolic pathway. Gene

expression is presented in "normalized" mode. **(C)** *t*-SNE visualization of *CsTSI, CsAlaDC* and

their overlap map. Blue color bar indicates *CsTSI* expression level, red color bar indicates

CsAlaDC expression level in scRNA-seq. **(D)**Tissue localization of *CsTSI* and *CsAlaDC* using

in situ PCR. Blue signal indicates gene expression in cells. The red boxes represent magnified

areas of root sections shown below. Scale bar = 200 μm.

Figure 6 A gene co-expression network predicted key regulators of *CsAlaDC*. (A) Using

bulk RNA-seq data, a gene co-expression network was generated for *CsAlaDC* (|correlation

coefficient| ≥ 0.85 and *p*-Value ≤ 0.05). Red and blue dots represent differential genes

positively and negatively correlated with *CsAlaDC*, respectively. Bulk RNA-seq data is from

different types of N treatment of tea plant roots (Yang et al., 2019). **(B)** Heatmap showing

expression of co-expressed genes with *CsAlaDC* in eight tea plant tissues. The black arrows

denote root specific co-expressed (RSCE) genes. **(C)** Heatmap showing the expression pattern

for RSCE genes under different types of N treatment (0 N, CK, EA, NH_4^+ and NO_3^-).

Correlation between RSCE genes and *CsAlaDC* is shown on the heatmap at left. **(D)** Heatmap

 showing the expression level of RSCE genes in each cell cluster. **(B-D)** Row normalization of

 gene expression is according to the "normalized" mode.

Figure 7 Theanine synthesis regulated by the transcription factor CsLBD37. (A)
Schematic representation of the tea plant tissues used for gene detection. **(B)** Relative
expression of *CsLBD37* in different tea plants tissues. **(C)** The *t*-SNE visualization graph shows
expression levels of *CsLBD37* in various cell clusters. **(D)** Cell specificity of *CsLBD37*
expression in tea plant roots; blue signal represents location of gene expression. Scale bar =
200 μm. **(E)** Subcellular localization of CsLBD37; green fluorescence signal represents protein
localization. **(F)** The binding of CsLBD37 to the CsAlaDC promoter, tested by yeast

one-hybrid assay (Y1H). **(G)** EMSA assay showing the association of CsLBD37 with the

CsAlaDC promoter. The red arrow points to the binding position. **(H)** A schematic of the

effector and reportor constructs for transcriptional activity of the *CsAlaDC* promoter (top). LCI

assays in tobacco leaves show that the *CsAlaDC* promoter can interact with CsLBD37. The

color bar indicates the range of luminescence intensity (mid). The LUC/REN values represent

relative activity of CsLBD37 on *CsAlaDC* expression. Significant difference was evaluated by

two-tailed Student's *t*-test analysis (bottom). **(I-L)** *CsLBD37* gene silencing and overexpression

in tea plant roots. The expression level of *CsLBD37* and *CsAlaDC* in gene silenced **(I)** and

overexpression plants **(K).** Ethylamine and theanine contents in gene silenced **(J)** and

overexpression plants. **(L).** sODN, sense oligo nucleotides; asODN, antisense

oligonucleotides, silenced *CsLBD37* in tea plant root by asODN methods; EV, empty vector;

CsLBD37-OE/*Cs*, *CsLBD37* overexpression in transgenic hairy root system. Significant

difference was determined by Welch's t test.

Figure 8 Proposed model for theanine biosynthesis, transport and regulation at single

cell resolution. (A) Multicellular compartmentation of theanine biosynthesis. The high

expression of *CsTSI* in pericycle cells suggests theanine is mainly synthesized in pericycle

cells. *CsAlaDC* is highly expressed in cambium and protoxylem cells, suggesting that

ethylamine (EA) is mainly synthesized in these cells. Thus, EA may move from cambium and

protoxylem cells into pericycle cells, where it is a substrate for theanine biosynthesis. **(B)** In

tea plant roots, CsAAP1 likely mediates in theanine retrieval from apoplast for its transport

into cortical/endodermal cells. CsCAT2 imports theanine into vacuoles, especially in

pericycle cells, which are the main cells for theanine biosynthesis. Theanine is exported, by

an unidentified theanine exporter, from pericycle cells into the apoplastic pathway for entry

into the xylem. **(C)** CsLBD37 co-regulates theanine biosynthesis and lateral root development.

Increasing levels of nitrogen (N) promote theanine biosynthesis and accumulation. High

accumulation of theanine and N induces *CsLBD37* expression. CsLBD37 inhibits *CsAlaDC*

expression to reduce ethylamine synthesis, which finetunes theanine biosynthesis in feedback

loop. Besides, *CsLBD37* is expressed in pericycle cells and also inhibits lateral root

development. It is reported that apoplastic theanine can negatively regulate lateral root

development, under high nitrogen levels (Chen et al., 2022). Therefore, CsLBD37 may also be

involved in theanine-regulated lateral root development.

Supplemental Figure 1. Summary of the tea plant root scRNA-seq data. (A) Filter

delocalized cell maps fitted to generalized linear models. The horizontal axis is the unique

molecular identifiers (UMI) number in each cell, and the vertical axis is the gene number in

each cell. According to the linear relationship between the UMI and genes per cell data, the

distribution model is fitted. The yellow points represent delocalized cells, which will be

removed in the follow-up analysis. **(B)** The cell number in each cell cluster based on the tea

plant root scRNA-seq data.

Supplemental Figure 2. Cell cluster annotation. (A) t-SNE visualization of cell-type marker

genes (*CsGL3, CsSCR, CsIRX9 and CsWOX5*) of root cells. Color bar indicated gene

expression level. **(B)** Violin plot showing expression patterns of 6^{th} to 10^{th} in the top10

differential marker genes of each cell cluster. The height and width of the violin represent the

gene expression level and the proportion of cells expressing in the cluster, respectively.

Asterisks represent homologous marker genes that are present in PlantscRNAdb. **(C)** The

tissue-located heatmap of *LAC7, PME41, NRP1* and *TIM17*. The color bar illustrated gene

expression level for the scRNA-seq data. The *Arabidopsis thaliana* data and pictures were

generated from the Root Cell Atlas (https://rootcellatlas.org/).

Supplemental Figure 3. t-SNE visualization of Cluster 1 and Cluster 2 top 10 marker

genes. t-SNE plots show the transcript accumulation of the Cluster 1 and Cluster 2 top 10

marker genes in individual cells. Color intensity indicates the relative transcript level for the

indicated gene in each cell.

Supplemental Figure 4. t-SNE visualization of Cluster 3 and Cluster 4 top 10 marker

genes. t-SNE plots shows the transcript accumulation of the Cluster 3 and Cluster 4 top 10

marker genes in individual cells. Color intensity indicates the relative transcript level for the

indicated gene in each cell.

Supplemental Figure 5. t-SNE visualization of Cluster 5 and Cluster 6 top 10 marker

genes. t-SNE plots shows the transcript accumulation of the Cluster 5 and Cluster 6 top 10

marker genes in individual cells. Color intensity indicates the relative transcript level for the

indicated gene in each cell.

Supplemental Figure 6. *t*-SNE visualization of Cluster 7 and Cluster 8 top 10 marker genes. *t*-SNE plots shows the transcript accumulation of the Cluster 7 and Cluster 8 top 10 marker genes in individual cells. Color intensity indicates the relative transcript level for the indicated gene in each cell.

Supplemental Figure 7. UMAP visualization of eight clusters and differentiation trajectory atop the UMAP. **(A)** UMAP visualization plot of 10,435 tea plant root cells. Each dot denoted a single cell. **(B)** Simulation of the successive differentiation trajectory of tea plant root cells over pseudo-time. Colors of the dots represent the pseudo-time score. Red and blue line mark major differentiation trajectories.

Supplemental Figure 8. Cell cluster assay of regulators and target genes based on published *Arabidopsis thaliana* data. **(A)** Schematic diagram of reported gene regulatory relationships. A hierarchical regulatory network with IAA14 as the upstream regulator (top). MYB-bHLH-WD40 (MBW) complex regulates downstream gene (bottom). **(B) and (C)** Heatmap (B) and tissue-located heatmap (C) for cell cluster expression patterns of genes which were showed in schematic diagram of (A). The scRNA-seq data of *Arabidopsis thaliana* root was obtained from publicly available data (Ryu et al., 2019; Wendrich et al., 2020; Zhang et al., 2019; Denyer et al., 2019; Jean-Baptiste et al. 2019; Shulse et al., 2019; Shahan et al., 2022) and database (Root Cell Atlas, https://rootcellatlas.org/; BAR, https://bar.utoronto.ca/#GeneExpressionAndProteinTools). The pictures of (C) were generated from Root Cell Atlas web with slight modification.

Supplemental Figure 9. Protein purification of *CsLBD37*. Protein purification of *CsLBD37*. The CsLBD37-GST fusion protein has a molecular weight of 50.9 kDa (i.e., 24.9 kDa + 26 kDa).

Supplemental Figure 10. Overexpression of *CsLBD37* in *Arabidopsis* inhibited lateral root development. **(A)** qRT-PCR assay for *CsLBD37* transcriptional level in three transgenic *Arabidopsis* lines (*CsLBD37*-OE1/*At*, *CsLBD37*-OE2/*At*, *CsLBD37*-OE3/*At*) and wild-type *Arabidopsis* (WT) plants. **(B and C)** The plant phenotypes of *CsLBD37* transgenic *Arabidopsis* lines and WT grown in MS medium for 12 days. Scale bar = 1 cm. **(C)** Lateral

root number, each line contained 25 independent biological replicates. **(D-G)**

Microscopy-based lateral root developmental phenotypes of *CsLBD37* transgenic *Arabidopsis*

lines and WT, 5 days post germination. Scale bar = 25 μm in **(D)**. Lateral root initiation

number **(E)**; lateral root primordium number **(F)** and lateral root emergence number **(G)**, each

line contained 20 independent biological replicates. **(H)** qRT-PCR assay for *CsLBD37*

expression level in tea seedlings grown under control (CK) or 1 mM theanine treatment

(Theanine). **(C and E-H)** Significant differences were determined by two-sided Student's

t-test.