Root-specific secondary metabolism at the single-cell level: a case study of theanine metabolism and regulation in the roots of tea plants (*Camellia sinensis*)

Shijia Lin[1,†], Yiwen Zhang[1,†], Shupei Zhang[1], Yijie Wei[1], Mengxue Han[1], Yamei Deng[1], Jiayi Guo[1], Biying Zhu[1], Tianyuan Yang[1], Enhua Xia[1], Xiaochun Wan[1], William J. Lucas[2], Zhaoliang Zhang[1,*]

[1]State Key Laboratory of Tea Plant Biology and Utilization, Anhui Agricultural University, Hefei, Anhui 230036, China

[2]Department of Plant Biology, College of Biological Sciences, University of California, Davis, CA 95616, USA

[†]These authors contributed equally to this work

[*]Corresponding author (Email: zhlzhang@ahau.edu.cn)

Abstract

Root-synthesized secondary metabolites are critical quality-conferring compounds of foods, plant-derived medicines, and beverages. However, information at a single-cell level on root-specific secondary metabolism remains largely unexplored. L-theanine, an important quality component of tea, is primarily synthesized in roots, from which it is then transported to new tea shoots. In this study, we present a single-cell RNA sequencing (scRNA-seq)-derived map for the tea root, which enabled cell-type-specific analysis of glutamate and ethylamine (two precursors of theanine biosynthesis) metabolism, and theanine biosynthesis, storage, and transport. Our findings support a model in which the theanine biosynthesis pathway occurs via multicellular compartmentation and does not require high co-expression levels of transcription factors and their target genes within the same cell cluster. This study provides novel insights into theanine metabolism and regulation, at the single-cell level, and offers an example for studying root-specific secondary metabolism in other plant systems.

Key words: scRNA-seq, theanine, metabolism, regulation, root development, *Camellia sinensis*

Introduction

Different root cell types produce diverse and complex secondary metabolites in response to environmental and developmental clues (Hong et al., 2017; Shaw et al., 2021); furthermore, such secondary metabolites can also participate in environmental adaptation and development (Garcia-Lemos et al., 2020; Hartman et al., 2017; Santelia et al., 2008; Schulz-Bohm et al., 2018; Steppuhn et al., 2004; Wan et al., 2018; Xu et al., 2017). As examples, the alkaloid nicotine is synthesized, specifically, in the root and is transported to the shoot for resistance to biotic stresses (Steppuhn et al., 2004; Xu et al., 2017); root-synthesized flavonoids regulate root tip growth through affecting auxin transport and metabolism (Santelia et al., 2008; Wan et al., 2018); legume roots secrete flavonoids as signaling agents to attract symbiotic bacteria, such as *Rhizobium* for nitrogen fixation (Hartman et al., 2017); in *Abies nordmanniana*, volatile organic compounds (e.g., propanal, g-nonalactone, and dimethyl disulfide) function to recruit certain bacteria or fungi, such as *Paenibacillus* sp. S37, which produce high quantities of indole-3-acetic acid that can then promote plant root growth (Garcia-Lemos et al., 2020; Schulz-Bohm et al., 2018).

In many medicinal and beverage plants, root-synthesized secondary metabolites are critical quality compounds (Bailly, 2021; Hu et al., 2019; Schmid et al., 2018). Lobetyolin (LBT), a *Codonopsis pilosula* (known as Radix codonopsis or Dangshen) root-synthesized polyacetylene glycoside, exhibits activities against various cancers, notably, gastric cancer (Bailly, 2021). Polysaccharides and lectins isolated from *Pseudostellaria heterophylla* (known as Taizishen or Haiershen) roots, have multiple pharmacological activities (Hu et al., 2019). Root-synthesized polyphenols and triterpenoid saponins are crucial quality compounds extracted from *Glycyrrhiza glabra* (known as Licorice) and are employed in the food and beverage industry (Schmid et al., 2018).

Due to the importance of secondary metabolites, the biosynthesis, transport, and regulation of plant root-synthesized secondary metabolites have been extensively studied (Fernie and Tohge, 2017; Wang et al., 2018b). However, in most cases, whole root materials were employed with very few assessing secondary metabolite biosynthesis and regulation, at the single-cell level.

Tea plant (*Camellia sinensis* L.), a crop with high economic value, is a member of the

eudicot family, *Theaceae*, has diverse and active secondary metabolic activities (Yu et al.,

2020). Tea plants produce and accumulate extensive amounts of secondary metabolites,

including theanine, catechins and caffeine (Zhao et al., 2020). These secondary metabolites

confer tea with both unique flavor and multiple health benefits (Zhao et al., 2020). Theanine, a

non-protein amino acid, endows the "umami" taste and relaxing effect of the tea infusion (Lin

et al., 2022). It is the most abundant free amino acid in the tea plant, accounting for 1% - 2%

dry weight of the tender shoots (Lin et al., 2022). It is primarily synthesized in tea roots and is

then transported to the shoots, via the vasculature system (Figure 1A, B) (Lin et al., 2022).

Tea roots acquire inorganic nitrogen, especially ammonium, from the soil and assimilate it

into amino acids, such as glutamate (Glu), glutamine (Gln), and alanine (Ala) (Lin et al., 2022).

Ala can be decarboxylated, by alanine decarboxylase (CsAlaDC), to produce ethylamine (EA)

(Bai et al., 2019; Zhu et al., 2021), which can then be combined with Glu, through the action of

theanine synthase (CsTSI), to produce theanine (She et al., 2022; Wei et al., 2018). The high

level of EA availability is likely a key factor in the elevated accumulation of theanine in tea

plants (Cheng et al., 2017; Zhu et al., 2021). Importantly, in the tea plant, CsAlaDC and CsTSI

are both indispensable for theanine synthesis (Zhu et al., 2021).

Theanine synthesis is tightly regulated, at multiple levels, especially at the transcriptional

level (Lin et al., 2022, Guo et al., 2022); CsMYB6 bound to the *CsTSI* promoter can regulate

theanine synthesis (Zhang et al., 2021c). Furthermore, CsMYB40 and CsHHO3 bound to the

CsAlaDC promoter can regulate theanine synthesis in "accelerator" or "brake" mode,

respectively, in response to N levels (Guo et al., 2022).

Theanine   transporters,   including   the   tonoplast-localized   CsCAT2   and   plasma

membrane-localized CsAAPs, have been proposed to mediate in theanine storage within the

root and root-to-shoot transport, respectively (Dong et al., 2020; Feng et al., 2021). In

addition, in response to N status in tea roots, theanine may affect apoplastic H_2O_2

accumulation to regulate lateral root development (Chen et al., 2022). Taken together, these

studies have partially revealed the mechanisms for theanine synthesis, transport, and

regulation, but information on the specific root cell types involved in these processes is

lacking.

Recently, with the in-depth study of biological structure and function, it has become

increasingly clear that differences exist in gene expression levels between cells, even if they

appear to be within the same cell population. Bulk RNA-seq can be employed to investigate

the average level of gene expression within whole tissues, whereas single-cell RNA sequencing

(scRNA-seq) can capture single cells, from whole tissues, to detect the heterogeneity of cells,

and obtain information on gene expression within single cells (Shaw et al., 2021). Thus,

scRNA-seq provides a powerful approach to elucidate the potential involvement of cell

heterogeneity in secondary metabolism in tea plant roots.

Currently, scRNA-seq has been performed in several plants, including *Arabidopsis thaliana*

(Ryu et al., 2019), *Oryza sativa* (Zhang et al., 2021b), and *Zea mays* (Nelms and Walbot, 2019).

Importantly, algorithms are available to decode the cellular regulatory codes from scRNA-seq

datasets (Becht et al., 2018; Haghverdi et al., 2015; Haghverdi et al., 2016; Trapnell et al.,

2014), and these databases can provide a critical foundation for analyzing the cellular-based

information on gene expression (Chen et al., 2021; Xu et al., 2022).

In plants, scRNA-seq has yet to be used for studying root-specific secondary metabolism.

In this study, using high-throughput scRNA-seq, we analyzed the cell heterogeneity of

theanine metabolism and regulation in the tea plant root. Our findings offer important insights

into the molecular mechanisms underlying theanine synthesis, transport, and regulation, and

provide a basis for studies on root-specific secondary metabolism in other plant systems.

Results

Tea plant root cell clusters identified by scRNA-seq

Tea plant roots synthesize and accumulate large amount of theanine (Figure 1A and 1B). The

tea root cell types, including cortical (Co), xylem (Xy), phloem (Ph), cambial (Ca), pericycle

(Pe), endodermal (En), epidermal (Ep) and root hair (RH), are shown in Figure 1C. To

understand the cellular characteristics of theanine biosynthesis in the tea root, protoplasts were

isolated from seedling roots (from the root tip back ~2.0 cm) of the tea plant cultivar

'Shuchazao' (Figure 1A). These protoplasts were subjected to scRNA-seq, using the

commercial 10× Genomics platform (Figure 1D), and the sequencing data were pre-filtered, at

the cell and gene levels, resulting in a pool of 10,435 cells (Supplemental Figure 1A;

Supplemental Data 1). The *t*-distributed stochastic neighborhood embedding (*t*-SNE) algorithm

was employed to visualize local similarities (Haghverdi et al., 2015), resulting in these 10,435

cells being grouped into 8 cell clusters (Figure 2A; Supplemental Figure 1B).

The cell types within these 8 clusters were assessed by an analysis of homologs of cell

type marker genes according to protein sequence similarity, confirmed in other scRNA-seq

studies, or in the model plant *Arabidopsis thaliana*. In *A. thaliana,* GLABRA3 (GL3) is

involved in epidermal cell fate specification (Bernhardt et al., 2003). In our scRNA-seq,

CsGL3 transcripts were detected in Cluster 2, indicating the presence of epidermal cells

(Supplemental Figure 2A). The *SCARECROW* (*SCR*) gene is required for the asymmetric

division of the cortex/endodermis initial (CEI) cells, and its expression is confined to the

endodermis, CEI cell, and the quiescent center (QC) (Dong et al., 2021). In our tea root data,

CsSCR transcripts were detected primarily in Clusters 3 and 4 (Supplemental Figure 2A),

suggesting that cortical and endodermal cell division occurs in these two clusters. However, as

CsAAP1 transcripts were detected in Cluster 4 (Figure 2B), and the gene encodes for AMINO

ACID PERMEASE 1, which generally mediates in uptake of amino acids, into the cortex and

endodermis, this cluster likely contained a population of cortical or endodermal cells.

The stele system in the root is comprised of the pericycle and vasculature, in which

LONESOME HIGHWAY (*LHW*) has been established as a pericycle marker gene (De Rybel et

al., 2013). As the *CsLHW* transcripts were detected in Cluster 8, this cell cluster likely contains

a pericycle cell population (Figure 2B). The *WALLS ARE THIN LIKE 1* (*WAT1*) and

IRREGULAR XYLEM 9 (*IRX9*) are characterized vasculature and xylem development marker

genes, respectively (Li et al., 2021). Hence, the high level of *CsWAT1* transcripts detected in

Clusters 1 and Cluster 3 (Figure 2B), along with *CsIRX9* transcripts present in Cluster 3

(Supplemental Figure 2A) support the notion that these two clusters represent vasculature or

xylem-related cells in the tea root. Finally, the *PHLOEM PROTEIN 2* (*PP2*) gene participates

in phloem development and is a well-characterized phloem marker (Dinant et al., 2003) and

the presence in Cluster 7 of abundant *CsPP2* transcripts is consistent with phloem cells being

located within this Cluster 7 (Figure 2B).

To confirm the annotation of these cell clusters, we performed *in situ* RT-PCR analyses.

These assays indicated expression of the following genes in their respective cell types:

CsWAT1 in the vasculature (including pericycle, xylem parenchyma and cambium); *CsPP2* in

the protophloem; *CsAAP1* mainly in the cortex and endodermis; and *CsLHW* in the pericycle

(Figure 2C). These findings were consistent with the above annotation of these cell clusters

(Figure 2B and 2C).

Given the lack of more marker genes in tea plants, we next applied an unbiased approach,

developed by Denyer et al. (2019), in which cluster-enriched genes are used as markers to

define cell cluster identity. Cluster-enriched genes are defined as those being expressed, at

significantly higher level, in each cluster, than in all other clusters, and expressed in ⩾ 10%

of cells within this specific cluster. The information on the cell-specific expression of these

cluster-enriched genes was confirmed in *A. thaliana*. Therefore, we analyzed expression

patterns of the top 10 cluster-enriched genes (Figure 2D; Supplemental Figures 2-6;

Supplemental data 3). Here, we could establish enrichment of *LACCASE 7* (*CsLAC7*) and

PECTIN METHYLESTERASE 41 (*CsPME41*) in Cluster 2, *ASPARAGINE RICH PROTEIN1*

(*CsNRP1*) in Clusters 3 and 5, and *MITOCHONDRIAL IMPORT INNER MEMBRANE*

TRANSLOCASE subunit TIM17 (*CsTIM17*) in Cluster 5 (Figure 2D). Based on these

scRNA-seq results from *A. thaliana*, and previous studies showing that *AtLAC7* and *AtPME41*

are expressed mainly in epidermal cells, *AtNRP1* functions as a root meristem maker, including

stem cell niche (SCN) and root cap (RC), and *AtTIM17* is expressed in the root cap (Passarinho

et al., 2001; Zhu et al., 2006; Supplemental Figure 2C), we assigned Cluster 3 and 5 as SCN and

RC cells, respectively.

Taken together, according to the above results, Clusters 1, 2, 4 and 7 were annotated as

xylem, epidermal, cortex/endodermal, and phloem cells, respectively (Figure 2E). In addition,

Cluster 8 was annotated as pericycle cells (Figure 2E), whereas Cluster 6 included primarily

procambium or transition cells during differentiation (Figure 2E), and Cluster 3 was defined as

the SCN, which serves as the cluster containing founder cells for differentiation (Figure 2E).

Differentiation trajectories of tea root cells supported cell cluster annotations

During root cell differentiation, all root cell types originate from stem cells generated by the

root apical meristem (RAM). In the RAM, stem cells, including stelar initial (SI) cells, CEI

cells, epidermis/lateral root cap initial (ELI) cells and columella initial (CI) cells, surround the

QC (Efroni et al., 2016) (Figure 3A). The QC cells are required for specification of the SCN

and maintaining the undifferentiated state of the stem cell initials. Stem cells continuously

undergo asymmetric division to produce daughter cells, which then begin to differentiate into

outer, inner, and middle cell layers (Efroni et al., 2016) (Figure 3A). We noticed that Cluster 3

could probably be divided into 4 sub-cell states, C3.1, C3.2, C3.3 and C3.4 (Figure 3A), with

C3.1 expressing the QC marker gene, *WUS-RELATED HOMEOBOX 5* (*CsWOX5,* Kong et al.,

2015), C3.2 expressing the xylem development gene *CsIRX9*, C3.3 expressing the cortex and

endodermal cell fate decision gene, *CsSCR,* and C3.4 being close to the Cluster 2 epidermal cell

population (Figure 3A; Supplemental Figure 2A). Thus, these four Cluster 3 sub-cell states

were defined as QC, SI, CEI, and ELI cells, respectively (Figure 3A).

The differentiation trajectories of these clusters were next assessed to confirm our annotation

of the grouped cell clusters. To this end, we conducted pseudo-time analysis by ordering cells

of these clusters along a reconstructed trajectory (Figure 3B). This analysis showed that Cluster

3 was primarily located at the start of the differentiation trajectories, and that the trajectories of

these cells bifurcated into two major and one minor branch (Figure 3B). Here, the two major

branches reflected differentiation trajectories of the ELI and SI cells from the SCN (Cluster 3),

with the ELI cells differentiating into the outer cell layer (red branch) and Cluster 2 (epidermal

cells) and Cluster 5 (root cap cells) located at the end of this divergence (Figure 3B). The SI

cells differentiated to give an inner cell layer (blue branch): Cluster 7 (phloem cells) and

Cluster 1 (xylem cells) were arranged successively (Figure 3B). The minor branch resulted in

CEI cell trajectories that differentiated to give a middle cell layer (black branch); Cluster 4

(cortex and endodermal cells) were located at the end of this divergence. Importantly, Cluster 4

(cortex and endodermis), Cluster 6 (cambium) and Cluster 8 (pericycle) cells were also located

within the region where cell differentiation was initiated, thus suggesting these three cell

clusters can differentiate into other cell types (Figure 3B). Indeed, the cambium differentiates

into the xylem and the phloem, especially in woody plants (Li et al., 2021); endodermal cells

assume a SCI to generate a new epidermal layer (Efroni et al., 2016); the founder cells, derived

from the pericycle, can form lateral roots (Péret et al., 2009). The results of our pseudo-time

analyses matched well with these earlier established findings.

We next constructed a differentiation heatmap of xylem and epidermal cells (Figure 3C).

Consistently, highly expressed genes in Cluster 1 (xylem cells) were most enriched in cell wall

biogenesis, lignin catabolic process and amino sugar catabolic process, etc. (Figure 3C and

Supplemental data 4). Meanwhile, genes highly expressed in Cluster 2 (epidermal cells) were

enriched in trichoblast differentiation, root hair elongation and plant epidermal cell

differentiation, etc. (Figure 3C and Supplemental data 4). These in-depth differentiation

trajectories of root cells further indicated reliability of the cell cluster annotations.

Finally, a differentiation map of tea root cells was constructed, based on the pseudo-time

analyses (Figure 3D). This map was generated to assist in developing an understanding of tea

root development, to provide a basis for studying secondary metabolism, in tea roots, at the

single-cell level.

Cell heterogeneity of N transport and metabolism in tea roots

In tea roots, theanine accounts for approx. 60~80% of the total free amino acids (Yang et al.,

2020). Nitrogen (N) uptake and transport, in the form of nitrate and ammonium, and N

assimilation into amino acids, are prerequisites for theanine biosynthesis (Yang et al., 2020)

(Figure 4A). We therefore investigated the cell heterogeneity of these processes by analyzing

the scRNA-seq data.

The NITRATE TRANSPORTERS (CsNRTs) and AMMONIUM TRANSPORTERS

(CsAMTs) function in N uptake in tea plants (Zhang et al., 2021a; Zhang et al., 2020) (Figure

4A). The heatmap of relative expression revealed that most *CsNRTs* and *CsAMTs*, including

CsAMT1.1, *CsAMT1.2*, *CsNRT1.1*, *CsNRT1.5*, *CsNRT1.7*, *CsNRT3.1* and *CsNRT3.2*, were

expressed primarily in Cluster 4 (cortex and endodermal cells); *CsAMT1.2* and *CsNRT1.5*

were also highly expressed in Cluster 7 (phloem cells); *CsAMT3.1* was expressed primarily in

Cluster 8 (pericycle cells) and Cluster 6 (cambium cells)*; CsAMT3.1* and *CsNRT3.2* were

relatively specifically expressed in Cluster 2 (epidermal cells); *CsNRT2.4* was highly and

specifically expressed in Cluster 3 (SCN) (Figure 4B). At the same time, our *in situ* RT-PCR

results for *CsAMT1.1, CsAMT3.1*, *CsNRT1.1* and *CsNRT3.2* confirmed these scRNA-seq

findings (Figure 4B and 4C). Furthermore, these results were similar to previous reports in *A.*

thaliana and rice (Wang et al., 2018a; Wang et al., 2018b).

Intracellular NO_3^- is assimilated in the cytoplasm into NO_2^-, by NITRATE REDUCTASE

(NR), which then enters the plastids where it is further assimilated into NH_4^+, by NITRITE

REDUCTASE (NiR). Subsequently, NH_4^+ enters the amino acid metabolism pathway (Galili et

al., 2016; Xu et al., 2012) (Figure 4A). Our scRNA-seq assays revealed that expression of

CsNR1, *CsNR2*, *CsNiR1* and *CsNiR2* was more active in Cluster 8 (pericycle cells) (Figure

4B); *CsNR1* and *CsNR2* expression was also relatively high in Cluster 1 (xylem cells). These

data supported the notion that N assimilation occurs primarily within stelar cells in tea roots.

Previously, we grouped amino acid metabolism into five pathways (P1-P5), based on their

metabolic relationships (Yang et al., 2020): glutamate pathway (P1: Glu, Gln, arginine [Arg],

proline [Pro], citrulline [Cit]); asparate (Asp) pathway (P2: Asp, isoleucine [Ile], lysine [Lys],

threonine [Thr] and methionine [Met]); pyruvate pathway (P3: Ala, leucine [Leu] and valine

[Val]); aromatic amino acid pathway (P4: phenylalanine [Phe], tyrosine [Tyr] and tryptophan

[Trp]); and the 3-phosphoglycerate pathway (P5: serine [Ser], glycine [Gly] and cysteine

[Cys]) (Figure 4D and Supplemental data 5). Theanine is synthesized from Glu and

Ala-derived ethylamine (Lin et al., 2022).

Based on the scRNA-seq analysis, we observed that the expression level of genes encoding

for Glu biosynthetic pathway enzymes was the highest (especially in Clusters 1, 3, 4, 6 and 8)

within these five pathways (Figure 4D and Supplemental data 5), suggesting that biosynthetic

activity for Glu synthesis is highly active in the roots. Meanwhile, the expression level of

genes encoding alanine biosynthetic pathway enzymes was also high in all these clusters,

especially in Clusters 3 (SCN) and 8 (pericycle cells) (Figure 4D and Supplemental data 5).

These findings indicated that Glu and Ala metabolism represent the major amino acid

metabolism in tea roots. This is the basis for N metabolism into theanine in these roots.

Cell types participating in theanine biosynthesis, storage, and transport in tea roots

The genes encoding enzymes in the theanine biosynthetic pathway, theanine transporters, and

transcription factors regulating theanine biosynthesis, were recently identified (Dong et al.,

2020; Feng et al., 2021; Guo et al., 2022; She et al., 2022; Wei et al., 2018; Zhang et al.,

2021c) (Figure 5A). We analyzed the cell-specific expression patterns of these genes using

our scRNA-seq data (Figure 5B and Supplemental data 6). Impressively, although *CsTSI* was

highly expressed in all clusters (Figure 5C), it was expressed, more specifically, in Cluster 8

(pericycle cells) (Figure 5B). Here, our *in situ* RT-PCR assays also revealed that *CsTSI* is

expressed primarily in the pericycle (Figure 5D). Consistently, Glu biosynthetic pathway

genes, including *CsGOGAT1*, *CsGDH2* and *CsGDH3*, also had a relatively high expression level in Cluster 8 (Figure 5B), although some *CsGOGAT* and *CsGDH* members were also relatively highly expressed in Clusters 1 (xylem cells), 2 (epidermis cells), 3 (SCN), 4 (cortex and endodermal cells), 5 (root cap cells), 6 (cambium cells) and 7 (phloem cells) (Figure 5B). These findings suggested that theanine biosynthesis is more enriched in Cluster 8 (pericycle cells) compared with Glu biosynthesis.

The *CsTSI* and *CsAlaDC* are the most important genes for theanine synthesis (She et al., 2022; Zhu et al., 2021). However, we observed that *CsTSI* and *CsAlaDC* were not enriched in the same cluster. Different from *CsTSI*, *CsAlaDC* was most highly expressed in Cluster 1 (xylem cell) and was also relatively highly expressed in Cluster 6 (cambium cell) (Figure 5B and 5C). The *in situ* RT-PCR assays verified the expression pattern of *CsAlaDC* in the vasculature and cambium, and also showed that *CsAlaDC* was expressed, at a low level, in the pericycle (Figure 5D). Therefore, generally, *CsTSI* and *CsAlaDC* are co-expressed in the pericycle, where *CsTSI* is most highly expressed, whereas *CsAlaDC* is mostly highly expressed in the vasculature, where *CsTSI* has a lower expression level. These results suggest that the theanine biosynthesis pathway has a multicellular compartmentation characteristic, as EA would be synthesized primarily in the vasculature, whereas theanine would be more actively synthesized in the pericycle.

Generally, root-synthesized theanine is stored in the root cells, or transported from root to shoot (Dong et al., 2020; Lin et al., 2022). In this regard, CsAAPs (CsAAP1, CsAAP2, CsAAP4, CsAAP5, CsAAP6 and CsAAP8) and CsCAT2 were recently identified as theanine transporters (Dong et al., 2020; Feng et al., 2021). Among them, CsCAT2 probably mediates theanine storage in the vacuole, with CsAAP1 functioning in theanine loading into the xylem transpiration stream for root-to-shoot transport (Dong et al., 2020; Feng et al., 2021). Interestingly, *CsCAT2* was shown to be expressed most highly in the same cluster as *CsTSI*; i.e., in Cluster 8 (pericycle cells) (Figure 5B). *CsCAT2* was also relatively highly expressed in Cluster 4 (cortex and endodermal cells), suggesting that CsCAT2 may function in theanine storage in the pericycle, cortex and endodermis. In contrast, *CsAAP1* was expressed primarily in Cluster 4 (cortex and endodermal cells). This is consistent with its proposed role in retrieving extracellular theanine for its transport through the endodermis prior to xylem loading (Dong et al., 2020; Lin et al., 2022). The expression of other

297 *CsAAPs* was also shown to be highly cell cluster-specific; for example, *CsAAP2* in Cluster 4

298 (cortex and endodermal cells), *CsAAP4* in Cluster 6 (cambium cells), *CsAAP6* in Cluster 3 (SCN)

299 and Cluster 4 (cortex and endodermal cells), and *CsAAP8* in Cluster 5 (root cap cells) (Figure 5B),

300 suggesting cell type-specific functions for these CsAAPs in theanine transport.

Transcriptional regulation of theanine biosynthesis

Theanine biosynthesis is regulated at the transcriptional level, based on recent findings that

transcription factors (TFs) can control *CsTSI* or *CsAlaDC* expression to regulate theanine

biosynthesis (Zhang et al., 2021; Guo et al., 2022). Here, Zhang et al. reported that CsMYB6, a

root-specific expressed TF, promoted theanine biosynthesis through activation of *CsTSI*

expression. Thus, we were surprised that *CsMYB6* was expressed in Cluster 3 (SCN), Cluster 6

(cambium cells) and Cluster 1 (xylem cells), rather than in Cluster 8 (the high *CsTSI* expression

cell cluster), as shown in Figure 5B.

A similar situation was also observed in terms of *CsAlaDC* transcriptional regulation. We

previously identified CsMYB40 and CsHHO3 to be an activator and repressor of *CsAlaDC*

expression, respectively, in response to N levels (Guo et al., 2022). Analysis of our scRNA-seq

data indicated that *CsMYB40* and *CsHHO3* were not highly expressed in Cluster 1 (the high

CsAlaDC expression cell cluster); *CsMYB40* was relatively highly expressed in Cluster 4 (cortex

and endodermal cells) and Cluster 7 (phloem cells); and *CsHHO3* was relatively highly expressed

in Clusters 3 (SCN) and 6 (cambium cells) (Figure 5B).

Based on these findings, we hypothesized that, for the regulation of theanine biosynthesis, TFs

and target genes are not necessarily always highly expressed in the same cells. In the theanine

biosynthesis pathway, CsAlaDC evolved a tea plant-specific activity in catalyzing EA biosynthesis

(Cheng et al., 2017), which is essential for theanine biosynthesis (Zhu et al., 2021). *CsAlaDC* is

highly and specifically express in tea plant roots (Zhu et al., 2021), and its expression levels were

shown to be highly correlated with theanine content in the roots of various tea plant cultivars, and

further, was under TF regulation (Zhu et al., 2021). Thus, taking the transcriptional regulation of

CsAlaDC as an example, we next analyzed the TFs that were co-expressed with *CsAlaDC* to

test this notion.

Identification of CsLBD37, a novel TF that co-regulates both theanine synthesis

and lateral root development

In tea plant roots, both theanine biosynthesis and root development are regulated by N level and its form (Yang et al., 2020). Moreover, theanine is probably involved in N-regulated lateral root development (Chen et al., 2022). This implies the presence of TFs that can co-regulate theanine biosynthesis and development in the tea root. To test this notion, we identified genes co-expressed with *CsAlaDC*, using our previous transcriptome data obtained for roots that were supplied with various N treatments, including without N (0N), 1.4 mM nitrate (NO_3^-), 1.4 mM EA (EA), 1.4 mM ammonium (NH_4^+), or 1.4 mM nitrate plus ammonium (1:1) (CK) (Supplemental data 9 and 10). By filtering of these genes, using |correlation coefficients| $\geq$ 0.85 and $p \leq 0.05$, we constructed a *CsAlaDC* co-expression network that consisted of 133 genes (Figure 6A and Supplemental data 10).

The expression patterns of these *CsAlaDC* co-expressing genes were assessed in eight tissues, based on comprehensive tea plant transcriptome data (Wei et al., 2018; Xia et al., 2019) (Figure 6B). Given that *CsAlaDC* expression is root-specific, we chose 34 root-specific co-expressed genes (RSCGs) for further analysis (Figure 6B). Expression analyses of these RSCGs showed that the genes which were positively correlated with *CsAlaDC* expression were induced by N, especially by EA, and the negatively correlated genes were induced by a 0 N stress treatment (Figure 6C). Surprisingly, the results of the scRNA-seq analysis indicated that most of these genes, which were highly correlated with *CsAlaDC*, were not highly expressed in Cluster 1 (xylem cell) (Figure 6D).

Within these 34 root-specific genes, there were 7 genes potentially encoding for TFs LBD37, LBD38, bZIP, WRKY, AP2 and MYB18 (Figure 6C). Among them, *CsLBD37* (CSS0024718) had the most positive correlation with *CsAlaDC,* and was induced by N (CK, EA and NH_4^+) and was not expressed highly in Cluster 1 (Figure 6C and 6D). Du et al. (2021) predicted that *CsLBD38*, named *CsLBD37* in this study, is a highly credible TF that regulates theanine synthesis and is in an evolution node of theanine-associated regulatory module (TARM) in *Camellia* and Non-*Camellia* species. Therefore, CsLBD37 probably regulates *CsAlaDC* expression in tea plants, but its cell type expression pattern is different from that of *CsAlaDC*.

To further explore this situation, we first confirmed the root-specific expression for

CsLBD37 (Figure 7A and 7B). Next, we performed *in situ* RT-PCR assays, which indicated that the *CsLBD37* expression pattern did not exactly reflect that observed for *CsAlaDC*. As for *CsLBD37*, *CsAlaDC* was expressed in the cambium; however, it was also expressed in the pericycle, whereas *CsAlaDC* was barely expressed in these cells (Figure 5C and Figure 7C). As expected, CsLBD37 was localized in the nucleus (Figure 7E), and furthermore, yeast one-hybrid (Y1H) assays, combined with *CsAlaDC* promoter-*LUC* assays conducted in tobacco leaves, were consistent with CsLBD37 binding directly to and repressing the activity of the *CsAlaDC* promoter (Figure 7F and 7H). We further conducted an electrophoretic mobility shift assay (EMSA) with recombinant CsLBD37 protein and showed that it bound to the proximal region (-476 to -277 bp) of the *CsAlaDC* promoter (Figure 7G and Supplemental Figure S6).

To test whether *CsLBD37* was able to inhibit *CsAlaDC* expression, *in vivo*, we developed transiently *CsLBD37*-silenced or over-expression tea seedlings, by antisense oligonucleotide (asODN) interference (Figure 7I) and generation of transgenic hairy roots, respectively (Figure 7K). *CsLBD37* expression was down-regulated and *CsAlaDC* was up-regulated in these asODN-treated roots (Figure 7I), whereas EA and theanine levels were increased (Figure 7J). In contrast, in *CsLBD37* transgenic hairy roots, *CsLBD37* was up-regulated and *CsAlaDC* was down-regulated (Figure 7K), whereas EA and theanine levels were decreased (Figure 7L). These findings offered support for the hypothesis that *CsLBD37* regulates *CsAlaDC* expression to modulate EA and theanine synthesis. *CsLBD37* probably acts as a "brake" to maintain the expression level of *CsAlaDC* within a certain range, in response to high levels of N. This may be why *CsLBD37* is a repressor but is positively correlated with *CsAlaDC*.

Some studies have reported that the LBD family of TFs can regulate both plant secondary metabolism and root development (Goh et al., 2012; Okushima et al., 2007; Ye et al., 2021). Lateral root founder cells originate from the pericycle, located opposite to the xylem poles (Péret et al., 2009). We noted that *CsLBD37* was also expressed in the tea root pericycle (Figure 7C). Therefore, we overexpressed *CsLBD37* in *A. thaliana* (Supplemental Figure 8A and 8B) and observed that the number of lateral roots, in these overexpression (OE) lines, was significantly less than in the wild-type (WT) plants (Supplemental Figure 8B and 8C). We

further observed that *CsLBD37* overexpression repressed the initial steps of lateral root

development (Supplemental Figure 8D-8G). These results provided support for a model in

which CsLBD37 also regulates lateral root development in tea plants. Moreover, *CsLBD37*

expression was induced by exogenous theanine treatment and this result underscores once

again the importance of CsLBD37 in co-regulating theanine biosynthesis and root

development (Supplemental Figure 8H).

Collectively, our findings support a model in which transcriptional regulation of theanine

biosynthesis does not require high expression of both TFs and target genes in the same cell

cluster, but only requires partial co-expression. We also established that *CsLBD37*

co-regulates both theanine biosynthesis and lateral root development in tea plants under

N-rich conditions.

Discussion

Application of single-cell transcriptome sequencing to plant studies has afforded progress in

understanding root development in model plants, such as *A. thaliana* and rice (Ryu et al., 2019;

Zhang et al., 2021b). In this regard, important aspects of secondary and specialized

metabolism occur in the plant root system and are important for plant growth and adaption to

environmental challenges. These metabolic products are also essential for food and beverage

quality, and pharmaceutical products. Bird et al. (2005) reported that biosynthesis of the

narcotic analgesics, morphine and codeine, is localized to sieve elements in the opium poppy,

which extends the function of sieve elements beyond transport, and into the realm of

metabolism. However, significantly less information is currently available regarding both the

cell heterogeneity of these secondary metabolic pathways and the underlying regulatory

events that occur in the plant root system, especially in woody plants. The information

regarding gene expression, in different type cells, is important with respect to establishing

insights into both the mechanisms underlying these secondary metabolites and the regulatory

events that control their syntheses. In this study, we performed a scRNA-seq-based study of

tea roots, as an example, to elucidate the biosynthesis and regulatory processes for theanine, a

root-specific secondary metabolite. In this process, we established the first map of tea roots

comprised of 8 cell clusters (Figures 2A and 3D).

Cell cluster annotation is a critical step for scRNA-seq. In this regard, we employed

candidate marker genes to annotate cell clusters, as in previous studies (Ryu et al., 2019; Zhang

et al., 2021b). In some non-model plants, including tea, transgenic technologies are not

currently available and, hence, we used *in situ* RNA hybridization to establish the location(s)

for gene expression. In some studies, isolation of different cell types was combined with

q-RT-PCR to detect cell-type marker gene expression (Wang et al., 2022). However, this

approach has two limitations in that it cannot display the gene location directly and has only

low resolution.

After numerous trials, we were able to optimize *in situ* RT-PCR assays (detailed in the

Methods), which enabled a cell-specific characterization of gene expression in tea root cells,

prior to establishing a genetic transformation system for tea. However, we note the challenge

associated with weak calling of homologous marker genes, which may reflect the differences

between tea and *A. thaliana*, along with differences in sampling sites, but this only slightly

impacted our findings, as in the annotation of cell clusters, *in situ* RT-PCR was employed to

identify expression patterns for weakly expressed tea root genes, such as *CsAAP1, CsLHW,*

CsAMT1.1, CsNRT1.1. Importantly, the results of *in situ* RT-PCR and scRNA-seq analysis were

generally consistent.

N is one of the most critical mineral nutrients essential for crop growth and yield

performance and provides precursors for the biosynthesis of secondary metabolites. Roots

acquire NO_3^- -N and NH_4^+-N from the soil (Xu et al., 2012), with CsNRTs (CsNRT2.5 and

CsNRT3.2) and CsAMTs (CsAMT3.1)   acting in N uptake and transport in tea roots (Zhang

et al., 2021a; Zhang et al., 2020). Our current findings provided further insight into the

cellular locations where these transporters function. We show that many *CsNRTs* and *CsAMTs*

are in the cortex, endodermal and stelar cells, whereas the N assimilation genes, *CsNRs* and

CsNiRs, were primarily located within the stelar cells (Figure 4B). Our data support a model

in which these CsNRTs and CsAMTs function in N uptake, into the cytoplasm of cortical and

endodermal cells, but also play an important role in NO_3^- and NH_4^+ loading into the cells

within the stele for N assimilation and amino acids synthesis.

The Glu pathway is the most abundant and active pathway for amino acid biosynthesis in

all root cell types. However, our findings indicated that stelar cells and the SCN have a higher

capacity for amino acid synthesis, compared with epidermal, cortical, and endodermal cells

(Figure 4C). For 20 amino acids, their rates of synthesis appeared to varying in different tea

root cells; a similar situation was recently reported in *A. thaliana* leaves (Kim et al., 2021),

but has not yet been reported in the root systems of other plants. These findings deepen our

understanding of N metabolism and transport in plant roots.

Numerous studies have established that the processes underlying plant N sensing, uptake,

transport, and utilization are critical for the next novel green revolution (Liu et al., 2022b). In

this regard, high-throughput scRNA-seq studies should offer greater insights into the cell-type

specificity of many N-associated genes, which could help to select cell-type-specific genes

more efficiently for future designed breeding programs.

Compared with N metabolism in other plant species, theanine metabolism is a special

component of N metabolism in tea plants (Xu et al., 2012; Zhao et al., 2020). Tea plants prefer

NH_4^+ and synthesize large amounts of non-toxic theanine to decrease toxicity associated with

high NH_4^+ levels (Lin et al., 2022). In the theanine biosynthesis pathway, and consistent with

the findings of She et al. (She et al., 2022), our study established that *CsTSI* is expressed

primarily in the pericycle cells (Cluster 8). We further clarified that *CsAlaDC* is generally

expressed in the vasculature (Clusters 1). This differential enrichment of *CsTSI* and *CsAlaDC*

expression suggests that EA and theanine are synthesized in different cell types (Figure 5),

presumably reflecting multicellular compartmentation (Figure 8A).

It is well known that theanine is the most abundant amino acid in the tea plant, and that

theanine synthesis does not always occur, but is regulated to maintain growth and

development in balance or adapted to environmental conditions. Indeed, the observed

multicellular compartmentation may contribute to the regulation of theanine biosynthesis

through precursor syntheses and transport within different cells. Thus, we propose a model in

which multicellular compartmentation of theanine and its precursors functions like an innate

design for regulation of secondary metabolism (Figure 8A). Similar mechanisms may also

have evolved in other plants. We also cannot rule out that synthesis of theanine or EA can occur

in the same cell type; after all, *CsTSI* and *CsAlaDC* are two of the most highly expressed genes

in tea roots and their cell type-specific expression partially overlapped (Yang et al., 2020)

(Figure 4F).

On the other hand, multicellular compartmentation of theanine and its precursors EA implied

EA transport from its location of biosynthetic into theanine biosynthetic cells. EA is a

colorless gas, and hence it may be transported by way of simple diffusion, or via a gas channel.

Perhaps aquaporins (AQPs), a class of channel proteins that transport water, gas, and nutrients

(Hachez and Chaumont, 2010), may be involved. At present, the mechanism of EA transport

remains unknown, and the involvement of cell-type-specific AQPs would be worthy to explore

in the future.

Previously, we identified theanine transporters acting in theanine root-to-shoot transport

and theanine storage in the vacuole of tea root cells (Dong et al., 2020; Feng et al., 2021). In

the current study, we elucidated the cell heterogeneity of theanine transporters (Figure 5B and

8B), and among them *CsAAP1* is located primarily in cortical and endodermal cells (Figures 2E

and 5B), and *CsCAT2* is co-expressed with *CsTSI* in the same cell type. This nicely reflects that

CsAAP1 and CsCAT2 mediate in the physiological function of theanine transport and storage

(Dong et al., 2020; Feng et al., 2021). However, these theanine transporters are all importers

(Dong et al., 2020; Feng et al., 2021). The exporter(s) mediating in theanine transport from

the xylem parenchyma for loading into the xylem transpiration stream, within the roots, along

with other theanine export processes in the shoot, remain to be identified (Figure 8B).

Combining the cell type information, provided by scRNA-seq data, may assist in the

identification of these specific-cell-located theanine exporter(s), and may also provide

information for identifying amino acid exporters in other plant species. At present, all the

exporters identified are bidirectional transporters, such as UMAMITs (Besnard et al., 2016).

Therefore, specific-cell-located transporters, with bidirectional transport functions, could be

potential theanine exporters.

Interestingly, genes highly correlated with *CsTSI* or *CsAlaDC* were identified, including

some confirmed TFs (*CsMYB6*, *CsMYB40*, *CsHHO3*) (Guo et al., 2022; Zhang et al., 2021c),

which were not always highly co-expressed in the same cell type. The cell heterogeneity of

regulatory genes for *CsTSI* or *CsAlaDC* expression may reflect the "accelerator" and "brake"

regulatory patterns that might act to fine-regulate theanine biosynthesis in specific cells (Guo

et al., 2022). Another possibility is that these theanine-associated genes are promiscuous,

having many target genes and regulate multiple biological processes in tea plants. Thus, TFs

that regulate the theanine biosynthesis pathway may not always be co-express highly with

their target genes, in the same cell type. Indeed, this point was confirmed in a subsequent

study of CsLBD37 regulation of *CsAlaDC* expression.

To date, some identified TFs have been shown to play a role in the fine-regulation of

CsTSI and *CsAlaDC* expression; however, the TFs that impart *CsTSI* and *CsAlaDC*

root-specificity and high expression remain to be identified. Recently, it was established that,

in roots, hypomethylation in the promoters of *CsTSI* and *CsAlaDC* provided an epigenetic

basis for their high expression. Although hypomethylation in the promoters provides

accessibility for transcription, activating TFs is still required for their high expression. In this

regard, the current cell-type-specific expression database offers a valuable resource to identify

such unknown TF activators. For example, root-specific and Cluster 1-specific CsMYB18

could well be an activator of *CsAlaDC* expression (Figure 7B-D).

Recently, some TFs regulating *CsTSI* or *CsAlaDC* were identified and characterized and

indicted that theanine regulates lateral root development (Chen et al., 2022; Guo et al., 2022;

Zhang et al., 2021c); however, the molecular link between these processes was not elucidated.

Previous studies have shown that several LBD family members regulate proanthocyanidin

(PA) metabolism and are involved in root development in response to N in *Arabidopsis*

thaliana (Rubin et al., 2009). In tea roots, theanine, PA metabolism and root development are

also regulated by N (Wang et al., 2021). In addition, *CsLBD37* was also N-inducible and was

one of the most positively correlated TF encoding genes with *CsAlaDC* (Figure 6C). This

information provided important insights into the roles for *CsLBD37* in theanine biosynthesis

and root development. In this study, we identified CsLBD37 as a repressor of *CsAlaDC*

transcription and root development (Figure 7 and 8C; Supplemental Figure 8). CsLBD37 may

act in a similar manner to CsHHO3, namely through being a repressor of *CsAlaDC*

transcription to "brake" theanine biosynthesis under high N availability (Guo et al., 2022).

Here, we also revealed that *CsLBD37* probably inhibits lateral root development, at the root

initial stage (Supplemental Figure 8D-8G), but the underlying mechanism remains unknown.

CsLBD37 expression could be induced by exogenous theanine treatment in tea seedlings

(Supplemental Figure 8H). This implied that CsLBD37 may also be involved in a process by

which theanine regulates lateral root development. Is *CsLBD37* involved in regulating root

development through the auxin and ROS homeostasis pathways as well as theanine? This question is worthy of future investigation.

In summary, we generated a transcriptional map of tea roots, at the single cell level, which provided cell-type specific information on the expression of some 40,000 genes. Moreover, these datasets provide an important resource for future studies on secondary metabolism within tea roots, as well as other plant root systems. Recently, spatial transcriptomics and metabolomics techniques were applied for studies in both animals and plants (La Manno et al., 2021; Liu et al., 2022a). An integration of these techniques with scRNA-seq would afford a powerful approach to further explore cell specific secondary metabolism, at higher resolution, in various crop root systems.

Methods

Plant materials and growth conditions. Tea plants (*Camellia sinensis var. sinensis cv. Shuchazao*) seeds were soaked in water for 7 days until their seed coats burst: water was changed daily. Tea seeds were then cultured in vermiculite and watered (twice weekly) for 30 days, until radicle emergence. These tea seeds were then transplanted, singly, to new vermiculite, and watered (three times weekly) until the two-leaf stage, when they were transferred to nutrient solution for further growth; these hydroponic conditions and nutrient solution were as described previously (Konishi et al., 1985).

Preparation of root samples for scRNA-seq. The root tip regions (~2 cm in length from root tip) were cut, longitudinally, into 3 - 4 small strips and then processed for protoplast isolation. Here, the root strips were transferred immediately and gently into the prepared enzyme solution (0.1 mol·L^{-1} KCl, 0.08 mol·L^{-1} MES, 0.02 mol·L^{-1} CaCl$_2$, 1.5% cellulase R10 (Yakult, Japan), 1% Pectolyase (Yakult, Japan), 0.4 mol·l-1 mannitol, and 0.1% BSA), and digested at 25°C, with shaking (40 rpm), in the dark for the indicated period. An equal volume of washing buffer (WB) solution (0.1 mol·L^{-1} KCl, 0.08 mol·L^{-1} MES, 0.4 mol·L^{-1} mannitol and 0.1% BSA) was added in the enzyme solution and the mixture was then filtered using nylon mesh (200 mesh/inch). The flow-through was centrifuged at 200 x g for 2 min to pellet protoplasts. To evaluate protoplast yield, under different isolation conditions, protoplasts were re-suspended with suitable WB solution and counted with a hemocytometer.

Protoplast viability was determined by trypan blue staining, with the ratio of viable cells being

more than 85%. The concentration of protoplasts was adjusted to 700–1200 cells/μL and cells

were then processed with the 10x Genomics Single Cell Protocol (CG00052, RevC).

Construction and sequencing of the scRNA-seq library. Approx. 12,593 counted cells

were loaded onto a Single Cell Gchip. The libraries were constructed using $10\times$ Genomics

Chromium Next GEM Single Cell 3′ Reagent Kits v3.1(1000268). In brief, cell

suspensions in a chip were loaded on a Chromium Controller (10xGenomics, Pleasanton, CA)

to generate single-cell GEMs (gel beads in emulsion). The scRNA-seq libraries were then

prepared, according to the manufacturer's protocol. Qualitative analysis of the DNA library

was performed on an Agilent 2100 Bioanalyzer. The concentration of DNA library was

measured by Qubit (Invitrogen). Libraries were sequenced by an Illumina NovaSeq sequencer

(Genergy Biotechnology Shanghai) and the raw scRNA-seq dataset comprised Read1, Read2,

and i7 index read. The 26-bp read length of Read1 contained the sequence of the 16 bp

10xBarcode and 10 bp UMI (unique molecular identifiers). The 98-bp read length of Read2

was the sequence of the cDNA fragment. In total, 332,593,050 reads were obtained.

Pre-processing of raw scRNA-seq data. The raw files were analyzed by Cell Ranger 5.0.0

(10x Genomics). The tea plant genome and GTF annotation files, which excluded the organelle

genomes, were downloaded from the Tea plants Information Archive (TPIA) website

(http://tpia.teaplants.cn) (Xia et al., 2019). Running 'cellranger mkref' with ''–genome, –fasta

and –genes'' inputs was employed to build the reference. Then 'cellranger count' with ''–id, –

transcriptome, –fastqs, –sample" was run. Some 79.8% of the reads in all of the samples were

aligned to the tea plant genome by the aligner STAR (v.2.5.1b). Mean reads per cell were

26,411. Median genes per cell was in the range of 680.

Data integration, clustering, and annotation. Downstream analyses were mainly

performed with the Seurat package (v.3.1.2), as previously described (Butler et al., 2018). To

remove low quality cells and likely multiplet captures, which is a major concern in

microdroplet-based experiments, we applied a criteria to filter out cells with UMI/gene numbers

out of the limit of the mean value +/- 2-fold of the standard deviations, assuming a Guassian

distribution of the UMI/gene number for each cell. Additionally, we applied DoubletFinder

package (version 2.0.2) potential doublet identification (McGinnis et al., 2019).

Library size normalization was performed with the "NormalizeData" function (LogNormalize

method, scaling factor of 10,000). We then detected variable genes with the

"FindVariableFeatures" function (vst method, 4000 features), scaled data with "ScaleData"

function, performed PCA analysis with "RunPCA" function (14 principal components),

determined statistical significance of PCA scores by "JackStraw" function, constructed the

SNN graph, clustered cells based on Louvain ("FindNeighbors" and"FindClusters"), and

visualized data with non-linear dimensional reduction algorithms ("RunTSNE"). The

cluster-enriched genes were computed with the FindAllMarkers function in Seurat, using the

following parameters: a bimod test; above 1 – fold difference (logfc. Threshold $\geq$0) between

the two groups of cells; test genes that having a minimum fraction was at least 0.25.

Pseudo-time analysis. Monocle2 (V2.9.0) package was used to infer the trajectory of cell

differentiation (Trapnell et al., 2014), and the specific steps were as follows: First, the import

CDS function of Monocle2 package was used to transform the Seurat object into the Cell Data

Set object. 0.01), next the reduce Dimension function was used for dimensionality reduction

clustering, and finally, order Cells function was used to infer the differentiation trajectory.

***In situ* RT-PCR.** *In situ* RT-PCR was improved and performed based on previous methods

(Munns et al., 2012). The experimental material was the primary root of a 2-month-old

'shuchazao' tea cultivar. The young root tissue was soaked in fresh FAA solution,

vacuum-infiltrated for 15-20 min, then put at 4°C for 12 h. The samples were then eluted, three

times, for 10 min each, in a mixture solution (63% ethanol and 5% acetic acid), and twice for 5

min each in 1× PBS. Root tissue was then embedded in 5% low-melting point agarose (agarose

dissolved in 1× PBS) and the embedded blocks were sectioned using a Leica RM2255

microtome (Leica, Nussloch, Germany). The root tissue sections were separated from agarose

by rinsing, twice, with Rnase-Free water, 3 μg/ml proteinase K was then added for 30 min at

25°C. A heat treatment, at 85°C for 2 min, was given to inactivate proteinase K, and samples

were then washed, once, separately with 1×PBS and then Rnase-Free water for 5-10 min,

followed by the addition of 1 U/μl Dnase I, at 37°C, for 20 min, or overnight. Liquid was then

removed and 15 mM EDTA (pH 8.0) was added, at 75°C for 10 min, followed by 1-2 washes

with Rnase-Free water to remove excess liquid. Reverse transcription reactions were performed

using the PrimeScript II 1st strand cDNA Synthesis Kit (TAKARA, Cat. No. 9750), solution

was then removed, followed by a wash, once, with Rnase-Free water. Mixed tissue sections and

reaction reagents: 2 µl 10× Tag DNA polymerase (NEB, Cat. No. M0267V), 0.4 µL Buffer Tag

DNA polymerase (5 U/µL, NEB, Cat. No. B9004S), 1.6 µL dNTP (TAKARA, Cat. No. 4030),

0.32 µL DIG-11-dUTP (25 nmol, Sigma, Cat. No. 11093088910), 1 µL Forward Primer, 1 µL

Reverse Primer, 0.6 µL MgCl$_2$ (1.5 mM, Thermo Fisher, Cat. No. R0971) and 13.08 µL ddH$_2$O.

PCR program: 95°C for 30 s, 95°C for 30 s, 55°C for 50 s, 68°C for 45 s, 30-35 cycle, 68°C for

5 min 10°C. Washed twice with 1×PBS for 5 min each, and blocked with confining liquid (5%

skim milk) for 30 min. Alkaline phosphatase antibody was diluted in confining liquid at 1:500,

50 µL and was added and allowed to stand for 1 h. Wash twice for 15 min each time using 10×

Washing Buffer. Sample staining was performed using BM purple AP substrate, precipitating

for 30 min or 1 h, then washed twice with Rnase-Free water, and observed under a microscope

(Carl Zeiss, Gottingen, Germany) and photographed. All primers used for *in situ* PCR are listed

in Supplemental data 11.

Analysis of theanine and ethylamine content. Theanine and ethylamine were exacted

and detected as previously described, with slight modification (Yang et al., 2020; Cheng et al.,

2017). Theanine was analyzed by a Waters e2695 HPLC system equipped with 2489 UV/Vis

detector (Waters, U.S.A). Ethylamine was analyzed by GC-MS detecting (Agilent, Santa

Clara, U.S.A).

WGCNA analysis. The WGCNA package, in R software, was used in co-expression

analyses to detect relative relationships among genes (Langfelder and Horvath, 2008). A

weighted adjacency matrix was created, following unsupervised hierarchal clustering analysis

of genes, as follows: (1) removal of outliner genes and samples by the variation efficient and

hierarchical clustering analysis; (2) choosing of an applicable soft threshold (power = 12); (3)

identification of co-expressed gene modules using the function blockwise Modules; (4)

visualization of the co-expression network using Cytoscape3.9.0.

GO analysis. Tbtools (v1.098769) software was used to analyze the GO enrichment of

co-expression genes (Chen et al., 2020). GO annotation texts of tea plants were summarized

from TPIA database (http://tpia.teaplants.cn) (Xia et al., 2019).

Subcellular localization. Sequence information on *CsLBD37* was obtained from TPIA

(http://tpia.teaplants.cn) (Xia et al., 2019). The ORF of *CsLBD37* within the entry vector

pDONR207 was cloned into the destination binary vector, pK7WGF2.0, for subcellular localization

studies. The plasmid, pK7WGF2.0-*CsLBD37*-GFP was transformed into *Agrobacterium*

tumefaciens strain EHA105 to select for positive colonies for infiltration into *Nicotiana*

benthamiana. After 48h-72h post-infiltration, the GFP fluorescence was imaged with an Olympus

FV1000 confocal microscope (Olympus, Tokyo, Japan) to reveal the subcellular localization of

CsLBD37. GFP fluorescence signals were excited with a 488-nm laser, and the emitted light was

recorded from 500 to 530 nm to display the subcellular localization of *CsLBD37*.

Yeast one-hybrid assays (Y1H). We inserted *CsAlaDC* promoter fragments (24 approx. 2000

bp) into the pAbAi vector to generate the Pro*CsAlaDC*-pAbAi reporter construct, which was then

introduced into the yeast one-hybrid gold strain. The transformed cells were grown on synthetic

dropout medium (SD–Ura) (Coolaber, Beijing, China). The ORF of *CsLBD37* was inserted into

vector pGADT7 AD to generate AD-CsLBD37, which was then introduced into the yeast strain

with pro*CsAlaDC*-pAbAi. Transformants were grown on synthetic dropout medium (SD-Leu) with

200 ng ml^{-1} Aureobasidin A (AbA) (Coolaber, Beijing, China) for 3 days at 30°C. pGADT7 AD

empty vectors were used as controls.

Luciferase activity assay. Approx. 2 kb *CsAlaDC* promoter fragments were inserted into the

pGreenII 0800-LUC vector to generate the luciferase reporter construct, pro*CsAlaDC*-LUC. The

CsLBD37 ORF was inserted into the pGreenII 62-SK vector to generate the effector construct

35S-CsLBD37. Empty vectors were used as negative controls. The above-described constructs were

introduced into *A. tumefaciens* strain GV3101 with the pSoup plasmid, respectively. Then effector

and reporter cell suspensions were mixed, in a 5:1 (v/v) ratio, and infiltrated into 4-week-old *N.*

benthamiana leaves, followed by harvesting after 3 days. Sections of leaves were sprayed with 150

mg L^{-1} D-luciferin, potassium salt (Sciencelight, Shanghai, China) and LUC signals were captured

using a Tanon 5200 Multi charge-coupled device camera (Tanon, Shanghai, China). LUC and REN

activities of other *N benthamiana* leaves were measured using a dual-luciferase assay kit (Yeasen

Biotech, Shanghai, China) and a Spectra Max M2 (Molecular Devices, America) according to the

manufacturer's instructions.

Electrophoretic mobility shift assay (EMSA). The *CsLBD37* ORF was inserted into the

GST vector pGEX-4T-1 to generate GST-CsLBD37 construct, which was then transferred into

Escherichia coli BL21 cells. The positive BL21 strains were induced by 0.5 mM

isopropyl-beta-D-thiogalactopyranoside (IPTG), at 16°C for 20 h, and the GST-CsLBD37

recombinant proteins were purified by immobilized glutathione beads (TransGen Biotech, Beijing,

China). DNA probes were synthesized and labelled with biotin, at the 3' end, by using the EMSA

Probe Biotin Labeling Kit (Beyotime Biotech, Shanghai, China). The EMSA was performed using

the instructions of the Chemiluminescent EMSA Kit (Beyotime Biotech, Shanghai, China).

Antisense oligonucleotides assay. The antisense oligonucleotides assay was performed,

as previously described (Xie et al., 2014), to silence the target gene *CsLBD37*. AsOND was

designed and selected using the SOLIGO software

(https://sfold.wadsworth.org/cgi-bin/soligo.pl). The sequence of the sODN and AsODN

are listed in Supplemental data 11. To silence the genes, the primary roots of tea plant

cultivar 'Shuchazao' were wounded and then incubated in 10 ml 50 μM asODN-*CsLBD37*

solution for 6 h and 12 h. The solution with sense oligonucleotides (sOND) of the gene

serves as the control. After treatment, the roots were collected for RNA isolation and

metabolite detection.

Overexpression of *CsLBD37* in a tea plant hairy root system. The *CsLBD37* ORF

was inserted into pK7WGF2.0 driven by the *35S* promoter, using the BP and LR reaction

(Gateway system) to generate the pK7WGF2.0-CsLBD37 overexpression construct. The

resultant vector was transformed into Ar.A4 Chemically Competent Cell (Weidi Biotech,

Shanghai, China). The selected positive transformants were collected and resuspended in

1/2 Murashige and Skoog (MS) culture medium to an $OD_{600} = 0.6$ to transform 3-month-old

tea seedlings, as described previously (Alagarsamy et al., 2018).

Arabidopsis **transformation.** The Gateway system was employed to generate the

pB2GW7-CsLBD37 construct. *A. tumefaciens* strain GV3101, carrying pB2GW7-*CsLBD37*,

was used for the genetic complementation of *Arabidopsis* wild-type (Col-0) plants.

Transformants were selected on 0.25 ‰ Basta, and further verified by genomic PCR and

quantitative RT-PCR (qRT-PCR). The T3 progeny plants were used for these analyses. All

primers are listed in Supplemental data 11.

Theanine treatment of tea seedlings. Tea seedlings with a 5-cm primary root were

cultured in 1/2 nutrient solution for 1 week, and then treated with 1/2 nutrient solution

containing theanine for 2 weeks. The theanine concentration of the control group (CK)

was 0 mM and that for the treatment group (Theanine) was 1 mM.

RNA isolation and q-RT–PCR. RNA isolation was performed according to the manual of

RNAprep Pure Plant Plus Kit (TIANGEN, Beijing, China), and cDNA synthesis was

accomplished using the TransScript II One-Step gDNA Removal and cDNA Synthesis

SuperMix kit (TransGen Biotech, Beijing, China). A QuantStudio 6 Flex Real-Time PCR

System (Thermo, Singapore) was used to measure relative expression of genes, which were

calculated based on the $2^{-\Delta Ct}$ method (Schmittgen et al., 2000), using *CsGAPDH* as an internal

control. All primers used for qRT-PCR are listed in Supplemental data 11.

Reference

Alagarsamy, K., Shamala, L. F. & Wei, S. (2018). Protocol: high-efficiency in-planta

agrobacterium-mediated transgenic hairy root induction of *Camellia sinensis* var. *sinensis*.

Plant Methods. **14**:17.

Bai, P. et al. (2019). Identification of a novel gene encoding the specialized alanine

decarboxylase in tea (*Camellia sinensis*) plants. *Molecules* **24**:540.

Bailly, C. (2021). Anticancer properties of lobetyolin, an essential component of *Radix*

Codonopsis (Dangshen). *Nat. Prod. Bioprospect*. **11**:143-153.

Becht, E. et al. (2018). Dimensionality reduction for visualizing single-cell data using UMAP.

Nat. Biotechnol.

Bernhardt, C. et al. (2003). The bHLH genes *GLABRA3* (*GL3*) and *ENHANCER OF*

GLABRA3 (*EGL3*) specify epidermal cell fate in the Arabidopsis root. *Development*. **130**:

6431-6439.

Besnard, J. et al. (2016). UMAMIT14 is an amino acid exporter involved in phloem

unloading in Arabidopsis roots. *J. Exp. Bot.* **67**:6385-6397.

Bird, D. et al. (2003). A tale of three cell types: alkaloid biosynthesis is localized to sieve

elements in opium poppy. *Plant Cell.* **11**:2626-35.

Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. (2018). Integrating

single-cell transcriptomic data across different conditions, technologies, and species. *Nat.*

Biotechnol. **36**:411-420.

Chen, C. et al. (2020). TBtools: an integrative toolkit developed for interactive analyses of

big biological data. Mol. Plant **13**:1194-1202.

Chen, H. et al. (2021). PlantscRNAdb: a database for plant single-cell RNA analysis. Mol.

Plant **14**:855-857.

Chen, T. et al. (2022). Theanine, a tea plant specific non-proteinogenic amino acid, is

involved in the regulation of lateral root development in response to nitrogen status. Hortic.

Res. **10**:uhac267.

Cheng, S. et al. (2017). Studies on the biochemical formation pathway of the amino acid

l-theanine in tea (*Camellia sinensis*) and other plants. J. Agric. Food Chem. **65**:7210-7216.

De Rybel, B. et al. (2013). A bHLH complex controls embryonic vascular tissue

establishment and indeterminate growth in *Arabidopsis*. Dev. Cell **24**:426-437.

Denyer, T. et al. (2019). Spatiotemporal developmental trajectories in the *Arabidopsis* root

revealed using high-throughput single-cell RNA sequencing. Dev. Cell **48**:840-852.e845.

Dinant, S. et al. (2003). Diversity of the superfamily of phloem lectins (phloem protein 2) in

angiosperms. Plant Physiol. **131**: 114-128.

Dong, C. et al. (2020). Theanine transporters identified in tea plants (*Camellia sinensis* L.).

Plant J. **101**:57-70.

Dong, W. et al. (2021). An SHR–SCR module specifies legume cortical cell fate to enable

nodulation. Nature **589**:586-590.

Du, J. et al. (2021). Gene coexpression network reveals insights into the origin and evolution

of a theanine-associated regulatory module in Non-*Camellia* and *Camellia* Species. J. Agric.

Food Chem. **69**:615-626.

Efroni, I. et al. (2016). Root regeneration triggers an embryo-like sequence guided by

hormonal interactions. Cell **165**:1721-1733.

Feng, L. et al. (2021). Tonoplast-localized theanine transporter CsCAT2 may mediate

theanine storage in the root of tea plants (*Camellia sinensis* L.). Front. Plant Sci.

12:797854.

Fernie, A. R. & Tohge, T. (2017). The genetics of plant metabolism. Annu. Rev. Genet.

51:287-310.

Galili, G., Amir, R. & Fernie, A. R. (2016). The regulation of essential amino acid synthesis

and accumulation in plants. Annu. Rev. Plant Biol. **67**:153-178.

Garcia-Lemos, A. M. et al. (2020). Identification of root-associated bacteria that influence

plant physiology, increase seed germination, or promote growth of the christmas tree

species *Abies nordmanniana*. Front. Microbiol. **11**:566613.

Goh, T., Joi, S., Mimura, T. & Fukaki, H. (2012). The establishment of asymmetry in

Arabidopsis lateral root founder cells is regulated by LBD16/ASL18 and related LBD/ASL

proteins. Development **139**:883-893.

Guo, J. et al. (2022). Potential "accelerator" and "brake" regulation of theanine biosynthesis

in tea plant (*Camellia sinensis*). Hortic. Res. **9**:uhac169.

Hachez, C. & Chaumont, F. (2010). Aquaporins: a family of highly regulated

multifunctional channels. Adv. Exp. Med. Biol. **679**:1-17.

Haghverdi, L., Buettner, F. & Theis, F. J. (2015). Diffusion maps for high-dimensional

single-cell analysis of differentiation data. Bioinformatics **31**:2989-2998.

Haghverdi, L., Büttner, M., Wolf, F. A., Buettner, F. & Theis, F. J. (2016). Diffusion

pseudotime robustly reconstructs lineage branching. Nat. Methods **13**: 845-848.

Hartman, K., van der Heijden, M. G., Roussely-Provent, V., Walser, J. C. & Schlaeppi,

K. (2017). Deciphering composition and function of the root microbiome of a legume plant.

Microbiome **5**:2.

Hong, J. H. et al. (2017). A sacrifice-for-survival mechanism protects root stem cell niche

from chilling stress. Cell **170**:102-113.e114.

Hu, D. J., Shakerian, F., Zhao, J. & Li, S. P. (2019). Chemistry, pharmacology and analysis

of *Pseudostellaria heterophylla*: a mini-review. Chin. Med. **14**: 21.

Kim, J. Y. et al. (2021). Distinct identities of leaf phloem cells revealed by single cell

transcriptomics. Plant Cell **33**:511-530.

Kong, X. et al. (2015). WOX5 is shining in the root stem cell niche. Trends Plant Sci.
20:601-603.

Konishi, S. et al. (1985). Stimulatory effects of aluminum on tea plants grown under low and
high phosphorus supply. *Soil Sci. Plant Nutr.* **31**:361–368

La Manno, G. et al. (2021). Molecular architecture of the developing mouse brain. Nature
596:92-96.

Langfelder, P. & Horvath, S. (2008). WGCNA: an R package for weighted correlation
network analysis. BMC. Bioinformatics **9**:559.

Li, H. et al. (2021). Single-cell RNA sequencing reveals a high-resolution cell atlas of xylem
in populus. J. Integr. Plant Biol. **63**:1906-1921.

Lin, S. et al. (2022). Theanine metabolism and transport in tea plants (*Camellia sinensis* L.):
advances and perspectives. Crit. Rev. Biotechnol. 1-15.

Liu, C. et al. (2022). A spatiotemporal atlas of organogenesis in the development of orchid
flowers. Nucleic Acids Res. **50**:9724-9737.

Liu, Q. et al. (2022). Improving crop nitrogen use efficiency toward sustainable green
revolution. Annu. Rev. Plant Biol. **73**:523-551.

Liu, W., Xu, L., Lin, H. & Cao, J. (2021). Two expansin genes, *AtEXPA4* and *AtEXPB5*, are
redundantly required for pollen tube growth and *AtEXPA4* is involved in primary root
elongation in *Arabidopsis thaliana*. Genes (Basel). **12**:249.

McGinnis, C. S., Murrow, L. M. & Gartner, Z. J. (2019). DoubletFinder: doublet detection
in single-cell RNA sequencing data using artificial nearest neighbors. Cell Syst. **8**:329-337.

Munns, R. et al. (2012). Wheat grain yield on saline soils is improved by an ancestral Na^+
transporter gene. Nat. Biotechnol. **30**:360-364.

Nelms, B. & Walbot, V. (2019). Defining the developmental program leading to meiosis in
maize. Science **364**:52-56.

Okushima, Y., Fukaki, H., Onoda, M., Theologis, A. & Tasaka, M. (2007). ARF7 and
ARF19 regulate lateral root formation via direct activation of LBD/ASL genes in
Arabidopsis. Plant Cell **19**:118-130.

Passarinho, P. A., Van Hengel, A. J., Fransz, P. F. & de Vries, S. C. (2001). Expression

pattern of the *Arabidopsis thaliana AtEP3/AtchitIV* endochitinase gene. Planta

212:556-567.

Péret, B. et al. (2009). *Arabidopsis* lateral root development: an emerging story. Trends Plant

Sci. **14**:399-408.

Rubin, G., Tohge, T., Matsuda, F., Saito, K. & Scheible, W. R. (2009). Members of the

LBD family of transcription factors repress anthocyanin synthesis and affect additional

nitrogen responses in *Arabidopsis*. Plant Cell **21**:3567-3584.

Ryu, K. H., Huang, L., Kang, H. M. & Schiefelbein, J. (2019). Single-cell RNA sequencing

resolves molecular relationships among individual plant cells. Plant Physiol.

179:1444-1456.

Santelia, D. et al. (2008). Flavonoids redirect PIN-mediated polar auxin fluxes during root

gravitropic responses. J. Biol. Chem. **283**:31218-31226.

Schmid, C., Dawid, C., Peters, V. & Hofmann, T. (2018). Saponins from european licorice

roots (*Glycyrrhiza glabra*). J. Nat. Prod. **81**:1734-1744.

Schmittgen, T. D. et al. (2000). Quantitative reverse transcription-polymerase chain reaction

to study mRNA decay: comparison of endpoint and real-time methods. Anal. Biochem.

285:194-204.

Schulz-Bohm, K. et al. (2018). Calling from distance: attraction of soil bacteria by plant root

volatiles. Isme J. **12**:1252-1262.

Shaw, R., Tian, X. & Xu, J. (2021). Single-cell transcriptome analysis in plants: advances

and challenges. Mol. Plant **14**:115-126.

She, G. et al. (2022). Characterization of *CsTSI* in the biosynthesis of theanine in tea plants

(*Camellia sinensis*). J. Agric. Food Chem. **70**:826-836.

Silva, L. S., Seabra, A. R., Leitão, J. N. & Carvalho, H. G. (2015). Possible role of

glutamine synthetase of the prokaryotic type (GSI-like) in nitrogen signaling in *Medicago*

truncatula. Plant Sci. **240**:98-108.

Sláviková, S. et al. (2005). The autophagy-associated Atg8 gene family operates both under

favourable growth conditions and under starvation stresses in *Arabidopsis* plants. J. Exp.

Bot. **56**:2839-2849.

Steppuhn, A., Gase, K., Krock, B., Halitschke, R. & Baldwin, I. T. (2004). Nicotine's

defensive function in nature. PLoS Biol. **2**:E217.

Trapnell, C. et al. (2014). The dynamics and regulators of cell fate decisions are revealed by

pseudotemporal ordering of single cells. Nat. Biotechnol. **32**:381-386.

Wan, J. et al. (2018). UV-B radiation induces root bending through the flavonoid-mediated

auxin pathway in *Arabidopsis*. Front. Plant Sci. **9**:618.

Wang, Q. et al. (2022). Single-cell transcriptome atlas reveals developmental trajectories and

a novel metabolic pathway of catechin esters in tea leaves. Plant Biotechnol. J.

20:2089-2106.

Wang, W. et al. (2018). Expression of the nitrate transporter gene *OsNRT1.1A/OsNPF6.3*

confers high yield and early maturation in rice. Plant Cell **30**:638-651.

Wang, Y. et al. (2021). Nitrogen-regulated theanine and flavonoid biosynthesis in tea plant

roots: protein-level regulation revealed by multiomics analyses. J. Agric. Food Chem.

69:10002-10016.

Wang, Y. Y., Cheng, Y. H., Chen, K. E. & Tsay, Y. F. (2018). Nitrate transport, signaling,

and use efficiency. Annu. Rev. Plant. Biol. **69**:85-122.

Wei, C. et al. (2018). Draft genome sequence of *Camellia sinensis var. sinensis* provides

insights into the evolution of the tea genome and tea quality. Proc. Natl. Acad. Sci. U. S. A.

115:E4151-e4158.

Xia, E. H. et al. (2019). Tea plant information archive: a comprehensive genomics and

bioinformatics platform for tea plant. Plant Biotechnol. J. **17**:1938-1953.

Xie, Z. et al. (2014). A selection strategy in plant transformation based on antisense

oligodeoxynucleotide inhibition. Plant J. **77**:954-961.

Xu, G., Fan, X. & Miller, A. J. (2012). Plant nitrogen assimilation and use efficiency. Annu.

Rev. Plant Biol. **63**:153-182.

Xu, S. et al. (2017). Wild tobacco genomes reveal the evolution of nicotine biosynthesis. Proc.

Natl. Acad. Sci. U. S. A. **114**:6133-6138.

Xu, Z. et al. (2022). Plant single cell transcriptome hub (PsctH): an integrated online tool to

explore the plant single-cell transcriptome landscape. Plant Biotechnol. J. **20**:10-12.

Yang, T. et al. (2020). Transcriptional regulation of amino acid metabolism in response to

nitrogen deficiency and nitrogen forms in tea plant root (*Camellia sinensis* L.). Sci. Rep.

10:6868.

Ye, L. et al. (2021). Cytokinins initiate secondary growth in the *Arabidopsis* root through a
set of LBD genes. Curr. Biol. **31**:3365-3373.e3367.

Yu, X. et al. (2020). Metabolite signatures of diverse *Camellia sinensis* tea populations. Nat.
Commun. **11**:5586.

Zhang, F. et al. (2020). Identification of regulatory networks and hub genes controlling
nitrogen uptake in tea plants [*Camellia sinensis* (L.) O. Kuntze]. J. Agric. Food Chem.
68:2445-2456.

Zhang, F. et al. (2021). Transcriptome analysis identifies CsNRT genes involved in nitrogen
uptake in tea plants, with a major role of CsNRT2.4. Plant Physiol. Biochem. **167**:970-979.

Zhang, T. et al. (2021). Single-cell transcriptome atlas and chromatin accessibility landscape
reveal differentiation trajectories in the rice root. Nat. Commun. **12**:2053.

Zhang, Y. et al. (2021). Molecular basis of the distinct metabolic features in shoot tips and
roots of tea plants (*Camellia sinensis*): characterization of MYB regulator for root theanine
synthesis. J. Agric. Food Chem. **69**:3415-3429.

Zhao, J., Li, P., Xia, T. & Wan, X. (2020). Exploring plant metabolic genomics: chemical
diversity, metabolic complexity in the biosynthesis and transport of specialized metabolites
with the tea plant as a model. Crit. Rev. Biotechnol. **40**:667-688.

Zhu, B. et al. (2021). CsAlaDC and CsTSI work coordinately to determine theanine
biosynthesis in tea plants (*Camellia sinensis* L.) and confer high levels of theanine
accumulation in a non-tea plant. Plant Biotechnol. J. **19**:2395-2397.

Zhu, Y. et al. (2006). *Arabidopsis* NRP1 and NRP2 encode histone chaperones and are
required for maintaining postembryonic root growth. Plant Cell **18**:2879-2892.

Acknowledgements

This work was supported by the National Key R&D Program of China (2022YFF1003103,
2021YFD1601101), and grants from the National Natural Science Foundation of China
(32072624), Anhui Provincial Major Science and Technology Project (202103b06020024)
and Anhui Educational Committee Excellent Youth Talent Support project
(gxyqZD2022018). We want to thank Oebiotech company provide 10× scRNA-seq service.

Author contributions

ZZ and WX conceived the research and finalized the manuscript. SL, YZ, SZ, YW, MH, YD, JG and BZ collected materials and carried out the experiments. SL and YZ analyzed the data. TY and EX offered guidance on experimental techniques and bioinformatics analysis. SL and ZZ and wrote the manuscript. LW offered suggestions and revisions to the writing of this manuscript. All authors read and approved the manuscript.

Competing interests

The authors declare no competing interests.

Figures and legends

Figure 1 Process employed for scRNA-seq of tea plant roots. (A) Tea seedlings were used for theanine detection and scRNA-seq. Tea plant roots (~2.0 cm in length from the root tip) were harvested and used for scRNA-seq. **(B)** Theanine contents of leaf, stem, cotyledon, and root. **(C)** Cross section of tea roots (from the root tip back ~2.0 cm). Co, Cortex; Xy, Xylem; Ph, Phloem; Ca, Cambium; Pe, Pericycle; En, Endodermis; EP, Epidermis. Scale bar = 200 μm. **(D)** Flow chart of scRNA-seq analyses.

Figure 2 Annotation of cell clusters based on scRNA-seq analysis of tea plant roots. (A) *t*-SNE visualization plot of 10,435 tea root cells showed that they were grouped into 8 cell clusters. Each dot denoted a single cell. **(B)** *t*-SNE visualization of cell-type marker genes of root cells. Color bar indicates gene expression level. **(C)** *In situ* RT-PCR of *CsWAT1*, *CsPP2*, *CsAAP1* and *CsLHW*. The blue areas of sliced tissue represent regions where genes are expressed. The red box represents the magnified areas of root sections. Scale bar = 200 μm. **(D)** Violin plot showed the expression patterns for the top 5 marker genes of each cell clusters. The height of violin represents the gene expression level, and the width represents the proportion of cells expressing in the cluster. Asterisks represent homologous *Arabidopsis thaliana* marker genes that can be found in PlantscRNAdb. **(E)** Schematic of the tea root anatomy, cell types and their associated clusters.

Figure 3 Trajectory of root cell differentiation in the tea root. (A) Schematic representation

of root meristem organization (in median longitudinal section). *t*-SNE visualization and cell

type annotation of Cluster 3 (bottom). **(B)** Simulation of the successive differentiation

trajectory of the stem cell niche (SCN) over pseudo-time. Each dot denotes a unique cell.

Pseudo-time analysis of all 10,435 tea root cells (center), and the detailed distribution of 8 cell

clusters along pseudo-time trajectory (outer). Colors of the dots represent the pseudo-time

score. Red and blue line mark major differentiation trajectories, black lines mark minor

differentiation trajectories. **(C)** Heatmap illustrating the expression patterns of differential

genes, along the pseudo-time trajectory, during cell differentiation of the vasculature and

epidermis. Each row represents one gene. Color bars indicate the relative expression levels.

Biological processes are given on the right. **(D)** Proposed tea roots cell differentiation

952   trajectories for cell clusters shown on the t-SNE visualization plots. Dotted arrows represent

953   assumed cell differentiation routes.

Figure 4 Cell heterogeneity of nitrogen (N) transport and assimilation in tea plant roots.

(A) Schematic of N transport and metabolism pathway in tea roots. Membrane-located transporter NRTs and AMTs uptake nitrate and ammonium into root cells, respectively, and subsequently NO_3^- is assimilated into NH_4^+ and then metabolized into various amino acids. **(B)** Heatmap shows cell cluster expression patterns of N transport and assimilation genes in tea roots. Row normalization for gene expression according to "normalized" mode. **(C)** *In situ* RT-PCR of *CsAMT1.1*, *CsAMT3.1*, *CsNRT1.1* and *CsNRT3.2*. The blue areas of sectioned

tissues represent regions where genes are expressed. The red boxes represent the magnified

areas of root sections shown on the right. Scale bar = 200 μm. **(D)** Expression of amino acid

synthesis pathway genes in different cell clusters. Color bars indicate the expression levels of

amino acids synthesis pathway genes. Dot size indicates the percentage of amino acid synthesis

genes expressed in a cell cluster.

Figure 5 Cell heterogeneity of theanine synthesis, transport, and regulation in tea plant

roots. (A) Model of the putative theanine synthesis, transport, and regulation in tea plant root

cells. Black full lines represent unique metabolic processes in tea plants; grey full lines

represent common metabolic pathways in plants; dotted lines indicate the multi-step process.

(B) Heatmap for cell cluster expression patterns of genes encoding theanine transporters

CsAAPs, key enzymes and transcription factors in the theanine metabolic pathway. Gene

expression is presented in "normalized" mode. **(C)** *t*-SNE visualization of *CsTSI, CsAlaDC* and

their overlap map. Blue color bar indicates *CsTSI* expression level, red color bar indicates

CsAlaDC expression level in scRNA-seq. **(D)**Tissue localization of *CsTSI* and *CsAlaDC* using

in situ PCR. Blue signal indicates gene expression in cells. The red boxes represent magnified

areas of root sections shown below. Scale bar = 200 μm.

Figure 6 A gene co-expression network predicted key regulators of *CsAlaDC*. (A) Using

bulk RNA-seq data, a gene co-expression network was generated for *CsAlaDC* (|correlation

coefficient| ≥ 0.85 and *p*-Value ≤ 0.05). Red and blue dots represent differential genes

positively and negatively correlated with *CsAlaDC*, respectively. Bulk RNA-seq data is from

different types of N treatment of tea roots (Yang et al., 2019). **(B)** Heatmap showing expression

of co-expressed genes with *CsAlaDC* in eight tea plant tissues. The black arrows denote root

specific co-expressed (RSCE) genes. **(C)** Heatmap showing the expression pattern for RSCE

genes under different types of N treatment (0 N, CK, EA, NH_4^+ and NO_3^-). Correlation between

RSCE genes and *CsAlaDC* is shown on the heatmap at left. **(D)** Heatmap showing the

expression level of RSCE genes in each cell cluster. **(B-D)** Row normalization of gene

expression is according to the "normalized" mode.

Figure 7 Theanine synthesis regulated by the transcription factor CsLBD37. (**A**) Schematic representation of the tea plant tissues used for gene detection. (**B**) Relative expression of *CsLBD37* in different tea plants tissues. (**C**) Cell specificity of *CsLBD37* expression in tea plant roots; blue signal represents location of gene expression. Scale bar = 200 μm. (**D**) The *t*-SNE visualization graph shows expression levels of *CsLBD37* in various cell clusters. (**E**) Subcellular localization of CsLBD37; green fluorescence signal represents

protein localization. **(F)** The binding of CsLBD37 to the CsAlaDC promoter, tested by yeast

one-hybrid assay (Y1H). **(G)** EMSA assay showing the association of CsLBD37 with the

CsAlaDC promoter. The red arrow points to the binding position. **(H)** A schematic of the

effector and reportor constructs for transcriptional activity of the *CsAlaDC* promoter (top). LCI

assays in tobacco leaves show that the *CsAlaDC* promoter can interact with CsLBD37. The

color bar indicates the range of luminescence intensity (mid). The LUC/REN values represent

relative activity of CsLBD37 on *CsAlaDC* expression. Significant difference was evaluated by

two-tailed Student's *t*-test analysis (bottom). **(I-L)** *CsLBD37* gene silencing and overexpression

in tea plant roots. The expression level of *CsLBD37* and *CsAlaDC* in gene silenced **(I)** and

overexpression plants **(K).** Ethylamine and theanine contents in gene silenced **(J)** and

overexpression plants. **(L).** sODN, sense oligo nucleotides; asODN, antisense

oligonucleotides, silenced *CsLBD37* in tea root by asODN methods; EV, empty vector;

CsLBD37-OE/*Cs*, *CsLBD37* overexpression in transgenic hairy root system. Significant

difference was determined by Welch's t test.

Figure 8 Proposed model for theanine biosynthesis, transport and regulation at single

cell resolution. (A) Multicellular compartmentation of theanine biosynthesis. The high

expression of *CsTSI* in pericycle cells suggests theanine is mainly synthesized in pericycle

cells. *CsAlaDC* is highly expressed in cambium and protoxylem cells, suggesting that

ethylamine (EA) is mainly synthesized in these cells. Thus, EA may move from cambium and

protoxylem cells into pericycle cells, where it is a substrate for theanine biosynthesis. **(B)** In

tea plant roots, CsAAP1 likely mediates in theanine retrieval from apoplast for its transport

into cortical/endodermal cells. CsCAT2 imports theanine into vacuoles, especially in

pericycle cells, which are the main cells for theanine biosynthesis. Theanine is exported, by

an unidentified theanine exporter, from pericycle cells into the apoplastic pathway for entry

into the xylem. **(C)** CsLBD37 co-regulates theanine biosynthesis and lateral root development.

Increasing levels of nitrogen (N) promote theanine biosynthesis and accumulation. High

accumulation of theanine and N induces *CsLBD37* expression. CsLBD37 inhibits *CsAlaDC*

expression to reduce ethylamine synthesis, which finetunes theanine biosynthesis in feedback

loop. Besides, *CsLBD37* is expressed in pericycle cells and also inhibits lateral root

development. It is reported that apoplastic theanine can negatively regulate lateral root

development, under high nitrogen levels (Chen et al., 2022). Therefore, CsLBD37 may also be

involved in theanine-regulated lateral root development.

Supplemental Figure 1. Summary of the tea root scRNA-seq data. (A) Filter delocalized

cell maps fitted to generalized linear models. The horizontal axis is the unique molecular

identifiers (UMI) number in each cell, and the vertical axis is the gene number in each cell.

According to the linear relationship between the UMI and genes per cell data, the distribution

model is fitted. The yellow points represent delocalized cells, which will be removed in the

follow-up analysis. **(B)** The cell number in each cell cluster based on the tea root scRNA-seq

data.

Supplemental Figure 2. Cell cluster annotation. (A) *t*-SNE visualization of cell-type marker

genes (*CsGL3, CsSCR, CsIRX9 and CsWOX5*) of root cells. Color bar indicated gene

expression level. **(B)** Violin plot showing expression patterns of 6^{th} to 10^{th} in the top10

differential marker genes of each cell cluster. The height and width of the violin represent the

gene expression level and the proportion of cells expressing in the cluster, respectively.

Asterisks represent homologous marker genes that are present in PlantscRNAdb. **(C)** The

tissue-located heatmap of *LAC7, PME41, NRP1* and *TIM17*. The color bar illustrated gene

expression level for the scRNA-seq data. The *Arabidopsis thaliana* data were generated from

the Root Cell Atlas (https://rootcellatlas.org/).

Supplemental Figure 3. *t*-SNE visualization of Cluster 1 and Cluster 2 top 10 marker

genes. *t*-SNE plots show the transcript accumulation of the Cluster 1 and Cluster 2 top 10

marker genes in individual cells. Color intensity indicates the relative transcript level for the

indicated gene in each cell.

Supplemental Figure 4. *t*-SNE visualization of Cluster 3 and Cluster 4 top 10 marker

genes. *t*-SNE plots shows the transcript accumulation of the Cluster 3 and Cluster 4 top 10

marker genes in individual cells. Color intensity indicates the relative transcript level for the

indicated gene in each cell.

Supplemental Figure 5. *t*-SNE visualization of Cluster 5 and Cluster 6 top 10 marker

genes. *t*-SNE plots shows the transcript accumulation of the Cluster 5 and Cluster 6 top 10

marker genes in individual cells. Color intensity indicates the relative transcript level for the

indicated gene in each cell.

Supplemental Figure 6. *t*-SNE visualization of Cluster 7 and Cluster 8 top 10 marker genes. *t*-SNE plots shows the transcript accumulation of the Cluster 7 and Cluster 8 top 10 marker genes in individual cells. Color intensity indicates the relative transcript level for the indicated gene in each cell.

Supplemental Figure 7. Transcriptional activation assay and protein purification of *CsLBD37*. Protein purification of *CsLBD37*. The CsLBD37-GST fusion protein has a molecular weight of 50.9 kDa (i.e., 24.9 kDa + 26 kDa).

Supplemental Figure 8. Overexpression of *CsLBD37* **in** *Arabidopsis* **inhibited lateral root development.** **(A)** qRT-PCR assay for *CsLBD37* transcriptional level in three transgenic *Arabidopsis* lines (*CsLBD37*-OE1*/At*, *CsLBD37*-OE2*/At*, *CsLBD37*-OE3*/At*) and wild-type *Arabidopsis* (WT) plants. **(B and C)** The plant phenotypes of *CsLBD37* transgenic *Arabidopsis* lines and WT grown in MS medium for 12 days. Scale bar = 1 cm. **(C)** Lateral root number, each line contained 25 independent biological replicates. **(D-G)** Microscopy-based lateral root developmental phenotypes of *CsLBD37* transgenic *Arabidopsis* lines and WT, 5 days post germination. Scale bar = 25 μm in **(D)**. Lateral root initiation number **(E)**; lateral root primordium number **(F)** and lateral root emergence number **(G)**, each line contained 20 independent biological replicates. **(H)** qRT-PCR assay for *CsLBD37* expression level in tea seedlings grown under control (CK) or 1 mM theanine treatment (Theanine). **(C and E-H)** Significant differences were determined by two-sided Student's *t*-test.

A

Leaf

Stem

Cotyledon

2 cm

Root

B

Theanine

Content (mg/g DW)

Leaf   Stem   Cotyledon   Root

C   Cell types of tea root

Xy
Ph
Ca
Pe
En
Co
Ep
Root hair

D

Root sample cellection → Protoplast → 10 X GEM partitioning → Sequencing and data analysis

Single cell RNA-seq

Buffer   Oil

Cell types   Gene expression (Different cells)

Bulk RNA-seq dataset of five forms N treatment ($0\ N$, NH_4^+ and NO_3^-, EA, NH_4^+, NO_3^-)

Gene co-expression network

Joint analysis

Functional identified

A

QC Stem cells Vasculature

Stem cell niche   Inner cell layer

Middle cell layer   Outer cell layer

Cluster 3

C3.1   C3.2

C3.3

C3.4

Quiescent center (QC): Cluster 3.1

Stele initials (SI): Cluster 3.2

Co/En initials (CEI): Cluster 3.3

Col initials (CI): Cluster 3.4

Ep/Lateral root cap initials
(ELI): Cluster 3.4

B

Cluster 2
(Ep)

Cluster 5
(RC)

Outer cell layer

Pseudotime

0 5 10 15 20

Cluster 7
(Ph)

Cluster 3
(SCN)

Start

Inner cell layer

Cluster 1
(Xy)

Cluster 6
(Ca)

Cluster 8
(Pe)

Cluster 4
(Co/En)

Middle cell layer

C

Vasculature psudotime

Biological process

Plant-type cell wall biogenesis

Lignin catabolic process

Amino sugar catabolic process

Amino acid import

Phenylpropanoid metabolic process

Start   Cluster 1

Epidermis pseudotime

Trichoblast differentiation

Root hair elongation

Plant epidermal cell differentiation

Root epidermal cell differentiation

Response to salt stress

Start   Cluster 2

Expression

3
2
1
0
-1
-2
-3

D

SI

QC

CEI

Xy

Co/En

Pe

ELI
CI

Ep

RC

Ph

A

Cell membrane
Cytoplasm

Amino acid transporter

Plastid

NH₄⁺

CsGS   α-KG

Gln   CsGDH

CsGOGAT   Glu

NADPH

Glu

Thea   ?

CsCAT2

Thea   Thea

Vacuole

Glu   Pyruvate ← Glycolysis

CsALT

Ala

CsAlaDC

EA   ┈ Translation

CsTSI

Thea

Nucleus

CsMYB40

CsHHO3

DNA

mRNA

CsAlaDC

CsMYB6

DNA

mRNA

CsTSI

Cell membrane

CsAAPs   ?

Thea

B

Theanine synthesis

CsTSI
CsAlaDC
CsGS1.1
CsGS1.2
CsGS1.3
CsGS2
CsGOGAT1
CsGOGAT2
CsGOGAT3
CsGDH2
CsGDH3
CsGDH4
CsALT

Expression
(Row normalization)

High

Low

Theanine regulation

CsMYB40
CsHHO3
CsMYB6

Theanine transport

CsAAP1
CsAAP2
CsAAP5
CsAAP6
CsAAP8
CsCAT2

1 2 3 4 5 6 7 8
Cluster

C

CsTSI   *CsAlaDC*   Overlap

Expression
Low *CsTSI* High

Expression
Low *CsAlaDC* High

D

CsTSI   *CsAlaDC*   Negative control

A

Bud | 1st Leaf | 3rd Leaf | Stem | Root

B

CsLBD37

Relative expression

Bud | 1st Leaf | 3rd Leaf | Stem | Root

C

CsLBD37 | Negative control

D

CsLBD37

Expression — High / Low

Cambium | Pericycle

Expression

Cluster: 1 2 3 4 5 6 7 8

E

EGFP | DAPI | Bright field | Merge

35S: eGFP

CsLBD37: eGFP

F

ProCsAlaDC-p*AbAi*

10^0 10^{-1} 10^{-2} 10^{-3}

AD

AD-*CsLBD37*

SD-Leu

ProCsAlaDC-p*AbAi*

10^0 10^{-1} 10^{-2} 10^{-3}

AD

AD-*CsLBD37*

SD-Leu+AbA

G

ProCsAlaDC

Biotin probe	+	+	+	+
GST	+	-	-	-
CsLBD37-GST	-	+	+	+
Cold probe	-	-	5×	10×

Bound

Free

Cold probe : ACATTGACAATTACG
TGACATAAAAATATTTTATTTTATA

H

Reportor:

LUC empty   35S — REN — Ter — LUC — Ter

ProCsAlaDC: LUC   35S — REN — Ter — *ProCsAlaDC* — LUC — Ter

Effector:

SK empty   35S — Ter

35S: *CsLBD37*   35S — *CsLBD37* — Ter

High / Low

| *ProCsAlaDC*: LUC + 35S: *CsLBD37* | *ProCsAlaDC*: LUC + SK empty |
| LUC empty + 35S: *CsLBD37* | LUC empty + SK empty |

ProCsAlaDC: LUC + SK empty

ProCsAlaDC: LUC + 35S: *CsLBD37*

P=0.0003

Relative activity (LUC/REN)

I

CsLBD37 | *CsAlaDC*

Gene expression

P=0.0090   *P*=0.0081

sODN asODN   sODN asODN

J

Ethylamine | Theanine

Content (mg/g DW)

P=0.0451   *P*=0.0219

sODN asODN   sODN asODN

K

CsLBD37 | *CsAlaDC*

Gene expression

P=0.0375   *P*=0.0256

EV *CsLBD37-OE/Cs*   EV *CsLBD37-OE/Cs*

L

Ethylamine | Theanine

Content (mg/g DW)

P=0.0451   *P*=0.0001

EV *CsLBD37-OE/Cs*   EV *CsLBD37-OE/Cs*

A

Epidermis
Cortex
Endodermis
Pericycle
Cambium
Protoxylem

Expression

CsTSI
CsAlaDC

Cell type

Pericycle
Cambium and protoxylem

Ala
CsAlaDC
Glu
EA
CsTSI
Thea
Thea
Thea
Thea
Thea
Thea

Ala
CsAlaDC
EA
EA
EA
EA
Glu
CsTSI
Thea

PD

Glu, Glutamic acid
Thea, Theanine

Ala, Alanine
PD, Plasmodesmata

EA, Ethylamine

B

Epidermis   Cortex   Endodermis   Pericycle   Xylem

VC

Theanine

Casparian
strip

Theanine
importer

CsAAP1

CsCAT2

Theanine
exporter
(unidentified)

Theanine
translocation

VC, Vacuole

C

Epidermis
Cortex
Endodermis
Pericycle

Lateral root primordium

Lateral root development

Nitrogen level

Low   High

Theanine

Low   High

CsLBD37

CsAlaDC

Ala   CsAlaDC   Ethylamine

Cluster 7 top 10 marker gene

Cluster 8 top 10 marker gene