1 ORIGINAL ARTICLE

4 CXCR3-expressing myeloid cells recruited to the hypothalamus protect
5 against diet-induced body mass gain and metabolic dysfunction

Natalia F. Mendes[1,2], Ariane M. Zanesco[2], Cristhiane F. Aguiar[3], Gabriela F. Rodrigues-Luiz[4],
Dayana C. da Silva[2], Jonathan F. Campos[2], Niels O. S. Câmara[5], Pedro M. M. de Moraes-Vieira[3],
Eliana P. de Araújo[2,6,#], Licio A. Velloso[2,7,*,#]

[1]School of Medical Sciences, Department of Translational Medicine (Section of Pharmacology),
University of Campinas, Brazil.
[2]Laboratory of Cell Signaling, Obesity and Comorbidities Research Center, University of
Campinas, Brazil.
[3]Laboratory of Immunometabolism, Institute of Biology - University of Campinas, Brazil.
[4] Department of Microbiology, Immunology and Parasitology, Federal University of Santa
Catarina, Brazil.
[5]Laboratory for Transplantation Immunobiology, Institute of Biomedical Sciences, University of
Sao Paulo, Brazil.
[6]Faculty of Nursing, University of Campinas, Brazil.
[7]National Institute of Science and Technology on Neuroimmunomodulation, Rio de Janeiro,
Brazil.
23
24
25
26 * Corresponding Authors:
27 Licio A. Velloso. Electronic address: lavellos@unicamp.br
28 Natalia F. Mendes. Eletronic address: natalia.mendesss@gmail.com
29 Laboratory of Cell Signaling-Obesity and Comorbidities Research Center, University of
30 Campinas, 13083-864 Campinas, Brazil
31
32 # EPA and LAV contributed equally as supervisors of this study

Abstract

Microgliosis is an important component of diet-induced hypothalamic inflammation in obesity. A few hours after the introduction of a high-fat diet, the mediobasal hypothalamus resident microglia undergo morphological and functional changes toward an inflammatory phenotype. If the consumption of large amounts of dietary fats persists for long periods, bone marrow-derived myeloid cells are recruited and integrated into a new landscape of hypothalamic microglia. However, it is currently unknown what are the transcriptional signatures and specific functions exerted by either resident or recruited subsets of hypothalamic microglia. Here, the elucidation of the transcriptional signatures revealed that resident microglia undergo only minor changes in response to dietary fats; however, under the consumption of a high-fat diet, there are major transcriptional differences between resident and recruited microglia with a major impact on chemotaxis. In addition, in recruited microglia, there are major transcriptional differences between females and males with an important impact on transcripts involved in neurodegeneration and thermogenesis. The chemokine receptor CXCR3 emerged as one of the components of chemotaxis with the greatest difference between recruited and resident microglia, and thus, was elected for further intervention. The hypothalamic immunoneutralization of CXCL10, one of the ligands for CXCR3, resulted in increased body mass gain and reduced energy expenditure, particularly in females. Furthermore, the chemical inhibition of CXCR3 resulted in a much greater change in phenotype with increased body mass gain, reduced energy expenditure, increased blood leptin, glucose intolerance, and reduced insulin. Thus, this study has elucidated the transcriptional differences between resident and recruited hypothalamic microglia in diet-induced obesity, identifying chemokines as a relevant subset of genes undergoing regulation. In addition, we showed that a subset of recruited microglia expressing CXCR3 has a protective, rather than a detrimental role in the metabolic outcomes promoted by the consumption of a high-fat diet, thus, establishing a new concept in obesity-associated hypothalamic inflammation.

Keywords: Hypothalamus, obesity, monocytes, microglia, chemokines.

Introduction

Obesity affects over 600 million people worldwide and projections are pessimistic indicating that over a billion people will be diagnosed with obesity by the year 2030 (worldobesity.org). Obesity develops as a consequence of a chronic state of anabolism in which caloric intake overcomes energy expenditure [1]. Both, experimental and human studies have indicated that, at least in part, the chronic anabolic state leading to obesity develops as a consequence of a defective hypothalamic regulation of whole-body energy balance [2] [3] [4].

Experimental studies have shown that the consumption of large amounts of saturated fats triggers an inflammatory response in the hypothalamus affecting the function of key neurons involved in the regulation of food intake, energy expenditure, and systemic metabolism [5] [6] [7] [8]. In addition, human studies using magnetic resonance imaging have identified hypothalamic gliosis in adults and children with obesity, providing clinical evidence for the existence of an obesity-associated hypothalamic inflammation [4] [9] [10]. Microglia are key cellular components of this inflammatory response undergoing structural and functional changes that develop early after the introduction of a high-fat diet (HFD) [11] [12] [13] [14]. Distinct strategies used to inhibit hypothalamic microglia resulted in impaired diet-induced hypothalamic inflammation, increased leptin sensitivity, reduced spontaneous caloric intake, and improved systemic glucose tolerance [12] [15]. Thus, elucidating the mechanisms involved in the regulation of microglial response to dietary factors may provide an advance in the definition of the pathophysiology of obesity, and potentially identify new targets for interventions aimed at treating obesity and its metabolic comorbidities [16] [17] [18].

Resident microglia are derived from the yolk sac primitive hematopoietic cells and populate the neuroepithelium at embryonic day 9.5 [19]. Under baseline conditions, in the absence of infections, trauma, or other types of potentially harmful stimuli, resident microglia remain rather quiescent; however, under stimulus, they undergo rapid morphological and functional changes aimed at confronting the threat [20]. In the hypothalamus, differently from most parts of the brain, resident microglia are responsive to fluctuations in the blood levels of nutrients and hormones involved in metabolic control, such as leptin [12]. Thus, a simple meal can promote considerable changes in hypothalamic resident microglia indicating the involvement of these cells in the complex network of cells that regulate whole body metabolism [21]. Under the consumption of a nutritionally balanced diet, meal-induced changes in hypothalamic resident microglia are cyclic and completely reversible [21]. However, under consumption of an HFD, microglia present profound changes in morphology and function; moreover, upon prolonged consumption of this type of diet, there is recruitment of bone marrow-derived myeloid cells that will compose a new landscape of hypothalamic microglia [12] [15] [18]. Despite the considerable advance in the understanding of how hypothalamic microglia are involved in diet-induced obesity (DIO), the transcriptional landscapes of resident and recruited microglia are currently unknown; and, the specific functions of either subset of microglia remain elusive.

In this study, we first elucidated the transcriptional differences between resident and recruited hypothalamic microglia in DIO, identifying chemokines as a relevant subset of genes undergoing regulation. Next, we identified a subset of recruited microglia expressing CXCR3, which exerts a protective role against DIO.

Results

Elucidating the transcriptional signatures of hypothalamic resident and recruited microglia in diet-induced obesity. Double reporter mice were obtained by the crossing of CX3CR1[GFP] and CCR2[RFP] (Fig. 1a). At the age of 8 weeks, female and male mice were randomly selected for either chow or HFD feeding for 28 days, and then specimens were harvested for analysis (Fig. 1b). In flow cytometry, cells expressing CCR2 could not be detected in the hypothalamus of mice fed on chow (Fig. 1c), whereas in the white adipose tissue, virtually all cells expressing CX3CR1 were also expressing CCR2 (Fig. 1c). Conversely, in both female and male mice fed on HFD, CCR2 cells accounted for approximately 10% of hypothalamic microglia (Fig. 1d). Histological analysis confirmed the results obtained by flow cytometry (Fig. 1e); furthermore, it was shown that CD169, which is classically regarded as a marker of bone marrow-derived cells [22], is expressed both in resident as well as in recruited microglia (Fig. 1f), confirming data published elsewhere [18]. Next, to prepare the samples for RNA sequencing, we sorted hypothalamic microglia expressing either CX3CR1 or CCR2 (Fig. 2a). The quality of sorting was confirmed by determining positivity for CX3CR1 (Fig. 2b) and CCR2 (Fig. 2c), and also by determining the positivity for several markers of resident microglia (Fig. 2d-2m) and several markers of bone marrow-derived cells (Fig. 2n-2x). The elucidation of the transcriptional landscapes of CX3CR1 and CCR2 revealed that either diet or sex exerted only minor differences in the expression of transcripts in CX3CR1 cells (Fig. 3a-3b); nevertheless, sex exerted major differences in CCR2 cells (Fig. 3a-3b). The direct comparisons between CX3CR1 and CCR2 obtained from mice fed on the HFD revealed the vast differences in the transcriptional landscapes of either female (Fig. 3c) or male (Fig. 3d) mice. Furthermore, the direct comparisons between female and male mice revealed a considerable degree of sexual dimorphism in the transcriptional landscapes of recruited CCR2 cells (Fig. 3e). In CX3CR1 cells, the consumption of the HFD impacted on IL17, lipids, toll-like receptor signaling, tumor necrosis factor signaling and chemokines (Fig. 4a); whereas in CCR2 cells, the consumption of the HFD impacted on lipids, toll-like receptor signaling, tumor necrosis factor signaling, chemokines, neurotrophins signaling, reactive oxygen species, thermogenesis, and pathways related to neurodegeneration (Fig. 4a). Next, we asked what functions related to chemotaxis were predominantly regulated in CCR2 cells of mice fed on the HFD (Fig. 4b-4c). As cell chemotaxis emerged as an important function in both females (Fig. 4b) and males (Fig. 4c), we looked with greater detail into the expression of chemokines and chemokine receptors (Fig. 4d). Virtually all the transcripts evaluated showed diametrically opposite expression between CX3CR1 and CCR2 (Fig. 4d).

Cxcr3 is highly expressed in recruited microglia. As we were particularly interested in identifying factors involved in the recruitment of CCR2 to the hypothalamus, we evaluated cytokine receptors with high expression in CCR2 and low expression in CX3CR1. As depicted in Fig. 5, Ccr3 (Fig. 5a), Ccr7 (Fig. 5b), Ccr8 (Fig. 5c), Cxcr2 (Fig. 5d), Cxcr3 (Fig. 5e), Cxcr4 (Fig. 5f), Cxcr5 (Fig. 5g), Cxcr6 (Fig. 5h) and Cxcr7 (Fig. 5i) were all expressed in CCR2 cells and virtually absent from CX3CR1 cells. Cxcr3 (Fig. 5e) and Cxcr6 (Fig. 5h) presented the highest expressions, and therefore, we performed a search for previous studies looking at either of these chemokine receptors in the context of DIO hypothalamic inflammation. Using the terms, hypothalamus, hypothalamic, obesity, inflammation, Cxcr3, and Cxcr6, we could find no prior publications. As Cxcr3 is involved in interferon-gamma (IFN-γ) induction [23], and IFN-γ is expressed in the context of DIO hypothalamic inflammation [5], we looked into the IFN-γ-related pathways regulated in recruited microglia (Fig. 6). First, we asked if the canonical ligands for CXCR3, C*XCL9*, CCL10, and CXCL11, were expressed in either CX3CR1 or CCR2 cells. C*XCL11* was not

detected in either cell type (not shown). CXCL9 (Fig. 6a) was expressed in CX3CR1 cells, only,
whereas CXCL10 (Fig. 6b) was expressed in both CX3CR1 and CCR2 cells. In addition, in both
female and male mice, Ifng was expressed in CCR2, but not in CX3CR1 cells (Fig. 6c).
Furthermore, IFN-γ pathways were shown to be modulated in both female (Fig. 6d) and male
(Fig. 6e) CCR2 cells. Thus, we elected CXCR3 as a target for further intervention.

The immunoneutralization of hypothalamic CXCL10 leads to increased body mass gain in
female mice. As an attempt to interfere with CXCR3 actions in CCR2 cells, we targeted one of its
ligands, CXCL10. As depicted in Fig. 7a, mice were submitted to two intracerebroventricular
(icv) injections of an anti-CXCL10 antibody aimed at immunoneutralizing the target protein in
the hypothalamus. As a result of the immunoneutralization of CXCL10, there were smaller
numbers of CCR2-positive cells in the hypothalamus of both female and male mice fed on an
HFD (Fig. 7b). However, there were no major changes in the numbers of CXCR3-expressing cells
in the hypothalamus of either female (Fig. 7c) or male (Fig. 7d) mice fed on an HFD. This was
accompanied by no changes in the transcript levels of Cxcr3 and several other chemokine-
related transcripts (Fig. 7c-7d), except for a trend to decrease Cxcl11 and an increase of Cxcr4
in females (Fig. 7c); and a decrease of Cxcl10 and a trend to increase Cx3cl1 in males (Fig. 7d).
Nevertheless, the immunoneutralization of hypothalamic CXCL10 (Fig. 8 and Fig. 9) resulted in
increased body mass gain (Fig. 8a-8b), a trend to reduce blood triglycerides (Fig. 8f), reduced
blood cholesterol (Fig. 8g), reduced expression of Agrp transcript in the hypothalamus (Fig. 8k),
a trend to reduce blood insulin (Fig. 8n), trends to reduce Il1b and Il6 transcripts in the
hypothalamus (Fig. 8o), and a trend to reduce respiratory quotient (Fig. 8s) during the dark
cycle in female mice. Conversely, in male mice (Fig. 9), the inhibition of hypothalamic CXCL10
had only a minor effect, leading to a trend to reduce hypothalamic Npy (Fig. 9k), a trend to
reduce hypothalamic Il1b, and a trend to increase hypothalamic Il6 (Fig. 9o).

The inhibition of CXCR3 worsens body mass gain and the metabolic phenotype of mice fed on a
high-fat diet. CXCR3 was inhibited using a pharmacological antagonist, AMG487 [24] (Fig. 10a).
The intervention resulted in the reduction of CCR2 (Fig. 10b) and CXCR3 (Fig. 10c-10d) cells in
the hypothalamus of both female and male mice fed an HFD. In addition, there was a reduction
of Ccl2 and an increase of Cx3cl1 transcripts in the hypothalamus of females (Fig. 10e), and a
reduction of Ccl2 transcripts in the hypothalamus of male (Fig. 10f) mice fed an HFD. The
inhibition of CXCR3 had a major impact on metabolic phenotype; thus, in female mice fed an
HFD, there was an increase in body mass gain (Fig. 11a-11b), a trend to increase brown adipose
tissue mass (Fig. 11c), a trend to increase blood leptin (Fig. 11e), an increase in blood
triglycerides (Fig. 11f), a trend to reduce hypothalamic Pomc (Fig. 11k), an increase in
hypothalamic Npy (Fig. 11k), a worsen glucose tolerance (Fig. 11l), increased fasting blood
glucose (Fig. 11m), reduced blood insulin (Fig. 11n), and reduction of Il6 and Tlr4 transcripts in
the hypothalamus (Fig. 11o). In males, the inhibition of CXCR3 promoted an increased body
mass gain (Fig. 12a-12b), increased white adipose tissue mass (Fig. 12d), increased blood leptin
(Fig. 12e), increased hypothalamic Npy and Mch (Fig. 12k), and reduced hypothalamic Tnfa and
Nlrp3 (Fig. 12o).

Discussion

In this study, we elucidated the transcriptional landscapes of resident and recruited microglia in
the hypothalamus of mice. We showed that resident microglia undergo minor transcriptional
changes when mice are fed an HFD; however, there are vast differences when confronting

resident versus recruited microglia transcriptomes. Moreover, there is a considerable degree of
sexual dimorphism in the transcriptomes of recruited microglia in mice fed an HFD. Upon
exploration of the differences between resident and recruited microglia, we identified the
chemokine receptor Cxcr3 as an interesting candidate for intervention as it was highly
expressed in recruited cells. The inhibition of CXCR3 resulted in increased body mass gain,
worsening of glucose intolerance, and increased expression of hypothalamic Npy. Thus, the
study is the first to identify a subset of recruited microglia that has a protective role against the
deleterious outcomes of DIO, therefore establishing a new concept in obesity-associated
hypothalamic inflammation.

Early studies in this field have shown that dietary fats, particularly long-chain saturated fatty
acids, trigger an inflammatory response in the MBH that emerges a few hours after the
introduction of an HFD and progresses to chronicity if the consumption of the HFD persists for
long [5] [6] [7]. Microglia play an important role in this inflammatory response, and studies
have shown that during a prolonged consumption of an HFD, there is recruitment of bone
marrow-derived cells to compose a new hypothalamic microglia landscape [9] [15] [18].
However, it was previously unknown what are the transcriptional signatures of hypothalamic
resident and recruited microglia in DIO.

To elucidate the transcriptional landscapes of resident and recruited microglia, we initially
prepared a double reporter mouse for CX3CR1 and CCR2. CX3CR1 encodes for fractalkine
receptor and is highly expressed in resident microglia [25]. The creation of CX3CR1 reporter
mice was regarded as an important step toward the characterization of resident microglia, and
several studies in the field employ this model [26] [27]. Conversely, CCR2 is expressed only in
bone marrow-derived cells and is regarded as a good marker for studying recruited microglia
[25] [28]. The quality of our model was proven good as makers of resident microglia were
present in CX3CR1 cells, only, whereas markers of recruited microglia were present in CCCR2
cells, only. Moreover, as previously described, we could find no CCR2 cells in the hypothalamus
of mice fed chow [14] [18].

The first important, and previously unknown finding emerged from the comparison of the
transcriptomes of resident microglia in mice fed chow versus mice fed an HFD. In females,
there were only 34 transcripts undergoing significant changes between the two dietary
conditions, whereas in males, the number was greater, 412, but still quite small considering the
whole transcriptome of resident microglia [29]. In a study evaluating the single-cell
transcriptomics of hypothalamic cells, the consumption of an HFD resulted in minor changes in
the so-called macrophage-like cells [30]; however, the detailed transcriptional landscape of
these cells was not explored in depth, so it is uncertain if it contained both resident and
recruited microglia. In an experimental model of Alzheimer´s disease, which evaluated male
and female mice, there were hippocampal resident microglial transcriptional changes of the
same magnitude as the one we found in the hypothalamus, affecting approximately 300 genes
[31]. In an experimental model of cerebral hemorrhage evaluating only males, the impact on
microglia transcriptome was also small, affecting only 10% of the evaluated genes [32]. In
addition, in a study evaluating transcriptional changes of resident microglia during aging [33],
there were also important differences between female and male mice; and, the magnitude of
the transcriptional changes occurring during aging was about the same as we see in the dietary
intervention. Thus, it seems that in different brain regions and under distinct interventions, the
magnitude of resident microglia transcriptional changes is quite small; however, the number of
studies evaluating this question is not expressive; thus, further studies are needed to provide a

definitive view regarding the actual magnitude of plasticity of resident microglia. Despite the
small number of transcripts undergoing changes in our model, the greatest impact occurred in
the expression of genes related to IL-17 signaling, lipid metabolism, TLR signaling, TNF
signaling, and chemokine signaling, which strongly supports the role of dietary lipids in the
induction of an inflammatory response by the resident microglia, supporting previous studies
in the field [6] [9] [15] [18]. Interestingly, IL17 signaling emerged as a major pathway
modulated by DIO. In a recent study, we have shown that IL17 can act upon POMC neurons
promoting a reduction in calorie intake [34]. The current finding of a transcriptional modulation
of IL17-related genes in resident microglia opens a new perspective in the understanding of
how inflammatory signals modulate the function of the hypothalamus in obesity, which should
be explored in the future.

Next, we confronted the transcriptomes of resident and recruited microglia, and in this case,
there were huge differences, reaching over 7,000 transcripts. Per se, this finding reveals a
striking difference between resident and recruited microglia in the hypothalamus of mice fed
an HFD. Nevertheless, despite this is new information regarding the hypothalamus, similar
findings were reported in other brain regions submitted to distinct interventions, such as in an
experimental model of glioma [35], in flavivirus infection [36], and in COVID-19 [37]. The
evaluation of the main pathways differently expressed in the two cell subsets revealed major
differences in lipids, toll-like receptor signaling, tumor necrosis factor signaling, chemokines,
neurotrophins signaling, reactive oxygen species, thermogenesis, and pathways related to
neurodegeneration. The impact of the consumption of an HFD on the regulation of lipid-
related pathways, toll-like receptor, and tumor necrosis factor signaling has been widely
explored in several previous studies, and interventions in these systems are known to mitigate
the harmful effects of the diet [6] [9] [15] [18]. However, little has been done about the
characterization of the mechanisms of chemotaxis that drive the recruitment of bone marrow-
derived cells to the hypothalamus. Therefore, we looked with greater detail into the main
chemokines and chemokine receptors differentially expressed in the two subsets of cells. We
elected Cxcr3 because it was highly expressed in Ccr2 cells and presented low expression in
Cx3cr1 cells. CXCR3 is a chemokine receptor that is involved in the recruitment of distinct types
of bone marrow-derived monocytic cells, such as plasmacytoid monocytes [38], synovial tissue
monocytes [39], and dendritic cells [40]. In the brain, CXCR3-expressing cells have been
implicated in Alzheimer's disease and other age-dependent cognitive dysfunctions [41] [42],
multiple sclerosis [43], epilepsy [44], and stoke [45]. However, no previous study has evaluated
CXCR3 in the hypothalamus in the context of obesity.

First, we asked if the known ligands for CXCR3, and Infg, which is induced in response to the
activation of CXCR3, were present in either subset of cells. We found Cxcl11 in neither cell
subset; Cxcl9 was expressed in Cx3cl1 cells, only; Cxcl10 was expressed in both subsets of cells,
with greater expression in Ccr2 cells; and Infg was expressed in Ccr2 cells only. Moreover, there
was a considerable engagement of INF-γ-related pathways in Ccr2 cells of both female and
male mice fed an HFD. Thus, we considered CXCR3 as a promising target for intervention.

Next, we intervened in one of the ligands for CXCR3, CXCL10. For that, we performed icv
injections of an immunoneutralizing antibody, which resulted in smaller numbers of CCR2 cells
in the hypothalamus. However, the intervention promoted minimal changes in the
hypothalamic expression of transcripts encoding several components of the chemotaxis
machinery. Nevertheless, in females, the inhibition of CXCL10 resulted in increased body mass
gain, reduction of hypothalamic Agrp, trends to reduce hypothalamic Il1b and Il6, and a trend

to reduce blood insulin. In males, the phenotype was much milder, leading to minimal changes
in hypothalamic Npy, Il1b, and Il6. Little is known about the involvement of CXCL10 in
hypothalamic physiology and pathology. In a model of caloric restriction, there was an increase
in the hypothalamic expression of Cxcl10 [46], and this was regarded as a component of the
mechanism of neuroprotection induced by caloric restriction. In addition, in a model of
hypothalamic inflammation elicited by exogenous LPS, Cxcl10 emerged as one of the
transcripts undergoing the greatest increase in the hypothalamic paraventricular nucleus [47].

As CXCR3 can be engaged by distinct chemokines, we decided to use a broader intervention,
pharmacologically inhibiting CXCR3. AMG487 is a chemical inhibitor of CXCR3 with an IC_{50} value
of 8.0 nM [48]. Upon treatment with AMG487, there was a reduction of the migration of Ccr2
cells to the hypothalamus, which was accompanied by minimal changes in the expressions of
chemokines and chemokine receptors. Under inhibition of CXCR3, both female and male mice
presented increased body mass gain, which was accompanied by increased blood leptin,
increased fasting glucose, increased hypothalamic Npy, and a reduction in markers of
hypothalamic inflammation. There were no changes in caloric intake, however, there were
reductions in energy expenditure during some periods during the 24 hours of recording.

These findings reveal that at least one subset of recruited microglia, have a protective rather
than a harmful role in DIO-associated hypothalamic inflammation. This is a completely new
concept in the field because all previous studies evaluating hypothalamic microglia in DIO
reported that, once active in response to dietary fats, either resident or recruited microglia
exerted inflammatory actions that impacted negatively energy balance and glucose tolerance
[11] [12] [13] [15]. This concept has been recently explored in depth in a study that used
elegant models to either activate or inactivate microglia [14] and the results confirm that,
whenever manipulating microglia using approaches that are not specific for a given subset of
cells, the net result is worsening of the metabolic phenotype when microglia is activated and
improvement of the metabolic phenotype when microglia is inactivated. Thus, we believe that
the subset of recruited microglia herein identified plays a regulatory role in hypothalamic
inflammation.

In conclusion, this study elucidated the transcriptional landscapes of resident and recruited
hypothalamic microglia in DIO. In resident microglia, the consumption of an HFD resulted in
small changes in transcript expression, whereas the confrontation of the transcriptional
landscapes of resident versus recruited microglia revealed broad differences that encompass
lipids, toll-like receptor signaling, tumor necrosis factor signaling, chemokines, neurotrophins
signaling, reactive oxygen species, thermogenesis, and pathways related to neurodegeneration.
In addition, the study revealed a considerable sexual dimorphism in the transcriptional
landscape of both resident and recruited microglia. The study also identified a subset of
recruited microglia, expressing the chemokine receptor Cxcr3 that has a protective role against
the harmful metabolic effects of the HFD; thus, providing a new concept in DIO-associated
hypothalamic inflammation.

Materials and methods

Animal care and diets. All animal care and experimental procedures were conducted in
accordance with the guidelines of the Brazilian College for Animal Experimentation and
approved by the Institutional Animal Care and Use Committee (CEUA 5497-1/2020 and 6210-

1/2023). Heterozygous CX3CR1$^{GFP/+}$CCR2$^{RFP/+}$ mice were generated by mating CX3CR1^{GFP}

homozygous mice (JAX#005582) with CCR2^{RFP} homozygous mice (JAX#017586). Genotypes of

these mice were identified by polymerase chain reaction (PCR). Mice were fed on a standard

chow diet (Nuvilab; 3.76 kcal/g; 12.6% energy from protein, 77.7% energy from carbohydrate,

and 9.58% energy from fat) or high-fat diet (HFD) (5.28 kcal/g; 12.88% energy from protein,

27.1% energy from carbohydrate, and 60% energy from fat) according to the experimental

protocols. Food and water were available ad libitum throughout the experimental periods,

except for the protocols that required fasting. The room temperature was controlled (22-24 °C),

and a light-dark cycle was maintained on a 12-hour on-off cycle.

Flow cytometry. For the separation of CX3CR1^{GFP+} and CCR2^{RFP+} cells from the white adipose

tissue of CX3CR1$^{GFP/+}$CCR2$^{RFP/+}$ mice we collected the retroperitoneal fat depot of one animal

fed on a HFD for 4 weeks. It was minced and digested with type VIII collagenase (0.5 mg/mL,

Sigma-Aldrich) in PBS for 20 min at 37°C with shaking. After digestion, the suspension was

filtered using a 100 μm cell filter. For isolation of the same cells from the hypothalamus,

samples of five CX3CR1$^{GFP/+}$CCR2$^{RFP/+}$ mice fed on an HFD for 4 weeks were pooled together and

gently pressed through a cell strainer (100 μm). The cell solution was subjected to a Percoll

gradient (70/40%) for monocyte purification. Samples were acquired on a BDFacs Symphony

instrument (BD Biosciences, USA) and then analyzed using FlowJo software.

Cell sorting. For cell sorting of CX3CR1^{GFP+} and CCR2^{RFP+} cells from the hypothalamus, we

employed CX3CR1^{GFP} mice fed on chow diet and CX3CR1$^{GFP/+}$CCR2$^{RFP/+}$ mice fed on HFD for 4

weeks. Harvested hypothalami of 20-30 male or 20-30 female mice were pooled together for

each sample and gently pressed through a cell strainer (100 μm). The cell solution was

subjected to a Percoll gradient (70/40%) for monocyte purification. The sorting was conducted

on a BDFacs Melody instrument (BD Biosciences, USA).

RNA-sequencing (RNA-Seq) and analysis. Cell-sorted CX3CR1 GFP+ and CCR2 RFP+ cell samples

from the hypothalamus were lysed for RNA extraction using the RNAqueous Micro kit

(Invitrogen). RNA integrity was analyzed on a Bioanalyzer RNA Pico 6000 chip at the Core

Facility for Scientific Research – University of São Paulo (CEFAP-USP). Low input RNA-Seq library

preparation (Takara SMART-Seq v4) and sequencing by Illumina NovaSeq S2 PE150 Sequencing

Lane (40M read pairs/sample avg) were performed by Maryland Genomics (Institute for

Genome Sciences - IGS, University of Maryland School of Medicine – Baltimore, USA). Illumina

sequencing adapters and low-quality reads were removed with Trimmomatic. Trimmed reads

were aligned to the mouse reference genome (GRCm39) by STAR. Aligned reads were mapped

to features using HTSeq, and differential expression analyses were performed using the DESeq2

package. Genes having less than 3 CPM were excluded prior to statistical analysis, and

differentially expressed genes (DEGs) were selected using as cutoffs the adjusted p-value <

0.05. Heatmaps were performed using Pheatmap and a list of DEGs was passed to EnrichR and

cluster Profiler for enrichment analyses.

CXCL10 immunoneutralization. For central neutralization of CXCL10, 8-wk CCR2^{RFP} male and

female mice underwent a stereotaxic surgery for ICV injections of anti-CXCL10 Monoclonal

Antibody (2 ul, Cat# MA5-23774, Thermo Fischer). The control groups were ICV injected with

Mouse IgG2a Isotype Control (2 ul, Cat#02-6200, Thermo Fischer). Two distinct ICV injections

were performed on Day 0 and Day 14, respectively, of the experimental protocol. For that, mice

were anesthetized with ketamine (100 mg/kg) and xylazine (10 mg/kg) and submitted to

stereotaxic surgery (Ultra Precise–model 963, Kopf). ICV coordinates were [antero-

posterior/lateral/depth to bregma]: -0.46/-1.0/−2.3 mm. Immediately after the first surgery, at

Day 0, mice began to be fed on HFD for 4 weeks. From Day 0 to Day 28 food intake and body
weight were evaluated weekly.

CXCR3 antagonism. For systemic blockage of CXCR3, 8-wk CCR2[RFP] male and female mice
underwent a treatment with AMG487 (Tocris Bioscience, Bristol, UK), an active and selective
CXC chemokine receptor 3 (CXCR3) antagonist. The in vivo formulation of AMG487 was
prepared in 20% hydroxypropyl-β-cyclodextrin (Sigma, St. Louis, MO). A 50% hydroxypropyl-β-
cyclodextrin (Sigma, St. Louis, MO) solution was prepared and AMG487 was added to this
solution, it was incubated in a sonicating water bath for 2 hours with occasional vortexing.
Next, distilled water was added to give the appropriate final concentration of AMG487 in 20%
of hydroxypropyl-β-cyclodextrin. This solution at 20% served as the vehicle. Mice were treated
with AMG487 or vehicle (VEH group) intraperitoneally at 5 mg/kg every 48h throughout four
weeks. During this period, mice were fed on HFD, and food intake and body weight were
evaluated weekly.

Glucose tolerance test. On the 24th day of CXCR3 blockage and CXCL10 neutralization
experimental protocols, mice were fasted for 6 hours, and blood glucose was measured via tail
bleed at baseline and 15, 30, 60, 90, and 120 min after an intraperitoneal injection of glucose
(2.0 g/kg).

Indirect calorimetry. The oxygen consumption (VO_2), carbon dioxide production (VCO_2), energy
expenditure, and respiratory quotient (RQ) were measured using an indirect open-circuit
calorimeter (Oxylet M3 system; PanLab/Harvard Apparatus, MA, USA). Mice were allowed to
adapt for 12 hours before data were recorded for 24 hours (light and dark cycles).

Immunofluorescence. On the day 28th of CXCR3 blockage and CXCL10 neutralization
experimental protocols, male and female mice were perfused with 0,9% saline followed by 4%
formaldehyde by cardiac canulation. Brains were extracted and incubated in 4% formaldehyde
overnight at 4 °C for extended fixation. The brains were then incubated in 30% sucrose at 4 °C
for 48h. A series of 20 µm-thick frozen sections (4 series equally) were prepared using a
cryostat and stored in an anti-freezing solution. For the free-floating immunostaining, slices
were washed with 0.1 M phosphate-buffered saline (PBS) (3 times, 5 min each) and blocked
with 0.2% Triton X-100 and 5% donkey serum in 0.1 M PBS for 2 h at room temperature. Slices
were incubated overnight at 4 °C with Anti-Cxcr3 (1:200, Cat# NB100-56404, Novus Biologicals)
or Anti-Sialoadhesin/CD169 (1:200, ab18619, Abcam) in a blocking solution. After washing with
0.1 M phosphate-buffered saline (PBS) (3 times, 5 min each), sections were incubated with
fluorophore-labeled secondary antibody (donkey anti-rabbit Alexa Fluor 405, 1:500, Cat#
A48258, Invitrogen or goat anti-mouse Alexa Fluor 405, 1:500, Cat# A31553, Invitrogen) in a
blocking solution for 2 h at room temperature. After washing again with 0.1 M phosphate-
buffered saline (PBS) (3 times, 5 min each), brain slices were mounted onto slides with ProLong
Diamond antifade mountant (Cat# P36930, Thermo Fischer). Sections were visualized with a
Zeiss LSM780, confocal microscope (Carl Zeiss AG, Germany) at the National Institute of
Photonics Applied to Cell Biology (INFABIC) at the University of Campinas.

Quantitative reverse transcription-polymerase chain reaction (qRT-PCR). Total RNA was
extracted using a TRIzol reagent (Thermo Fisher Scientific) and synthesized cDNA with a High-
Capacity cDNA Reverse Transcription Kit (HighCapacity cDNA Reverse Transcription Kit, Life
Technologies). Real-time PCR reactions were run using the TaqMan system (Applied
Biosystems). Primers used were Cxcr3 (Mm99999054_s1); Cxcl9 (Mm00434946_m1), Cxcl10
(Mm00445235_m1); Cxcl11 (Mm00444662_m1); Ccl2 (Mm00441242_m1); Cxcr4

(Mm01996749_s1); Cxcl12 (Mm00445553_m1), Cxcr6(Mm02620517_s1), Cxcl16
(Mm00469712_m1), Cx3cl1(Mm00436454_m1), Pomc (Mm00435874_m1), Agrp
(Mm00475829_g1); Npy (Mm00445771_m1); Cartpt (Mm04210469_m1), Pmch
(Mm01242886_g1), Tnfa (Mm00443258_m1), Il1b (Mm00434228_m1), Il6
(Mm00446190_m1), Nlrp3 (Mm00840904_m1), Tlr4 (Mm00445273_m1), Infg
(Mm01168134_m1). GAPDH (Mm99999915_g1) was employed as a reference gene. Gene
expression was obtained using the QuantStudio 6 (Thermo Fischer Scientific).

Hormonal and biochemical determinations. Serum insulin and leptin were measured by an
enzyme-linked immunosorbent assay (ELISA) kit (#EZRMI-13K and #EZML-82K; Millipore).
Serum triglyceride levels and total cholesterol were measured using a commercial colorimetric
assay kit (LaborLab®, Guarulhos - SP, Brazil) following the manufacturer's instructions.

Statistical analysis. Data are presented as means ± standard error of the mean (SEM). The
statistical analyses were carried out using a non-parametric Mann-Whitney test or two-way
analysis of variance (ANOVA) when appropriate. Post hoc comparisons were performed using
Sidak's test. Statistical significances were analyzed using Prism 8.0 software (GraphPad
Software, La Jolla, CA). A p-value ≤0.05 was considered statistically significant.

Declaration of Interests/Relevant conflicts of interests/Financial disclosures. All authors declare no
competing financial or other interests regarding this study.

Authors' contributions. NFM, EPA and LAV designed and planned the study; NFM, AMZ, DCS, and JFC
performed most of the experiments; NFM and CFA performed FACS and cytometry analysis; GFL
conducted bioinformatic analysis; NOSC provided animals models; PMMM-V supervised the work of CFA
and provided equipment and methods for FACS and cytometry; NFM, EPA, and LAV discussed the data.
NFM and LAV wrote the manuscript and prepared figures; EPA supervised NFM during her doctorate
program. All authors read and approved the final manuscript.

Funding. This research was funded by The Sao Paulo Research Foundation (FAPESP): 2013/07607-8;
2017/22511-8; and 2021/00443-6.

Data Availability. Data will be available upon request to the corresponding author, L.A.V
(lavellos@unicamp.br).

Acknowledgments. We are grateful to Erika Roman, Joseane Morari, Marcio Cruz, and Gerson Ferraz for
laboratory management.

References

464   [1]   Theilade, S., Christensen, M.B., Vilsbøll, T., Knop, F.K., 2021. An overview of obesity mechanisms in
465   humans: Endocrine regulation of food intake, eating behaviour and common determinants of body
466   weight. Diabetes, Obesity & Metabolism 23 Suppl 1: 17–35, Doi: 10.1111/dom.14270.
467   [2]   Cavadas, C., Aveleira, C.A., Souza, G.F.P., Velloso, L.A., 2016. The pathophysiology of defective
468   proteostasis in the hypothalamus - from obesity to ageing. Nature Reviews. Endocrinology 12(12):
469   723–33, Doi: 10.1038/nrendo.2016.107.
470   [3]   Sonnefeld, L., Rohmann, N., Geisler, C., Laudes, M., 2023. Is human obesity an inflammatory
471   disease of the hypothalamus? European Journal of Endocrinology 188(3): R37–45, Doi:
472   10.1093/ejendo/lvad030.

[4] van de Sande-Lee, S., Melhorn, S.J., Rachid, B., Rodovalho, S., De-Lima-Junior, J.C., Campos, B.M.,
et al., 2020. Radiologic evidence that hypothalamic gliosis is improved after bariatric surgery in
obese women with type 2 diabetes. International Journal of Obesity (2005) 44(1): 178–85, Doi:
10.1038/s41366-019-0399-8.

[5] De Souza, C.T., Araujo, E.P., Bordin, S., Ashimine, R., Zollner, R.L., Boschero, A.C., et al., 2005.
Consumption of a fat-rich diet activates a proinflammatory response and induces insulin resistance
in the hypothalamus. Endocrinology 146(10): 4192–9, Doi: 10.1210/en.2004-1520.

[6] Milanski, M., Degasperi, G., Coope, A., Morari, J., Denis, R., Cintra, D.E., et al., 2009. Saturated fatty
acids produce an inflammatory response predominantly through the activation of TLR4 signaling in
hypothalamus: implications for the pathogenesis of obesity. The Journal of Neuroscience: The
Official Journal of the Society for Neuroscience 29(2): 359–70, Doi: 10.1523/JNEUROSCI.2760-
08.2009.

[7] Zhang, X., Zhang, G., Zhang, H., Karin, M., Bai, H., Cai, D., 2008. Hypothalamic IKKbeta/NF-kappaB
and ER stress link overnutrition to energy imbalance and obesity. Cell 135(1): 61–73, Doi:
10.1016/j.cell.2008.07.043.

[8] Milanski, M., Arruda, A.P., Coope, A., Ignacio-Souza, L.M., Nunez, C.E., Roman, E.A., et al., 2012.
Inhibition of hypothalamic inflammation reverses diet-induced insulin resistance in the liver.
Diabetes 61(6): 1455–62, Doi: 10.2337/db11-0390.

[9] Thaler, J.P., Yi, C.-X., Schur, E.A., Guyenet, S.J., Hwang, B.H., Dietrich, M.O., et al., 2012. Obesity is
associated with hypothalamic injury in rodents and humans. The Journal of Clinical Investigation
122(1): 153–62, Doi: 10.1172/JCI59660.

[10] Sewaybricker, L.E., Schur, E.A., Melhorn, S.J., Campos, B.M., Askren, M.K., Nogueira, G.A.S., et al.,
2019. Initial evidence for hypothalamic gliosis in children with obesity by quantitative T2 MRI and
implications for blood oxygen-level dependent response to glucose ingestion. Pediatric Obesity
14(2): e12486, Doi: 10.1111/ijpo.12486.

[11] Tapia-González, S., García-Segura, L.M., Tena-Sempere, M., Frago, L.M., Castellano, J.M., Fuente-
Martín, E., et al., 2011. Activation of microglia in specific hypothalamic nuclei and the cerebellum
of adult rats exposed to neonatal overnutrition. Journal of Neuroendocrinology 23(4): 365–70,
Doi: 10.1111/j.1365-2826.2011.02113.x.

[12] Valdearcos, M., Robblee, M.M., Benjamin, D.I., Nomura, D.K., Xu, A.W., Koliwad, S.K., 2014.
Microglia dictate the impact of saturated fat consumption on hypothalamic inflammation and
neuronal function. Cell Reports 9(6): 2124–38, Doi: 10.1016/j.celrep.2014.11.018.

[13] Fernández-Arjona, M.D.M., León-Rodríguez, A., Grondona, J.M., López-Ávalos, M.D., 2022. Long-
term priming of hypothalamic microglia is associated with energy balance disturbances under diet-
induced obesity. Glia 70(9): 1734–61, Doi: 10.1002/glia.24217.

[14] Valdearcos, M., Douglass, J.D., Robblee, M.M., Dorfman, M.D., Stifler, D.R., Bennett, M.L., et al.,
2017. Microglial Inflammatory Signaling Orchestrates the Hypothalamic Immune Response to
Dietary Excess and Mediates Obesity Susceptibility. Cell Metabolism 26(1): 185-197.e3, Doi:
10.1016/j.cmet.2017.05.015.

[15] Morari, J., Anhe, G.F., Nascimento, L.F., de Moura, R.F., Razolli, D., Solon, C., et al., 2014.
Fractalkine (CX3CL1) is involved in the early activation of hypothalamic inflammation in
experimental obesity. Diabetes 63(11): 3770–84, Doi: 10.2337/db13-1495.

[16] Mendes, N.F., Kim, Y.-B., Velloso, L.A., Araújo, E.P., 2018. Hypothalamic Microglial Activation in
Obesity: A Mini-Review. Frontiers in Neuroscience 12: 846, Doi: 10.3389/fnins.2018.00846.

[17] Mendes, N.F., Velloso, L.A., 2022. Perivascular macrophages in high-fat diet-induced hypothalamic
inflammation. Journal of Neuroinflammation 19(1): 136, Doi: 10.1186/s12974-022-02519-6.

[18] Valdearcos, M., Myers, M.G., Koliwad, S.K., 2019. Hypothalamic microglia as potential regulators of
metabolic physiology. Nature Metabolism 1(3): 314–20, Doi: 10.1038/s42255-019-0040-0.

[19] Ginhoux, F., Greter, M., Leboeuf, M., Nandi, S., See, P., Gokhan, S., et al., 2010. Fate Mapping
Analysis Reveals That Adult Microglia Derive from Primitive Macrophages. Science 330(6005):
841–5, Doi: 10.1126/science.1194637.

[20] Saijo, K., Glass, C.K., 2011. Microglial cell origin and phenotypes in health and disease. Nature
Reviews Immunology 11(11): 775–87, Doi: 10.1038/nri3086.

[21] Gao, Y., Ottaway, N., Schriever, S.C., Legutko, B., García-Cáceres, C., de la Fuente, E., et al., 2014.
Hormones and diet, but not body weight, control hypothalamic microglial activity. Glia 62(1): 17–
25, Doi: 10.1002/glia.22580.

[22] Chávez-Galán, L., Olleros, M.L., Vesin, D., Garcia, I., 2015. Much More than M1 and M2
Macrophages, There are also CD169(+) and TCR(+) Macrophages. Frontiers in Immunology 6: 263,
Doi: 10.3389/fimmu.2015.00263.

[23] Cole, K.E., Strick, C.A., Paradis, T.J., Ogborne, K.T., Loetscher, M., Gladue, R.P., et al., 1998.
Interferon-inducible T cell alpha chemoattractant (I-TAC): a novel non-ELR CXC chemokine with
potent activity on activated T cells through selective high affinity binding to CXCR3. The Journal of
Experimental Medicine 187(12): 2009–21, Doi: 10.1084/jem.187.12.2009.

[24] Walser, T.C., Rifat, S., Ma, X., Kundu, N., Ward, C., Goloubeva, O., et al., 2006. Antagonism of CXCR3
inhibits lung metastasis in a murine model of metastatic breast cancer. Cancer Research 66(15):
7701–7, Doi: 10.1158/0008-5472.CAN-06-0709.

[25] Greter, M., Lelios, I., Croxford, A.L., 2015. Microglia Versus Myeloid Cell Nomenclature during Brain
Inflammation. Frontiers in Immunology 6, Doi: 10.3389/fimmu.2015.00249.

[26] Yona, S., Kim, K.-W., Wolf, Y., Mildner, A., Varol, D., Breker, M., et al., 2013. Fate Mapping Reveals
Origins and Dynamics of Monocytes and Tissue Macrophages under Homeostasis. Immunity 38(1):
79–91, Doi: 10.1016/j.immuni.2012.12.001.

[27] Goldmann, T., Wieghofer, P., Müller, P.F., Wolf, Y., Varol, D., Yona, S., et al., 2013. A new type of
microglia gene targeting shows TAK1 to be pivotal in CNS autoimmune inflammation. Nature
Neuroscience 16(11): 1618–26, Doi: 10.1038/nn.3531.

[28] Mildner, A., Schmidt, H., Nitsche, M., Merkler, D., Hanisch, U.-K., Mack, M., et al., 2007. Microglia
in the adult brain arise from Ly-6ChiCCR2+ monocytes only under defined host conditions. Nature
Neuroscience 10(12): 1544–53, Doi: 10.1038/nn2015.

[29] Young, A.M.H., Kumasaka, N., Calvert, F., Hammond, T.R., Knights, A., Panousis, N., et al., 2021. A
map of transcriptional heterogeneity and regulatory variation in human microglia. Nature Genetics
53(6): 861–8, Doi: 10.1038/s41588-021-00875-2.

[30] Campbell, J.N., Macosko, E.Z., Fenselau, H., Pers, T.H., Lyubetskaya, A., Tenen, D., et al., 2017. A
molecular census of arcuate hypothalamus and median eminence cell types. Nature Neuroscience
20(3): 484–96, Doi: 10.1038/nn.4495.

[31] Rivera-Escalera, F., Pinney, J.J., Owlett, L., Ahmed, H., Thakar, J., Olschowka, J.A., et al., 2019. IL-1β-
driven amyloid plaque clearance is associated with an expansion of transcriptionally
reprogrammed microglia. Journal of Neuroinflammation 16(1): 261, Doi: 10.1186/s12974-019-
1645-7.

[32] Taylor, R.A., Chang, C.-F., Goods, B.A., Hammond, M.D., Mac Grory, B., Ai, Y., et al., 2017. TGF-β1
modulates microglial phenotype and promotes recovery after intracerebral hemorrhage. The
Journal of Clinical Investigation 127(1): 280–92, Doi: 10.1172/JCI88647.

[33] Li, X., Li, Y., Jin, Y., Zhang, Y., Wu, J., Xu, Z., et al., 2023. Transcriptional and epigenetic decoding of
the microglial aging process. Nature Aging, Doi: 10.1038/s43587-023-00479-x.

[34] Nogueira, G., Solon, C., Carraro, R.S., Engel, D.F., Ramalho, A.F., Sidarta-Oliveira, D., et al., 2020.
Interleukin-17 acts in the hypothalamus reducing food intake. Brain, Behavior, and Immunity 87:
272–85, Doi: 10.1016/j.bbi.2019.12.012.

[35] Ochocka, N., Segit, P., Walentynowicz, K.A., Wojnicki, K., Cyranowski, S., Swatler, J., et al., 2021.
Single-cell RNA sequencing reveals functional heterogeneity of glioma-associated brain
macrophages. Nature Communications 12(1): 1151, Doi: 10.1038/s41467-021-21407-w.

[36] Spiteri, A.G., Wishart, C.L., Ni, D., Viengkhou, B., Macia, L., Hofer, M.J., et al., 2023. Temporal
tracking of microglial and monocyte single-cell transcriptomics in lethal flavivirus infection. Acta
Neuropathologica Communications 11(1): 60, Doi: 10.1186/s40478-023-01547-4.

[37] Schwabenland, M., Salié, H., Tanevski, J., Killmer, S., Lago, M.S., Schlaak, A.E., et al., 2021. Deep
spatial profiling of human COVID-19 brains reveals neuroinflammation with distinct
microanatomical microglia-T-cell interactions. Immunity 54(7): 1594-1610.e11, Doi:
10.1016/j.immuni.2021.06.002.

[38] Cella, M., Jarrossay, D., Facchetti, F., Alebardi, O., Nakajima, H., Lanzavecchia, A., et al., 1999.
Plasmacytoid monocytes migrate to inflamed lymph nodes and produce large amounts of type I
interferon. Nature Medicine 5(8): 919–23, Doi: 10.1038/11360.

[39] Katschke, K.J., Rottman, J.B., Ruth, J.H., Qin, S., Wu, L., LaRosa, G., et al., 2001. Differential
expression of chemokine receptors on peripheral blood, synovial fluid, and synovial tissue
monocytes/macrophages in rheumatoid arthritis. Arthritis and Rheumatism 44(5): 1022–32, Doi:
10.1002/1529-0131(200105)44:5<1022::AID-ANR181>3.0.CO;2-N.

[40]   Padovan, E., Spagnoli, G.C., Ferrantini, M., Heberer, M., 2002. IFN-alpha2a induces IP-10/CXCL10
586   and MIG/CXCL9 production in monocyte-derived dendritic cells and enhances their capacity to
587   attract and stimulate CD8+ effector T cells. Journal of Leukocyte Biology 71(4): 669–76.

588   [41]   Jorfi, M., Park, J., Hall, C.K., Lin, C.-C.J., Chen, M., von Maydell, D., et al., 2023. Infiltrating CD8+ T
589   cells exacerbate Alzheimer's disease pathology in a 3D human neuroimmune axis model. Nature
590   Neuroscience 26(9): 1489–504, Doi: 10.1038/s41593-023-01415-3.

591   [42]   Schroer, A.B., Ventura, P.B., Sucharov, J., Misra, R., Chui, M.K.K., Bieri, G., et al., 2023. Platelet
592   factors attenuate inflammation and rescue cognition in ageing. Nature 620(7976): 1071–9, Doi:
593   10.1038/s41586-023-06436-3.

594   [43]   Bogers, L., Engelenburg, H.J., Janssen, M., Unger, P.-P.A., Melief, M.-J., Wierenga-Wolf, A.F., et al.,
595   2023. Selective emergence of antibody-secreting cells in the multiple sclerosis brain. EBioMedicine
596   89: 104465, Doi: 10.1016/j.ebiom.2023.104465.

597   [44]   Liang, P., Zhang, X., Zhang, Y., Wu, Y., Song, Y., Wang, X., et al., 2023. Neurotoxic A1 astrocytes
598   promote neuronal ferroptosis via CXCL10/CXCR3 axis in epilepsy. Free Radical Biology & Medicine
599   195: 329–42, Doi: 10.1016/j.freeradbiomed.2023.01.002.

600   [45]   Cai, W., Shi, L., Zhao, J., Xu, F., Dufort, C., Ye, Q., et al., 2022. Neuroprotection against ischemic
601   stroke requires a specific class of early responder T cells in mice. The Journal of Clinical
602   Investigation 132(15): e157678, Doi: 10.1172/JCI157678.

603   [46]   Matthews, D., Diskin, M.G., Kenny, D.A., Creevey, C.J., Keogh, K., Waters, S.M., 2017. Effect of short
604   term diet restriction on gene expression in the bovine hypothalamus using next generation RNA
605   sequencing technology. BMC Genomics 18(1): 857, Doi: 10.1186/s12864-017-4265-6.

606   [47]   Reyes, T.M., Walker, J.R., DeCino, C., Hogenesch, J.B., Sawchenko, P.E., 2003. Categorically distinct
607   acute stressors elicit dissimilar transcriptional profiles in the paraventricular nucleus of the
608   hypothalamus. The Journal of Neuroscience: The Official Journal of the Society for Neuroscience
609   23(13): 5607–16, Doi: 10.1523/JNEUROSCI.23-13-05607.2003.

610   [48]   Johnson, M., Li, A.-R., Liu, J., Fu, Z., Zhu, L., Miao, S., et al., 2007. Discovery and optimization of a
611   series of quinazolinone-derived antagonists of CXCR3. Bioorganic & Medicinal Chemistry Letters
612   17(12): 3339–43, Doi: 10.1016/j.bmcl.2007.03.106.

617   **Table**

618   **Table 1.** Number of DEGs for each comparison

		↑ Up	↓ Down
Effect of the diet in brain resident microglia cells	**hfd_cx_f** Cx3cr1_hfd_female *vs.* Cx3cr1_chow_female	25	9
	hfd_cx_m Cx3cr1_hfd_male *vs.* Cx3cr1_chow_male	261	151
Effect of the sex in brain resident microglia cells	**chow_cx** Cx3cr1_chow_male *vs.* Cx3cr1_chow_female	9	11
	hfd_cx Cx3cr1_hfd_male *vs.* Cx3cr1_hfd_female	7	5
Brain resident microglia cells *vs.* infiltrating cells	**hfd_cc_f** Cx3cr1_hfd_female *vs.* Ccr2_hfd_female	4036	3569
	hfd_cc_m Cx3cr1_hfd_male *vs.* Ccr2_hfd_male	3838	3350
Effect of the sex in infiltrating cells	**hfd_cc** Ccr2_hfd_male *vs.* Ccr2_hfd_female	1598	1676

Figures

Figure 1. CCR2+ cells infiltrate the hypothalamus of mice fed a high-fat diet. a) CX3CR1$^{GFP/+}$CCR2$^{RFP/+}$
dual-reporter mutant mice generation. b) Schematic representation of the experimental protocol for
analysis of HFD-induced CCR2+ peripheral-cell chemotaxis towards the hypothalamus. c) Flow cytometry
analysis of CX3CR1^{GFP+} and CCR2^{RFP+} cells in the white adipose tissue and in the hypothalamus of HFD-fed
mice. d) Measured event rate detected by flow cytometer of CX3CR1^{GFP+} and CCR2^{RFP+} cells isolated from
the hypothalamus of HFD-fed male and female mice. e) Coronal brain sections of mediobasal
hypothalamus (MBH) from chow- and 4 weeks HFD-fed mice CX3CR1$^{GFP/+}$CCR2$^{RFP/+}$. f) Coronal brain
sections of MBH from 4 weeks HFD-fed mice CX3CR1$^{GFP/+}$CCR2$^{RFP/+}$ immunostained for CD169
(Sialoadhesin). White arrows indicate overlap between CD169+ cells with CX3CR1+ cells or with CCR2+
cells. 3V = Third ventricle, Scale Bar = 50 μm.

Figure 2. CX3CR1+ resident and CCR2+ recruited cells sorted from the hypothalamus of HFD-fed mice express classical markers of microglia and other immune cells. a) Schematic representation of the experimental protocol for sorting and sequencing CX3CR1+ and CCR2+ cells from the hypothalamus of chow- and HFD-fed mice. b) Cx3cr1 gene expression and c) Ccr2 gene expression of CX3CR1+ and CCR2+ cells sorted from the hypothalamus of HFD-fed mice. Analysis of d-m) classical microglial markers and n-x) bone marrow-derived immune cell markers in the transcriptome of CX3CR1+ and CCR2+ cells sorted from the hypothalamus of HFD-fed mice. To perform RNA-sequencing we have employed a total of 200 mice of each sex fed on HFD and 100 mice of each sex fed on chow diet. They were divided in 5 independent experiments. In order to get total RNA amount from CCR2+ cells in the hypothalamus of HFD-fed mice that was enough for library construction and RNA-seq, CCR2+ samples were pooled together in 3 samples, but only 2 samples could be sequenced due the final RNA integrity and amount.

Figure 3. Differential gene expression (DGE) analysis of CX3CR1+ resident microglia and CCR2+
infiltrating cells sorted from the hypothalamus of HFD-fed mice show an enormous difference in their
transcriptomic signature. A-b) Venn diagram showing the number of DGEs for chow and HFD diet
comparison and sex comparison, respectively. c) Heatmap of up and downregulated DEGs when
comparing CX3CR1+ resident and CCR2+ infiltrating cells from HFD-fed female mice. d) Heatmap of up
and downregulated DEGs when comparing CX3CR1+ resident and CCR2+ infiltrating cells from HFD-fed
male mice. e) Heatmap of up and downregulated DEGs when comparing CCR2+ infiltrating cells from
HFD-fed male and female mice. hfd_cx_f = Cx3cr1_hfd_female vs. Cx3cr1_chow_female; hfd_cx_m =
Cx3cr1_hfd_male vs. Cx3cr1_chow_male; chow_cx = Cx3cr1_chow_male vs. Cx3cr1_chow_female;
hfd_cx = Cx3cr1_hfd_male vs. Cx3cr1_hfd_female; hfd_cc_f = Cx3cr1_hfd_female vs. Ccr2_hfd_female;
hfd_cc_m = Cx3cr1_hfd_male vs. Ccr2_hfd_male; hfd_cc = Ccr2_hfd_male vs. Ccr2_hfd_female.

Figure 4. Various differential gene expression (DGE) found in CCR2+ infiltrating cells from hypothalamus of HFD-fed mice belongs to chemotaxis pathways. a) KEGG enrichment analysis shows the distribution of DGEs in distinct metabolic pathways. b-c) Ontology analysis for DGEs related to chemotaxis from CCR2+ cells sorted from the hypothalamus of HFD-fed female and male mice, respectively. d) Heatmap of up and downregulated DEGs related to chemotaxis when comparing CX3CR1+ resident and CCR2+ infiltrating cells from HFD-fed male and female mice.

682 **Figure 5. CCR2+ infiltrating cells from the hypothalamus of HFD-fed mice express a broad of**
683 **chemokine receptors.** a-i) Chemokine receptors gene expression in the transcriptome of CX3CR1+ and
684 CCR2+ cells sorted from the hypothalamus of HFD-fed mice.

**Figure 6. CXCL10/interferon γ-induced protein 10 kDa (IP-10) is highly expressed in CCR2+ infiltrating
cells from the hypothalamus of HFD-fed mice.** a-c) Cxcl9, Cxcl10 and Ifng gene expression in the
transcriptome of CX3CR1+ and CCR2+ cells sorted from the hypothalamus of HFD-fed mice. d-e)
Ontology analysis for DGEs related to interferon signaling pathways from CCR2+ cells sorted from the
hypothalamus of HFD-fed female and male mice, respectively.

Figure 7. CXCL10 neutralization has a mild impact on reducing CCR2+ and CXCR3+ cell chemotaxis
towards the hypothalamus of HFD-fed mice. a) Schematic representation of the experimental protocol
for CXCL10 central neutralization. b) Coronal brain sections from 4 weeks HFD-fed CCR2[RFP] mice showing
the CCR2+ cells distribution in the hypothalamic parenchyma upon CXCL10 central neutralization. 3V =
Third ventricle, Scale Bars = 50 μm. c) Coronal brain sections from 4 weeks HFD-fed female CCR2[RFP] mice
immunostained for CXCR3 upon CXCL10 central neutralization. d) Coronal brain sections from 4 weeks
HFD-fed male CCR2[RFP] mice immunostained for CXCR3 upon CXCL10 central neutralization. 3V = Third
ventricle, Scale Bars = 25 μm. e-f) Hypothalamic mRNA levels of several chemokine receptors and
chemokines in HFD-fed female (light gray and purple bars) and male (light gray and green bars) CCR2[RFP]
mice upon CXCL10 central neutralization. For qualitative confocal image analysis, we have used 3
samples per group. For RT-qPCR of hypothalamus we have used 5-6 samples per group. Two-tailed
Mann-Whitney tests were used for statistical analyses. *p<0.05 and **p<0.01 in comparison with
respective Anti-IgG treated groups.

Figure 8. CXCL10 central neutralization in HFD-fed female mice. a) Percentual of body weight gain from Day 0 to Day 28 of the experimental protocol. b) Delta body weight during the experimental period. c) Brown adipose tissue weight and d) White adipose tissue (retroperitoneal depot) weight at Day 28. e) Leptin, f) Triglycerides and g) Total cholesterol plasma levels at Day 28. h) Weekly food intake measurement during experimental period. i) Cumulative food intake during the experimental period. j) 90 min food intake measurement after 16h-fasting. k) Hypothalamic mRNA levels of neuropeptides involved in food intake control. l) Intraperitoneal glucose tolerance test at Day 24. m) 6h-fasting blood glucose levels. n) Insulin plasma levels at Day 28. o) Hypothalamic mRNA levels of inflammatory genes. p) O_2 consumption; q) CO_2 production; r) Energy Expenditure and s) Respiratory Quotient at Day 24. Data were expressed as mean ± SEM of 8-10 mice per group (in two independent experiments). To perform qRT-PCR we have used 6 mice/group. To perform biochemical analysis in plasma we have used 8-10 mice/group. To perform ipGTT we have used 4 mice/group. To perform indirect calorimetry, we have used 4 mice/group. Two-way ANOVA following by Sidak's post-hoc test and Mann-Whitney test were used for statistical analyses. *$p < 0.05$, **$p < 001$, ***$p < 0.001$, ****$p < 0.0001$ in comparison with IgG treated group.

Figure 9. CXCL10 central neutralization in HFD-fed male mice. a) Percentual of body weight gain from Day 0 to Day 28 of the experimental protocol. b) Delta body weight during the experimental period. c) Brown adipose tissue weight and d) White adipose tissue (retroperitoneal depot) weight at Day 28. e) Leptin, f) Triglycerides and g) Total cholesterol plasma levels at Day 28. h) Weekly food intake measurement during experimental period. i) Cumulative food intake during the experimental period. j) 90 min food intake measurement after 16h-fasting. k) Hypothalamic mRNA levels of neuropeptides involved in food intake control. l) Intraperitoneal glucose tolerance test at Day 24. m) 6h-fasting blood glucose levels. n) Insulin plasma levels at Day 28. o) Hypothalamic mRNA levels of inflammatory genes. p) O_2 consumption; q) CO_2 production; r) Energy Expenditure and s) Respiratory Quotient at Day 24. Data were expressed as mean ± SEM of 8-10 mice per group (in two independent experiments). To perform qRT-PCR we have used 6 mice/group. To perform biochemical analysis in plasma we have used 8-10 mice/group. To perform ipGTT we have used 4 mice/group. To perform indirect calorimetry, we have used 4 mice/group. Two-way ANOVA following by Sidak's post-hoc test and Mann-Whitney test were used for statistical analyses. *$p < 0.05$, **$p < 001$, ***$p < 0.001$, ****$p < 0.0001$ in comparison with IgG treated group.

Figure 10. AMG487 treatment attenuates CCR2+ and CXCR3+ cell chemotaxis towards the
hypothalamus of HFD-fed mice. a) Schematic representation of the experimental protocol for CXCR3
systemic blockage. b) Coronal brain sections from 4 weeks HFD-fed CCR2[RFP] mice showing the CCR2+
cells distribution in the hypothalamic parenchyma upon AMG487 treatment. 3V = Third ventricle, Scale
Bars = 50 μm. c) Coronal brain sections from 4 weeks HFD-fed female CCR2[RFP] mice immunostained for
CXCR3 upon AMG487 treatment. d) Coronal brain sections from 4 weeks HFD-fed male CCR2[RFP] mice
immunostained for CXCR3 upon AMG487 treatment. 3V = Third ventricle, Scale Bars = 25 μm. e-f)
Hypothalamic mRNA levels of several chemokine receptors and chemokines in HFD-fed female (light gray
and pink bars) and male (light gray and blue bars) CCR2[RFP] mice upon AMG487 treatment. For qualitative
confocal image analysis, we have used 3 samples per group. For RT-qPCR of hypothalamus we have used
7-8 samples per group. Two-tailed Mann-Whitney tests were used for statistical analyses. *p<0.05 and
**p<0.01 in comparison with respective VEH treated groups.

Figure 11. CXCR3 systemic blockage in HFD-fed female mice. a) Percentual of body weight gain from Day 0 to Day 28 of the experimental protocol. b) Delta body weight during the experimental period. c) Brown adipose tissue weight and d) White adipose tissue (retroperitoneal depot) weight at Day 28. e) Leptin, f) Triglycerides and g) Total cholesterol plasma levels at Day 28. h) Weekly food intake measurement during experimental period. i) Cumulative food intake during the experimental period. j) 90 min food intake measurement after 16h-fasting. k) Hypothalamic mRNA levels of neuropeptides involved in food intake control. l) Intraperitoneal glucose tolerance test at Day 24. m) 6h-fasting blood glucose levels. n) Insulin plasma levels at Day 28. o) Hypothalamic mRNA levels of inflammatory genes. p) O_2 consumption; q) CO_2 production; r) Energy Expenditure and s) Respiratory Quotient at Day 24. Data were expressed as mean ± SEM of 14-16 mice per group (in four independent experiments). To perform qRT-PCR we have used 8 mice/group. To perform biochemical analysis in plasma we have used 8-10 mice/group. To perform ipGTT we have used 5 mice/group. To perform indirect calorimetry, we have used 4-5 mice/group. Two-way ANOVA following by Sidak's post-hoc test and Mann-Whitney test were used for statistical analyses. *$p<0.05$, **$p<001$, ***$p<0.001$, ****$p<0.0001$ in comparison with VEH treated group.

Figure 12. CXCR3 systemic blockage in HFD-fed male mice. a) Percentual of body weight gain from Day 0
to Day 28 of the experimental protocol. b) Delta body weight during the experimental period. c) Brown
adipose tissue weight and d) White adipose tissue (retroperitoneal depot) weight at Day 28. e) Leptin, f)
Triglycerides and g) Total cholesterol plasma levels at Day 28. h) Weekly food intake measurement during
experimental period. i) Cumulative food intake during the experimental period. j) 90 min food intake
measurement after 16h-fasting. k) Hypothalamic mRNA levels of neuropeptides involved in food intake
control. l) Intraperitoneal glucose tolerance test at Day 24. m) 6h-fasting blood glucose levels. n) Insulin
plasma levels at Day 28. o) Hypothalamic mRNA levels of inflammatory genes. p) O_2 consumption; q)
CO_2 production; r) Energy Expenditure and s) Respiratory Quotient at Day 24. Data were expressed as
mean ± SEM of 14-16 mice per group (in four independent experiments). To perform qRT-PCR we have
used 8 mice/group. To perform biochemical analysis in plasma we have used 8-10 mice/group. To
perform ipGTT we have used 5 mice/group. To perform indirect calorimetry, we have used 4-5
mice/group. Two-way ANOVA following by Sidak's post-hoc test and Mann-Whitney test were used for
statistical analyses. *p<0.05, **p<001, ***p<0.001, ****p<0.0001 in comparison with VEH treated group