MAIT Cells Modulate Innate Immune Cells and Inhibit Colon Cancer Growth

Olivia J. Cheng[1,2], Eric J. Lebish[3], Owen Jensen[1,2], Damian Jacenik[3,4], Shubhanshi Trivedi[2],

Jackson Cacioppo[6], Jeffrey Aubé[6], Ellen J. Beswick*[3,5], Daniel T. Leung*[1,2]

[1]Division of Microbiology & Immunology, Department of Pathology, University of Utah, Salt

Lake City, Utah, United States

[2]Division of Infectious Disease, Department of Internal Medicine, University of Utah, Salt Lake

City, Utah, United States

[3]Division of Gastroenterology, Department of Internal Medicine, University of Utah, Salt Lake

City, Utah, United States

[4]Department of Cytobiochemistry, Faculty of Biology and Environmental Protection, University

of Lodz, Lodz, Poland

[5]Division of Digestive Diseases and Nutrition, Department of Internal Medicine, University of

Kentucky, Lexington, Kentucky, United States

[6]Department of Chemistry, University of North Carolina at Chapel Hill, Chapel Hill, North

Carolina, United States

Corresponding Authors: Daniel T. Leung, MD

26 N Medical Drive Room 517, Salt Lake City, Utah 84132

+1 801-581-8812, Daniel.leung@utah.edu

The authors have declared that no conflict of interest exists.

*Co-senior authors

Funding

The study was funded by the National Institutes of Health (R01CA207051 to EJB, R01AI130378

to DTL, and K24AI166087 to DTL).

Abstract

Mucosal-associated invariant T (MAIT) cells are innate-like T cells that can be activated by

microbial antigens and cytokines and are abundant in mucosal tissues including the colon. MAIT

cells have cytotoxic and pro-inflammatory functions. Despite their potential for adoptive cell

therapy, studies into their anti-cancer activity, including their role in colon cancer, are limited.

We expanded MAIT cells *in vivo* and injected them into RAG1$^{-/-}$ mice with MC38-derived tumors

and tumor growth was assessed. Peritumoral injection of MAIT cells inhibits tumor growth

compared to control. Genomic and multiplex cytokine analyses show that tumors from the

MAIT cell-treated group have higher expression of markers for eosinophil-activating cytokines,

suggesting an association between eosinophil recruitment and tumor inhibition. Multiplex

cytokine and flow cytometry analyses of human peripheral leukocytes stimulated with MAIT

ligand show an increase in eotaxin-1 production and activation of eosinophils, associated with

increased cancer cell killing. We documented that MAIT cells have a protective role in a murine

colon cancer model, potentially involving novel eosinophil-associated mechanisms. This study

provides evidence of the immune modulating potential of MAIT cells within the tumor

microenvironment, leading to compositional and functional changes. Our results highlight the

potential of MAIT cells for colon cancer immunotherapy.

Brief summary

In models of colon cancer, MAIT cells have anti-tumor activity, associated with increased

production of proinflammatory and eosinophil-modulating cytokines.

Introduction

Mucosal-associated invariant T (MAIT) cells are innate-like T cells that recognize microbial-derived products in a TCR-dependent manner via MR-1, a highly conserved MHC class I-related molecule (1, 2). MAIT cells can be activated with the ligand 5-OP-RU, a microbial riboflavin metabolite, for specific stimulation (3, 4). MAIT cells can also be activated in a TCR-independent manner by cytokines, such as interleukins (IL)-7, IL-12, IL-15, and IL-18 (5-7), broadening the modes of activation and the target cell types. MAIT cells are known for their rapid effector function and cytotoxicity upon activation and have been shown to kill infected target cells (2, 8, 9). Additionally, MAIT cells have the capacity to interact with and modulate other immune cell types, including B cells, neutrophils, and monocytes (10-13). In humans, MAIT cells are abundant in the peripheral blood (14), as well as in mucosal tissues such as the liver (15), lung (16), and gut (17, 18). In mice, MAIT cell frequencies vary by strain and tissue type, but overall have lower frequencies than in humans (19, 20).

Due to the abundance of MAIT cells in various mucosal tissues, interest in MAIT cells has expanded beyond its anti-microbial functions, and recent studies have examined their potential role in mucosal tumors. Reports have suggested both pro- and anti-tumor activity of MAIT cells, with some studies demonstrating a modulating effect on Natural Killer (NK) cell anti-tumor activity (21-23). In addition, due to their lack of alloreactive potential and cytotoxic capacity, MAIT cells have been considered for use in immunotherapy (24-26). Evidently, the role of MAIT cells in cancer is complex and context-dependent, warranting further studies in various types of cancer, particularly their potential to interact with other immune cell types.

As of 2020, human colorectal cancer (CRC) represents about 10% of new cancer cases and

death worldwide (27). In addition to surgical resection and adjuvant chemotherapy, the use of

immunotherapies is available but limited in efficacy; thus, further understanding of the immune

cells involved could help advance therapy (28-30). In human CRC, studies have demonstrated

both positive and negative associations between tumor-infiltrating MAIT cells and overall

survival (31, 32). MAIT cells accumulate in CRC tumors (17, 31, 33) and tumor-infiltrating MAIT

cells display bacterial antigen recognition signatures (34), but the impact of these cells in the

tumor microenvironment is not well understood. Tumor-infiltrating MAIT cells retain their

cytotoxicity and production of cytokines (17, 35), but have impaired interferon- γ (IFN-γ)

production (17, 36) and high levels of exhaustion markers (33, 36). A multi-omics analysis of

CRC demonstrated an abundance of intratumor microbes and their immune modulation

potential on tumor-infiltrating immune cells, including MAIT cells (32). The prevalence of

intratumoral bacteria in CRC (32, 34) and the specificity of MAIT cells for microbial antigens

warrant further investigation into the role of MAIT cells in CRC.

Here, we show that peritumoral injection of purified MAIT cells inhibits MC38 cells-derived

tumor growth in a mouse model of colon cancer. We show that this tumor inhibition is

associated with changes in the immune cell profile and functional changes, including higher

levels of myeloid cell-modulating cytokines and chemokines. Culture of human peripheral

leukocytes with colon cancer cell line resulted in the activation of eosinophils upon MAIT cell

stimulation. Together, our data suggest a novel mechanism by which MAIT cells can modulate

innate immunity affecting eosinophils n a mixed immune cell environment and provide

protection in a murine model of colon cancer, highlighting the potential of MAIT cells to be a

candidate for CRC immunotherapy.

Results

Peritumoral injection of MAIT cells leads to inhibition of tumor growth in the MC38 model of

CRC

To test the effect of MAIT cells on colon cancer growth, MAIT cells were expanded in C57BL/6J

wildtype (WT) mice for 7 days, isolated (Supplemental Fig. 1), and were peritumorally injected

into RAG1$^{-/-}$ mice with MC38 cells-derived tumor 1 day after cancer cell injection (Fig. 1A). We

observed that the growth of MC38 cells-derived tumors was significantly lower ($p<0.05$) when

mice were treated with MAIT cells, as shown in the final tumor volume and weight at harvest

(Fig. 1B-E). The same effect was also observed when MAIT cells were injected with ligand 5-A-

RU and methylglyoxal (MGO) into both MC38-bearing RAG1$^{-/-}$ and C57BL/6J WT mice

(Supplemental Fig. 2). These data suggest that MAIT cells provide a protective role in this

murine subcutaneous model of colon cancer.

MAIT cell peritumoral injection leads to changes in the tumor microenvironment.

To examine potential functional and compositional changes induced by MAIT cell peritumoral

injection, RNA was isolated from extracted tumors and quantitative real-time reverse-

transcription PCR (qRT-PCR) was performed to evaluate the expression of apoptosis- and

oxidative stress-related genes relative to saline treated control group. We observed higher gene

expression of *Caspase3* (p < 0.01) *and Fas* (p < 0.05) in tumors treated with MAIT cells

(Supplemental Fig. 3A). Consistently, caspase 3/7 protein activity was detected to be

significantly higher in tumors in the MAIT cells-treated group compared to the control group

(p<0.01; Fig. 2A), supporting the tumor inhibition effect observed *in vivo*. Using gene markers

for various innate immune cells, we observed a non-significant increase in *Siglecf* and *Ccr3*

expression in tumors in mice from the MAIT treatment group compared to control, suggesting

an impact on eosinophils, while there was no evidence of changes in NK (*Klrb1c*) or macrophage

(Adgre1) cell markers (Supplemental Fig. 3B). Consistent with the increase in eosinophil

markers mentioned above, an increased expression of eotaxin-2 (also known as *Ccl24)* and

cytokine *Il5* (p = 0.052) was observed in the MAIT cells treatment group (Supplemental Fig 3C).

Overall, our results suggest that the treatment of MC38-bearing mice with MAIT cells leads to

increased cell death and eosinophil-activating cytokine.

To confirm functional changes at the protein level, we analyzed released molecules from the

extracted tumor. Tumor pieces (8 mg) were incubated in complete media for 18 hours and the

resulting supernatants were analyzed using Luminex multiplex cytokine and chemokine

analysis. We observed a significant increase in the production of various cytokines, including

the proinflammatory cytokines IFN-γ, and IL-17, and GM-CSF (p<0.05; Fig. 2B, C, and D), which

provides further evidence that the introduction of MAIT cells into MC38 cells-derived tumors

leads to the reprogramming of the tumor microenvironment. Consistent with the increase in

eosinophil gene markers mentioned above, a higher level of eotaxin-1 was detected in the

tumors extracted from mice in the MAIT treatment group (Fig. 2E). To determine the

plausibility that these cytokines could be produced by activated MAIT cells, we cultured sorted

mouse MAIT cells alone *in vitro* with a T cell activation cocktail. We confirmed significantly

higher levels of each of the above cytokines compared to unstimulated conditions

(Supplemental Fig. 4). Together, we observed functional changes of immune cells and

reprogramming of the tumor microenvironment associated with *in vivo* tumor inhibition.

Activation of human MAIT cells with a MAIT ligand enhances the killing of colon cancer cells

To model the effect of human MAIT cells on colon cancer cells in the presence of other immune

cells, we utilized human whole leukocytes co-cultured with the human colon cancer cell line

COLO 205 (Fig. 3A). Calcein release assay was used to examine cytotoxicity and culturing with

human whole leukocytes for 4 hours leads to specific lysis of COLO 205 cells (Fig. 3B). Activating

MAIT cells with the MAIT ligand, 5-A-RU and MGO, significantly increased the specific lysis

capacity of the leukocytes ($p<0.05$; Fig. 3B). Flow cytometry analysis also demonstrates

enhanced killing of COLO 205 by whole leukocytes with the addition of the MAIT ligand in a 16-

hour co-culture, with a significant reduction in the frequency of CFSE-labeled cells ($p<0.05$; Fig.

3C). Together, these data show that MAIT cell-specific activation with 5-A-RU enhances the

killing of colon cancer cells.

Stimulation of MAIT cells in human peripheral leukocytes induces eotaxin-1 production in

MAIT cells and activation of eosinophils in an MR-1 dependent manner

To determine the immune modulating potential of human MAIT cell in a mixed cell population,

we used human whole leukocyte cultures to examine the effect of MAIT cell activation in a cell

culture model of CRC. MAIT cells were selectively stimulated with 5-A-RU overnight and culture

supernatant was analyzed using multiplex cytokine and chemokine assay. In 5-A-RU stimulated

cultures, we found a higher level of Macrophage inflammatory protein (MIP)-1α, MIP-1β, and

Monocyte chemoattractant protein (MCP)-1 ($p < 0.05$ and $p < 0.01$; Fig. 4A), suggesting a capacity

of MAIT cells to modulate macrophages and monocytes and granulocytes (37-40). Consistent

with the recruitment of eosinophils with MAIT treatment in our murine model (Fig 2A), we

found that concentrations of IL-5, GM-CSF, and eotaxin-1 levels are significantly higher in

cultures stimulated with the MAIT ligand 5-A-RU/MGO compared to unstimulated leukocytes

($p < 0.05$ and $p < 0.01$; Fig. 4A). In addition, we observed that cytokines known to activate and

attract eosinophils, including IL-4 ($p < 0.01$) and IL-8 ($p = 0.0547$) (41, 42), are also significantly

higher in 5-A-RU/MGO stimulated cultures compared to control (Fig. 4A). Together, these data

demonstrate the immune modulating capacity of human MAIT cell activation in a mixed cell

population, specifically the potential to modulate eosinophils.

Previous studies have demonstrated the ability of eosinophils to kill COLO205 cells in a

granzyme A-dependent manner (43). To determine if MAIT cell activation can induce eosinophil

activation and cytotoxic function in a setting with mixed cell population of leukocytes, we

cultured human peripheral leukocytes with or without the MAIT ligand 5-A-RU/MGO overnight

and analyzed cytokine expression and levels of activation of MAIT cell and eosinophil using flow

cytometry (Supplemental Fig. 1). Ligand stimulated MAIT cells had significantly higher

expression of the activation marker CD69 ($p < 0.01$) and of eotaxin-1 ($p < 0.001$) (Fig. 4B).

Compared to unstimulated cultures, whole leukocyte cultures stimulated with the MAIT ligand

also shows a significant increase in eosinophil activation, marked by CD69 and granzyme A

expression (p<0.0001; Figure 4C). These data show that in peripheral leukocyte populations, the

addition of the MAIT ligand induces the expression of eosinophil-activating cytokines in MAIT

cells and correspondingly, eosinophil activation.

To confirm if eosinophil activation upon 5-A-RU/MGO stimulation is mediated via MR1 or a

broad consequence of antigen processing, anti-MR1 neutralizing antibody was added to the

human leukocyte cultures 1 hour before stimulation with 5-A-RU/MGO. In the presence of anti-

MR1, MAIT cell activation and the expression of eotaxin-1 are significant reduced ($p < 0.001$ and

$p < 0.0001$; Fig. 4B). Similarly, the activation and the induction of effector molecule granzyme A

in eosinophils induced by MAIT cell stimulation are significantly reduced in the presence of anti-

MR1 antibody ($p < 0.001$; Fig. 4C), suggesting that the effect of 5-A-RU/MGO on eosinophils is

dependent on MR1.

Together, these data show that the addition of 5-A-RU/MGO, a MAIT-specific ligand, leads to an

increased production of eosinophil-recruiting chemokines and cytokines by MAIT cells and

eosinophil activation, highlighting the ability of MAIT cells to modulate eosinophils in a MR1-

dependent manner.

Discussion

While MAIT cells are primarily known for their ability to kill microbially infected cells, other

potential roles have emerged. Specifically, due to their abundance at mucosal tissues, there has

been an increasing interest in the role of MAIT cells in mucosal tissue-associated cancers. MAIT

cells are a promising potential candidate for cancer immunotherapies: MAIT cells lack

alloreactive potential, thus having an advantage over the use of conventional T cells with HLA-

restrictions for the application of universal, non-donor-restrictive adoptive cell therapy (24).

MAIT cells also express higher levels of the multi-drug efflux protein MDR1, conferring higher

persistence against certain chemotherapy drugs compared to conventional CD8$^+$ T cells and

thus an ideal target for use in immunotherapy coupled with chemotherapy (25, 26, 44).

Additionally, the checkpoint inhibitor, anti-PD1, has been shown to enhance the cytotoxicity of

MAIT cells in patients with prostate cancer (45), basal and squamous cell carcinoma, and

hepatocellular carcinoma (46-48). Recent studies have demonstrated the ability to generate

chimeric antigen receptor (CAR)-MAIT cells, which retain their anti-tumor capacity even in the

presence of immunosuppressive tumor-associated macrophages (49, 50). Furthermore,

protocols for MAIT cell expansion have been established, including demonstration that, when

expanded with a human serum replacement, MAIT cells display enhanced functionality and low

levels of exhaustion markers, making them optimized for adoptive therapy (51, 52).

In this study, we provided evidence that MAIT cells have an anti-tumoral role both in an MC38

cell-derived allograft tumor model in RAG1$^{-/-}$ mice and against human colon cancer cells. We

used RAG deficient mice which are characterized by the lack of mature B and T cells to exclude

the potential impact of the presence of endogenous T cells. In the murine model of colon

cancer, we observed tumor inhibition, associated with a significant increase in the production

of anti-tumor cytokines, and evidence suggesting the recruitment of eosinophils with increases

in both gene and protein expression of eotaxin, IL-5, and SiglecF, among others. In human

leukocyte and human colon cancer cell co-culture, we observed enhanced levels of COLO 205

lysis after stimulation with MAIT ligand 5-A-RU/MGO, demonstrating their cytotoxic capacity

against colon cancer cells. In addition, MAIT cell activation in human whole leukocytes leads to

the production of a wide range of cytokines and chemokines, including eotaxin-1, GM-CSF, and

Mip-1, highlighting the potential for anti-tumor effects through their immune-modulating

properties (10). Particularly, we observed that MAIT cell activation resulted in an increase in the

production of IL-5 and eotaxin-1, and GM-CSF, and an increase in activation and effector

function of eosinophils, the effect of which was MR1-dependent.

We observed the capacity of MAIT cell to modulate innate cell populations in our models of

colon cancer. We found significantly higher levels of eotaxin-1, known to recruit and activate

eosinophils, in both mouse *in vivo* and human *in vitro* models. *In vitro* studies have

demonstrated that eosinophils have the ability to induce direct killing on colon cancer cells (43,

53), supported by an *in vivo* study of two colon cancer models showing that eosinophil

contributes to tumor suppression (53). Another study using a subcutaneous MC38 cells-derived

tumor model in WT mice has shown that eosinophil inversely correlates with tumor growth and

the observed tumor suppression was dependent on GM-CSF and eosinophil recruitment (54).

This is consistent with our experiment where we observed an association between MAIT cell-

mediated inhibition of MC38-derived tumors and an increase in tumor production of GM-CSF

and eotaxin-1 production in the tumors. With eosinophils' known ability to release effector

molecules like major basic protein (MBP) into the tumor microenvironment to induce cancer

cell lysis (55), these data support the involvement of eosinophils in MAIT-mediated tumor

suppression.

In addition to the production of eosinophil modulating effectors, we report MAIT cell

production of other pro-inflammatory and immune-modulating cytokines, GM-CSF, IFN-γ, and

IL-17. GM-CSF regulates the proliferation, differentiation, recruitment, and effector functions of

myeloid cell populations (11, 12, 56), and its addition of GM-CSF to human primary monocytes

inhibits COLO 205 proliferation (57). In murine colon cancer, neutrophils play a role in limiting

tumor progression, invasion and proliferation (58). We found higher GM-CSF in tumors from

the MAIT cell-injection group, suggesting a potential role for myeloid cell populations in

mediating the observed anti-tumor effect. We also found increased expression of IFN-γ in

MAIT-treated tumors. IFN-γ is a cytotoxic cytokine that can act directly on tumor cells to induce

tumor elimination (59-62). Specifically, IFN-γ inhibits MC38-derived tumor growth in mice (63)

and COLO 205 in vitro (64). In addition to the direct effects on tumor cells, IFN-γ also has

immune modulating effects on innate immune cells, including the enhancement of the anti-

tumor activity of eosinophils against colon cancer cell lines, including MC38 (53). Finally, our

demonstration of increased IL-13 and IL-17 in MAIT-treated tumors is consistent with studies

showing their production by MAIT cells (26, 65, 66). In line with our murine model of colon

cancer, a prior study showed that treatment with recombinant IL-17 in MC38-bearing mice

results in tumor inhibition (63).

Taken together, our study employing both murine model and human cell cultures demonstrates

a protective role of MAIT cells in colon cancer. While further analysis of immune cell activity in

the tumor microenvironment is needed to fully understand the interaction between MAIT cells

and other immune cells, our findings suggest a novel role of MAIT cells in recruitment and

activation of eosinophils. The mechanism by which MAIT cell mediates MC38-derived tumor

inhibition, particularly the role of eosinophils, needs further study, including how MAIT cells

could be harnessed in future therapeutic applications against colon cancer.

Methods and Materials

Mouse and cell lines

CL57BL/6 wild type and B6.129S7-$Rag1^{tm1Mom}$/J ($Rag1^{-/-}$) mice were purchased from the

Jackson Laboratory (Bar Harbor, MA, USA). Mice were housed at the University of Utah CMC

Animal Facility on a 12h – 12h light-dark cycle and were provided with water and normal rodent

chow ad libitum. Mouse and human cell lines, MC38 and Colo205, were purchased from the

American Type Culture Collection (ATCC, Manassas, Virginia, United States) and maintained in

RPMI complete media supplemented with 10% fetal bovine serum (FBS, Thermo Fisher

Scientific, Waltham, MA, USA), 1% penicillin/streptomycin cocktail (Corning, Tewksbury, MA,

USA) and 25mM HEPES (Thermo Fisher Scientific, Waltham, MA, USA).

In vivo MAIT cell expansion and isolation

On day 0, C57BL/6J mice were anesthetized with isoflurane and intranasally inoculated with a

50 mL solution (25 mL per nare) of PBS containing 200 nmol 5-A-RU (67), 50 mM methylglyoxal

(Sigma-Aldrich, St. Louis, MO, USA), and 16.67 mg Pam2CSK4 (InvivoGen, San Diego, CA, USA).

The reaction of methylglyoxal with 5-A-RU results in the formation of the MAIT-activating ligand

5-(2-oxopropylideneamino)-6-D-ribitylaminouracil (5-OP-RU) (68). On days 1, 2 and 4, mice

were again inoculated with 200 nmol 5-A-RU and 50 mM methylglyoxal. On day 7, mice were

euthanized, lungs were perfused with 5 mL of PBS, and then single-cell lung suspensions were

prepared using the gentleMACS Mouse Lung Dissociation Kit (Miltenyi Biotech, Bergisch

Gladbach, North Rhine-Westphalia, Germany) according to the manufacturer's protocol. Lung

suspensions were then pooled into male and female groups and stained with fluorochrome-

conjugated antibodies for MAIT isolation using fluorescent-associated cell sorting (FACS). Lung

suspensions were first incubated with Fixable Viability Dye eFluor 780 (BD Biosciences, Franklin

Lakes, NJ, USA) and then blocked with anti-mouse CD16/CD32 Fc block (Biolegend, San Diego,

CA, USA) and unlabeled MR1-6-formylpterin-tetramer (6-FP) (NIH Tetramer Core Facility at

Emory University, Atlanta, GA, USA) to minimize non-specific binding. Suspensions were then

stained for 30 minutes at room temperature with the MAIT cell TCR binding MR1-5-OP-RU-

tetramer (NIH Tetramer Core Facility at Emory University, Atlanta, GA, USA) and the following

antibodies: anti-CD3-PE-Dazzle594, anti-TCRb-BV421, anti-CD4-FITC, anti-CD8-APC, anti-TCRgd-

PE-Cy7, anti-CD44-BV650, and anti-CD45R-PE-Cy5 (Supp. Table 1). MAIT cells were gated as

single cell lymphocyte, live, B220-, TCR γδ-, CD44^{hi}, CD3+, TCRβ+ and MR1-5OPRU+. Following

FACS, sorted MAIT cells were washed in sterile PBS and resuspended at 1×10^6 cells per ml for

use for peritumoral injection in the MC38 cancer model.

MAIT cell peritumoral injection in MC38 mouse model.

In Rag1-/- mice, 1×10^6 MC38 cells were injected subcutaneously on Day 0. 1×10^5 FACS-sorted,

sex-matched MAIT cells were then injected peritumorally on Day 1. For control group, mice

were injected with an equivalent volume of PBS. Tumor growth was monitored until Day 15-21

and the mice were sacrificed to obtain the tumors. Extracted tumors were weighed at the end

of the experiment.

Mouse MAIT cell in vitro stimulation

Expanded lung MAIT cells from C57BL/6J donor mice were cultured in vitro at 50,000 cells/well.

Cells were either left unstimulated or stimulated with 1X eBioscience Cell Stimulation Cocktail

(500X, Invitrogen, Thermo Fisher Scientific, Waltham, MA, USA) in 100uL complete RPMI

(Gibco, Thermo Fisher Scientific, Waltham, MA, USA, supplemented with 10% FBS, 1%

penicillin/streptomycin, 1mM HEPES). After overnight culture, supernatants were analyzed

using MILLIPLEX MAP Mouse Cytokine/Chemokine Magnetic Bead Panel - Premixed 25 Plex -

Immunology Multiplex Assay (EMD Millipore Corporation, Billerica, MA, USA). and custom FAS

panel on the Luminex MAGPIX.

Detection of tumor-released cytokines

8 mg (± 0.5mg) of each mouse tumor was incubated in 500 µL of complete RMPI medium for 18

hours and 25 µL of the resulting supernatant was analyzed using Luminex bead assay for

various soluble markers using MILLIPLEX MAP Mouse Cytokine/Chemokine Magnetic Bead

Panel - Premixed 25 Plex - Immunology Multiplex Assay (EMD Millipore Corporation, Billerica,

MA, USA). and custom FAS panel on the Luminex MAGPIX. Measurements below detectable

limits were considered as 0.

Caspase activity measurement

Apo-ONE® Homogeneous Caspase-3/7 Assay (Promega Corporation, Madison, WI, USA) was

used to determine the caspase activity in the extracted tumors. 8 mg tumor pieces were

homogenized in Apo-ONE® Caspase-3/7 Reagent using a bead beater. After 1 freeze-thaw to

ensure lysis, samples were incubated for 30 minutes on a shaker and analyzed according to

manufacturer's protocol. Caspase 3/7 activity is measured as arbitrary fluorescent unit (AFU)

using a SpectraMax Mini plate reader (Molecular Devices, San Jose, CA, USA).

RNA isolation and qRT-PCR

RNA was extracted from tumor pieces stored in RNAlater or flash-frozen using Trizol (Thermo

Fisher Scientific, Waltham, MA, USA). Extracted RNA was converted to cDNA using High-

Capacity RNA-to-cDNA™ Kit (Applied Biosystems, Thermo Fisher Scientific, Waltham, MA, USA)

according to the manufacturer's protocol. qRT-PCR was performed using TaqMan Real Time

PCR Assays (Thermo Fisher Scientific, Waltham, MA, USA) with comparative Ct (ΔΔCt) analysis

on QuantStudio5.

Human whole leukocyte preparation and culture

Human whole leukocytes from anonymous healthy donors were isolated from apheresis cones

from ARUP using RBC sedimentation with HetaSep (STEMCELL Technologies, Vancouver, British

Columbia). Freshly isolated whole leukocytes were used fresh for *in vitro* culture.

Whole leukocytes were cultured with or without 5-A-RU (50 nM) and MGO (50 µM) in 200 µL

RPMI complete media and incubated at 37 °C overnight. CD69, and eotaxin, granzyme A, and in

MAIT cells and Eosinophils were analyzed using flow cytometry. To detect released cytokines in

the medium, 25 µL of supernatant was analyzed using Cytokine 35-Plex Human Panel for the

Luminex™ platform (Invitrogen, Thermo Fisher Scientific, Waltham, MA, USA) on the Luminex

MAGPIX. To block interactions activation of MAIT cell via MR1, anti-MR1 (Clone 26.5, Biolegend,

San Diego, CA, USA) or an isotype control was used at 10 µg/ml.

COLO 205 Killing Assays

For Calcein Release Assay, 50,000 Calcein-labeled (2 µM, Biolegend, San Diego, CA, USA) COLO

205 cells were co-cultured with 500,000 human whole leukocytes for 4 hours with or without 5-

A-RU (50 nM) and MGO (50 µM) in 200 µL RPMI complete media at 37 °C. Maximum release

was set up with 2% Triton X-100 (Acros Organics, Thermo Fisher Scientific, Waltham, MA, USA).

All samples and controls were set up in triplicates. Cultures were gently mixed and centrifuged,

and 100 µL of culture supernatant was transferred to a new plate and analyzed for fluorescence

intensity using BioTek Synergy H1 Multimode Reader (BioTek Instruments, Inc, Winooski, VT,

USA). Percent Specific Lysis was expressed as a percent relative to maximum release control,

calculated as [(Fluorescence $_{Sample}$ – Fluorescence $_{Untreated}$)/ (Fluorescence $_{max}$ – Fluorescence

_{Untreated})] x 100. For flow cytometry-based killing assay, 100,000 CFSE-stained (Immunochemistry

Technologies, Davis, California, CA, USA) COLO 205 was co-cultured with 10^6 human whole

leukocytes for 16 hours and the frequency of CFSE+ cells was analyzed.

Flow Cytometry and FACS

Isolated mouse lung cells were stained and MAIT cells were sorted using FACS Aria III Cell Sorter

(BD Bioscience). Stained human *in vitro* samples were analyzed using Cytek Aurora (Cytek

Biosciences, Fremont, CA, USA). FCS files were analyzed using FlowJo^{TM} v10.8 Software (BD Life

Sciences, Franklin Lakes, NJ, USA). MAIT cells were defined from the lymphocyte as live singlet,

CD45$^+$, CD3$^+$, MR1-5-OP-RU-Tetramer$^+$, Vα7.2$^+$ cells. Eosinophils were gated from the

granulocyte population as live singlet, CD45$^+$, CD3$^-$, CD44$^+$, Siglec8$^+$ cells. Gating strategies are

included in Supplemental Figure 1.

Antibodies used for FACS and Flow Cytometry

List of antibodies used in FACS of mouse MAIT cells.

Antibody/Viability dye	Clone	Vendor
eBioscience Fixable Viability Dye eFluor™ 780	NA	Invitrogen, Thermo Fisher Scientific
Anti-CD45R/B220	RA3-6B2	Biolegend
Anti-gd	GL3	Biolegend

Anti-CD3	17A2	Biolegend
Anti-CD44	IM7	Biolegend
MR1-5-OP-RU	NA	NIH Tetramer Core
Anti-TCRb	H57-597	Biolegend

389   List of antibodies used in flow cytometry analysis of human leukocyte cultures.

Antibody/Tetramer	Clone	Vendor
Zombie UV	NA	Biolegend
Anti-CD45	HI30	Biolegend
Anti-CD45RO	UCHL1	BD Biosciences
Anti-CD3	SK7	BD Biosciences
Anti-CD4	SK3	BD Biosciences
Anti-CD8	RPA-T8	Biolegend
MR1-5OPRU	NA	NIH Tetramer Core
Anti-Vα7.2	3C10	Biolegend
Anti-CD161	HP-3G10	Biolegend
Anti-CD25	BC96	Biolegend
Anti-CD69	FN50	BD Biosciences
Anti-CD44	IM7	Biolegend
Anti-Siglec8	7C9	Biolegend
Anti-Granzyme A	CB9	Biolegend

Anti-Eotaxin-1	IC3201U	R&D Systems

Statistical Analysis

Statistical analysis was performed using GraphPad Prism version 8 for Mac (GraphPad Software,

Boston, Massachusetts USA). Two-tailed Mann-Whitney U test was used to compare tumor

weights between groups, and Wilcoxon ranked t test was used to compare variables between

treatment groups in experiment involving paired human samples.

Study Approval

Mouse studies were approved by the University of Utah IACUC, protocol numbers 21-09-16.

The protocol to use human whole leukocytes from anonymous blood donors was reviewed by

the Institutional Review Board of the University of Utah and determined to be exempt

(IRB_00097389)

Author Contributions

OJC, EJB, and DTL designed the experiments. OJC, EJL, OJ, DJ, and ST performed the

experiments. OJC analyzed the data. EJB and DTL supervised the work. OJC, OJ, EJB and DTL

wrote the paper. EJB and DTL obtained funding. All authors reviewed and approved the final

manuscript.

Acknowledgement

The research was supported by the National Institutes of Health (R01CA207051 to EJB,

R01AI130378 to DTL, and K24AI166087 to DTL). We would like to thank the University of Utah

Flow Cytometry Core.

Reference

1.   Treiner E, Duban L, Bahram S, Radosavljevic R, Wanner V, Tilloy F, Affaticati P, Gilfillan S,

Lantz O. Selection of evolutionarily conserved mucosal-associated invariant T cells by MR1.

Nature. 2003;422.

2.   Le Bourhis L, Martin E, Peguillet I, Guihot A, Froux N, Core M, Levy E, Dusseaux M,

Meyssonnier V, Premel V, Ngo C, Riteau B, Duban L, Robert D, Huang S, Rottman M, Soudais C,

Lantz O. Antimicrobial activity of mucosal-associated invariant T cells. Nat Immunol.

2010;11(8):701-8.

3.   Mak JYW, Liu L, Fairlie DP. Chemical Modulators of Mucosal Associated Invariant T Cells.

Acc Chem Res. 2021;54(17):3462-75.

4.   Hagel J, Garner L, Bilton M, Mehta H, Leng T, Hackstein C, Phalora P, Amini A, Akther H,

Provine N, Edmans M, Willberg C, Klenerman P. Human MAIT Cell Activation. Methods in

Moelcular Biology. 2020;2098:97-124.

5.   Ussher JE, Bilton M, Attwod E, Shadwell J, Richardson R, de Lara C, Mettke E, Kurioka A,

Hansen TH, Klenerman P, Willberg CB. CD161++ CD8+ T cells, including the MAIT cell subset, are

specifically activated by IL-12+IL-18 in a TCR-independent manner. Eur J Immunol.

2014;44(1):195-203.

6.   Leeansyah E, Svard J, Dias J, Buggert M, Nystrom J, Quigley MF, Moll M, Sonnerborg A,

Nowak P, Sandberg JK. Arming of MAIT Cell Cytolytic Antimicrobial Activity Is Induced by IL-7

and Defective in HIV-1 Infection. PLoS Pathog. 2015;11(8):e1005072.

7.   Rha MS, Han JW, Kim JH, Koh JY, Park HJ, Kim SI, Kim MS, Lee JG, Lee HW, Lee DH, Kim

W, Park JY, Joo DJ, Park SH, Shin EC. Human liver CD8(+) MAIT cells exert TCR/MR1-independent

innate-like cytotoxicity in response to IL-15. J Hepatol. 2020;73(3):640-50.

8.   Lamichhane R, Schneider M, de la Harpe SM, Harrop TWR, Hannaway RF, Dearden PK,

Kirman JR, Tyndall JDA, Vernall AJ, Ussher JE. TCR- or Cytokine-Activated CD8(+) Mucosal-

Associated Invariant T Cells Are Rapid Polyfunctional Effectors That Can Coordinate Immune

Responses. Cell Rep. 2019;28(12):3061-76 e5.

9.   van Wilgenburg B, Scherwitzl I, Hutchinson EC, Leng T, Kurioka A, Kulicke C, de Lara C,

Cole S, Vasanawathana S, Limpitikul W, Malasit P, Young D, Denney L, consortium S-H, Moore

MD, Fabris P, Giordani MT, Oo YH, Laidlaw SM, Dustin LB, Ho LP, Thompson FM, Ramamurthy

N, Mongkolsapaya J, Willberg CB, Screaton GR, Klenerman P. MAIT cells are activated during

human viral infections. Nat Commun. 2016;7:11653.

10.   Ioannidis M, Cerundolo V, Salio M. The Immune Modulating Properties of Mucosal-

Associated Invariant T Cells. Front Immunol. 2020;11:1556.

11.   Davey MS, Morgan MP, Liuzzi AR, Tyler CJ, Khan MWA, Szakmany T, Hall JE, Moser B,

Eberl M. Microbe-specific unconventional T cells induce human neutrophil differentiation into

antigen cross-presenting cells. J Immunol. 2014;193(7):3704-16.

12.   Meierovics AI, Cowley SC. MAIT cells promote inflammatory monocyte differentiation

into dendritic cells during pulmonary intracellular infection. J Exp Med. 2016;213(12):2793-809.

13.   Bennett MS, Trivedi S, Iyer AS, Hale JS, Leung DT. Human mucosal-associated invariant T

(MAIT) cells possess capacity for B cell help. J Leukoc Biol. 2017;102(5):1261-9.

14.   Gherardin NA, Souter MN, Koay HF, Mangas KM, Seemann T, Stinear TP, Eckle SB,

Berzins SP, d'Udekem Y, Konstantinov IE, Fairlie DP, Ritchie DS, Neeson PJ, Pellicci DG, Uldrich

AP, McCluskey J, Godfrey DI. Human blood MAIT cell subsets defined using MR1 tetramers.

Immunol Cell Biol. 2018;96(5):507-25.

15.   Tang XZ, Jo J, Tan AT, Sandalova E, Chia A, Tan KC, Lee KH, Gehring AJ, De Libero G,

Bertoletti A. IL-7 licenses activation of human liver intrasinusoidal mucosal-associated invariant

T cells. J Immunol. 2013;190(7):3142-52.

16.   Khuzwayo S, Mthembu M, Meermeier EW, Prakadan SM, Kazer SW, Bassett T,

Nyamande K, Khan DF, Maharaj P, Mitha M, Suleman M, Mhlane Z, Ramjit D, Karim F, Shalek

AK, Lewinsohn DM, Ndung'u T, Wong EB. MR1-Restricted MAIT Cells From The Human Lung

Mucosal Surface Have Distinct Phenotypic, Functional, and Transcriptomic Features That Are

Preserved in HIV Infection. Front Immunol. 2021;12:631410.

17.   Sundstrom P, Ahlmanner F, Akeus P, Sundquist M, Alsen S, Yrlid U, Borjesson L, Sjoling A,

Gustavsson B, Wong SB, Quiding-Jarbrink M. Human Mucosa-Associated Invariant T Cells

Accumulate in Colon Adenocarcinomas but Produce Reduced Amounts of IFN-gamma. J

Immunol. 2015;195(7):3472-81.

18.   Booth JS, Salerno-Goncalves R, Blanchard TG, Patil SA, Kader HA, Safta AM, Morningstar

LM, Czinn SJ, Greenwald BD, Sztein MB. Mucosal-Associated Invariant T Cells in the Human

Gastric Mucosa and Blood: Role in Helicobacter pylori Infection. Front Immunol. 2015;6:466.

19.   Rahimpour A, Koay HF, Enders A, Clanchy R, Eckle SB, Meehan B, Chen Z, Whittle B, Liu

L, Fairlie DP, Goodnow CC, McCluskey J, Rossjohn J, Uldrich AP, Pellicci DG, Godfrey DI.

Identification of phenotypically and functionally heterogeneous mouse mucosal-associated

invariant T cells using MR1 tetramers. J Exp Med. 2015;212(7):1095-108.

20. Kurioka A, Walker LJ, Klenerman P, Willberg CB. MAIT cells: new guardians of the liver.

Clin Transl Immunology. 2016;5(8):e98.

21. Yan J, Allen S, McDonald E, Das I, Mak JYW, Liu L, Fairlie DP, Meehan BS, Chen Z, Corbett

AJ, Varelias A, Smyth MJ, Teng MWL. MAIT Cells Promote Tumor Initiation, Growth, and

Metastases via Tumor MR1. Cancer Discov. 2020;10(1):124-41.

22. Ruf B, Catania VV, Wabitsch S, Ma C, Diggs LP, Zhang Q, Heinrich B, Subramanyam V, Cui

LL, Pouzolles M, Evans CN, Chari R, Sakai S, Oh S, Barry CE, 3rd, Barber DL, Greten TF. Activating

Mucosal-Associated Invariant T Cells Induces a Broad Antitumor Response. Cancer Immunol

Res. 2021;9(9):1024-34.

23. Petley EV, Koay HF, Henderson MA, Sek K, Todd KL, Keam SP, Lai J, House IG, Li J,

Zethoven M, Chen AXY, Oliver AJ, Michie J, Freeman AJ, Giuffrida L, Chan JD, Pizzolla A, Mak

JYW, McCulloch TR, Souza-Fonseca-Guimaraes F, Kearney CJ, Millen R, Ramsay RG, Huntington

ND, McCluskey J, Oliaro J, Fairlie DP, Neeson PJ, Godfrey DI, Beavis PA, Darcy PK. MAIT cells

regulate NK cell-mediated tumor immunity. Nat Commun. 2021;12(1):4746.

24. Tourret M, Talvard-Balland N, Lambert M, Ben Youssef G, Chevalier MF, Bohineust A,

Yvorra T, Morin F, Azarnoush S, Lantz O, Dalle JH, Caillat-Zucman S. Human MAIT cells are

devoid of alloreactive potential: prompting their use as universal cells for adoptive immune

therapy. J Immunother Cancer. 2021;9(10).

25.   Fergusson JR, Ussher JE, Kurioka A, Klenerman P, Walker LJ. High MDR-1 expression by

MAIT cells confers resistance to cytotoxic but not immunosuppressive MDR-1 substrates. Clin

Exp Immunol. 2018;194(2):180-91.

26.   Dusseaux M, Martin E, Serriari N, Peguillet I, Premel V, Louis D, Milder M, Le Bourhis L,

Soudais C, Treiner E, Lantz O. Human MAIT cells are xenobiotic-resistant, tissue-targeted,

CD161hi IL-17-secreting T cells. Blood. 2011;117(4):1250-9.

27.   Morgan E, Arnold M, Gini A, Lorenzoni V, Cabasag CJ, Laversanne M, Vignat J, Ferlay J,

Murphy N, Bray F. Global burden of colorectal cancer in 2020 and 2040: incidence and mortality

estimates from GLOBOCAN. Gut. 2023;72(2):338-44.

28.   Golshani GaZ, Y. Advances in immunotherapy for colorectal cancer-a review.

Therapeutic Advances in Gastroenterology. 2020;13:1-11.

29.   Elez E, Baraibar I. Immunotherapy in colorectal cancer: an unmet need deserving of

change. Lancet Oncol. 2022;23(7):830-1.

30.   Carlsen L, Huntington KE, El-Deiry WS. Immunotherapy for Colorectal Cancer:

Mechanisms and Predictive Biomarkers. Cancers (Basel). 2022;14(4).

31.   Zabijak L, Attencourt C, Guignant C, Chatelain D, Marcelo P, Marolleau JP, Treiner E.

Increased tumor infiltration by mucosal-associated invariant T cells correlates with poor survival

in colorectal cancer patients. Cancer Immunol Immunother. 2015;64(12):1601-8.

32.   Liu Z, Zhang X, Zhang H, Zhang H, Yi Z, Zhang Q, Liu Q, Liu X. Multi-Omics Analysis

Reveals Intratumor Microbes as Immunomodulators in Colorectal Cancer. Microbiology

Spectrum. 2023;11.

33.   Mo J, Zheng L, Gao Z, Wu J, Bao Y. The Study of Mucosal-Associated Invariant T Cells in

Colon Cancer and Roles in Immune Activities. Onco Targets Ther. 2021;14:5263-73.

34.   Li S, Simoni Y, Becht E, Loh CY, Li N, Lachance D, Koo SL, Lim TP, Tan EKW, Mathew R,

Nguyen A, Golovato J, Berkson JD, Prlic M, Lee B, Minot SS, Nagarajan N, Dey N, Tan DSW, Tan

IB, Newell EW. Human Tumor-Infiltrating MAIT Cells Display Hallmarks of Bacterial Antigen

Recognition in Colorectal Cancer. Cell Rep Med. 2020;1(3):100039.

35.   Sundström PS, L.; Ahlmanner, F.;Sundquist, M.; Wong, J.; Lindskog, E.; Gustafsson, B.;

Quiding-Järbrink, M. Tumor-infiltrating mucosal-associated invariant T (MAIT) cells retain

expression of cytotoxic effector molecules. Oncotarget. 2019;10.

36.   Rodin W, Sundstrom P, Ahlmanner F, Szeponik L, Zajt KK, Wettergren Y, Bexe Lindskog E,

Quiding Jarbrink M. Exhaustion in tumor-infiltrating Mucosal-Associated Invariant T (MAIT) cells

from colon cancer patients. Cancer Immunol Immunother. 2021;70(12):3461-75.

37.   Zheng Y, Wang Z, Wei S, Liu Z, Chen G. Epigenetic silencing of chemokine CCL2 represses

macrophage infiltration to potentiate tumor development in small cell lung cancer. Cancer Lett.

2021;499:148-63.

38.   Dorner BG, Scheffold A, Rolph MS, Huser MB, Kaufmann SH, Radbruch A, Flesch IE,

Kroczek RA. MIP-1alpha, MIP-1beta, RANTES, and ATAC/lymphotactin function together with

IFN-gamma as type 1 cytokines. Proc Natl Acad Sci U S A. 2002;99(9):6181-6.

39.   Ramos CD, Canetti C, Souto JT, Silva JS, Hogaboam CM, Ferreira SH, Cunha FQ. MIP-

1alpha[CCL3] acting on the CCR1 receptor mediates neutrophil migration in immune

inflammation via sequential release of TNF-alpha and LTB4. J Leukoc Biol. 2005;78(1):167-77.

40.   Oliveira SHP, Lira S, Martinez-A C, Wiekowski M, Sullivan L, Lukacs NW. Increased

responsiveness of murine eosinophils to MIP-1β (CCL4) and TCA-3 (CCL1) is mediated by their

specific receptors, CCR5 and CCR8. Journal of Leukocyte Biology. 2002;71(6):1019-25.

41.   W. L, T. M, H. M, Virchow J J. Synergistic effects of interleukin-4 or interleukin-13 and

tumor necrosis factor-alpha on eosinophil activation in vitro. Am J Respir Cell Mol Biol. 1999;20.

42.   Erger R, TB. C. Interleukin-8 is a potent mediator of eosinophil chemotaxis through

endothelium and epithelium. Am J Physiol. 1995;268.

43.   Legrand F, Driss V, Delbeke M, Loiseau S, Hermann E, Dombrowicz D, Capron M. Human

eosinophils exert TNF-alpha and granzyme A-mediated tumoricidal activity toward colon

carcinoma cells. J Immunol. 2010;185(12):7443-51.

44.   Li YR, Zhou K, Wilson M, Kramer A, Zhu Y, Dawson N, Yang L. Mucosal-associated

invariant T cells for cancer immunotherapy. Mol Ther. 2023;31(3):631-46.

45.   Jarvis EM, Collings S, Authier-Hall A, Dasyam N, Luey B, Nacey J, Painter GF, Delahunt B,

Hermans IF, Weinkove R. Mucosal-Associated Invariant T (MAIT) Cell Dysfunction and PD-1

Expression in Prostate Cancer: Implications for Immunotherapy. Front Immunol.

2021;12:748741.

46.   Yao T, Shooshtari P, Haeryfar SMM. Leveraging Public Single-Cell and Bulk

Transcriptomic Datasets to Delineate MAIT Cell Roles and Phenotypic Characteristics in Human

Malignancies. Frontiers in Immunology. 2020;11.

47.   Yost KE, Satpathy AT, Wells DK, Qi Y, Wang C, Kageyama R, McNamara KL, Granja JM,

Sarin KY, Brown RA, Gupta RK, Curtis C, Bucktrout SL, Davis MM, Chang ALS, Chang HY. Clonal

replacement of tumor-specific T cells following PD-1 blockade. Nature Medicine.

2019;25(8):1251-9.

48.   Ruf B, Bruhns M, Babaei S, Kedei N, Ma L, Revsine M, Benmebarek MR, Ma C, Heinrich B,

Subramanyam V, Qi J, Wabitsch S, Green BL, Bauer KC, Myojin Y, Greten LT, McCallen JD, Huang

P, Trehan R, Wang X, Nur A, Murphy Soika DQ, Pouzolles M, Evans CN, Chari R, Kleiner DE,

Telford W, Dadkhah K, Ruchinskas A, Stovroff MK, Kang J, Oza K, Ruchirawat M, Kroemer A,

Wang XW, Claassen M, Korangy F, Greten TF. Tumor-associated macrophages trigger MAIT cell

dysfunction at the HCC invasive margin. Cell. 2023;186(17):3686-705 e32.

49.   Li YR, Brown J, Yu Y, Lee D, Zhou K, Dunn ZS, Hon R, Wilson M, Kramer A, Zhu Y, Fang Y,

Yang L. Targeting Immunosuppressive Tumor-Associated Macrophages Using Innate T Cells for

Enhanced Antitumor Reactivity. Cancers (Basel). 2022;14(11).

50.   Dogan M, Karhan E, Kozhaya L, Placek L, Chen X, Yigit M, Unutmaz D. Engineering

Human MAIT Cells with Chimeric Antigen Receptors for Cancer Immunotherapy. J Immunol.

2022;209(8):1523-31.

51.   Labuz D, Cacioppo J, Li K, Aube J, Leung DT. Enhancing Mucosal-Associated Invariant T

Cell Function and Expansion with Human Selective Serum. Immunohorizons. 2023;7(1):116-24.

52.   Parrot T, Healy K, Boulouis C, Sobkowiak MJ, Leeansyah E, Aleman S, Bertoletti A,

Sallberg Chen M, Sandberg JK. Expansion of donor-unrestricted MAIT cells with enhanced

cytolytic function suitable for TCR redirection. JCI Insight. 2021;6(5).

53.   Reichman H, Itan M, Rozenberg P, Yarmolovski T, Brazowski E, Varol C, Gluck N, Shapira

S, Arber N, Qimron U, Karo-Atar D, Lee JJ, Munitz A. Activated Eosinophils Exert Antitumorigenic

Activities in Colorectal Cancer. Cancer Immunol Res. 2019;7(3):388-400.

54.   Arnold IC, Artola-Boran M, Gurtner A, Bertram K, Bauer M, Frangez Z, Becher B, Kopf M,

Yousefi S, Simon HU, Tzankov A, Muller A. The GM-CSF-IRF5 signaling axis in eosinophils

promotes antitumor immunity through activation of type 1 T cell responses. J Exp Med.

2020;217(12).

55.   Robinson I, Santa Lucia G, Li A, Oberholtzer N, Plante J, Quinn KM, Reuben D, Mehrotra

S, Valdebran M. Eosinophils and melanoma: Implications for immunotherapy. Pigment Cell

Melanoma Res. 2022;35(2):192-202.

56.   Ushach I, Zlotnik A. Biological role of granulocyte macrophage colony-stimulating factor

(GM-CSF) and macrophage colony-stimulating factor (M-CSF) on cells of the myeloid lineage. J

Leukoc Biol. 2016;100(3):481-9.

57.   Nebiker CA, Han J, Eppenberger-Castori S, Iezzi G, Hirt C, Amicarella F, Cremonesi E,

Huber X, Padovan E, Angrisani B, Droeser RA, Rosso R, Bolli M, Oertli D, von Holzen U, Adamina

M, Muraro MG, Mengus C, Zajac P, Sconocchia G, Zuber M, Tornillo L, Terracciano L, Spagnoli

GC. GM-CSF Production by Tumor Cells Is Associated with Improved Survival in Colorectal

Cancer. Clin Cancer Res. 2014;20(12):3094-106.

58.   Triner D, Devenport SN, Ramakrishnan SK, Ma X, Frieler RA, Greenson JK, Inohara N,

Nunez G, Colacino JA, Mortensen RM, Shah YM. Neutrophils Restrict Tumor-Associated

Microbiota to Reduce Growth and Invasion of Colon Tumors in Mice. Gastroenterology.

2019;156(5):1467-82.

59.   Ni L, Lu J. Interferon gamma in cancer immunotherapy. Cancer Med. 2018;7(9):4509-16.

60.   Gocher AM, Workman CJ, Vignali DAA. Interferon-gamma: teammate or opponent in the

tumour microenvironment? Nat Rev Immunol. 2022;22(3):158-72.

61.   Briesemeister D, Sommermeyer D, Loddenkemper C, Loew R, Uckert W, Blankenstein T,

Kammertoens T. Tumor rejection by local interferon gamma induction in established tumors is

associated with blood vessel destruction and necrosis. Int J Cancer. 2011;128(2):371-8.

62.   Kammertoens T, Friese C, Arina A, Idel C, Briesemeister D, Rothe M, Ivanov A,

Szymborska A, Patone G, Kunz S, Sommermeyer D, Engels B, Leisegang M, Textor A, Fehling HJ,

Fruttiger M, Lohoff M, Herrmann A, Yu H, Weichselbaum R, Uckert W, Hubner N, Gerhardt H,

Beule D, Schreiber H, Blankenstein T. Tumour ischaemia by interferon-gamma resembles

physiological blood vessel regression. Nature. 2017;545(7652):98-102.

63.   Karagiannidis I, Jerman SJ, Jacenik D, Phinney BB, Yao R, Prossnitz ER, Beswick EJ. G-CSF

and G-CSFR Modulate CD4 and CD8 T Cell Responses to Promote Colon Tumor Growth and Are

Potential Therapeutic Targets. Front Immunol. 2020;11:1885.

64.   Wang QS, Shen SQ, Sun HW, Xing ZX, Yang HL. Interferon-gamma induces autophagy-

associated apoptosis through induction of cPLA2-dependent mitochondrial ROS generation in

colorectal cancer cells. Biochem Biophys Res Commun. 2018;498(4):1058-65.

65.   Kelly J, Minoda Y, Meredith T, Cameron G, Philipp MS, Pellicci DG, Corbett AJ, Kurts C,

Gray DH, Godfrey DI, Kannourakis G, Berzins SP. Chronically stimulated human MAIT cells are

unexpectedly potent IL-13 producers. Immunol Cell Biol. 2019;97(8):689-99.

66.   Cole S, Murray J, Simpson C, Okoye R, Tyson K, Griffiths M, Baeten D, Shaw S, Maroof A.

Interleukin (IL)-12 and IL-18 Synergize to Promote MAIT Cell IL-17A and IL-17F Production

Independently of IL-23 Signaling. Front Immunol. 2020;11:585134.

67.   Li K, Vorkas CK, Chaudhry A, Bell DL, Willis RA, Rudensky A, Altman JD, Glickman MS,

Aube J. Synthesis, stabilization, and characterization of the MR1 ligand precursor 5-amino-6-D-

ribitylaminouracil (5-A-RU). PLoS One. 2018;13(2):e0191837.

68.   Corbett AJ, Eckle SB, Birkinshaw RW, Liu L, Patel O, Mahony J, Chen Z, Reantragoon R,

Meehan B, Cao H, Williamson NA, Strugnell RA, Van Sinderen D, Mak JY, Fairlie DP, Kjer-Nielsen

L, Rossjohn J, McCluskey J. T-cell activation by transitory neo-antigens derived from distinct

microbial pathways. Nature. 2014;509(7500):361-5.

Figure 1. MAIT cell peritumoral injection leads to inhibition of MC38 cells-derived tumor growth

A) Experimental setup of MAIT cell expansion in C57BL/6J mice and MAIT cell peritumoral injection in MC38-bearing RAG1$^{-/-}$ mice. MAIT cells were expanded in C57BL/6J mice for 7 days and sorted MAIT cells were injected peritumorally in RAG1$^{-/-}$ mice 1 day after MC38 cell injection. Control: n = 11 males. MAIT: n = 10 (9 males and 1 female). Tumor volume was monitored during the course of the experiment. Tumors were extracted at the end of the experiment for weight measurement, qRT-PCR analysis and Luminex multiplex analysis. B) MC38 cells-derived tumor volume in mm^3 over time from 2 independent experiments. C-D) Tumor measurements. Each dot represents a tumor sample and data is shown as median with interquartile range. C) Volume of extracted tumors in mm^3 and D) Weight of extracted tumors in mg. *$p<0.05$ by two-tailed Mann-Whitney U test. E) Representative photos of extracted tumors from each experimental group. Figure created using BioRender.com

Figure 2. MAIT cell peritumoral injection induces changes in cytokines produced within

tumors.

Comparison of (A) caspase activity and (B – E) concentration of various cytokines and

chemokines released from extracted tumors between control group and MAIT group. A)

Caspase activity is measured in arbitrary fluorescent unit (AFU). B) GM-CSF, C) eotaxin, D) IFN-γ,

and E) IL-17. 25 μL of supernatants from 18-hour culture of 8 mg tumor pieces were analyzed

using Luminex multiplex assay. Each dot represents an individual sample of supernatant and

data is shown as median with interquartile range. *p<0.05 **p<0.01 by two-tailed Mann-

Whitney U test.

Figure 3. Activation of MAIT cells enhance killing against colon cancer cells.

A) Experimental setup of human whole leukocyte isolation and killing assay co-culture with

COLO 205. B) Percent specific lysis of COLO 205 co-cultured with human whole leukocytes with

or without 5-A-RU/MGO stimulation using a 4-hour Calcein Release Assay. C) Frequency of

CFSE+ cells after overnight co-culture of human whole leukocytes with COLO 205 with or

without 5-A-RU/MGO stimulation for 16 hours by flow cytometry. Figure created using

BioRender.com. *p<0.05 by Wilcoxon ranked test.

Figure 4. Stimulation of human peripheral leukocytes with MAIT ligand induces eotaxin-1

production in MAIT cells and activation of eosinophils in an MR-1 dependent manner.

A) Supernatant concentration of various cytokines and chemokines detected in overnight cultures of human whole leukocytes with or without 5-A-RU/MGO stimulation analyzed by Luminex multiplex assay. Each dot represents an individual sample of supernatant from a healthy donor and data is shown as median with interquartile range. B) Flow cytometry analysis of CD69 and eotaxin-1 expression in MAIT cells. C) Flow cytometry analysis of CD69 and granzyme A expression in eosinophils. (B and C) Expression of each marker is shown as MFI (median fluorescence intensity). Each dot represents a sample from an individual healthy donor.

**p<0.01 *** p<0.001 **** p<0.0001 by Wilcoxon ranked test.